

### Electrovalent bonding

- Which forms a crystal of  $NaCl$  [CPMT 1972; NCERT 1976; DPMT 1996]
  - $NaCl$  molecules
  - $Na^+$  and  $Cl^-$  ions
  - $Na$  and  $Cl$  atoms
  - None of the above
- When sodium and chlorine reacts then [NCERT 1973]
  - Energy is released and ionic bond is formed
  - Energy is released and a covalent bond is formed
  - Energy is absorbed and ionic bond is formed
  - Energy is absorbed and covalent bond is formed
- Which one is least ionic in the following compounds [CPMT 1976; BHU 1998]
  - $AgCl$
  - $KCl$
  - $BaCl_2$
  - $CaCl_2$
- The electronic configuration of four elements  $L$ ,  $P$ ,  $Q$  and  $R$  are given in brackets  
 $L(1s^2, 2s^2 2p^4)$ ;  $Q(1s^2, 2s^2 2p^6, 3s^2 3p^5)$   
 $P(1s^2, 2s^2 2p^6, 3s^1)$ ;  $R(1s^2, 2s^2 2p^6, 3s^2)$   
 The formulae of ionic compounds that can be formed between these elements are [NCERT 1983]
  - $L_2P$ ,  $RL$ ,  $PQ$  and  $R_2Q$
  - $LP$ ,  $RL$ ,  $PQ$  and  $RQ$
  - $P_2L$ ,  $RL$ ,  $PQ$  and  $RQ_2$
  - $LP$ ,  $R_2L$ ,  $P_2Q$  and  $RQ$
- Electrovalent compound's [MP PMT 1984]
  - Melting points are low
  - Boiling points are low
  - Conduct current in fused state
  - Insoluble in polar solvent
- A electrovalent compound is made up of [CPMT 1978, 81; MNR 1979]
  - Electrically charged molecules
  - Neutral molecules
  - Neutral atoms
  - Electrically charged atoms or group of atoms
- Electrovalent bond formation depends on
  - Ionization energy
  - Electron affinity
  - Lattice energy
  - All the three above
- In the following which substance will have highest boiling point [NCERT 1973; MP PMT 1987]
  - $He$
  - $CsF$
  - $NH_3$
  - $CHCl_3$
- An atom of sodium loses one electron and chlorine atom accepts one electron. This result the formation of sodium chloride molecule. This type of molecule will be [MP PMT 1987]
  - Coordinate
  - Covalent
  - Electrovalent
  - Matallic bond
- Formula of a metallic oxide is  $MO$ . The formula of its phosphate will be [CPMT 1986, 93]
  - $M_2(PO_4)_2$
  - $M(PO_4)$
  - $M_2PO_4$
  - $M_3(PO_4)_2$
- From the following which group of elements easily forms cation
  - $F$ ,  $Cl$ ,  $Br$
  - $Li$ ,  $Na$ ,  $K$
  - $O$ ,  $S$ ,  $Se$
  - $N$ ,  $P$ ,  $As$
- Which type of compounds show high melting and boiling points
  - Electrovalent compounds
  - Covalent compounds

- (c) Coordinate compounds  
(d) All the three types of compounds have equal melting and boiling points
13. Lattice energy of an ionic compound depends upon [AIEEE 2005]  
(a) Charge on the ion only  
(b) Size of the ion only  
(c) Packing of ions only  
(d) Charge on the ion and size of the ion
14. In the given bonds which one is most ionic [EAMCET 1980]  
(a)  $Cs-Cl$  (b)  $Al-Cl$   
(c)  $C-Cl$  (d)  $H-Cl$
15. Element  $x$  is strongly electropositive and  $y$  is strongly electronegative. Both elements are univalent, the compounds formed from their combination will be [IIT 1980]  
(a)  $x^+y^-$  (b)  $x^-y^+$   
(c)  $x-y$  (d)  $x \rightarrow y$
16. In the formation of  $NaCl$  from  $Na$  and  $Cl$  [CPMT 1985]  
(a) Sodium and chlorine both give electrons  
(b) Sodium and chlorine both accept electrons  
(c) Sodium loses electron and chlorine accepts electron  
(d) Sodium accepts electron and chlorine loses electron
17. Which of the following is an electrovalent linkage [CPMT 1974; DPMT 1984, 91; AFMC 1988]  
(a)  $CH_4$  (b)  $MgCl_2$   
(c)  $SiCl_4$  (d)  $BF_3$
18. Electrovalent compounds do not have [CPMT 1991]  
(a) High M.P. and Low B.P. (b) High dielectric constant  
(c) High M.P. and High B.P. (d) High polarity
19. Many ionic crystals dissolve in water because [NCERT 1982]  
(a) Water is an amphiprotic solvent  
(b) Water is a high boiling liquid  
(c) The process is accompanied by a positive heat of solution  
(d) Water decreases the interionic attraction in the crystal lattice due to solvation
20. The electronic structure of four elements A, B, C, D are  
(A)  $1s^2$  (B)  $1s^2, 2s^2 2p^2$   
(C)  $1s^2, 2s^2 2p^5$  (D)  $1s^2, 2s^2 2p^6$   
The tendency to form electrovalent bond is largest in [MNR 1987, 95]  
(a)  $A$  (b)  $B$   
(c)  $C$  (d)  $D$
21. Chloride of metal is  $MCl_2$ . The formula of its phosphate will be  
(a)  $M_2PO_4$  (b)  $M_3(PO_4)_2$   
(c)  $M_2(PO_4)_3$  (d)  $MPO_4$
22. The phosphate of a metal has the formula  $MPO_4$ . The formula of its nitrate will be [CPMT 1971; MP PMT 1996]  
(a)  $MNO_3$  (b)  $M_2(NO_3)_2$   
(c)  $M(NO_3)_2$  (d)  $M(NO_3)_3$
23. In the transition of  $Zn$  atoms to  $Zn^{++}$  ions there is a decrease in the [CPMT 1972]  
(a) Number of valency electrons  
(b) Atomic weight  
(c) Atomic number  
(d) Equivalent weight
24. Phosphate of a metal  $M$  has the formula  $M_3(PO_4)_2$ . The formula for its sulphate would be [CPMT 1973; MP PMT 1996]  
(a)  $MSO_4$  (b)  $M(SO_4)_2$   
(c)  $M_2(SO_4)_3$  (d)  $M_3(SO_4)_2$
25. The molecular formula of chloride of a metal  $M$  is  $MCl_3$ . The formula of its carbonate would be [CPMT 1987]  
(a)  $MCO_3$  (b)  $M_2(CO_3)_3$   
(c)  $M_2CO_3$  (d)  $M(CO_3)_2$
26. Sodium chloride easily dissolves in water. This is because [NCERT 1972; BHU 1973]  
(a) It is a covalent compound  
(b) Salt reacts with water  
(c) It is a white substance  
(d) Its ions are easily solvated
27. When  $NaCl$  is dissolved in water the sodium ion becomes [NCERT 1974; CPMT 1989; MP PMT 1999]  
(a) Oxidized (b) Reduced  
(c) Hydrolysed (d) Hydrated
28. Solid  $NaCl$  is a bad conductor of electricity since [AFMC 1980]  
(a) In solid  $NaCl$  there are no ions  
(b) Solid  $NaCl$  is covalent  
(c) In solid  $NaCl$  there is no motion of ions  
(d) In solid  $NaCl$  there are no electrons
29. Favourable conditions for electrovalency are  
(a) Low charge on ions, large cation, small anion  
(b) High charge on ions, small cation, large anion  
(c) High charge on ions, large cation, small anion  
(d) Low charge on ions, small cation, large anion
30. The sulphate of a metal has the formula  $M_2(SO_4)_3$ . The formula for its phosphate will be [DPMT 1982; CPMT 1972; MP PMT 1995]  
(a)  $M(HPO_4)_2$  (b)  $M_3(PO_4)_2$   
(c)  $M_2(PO_4)_3$  (d)  $MPO_4$
31. Ionic bonds are usually formed by combination of elements with [CBSE PMT 1999]  
(a) High ionisation potential and low electron affinity  
(b) Low ionisation potential and high electron affinity  
(c) High ionisation potential and high electron affinity  
(d) Low ionisation potential and low electron affinity
32. Molten sodium chloride conducts electricity due to the presence of  
(a) Free electrons  
(b) Free ions  
(c) Free molecules  
(d) Atoms of sodium and chlorine



33. The phosphate of a metal has the formula  $MHPO_4$ . The formula of its chloride would be  
[NCERT 1974; CPMT 1977]  
(a)  $MCl$  (b)  $MCl_2$   
(c)  $MCl_3$  (d)  $M_2Cl_3$
34. A number of ionic compounds e.g.  $AgCl$ ,  $CaF_2$ ,  $BaSO_4$  are insoluble in water. This is because [NCERT 1984]  
(a) Ionic compounds do not dissolve in water  
(b) Water has a high dielectric constant  
(c) Water is not a good ionizing solvent  
(d) These molecules have exceptionally high attractive forces in the lattice
35. What is the nature of chemical bonding between  $Cs$  and  $F$   
[MP PMT 1987; CPMT 1976]  
(a) Covalent (b) Ionic  
(c) Coordinate (d) Metallic
36. Which one of the following compound is ionic [MNR 1985]  
(a)  $KCl$  (b)  $CH_4$   
(c) Diamond (d)  $H_2$
37. Which of the following compound has electrovalent linkage [CPMT 1983, 84, 93]  
(a)  $CH_3Cl$  (b)  $NaCl$   
(c)  $CH_4$  (d)  $Cl_2$
38. An ionic compound is generally a [MADT Bihar 1981]  
(a) Good electrolyte (b) Weak electrolyte  
(c) Non-electrolyte (d) Neutral
39. What metals combine with non-metals, the metal atom tends to  
(a) Lose electrons  
(b) Gain electrons  
(c) Remain electrically neutral  
(d) None of these
40. Chemical formula for calcium pyrophosphate is  $Ca_2P_2O_7$ . The formula for ferric pyrophosphate will be [NCERT 1977]  
(a)  $Fe_3(P_2O_7)_3$  (b)  $Fe_4P_4O_{14}$   
(c)  $Fe_4(P_2O_7)_3$  (d)  $Fe_3PO_4$
41. Among the bonds formed by a chlorine atom with atoms of hydrogen, chlorine, sodium and carbon, the strongest bond is formed between [EAMCET 1988; MP PMT 1993]  
(a)  $H-Cl$  (b)  $Cl-Cl$   
(c)  $Na-Cl$  (d)  $C-Cl$
42. Which of the following is least soluble [CPMT 1989]  
(a)  $BeF_2$  (b)  $SrF_2$   
(c)  $CaF_2$  (d)  $MgF_2$
43. Which of the following halides has maximum melting point  
(a)  $NaCl$  (b)  $NaBr$   
(c)  $NaI$  (d)  $NaF$
44. The high melting point and insolubility in organic solvents of sulphuric acid are due to its ..... structure. [IIT 1994]  
(a) Simple ionic (b) Bipolar ionic  
(c) Cubic (d) Hexagonal
45. Out of the following, which compound will have electrovalent bonding  
(a) Ammonia (b) Water  
(c) Calcium chloride (d) Chloromethane
46. The force which holds atoms together in an electrovalent bond is  
(a) Vander Waal's force  
(b) Dipole attraction force  
(c) Electrostatic force of attraction  
(d) All the above
47. The main reaction during electrovalent bond formation is  
(a) Redox reaction (b) Substitution reaction  
(c) Addition reaction (d) Elimination reaction
48. Electrovalent compounds are [CPMT 1996]  
(a) Good conductor of electricity  
(b) Polar in nature  
(c) Low M.P. and low B.P.  
(d) Easily available
49. Ionic compounds do not have [RPMT 1997]  
(a) Hard and brittle nature  
(b) High melting and boiling point  
(c) Directional properties  
(d) Soluble in polar solvents
50. Highest melting point would be of [RPMT 1999]  
(a)  $He$  (b)  $CsCl$   
(c)  $NH_3$  (d)  $CHCl_3$
51. What is the effect of more electronegative atom on the strength of ionic bond [AMU 1999]  
(a) Decreases (b) Increases  
(c) Decreases slowly (d) Remains the same
52. An element  $X$  with the electronic configuration  $1s^2, 2s^2 2p^6, 3s^2$  would be expected to form the chloride with the formula  
(a)  $XCl_3$  (b)  $XCl_2$   
(c)  $XCl$  (d)  $X_2Cl$
53. Two elements have electronegativity of 1.2 and 3.0. Bond formed between them would be [CPMT 1982; DCE 2000]  
(a) Ionic (b) Polar covalent  
(c) Co-ordinate (d) Metallic
54. Which of the following is least ionic [MP PET 2002]  
(a)  $C_2H_5Cl$  (b)  $KCl$   
(c)  $BaCl_2$  (d)  $C_6H_5N^+H_3Cl^-$
55. Which type of bonding exists in  $Li_2O$  and  $CaF_2$  respectively [RPET 2000]  
(a) Ionic, ionic (b) Ionic, covalent  
(c) Covalent, ionic (d) Coordinate, ionic
56. An atom with atomic number 20 is most likely to combine chemically with the atom whose atomic number is [BHU 2000]  
(a) 11 (b) 14  
(c) 16 (d) 10
57. Bond formed in crystal by anion and cation is [CBSE PMT 2000]  
(a) Ionic (b) Metallic

- (c) Covalent (d) Dipole
58. Atoms or group of atoms which are electrically charged are known  
(a) Anions (b) Cations  
(c) Ions (d) Atoms
59. Which one is the strongest bond [Pb. PMT 2001]  
(a)  $Br-F$  (b)  $F-F$   
(c)  $Cl-F$  (d)  $Br-Cl$
60. The interionic attraction depends on interaction of [Kerala CET (Med.) 2002]  
(a) Solute-Solute (b) Solvent-Solvent  
(c) The charges (d) Molecular properties
61. Which of the following compounds is ionic [UPSEAT 2002]  
(a)  $KI$  (b)  $CH_4$   
(c) Diamond (d)  $H_2$
62. Which of the following pairs of species has same electronic configuration [UPSEAT 2002]  
(a)  $Zn^{2+}$  and  $Ni^{2+}$  (b)  $Co^{+3}$  and  $Ni^{4+}$   
(c)  $Co^{2+}$  and  $Ni^{2+}$  (d)  $Ti^{4+}$  and  $V^{3+}$
63. The energy that opposes dissolution of a solvent is [CPMT 2002]  
(a) Hydration energy (b) Lattice energy  
(c) Internal energy (d) Bond energy
64. Which of the following has highest melting point [RPET 2003]  
(a)  $BeCl_2$  (b)  $MgCl_2$   
(c)  $CaCl_2$  (d)  $BaCl_2$
65. Which of the following statements is not true for ionic compounds [RPET 2003]  
(a) High melting point  
(b) Least lattice energy  
(c) Least solubility in organic compounds  
(d) Soluble in water
66. Electrolytes are compound containing [MADT Bihar 1981]  
(a) Electrovalent bond (b) Covalent bond  
(c) Coordinate bond (d) Hydrogen bond
67. Which of the following hydrides are ionic [Roorkee 1999]  
(a)  $CaH_2$  (b)  $BaH_2$   
(c)  $SrH_2$  (d)  $BeH_2$
68. Which of the following conduct electricity in the fused state [Roorkee 2000]  
(a)  $BeCl_2$  (b)  $MgCl_2$   
(c)  $SrCl_2$  (d)  $BaCl_2$
- (c) 6 (d) 10
3. The electronic configuration of four elements are given in brackets  
 $L(1s^2, 2s^2 2p^1)$ ;  $M(1s^2, 2s^2 2p^5)$   
 $Q(1s^2, 2s^2 2p^6, 3s^1)$ ;  $R(1s^2, 2s^2 2p^2)$   
The element that would most readily form a diatomic molecule is [NCERT 1983]  
(a)  $Q$  (b)  $M$   
(c)  $R$  (d)  $L$
4. In covalency [CPMT 1974, 76, 78, 81; AFMC 1982]  
(a) Electrons are transferred  
(b) Electrons are equally shared  
(c) The electron of one atom are shared between two atoms  
(d) None of the above
5. Which compound is highest covalent  
(a)  $LiCl$  (b)  $LiF$   
(c)  $LiBr$  (d)  $LiI$
6. The nature of bonding in graphite is [DPMT 1986; CPMT 1986]  
(a) Covalent (b) Ionic  
(c) Metallic (d) Coordinate
7. Which of the following substances has giant covalent structure [DPMT 1985, 86; NCERT 1975]  
(a) Iodine crystal (b) Solid  $CO_2$   
(c) Silica (d) White phosphorus
8. With which of the given pairs  $CO$  resembles [BHU 2005]  
(a)  $HgCl, CH_4$  (b)  $HgCl, SnCl$   
(c)  $CH, NO$  (d)  $NO, NO_2$
9. The electron pair which forms a bond between two similar non-metallic atoms will be [IIT 1986]  
(a) Dissimilar shared between the two  
(b) By complete transfer from one atom to other  
(c) In a similar spin condition  
(d) Equally shared in between the two
10. For the formation of covalent bond, the difference in the value of electronegativities should be [EAMCET 1982]  
(a) Equal to or less than 1.7 (b) More than 1.7  
(c) 1.7 or more (d) None of these
11. Which type of bond is formed between similar atoms  
(a) Ionic (b) Covalent  
(c) Coordinate (d) Metallic
12. Covalent compounds are generally ..... in water [CPMT 1987]  
(a) Soluble (b) Insoluble  
(c) Dissociated (d) Hydrolysed
13. Which one is the electron deficient compound [AIIMS 1982]  
(a)  $ICl$  (b)  $NH_3$   
(c)  $BCl_3$  (d)  $PCl_3$
14. Which among the following elements has the tendency to form covalent compounds  
(a)  $Ba$  (b)  $Be$   
(c)  $Mg$  (d)  $Ca$
15. Silicon has 4 electrons in the outermost orbit. In forming the bonds  
(a) It gains electrons (b) It loses electrons  
(c) It shares electrons (d) None of these
16. Which of the following occurs when two hydrogen atoms bond with each others  
(a) Potential energy is lowered  
(b) Kinetic energy is lowered

### Covalent bonding

1. The valency of sulphur in sulphuric acid is [NCERT 1974]  
(a) 2 (b) 4  
(c) 6 (d) 8
2. The number of electrons involved in the bond formation of  $N_2$  molecule [IIT 1980; CPMT 1983, 84, 85; CBSE PMT 1992]  
(a) 2 (b) 4
15. Silicon has 4 electrons in the outermost orbit. In forming the bonds  
(a) It gains electrons (b) It loses electrons  
(c) It shares electrons (d) None of these
16. Which of the following occurs when two hydrogen atoms bond with each others  
(a) Potential energy is lowered  
(b) Kinetic energy is lowered

- (c) Electronic motion ceases  
(d) Energy is absorbed
17. A bond with maximum covalent character between non-metallic elements is formed [NCERT 1982]  
(a) Between identical atoms  
(b) Between chemically similar atoms  
(c) Between atoms of widely different electronegativities  
(d) Between atoms of the same size
18. Amongst the following covalent bonding is found in [CPMT 1973]  
(a) Sodium chloride (b) Magnesium chloride  
(c) Water (d) Brass
19. Indicate the nature of bonding in diamond [EAMCET 1980; BHU 1996; KCET 2000]  
(a) Covalent (b) Ionic  
(c) Coordinate (d) Hydrogen
20. Octet rule is not valid for the molecule [IIT 1979; MP PMT 1995]  
(a)  $CO_2$  (b)  $H_2O$   
(c)  $CO$  (d)  $O_2$
21. Which of the following compounds are covalent [IIT 1980; MLNR 1982]  
(a)  $H_2$  (b)  $CaO$   
(c)  $KCl$  (d)  $Na_2S$
22. Indicate the nature of bonding in  $CCl_4$  and  $CaH_2$  [NCERT 1973]  
(a) Covalent in  $CCl_4$  and electrovalent in  $CaH_2$   
(b) Electrovalent in both  $CCl_4$  and  $CaH_2$   
(c) Covalent in both  $CCl_4$  and  $CaH_2$   
(d) Electrovalent in  $CCl_4$  and covalent in  $CaH_2$
23. If the atomic number of element  $X$  is 7, the best electron dot symbol for the element is [NCERT 1973; CPMT 2003]  
(a)  $X \cdot$  (b)  $\cdot X \cdot$   
(c)  $\cdot \overset{\cdot}{X} :$  (d)  $:\overset{\cdot\cdot}{X} \cdot$
24. Which is the most covalent [AFMC 1982]  
(a)  $C-O$  (b)  $C-Br$   
(c)  $C-S$  (d)  $C-F$
25. The covalent compound  $HCl$  has the ionic character as [EAMCET 1980]  
(a) The electronegativity of hydrogen is greater than that of chlorine  
(b) The electronegativity of hydrogen is equal to that of chlorine  
(c) The electronegativity of chlorine is greater than that of hydrogen  
(d) Hydrogen and chlorine are gases
26. The correct sequence of increasing covalent character is represented by [CBSE PMT 2005]  
(a)  $LiCl < NaCl < BeCl_2$  (b)  $BeCl_2 < NaCl < LiCl$   
(c)  $NaCl < LiCl < BeCl_2$  (d)  $BeCl_2 < LiCl < NaCl$
27. Bond energy of covalent  $O-H$  bond in water is [EAMCET 1982]  
(a) Greater than bond energy of  $H-H$  bond  
(b) Equal to bond energy of  $H-H$  bond  
(c) Less than bond energy of  $H-H$  bond  
(d) None of these
28. Solid  $CH_4$  is [DPMT 1983]  
(a) Molecular solid (b) Ionic solid  
(c) Pseudo solid (d) Does not exist
29. A covalent bond is likely to be formed between two elements which  
(a) Have similar electronegativities  
(b) Have low ionization energies  
(c) Have low melting points  
(d) Form ions with a small charge
30. The bond between two identical non-metal atoms has a pair of electrons [CPMT 1986]  
(a) Unequally shared between the two  
(b) Transferred fully from one atom to another  
(c) With identical spins  
(d) Equally shared between them
31. The valency of phosphorus in  $H_3PO_4$  is [DPMT 1984]  
(a) 2 (b) 5  
(c) 4 (d) 1
32. Which of the following substances has covalent bonding [AMU 1985]  
(a) Germanium (b) Sodium chloride  
(c) Solid neon (d) Copper
33. The covalency of nitrogen in  $HNO_3$  is [CPMT 1987]  
(a) 0 (b) 3  
(c) 4 (d) 5
34. Hydrogen chloride molecule contains a [CPMT 1984]  
(a) Covalent bond (b) Double bond  
(c) Coordinate bond (d) Electrovalent bond
35. As compared to covalent compounds, electrovalent compounds generally have [CPMT 1990, 94; MP PMT 1997]  
(a) Low melting points and low boiling points  
(b) Low melting points and high boiling points  
(c) High melting points and low boiling points  
(d) High melting points and high boiling points
36. The interatomic distances in  $H_2$  and  $Cl_2$  molecules are 74 and 198 pm respectively. The bond length of  $HCl$  is [MP PET 1993]  
(a) 272 pm (b) 136 pm  
(c) 124 pm (d) 248 pm
37. On analysis, a certain compound was found to contain iodine and oxygen in the ratio of 254 gm of iodine and 80 gm of oxygen.

The atomic mass of iodine is 127 and that of oxygen is 16. Which of the following is the formula of the compound

- (a)  $IO$  (b)  $I_2O$   
(c)  $I_5O_2$  (d)  $I_2O_5$

38. Ionic and covalent bonds are present in  
[CBSE PMT 1990; MNR 1990; KCET 2000; UPSEAT 2001]

- (a)  $CCl_4$  (b)  $CaCl_2$   
(c)  $NH_4Cl$  (d)  $H_2O$

39. Highest covalent character is found in [EAMCET 1992]

- (a)  $CaF_2$  (b)  $CaCl_2$   
(c)  $CaBr_2$  (d)  $CaI_2$

40. Among the following which property is commonly exhibited by a covalent compound [MP PET 1994]

- (a) High solubility in water  
(b) High electrical conductance  
(c) Low boiling point  
(d) High melting point

41. Atoms in the water molecule are linked by [MP PAT 1996]

- (a) Electrovalent bond  
(b) Covalent bond  
(c) Coordinate covalent bond  
(d) Odd electron bond

42. Which is the correct electron dot structure of  $N_2O$  molecule  
[MP PET 1996]

- (a)  $:N = N = \ddot{O}$  (b)  $:N \equiv N^+ - \ddot{O}^-$   
(c)  $\ddot{N} = \ddot{N} = \ddot{O}$  (d)  $:N = N = \ddot{O}:$

43. A covalent bond between two atoms is formed by which of the following [MP PMT 1996]

- (a) Electron nuclear attraction  
(b) Electron sharing  
(c) Electron transfer  
(d) Electrostatic attraction

44. The electronic configuration of a metal  $M$  is  $1s^2, 2s^2 2p^6, 3s^1$ . The formula of its oxides will be [MP PET/PMT 1998]

- (a)  $MO$  (b)  $M_2O$   
(c)  $M_2O_3$  (d)  $MO_2$

45. Which of the following statements regarding covalent bond is not true [MP PET/PMT 1998]

- (a) The electrons are shared between atoms  
(b) The bond is non-directional  
(c) The strength of the bond depends upon the extent of overlapping  
(d) The bond formed may or may not be polar

46. If the electronic configuration of  $M = 2, 8, 3$  and that of  $A = 2, 8, 7$ , the formula of the compound is [Bihar MEE 1996]

- (a)  $M_2A_3$  (b)  $MA_2$   
(c)  $M_2A$  (d)  $MA_3$

(e)  $M_3A$

47. The table shown below gives the bond dissociation energies ( $E_{diss}$ ) for single covalent bonds of carbon (C) atoms with element A, B, C and D. Which element has the smallest atoms [CBSE PMT 1994]

Bond	$E_{diss} (kJ mol^{-1})$
$C-A$	240
$C-B$	328
$C-C$	276
$C-D$	485

- (a) A (b) B  
(c) C (d) D

48. If a molecule  $X_2$  has a triple bond, then  $X$  will have the electronic configuration [CET Pune 1998]

- (a)  $1s^2 2s^2 2p^5$  (b)  $1s^2 2s^2 2p^3$   
(c)  $1s^2 2s^1$  (d)  $1s^2 2s^2 2p^1$

49. Which of the following compounds does not follow the octet rule for electron distribution [CET Pune 1998]

- (a)  $PCl_5$  (b)  $PCl_3$   
(c)  $H_2O$  (d)  $PH_3$

50. The valency of  $A = 3$  and  $B = 2$ , then the compound is [Bihar MEE 1997]

- (a)  $A_2B_3$  (b)  $A_3B_2$   
(c)  $A_3B_3$  (d)  $A_2B_2$   
(e) None of these

51. The number of electrons shared by each outermost shell of  $N_2$  is

- (a) 2 (b) 3  
(c) 4 (d) 5

52. Which of the following substances when dissolved in water will give a solution that does not conduct electricity [JIPMER 1999]

- (a) Hydrogen chloride (b) Potassium hydroxide  
(c) Sodium acetate (d) Urea

53. Which of the following atoms has minimum covalent radius [DPMT 2000]

- (a) B (b) C  
(c) N (d) Si

54. Boron form covalent compound due to [Pb. PMT 2000]

- (a) Small size (b) Higher ionization energy  
(c) Lower ionization energy (d) Both (a) and (b)

55. Two elements X and Y have following electron configurations

$$X = 1s^2, 2s^2 2p^6, 3s^2 3p^6, 4s^2$$

$$\text{and } Y = 1s^2, 2s^2 2p^6, 3s^2 3p^6$$

The compound formed by combination of X and Y is [DPMT 2001]

- (a)  $XY_5$  (b)  $X_2Y_5$   
(c)  $X_5Y_3$  (d)  $XY_2$

56. Covalent compounds have low melting point because [KCET 2002]

- (a) Covalent bond is less exothermic  
(b) Covalent molecules have definite shape  
(c) Covalent bond is weaker than ionic bond



- (d) Covalent molecules are held by weak Vander Waal's force of attraction
57. *p* and *n*-type of semiconductors are formed due to [UPSEAT 2002]  
 (a) Covalent bonds (b) Metallic bonds  
 (c) Ionic bonds (d) Co-ordinate bond
58. Which of the following is Lewis acid [RPET 2003]  
 (a)  $BF_3$  (b)  $NH_3$   
 (c)  $PH_3$  (d)  $SO_2$
59. Among the species :  $CO_2$ ,  $CH_3COO^-$ ,  $CO$ ,  $CO_3^{2-}$ ,  $HCHO$  which has the weakest carbon-oxygen bond [Kerala PMT 2004]  
 (a)  $CO_2$  (b)  $CH_3COO^-$   
 (c)  $CO$  (d)  $CO_3^{2-}$   
 (e)  $HCHO$
60. Valency of sulphur in  $Na_2S_2O_3$  is [DPMT 1984]  
 (a) Two (b) Three  
 (c) Four (d) Six
61. The acid having  $O-O$  bond is [IIT JEE Screening 2004]  
 (a)  $H_2S_2O_3$  (b)  $H_2S_2O_6$   
 (c)  $H_2S_2O_8$  (d)  $H_2S_4O_6$
62. The following salt shows maximum covalent character [UPSEAT 2004]  
 (a)  $AlCl_3$  (b)  $MgCl_2$   
 (c)  $CsCl$  (d)  $LaCl_3$
63. Which type of bond is present in  $H_2S$  molecule [MHCET 2003; Pb CET 2001]  
 (a) Ionic bond (b) Covalent bond  
 (c) Co-ordinate (d) All of three
64.  $H_2S$  is more acidic than  $H_2O$ , due to [BVP 2004]  
 (a)  $O$  is more electronegative than  $S$   
 (b)  $O-H$  bond is stronger than  $S-H$  bond  
 (c)  $O-H$  bond is weaker than  $S-H$  bond  
 (d) None of these
65. Which of the following has covalent bond [AFMC 1988; DCE 2004]  
 (a)  $Na_2S$  (b)  $AlCl_3$   
 (c)  $NaH$  (d)  $MgCl_2$
66. The following element forms a molecule with eight its own weight atoms [MHCET 2004]  
 (a)  $Si$  (b)  $S$   
 (c)  $Cl$  (d)  $P$
67. In  $H_2O_2$ , the two oxygen atoms have  
 (a) Electrovalent bond (b) Covalent bond  
 (c) Coordinate bond (d) No bond
68. Carbon has a valency of 2 in  $CO$  and 4 in  $CO_2$  and  $CH_4$ . Its valency in acetylene ( $C_2H_2$ ) is [NCERT 1971]  
 (a) 1 (b) 2  
 (c) 3 (d) 4
69. Number of electrons in the valence orbit of nitrogen in an ammonia molecule are  
 (a) 8 (b) 5  
 (c) 6 (d) 7
70. Hydrogen atoms are held together to form hydrogen molecules by  
 (a) Hydrogen bond (b) Ionic bond  
 (c) Covalent bond (d) Dative bond
71. Strongest bond is [AFMC 1987]  
 (a)  $C-C$  (b)  $C-H$   
 (c)  $C-N$  (d)  $C-O$
72. The major binding force of diamond, silicon and quartz is [Kerala CET (Med.) 2002]  
 (a) Electrostatic force (b) Electrical attraction  
 (c) Co-valent bond force (d) Non-covalent bond force
73. Multiple covalent bonds exist in a molecule of [NCERT 1973]  
 (a)  $H_2$  (b)  $F_2$   
 (c)  $C_2H_4$  (d)  $N_2$
74. Which of the following does not obey the octet rule [EAMCET 1993]  
 (a)  $CO$  (b)  $NH_3$   
 (c)  $H_2O$  (d)  $PCl_5$
75. Which of the following statements is correct for covalent bond  
 (a) Electrons are shared between two atoms  
 (b) It may be polar or non-polar  
 (c) Direction is non-polar  
 (d) Valency electrons are attracted
76. Among  $CaH_2$ ,  $NH_3$ ,  $NaH$  and  $B_2H_6$ , which are covalent hydride [Orissa JEE 2005]  
 (a)  $NH_3$  and  $B_2H_6$  (b)  $NaH$  and  $CaH_2$   
 (c)  $NaH$  and  $NH_3$  (d)  $CaH_2$  and  $B_2H_6$

### Co-ordinate or Dative bonding

1. Which species has the maximum number of lone pair of electrons on the central atom? [IIT 2005]  
 (a)  $[ClO]^-$  (b)  $XeF_4$   
 (c)  $SF_6$  (d)  $[I]^-$
2. A simple example of a coordinate covalent bond is exhibited by  
 (a)  $C_2H_2$  (b)  $H_2SO_4$   
 (c)  $NH_3$  (d)  $HCl$
3. The bond that exists between  $NH_3$  and  $BF_3$  is called [AFMC 1982; MP PMT 1985; MNR 1994; KCET 2000; MP PET 2001; UPSEAT 2001]  
 (a) Electrovalent (b) Covalent  
 (c) Coordinate (d) Hydrogen
4. Which of the following does not have a coordinate bond [MADT Bihar 1984]  
 (a)  $SO_2$  (b)  $HNO_3$   
 (c)  $H_2SO_3$  (d)  $HNO_2$
5. Coordinate covalent compounds are formed by [CPMT 1990, 94]  
 (a) Transfer of electrons (b) Sharing of electrons  
 (c) Donation of electrons (d) None of these process
6. In the coordinate valency [CPMT 1989]  
 (a) Electrons are equally shared by the atoms  
 (b) Electrons of one atom are shared with two atoms  
 (c) Hydrogen bond is formed

- (d) None of the above [DPMT 1985]
7. Which of the following contains a coordinate covalent bond [MNR 1990; IIT 1986]
- (a)  $N_2O_5$  (b)  $BaCl_2$   
(c)  $HCl$  (d)  $H_2O$
8. A coordinate bond is formed when an atom in a molecule has
- (a) Electric charge on it  
(b) All its valency electrons shared  
(c) A single unshared electron  
(d) One or more unshared electron pair
9. Which has a coordinate bond [RPMT 1997]
- (a)  $SO_3^{2-}$  (b)  $CH_4$   
(c)  $CO_2$  (d)  $NH_3$
10. The compound containing co-ordinate bond is [AFMC 1999; Pb. CET 2002]
- (a)  $O_3$  (b)  $SO_3$   
(c)  $H_2SO_4$  (d) All of these
11. The number of dative bonds in sulphuric acid molecules is [MP PET 2002]
- (a) 0 (b) 1  
(c) 2 (d) 4
12. Which of the following compounds has coordinate (dative) bond
- (a)  $CH_3NC$  (b)  $CH_3OH$   
(c)  $CH_3Cl$  (d)  $NH_3$
13. The structure of orthophosphoric acid is [KCET 2003]
- (a) 
$$\begin{array}{c} O \\ \uparrow \\ H-O-P-O-H \\ | \\ O \\ | \\ H \end{array}$$
 (b) 
$$\begin{array}{c} H \\ | \\ O \leftarrow P-O-H \\ | \\ O \\ | \\ H \end{array}$$
  
(c) 
$$\begin{array}{c} H \\ | \\ O \leftarrow P-O-H \\ | \\ H \end{array}$$
 (d) 
$$\begin{array}{c} O \\ \uparrow \\ H-O-P=O \end{array}$$
14. What is the nature of the bond between B and O in  $(C_2H_5)_2OBH_3$  [Orissa JEE 2003]
- (a) Covalent (b) Co-ordinate covalent  
(c) Ionic bond (d) Banana shaped bond
15. Sulphuric acid provides a example of [Kerala CET (Med.) 2002]
- (a) Co-ordinate bonds  
(b) Non-covalent compound  
(c) Covalent and co-ordinate bond  
(d) Non-covalent ion
- (a)  $BF_3$  (b)  $CCl_4$   
(c)  $BeCl_2$  (d) All of these
3. Which molecule has the largest dipole moment [CPMT 1991]
- (a)  $HI$   
(c)  $HBr$  (d)  $HF$
4. The unequal sharing of bonded pair of electrons between two atoms in a molecule causes [EAMCET 1986]
- (a) Dipole  
(b) Radical formation  
(c) Covalent bond  
(d) Decomposition of molecule
5. Which of the following will show least dipole character [NCERT 1975; Kurukshetra CEE 1998]
- (a) Water (b) Ethanol  
(c) Ethane (d) Ether
6. Which of the following molecules will show dipole moment [NCERT 1972, 74; DPMT 1985]
- (a) Methane (b) Carbon tetrachloride  
(c) Chloroform (d) Carbon dioxide
7. Which of the following compounds possesses the dipole moment [NCERT 1978; EAMCET 2003]
- (a) Water (b) Boron trifluoride  
(c) Benzene (d) Carbon tetrachloride
8. Which bond angle  $\theta$  would result in the maximum dipole moment for the triatomic molecule  $YXY$  [AIIMS 1980]
- (a)  $\theta = 90^\circ$  (b)  $\theta = 120^\circ$   
(c)  $\theta = 150^\circ$  (d)  $\theta = 180^\circ$
9. Which of the following would have a permanent dipole moment [CBSE PMT 2000]
- (a)  $BF_3$  (b)  $SiF_4$   
(c)  $SF_4$  (d)  $XeF_4$
10. Carbon tetrachloride has no net dipole moment because of [IIT 1982, 83; MP PMT 1985, 91; EAMCET 1988; AMU 1999]
- (a) Its planar structure  
(b) Its regular tetrahedral structure  
(c) Similar sizes of carbon and chlorine atoms  
(d) Similar electron affinities of carbon and chlorine
11. The molecule which has the largest dipole moment amongst the following [MNR 1983]
- (a)  $CH_4$  (b)  $CHCl_3$   
(c)  $CCl_4$  (d)  $CHI_3$
12. Positive dipole moment is present in [MNR 1986; MP PET 2000]
- (a)  $CCl_4$  (b)  $C_6H_6$   
(c)  $BF_3$  (d)  $HF$
13. The polarity of a covalent bond between two atoms depends upon
- (a) Atomic size (b) Electronegativity  
(c) Ionic size (d) None of the above
14. Pick out the molecule which has zero dipole moment [CPMT 1989; EAMCET 1993; MP PMT 1999]
- (a)  $NH_3$  (b)  $H_2O$   
(c)  $BCl_3$  (d)  $SO_2$

## Dipole moment

1. Which molecules has zero dipole moment [AIIMS 1980, 82, 91; Roorkee 2000; MH CET 2001]
- (a)  $H_2O$  (b)  $CO_2$   
(c)  $HF$  (d)  $HBr$
2. In the following which one have zero dipole moment



15. Zero dipole moment is present in [DPMT 1986; IIT 1987]  
 (a)  $NH_3$  (b)  $H_2O$   
 (c) *cis* 1, 2-dichloroethene (d) *trans* 1, 2-dichloroethene
16. Which of the following is the most polar [AFMC 1988]  
 (a)  $CCl_4$  (b)  $CHCl_3$   
 (c)  $CH_3OH$  (d)  $CH_3Cl$
17. Which one has minimum (nearly zero) dipole moment [IIT Screening 1994; CBSE PMT 1996]  
 (a) Butene-1 (b) *cis* butene-2  
 (c) *trans* butene-2 (d) 2-methyl-1-propene
18. Which one of the following is having zero dipole moment [RPMT 1997; EAMCET 1988; MNR 1991]  
 (a)  $CCl_4$  (b)  $CH_3Cl$   
 (c)  $CH_3F$  (d)  $CHCl_3$
19. Which of the following molecules does not possess a permanent dipole moment [CBSE PMT 1994]  
 (a)  $H_2S$  (b)  $SO_2$   
 (c)  $CS_2$  (d)  $SO_3$
20. Which of the following has zero dipole moment [CPMT 1997; AFMC 1998; CBSE PMT 2001]  
 (a)  $CH_2Cl_2$  (b)  $CH_4$   
 (c)  $NH_3$  (d)  $PH_3$
21. Fluorine is more electronegative than either boron or phosphorus. What conclusion can be drawn from the fact that  $BF_3$  has no dipole moment but  $PF_3$  does [Pb. PMT 1998]  
 (a)  $BF_3$  is not spherically symmetrical but  $PF_3$  is  
 (b)  $BF_3$  molecule must be linear  
 (c) The atomic radius of  $P$  is larger than the atomic radius of  $B$   
 (d) The  $BF_3$  molecule must be planar triangular
22. Which molecule does not show zero dipole moment [RPET 1997, 99]  
 (a)  $BF_3$  (b)  $NH_3$   
 (c)  $CCl_4$  (d)  $CH_4$
23. The dipole moment of  $HBr$  is  $1.6 \times 10^{-30}$  cm and interatomic spacing is  $1\text{Å}$ . The % ionic character of  $HBr$  is [MP PMT 2000]  
 (a) 7 (b) 10  
 (c) 15 (d) 27
24. Non-polar solvent is [RPET 2000]  
 (a) Dimethyl sulphoxide (b) Carbon tetrachloride  
 (c) Ammonia (d) Ethyl alcohol
25. Which shows the least dipole moment [UPSEAT 2001; DPMT 1982]  
 (a)  $CCl_4$  (b)  $CHCl_3$   
 (c)  $CH_3CH_2OH$  (d)  $CH_3COCH_3$
26. Which molecule has zero dipole moment [UPSEAT 2001]  
 (a)  $H_2O$  (b)  $AgI$   
 (c)  $PbSO_4$  (d)  $HBr$
27. The dipole moment is zero for the molecule [IIT 1989; MP PMT 2002]  
 (a) Ammonia (b) Boron trifluoride  
 (c) Sulphur dioxide (d) Water
28.  $N_2$  is less reactive than  $CN^-$  due to [UPSEAT 2003]  
 (a) Presence of more electrons in orbitals  
 (b) Absence of dipole moment  
 (c) Difference in spin quantum no  
 (d) None of these
29. In a polar molecule, the ionic charge is  $4.8 \times 10^{-10}$  e.s.u. If the inter ionic distance is one Å unit, then the dipole moment is  
 (a) 41.8 debye (b) 4.18 debye  
 (c) 4.8 debye (d) 0.48 debye
30. Which of the following is a polar compound [Pb. CET 2000]  
 (a)  $HCl$  (b)  $H_2Se$   
 (c)  $CH_4$  (d)  $HI$
31. Which of the following has no dipole moment [DCE 2002]  
 (a)  $CO_2$  (b)  $SO_3$   
 (c)  $O_3$  (d)  $H_2O$
32. Which of the following is non-polar [DCE 2002]  
 (a)  $PCl_5$  (b)  $PCl_3$   
 (c)  $SF_6$  (d)  $IF_7$
33. Identify the non-polar molecule in the set of compounds given :  $HCl, HF, H_2, HBr$  [UPSEAT 2004]  
 (a)  $H_2$  (b)  $HCl$   
 (c)  $HF, HBr$  (d)  $HBr$
34. Dipole moment is shown by [IIT 1986]  
 (a) 1, 4-dichlorobenzene  
 (b) *cis* 1, 2-dichloroethene  
 (c) *trans* 1, 2-dichloroethene  
 (d) *trans* 1, 2-dichloro-2-pentene
35. If  $HCl$  molecule is completely polarized, so expected value of dipole moment is 6.12D (deby), but experimental value of dipole moment is 1.03D. Calculate the percentage ionic character [Kerala CET 2005]  
 (a) 17 (b) 83  
 (c) 50 (d) Zero  
 (e) 90

### Polarisation and Fajan's rule

1.  $BF_3$  and  $NF_3$  both molecules are covalent, but  $BF_3$  is non-polar and  $NF_3$  is polar. Its reason is [CPMT 1989; NCERT 1980]  
 (a) In uncombined state boron is metal and nitrogen is gas  
 (b)  $B-F$  bond has no dipole moment whereas  $N-F$  bond has dipole moment  
 (c) The size of boron atom is smaller than nitrogen  
 (d)  $BF_3$  is planar whereas  $NF_3$  is pyramidal
2. Which one is polar molecule among the following  
 (a)  $CO_2$  (b)  $CCl_4$

- (c)  $H_2O$  (d)  $CH_4$
3. If the electron pair forming a bond between two atoms  $A$  and  $B$  is not in the centre, then the bond is [AIIMS 1984]  
 (a) Single bond (b) Polar bond  
 (c) Non-polar bond (d)  $\pi$  bond
4. Which of the following liquids is not deflected by a non-uniform electrostatic field [NCERT 1978]  
 (a) Water (b) Chloroform  
 (c) Nitrobenzene (d) Hexane
5. Which of the following is non-polar [EAMCET 1983]  
 (a)  $H_2S$  (b)  $NaCl$   
 (c)  $Cl_2$  (d)  $H_2SO_4$
6. Polarization is the distortion of the shape of an anion by an adjacently placed cation. Which of the following statements is correct [NCERT 1982]  
 (a) Maximum polarization is brought about by a cation of high charge  
 (b) Minimum polarization is brought about by a cation of low radius  
 (c) A large cation is likely to bring about a large degree of polarization  
 (d) A small anion is likely to undergo a large degree of polarization
7. The bonds between  $P$  atoms and  $Cl$  atoms in  $PCl_5$  are likely to be [MP PMT 1987]  
 (a) Ionic with no covalent character  
 (b) Covalent with some ionic character  
 (c) Covalent with no ionic character  
 (d) Ionic with some metallic character
8. Two electrons of one atom  $A$  and two electrons of another atom  $B$  are utilized to form a compound  $AB$ . This is an example of  
 (a) Polar covalent bond (b) Non-polar covalent bond  
 (c) Polar bond (d) Dative bond
9. In which of the following molecule is the covalent bond most polar [AMU 1985; MP PET 2001]  
 (a)  $HI$  (b)  $HBr$   
 (c)  $HCl$  (d)  $H_2$
10. Amongst  $ClF_3$ ,  $BF_3$  and  $NH_3$  molecules the one with non-planar geometry is [MP PMT 1999]  
 (a)  $ClF_3$  (b)  $NH_3$   
 (c)  $BF_3$  (d) None of these
11. Which of the following possesses highest melting point [CPMT 1999]  
 (a) Chlorobenzene (b)  $o$ -dichlorobenzene  
 (c)  $m$ -dichlorobenzene (d)  $p$ -dichlorobenzene
12. The polar molecule among the following is [Orissa JEE 1997]  
 (a)  $CCl_4$  (b)  $CO_2$   
 (c)  $CH_2Cl_2$  (d)  $CH_2 = CH_2$
13. Which of the following have both polar and non-polar bonds  
 (a)  $C_2H_6$  (b)  $NH_4Cl$   
 (c)  $HCl$  (d)  $AlCl_3$
14. Which of the following has a high polarising power [CET Pune 1998]  
 (a)  $Mg^{2+}$  (b)  $Al^{3+}$
- (c)  $Na^+$  (d)  $Ca^{2+}$
15. Maximum covalent character is associated with the compound [RPMT 1999]  
 (a)  $NaI$  (b)  $MgI_2$   
 (c)  $AlCl_3$  (d)  $AlI_3$
16. Polarisability of halide ions increases in the order [DCE 1999]  
 (a)  $F^-, I^-, Br^-, Cl^-$  (b)  $Cl^-, Br^-, I^-, F^-$   
 (c)  $I^-, Br^-, Cl^-, F^-$  (d)  $F^-, Cl^-, Br^-, I^-$
17. According to Fajan's rule, covalent bond is favoured by [AIIMS 1999]  
 (a) Large cation and small anion  
 (b) Large cation and large anion  
 (c) Small cation and large anion  
 (d) Small cation and small anion
18. Which of the following statements is correct [AMU 1999]  
 (a)  $SF_4$  is polar and non-reactive  
 (b)  $SF_6$  is non-polar and very reactive  
 (c)  $SF_6$  is a strong fluorinating agent  
 (d)  $SF_4$  is prepared by fluorinating  $SCl_2$  with  $NaF$
19. Choose the correct statement [RPMT 2000]  
 (a) Amino polarisation is more pronounced by highly charged cation  
 (b) Small cation has minimum capacity to polarise an anion.  
 (c) Small anion has maximum polarizability  
 (d) None of these
20. The  $ICl$  molecule is [DPMT 2001]  
 (a) Purely electrovalent  
 (b) Purely covalent  
 (c) Polar with negative end on iodine  
 (d) Polar with negative end on chlorine
21. Which of the following is a polar compound [AIIMS 2001]  
 (a)  $HF$  (b)  $HCl$   
 (c)  $HNO_3$  (d)  $H_2SO_4$
22. Which of the following has zero dipole moment [MP PMT 2002]  
 (a)  $ClF$  (b)  $PCl_3$   
 (c)  $SiF_4$  (d)  $CFCl_3$
23. Which of the following compounds has least dipole moment [RPET 2003]  
 (a)  $PH_3$  (b)  $CHCl_3$   
 (c)  $NH_3$  (d)  $BF_3$
24. Pauling's electronegativity values for elements are useful in predicting [UPSEAT 2004]  
 (a) Polarity of bonds in molecules  
 (b) Position of elements in electrochemical series  
 (c) Co-ordination number  
 (d) Dipole moment of various molecules
25. Amongst  $LiCl$ ,  $RbCl$ ,  $BeCl_2$  and  $MgCl_2$  the compounds with the greatest and the least ionic character, respectively, are [UPSEAT 2002]  
 (a)  $LiCl$  and  $RbCl$  (b)  $RbCl$  and  $BeCl_2$   
 (c)  $RbCl$  and  $MgCl_2$  (d)  $MgCl_2$  and  $BeCl_2$
26. Bond polarity of diatomic molecule is because of [UPSEAT 2002]

- (a) Difference in electron affinities of the two atoms
- (b) Difference in electronegativities of the two atoms
- (c) Difference in ionisation potential
- (d) All of these

### Overlapping- $\sigma$ and $\pi$ bonds




1. Triple bond in ethyne is formed from  
[MP PMT 1990; NCERT 1979; EAMCET 1978; AMU 1985; CPMT 1988; MADT Bihar 1982; MH CET 2000]
  - (a) Three sigma bonds
  - (b) Three pi bonds
  - (c) One sigma and two pi bonds
  - (d) Two sigma and one pi bond
2. The bond in the formation of fluorine molecule will be  
[MP PMT 1987]
  - (a) Due to  $s-s$  overlapping
  - (b) Due to  $s-p$  overlapping
  - (c) Due to  $p-p$  overlapping
  - (d) Due to hybridization
3. Which type of overlapping results the formation of a  $\pi$  bond  
[DPMT 1981]
  - (a) Axial overlapping of  $s-s$  orbitals
  - (b) Lateral overlapping of  $p-p$  orbitals
  - (c) Axial overlapping of  $p-p$  orbitals
  - (d) Axial overlapping of  $s-p$  orbitals
4. The number and type of bonds between two carbon atoms in calcium carbide are  
[AIEEE 2005]
  - (a) One sigma, one pi
  - (b) One sigma, two pi
  - (c) Two sigma, one pi
  - (d) Two sigma, two pi
5. In a double bond connecting two atoms, there is a sharing of  
[CPMT 1977, 80, 81; NCERT 1975; Bihar MEE 1980; MP PET 1999]
  - (a) 2 electrons
  - (b) 1 electron
  - (c) 4 electrons
  - (d) All electrons
6. Strongest bond is  
[DPMT 1990]
  - (a)  $C-C$
  - (b)  $C=C$
  - (c)  $C\equiv C$
  - (d) All are equally strong
7.  $\pi$  bond is formed  
[JIPMER 2002]
  - (a) By overlapping of atomic orbitals on the axis of nuclei
  - (b) By mutual sharing of pi electron
  - (c) By sidewise overlapping of half filled  $p$ -orbitals
  - (d) By overlapping of  $s$ -orbitals with  $p$ -orbitals
8. The double bond between the two carbon atoms in ethylene consists of  
[NCERT 1981; EAMCET 1979]
  - (a) Two sigma bonds at right angles to each other
  - (b) One sigma bond and one pi bond
  - (c) Two pi bonds at right angles to each other
  - (d) Two pi bonds at an angle of  $60^\circ$  to each other
9. In the series ethane, ethylene and acetylene, the  $C-H$  bond energy is  
[NCERT 1977]
  - (a) The same in all the three compounds
  - (b) Greatest in ethane
  - (c) Greatest in ethylene
  - (d) Greatest in acetylene
10. In a sigma bond
  - (a) Sidewise as well as end to end overlap of orbitals take place

- (b) Sidewise overlap of orbitals takes place
- (c) End to end overlap of orbitals takes place
- (d) None of the above

11. The number of sigma and pi bonds in 1-butene-3-yne are  
[IIT 1989]
  - (a) 5 sigma and 5 pi
  - (b) 7 sigma and 3 pi
  - (c) 8 sigma and 2 pi
  - (d) 6 sigma and 4 pi
12. The most acidic compound among the following is  
[MP PET 1993]
  - (a)  $CH_3CH_2OH$
  - (b)  $C_6H_5OH$
  - (c)  $CH_3COOH$
  - (d)  $CH_3CH_2CH_2OH$
13. Which of the following is not correct  
[CBSE PMT 1990]
  - (a) A sigma bond is weaker than  $\pi$  bond
  - (b) A sigma bond is stronger than  $\pi$  bond
  - (c) A double bond is stronger than a single bond
  - (d) A double bond is shorter than a single bond
14. Strongest bond formed, when atomic orbitals
  - (a) Maximum overlap
  - (b) Minimum overlap
  - (c) Overlapping not done
  - (d) None of them
15. The  $p-p$  orbital overlapping is present in the following molecule
  - (a) Hydrogen
  - (b) Hydrogen bromide
  - (c) Hydrogen chloride
  - (d) Chlorine
16. In  $N_2$  molecule, the atoms are bonded by  
[MP PET 1996; UPSEAT 2001]
  - (a) One  $\sigma$ , Two  $\pi$
  - (b) One  $\sigma$ , One  $\pi$
  - (c) Two  $\sigma$ , One  $\pi$
  - (d) Three  $\sigma$  bonds
17. In which of following there exists a  $p\pi-d\pi$  bonding  
[AFMC 2001]
  - (a) Diamond
  - (b) Graphite
  - (c) Dimethyl amine
  - (d) Trisilylamine
18. Number of bonds in  $SO_2$   
[DCE 2001]
  - (a) Two  $\sigma$  and two  $\pi$
  - (b) Two  $\sigma$  and one  $\pi$
  - (c) Two  $\sigma$ , two  $\pi$  and one lone pair
  - (d) None of these
19. Which of the following has  $p\pi-d\pi$  bonding  
[CBSE 2002]
  - (a)  $NO_3^-$
  - (b)  $CO_3^{2-}$
  - (c)  $BO_3^{3-}$
  - (d)  $SO_3^{2-}$
20. Number of sigma bonds in  $P_4O_{10}$  is  
[AIEEE 2002]
  - (a) 6
  - (b) 7
  - (c) 17
  - (d) 16

### Hybridisation

1. Which molecule is not linear  
[CPMT 1994]
  - (a)  $BeF_2$
  - (b)  $BeH_2$
  - (c)  $CO_2$
  - (d)  $H_2O$
2. The bond angle in water molecule is nearly or Directed bonds in water forms an angle of  
[NCERT 1980; EAMCET 1981; MNR 1983, 85; AIIMS 1982; CPMT 1989; MP PET 1994, 96; MP PET/PMT 1998]
  - (a)  $120^\circ$
  - (b)  $180^\circ$

- (c)  $109^{\circ}28'$  (d)  $104^{\circ}30'$
3. The central atom in a molecule is in  $sp^2$  hybrid state. The shape of molecule will be [MP PMT 1987; CBSE PMT 1989]  
 (a) Pyramidal (b) Tetrahedral  
 (c) Octahedral (d) Trigonal planar
4. Which molecule is linear [MP PMT 1984; IIT 1982, 88; EAMCET 1993; CBSE PMT 1992; MP PET 1995; RPMT 1997]  
 (a)  $NO_2$  (b)  $ClO_2$   
 (c)  $CO_2$  (d)  $H_2S$
5. Which of the following molecules has trigonal planer geometry [CBSE PMT 2005]  
 (a)  $IF_3$  (b)  $PCl_3$   
 (c)  $NH_3$  (d)  $BF_3$
6. A  $sp^3$  hybridized orbital contains [DPMT 1984; BHU 1985; CPMT 1976]  
 (a)  $\frac{1}{4}$  s - character (b)  $\frac{1}{2}$  s - character  
 (c)  $\frac{2}{3}$  s - character (d)  $\frac{3}{4}$  s - character
7. Structure of ammonia is [MP PMT 1987, 89, 91; CPMT 1975, 82; RPMT 1999; JIPMER 2002]  
 (a) Trigonal (b) Tetrahedral  
 (c) Pyramidal (d) Trigonal pyramidal
8. The bond angle in ethylene is [CPMT 1987]  
 (a)  $180^{\circ}$  (b)  $120^{\circ}$   
 (c)  $109^{\circ}$  (d)  $90^{\circ}$
9. Compound formed by  $sp^3d$  hybridization will have structure [BHU 1982; RPMT 1999]  
 (a) Planar (b) Pyramidal  
 (c) Angular (d) Trigonal bipyramidal
10. Which of the following formula does not correctly represent the bonding capacity of the atom involved [CBSE PMT 1990]
- (a)  $\left[ \begin{array}{c} H \\ | \\ H - P - H \\ | \\ H \end{array} \right]$  (b) 
- (c)  (d) 
11. Which of the following statement is not correct [AIIMS 1983]  
 (a) Hybridization is the mixing of atomic orbitals prior to their combining into molecular orbitals  
 (b)  $sp^2$  hybrid orbitals are formed from two  $p$  atomic orbitals and one  $s$  atomic orbital  
 (c)  $d^2sp^3$  hybrid orbitals are directed towards the corners of a regular octahedron  
 (d)  $dsp^3$  hybrid orbitals are all at  $90^{\circ}$  to one another
12. The mode of hybridisation of carbon in  $CO_2$  is [CPMT 1991]  
 (a)  $sp$  (b)  $sp^2$   
 (c)  $sp^3$  (d) None of these
13. In which of the following the central atom does not use  $sp^3$  hybrid orbitals in its bonding [MNR 1992]  
 (a)  $BeF_3^-$  (b)  $OH_3^+$   
 (c)  $NH_2^-$  (d)  $NF_3$
14.  $XeF_2$  involves hybridisation [DPMT 1990]  
 (a)  $sp^3$  (b)  $sp^3d$   
 (c)  $sp^3d^2$  (d) None of these
15. Which of the following hybridisation results in non-planar orbitals  
 (a)  $sp^3$  (b)  $dsp^2$   
 (c)  $sp^2$  (d)  $sp$
16. Octahedral molecular shape exists in ..... hybridisation [DPMT 1990]  
 (a)  $sp^3d$  (b)  $sp^3d^2$   
 (c)  $sp^3d^3$  (d) None of these
17. The electronic structure of molecule  $OF_2$  is a hybrid of  
 (a)  $sp$  (b)  $sp^2$   
 (c)  $sp^3$  (d)  $sd^3$
18. Percentage of s-character in  $sp^3$  hybrid orbital is  
 (a) 25 (b) 50  
 (c) 66 (d) 75
19. Shape of  $XeF_4$  molecule is [BHU 1987; AFMC 1992; CET Pune 1998; Roorkee Qualifying 1998; DCE 2002]  
 (a) Linear (b) Pyramidal  
 (c) Tetrahedral (d) Square planar
20. For which of the following hybridisation the bond angle is maximum  
 (a)  $sp^2$  (b)  $sp$   
 (c)  $sp^3$  (d)  $dsp^2$
21. The  $C-H$  bond distance is the longest in [MNR 1990]  
 (a)  $C_2H_2$  (b)  $C_2H_4$   
 (c)  $C_2H_4Br_2$  (d)  $C_6H_6$
22. The nature of hybridization in  $CH_2Cl-CH_2Cl$  for carbon is  
 (a)  $sp$  (b)  $sp^2$   
 (c)  $sp^3$  (d)  $sp^2d$
23. Shape of methane molecule is [MNR 1983]  
 (a) Tetrahedral (b) Pyramidal  
 (c) Octahedral (d) Square planar
24. Which one amongst the following possesses an  $sp$  hybridized carbon in its structure [CBSE PMT 1989]  
 (a)  $CH_2 = C.Cl - CH = CH_2$

- (b)  $C.Cl_2 = C.Cl_2$   
 (c)  $CH_2 = C = CH_2$   
 (d)  $CH_2 = CH - CH = CH_2$
25. Which of the following is the correct electronic formula of chlorine molecule [CPMT 1971]
- (a)  $:\ddot{Cl}:\ddot{Cl}:$  (b)  $:\ddot{Cl}^-::\ddot{Cl}^+:$   
 (c)  $:\ddot{Cl}:\ddot{Cl}:$  (d)  $:\ddot{Cl}::\ddot{Cl}:$
26. In  $XeF_4$  hybridization is
- (a)  $sp^3d^2$  (b)  $sp^3$   
 (c)  $sp^3d$  (d)  $sp^2d$
27. In  $HCHO$ , 'C' has hybridization [AIIMS 1987]
- (a)  $sp$  (b)  $sp^2$   
 (c)  $sp^3$  (d) All the above
28. Which has the shortest C - C bond length [NCERT 1982; CPMT 1989]
- (a)  $C_2H_5OH$  (b)  $C_2H_6$   
 (c)  $C_2H_2$  (d)  $C_2H_4$
29. The hybridization of Ag in the linear complex  $[Ag(NH_3)_2]^+$  is
- (a)  $dsp^2$  (b)  $sp$   
 (c)  $sp^2$  (d)  $sp^3$
30. Experiment shows that  $H_2O$  has a dipole moment while  $CO_2$  has not. Point out the structures which best illustrate these facts [DPMT 1984; NCERT 1983; CPMT 1984]
- (a)  $O = C = O$ ;  $H \begin{array}{c} \diagup O \\ \diagdown \end{array} H$  (b)  $O = C = O$ ;  $H - O - H$   
 (c)  $\begin{array}{c} O \\ // \\ C \\ // \\ O \end{array}$ ;  $H - H - O$  (d)  $\begin{array}{c} O \\ || \\ C = O \\ | \\ O - H \end{array}$  ;  $\begin{array}{c} H \\ | \\ O - H \end{array}$
31. Which species do not have  $sp^3$  hybridization [DPMT 1985]
- (a) Ammonia (b) Methane  
 (c) Water (d) Carbon dioxide
32. As compared to pure atomic orbitals, hybrid orbitals have
- (a) Low energy (b) Same energy  
 (c) High energy (d) None of these
33. The compound 1, 2-butadiene has [IIT 1983; MP PMT 1996]
- (a) Only  $sp$  hybridized carbon atoms  
 (b) Only  $sp^2$  hybridized carbon atoms  
 (c) Both  $sp$  and  $sp^2$  hybridized carbon atoms  
 (d)  $sp$ ,  $sp^2$  and  $sp^3$  hybridized carbon atoms
34. The number of unpaired electrons in  $O_2$  molecule is [MNR 1983; Kerala PET 2002]
- (a) 0 (b) 1  
 (c) 2 (d) 3
35. In the following molecule, the two carbon atoms marked by asterisk (\*) possess the following type of hybridized orbitals  $H_3C - C^* \equiv C^* - CH_3$  [NCERT 1984]
- (a)  $sp^3$  orbital (b)  $sp^2$  orbital  
 (c)  $sp$  orbital (d)  $s$  orbital
36. The bond angle in carbon tetrachloride is approximately [MNR 1981; MP PMT 1987]
- (a)  $90^\circ$  (b)  $109^\circ$   
 (c)  $120^\circ$  (d)  $180^\circ$
37. When two pairs of electrons are shared, bond is [MNR 1979]
- (a) Single covalent bond (b) Double covalent bond  
 (c) Dative bond (d) Triple bond
38. The nature of hybridization in the  $NH_3$  molecule is [EAMCET 1982]
- (a)  $sp$  (b)  $sp^2$   
 (c)  $sp^3$  (d)  $sp^3d$
39. Which one of the following compounds has bond angle as nearly  $90^\circ$  [MP PMT 1985]
- (a)  $NH_3$  (b)  $H_2S$   
 (c)  $H_2O$  (d)  $CH_4$
40. In ethene, the bond angle(s) is/are [CPMT 1985; BHU 1981] [CPMT 1976; AMU 1984; MP PMT 1985]
- (a)  $109^\circ 28'$  (b)  $120^\circ$   
 (c)  $180^\circ$  (d) Different
41. Structure formula of  $H_2O_2$  is [CPMT 1993]
- (a)  $\begin{array}{c} H \\ \diagup \\ O \rightarrow O \\ \diagdown \\ H \end{array}$   
 (b)  $H - O - O - H$  (straight line)  
 (c)  $\begin{array}{c} H' \\ | \\ O - O \\ | \\ H \end{array}$   
 (d)  $\begin{array}{c} H' \\ | \\ O - O \\ | \\ H \end{array}$
- Where  $\angle H - O - O = \angle O - O - H' = 101.5^\circ$  and all the four atoms are in the same plane
- (d)  $\begin{array}{c} H' \\ | \\ O - O \\ | \\ H \end{array}$
- Where  $\angle H - O - O = \angle O - O - H' = 97^\circ$  and the angle between  $H - O - O$  plane and  $O - O - H'$  plane is  $101^\circ$
42. Number of shared electrons in between carbon-carbon atoms in ethylene molecule is [MADT Bihar 1983]
- (a) 2 (b) 4  
 (c) 6 (d) 3
43. The structural formula of a compound is  $CH_3 - CH = C = CH_2$ . The type of hybridization at the four carbons from left to right are [CBSE PMT 1989]
- (a)  $sp^2$ ,  $sp$ ,  $sp^2$ ,  $sp^3$  (b)  $sp^2$ ,  $sp^3$ ,  $sp^2$ ,  $sp$

- (c)  $sp^3, sp^2, sp, sp^2$  (d)  $sp^3, sp^2, sp^2, sp^2$
44. Acetate ion contains [AMU 1983]  
 (a) One C, O single bond and one C, O double bond  
 (b) Two C, O single bonds  
 (c) Two C, O double bonds  
 (d) None of the above
45. The two carbon atoms in acetylene are [AMU 1984; MADT Bihar 1982]  
 (a)  $sp^3$  hybridized (b)  $sp^2$  hybridized  
 (c)  $sp$  hybridized (d) Unhybridized
46. Among the following compounds which is planar in shape [AMU 1992]  
 (a) Methane (b) Acetylene  
 (c) Benzene (d) Isobutene
47. In methane the bond angle is [AMU 1983]  
 (a)  $180^\circ$  (b)  $90^\circ$   
 (c)  $120^\circ$  (d)  $109^\circ$
48. The angle between  $sp^2$  orbitals in ethylene is [BHU 1987, 95; AMU 1985]  
 (a)  $90^\circ$  (b)  $120^\circ$   
 (c)  $180^\circ$  (d)  $109.5^\circ$
49. The species in which the central atom uses  $sp^2$  hybrid orbitals in its bonding is [IIT 1988]  
 (a)  $PH_3$  (b)  $NH_3$   
 (c)  $H_3C^+$  (d)  $SbH_3$
50. Carbon atoms in diamond are bonded to each other in a configuration [CPMT 1981]  
 (a) Tetrahedral (b) Planar  
 (c) Linear (d) Octahedral
51. Which of the following molecules can central atom said to adopt  $sp^2$  hybridization [CBSE PMT 1989; MP PET 1994]  
 (a)  $BeF_2$  (b)  $BCl_3$   
 (c)  $C_2H_2$  (d)  $NH_3$
52. In  $[Cu(NH_3)_4]SO_4$ , ; Cu has following hybridization [AIIMS 1988; UPSEAT 2001]  
 (a)  $dsp^2$  (b)  $sp^3$   
 (c)  $sp^2$  (d)  $sp^3d^2$
53. The hybridization of carbon atoms in C – C single bond of  $HC \equiv C - CH = CH_2$  is [IIT 1991; MP PET 1995]  
 (a)  $sp^3 - sp^3$  (b)  $sp^2 - sp^3$   
 (c)  $sp - sp^2$  (d)  $sp^3 - sp$
54. The compound in which  $C^*$  uses  $sp^3$  hybrids for bond formation is [IIT 1989]  
 (a)  $HCOOH^+$  (b)  $(NH_2)_2CO^+$   
 (c)  $(NH_3)_3COH^+ HgCl_2$  (d)  $CH_3CHO^+$
55. In diborane, the  $H - B - H$  bond angle is  $120^\circ$ . The hybridization of boron is likely to be [BHU 1981; CBSE PMT 1999]  
 (a)  $sp$  (b)  $sp^2$   
 (c)  $sp^3$  (d)  $dsp^2$
56. The number of shared pairs of electrons in propane is [BHU 1981]  
 (a) 2 (b) 4  
 (c) 6 (d) 10
57. s-character in  $sp$  hybridised orbitals are  
 (a)  $\frac{1}{3}$  (b)  $\frac{1}{2}$   
 (c)  $\frac{1}{4}$  (d)  $\frac{2}{3}$
58. The two types of bonds present in  $B_2H_6$  are covalent and [IIT 1994]  
 (a) Three centre bond (b) Hydrogen bond  
 (c) Two centre bond (d) None of the above
59. In the compound  $CH_3 \textcircled{C} OCl$ , which type of orbitals have been used by the circled carbon in bond formation [MP PET 1994]  
 (a)  $sp^3$  (b)  $sp^2$   
 (c)  $sp$  (d)  $p$
60. The correct order of the O – O bond length in  $O_2, H_2O_2$  and  $O_3$  is [CBSE PMT 1995]  
 (a)  $O_2 > O_3 > H_2O_2$  (b)  $O_3 > H_2O_2 > O_2$   
 (c)  $H_2O_2 > O_3 > O_2$  (d)  $O_2 > H_2O_2 > O_3$
61. The structure of  $PF_5$  molecule is [AFMC 1995; JIPMER 2001]  
 (a) Tetrahedral (b) Trigonal bipyramidal  
 (c) Square planar (d) Pentagonal bipyramidal
62. Which of the following hybridisation has maximum s-characters  
 (a)  $sp^3$  (b)  $sp^2$   
 (c)  $sp$  (d) None of these
63. The  $PCl_5$  molecule is a result of the hybridisation of [MP PET 1995; DCE 2000; MP PMT 2002]  
 (a)  $sp^2d^2$  (b)  $sp^3d$   
 (c)  $sp^3d^3$  (d)  $sp^2d^3$
64. Hybridisation involves [MP PMT 1996]  
 (a) Addition of an electron pair  
 (b) Mixing up of atomic orbitals  
 (c) Removal of an electron pair  
 (d) Separation of orbitals
65. The geometry of sulphur trioxide molecule is  
 (a) Tetrahedral (b) Trigonal planar  
 (c) Pyramidal (d) Square planar
66. The shapes of  $BCl_3, PCl_3$  and  $ICl_3$  molecules are all  
 (a) Triangular (b) Pyramidal  
 (c) T – shaped (d) All above are incorrect
67. In benzene molecule all C – C bond lengths are equal because



- (a) All carbon atoms are equivalent  
 (b) All carbon atoms are  $sp^2$  hybridised  
 (c) All  $C - C$  bonds in benzene, have same order  
 (d) All  $C - C$  bonds are single covalent bond
68. Which one is false in the following statements [MP PET 1997]  
 (a) Each carbon in ethylene is in  $sp^2$  hybridisation  
 (b) Each carbon in acetylene is in  $sp^3$  hybridisation  
 (c) Each carbon in benzene is in  $sp^2$  hybridisation  
 (d) Each carbon in ethane is in  $sp^3$  hybridisation
69. Out of the following hybrid orbitals, the one which forms the bond at angle  $120^\circ$ , is [MP PMT 1997]  
 (a)  $d^2sp^3$  (b)  $sp^3$   
 (c)  $sp^2$  (d)  $sp$
70. As the  $p$ -character increases, the bond angle in hybrid orbitals formed by  $s$  and atomic orbitals [MP PMT 1997]  
 (a) Decreases (b) Increases  
 (c) Doubles (d) Remains unchanged
71.  $sp^3$  hybridization leads to which shape of the molecule [MP PET/PMT 1998]  
 (a) Tetrahedron (b) Octahedron  
 (c) Linear (d) Plane triangle
72. Which of the following will be octahedral [MP PET 1999]  
 (a)  $SF_6$  (b)  $BF_4^-$   
 (c)  $PCl_5$  (d)  $BO_3^{3-}$
73. The hybrid orbitals used by central atoms in  $BeCl_2$ ,  $BCl_3$  and  $CCl_4$  molecules are respectively [MP PMT 1999]  
 (a)  $sp^2$ ,  $sp^3$  and  $sp$  (b)  $sp$ ,  $sp^2$  and  $sp^3$   
 (c)  $sp^3$ ,  $sp$  and  $sp^2$  (d)  $sp^2$ ,  $sp$  and  $sp^3$
74. The structure of  $H_2O_2$  is [CBSE PMT 1999; AFMC 2003]  
 (a) Planar (b) Non-planar  
 (c) Spherical (d) Linear
75. Which of the following is isoelectronic as well as has same structure as that of  $N_2O$  [CPMT 1999]  
 (a)  $N_3H$  (b)  $H_2O$   
 (c)  $NO_2$  (d)  $CO_2$
76.  $CCl_4$  has the hybridisation [DPMT 1996]  
 (a)  $sp^3d$  (b)  $dsp^2$   
 (c)  $sp$  (d)  $sp^3$
77. Compound having planar symmetry is [DPMT 1996]  
 (a)  $H_2SO_4$  (b)  $H_2O$   
 (c)  $HNO_3$  (d)  $CCl_4$
78. Which of the following compounds is not linear [CPMT 1996]  
 (a)  $SnCl_2$  (b)  $HCl$   
 (c)  $CO_2$  (d)  $HgCl_2$
79. Which one of the following statements is true for ammonium ion  
 (a) All bonds are ionic  
 (b) All bonds are coordinate covalent  
 (c)  $H$  atoms are situated at the corners of a square  
 (d)  $H$  atoms are situated at the corners of a tetrahedron
80. The bond angle in  $sp^2$  hybridisation is [RPMT 1997]  
 (a)  $180^\circ$  (b)  $120^\circ$   
 (c)  $90^\circ$  (d)  $109^\circ 2'$
81. The correct order towards bond angle is [RPMT 1997]  
 (a)  $sp < sp^2 < sp^3$   
 (b)  $sp^2 < sp < sp^3$   
 (c)  $sp^3 < sp^2 < sp$   
 (d) Bond angle does not depend on hybridisation
82. The geometry and the type of hybrid orbital present about the central atom in  $BF_3$  is [IIT 1998; BHU 2001]  
 (a) Linear,  $sp$  (b) Trigonal planar,  $sp^2$   
 (c) Tetrahedral,  $sp^3$  (d) Pyramidal,  $sp^3$
83. In graphite, electrons are [CBSE PMT 1997]  
 (a) Localised on every third  $C$  atom  
 (b) Present in antibonding orbital  
 (c) Localised on each  $C$  atom  
 (d) Spread out between the structure
84. The ammonium ion is [CET Pune 1998]  
 (a) Tetrahedral (b) Trigonal pyramidal  
 (c) Square planar (d) Square pyramidal
85. In  $sp$  hybridisation, shape is [Bihar MEE 1997]  
 (a) Angular (b) Tetrahedral  
 (c) Bipyramidal (d) Linear  
 (e) None of these
86. When the hybridisation state of carbon atom changes from  $sp^3$  to  $sp^2$  to  $sp$ , the angle between the hybridised orbitals [AIIMS 1998]  
 (a) Decreases gradually (b) Increases gradually  
 (c) Decreases considerably (d) All of these
87. The structure and hybridisation of  $Si(CH_3)_4$  is [CBSE PMT 1996]  
 (a) Bent,  $sp$  (b) Trigonal,  $sp^2$   
 (c) Octahedral,  $sp^3d$  (d) Tetrahedral,  $sp^3$
88. The type of hybridisation of boron in diborane is [BHU 1999]  
 (a)  $sp$ -hybridisation (b)  $sp^2$ -hybridisation  
 (c)  $sp^3$ -hybridisation (d)  $sp^3d^2$ -hybridisation
89. Which compound does not possess linear geometry

- (a)  $CH_2 = CH_2$  (b)  $HC \equiv CH$  [RPET 1999]
- (c)  $BeCl_2$  (d)  $CO_2$
90. Which of the following molecule does not show tetrahedral shape [RPET 1999]
- (a)  $CCl_4$  (b)  $SiCl_4$
- (c)  $SF_4$  (d)  $CF_4$
91. Pyramidal shape would be of [RPET 1999]
- (a)  $NO_3^-$  (b)  $H_2O$
- (c)  $H_3O^+$  (d)  $NH_4^+$
92. What is the correct mode of hybridization of the central atom in the following compounds :  $NO_2^+, SF_4, PF_6^-$  [AMU 1999]
- (a)  $sp^2, sp^3, d^2sp^3$  (b)  $sp^3, sp^3d^2, sp^3d^2$
- (c)  $sp, sp^3d, sp^3d^2$  (d)  $sp, sp^2, sp^3$
93. The hybridization in  $PF_3$  is [DCE 2000]
- (a)  $sp^3$  (b)  $sp^2$
- (c)  $dsp^3$  (d)  $d^2sp^3$
94. Which of the following molecule is linear [MP PMT 2000]
- (a)  $SO_2$  (b)  $NO_2^+$
- (c)  $NO_2^-$  (d)  $SCL_2$
95. The geometry of the molecule with  $sp^3d^2$  hybridised central atom is [NCERT 1981; AFMC 1982; RPMT 2000]
- (a) Square planar (b) Trigonal bipyramidal
- (c) Octahedral (d) Square pyramidal
96. The bond angle in  $PH_3$  is [RPMT 2000]
- (a) Much less than  $NH_3$
- (b) Equal to that of  $NH_3$
- (c) Much greater than  $NH_3$
- (d) Slightly greater than  $NH_3$
97. Which of the following has tetrahedral structure [CPMT 2000]
- (a)  $CO_3^-$  (b)  $NH_4^+$
- (c)  $K_4[Fe(CN)_6]$  (d) None of these
98. The single, double and triple bond lengths of carbon in carbon dioxide are respectively [AIIMS 2000]
- (a) 1.15, 1.22 and 1.10 Å (b) 1.22, 1.15 and 1.10 Å
- (c) 1.10, 1.15 and 1.22 Å (d) 1.15, 1.10 and 1.22 Å
99. Shape of  $BF_3$  molecule is [CPMT 2000; Pb. CET 2002]
- (a) Linear (b) Planar
- (c) Tetrahedral (d) Square pyramidal
100. In the complex  $[SbF_5]^{2-}$ ,  $sp^3d$  hybridization is present. Geometry of the complex is [Pb. PMT 2000]
- (a) Square (b) Square pyramidal
- (c) Square bipyramidal (d) Tetrahedral
101. The bond angle is minimum in [Pb. PMT 2001; MP PET 2003; UPSEAT 2004]
- (a)  $H_2Te$  (b)  $H_2Se$
- (c)  $H_2O$  (d)  $H_2S$
102. The correct order of hybridization of the central atom in the following species  $NH_3$ ,  $[PtCl_4]^{2-}$ ,  $PCl_5$  and  $BCl_3$  is [IIT Screening 2001; BHU 2005]
- (a)  $dsp^2, dsp^3, sp^2$  and  $sp^3$  (b)  $sp^3, dsp^2, dsp^3, sp^2$
- (c)  $dsp^2, sp^2, sp^3, dsp^3$  (d)  $dsp^2, sp^3, sp^2, dsp^3$
103. Which of the following pairs has same structure [BHU 2001]
- (a)  $PH_3$  and  $BCL_3$  (b)  $SO_2$  and  $NH_3$
- (c)  $PCl_5$  and  $SF_6$  (d)  $NH_4^+$  and  $SO_4^{2-}$
104. The smallest bond angle is found in [AIIMS 2001]
- (a)  $IF_7$  (b)  $CH_4$
- (c)  $BeF_2$  (d)  $BF_3$
105. Which of the following is not linear [DCE 2001]
- (a)  $CO_2$  (b)  $ClO_2$
- (c)  $I_3^-$  (d) None of these
106. Which of the following is not tetrahedral [MP PMT 2001]
- (a)  $SCL_4$  (b)  $SO_4^{2-}$
- (c)  $Ni(CO)_4$  (d)  $NiCl_4^{2-}$
107. As the  $s$ -character of hybridisation orbital increases, the bond angle [BHU 2002; AFMC 2002; MHCET 2003]
- (a) Increases (b) Decreases
- (c) Becomes zero (d) Does not change
108. The shape of  $IF_7$  molecule is [AFMC 2002; MHCET 2003]
- (a) Octahedral (b) Pentagonal bipyramidal
- (c) Trigonal bipyramidal (d) Tetrahedral
109. A completely filled  $d$  orbital ( $d^{10}$ ) [UPSEAT 2002]
- (a) Spherically symmetrical
- (b) Has octahedral symmetry
- (c) Has tetrahedral symmetry
- (d) Depends on the atom
110. Which has  $sp^3$  hybridization of central atom [UPSEAT 2002]
- (a)  $PCl_3$  (b)  $SO_3$
- (c)  $BF_3$  (d)  $NO_3^-$
111. In which of the following species is the interatomic bond angle is  $109^\circ 28'$  [AIEEE 2002]
- (a)  $NH_3, (BF_4)^-$  (b)  $(NH_4)^+, BF_3$
- (c)  $NH_3, BF_4$  (d)  $(NH_2)^-, BF_3$
112. A square planar complex is formed by hybridisation of which atomic orbitals [AIEEE 2002]
- (a)  $s, p_x, p_y, d_{yz}$  (b)  $s, p_x, p_y, d_{x^2-y^2}$
- (c)  $s, p_x, p_y, d_{z^2}$  (d)  $s, p_y, p_z, d_{xy}$
113. In benzene, all the six  $C-C$  bonds have the same length because of [MP PET 2002]
- (a) Tautomerism (b)  $sp^2$  hybridisation

- (c) Isomerism (d) Inductive effect
114. The bond energies of  $H-H$  and  $Cl-Cl$  are  $430 \text{ kJ mol}^{-1}$  and  $242 \text{ kJ mol}^{-1}$  respectively,  $\Delta H_f$  for  $HCl$  is  $91 \text{ kJ mol}$ . The bond energy of  $HCl$  will be [MP PET 2003]  
 (a)  $427 \text{ kJ}$  (b)  $766 \text{ kJ}$   
 (c)  $285 \text{ kJ}$  (d)  $245 \text{ kJ}$
115. Which of the following has  $dsp^2$  hybridization [MP PET 2003]  
 (a)  $NiCl_4^{2-}$  (b)  $SCl_4$   
 (c)  $NH_4^+$  (d)  $PtCl_4^{2-}$
116. Which one of the following is a planar molecule [EAMCET 2003]  
 (a)  $NH_3$  (b)  $H_3O^+$   
 (c)  $BCl_3$  (d)  $PCl_3$
117. Which one of the following is a correct set with respect to molecule, hybridisation and shape [EAMCET 2003]  
 (a)  $BeCl_2$ ,  $sp^2$ , linear  
 (b)  $BeCl_2$ ,  $sp^2$ , triangular planar  
 (c)  $BCl_3$ ,  $sp^2$ , triangular planar  
 (d)  $BCl_3$ ,  $sp^3$ , tetrahedral
118. Which of the following compounds doesn't have linear structure [RPET 1997, 2003]  
 (a)  $CO_2$  (b)  $SO_2$   
 (c)  $BeCl_2$  (d)  $C_2H_2$
119. Which of the following bonds require the largest amount of bond energy to dissociate the atom concerned [UPSEAT 2003]  
 (a)  $H-H$  bond in  $H_2$  (b)  $C-C$  bond in  $CH_4$   
 (c)  $N \equiv N$  bond in  $N_2$  (d)  $O=O$  bond in  $O_2$   
 (e)  $C-C$  bond in ethane
120. The percentage  $s$ -character of the hybrid orbitals in methane, ethene and ethyne are respectively [KCET 2003]  
 (a) 25, 33, 50 (b) 25, 50, 75  
 (c) 50, 75, 100 (d) 10, 20, 40
121. Arrange the hydra-acids of halogens in increasing order of acidity  
 (a)  $HF < HCl < HBr < HI$  (b)  $HI < HBr < HCl < HF$   
 (c)  $HF < HBr < HI < HCl$  (d)  $HF < HI < HBr < HCl$
122. Which one has  $sp^2$ -hybridisation [MP PMT 2004]  
 (a)  $CO_2$  (b)  $N_2O$   
 (c)  $SO_2$  (d)  $CO$
123. Among the following compounds the one that is polar and has central atom with  $sp^2$ -hybridization is [MP PMT 2004; IIT 1997]  
 (a)  $H_2CO_3$  (b)  $BF_3$   
 (c)  $SiF_4$  (d)  $HClO_2$
124. The molecule which is pyramid shape is [MP PMT 2004; EAMCET 1985; IIT 1989]  
 (a)  $PCl_3$  (b)  $CO_3^{2-}$
- (c)  $SO_3$  (d)  $NO_3^-$
125. Which of the following has a linear structure [MP PMT 2004]  
 (a)  $CCL_4$  (b)  $C_2H_2$   
 (c)  $SO_2$  (d)  $C_2H_4$
126. In a regular octahedral molecule,  $MX_6$ , the number  $X-M-X$  bonds at  $180^\circ$  is [CBSE PMT 2004]  
 (a) Six (b) Four  
 (c) Three (d) Two
127.  $sp^3d^2$  hybrid orbitals are [MP PET 2004]  
 (a) Linear bipyramidal (b) Pentagonal  
 (c) Trigonal bipyramidal (d) Octahedral
128. In an octahedral structure, the pair of  $d$  orbitals involved in  $d^2sp^3$  hybridization is [CBSE PMT 2004]  
 (a)  $d_{x^2}, d_{xz}$  (b)  $d_{xy}, d_{yz}$   
 (c)  $d_{x^2-y^2}, d_{z^2}$  (d)  $d_{xz}, d_{x^2-y^2}$
129. The correct order of bond angles (smallest first) in  $H_2S, NH_3, BF_3$  and  $SiH_4$  is [AIEEE 2004]  
 (a)  $H_2S < NH_3 < SiH_4 < BF_3$   
 (b)  $NH_3 < H_2S < SiH_4 < BF_3$   
 (c)  $H_2S < SiH_4 < NH_3 < BF_3$   
 (d)  $H_2S < NH_3 < BF_3 < SiH_4$
130. Which one of the following has the regular tetrahedral structure  
 (a)  $BF_4^-$  (b)  $SF_4$   
 (c)  $XeF_4$  (d)  $[Ni(CN)_4]^{2-}$   
 (Atomic no. :  $B = 5, S = 16, Ni = 28, Xe = 54$ )
131. The states of hybridization of boron and oxygen atoms in boric acid ( $H_3BO_3$ ) are respectively [AIEEE 2004]  
 (a)  $sp^3$  and  $sp^2$  (b)  $sp^2$  and  $sp^3$   
 (c)  $sp^2$  and  $sp^2$  (d)  $sp^3$  and  $sp^3$
132. The hybridisation in  $BF_3$  molecule is [Pb. PMT 2004]  
 (a)  $sp$  (b)  $sp^2$   
 (c)  $sp^3$  [Orissa JEE 2003] (d)  $sp^3d$
133. Among the compounds,  $BF_3, NCl_3, H_2S, SF_4$  and  $BeCl_2$ , identify the ones in which the central atom has the same type of hybridisation [Kerala PMT 2004]  
 (a)  $BF_3$  and  $NCl_3$  (b)  $H_2S$  and  $BeCl_2$   
 (c)  $BF_3, NCl_3$  and  $H_2S$  (d)  $SF_4$  and  $BeCl_2$   
 (e)  $NCl_3$  and  $H_2S$
134. The molecule of  $CO_2$  has  $180^\circ$  bond angle. It can be explained on the basis of [AFMC 2004]  
 (a)  $sp^3$  hybridisation (b)  $sp^2$  hybridisation  
 (c)  $sp$  hybridisation (d)  $d^2sp^3$  hybridisation
135.  $sp^3$  hybridisation is found in [Pb. CET 2003; Orissa JEE 2005]





- (a)  $CO_3^{2-}$  (b)  $BF_3$   
(c)  $NO_3^-$  (d)  $NH_3$
136. Which set hybridisation is correct for the following compounds [Pb. CET 2003]  
 $NO_2$ ,  $SF_4$ ,  $PF_6^-$   
(a)  $sp$ ,  $sp^2$ ,  $sp^3$   
(b)  $sp$ ,  $sp^3d$ ,  $sp^3d^2$   
(c)  $sp^2$ ,  $sp^3$ ,  $d^2sp^3$   
(d)  $sp^3$ ,  $sp^3d^2$ ,  $sp^3d^2$
137. The state of hybridisation of B in  $BCl_3$  is [Pb. CET 2000; BHU 2004]  
(a)  $sp$  (b)  $sp^2$   
(c)  $sp^3$  (d)  $sp^2d^2$
138. The hybrid state of sulphur in  $SO_3$  molecule is [DCE 2004]  
(a)  $sp^3d$  (b)  $sp^3$   
(c)  $sp^3d^2$  (d)  $sp^2$
139. Which of the following molecules has pyramidal shape [DCE 2004; J&K CET 2005]  
(a)  $PCl_3$  (b)  $SO_3$   
(c)  $CO_3^{2-}$  (d)  $NO_3^-$
140. The hybridization of  $IF_7$  is [Pb. CET 2001]  
(a)  $sp^3d^3$  (b)  $sp^2d$   
(c)  $d^2sp^3$  (d)  $sp^3$
141. In which compound, the hydrogen bonding is the strongest in its liquid phase [Pb. CET 2001]  
(a)  $HF$  (b)  $HI$   
(c)  $CH_4$  (d)  $PH_3$
142. Geometry of ammonia molecule and the hybridization of nitrogen involved in it are [MH CET 2004]  
(a)  $sp^3$ -hybridization and tetrahedral geometry  
(b)  $sp^3$ -hybridization and distorted tetrahedral geometry  
(c)  $sp^2$ -hybridization and triangular geometry  
(d) None of these
143. Be in  $BeCl_2$  undergoes [MH CET 2004]  
(a) Diagonal hybridization  
(b) Trigonal hybridization  
(c) Tetrahedral hybridization  
(d) No hybridization
144. Which of the following is non-linear molecule [DCE 2003]  
(a)  $CO_3$  (b)  $CO_2$   
(c)  $CS_2$  (d)  $BeCl_2$
145. The trigonal bipyramidal geometry results from the hybridisation [UPSEAT 2004]  
(a)  $dsp^3$  or  $sp^3d$  (b)  $dsp^2$  or  $sp^2d$   
(c)  $d^2sp^3$  or  $sp^3d^2$  (d)  $d^3sp^2$  or  $d^2sp^3$
146. The valency of carbon is four. On what principle it can be explained in a better way  
(a) Resonance (b) Hybridization  
(c) Electron transfer (d) None of the above
147. Hybridization is due to the overlapping of [MADT Bihar 1983]  
(a) Orbitals of different energy levels  
(b) Orbitals of different energy content  
(c) Orbitals of same energy content  
(d) None of the above
148. If a molecule  $MX_3$  has zero dipole moment, the sigma bonding orbital used by M are [IIT 1981; MP PMT 1994; Kerala PMT 2004]  
(a)  $sp^3d$  - hybrid (b)  $sp$  - hybrid  
(c)  $sp^3d^2$  - hybrid (d)  $sp^2$  - hybrid
149. The linear structure is assumed by [IIT 1991]  
(a)  $SnCl_2$  (b)  $NCO^-$   
(c)  $CS_2$  (d)  $NO_2^+$
150. Hybridisation of central atom in  $NF_3$  is [Orissa JEE 2005]  
(a)  $sp^3$  (b)  $sp$   
(c)  $sp^2$  (d)  $dsp^2$
151. The pair having similar geometry is [J&K CET 2005]  
(a)  $PCl_3$ ,  $NH_3$  (b)  $BeCl_2$ ,  $H_2O$   
(c)  $CH_4$ ,  $CCl_4$  (d)  $IF_5$ ,  $PF_5$
152. The d-orbital involved in  $sp^3d$  hybridisation is [J&K CET 2005]  
(a)  $d_{x^2-y^2}$  (b)  $d_{xy}$   
(c)  $d_{z^2}$  (d)  $d_{zx}$

## Resonance

- Which one in the following is not the resonance structure of  $CO_2$   
(a)  $O=C=O$  (b)  $^-O-C\equiv O^+$   
(c)  $^+O\equiv C-O^-$  (d)  $O\equiv C=O$
- Which of the following molecule contains one pair of non-bonding electrons  
(a)  $CH_4$  (b)  $NH_3$   
(c)  $H_2O$  (d)  $HF$
- Resonance is due to [NCERT 1981; Kurukshetra CEE 1998]  
(a) Delocalization of  $\sigma$  electrons  
(b) Delocalization of  $\pi$  electrons  
(c) Migration of H atoms  
(d) Migration of protons
- Resonating structures have different [AMU 1983]  
(a) Atomic arrangements (b) Electronic arrangements  
(c) Functional groups (d) Alkyl groups

In the cyanide ion, the formal negative charge is on

[AMU 1984]

- (a) C  
(b) N  
(c) Both C and N  
(d) Resonate between C and N
6. Which does not show resonance [CPMT 1990]  
(a) Benzene (b) Aniline  
(c) Ethyl amine (d) Toluene
7. The enolic form of acetone contains [IIT 1990; Bihar MEE 1997]  
(a) 9 sigma bonds, 1 pi bond and 2 lone pairs  
(b) 8 sigma bonds, 2 pi bonds and 2 lone pairs  
(c) 10 sigma bonds, 1 pi bond and 1 lone pair  
(d) 9 sigma bonds, 2 pi bonds and 1 lone pair
8. Point out incorrect statement about resonance [MP PET 1997]  
(a) Resonance structures should have equal energy  
(b) In resonance structures, the constituent atoms should be in the same position  
(c) In resonance structures, there should not be the same number of electron pairs  
(d) Resonance structures should differ only in the location of electrons around the constituent atoms
9. The number of possible resonance structures for  $CO_3^{2-}$  is [MP PMT 2000]  
(a) 2 (b) 3  
(c) 6 (d) 9
10. Resonance hybrid of nitrate ion is [RPET 2000]  
(a)  (b)   
(c)  (d) 
11.  $CO_3^{2-}$  anion has which of the following characteristics [Roorkee 1999]  
(a) Bonds of unequal length  
(b)  $sp^2$  hybridization of C atom  
(c) Resonance stabilization  
(d) Same bond angles
- (c)  $PF_3$  (d)  $NH_3$
4. Which has the least bond angle [NCERT 1973; DPMT 1990; CBSE PMT 1990; UPSEAT 2003]  
(a)  $NH_3$  (b)  $BeF_2$   
(c)  $H_2O$  (d)  $CH_4$
5. In compound X, all the bond angles are exactly  $109^\circ 28'$ , X is  
(a) Chloromethane (b) Iodoform  
(c) Carbon tetrachloride (d) Chloroform
6. The shape of  $SO_4^{2-}$  ion is [CPMT 1982; DPMT 1983, 84, 96; Bihar MEE 1997]  
(a) Square planar (b) Tetrahedral  
(c) Trigonal bipyramidal (d) Hexagonal
7. Which of the following molecules has one lone pair of electrons on the central atom [EAMCET 1980; AMU 1982; MNR 1989]  
(a)  $H_2O$  (b)  $NH_3$   
(c)  $CH_4$  (d)  $PCl_5$
8. Of the following compounds, the one having a linear structure is [NCERT 1981; MP PMT 1985; AIIMS 1996]  
(a)  $NH_2$  (b)  $CH_4$   
(c)  $C_2H_2$  (d)  $H_2O$
9.  $XeF_6$  is  
(a) Octahedral (b) Distorted octahedral  
(c) Planar (d) Tetrahedral
10. Which has maximum bond angle [CPMT 1993]  
(a)  $CHF_3$   
(b)  $CHCl_3$   
(c)  $CHBr_3$   
(d) All have maximum bond angle
11. Of the following species the one having a square planar structure is [NCERT 1981; MP PMT 1994]  
(a)  $NH_4^+$  (b)  $BF_4^-$   
(c)  $XeF_4$  (d)  $SrCl_4$
12. In which of the following is the angle between the two covalent bonds greatest [NCERT 1975; AMU 1982; MNR 1987; IIT 1981; CPMT 1988; MP PMT 1994]  
(a)  $CO_2$  (b)  $CH_4$   
(c)  $NH_3$  (d)  $H_2O$
13. As the s-character of hybridized orbital decreases, the bond angle  
(a) Decreases (b) Increases  
(c) Does not change (d) Becomes zero
14.  $XeF_2$  molecule is [BHU 1982]  
(a) Linear (b) Triangular planar  
(c) Pyramidal (d) Square planar
15. Of the following sets which one does NOT contain isoelectronic species [AIIEE 2005]  
(a)  $PO_4^{3-}, SO_4^{2-}, ClO_4^-$  (b)  $CN^-, N_2, C_2^{2-}$   
(c)  $SO_3^{2-}, CO_3^{2-}, NO_3^-$  (d)  $BO_3^{3-}, CO_3^{2-}, NO_3^-$
16. A molecule which contains unpaired electrons is

## VSEPR Theory

1. The structure of  $[Cu(H_2O)_4]^{++}$  ion is [NCERT 1983; MP PMT 1983]  
(a) Square planar (b) Tetrahedral  
(c) Distorted rectangle (d) Octahedral
2. The bond angle in  $PH_3$  would be expected to be close to  
(a)  $90^\circ$  (b)  $105^\circ$   
(c)  $109^\circ$  (d)  $120^\circ$
3. In which molecule are all atoms coplanar [MP PMT 1994]  
(a)  $CH_4$  (b)  $BF_3$

- [NCERT 1982]
- (a) Carbon monoxide                      (b) Molecular nitrogen  
(c) Molecular oxygen                      (d) Hydrogen peroxide
17.  $H_2O$  is [MADT Bihar 1983]
- (a) A linear triatomic molecule  
(b) A bent (angular) triatomic molecule  
(c) Both of these  
(d) None of these
18. Bond angle between two hybrid orbitals is  $105^\circ$ . % s-orbital character of hybrid orbital is [MP PMT 1986]
- (a) Between 20 – 21%                      (b) Between 19 – 20%  
(c) Between 21 – 22%                      (d) Between 22 – 23%
19. The bond angle between  $H - O - H$  in ice is closest to [CPMT 1989; UPSEAT 2002]
- (a)  $120^\circ 28'$                                   (b)  $60^\circ$   
(c)  $90^\circ$     (d)  $105^\circ$
20. Which of the following molecules does not have a linear arrangement of atoms [CBSE PMT 1989]
- (a)  $H_2S$     (b)  $C_2H_2$   
(c)  $BeH_2$     (d)  $CO_2$
21.  $BCl_3$  is a planar molecule while  $NCl_3$  is pyramidal, because
- (a)  $BCl_3$  has no lone pair of electrons but  $NCl_3$  has a lone pair of electrons  
(b)  $B - Cl$  bond is more polar than  $N - Cl$  bond  
(c) Nitrogen atom is smaller than boron atom  
(d)  $N - Cl$  bond is more covalent than  $B - Cl$  bond
22. The isoelectronic pair is [AIIMS 2005]
- (a)  $Cl_2O, ICl_2^-$                                   (b)  $ICl_2^-, ClO_2^-$   
(c)  $IF_2^+, I_3^-$                                       (d)  $ClO_2^-, ClF_2^+$
23. According to VSEPR theory, the most probable shape of the molecule having 4 electron pairs in the outer shell of the central atom is [MP PET 1996, 2001]
- (a) Linear    (b) Tetrahedral  
(c) Hexahedral                                      (d) Octahedral
24. The molecular shapes of  $SF_4, CF_4$  and  $XeF_4$  are [AIEEE 2005]
- (a) The same with 2, 0 and 1 lone pairs of electrons on the central atom, respectively  
(b) The same with 1, 1 and 1 lone pair of electrons on the central atoms, respectively  
(c) Different with 0, 1 and 2 lone pairs of electrons on the central atom, respectively  
(d) Different with 1, 0 and 2 lone pairs of electrons on the central atom, respectively
25. Which of the following species is planar [JIPMER 1997]
- (a)  $CO_3^{2-}$     (b)  $NH_2$   
(c)  $PCl_3$     (d) None of these
26. The shape of  $CH_3^+$  species is [RPET 1999]
- (a) Tetrahedral                                      (b) Square planar  
(c) Trigonal planar                                  (d) Linear
27. Which of the following is the correct reducing order of bond-angle
- (a)  $NH_3 < CH_4 < C_2H_2 < H_2O$                       (b)  $C_2H_2 > NH_3 > H_2O < CH_4$   
(c)  $NH_3 > H_2O > CH_4 < C_2H_2$                       (d)  $H_2O < NH_3 > CH_4 < C_2H_2$
28. Which compound has bond angle nearly to  $90^\circ$  [Pb. PMT 2001]
- (a)  $H_2O$     (b)  $H_2S$   
(c)  $NH_3$     (d)  $CH_4$
29. A lone pair of electrons in an atom implies [KCET 2002]
- (a) A pair of valence electrons not involved in bonding  
(b) A pair of electrons involved in bonding  
(c) A pair of electrons  
(d) A pair of valence electrons
30. The bond angle of water is  $104.5^\circ$  due to [CPMT 2002]
- (a) Repulsion between lone pair and bond pair  
(b)  $sp^3$  hybridization of O  
(c) Bonding of  $H_2O$   
(d) Higher electronegativity of O
31. The correct sequence of decrease in the bond angle of the following hydrides is [MP PET 2002]
- (a)  $NH_3 > PH_3 > AsH_3 > SbH_3$   
(b)  $NH_3 > AsH_3 > PH_3 > SbH_3$  [CBSE PMT 1995]  
(c)  $SbH_3 > AsH_3 > PH_3 > NH_3$   
(d)  $PH_3 > NH_3 > AsH_3 > SbH_3$
32. Central atom of the following compound has one lone pair of electrons and three bond pairs of electrons [JIPMER 2002]
- (a)  $H_2S$     (b)  $AlCl_3$   
(c)  $NH_3$     (d)  $BF_3$
33. Among  $KO_2, AlO_2^-, BaO_2$  and  $NO_2^+$  unpaired electron is present in [MP PET 2003]
- (a)  $NO_2^+$  and  $BaO_2$                                   (b)  $KO_2$  and  $AlO_2^-$   
(c)  $KO_2$  only    (d)  $BaO_2$  only
34. True order of bond angle is [RPET 2003]
- (a)  $H_2O > H_2S > H_2Se > H_2Te$   
(b)  $H_2Te > H_2Se > H_2S > H_2O$   
(c)  $H_2S > H_2O > H_2Se > H_2Te$   
(d)  $H_2O > H_2S > H_2Te > H_2Se$
35. Which of the following has not a lone pair over the central atom
- (a)  $NH_3$     (b)  $PH_3$   
(c)  $BF_3$     (d)  $PCl_3$
36. In  $BrF_3$  molecule, the lone pairs occupy equatorial positions to minimize [CBSE PMT 2004]
- (a) Lone pair- lone pair repulsion and lone pair-bond pair repulsion  
(b) Lone pair- lone pair repulsion only  
(c) Lone pair- bond pair repulsion only  
(d) Bond pair- bond pair repulsion only
37.  $H_2O$  is dipolar, whereas  $BeF_2$  is not. It is because [BHU 2000] [CBSE PMT 1989; 2004]
- (a)  $H_2O$  is linear and  $BeF_2$  is angular



- (b)  $H_2O$  is angular and  $BeF_2$  is linear  
(c) The electronegativity of  $F$  is greater than that of  $O$   
(d)  $H_2O$  involves hydrogen bonding whereas  $BeF_2$  is a discrete molecule
38. Maximum bond angle is present in [BVP 2004]  
(a)  $BCl_3$  (b)  $BBr_3$   
(c)  $BF_3$  (d) Same for all
39. The shape of a molecule of  $NH_3$ , in which central atoms contains lone pair of electron, is [MHCET 2003]  
(a) Tetrahedral (b) Planar trigonal  
(c) Square planar (d) Pyramidal
40. The largest bond angle is in [DCE 2002; MNR 1984]  
(a)  $AsH_3$  (b)  $NH_3$   
(c)  $H_2O$  (d)  $PH_3$
41. The bond angle in ammonia molecule is [EAMCET 1980]  
(a)  $91^\circ 8'$  (b)  $93^\circ 3'$   
(c)  $106^\circ 45'$  (d)  $109^\circ 28'$
42. Which of the following gives correct arrangement of compounds involved based on their bond strength [BHU 2005]  
(a)  $HF > HCl > HBr > HI$   
(b)  $HI > HBr > HCl > HF$   
(c)  $HF > HBr > HCl > HI$   
(d)  $HCl > HF > HBr > HI$
43. Which one has a pyramidal structure [CBSE PMT 1990]  
(a)  $CH_4$  (b)  $NH_3$   
(c)  $H_2O$  (d)  $CO_2$
44. Among the following the pair in which the two species are not isostructural is [CBSE PMT 2004]  
(a)  $BH_4^-$  and  $NH_4^+$  (b)  $PF_6^-$  and  $SF_6$   
(c)  $SiF_4$  and  $SF_4$  (d)  $IO_3^-$  and  $XeO_3$
45. The maximum number of  $90^\circ$  angles between bond pair-bond pair of electrons is observed in [AIEEE 2004]  
(a)  $dsp^2$  hybridization (b)  $sp^3d$  hybridization  
(c)  $dsp^3$  hybridization (d)  $sp^3d^2$  hybridization
- (a)  $O_2$  (b)  $O_2^{-2}$   
(c)  $O_2^{+1}$  (d)  $O_2^{-1}$
5. The bond order is maximum in [AIIMS 1983, 85; CBSE PMT 1994; MP PET 2002]  
(a)  $O_2$  (b)  $O_2^{-1}$   
(c)  $O_2^{+1}$  (d)  $O_2^{-2}$
6. Which of the following compounds of boron does not exist in the free form  
(a)  $BCl_3$  (b)  $BF_3$   
(c)  $BBr_3$  (d)  $BH_3$
7. Molecular orbital theory was developed mainly by [BHU 1987; Pb. CET 2003]  
(a) Pauling (b) Pauling and Slater  
(c) Mulliken (d) Thomson
8. The bond order of a molecule is given by [NCERT 1984]  
(a) The difference between the number of electrons in bonding and antibonding orbitals  
(b) Total number of electrons in bonding and antibonding orbitals  
(c) Twice the difference between the number of electrons in bonding and antibonding electrons  
(d) Half the difference between the number of electrons in bonding and antibonding electrons
9. Oxygen molecule is paramagnetic because [NCERT 1984; IIT 1984]  
(a) Bonding electrons are more than antibonding electrons  
(b) Contains unpaired electrons  
(c) Bonding electrons are less than antibonding electrons  
(d) Bonding electrons are equal to antibonding electrons
10. Which one is paramagnetic from the following [IIT 1989; CBSE PMT 1995]  
(a)  $O_2^-$  (b)  $NO$   
(c) Both (a) and (b) (d)  $CN^-$
11. The bond order in  $N_2^+$  ion is [Pb. CET 2004]  
(a) 1 (b) 2  
(c) 2.5 (d) 3
12. Out of the following which has smallest bond length [RPMT 1997]  
(a)  $O_2$  (b)  $O_2^+$   
(c)  $O_2^-$  (d)  $O_2^{2-}$

## Molecular orbital theory

1. Bond order is a concept in the molecular orbital theory. It depends on the number of electrons in the bonding and antibonding orbitals. Which of the following statements is true about it? The bond order [AIIMS 1980]  
(a) Can have a negative quantity  
(b) Has always an integral value  
(c) Can assume any positive or integral or fractional value including zero  
(d) Is a non zero quantity
2. The bond order of  $NO$  molecule is [MP PET 1996]  
(a) 1 (b) 2  
(c) 2.5 (d) 3
3. When two atomic orbitals combine they form  
(a) One molecular orbital (b) Two molecular orbital  
(c) Three molecular orbital (d) Four molecular orbital
4. Which of the following species is the least stable
13. Which of the following molecule is paramagnetic [CPMT 1980; RPET 1999; MP PMT 1999; RPMT 2000]  
(a) Chlorine (b) Nitrogen  
(c) Oxygen (d) Hydrogen
14. Which molecule has the highest bond order  
(a)  $N_2$  (b)  $Li_2$   
(c)  $He_2$  (d)  $O_2$
15. The molecular electronic configuration of  $H_2^-$  ion is  
(a)  $(\sigma 1s)^2$  (b)  $(\sigma 1s)^2(\sigma^* 1s)^2$   
(c)  $(\sigma 1s)^2(\sigma^* 1s)^1$  (d)  $(\sigma 1s)^3$
16. The paramagnetic nature of oxygen molecule is best explained on the basis of [BHU 1996]  
(a) Valence bond theory (b) Resonance

- (c) Molecular orbital theory (d) Hybridization
17. In which case the bond length is minimum between carbon and nitrogen  
(a)  $CH_3NH_2$  (b)  $C_6H_5CH=NOH$   
(c)  $CH_3CONH_2$  (d)  $CH_3CN$
18. Which one of the following species is diamagnetic in nature [AIEEE 2005]  
(a)  $He_2^+$  (b)  $H$   
(c)  $H_2^+$  (d)  $H_2^-$
19. Which one of the following oxides is expected exhibit paramagnetic behaviour [CBSE PMT 2005]  
(a)  $CO_2$  (b)  $SO_2$   
(c)  $ClO_2$  (d)  $SiO_2$
20. The bond order in  $N_2$  molecule is [CBSE 1995; Pb. PMT 1999; MP PET 1997]  
(a) 1 (b) 2  
(c) 3 (d) 4
21. Which one is paramagnetic and has the bond order 1/2 [NCERT 1983]  
(a)  $O_2$  (b)  $N_2$   
(c)  $F_2$  (d)  $H_2^+$
22. When two atoms of chlorine combine to form one molecule of chlorine gas, the energy of the molecule [AMU 1982]  
(a) Greater than that of separate atoms  
(b) Equal to that of separate atoms  
(c) Lower than that of separate atoms  
(d) None of the above statement is correct
23. An atom of an element  $A$  has three electrons in its outermost shell and that of  $B$  has six electrons in the outermost shell. The formula of the compound between these two will be [CPMT 1974, 84; RPMT 1999]  
(a)  $A_3B_4$  (b)  $A_2B_3$   
(c)  $A_3B_2$  (d)  $A_2B$
24. The bond order of individual carbon-carbon bonds in benzene is  
(a) One (b) Two  
(c) Between 1 and 2 (d) One and two alternately
25.  $PCl_5$  exists but  $NCl_5$  does not because [EAMCET 1977; MP PET/PMT 1988]  
(a) Nitrogen has no vacant d-orbitals  
(b)  $NCl_5$  is unstable  
(c) Nitrogen atom is much smaller  
(d) Nitrogen is highly inert
26. Paramagnetism is exhibited by molecules [NCERT 1979; MP PET 2002]  
(a) Not attracted into a magnetic field  
(b) Containing only paired electrons  
(c) Carrying a positive charge  
(d) Containing unpaired electrons
27. Which one of the following is paramagnetic [DPMT 1985]  
(a)  $H_2O$  (b)  $NO_2$   
(c)  $SO_2$  (d)  $CO_2$
28. The energy of a  $2p$  orbital except hydrogen atom is [AMU 1983]  
(a) Less than that of  $2s$  orbital  
(b) More than that of  $2s$  orbital  
(c) Equal to that of  $2s$  orbital  
(d) Double that of  $2s$  orbital
29. In the electronic structure of acetic acid, there are [AMU 1983]  
(a) 16 shared and 8 unshared electrons  
(b) 8 shared and 16 unshared electrons  
(c) 12 shared and 12 unshared electrons  
(d) 18 shared and 6 unshared electrons
30. Which of the following does not exist on the basis of molecular orbital theory [AFMC 1990; MP PMT 1996]  
(a)  $H_2^+$  (b)  $He_2^+$   
(c)  $He_2$  (d)  $Li_2$
31. In  $P_4O_{10}$ , the number of oxygen atoms attached to each phosphorus atom is [IIT 1995]  
(a) 2 (b) 3  
(c) 4 (d) 2.5
32. Of the following statements which one is correct  
(a) Oxygen and nitric oxide molecules are both paramagnetic because both contain unpaired electrons  
(b) Oxygen and nitric oxide molecules are both diamagnetic because both contain no unpaired electrons  
(c) Oxygen is paramagnetic because it contains unpaired electrons, while nitric oxide is diamagnetic because it contains no unpaired electrons  
(d) Oxygen is diamagnetic because it contains no unpaired electrons, while nitric oxide is paramagnetic because it contains an unpaired electron
33. According to the molecular orbital theory, the bond order in  $C_2$  molecule is  
(a) 0 (b) 1  
(c) 2 (d) 3
34. The molecular orbital configuration of a diatomic molecule is [IIT 1980]  
 $\sigma 1s^2 \sigma^* 1s^2 \sigma 2s^2 \sigma^* 2s^2 \sigma 2p_x^2 \begin{cases} \pi 2p_y^2 \\ \pi 2p_z^2 \end{cases}$
- Its bond order is  
(a) 3 (b) 2.5  
(c) 2 (d) 1
35. The difference in energy between the molecular orbital formed and the combining atomic orbitals is called  
(a) Bond energy (b) Activation energy  
(c) Stabilization energy (d) Destabilization energy
36. According to molecular orbital theory, the paramagnetism of  $O_2$  molecule is due to presence of [MP PMT 1997]  
(a) Unpaired electrons in the bonding  $\sigma$  molecular orbital  
(b) Unpaired electrons in the antibonding  $\sigma$  molecular orbital  
(c) Unpaired electron in the bonding  $\pi$  molecular orbital  
(d) Unpaired electrons in the antibonding  $\pi$  molecular orbital
37. The bond order in  $O_2^+$  is [MP PET 1999; BHU 2001]  
(a) 2 (b) 2.5  
(c) 1.5 (d) 3

38. Which of the following is paramagnetic [MP PET 1999]  
 (a)  $O_2$  (b)  $CN^-$   
 (c)  $CO$  (d)  $NO^+$
39. If  $N_x$  is the number of bonding orbitals of an atom and  $N_y$  is the number of antibonding orbitals, then the molecule/atom will be stable if [DPMT 1996]  
 (a)  $N_x > N_y$  (b)  $N_x = N_y$   
 (c)  $N_x < N_y$  (d)  $N_x \leq N_y$
40. Which of the following molecular orbitals has two nodal planes  
 (a)  $\sigma 2s$  (b)  $\pi 2p_y$   
 (c)  $\pi^* 2p_y$  (d)  $\sigma^* 2p_x$
41. The number of nodal planes 'd' orbital has [KCET 1996]  
 (a) Zero (b) One  
 (c) Two (d) Three
42. Atomic number of an element is 26. The element shows [CPMT 1996]  
 (a) Ferromagnetism (b) Diamagnetism  
 (c) Paramagnetism (d) None of these
43. What is correct sequence of bond order [BHU 1997]  
 (a)  $O_2^+ > O_2^- > O_2$  (b)  $O_2^+ > O_2 > O_2^-$   
 (c)  $O_2 > O_2^- > O_2^+$  (d)  $O_2^- > O_2^+ > O_2$
44. Which bond is strongest [RPMT 1997]  
 (a)  $F-F$  (b)  $Br-F$   
 (c)  $Cl-F$  (d)  $I-F$
45. Which of the following is not paramagnetic [AIIMS 1997]  
 (a)  $S^{-2}$  (b)  $N_2^-$   
 (c)  $O_2^-$  (d)  $NO$
46. Which one of the following molecules is paramagnetic [Pb. PMT 1998]  
 (a)  $CO_2$  (b)  $SO_2$   
 (c)  $NO$  (d)  $H_2O$
47.  $N_2$  and  $O_2$  are converted into monoanions  $N_2^-$  and  $O_2^-$  respectively, which of the following statements is wrong [CBSE PMT 1997]  
 (a) In  $N_2$ , the  $N-N$  bond weakens  
 (b) In  $O_2$ , the  $O-O$  bond order increases  
 (c) In  $O_2$ , bond length increases  
 (d)  $N_2^-$  becomes diamagnetic
48. With increasing bond order, stability of a bond [CET Pune 1998]  
 (a) Remains unaltered (b) Decreases  
 (c) Increases (d) None of these
49. Which is not paramagnetic [DCE 1999, 2000]  
 (a)  $O_2$  (b)  $O_2^+$   
 (c)  $O_2^{2-}$  (d)  $O_2^-$
50. The number of antibonding electron pairs in  $O_2^{2-}$  molecular ion on the basis of molecular orbital theory [Pb. PMT 2000]  
 (a) 4 (b) 3  
 (c) 2 (d) 5
51. The bond order of  $He_2^+$  molecule ion is [Pb. PMT 2000; Pb CET 2001]  
 (a) 1 (b) 2  
 (c)  $\frac{1}{2}$  (d)  $\frac{1}{4}$
52. Which of the following exhibit paramagnetism [DPMT 2000]  
 (a)  $ClO_2$  (b)  $ClO_2^-$   
 (c)  $NO_2$  (d)  $NO$
53. In which of the following pairs the two molecules have identical bond order [MP PMT 2000]  
 (a)  $N_2, O_2^{2+}$  (b)  $N_2, O_2^-$   
 (c)  $N_2^-, O_2$  (d)  $O_2^+, N_2$
54. The bond order is not three for [MP PMT 2001]  
 (a)  $N_2^+$  (b)  $O_2^{2+}$   
 (c)  $N_2$  (d)  $NO^+$
55. In  $H_2O_2$  molecule, the angle between the two  $O-H$  planes is  
 (a)  $90^\circ$  (b)  $101^\circ$   
 (c)  $103^\circ$  (d)  $105^\circ$
56. Which of the following molecule has highest bond energy [AIIMS 2002]  
 (a)  $F-F$  (b)  $C-C$   
 (c)  $N-N$  (d)  $O-O$
57. Which of the following species would be expected paramagnetic  
 (a) Copper crystals (b)  $Cu^+$   
 (c)  $Cu^{++}$  (d)  $H_2$
58. Which of the following is correct for  $N_2$  triple bond [CPMT 2002]  
 (a)  $3s$  (b)  $1p, 2s$   
 (c)  $2p, 1s$  (d)  $3p$
59. In which of the following pairs molecules have bond order three and are isoelectronics [MP PET 2003]  
 (a)  $CN^-, CO$  (b)  $NO^+, CO^+$   
 (c)  $CN^-, O_2^+$  (d)  $CO, O_2^+$
60. Which of the following is paramagnetic [MP PET 2003]  
 (a)  $O_2^+$  (b)  $CN^-$   
 (c)  $CO$  (d)  $N_2$
61. How many bonding electron pairs are there in white phosphorous  
 (a) 6 (b) 12  
 (c) 4 (d) 8
62. The atomicity of phosphorus is  $X$  and the  $\hat{P}PP$  bond angle in the molecule is  $Y$ . What are  $X$  and  $Y$  [EAMCET 2003]  
 (a)  $X = 4, Y = 90^\circ$  (b)  $X = 4, Y = 60^\circ$   
 (c)  $X = 3, Y = 120^\circ$  (d)  $X = 2, Y = 180^\circ$

63. From elementary molecular orbital theory we can give the electronic configuration of the singly positive nitrogen molecular ion  $N_2^+$  as
- $\sigma(1s)^2 \sigma^*(1s)^2 \sigma(2s)^2 \sigma^*(2s)^2 \pi(2p)^4 \sigma(2p)^1$
  - $\sigma(1s)^2 \sigma^*(1s)^2 \sigma(2s)^2 \sigma^*(2s)^2 \sigma(2p)^1 \pi(2p)^3$
  - $\sigma(1s)^2 \sigma^*(1s)^2 \sigma(2s)^2 \sigma^*(2p)^2 \pi(2p)^4$
  - $\sigma(1s)^2 \sigma^*(1s)^2 \sigma(2s)^2 \sigma^*(2s)^2 \sigma(2p)^2 \pi(2p)^2$
64. The paramagnetic property of the oxygen molecule due to the presence of unpaired electrons present in [Kerala PMT 2004]
- $(\sigma 2p_x)^1$  and  $(\sigma^* 2p_x)^1$
  - $(\sigma 2p_x)^1$  and  $(\pi 2p_y)^1$
  - $(\pi^* 2p_y)^1$  and  $(\pi^* 2p_z)^1$
  - $(\pi^* 2p_y)^1$  and  $(\pi 2p_y)^1$
  - $(\pi^* 2p_z)^1$  and  $(\pi 2p_z)^1$
65. In  $PO_4^{3-}$  ion, the formal charge on each oxygen atom and  $P-O$  bond order respectively are [DPMT 2004]
- 0.75, 1.25
  - 0.75, 1.0
  - 0.75, 0.6
  - 3, 1.25
66. The bond order in  $CO_3^{2-}$  ion between  $C-O$  is [Pb. PMT 2004]
- Zero
  - 0.88
  - 1.33
  - 2
67. The bond order of  $O_2^+$  is the same as in [CPMT 2004]
- $N_2^+$
  - $CN^-$
  - $CO$
  - $NO^+$
68. Bond order of  $O_2$  is [DPMT 2004]
- 2
  - 1.5
  - 3
  - 3.5
69. The total number of electron that takes part in forming bonds in  $N_2$  is [MP PET 2004]
- 2
  - 4
  - 6
  - 10
70. The bond length the species  $O_2, O_2^+$  and  $O_2^-$  are in the order of
- $O_2^+ > O_2 > O_2^-$
  - $O_2^+ > O_2^- > O_2$
  - $O_2 > O_2^+ > O_2^-$
  - $O_2^- > O_2 > O_2^+$
71. According to molecular orbital theory which of the following statement about the magnetic character and bond order is correct regarding  $O_2^+$  [IIT JEE Screening 2004]
- Paramagnetic and bond order  $< O_2$
  - Paramagnetic and bond order  $> O_2$
  - Dimagnetic and bond order  $< O_2$
  - Dimagnetic and bond order  $> O_2$
72. The bond order in  $NO$  is 2.5 while that in  $NO^+$  is 3. Which of the following statements is true for these two species [AIEEE 2004]
- Bond length in  $NO^+$  is equal to that in  $NO$
  - Bond length in  $NO$  is greater than in  $NO^+$
  - Bond length in  $NO^+$  is greater than in  $NO$  [UPSEAT 2003]
  - Bond length is unpredictable
73. Which of the following is diamagnetic [BVP 2004]
- Oxygen molecule
  - Boron molecule
  - $N_2^+$
  - None
74. Bond energies in  $NO, NO^+$  and  $NO^-$  are such as [Pb. CET 2004]
- $NO^- > NO > NO^+$
  - $NO > NO^- > NO^+$
  - $NO^+ > NO > NO^-$
  - $NO^+ > NO^- > NO$
75. Which of the following is paramagnetic [UPSEAT 2004]
- $B_2$
  - $C_2$
  - $N_2$
  - $F_2$
76. The paramagnetic molecule at ground state among the following is [UPSEAT 2004]
- $H_2$
  - $O_2$
  - $N_2$
  - $CO$
77. Which has the highest bond energy [DCE 2002]
- $F_2$
  - $Cl_2$
  - $Br_2$
  - $I_2$
78. In  $O_2^-, O_2$  and  $O_2^{2-}$  molecular species, the total number of antibonding electrons respectively are [DCE 2003]
- 7, 6, 8
  - 1, 0, 2
  - 6, 6, 6
  - 8, 6, 8
79. Which of the following is not paramagnetic [DCE 2002]
- $O_2$
  - $O_2^{2+}$
  - $O_2^{2-}$
  - $O_2^-$
80. Which of the following species have maximum number of unpaired electrons [AIIMS 1983]
- $O_2$
  - $O_2^+$
  - $O_2^-$
  - $O_2^{2-}$
81. The correct order in which the  $O-O$  bond length increases in the following is [BHU 2000; CBSE PMT 2005]
- $H_2O_2 < O_2 < O_3$
  - $O_2 < H_2O_2 < O_3$
  - $O_2 < O_3 < H_2O_2$
  - $O_3 < H_2O_2 < O_2$
82. Correct order of bond length is [MP PET 2004] [Orissa JEE 2005]
- $CO_3^{2-} > CO_2 > CO$
  - $CO_2 > CO > CO_3^{2-}$
  - $CO > CO_2 > CO_3^{2-}$
  - None of these
83. Which of the following is paramagnetic [DPMT 2005]
- $N_2$
  - $C_2$
  - $N_2^+$
  - $O_2^{2-}$
84. Among the following molecules which one have smallest bond angle [Orissa JEE 2005]
- $NH_3$
  - $PH_3$
  - $H_2O$
  - $H_2S$

## Hydrogen bonding

1. In the following which bond will be responsible for maximum value of hydrogen bond
- $O-H$
  - $N-H$

- (c)  $S-H$  (d)  $F-H$
2. In which of the following hydrogen bond is present  
(a)  $H_2$  (b) Ice  
(c) Sulphur (d) Hydrocarbon
3. In the following which has highest boiling point  
[MP PMT 1989; RPMT 1997]  
(a)  $HI$  (b)  $HF$   
(c)  $HBr$  (d)  $HCl$
4. Which contains hydrogen bond [MP PMT 1989]  
(a)  $HF$  (b)  $HCl$   
(c)  $HBr$  (d)  $HI$
5. Contrary to other hydrogen halides, hydrogen fluoride is a liquid because [MP PMT 1990; AMU 1983; EAMCET 1980]  
(a) Size of  $F$  atom is small  
(b)  $HF$  is a weak acid  
(c)  $HF$  molecule are hydrogen bonded  
(d) Fluorine is highly reactive
6. In the following which species does not contain  $sp^3$  hybridization  
(a)  $NH_3$  (b)  $CH_4$   
(c)  $H_2O$  (d)  $CO_2$
7. As a result of  $sp$  hybridization, we get [IIT 1984]  
(a) Two mutual perpendicular orbitals  
(b) Two orbitals at  $180^\circ$   
(c) Four orbitals in tetrahedral directions  
(d) Three orbitals in the same plane
8. The reason for exceptionally high boiling point of water is [DPMT 1986; NCERT 1976; AMU 1984; EAMCET 1979; MP PMT 1993; AIIMS 1996; KCET 2001; CPMT 2003]  
(a) Its high specific heat  
(b) Its high dielectric constant  
(c) Low ionization of water molecules  
(d) Hydrogen bonding in the molecules of water
9. Which concept best explains that *o*-nitrophenol is more volatile than *p*-nitrophenol [AIIMS 1980, 82; Kurukshetra CEE 1998; MP PET 2002]  
(a) Resonance (b) Hyperconjugation  
(c) Hydrogen bonding (d) Steric hindrance
10. Which contains strongest  $H-H$  bond [IIT 1986; MP PET 1997, 2003; UPSEAT 2001, 03]  
(a)  $O-H \dots S$  (b)  $S-H \dots O$   
(c)  $F-H \dots F$  (d)  $F-H \dots O$
11. Which of the following compound can form hydrogen bonds [NCERT 1978; MP PMT 1997]  
(a)  $CH_4$  (b)  $NaCl$   
(c)  $CHCl_3$  (d)  $H_2O$
12. Of the following hydrides which has the lowest boiling point [CBSE PMT 1987]  
(a)  $NH_3$  (b)  $PH_3$   
(c)  $SbH_3$  (d)  $AsH_3$
13. The pairs of bases in DNA are held together by [NCERT 1978; DPMT 1985; CBSE PMT 1992]  
(a) Hydrogen bonds (b) Ionic bonds  
(c) Phosphate groups (d) Deoxyribose groups
14. Water has high heat of vaporisation due to [AFMC 1982]  
(a) Covalent bonding (b)  $H-H$  bonding  
(c) Ionic bonding (d) None of the above
15. In which of the following compounds does hydrogen bonding occur  
(a)  $SiH_4$  (b)  $LiH$   
(c)  $HI$  (d)  $NH_3$
16. Which among the following compounds does not show hydrogen bonding [MP PMT 1989]  
(a) Chloroform (b) Ethyl alcohol  
(c) Acetic acid (d) Ethyl ether
17. Acetic acid exists as dimer in benzene due to [CPMT 1982]  
(a) Condensation reaction  
(b) Hydrogen bonding [DPMT 1985]  
(c) Presence of carboxyl group  
(d) Presence of hydrogen atom at  $\alpha$ -carbon
18. Which one among the following does not have the hydrogen bond [UPSEAT 2001]  
(a) Phenol (b) Liquid  $NH_3$   
(c) Water (d) Liquid  $HCl$
19. The bond that determines the secondary structure of a protein is [NCERT 1984; MP PMT 1993; KCET 2001; CPMT 2003]  
(a) Coordinate bond (b) Covalent bond  
(c) Hydrogen bond (d) Ionic bond
20.  $HCl$  is a gas but  $HF$  is a low boiling liquid. This is because [NCERT 1984; MP PMT 2001]  
(a)  $H-F$  bond is strong  
(b)  $H-F$  bond is weak  
(c) Molecules aggregate because of hydrogen bonding  
(d)  $HF$  is a weak acid
21. The relatively high boiling point of  $HF$  is due to [NCERT 1984]  
(a) Hydrogen bonding  
(b) Covalent bonding  
(c) Unshared electron pair on  $F$   
(d) Being a halogen acid
22. Water is liquid due to [MADT Bihar 1983]  
(a) Hydrogen bonding (b) Covalent bond  
(c) Ionic bond (d) Vander Waals forces
23. The maximum possible number of hydrogen bonds in which an  $H_2O$  molecule can participate is [MP PMT 1986; MNR 1991; IIT 1992; MP PET 1999]  
(a) 1 (b) 2  
(c) 3 (d) 4
24. Hydrogen bonding is maximum in [IIT 1987; MP PMT 1991; MP PET 1993, 2001; MNR 1995; CPMT 1999; KCET (Med.) 2002]  
(a) Ethanol (b) Diethyl ether

- (c) Ethyl chloride (d) Triethyl amine
25. The hydrogen bond is strongest in [BHU 1987; CBSE PMT 1990, 92]  
 (a) Water (b) Ammonia  
 (c) Hydrogen fluoride (d) Acetic acid
26. The high boiling point of ethanol ( $78.2^\circ\text{C}$ ) compared to dimethyl ether ( $-23.6^\circ\text{C}$ ), though both having the same molecular formulae  $\text{C}_6\text{H}_6\text{O}$ , is due to [MP PMT 1993]  
 (a) Hydrogen bonding  
 (b) Ionic bonding  
 (c) Coordinate covalent bonding  
 (d) Resonance
27. Methanol and ethanol are miscible in water due to [CPMT 1989]  
 (a) Covalent character  
 (b) Hydrogen bonding character  
 (c) Oxygen bonding character  
 (d) None of these
28. B.P. of  $\text{H}_2\text{O}$  ( $100^\circ\text{C}$ ) and  $\text{H}_2\text{S}$  ( $-42^\circ\text{C}$ ) explained by  
 (a) Vander Waal's forces (b) Covalent bond  
 (c) Hydrogen bond (d) Ionic bond
29. Strength of hydrogen bond is intermediate between [DPMT 1991]  
 (a) Vander Waal and covalent  
 (b) Ionic and covalent  
 (c) Ionic and metallic  
 (d) Metallic and covalent
30. In which of the following compounds intramolecular hydrogen bond is present [MP PET 1994]  
 (a) Ethyl alcohol (b) Water  
 (c) Salicylaldehyde (d) Hydrogen sulphide
31. Hydrogen bonding is formed in compounds containing hydrogen and [MP PET 1995]  
 (a) Highly electronegative atoms  
 (b) Highly electropositive atoms  
 (c) Metal atoms with  $d$ -orbitals occupied  
 (d) Metalloids
32. Which of the following compounds in liquid state does not have hydrogen bonding [MP PMT 1996]  
 (a)  $\text{H}_2\text{O}$  (b)  $\text{HF}$   
 (c)  $\text{NH}_3$  (d)  $\text{C}_6\text{H}_6$
33. Compounds showing hydrogen bonding among  $\text{HF}$ ,  $\text{NH}_3$ ,  $\text{H}_2\text{S}$  and  $\text{PH}_3$  are  
 (a) Only  $\text{HF}$ ,  $\text{NH}_3$  and  $\text{PH}_3$   
 (b) Only  $\text{HF}$  and  $\text{NH}_3$   
 (c) Only  $\text{NH}_3$ ,  $\text{H}_2\text{S}$  and  $\text{PH}_3$   
 (d) All the four
34. The high density of water compared to ice is due to [CBSE PMT 1997; BHU 1999; AFMC 2001]  
 (a) Hydrogen bonding interactions  
 (b) Dipole-dipole interactions  
 (c) Dipole-induced dipole interactions  
 (d) Induced dipole-induced dipole interactions
35. Ethanol and dimethyl ether form a pair of functional isomers. The boiling point of ethanol is higher than that of dimethyl ether due to the presence of [AIIMS 1998]  
 (a) Hydrogen bonding in ethanol  
 (b) Hydrogen bonding in dimethyl ether  
 (c)  $\text{CH}_3$  group in ethanol  
 (d)  $\text{CH}_3$  group in dimethyl ether
36. Which of the following hydrogen bonds are strongest in vapour phase [AMU 1999]  
 (a)  $\text{HF} \cdots \text{HF}$  (b)  $\text{HF} \cdots \text{HCl}$   
 (c)  $\text{HCl} \cdots \text{HCl}$  (d)  $\text{HF} \cdots \text{HI}$
37. Which of the following shows hydrogen bonding [CPMT 2000]  
 (a)  $\text{NH}_3$  (b)  $\text{P}$   
 (c)  $\text{As}$  (d)  $\text{Sb}$
38. The boiling point of a compound is raised by [DPMT 2001]  
 (a) Intramolecular hydrogen bonding  
 (b) Intermolecular hydrogen bonding  
 (c) Covalent bonding  
 (d) Ionic covalent
39. The boiling point of water is exceptionally high because [KCET 2001]  
 (a) Water molecule is linear  
 (b) Water molecule is not linear  
 (c) There is covalent bond between  $\text{H}$  and  $\text{O}$   
 (d) Water molecules associate due to hydrogen bonding
40.  $\text{NH}_3$  has a much higher boiling point than  $\text{PH}_3$  because [UPSEAT 2002; MNR 1994]  
 (a)  $\text{NH}_3$  has a larger molecular weight  
 (b)  $\text{NH}_3$  undergoes umbrella inversion  
 (c)  $\text{NH}_3$  forms hydrogen bond  
 (d)  $\text{NH}_3$  contains ionic bonds whereas  $\text{PH}_3$  contains covalent bonds
41. Which one has the highest boiling point [MP PET 2002]  
 (a) Acetone (b) Ethyl alcohol  
 (c) Diethyl ether (d) Chloroform
42. Which of the following compounds has the highest boiling point  
 (a)  $\text{HCl}$  (b)  $\text{HBr}$   
 (c)  $\text{H}_2\text{SO}_4$  (d)  $\text{HNO}_3$
43. Which of the following has minimum melting point [UPSEAT 2003]  
 (a)  $\text{CsF}$  (b)  $\text{HCl}$   
 (c)  $\text{HF}$  (d)  $\text{LiF}$
44. Hydrogen bond energy is equal to [UPSEAT 2003]  
 (a)  $3 - 7 \text{ cal}$  (b)  $30 - 70 \text{ cal}$   
 (c)  $3 - 10 \text{ kcal}$  (d)  $30 - 70 \text{ kcal}$
45.  $\text{H}_2\text{O}$  is a liquid while  $\text{H}_2\text{S}$  is gas due to [BHU 2003]  
 (a) Covalent bonding  
 (b) Molecular attraction  
 (c)  $\text{H} - \text{bonding}$   
 (d)  $\text{H} - \text{bonding}$  and molecular attraction



46.  $H$ -bonding is maximum in [BHU 2003]  
 (a)  $C_6H_5OH$  (b)  $C_6H_5COOH$   
 (c)  $CH_3CH_2OH$  (d)  $CH_3COCH_3$
47. Select the compound from the following which dissolves in water  
 (a)  $CCl_4$  (b)  $CS_2$   
 (c)  $CHCl_3$  (d)  $C_2H_5OH$
48. When two ice cubes are pressed over each other, they unit to form one cube. Which of the following force is responsible for holding them together [NCERT 1978]  
 (a) Vander Waal's forces  
 (b) Hydrogen bond formation  
 (c) Covalent attraction  
 (d) Dipole-dipole attraction
49. Which is the weakest among the following types of bond [NCERT 1979; MADT Bihar 1984]  
 (a) Ionic bond (b) Metallic bond  
 (c) Covalent bond (d) Hydrogen bond
50.  $H$ -bond is not present in [BCECE 2005]  
 (a) Water (b) Glycerol  
 (c) Hydrogen fluoride (d) Hydrogen Sulphide
51. In melting process, structure of solid [CPMT 1982]  
 (a) Remains unchanged (b) Changes  
 (c) Becomes compact (d) None of the above
52. Which of the following has the highest melting point [CPMT 1994]  
 (a)  $Pb$  (b) Diamond  
 (c)  $Fe$  (d)  $Na$
53. In the formation of a molecule by an atom [AFMC 1995]  
 (a) Attractive forces operate  
 (b) Repulsive forces operate  
 (c) Both attractive and repulsive forces operate  
 (d) None of these
54. Which has weakest bond [RPMT 1997]  
 (a) Diamond (b) Neon (Solid)  
 (c)  $KCl$  (d) Ice
55. Which of the following exhibits the weakest intermolecular forces [AIIMS 1999; B]  
 (a)  $He$  (b)  $HCl$   
 (c)  $NH_3$  (d)  $H_2O$
56. Glycerol has strong intermolecular bonding therefore it is [RPET 2000]  
 (a) Sweet (b) Reactive  
 (c) Explosive (d) Viscous
57. Among the following the weakest one is [Pb. PMT 2004; CPMT 2002]  
 (a) Metallic bond (b) Ionic bond  
 (c) Van der Waal's force (d) Covalent bond
58. Lattice energy of alkali metal chlorides follows the order [DPMT 2004]  
 (a)  $LiCl > NaCl > KCl > RbCl > CsCl$   
 (b)  $CsCl > NaCl > KCl > RbCl > LiCl$   
 (c)  $LiCl > CsCl > NaCl > KCl > RbCl$   
 (d)  $NaCl > LiCl > KCl > RbCl > CsCl$
59. In the following which molecule or ion possesses electrovalent, covalent and co-ordinate bond at the same time  
 (a)  $HCl$  (b)  $NH_4^+$   
 (c)  $Cl^-$  (d)  $H_2O_2$
60. Both ionic and covalent bond is present in the following [MNR 1986; MP PMT 2004]  
 (a)  $CH_4$  (b)  $KCl$   
 (c)  $SO_2$  (d)  $NaOH$
61. The formation of a chemical bond is accompanied by [MP PET 1995]  
 (a) Decrease in energy  
 (b) Increase in energy  
 (c) Neither increase nor decrease in energy  
 (d) None of these
62. Chemical bond implies [KCET 2002]

### Types of bonding and Forces in solid

1. In a crystal cations and anions are held together by [EAMCET 1982]  
 (a) Electrons (b) Electrostatic forces  
 (c) Nuclear forces (d) Covalent bonds
2. In the following metals which one has lowest probable interatomic forces [MP PMT 1990]  
 (a) Copper (b) Silver  
 (c) Zinc (d) Mercury
3. In solid argon, the atoms are held together by [NCERT 1981; MP PET 1995]  
 (a) Ionic bonds (b) Hydrogen bonds  
 (c) Vander Waals forces (d) Hydrophobic forces
4. Which one is the highest melting halide [AIIMS 1980]  
 (a)  $NaCl$  (b)  $NaBr$   
 (c)  $NaF$  (d)  $NaI$
5. The enhanced force of cohesion in metals is due to [NCERT 1972]  
 (a) The covalent linkages between atoms  
 (b) The electrovalent linkages between atoms  
 (c) The lack of exchange of valency electrons  
 (d) The exchange energy of mobile electrons
6. Which one of the following substances consists of small discrete molecules [CPMT 1987]  
 (a)  $NaCl$  (b) Graphite  
 (c) Copper (d) Dry ice
7. Which of the following does not apply to metallic bond [CBSE PMT 1989]

[UPSEAT 2001]

- (a) Attraction  
(b) Repulsion  
(c) Neither attraction nor repulsion  
(d) Both (a) and (b)
20. Which of the following statements is true [AIEEE 2002]  
(a)  $HF$  is less polar than  $HBr$   
(b) Absolutely pure water does not contain any ions  
(c) Chemical bond formation take place when forces of attraction overcome the forces of repulsion  
(d) In covalency transference of electron takes place
21. Which of the following statements is true about  $[Cu(NH_3)_4]SO_4$   
(a) It has coordinate and covalent bonds  
(b) It has only coordinate bonds  
(c) It has only electrovalent bonds  
(d) It has electrovalent, covalent and coordinate bonds
22. Blue vitriol has  
(a) Ionic bond (b) Coordinate bond  
(c) Hydrogen bond (d) All the above
23. The number of ionic, covalent and coordinate bonds in  $NH_4Cl$  are respectively [MP PMT 1999]  
(a) 1, 3 and 1 (b) 1, 3 and 2  
(c) 1, 2 and 3 (d) 1, 1 and 3
24. Covalent molecules are usually held in a crystal structure by [MP PET 1995]  
(a) Dipole-dipole attraction  
(b) Electrostatic attraction  
(c) Hydrogen bonds  
(d) Vander Waal's attraction
6. Which combination is best explained by the co-ordinate covalent bond [JIPMER 2001; CBSE PMT 1990]  
(a)  $N_2H_5^+$  (b)  $BaCl_2$   
(c)  $HCl$  (d)  $H_2O$
7. Arrange the following compounds in order of increasing dipole moment. [CPMT 1988]  
(I) Toluene (II)  $m$ -dichlorobenzene  
(III)  $o$ -dichlorobenzene (IV)  $p$ -dichlorobenzene [IIT 1996]  
(a)  $I < IV < II < III$  (b)  $IV < I < II < III$   
(c)  $IV < I < III < II$  (d)  $IV < II < I < III$
8. The correct order of dipole moment is [Roorkee 1999]  
(a)  $CH_4 < NF_3 < NH_3 < H_2O$   
(b)  $NF_3 < CH_4 < NH_3 < H_2O$   
(c)  $NH_3 < NF_3 < CH_4 < H_2O$   
(d)  $H_2O < NH_3 < NF_3 < CH_4$
9. Which of the following has the highest dipole moment [AIIMS 2002]  
(a)  $\begin{array}{c} H \\ \diagdown \\ C = O \\ \diagup \\ H \end{array}$  (b)  $\begin{array}{c} H & CH_3 \\ | & | \\ C & = C \\ | & | \\ CH_3 & H \end{array}$   
(c)  $\begin{array}{c} CH_3 & H \\ | & | \\ C & = C \\ | & | \\ CH_3 & H \end{array}$  (d)  $\begin{array}{c} Cl & CH_3 \\ | & | \\ C & = C \\ | & | \\ CH_3 & Cl \end{array}$
10. Which of the following arrangement of molecules is correct on the basis of their dipole moments [AIIMS 2002]  
(a)  $BF_3 > NF_3 > NH_3$  (b)  $NF_3 > BF_3 > NH_3$   
(c)  $NH_3 > BF_3 > NF_3$  (d)  $NH_3 > NF_3 > BF_3$  [MP PET 2003]
11. The type of hybrid orbitals used by the chlorine atom in  $ClO_2^-$  is  
(a)  $sp^3$  (b)  $sp^2$   
(c)  $sp$  (d) None of these
12. Among the following species, identify the isostructural pairs,  $NF_3, NO_3^-, BF_3, H_3O^+, HN_3$  [IIT 1996]  
(a)  $[NF_3, NO_3^-]$  and  $[BF_3, H_3O^+]$  [IIT 1995]  
(b)  $[NF_3, HN_3]$  and  $[NO_3^-, BF_3]$   
(c)  $[NF_3, H_3O^+]$  and  $[NO_3^-, BF_3]$   
(d)  $[NF_3, H_3O^+]$  and  $[HN_3, BF_3]$
13. In the compound  $CH_2 = CH - CH_2 - CH_2 - C \equiv CH$ , the  $C_2 - C_3$  bond is of the type [IIT 1999]  
(a)  $sp - sp^2$  (b)  $sp^3 - sp^3$

## Critical Thinking

### Objective Questions

1. The values of electronegativity of atoms  $A$  and  $B$  are 1.20 and 4.0 respectively. The percentage of ionic character of  $A - B$  bond is  
(a) 50 % (b) 43 %  
(c) 55.3 % (d) 72.24%
2.  $O_2^{2-}$  is the symbol of ..... ion [EAMCET 2003]  
(a) Oxide (b) Superoxide  
(c) Peroxide (d) Monoxide
3. The number of electrons that are paired in oxygen molecule is  
(a) 7 (b) 8  
(c) 14 (d) 16
4. When  $N_2$  goes to  $N_2^+$ , the  $N - N$  bond distance ..... and when  $O_2$  goes to  $O_2^+$ , the  $O - O$  bond distance ..... [IIT 1996]  
(a) Decrease, increase (b) Increase, decrease  
(c) Increase, increase (d) None of these
5. Which of the following contains a coordinate covalent bond



- (c)  $sp - sp^3$  (d)  $sp^2 - sp^3$
14. The correct order of increasing  $C - O$  bond length of  $CO$ ,  $CO_3^{2-}$ ,  $CO_2$  is [IIT 1999]  
 (a)  $CO_3^{2-} < CO_2 < CO$  (b)  $CO_2 < CO_3^{2-} < CO$   
 (c)  $CO < CO_3^{2-} < CO_2$  (d)  $CO < CO_2 < CO_3^{2-}$
15. In the dichromate dianion [IIT 1999]  
 (a) 4  $Cr - O$  bonds are equivalent  
 (b) 6  $Cr - O$  bonds are equivalent  
 (c) All  $Cr - O$  bonds are equivalent  
 (d) All  $Cr - O$  bonds are non-equivalent
16. Bond length of ethane (I), ethene (II), acetylene (III) and benzene (IV) follows the order [CPMT 1999]  
 (a)  $I > II > III > IV$  (b)  $I > II > IV > III$   
 (c)  $I > IV > II > III$  (d)  $III > IV > II > I$
17. Hybridisation state of chlorine in  $ClF_3$  is [RPET 1999]  
 (a)  $sp^3$  (b)  $sp^3 d$   
 (c)  $sp^3 d^2$  (d)  $sp^3 d^3$
18. Molecular shapes of  $SF_4$ ,  $CF_4$  and  $XeF_4$  are [IIT Screening 2000]  
 (a) The same with 2, 0 and 1 lone pairs of electrons respectively  
 (b) The same, with 1, 1 and 1 lone pairs of electrons respectively  
 (c) Different, with 0, 1 and 2 lone pairs of electrons respectively  
 (d) Different, with 1, 0 and 2 lone pairs of electrons respectively
19. Structure of  $IF_4^+$  and hybridization of iodine in this structure are [UPSEAT 2001]  
 (a)  $sp^3 d$ , Linear  
 (b)  $sp^3 d^2$ , T-shaped  
 (c)  $sp^3 d$ , Irregular tetrahedral  
 (d)  $sp^3 d^2$ , Octahedral
20. In which of the following the central atom does not use  $sp$  hybrid orbitals in its bonding [UPSEAT 2001, 02]  
 (a)  $BeF_3^-$  (b)  $OH_3^+$   
 (c)  $NH_2^-$  (d)  $NF_3$
21. The magnetic moment of  $K_3[Fe(CN)_6]$  is found to be 1.7  $B.M.$ . How many unpaired electron (s) is/are present per molecule [Orissa JEE 2003]  
 (a) 1 (b) 2  
 (c) 3 (d) 4
22.  $N_2$  and  $O_2$  are converted into monocations  $N_2^+$  and  $O_2^+$  respectively. Which is wrong [CBSE PMT 1997]  
 (a) In  $N_2$ , the  $N - N$  bond weakens  
 (b) In  $O_2$ , the  $O - O$  bond order increases  
 (c) In  $O_2$ , paramagnetism decreases  
 (d)  $N_2^+$  becomes diamagnetic
23. The common features among the species  $CN^-$ ,  $CO$  and  $NO^+$  are [IIT Screening 2001]  
 (a) Bond order three and isoelectronic  
 (b) Bond order three and weak field ligands  
 (c) Bond order two and  $\pi$ -acceptors  
 (d) Isoelectronic and weak field ligands
24. The number of  $S - S$  bonds in sulphur trioxide trimer  $S_3O_9$  is  
 (a) Three (b) Two  
 (c) One (d) Zero
25. Strongest intermolecular hydrogen bond is present in the following molecules pairs [IIT 1981; DCE 2000]  
 (a)  $SiH_4$  and  $SiF$   
 (b)  $CH_3 - \overset{O}{\parallel} C - CH_3$  and  $CHCl_3$   
 (c)  $H - \overset{O}{\parallel} C - OH$  and  $CH_3 - \overset{O}{\parallel} C - OH$   
 (d)  $H_2O$  and  $H_2O_2$
26. A compound contains atoms  $X$ ,  $Y$ ,  $Z$ . The oxidation number of  $X$  is +2,  $Y$  is +5 and  $Z$  is -2. Therefore, a possible formula of the compound is [CPMT 1988]  
 (a)  $XYZ_2$  (b)  $X_2(YZ_3)_2$   
 (c)  $X_3(YZ_4)_2$  (d)  $X_3(Y_4Z)_2$
27. Bonds present in  $CuSO_4 \cdot 5H_2O$  is [IIT 1983; DCE 2001]  
 (a) Electrovalent and covalent  
 (b) Electrovalent and coordinate  
 (c) Electrovalent, covalent and coordinate  
 (d) Covalent and coordinate
28. The ionization of hydrogen atom would give rise to [UPSEAT 2001]  
 (a) Hybrid ion (b) Hydronium ion  
 (c) Proton (d) Hydroxyl ion
29. Which can be described as a molecule with residual bonding capacity [JIPMER 2000]  
 (a)  $BeCl_2$  (b)  $NaCl$   
 (c)  $CH_4$  (d)  $N_2$



## Assertion & Reason

For AIIMS Aspirants

Read the assertion and reason carefully to mark the correct option out of the options given below :

- (a) If both assertion and reason are true and the reason is the correct explanation of the assertion.  
 (b) If both assertion and reason are true but reason is not the correct explanation of the assertion.  
 (c) If assertion is true but reason is false.  
 (d) If the assertion and reason both are false.  
 (e) If assertion is false but reason is true.

1. Assertion : Water is a good solvent for ionic compounds but poor one for covalent compounds.  
 Reason : Hydration energy of ions releases sufficient energy to overcome lattice energy and break hydrogen bonds in water, while covalent bonded compounds interact so weakly that even Vander

- Wall's forces between molecules of covalent compounds cannot be broken. [AIIMS 1996]
2. Assertion : The atoms in a covalent molecule are said to share electrons, yet some covalent molecules are polar.  
Reason : In a polar covalent molecule, the shared electrons spend more time on the average near one of the atoms. [AIIMS 1996]
3. Assertion : Diborane is electron deficient  
Reason : There are not enough valence electrons to form the expected number of covalent bonds [AIIMS 2001]
4. Assertion : A resonance hybrid is always more stable than any of its canonical structures  
Reason : This stability is due to delocalization of electrons [AIIMS 1999]
5. Assertion : All  $F-S-F$  angle in  $SF_4$  greater than  $90^\circ$  but less than  $180^\circ$   
Reason : The lone pair-bond pair repulsion is weaker than bond pair-bond pair repulsion [AIIMS 2004]
6. Assertion : The electronic structure of  $O_3$  is   
Reason :  structure is not allowed because octet around cannot be expanded. [IIT 1998]
7. Assertion : Bond order can assume any value number including zero  
Reason : Higher the bond order, shorter is bond length and greater is bond energy [AIIMS 1999]
8. Assertion : Ortho nitrophenol molecules are associated due to the presence of intermolecular hydrogen bonding while paranitrophenol involves intramolecular, hydrogen bonding  
Reason : Ortho nitrophenol is more volatile than the para nitrophenol [AIIMS 1999]
9. Assertion : Nitrogen molecule diamagnetic.  
Reason :  $N_2$  molecule have unpaired electrons.
10. Assertion : Ice is less dense than liquid water.  
Reason : There are vacant spaces between hydrogen bonded water molecules in ice.
11. Assertion : Water is liquid but  $H_2S$  is a gas.  
Reason : Oxygen is paramagnetic.
12. Assertion : Iodine is more soluble in water than in carbon tetrachloride.  
Reason : Iodine is a polar compound.
13. Assertion : *o* and *p*-nitrophenols can be separated by steam distillation.  
Reason : *o*-nitrophenol have intramolecular hydrogen bonding while *p*-nitrophenol exists as associated molecules.
14. Assertion : The fluorine has lower reactivity.  
Reason :  $F-F$  bond has low bond dissociation energy.
15. Assertion :  $\sigma$  is strong while  $\pi$  is a weak bond.  
Reason : Atoms rotate freely about  $\pi$  bond.
16. Assertion : The crystal structure gets stabilized even though the sum of electron gain enthalpy and ionization enthalpy is positive.  
Reason : Energy is absorbed during the formation of crystal lattice.
17. Assertion : Order of lattice energy for same halides are as  $LiX > NaX > KX$ .  
Reason : Size of alkaline - earth metal increases from *Li* to *K*.
18. Assertion : Born-Haber cycle is based on Hess's law.  
Reason : Lattice enthalpy can be calculated by Born-Haber cycle.
19. Assertion : Bond energy has order like  $C-C < C=C < C\equiv C$ .  
Reason : Bond energy increases with increase in bond order.
20. Assertion : Electron affinity refers to an isolated atom's attraction for an additional electron while electronegativity is the ability of an element to attract electrons towards itself in a shared pair of electrons.  
Reason : Electron affinity is a relative number and electronegativity is experimentally measurable.
21. Assertion : Geometry of  $SF_4$  molecule can be termed as distorted tetrahedron, a folded square or see saw.  
Reason : Four fluorine atoms surround or form bond with sulphur molecule.
22. Assertion :  $BF_3$  has greater dipole moment than  $H_2S$ .  
Reason : Fluorine is more electronegative than sulphur.
23. Assertion : The bond between two identical nonmetal atoms has a pair of electrons with identical spin.  
Reason : Electrons are transferred fully from one atom to another.
24. Assertion : *B* molecule is diamagnetic.  
Reason : The highest occupied molecular orbital is of  $\sigma$  type. [AIIMS 2005]
25. Assertion : The nearly tetrahedral arrangement of the orbitals about the oxygen atom allows each water molecule to form hydrogen bonds with as many as four neighbouring water molecules.  
Reason : In ice each molecule forms four hydrogen bonds as each molecule is fixed in the space.
26. Assertion : The bond order of helium is always zero.  
Reason : The number of electrons in bonding molecular orbital and antibonding molecular orbital is equal.

# Answers

## Electrovalent bonding

1	b	2	a	3	a	4	c	5	c
6	d	7	d	8	b	9	c	10	d
11	b	12	a	13	d	14	a	15	a
16	c	17	b	18	a	19	d	20	c

21	b	22	d	23	a	24	a	25	b
26	d	27	d	28	c	29	a	30	d
31	b	32	b	33	b	34	d	35	b
36	a	37	b	38	a	39	a	40	c
41	c	42	b	43	d	44	b	45	c
46	c	47	a	48	b	49	c	50	b
51	b	52	b	53	a	54	a	55	a
56	c	57	a	58	c	59	a	60	c
61	a	62	b	63	d	64	d	65	b
66	a	67	abc	68	bd				

### Covalent bonding

1	c	2	c	3	B	4	b	5	d
6	a	7	c	8	a	9	d	10	a
11	b	12	b	13	c	14	b	15	c
16	a	17	a	18	c	19	a	20	b
21	a	22	a	23	c	24	c	25	c
26	c	27	a	28	a	29	a	30	d
31	b	32	a	33	d	34	a	35	d
36	b	37	d	38	c	39	d	40	c
41	b	42	b	43	b	44	b	45	b
46	d	47	d	48	b	49	a	50	a
51	b	52	d	53	c	54	d	55	d
56	d	57	a	58	a	59	d	60	a
61	c	62	a	63	b	64	b	65	b
66	b	67	b	68	d	69	b	70	c
71	c	72	c	73	cd	74	ad	75	ab
76	a								

### Co-ordinate or Dative bonding

1	d	2	b	3	c	4	d	5	c
6	b	7	a	8	d	9	a	10	d
11	c	12	a	13	a	14	b	15	c

### Dipole moment

1	b	2	d	3	d	4	a	5	c
6	c	7	a	8	a	9	c	10	b
11	b	12	d	13	b	14	c	15	d
16	c	17	c	18	a	19	c	20	b
21	d	22	b	23	b	24	b	25	a
26	b	27	b	28	b	29	c	30	a
31	a	32	c	33	a	34	bd	35	a

### Polarisation and Fajan's rule

1	d	2	c	3	b	4	d	5	c
6	a	7	b	8	a	9	c	10	b
11	d	12	c	13	b	14	b	15	d
16	d	17	c	18	b	19	a	20	d
21	a	22	c	23	d	24	a	25	b
26	b								

### Overlapping - $\sigma$ and $\pi$ - bonds

1	c	2	c	3	b	4	b	5	c
6	c	7	c	8	b	9	d	10	c
11	b	12	c	13	a	14	a	15	d
16	a	17	d	18	c	19	d	20	d

### Hybridisation

1	d	2	d	3	d	4	c	5	d
6	a	7	c	8	b	9	d	10	d
11	d	12	a	13	a	14	b	15	a
16	b	17	c	18	a	19	d	20	b
21	c	22	c	23	a	24	c	25	a
26	a	27	b	28	c	29	b	30	a
31	d	32	a	33	d	34	c	35	c
36	b	37	b	38	c	39	b	40	b
41	d	42	b	43	c	44	a	45	c
46	c	47	d	48	b	49	c	50	a
51	b	52	a	53	c	54	c	55	c
56	d	57	b	58	a	59	b	60	c
61	b	62	c	63	b	64	b	65	b
66	a	67	c	68	b	69	c	70	a
71	a	72	a	73	b	74	b	75	d
76	d	77	c	78	a	79	d	80	b
81	c	82	b	83	d	84	a	85	d
86	b	87	d	88	c	89	a	90	c
91	c	92	c	93	a	94	b	95	c
96	a	97	b	98	b	99	b	100	b
101	a	102	b	103	d	104	a	105	b
106	a	107	a	108	b	109	b	110	a
111	a	112	b	113	b	114	d	115	d
116	c	117	c	118	b	119	c	120	a
121	a	122	c	123	a	124	a	125	b
126	c	127	d	128	c	129	c	130	a
131	b	132	b	133	e	134	c	135	d

136	b	137	b	138	d	139	a	140	a
141	a	142	b	143	a	144	a	145	a
146	b	147	c	148	d	149	bcd	150	a
151	ac	152	a						

### Resonance

1	d	2	b	3	b	4	b	5	b
6	c	7	a	8	c	9	b	10	c
11	abcd								

### VSEPR Theory

1	a	2	a	3	b	4	c	5	c
6	b	7	b	8	c	9	b	10	a
11	c	12	a	13	a	14	a	15	c
16	c	17	b	18	d	19	d	20	a
21	a	22	d	23	b	24	d	25	a
26	c	27	b	28	b	29	a	30	a
31	a	32	c	33	c	34	a	35	c
36	b	37	b	38	d	39	d	40	b
41	c	42	a	43	b	44	c	45	d

### Molecular orbital theory

1	a	2	c	3	b	4	b	5	c
6	d	7	c	8	b	9	c	10	b
11	c	12	b	13	c	14	a	15	c
16	c	17	d	18	b	19	c	20	c
21	d	22	c	23	b	24	c	25	a
26	d	27	b	28	b	29	a	30	c
31	c	32	a	33	c	34	a	35	c
36	d	37	b	38	a	39	a	40	c
41	c	42	a	43	b	44	a	45	a
46	c	47	b	48	c	49	c	50	a
51	c	52	b	53	a	54	a	55	a
56	c	57	c	58	c	59	a	60	a
61	a	62	b	63	a	64	c	65	a
66	c	67	a	68	a	69	c	70	a
71	b	72	b	73	d	74	c	75	a
76	b	77	b	78	a	79	c	80	a
81	c	82	a	83	c	84	d		

### Hydrogen bonding

1	d	2	b	3	b	4	a	5	c
---	---	---	---	---	---	---	---	---	---

6	d	7	b	8	d	9	c	10	c
11	d	12	b	13	a	14	b	15	d
16	d	17	b	18	d	19	c	20	c
21	a	22	a	23	d	24	a	25	c
26	a	27	b	28	c	29	a	30	c
31	a	32	b	33	d	34	a	35	a
36	a	37	a	38	b	39	d	40	c
41	a	42	c	43	b	44	c	45	c
46	b	47	d	48	b	49	d	50	d

### Types of bonding and Forces in solid

1	b	2	d	3	c	4	c	5	d
6	d	7	d	8	b	9	b	10	c
11	d	12	a	13	d	14	c	15	a
16	b	17	d	18	a	19	d	20	c
21	d	22	d	23	a	24	d		

### Critical Thinking Question

1	d	2	c	3	c	4	b	5	a
6	a	7	b	8	a	9	a	10	d
11	a	12	c	13	d	14	d	15	b
16	c	17	b	18	d	19	c	20	a
21	a	22	d	23	a	24	d	25	c
26	c	27	c	28	c	29	a		

### Assertion & Reason


1	a	2	a	3	a	4	a	5	c
6	b	7	b	8	e	9	c	10	a
11	b	12	d	13	a	14	e	15	c
16	c	17	c	18	b	19	a	20	c
21	b	22	e	23	d	24	d	25	a
26	a								



# AS Answers and Solutions

## Electrovalent bonding

- (b)  $NaCl$  is ionic crystal so it is formed by  $Na^+$  and  $Cl^-$  ions.
- (a) Bond formation is always exothermic. Compounds of sodium are ionic.
- (a) According to Fajan's rule ionic character is less.
- (c) Valencies of  $L$ ,  $Q$ ,  $P$  and  $R$  is  $-2$ ,  $-1$ ,  $+1$  and  $+2$  respectively so they will form  $P_2L$ ,  $RL$ ,  $PQ$  and  $RQ_2$ .
- (c) Electrovalent compounds are good conductor of heat and electricity in molten state or in aqueous solution.
- (d) Electrovalent bond formation depends on ionization energy of cation, electron affinity of anion and on lattice energy.
- (b) Because  $CsF$  is electrovalent compound.
- (c)  $NaCl$  is formed by electrovalent bonding.
- (d) Valency of metal is  $+2$  by formula  $MO$  so its phosphate would be  $M_3(PO_4)_2$  because valency of  $[PO_4]$  is  $-3$ .
- (b)  $Li$ ,  $Na$  and  $K$  are alkali metals with low ionization energy and one electron in their outermost shell so they will form cation easily.
- (a) Melting point and boiling point of electrovalent compounds are high due to strong electrostatic force of attraction between the ions.
- (d) The value of lattice energy depends on the charges present on the two ions and distance between them. It shall be high if charges are high and ionic radii are small.
- (a)  $Cs$  is more electropositive.
- (a)  $X$  loses electron,  $Y$  gains it.
- (c) Formation of  $NaCl$  occurs by  $Na_{ion}^+$  and  $Cl_{ion}^-$ .
- (b)  $MgCl_2$  has electrovalent linkage because magnesium is electropositive metal while chlorine is electronegative.
- (a) Electrovalent compounds generally have high m.pt and high b.pt due to stronger coulombic forces of attractions.
- (d) Water is a polar solvent so it decreases the interionic attraction in the crystal lattice due to solvation.
- (c) Element  $C$  has electronic structure  $1s^2, 2s^2 2p^5$ , it requires only one electron to complete its octet and it will form anion so it will form electrovalent bond.
- (b) Since the chloride of a metal is  $MCl_2$  therefore metal ' $M$ ' must be divalent i.e.  $M^{2+}$ . As a result the formula of its phosphate is  $M_3(PO_4)_2$ .
- (d) In  $MPO_4$  the oxidation state of  $M$  is  $+3$ . Hence, the formula of nitrate is  $M(NO_3)_3$ .
- (a) Ion is formed by gaining or losing electrons. To form cation electron are lost from the valency shell, so  $Zn$  atoms to  $Zn^{++}$  ions there is a decrease in the no. of valency electron.
- (a)  $M_3(PO_4)_2$  means  $M$  is divalent so formula of its sulphate is  $M_2SO_4$ .
- (b) As the molecular formula of chloride of a metal  $M$  is  $MCl_3$ , it is trivalent so formula of its carbonate will be  $M_2(CO_3)_3$ .
- (d) Sodium chloride is electrovalent compound so it dissolves in water which is a polar solvent.
- (d) When sodium chloride is dissolved in water, the sodium ion is hydrated.
- (d) Yet the formula of sulphate of a metal ( $M$ ) is  $M_2(SO_4)_3$ , it is  $M^{3+}$  ion so formula of its phosphate would be  $MPO_4$ .
- (b) Molten sodium chloride conducts electricity due to the presence of free ions.
- (b) The phosphate of a metal has the formula  $MHPO_4$  it means metal is divalent so its chloride would be  $MCl_2$ .
- (d)
- (b)  $Cs$  is highly electropositive while  $F$  is highly electronegative so they will form ionic bond.
- (b)  $Na$  is highly electropositive while  $Cl$  is highly electronegative so they will form ionic bond.
- (a) Ionic compounds are good conductors of heat and electricity so they are good electrolyte.
- (a) Metal tends to lose electrons due to low ionization energy.
- (c) As the formula of calcium pyrophosphate is  $Ca_2P_2O_7$  means valency of pyrophosphate radical is  $-4$  so formula of ferric pyrophosphate is  $Fe_4(P_2O_7)_3$ .
- (c)  $M-X$  bond is a strongest bond so between  $Na-Cl$  is a strongest bond.
- (b) The solubility order is :  
 $BeF_2 > MgF_2 > CaF_2 > SrF_2$  so  $SrF_2$  is least soluble.
- (d)  $NaF$  has maximum melting point, melting point decreases of sodium halide with increase in size of halide their bond energy get lower.
- (b) Sulphanilic acids have bipolar structure so their melting point is high and insoluble in organic solvents.
- (c)  $CaCl_2$  will have electrovalent bonding because calcium is electropositive metal while chlorine is electronegative so they will combined with electrovalent bond.
- (a) Electrovalent bond is formed by losing electrons from one atom and gaining electron by other atom i.e. redox reaction.
- (b) Electrovalent compound are polar in nature because they are formed by ions.
- (b)  $CsCl$  has ionic bonding.
- (b) As soon as the electronegativity increases, ionic bond strength increases.
- (b) This  $X$  element is a second group element so its chloride will be  $XCl_2$ .
- (a) When electronegativity difference is from 1.7 to 3.0. This bond is called as ionic bond.
- (a) Ethyl chloride is an organic compound so it will be covalent.
- (a) Lithium oxide and calcium fluoride show ionic characters.
- (a) Generally cation and anion form ionic bond.
- (c) Those atoms which contain  $+ve$  and  $-ve$  sign are known as ion.
- (a) Generally  $Br-F$  contain maximum electronegativity difference compare to other compound.
- (a) Due to greater electronegativity difference.




62. (b)  $Co^{3+} = 3d^6 4s^0$ ,

$$Ni^{4+} = 3d^6 4s^0,$$

64. (d)  $BaCl_2$  contain higher ionic character.  
 66. (a) Electrolytes are compound which get dissociated into their ion in water so it contains electrovalent bond.  
 67. (abc)  $CaH_2, BaH_2, SrH_2$  are ionic hydride.  
 68. (bcd) Generally  $MgCl_2, SrCl_2, BaCl_2$  are ionic compounds so they conduct electricity in fused state.

### Covalent bonding

2. (c) In  $N_2$  molecule each Nitrogen atom contribute  $3e^-$  so total no. of electron's are 6.  
 3. (b) Non-metals readily form diatomic molecules by sharing of electrons. Element  $M(1s^2 2s^2 2p^5)$  has seven electrons in its valence shell and thus needs one more electron to complete its octet. Therefore, two atoms share one electron each to form a diatomic molecule ( $M_2$ )



5. (d) Covalent character depend on the size of cation and anion.  
 6. (a) In graphite all carbon atoms are  $sp^2$ -hybridised and have covalent bond.  
 7. (c) Silica has tendency to form long chain covalent structure such as carbon so it has giant covalent structure.  
 8. (a) All have linear structure.  
 $O = C = O, Cl - Hg - Cl, HC \equiv CH$   
 9. (d) Similar atoms form covalent bond.  
 10. (a) Covalent bond forms when electronegativity difference of two atom is equal to 1.7 or less than 1.7  
 11. (b) Similar atoms form covalent bond.  
 12. (b) Water is a polar solvent while covalent compounds are non-polar so they usually insoluble in water.  
 13. (c)  $BCl_3$  is electron deficient compound because it has only '6' electrons after forming bond.  
 14. (b) Due to its small size and 2 electrons in  $s$ -orbital  $Be$  forms covalent compound.  
 18. (c)  $H_2O$  will formed by covalent bonding.  
 21. (a) Two identical atoms are joined with covalent bond so  $H_2$  will be covalent.  
 23. (c) Element 'X' has atomic no. 7 so its electronic configuration will be 2, 5. So its electron dot symbol would be  $\cdot\cdot\ddot{X}\cdot$ .  
 24. (c)  $C-S$  will be most covalent. Covalent character depend on the size of cation and anion.  
 25. (c)  $HCl$  has ionic character yet it has covalent compound because electronegativity of chlorine is greater than that of hydrogen.  
 26. (c) Order of polarising power  $Be^{++} > Li^+ > Na^+$   
 Hence order of covalent character  $BeCl_2 > LiCl > NaCl$ .

31. (b) Valency of phosphorus in  $H_3PO_4$  is supposed 'x' then  
 $3 + x - 8 = 0, x - 5 = 0, x = 5$ .

33. (d)  $(+1) + x + 3(-2) = 0 \Rightarrow 1 + x - 6 = 0 \Rightarrow x = 6 - 1 = 5$ .

34. (a)  $HCl$  molecule has covalent bond.

35. (d) Electrovalent compounds have high melting point and high boiling point.

36. (b) Middle length of  $H_2 = 74 pm$

$$\text{Length of } H = \frac{74}{2} = 37 pm$$

$$\text{Middle length of } Cl_2 = 198 pm$$

$$\text{Length of } Cl = \frac{198}{2} = 99 pm$$


$$\begin{aligned} \text{Bond length of } HCl &= \text{Length of } H + \text{Length of } Cl \\ &= 37 + 99 = 136 pm \end{aligned}$$

37. (d) Compound has 254 gm of  $I_2$  means  $\frac{254}{127} = 2$  mole, while

$$80 gm O_2 \text{ means } \frac{80}{16} = 5 \text{ mole so they will form compound}$$




38. (c)  $NH_4Cl$  has covalent as well as ionic bond.



39. (d) Covalent character increases when we come down a group so  $CaI_2$  will have highest covalent character.

41. (b) In water molecule three atom are linked by covalent bond.



42. (b)  $:N \equiv N^+ - \ddot{O}:^-$  or  $N \equiv N \rightarrow O$ .

44. (b) The electronic configuration of  $Na(Z = 11)$  is  $1s^2, 2s^2 2p^6, 3s^1$ . The oxide of  $Na$  is  $Na_2O$ .

45. (b) Covalent bond is directional.

47. (d) Bond dissociation energy decreases with increase in size. So  $D$  is smallest.

48. (b) Molecule  $X$  is nitrogen because nitrogen molecule has triple bond. It's configuration will be  $1s^2, 2s^2 2p^3$ .

49. (a)  $PCl_5$  does not follow octet rule, it has 10 electrons in its valence shell.

50. (a) The compound will be  $A_2B_3$  (By criss cross rule).

51. (b) Each nitrogen share 3 electrons to form triple bond.

52. (d) Urea solution does not conduct electricity because it is a covalent compound.


54. (d) Due to the small size and higher ionization energy, boron forms covalent compound.

58. (a)  $BF_3$  contain 6 electron so it is lewis acid.

59. (d) Among the given species. The bond dissociation energy of  $C-O$  bond is minimum in case of  $CO_3^{2-}$  by which

C-O bond become more weaker in  $CO_3^{2-}$  or the bond order of  $CO_3^{2-}$  (1.33) is minimum so the bond become weaker.


60. (a) Valency of  $Na_2S_2O_3$  is supposed to be  $x$ , then  $2 + 2x + (-6) = 0$ ,  $2x - 4 = 0$ ,  $x = 2$ .



62. (a) Among the given choice  $Al$  is least electropositive therefore, the bond between  $Al$  and  $Cl$  will be least ionic or most covalent or the difference in electronegativity of two atom is less than 1.8.

63. (b) Electronic configuration of  $_{16}S^{32} = 1s^2, 2s^2, 2p^6, 3s^2, 3p^4$ . In the last orbit it has only 6 electron. So it require 2 electron to complete its octet, therefore it share 2 electron with two hydrogen atom and forms 2 covalent bond with it.


64. (b) The acidity of hydrides of VI group elements increase from top to bottom as the bond strength  $X-H$  decrease from top to bottom



65. (b) We know that  $Al^{3+}$  cation is smaller than  $Na^+$  (because of greater nuclear charge) According to Fajan's rule, small cation polarise anion upto greater extent. Hence  $Al^{3+}$  polarise  $Cl^-$  ion upto greater extent, therefore  $AlCl_3$  has covalent bond between  $Al$  and  $Cl$  atoms.


66. (b) Sulphur has the second highest catenation property after carbon. Its molecule has eight atom bonded together (i.e.  $S_8$ )

67. (b)  $H_2O_2$  has open book structure.



69. (b) The electronic configuration of nitrogen is  ${}_7N = 1s^2, 2s^2, 2p^3$


It has 5 electrons in valency shell, hence in ammonia molecule it complete its octet by sharing of three electron with three  $H$  atom, therefore it has 8 electrons in its valence shell in ammonia molecule




71. (c) Multiple bonds have more bond energy so  $C \equiv N$  will be the strongest.
72. (c) Diamond, silicon and quartz molecule bounded by covalent bond.
73. (cd)  $C_2H_4$  and  $N_2$  has multiple bonds.
74. (ad)  $CO$  has only 6 electrons while  $PCl_5$  has 10 electrons after sharing so both don't follow octet rule.
76. (a) Among these,  $NaH$  and  $CaH_2$  are ionic hydrides and  $B_2H_6$  and  $NH_3$  are covalent hydrides.

### Co-ordinate or Dative bonding

1. (d)



2. (b)  $H_2SO_4$  has co-ordinate covalent bond.



3. (c)  $NH_3$  has lone pair of electron while  $BF_3$  is electron deficient compound so they form a co-ordinate bond.  $NF_3 \rightarrow BF_3$

4. (d)  $HNO_2$  does not have co-ordinate bond. Structure is  $H-O-N=O$ .


7. (a) Structure of  $N_2O_5$  is  $O = \overset{\downarrow}{N} - O - \overset{\downarrow}{N} = O$ .

9. (a)  $SO_3^{2-}$  has one coordinate bond.  $^-O - \overset{\downarrow}{S} - O^-$

10. (d) Co-ordinate bond is a special type of covalent bond which is formed by sharing of electrons between two atoms, where both the electrons of the shared pair are contributed by one atom. Since this type of sharing of electrons exists in  $O_3$ ,  $SO_3$  and  $H_2SO_4$ . Therefore all these contains coordinate bond.

12. (a)  $CH_3N \equiv C$  contain dative bond.

13. (a)  $H_3PO_4$  is orthophosphoric acid.




15. (c) Sulphuric acid contain, covalent and co-ordinate bond.

### Dipole moment

1. (b)  $CO_2$  is a symmetrical molecule so its dipole moment is zero.
2. (d) These all have zero dipole moment.
3. (d)  $HF$  has largest dipole moment because electronegativity difference of both is high so it is highly polar.
5. (c) Due to its symmetrical structure.
6. (c) Chloroform has 3 chlorine atom and one hydrogen atom attached to the carbon so it is polarised and it will show dipole moment.
8. (a) The dipole moment of two dipoles inclined at an angle  $\theta$  is given by the equation  $\mu = \sqrt{X^2 + Y^2 + 2XY \cos \theta}$   $\cos 90^\circ = 0$ . Since the angle increases from  $90 - 180$ , the value of  $\cos \theta$  becomes more and more  $-ve$  and hence resultant decreases. Thus, dipole moment is maximum when  $\theta = 90^\circ$ .

9. (c) Due to distorted tetrahedral geometry  $SF_4$  has permanent dipole moment



10. (b)  $CCl_4$  has no net dipole moment because of its regular tetrahedral structure.
12. (d)  $HF$  is polar due to difference of electronegativity of hydrogen and fluorine so it shows positive dipole moment.
14. (c)  $BCl_3$  has zero dipole moment because of its trigonal planar geometry.
16. (c) Dipole moment of  $CH_3OH$  is maximum in it.
20. (b)  $CH_4$  have regular tetrahedron so its dipole moment is zero.
22. (b) Ammonia have some dipole moment.
23. (b) Charge of  $e^- = 1.6 \times 10^{-19}$

$$\text{Dipole moment of } HBr = 1.6 \times 10^{-30}$$

$$\text{Inter atomic spacing} = 1 \text{ \AA} = 1 \times 10^{-10} \text{ m}$$

% of ionic character in

$$HBr = \frac{\text{dipole moment of } HBr \times 100}{\text{interspacing distance} \times q}$$

$$= \frac{1.6 \times 10^{-30}}{1.6 \times 10^{-19} \times 10^{-10}} \times 100$$

$$= 10^{-30} \times 10^{29} \times 100 = 10^{-1} \times 100 = 0.1 \times 100 = 10\%$$

25. (a) Carbon tetrachloride has a zero dipole moment because of its regular tetrahedral structure.
27. (b)  $BF_3$  has zero dipole moment.
29. (c) Given ionic charge  $= 4.8 \times 10^{-10}$  e.s.u. and ionic distance  $= 1 \text{ \AA} = 10^{-8} \text{ cm}$  we know that dipole moment = ionic charge  $\times$  ionic distance  $= 4.8 \times 10^{-10} \times 10^{-8}$   
 $= 4.8 \times 10^{-8}$  e.s.u. per cm = 4.8 debye.
30. (a) Higher is the difference in electronegativity of two covalently bonded atoms, higher is the polarity. In  $HCl$  there is high difference in the electronegativity of  $H$  and  $Cl$  atom so it is a polar compound.
31. (a) Linear molecular has zero dipole moment  $CO_2$  has linear structure so it does not have the dipole moment  $O=C=O$ .
32. (c)  $SF_6$  is symmetrical and hence non polar because its net dipole moment is zero.
33. (a) Polarity create due to the difference in electronegativity of both atom in a molecule except  $H_2$  all other molecule have the different atom so they will have the polarity while  $H_2$  will be non polar.
34. (bd) *cis* isomer shows dipole moment while that of *trans* is zero or very low value. *Trans* 1, 2 di-chloro-2-pentene will also show dipole moment due to unsymmetry.


35. (a) % of ionic character

$$= \frac{\text{Experimental value of dipole moment}}{\text{Expected value of dipole moment}}$$

$$= \frac{1.03}{6.12} \times 100 = 16.83\% \approx 17\%$$

## Polarisation and Fajan's rule

- (d)  $BF_3$  is planar while  $NF_3$  is pyramidal due to the presence of lone pair of electron on nitrogen in  $NF_3$ .
- (c)  $H_2O$  is a polar molecule due to electronegativity difference of hydrogen and oxygen.
- (b) When electronegativity difference is more between two joined atoms then covalent bond becomes polar and electron pair forming a bond don't remain in the centre.
- (d) Hexane has symmetrical structure so does not have polarity.
- (c) When two identical atoms form a bond, bond is non-polar.
- (a) According to Fajan's rule, polarisation of anion is influenced by charge and size of cation more is the charge on cation, more is polarisation of anion.
- (a) When two atoms shares two electrons it is an example of covalent bond. This covalent bond may be polar or may be non-polar depends on the electronegativity difference. In given example formula is  $AB$ . So it is polar.
- (c)  $HCl$  is most polar due to high electronegativity of  $Cl$ .
- (b)  $NH_3$  has  $sp^3$  hybridised central atom so it is non planar.
- (d) *p*-dichloro benzene have highest melting point.
- (b)  $NH_4Cl$  has both types of bonds polar and non polar




- (b) Greater the charge of cation more will be its polarising power (according to Fajan's rule).
- (d)  $AlI_3$  Aluminium triiodide shows covalent character. According to Fajan's rule.
- (d) As the size of anion increases, polarity character increases.
- (d) Due to the electronegativity difference.
- (a) We know that greater the difference in electronegativity of two atoms forming a covalent bond. More is its polar nature. In  $HF$  there is a much difference in the electronegatives of hydrogen and fluorine. Therefore ( $HF$ ) is a polar compound.
- (c) Silicon tetrafluoride has a centre of symmetry.
- (d)  $BF_3$  have zero dipole moment.
- (b) According to Fajan's rule largest cation and smallest anion form ionic bond.
- (b) Polarity character is due to the difference in electronegativity of two atoms or molecule.

## Overlapping- $\sigma$ and $\pi$ - bonds

- (c)
- (c) In fluorine molecule formation *p-p* orbitals take part in bond formation.



3. (b)  $\pi$ -bond is formed by lateral overlapping of unhybridised  $p$ - $p$  orbitals.




5. (c) In a double bond connecting two atom sharing of 4 electrons take place as in  $H_2C = CH_2$ .

6. (c)  $C \equiv C$  is a multiple bond so it is strongest.

9. (d) As the bond order increases,  $C-H$  bond energy also increases so it will be greatest in acetylene because its B.O. is 3.




17. (d) We know that trisilylamine is  $sp^2$ -hybridized therefore  $p\pi-d\pi$  bonding is possible due to the availability of vacant  $d$ -orbitals with silicon.



19. (d)  $\ddot{O} = S = \ddot{O}$ : 5 atoms has 12 electrons in its outermost shell. One  $(S-O)\pi$  bond will be  $(p-p)$   $\pi$  bond while two  $(S-O)\pi$  bond will be  $(p-d)$   $\pi$  bond.


20. (d) Structure of  $P_4O_{10}$  is



Each phosphorus is attached to 4 oxygen atoms.

## Hybridisation


1. (d)  $H_2O$  is not linear because oxygen is  $sp^3$  hybridised in  $H_2O$ .



4. (c)  $CO_2$  has  $sp$ - hybridization and is linear.

5. (d)  $No. of e pair = 3 + \frac{1}{2}[3 - 3] = 0$

$No. of e pair = 3 + 0$



$No. of atom bonded to the central atom = 3$   
In case of 3, 3 geometry is Trigonal planar.

6. (a) In  $sp^3$ -hybridisation each  $sp^3$  hybridised orbital has 1/4  $s$ -character.

8. (b) In ethylene both Carbon atoms are  $sp^2$ - hybridised so  $120^\circ$ .

9. (d) Structure of  $sp^3d$  hybridized compound is Trigonal bipyramidal.

10. (d) In  $H-C \equiv C^* - O - H$  the asterisked carbon has a valency of 5 and hence this formula is not correct.

11. (d)  $dsp^3$  hybrid orbitals have bond angles  $120^\circ, 90^\circ$ .

13. (a) In  $BeF_3^-$ ,  $Be$  is not  $sp^3$ -hybridised it is  $sp^2$  hybridised.

17. (c) In molecule  $OF_2$  oxygen is  $sp^3$  hybridised.

18. (a) In  $sp^3$  hybrid orbitals  $s$ -character is 1/4 means 25%.

19. (d)  $XeF_4$  molecule has 'Xe'  $sp^3d^2$  hybridised and its shape is square planar.

20. (b) The bond angle is maximum for  $sp$  hybridisation because two  $sp$  hybridised orbitals lies at angle of  $180^\circ$ .

21. (c)  $C_2H_4Br_2$  has all single bonds so  $C-H$  bond distance is the largest.


23. (a) In methane molecule  $C$  is  $sp^3$  hybridised so its shape will be tetrahedral.

24. (c) In compound  ${}^3CH_2 = {}^2C = {}^1CH_2$  the second carbon  $sp$ -hybridised.

25. (a)  $:\ddot{Cl} \cdot \cdot \ddot{Cl}:$  is the correct electronic formula of  $Cl_2$  molecule because each chlorine has 7 electrons in its valence shell.


26. (a)  $XeF_4$  has  $sp^3d^2$  hybridisation, its shape is square planar.

27. (b) In  $HCHO$ , carbon is  $sp^2$  hybridized



28. (c) Because of the triple bond, the carbon-carbon bond distance in ethyne is shortest.

29. (b) The hybridisation of  $Ag$  in complex  $[Ag(NH_3)_2]^+$  will be  $sp$  because it is a Linear complex.

30. (a) Structure of  $CO_2$  is linear  $O = C = O$  while that of  $H_2O$  is  i.e. bent structure so in  $CO_2$  resultant dipole moment is zero while that of  $H_2O$  has some value.

31. (d)  $CO_2$  is not  $sp^3$  hybridised, it is  $sp$  hybridised.

32. (a) As compare to pure atomic orbitals, hybrid orbitals have low energy.
33. (d)  $CH_2 = \overset{sp^2}{C} = \overset{sp}{C} - \overset{sp^2}{CH} - \overset{sp^3}{CH_3}$  1, 2-butadiene.
36. (b)  $CCl_4$  is  $sp^3$  hybridised so bond angle will be approximately  $109^\circ$ .
40. (b) Ethene has  $sp^2$  hybridised carbon so bond angles are  $120^\circ$ .
44. (a) Acetate ion is  $CH_3 - \overset{\ominus}{O} \begin{array}{l} // \\ \backslash \end{array} O^-$  i.e. one  $C-O$  single bond and one  $C=O$  double bond.
46. (c) Benzene has all carbons  $sp^2$  hybridised and planar in shape.
47. (d) In methane  $C$  is  $sp^3$  hybridized and bond angle is  $109^\circ$ .
56. (d) 
$$\begin{array}{c} H & H & H \\ | & | & | \\ H - C - C - C - H \\ | & | & | \\ H & H & H \end{array}$$
 There are 10 shared pairs of electrons.
58. (a) The diborane molecule has two types of  $B-H$  bond :  
 (i)  $B-H_t$  - It is a normal covalent bond.  
 (ii)  $B-H_b$  - It is a three centred bond.
- 
61. (b)  $PF_5$  involves  $sp^3d$  hybridization and hence has trigonal bipyramidal structure.
62. (c)  $s$ -character in  $sp = \frac{1}{2} \times 100 = 50\%$   
 $s$ -character in  $sp^2 = \frac{1}{3} \times 100 = 33.3\%$   
 $s$ -character in  $sp^3 = \frac{1}{4} \times 100 = 25\%$   
 Hence, maximum  $s$ -character is found in  $sp$ -hybridisation.
63. (b) The molecule of  $PCl_5$  has  $sp^3d$  hybridisation, structure is trigonal bipyramidal.
64. (b) Merging (mixing) of dissimilar orbitals of different energies to form new orbitals is known as hybridisation and the new orbital formed are known as hybrid orbitals. They have similar energy.
65. (b) In  $SO_3$  sulphur is  $sp^2$  hybridized so its shape will be trigonal planar.
66. (a) These all are triangular with  $sp^2$  hybridization.
67. (c) Bond length depends upon bond order and in benzene all  $C-C$  bonds have same bond order.
68. (b) In  $C_2H_2$  each carbon has  $sp$ -hybridization  

$$H - \underset{sp}{C} \equiv \underset{sp}{C} - H$$
70. (a) As  $p$ -character increases the bond angle decreases.  
 In  $sp$  -  $p$ -character  $\frac{1}{2}$ , bond angle -  $180^\circ$   
 In  $sp^2$  -  $p$ -character  $\frac{2}{3}$ , bond angle -  $120^\circ$   
 In  $sp^3$  -  $p$ -character  $\frac{3}{4}$ , bond angle -  $109^\circ$
71. (a)  $sp^3$ -hybridization called tetrahedral because it provides tetrahedral shape to the molecule.
72. (a)  $S$ -atom in  $SF_6$  has  $sp^3d^2$  hybridisation. So, the structure of  $SF_6$  will be octahedral.
74. (b) Structure of  $H_2O_2$  is non-planar. It has open book structure.
75. (d) Structure of  $N_2O$  is similar to  $CO_2$  both have linear structure.
78. (a)  $SnCl_2$  is V-shaped.
79. (d) In  $NH_4^+$  nitrogen is  $sp^3$  hybridised so 4 hydrogen situated at the corners of a tetrahedron.
81. (c) Increasing order of bond angle is  $sp^3 < sp^2 < sp$ .  
 $109^\circ < 120^\circ < 180^\circ$
84. (a)  $NH_4^+$  has  $sp^3$ -hybridized nitrogen so its shape is tetrahedral.
86. (b) Bond angle increases with change in hybridisation in following order  $sp^3 < sp^2 < sp$ .
88. (c) In Diborane boron shows  $sp^3$ -hybridization.
89. (a) Alkene does not show linear structure but it has planar structure due to  $sp^2$ -hybridisation.
90. (c) Generally  $SF_4$  consist of 10 electrons, 4 bonding electron pair and one lone pair of electron, hence it shows  $sp^3d$  hybridization.
92. (c) Atom/Ion Hybridisation  
 $NO_2^+$   $sp$   
 $SF_4$   $sp^3d$  with one lone pair of electron  
 $PF_6^-$   $sp^3d^2$
93. (a)  $PF_3$  consist of three bonding pair electrons and one lone pair of electron hence it shows  $sp^3$ -hybridization.
94. (b)  $NO_2^+$  shows  $sp$ -hybridization. So its shape is linear.
95. (c) Generally octahedral compound show  $sp^3d^2$ -hybridization.
96. (a) In fifth group hydride bond angle decreases from top to bottom  
 $NH_3 > PH_3 > AsH_3 > SbH_3 > BiH_3$ .
97. (b) Generally  $NH_4^+$  shows  $sp^3$  hybridization.
98. (b) We know that single, double and triple bond lengths of carbon in carbon dioxide are  $1.22 \text{ \AA}$ ,  $1.15 \text{ \AA}$  and  $1.10 \text{ \AA}$  respectively.
99. (b) It shows  $sp^2$ -hybridization so it is planar.
101. (a) Bond angle of hydrides decreases down the group.



102. (b) Hybridization of  $N$  in  $NH_3$  is  $sp^3$  that of  $Pt$  in  $[PtCl_4]^{2-}$  is  $dsp^2$  that  $P$  in  $PCl_5$  is  $sp^3d$  and that of  $B$  in  $BCl_3$  is  $sp^2$ .

103. (d)  $NH_4^+$  and  $SO_4^{2-}$  both show  $sp^3$ -hybridization and tetrahedral structure.

104. (a) It shows  $sp^3d^3$ -hybridization. Hence the bond angle is about  $72^\circ$ .

107. (a)  $s$ -character increases with increase in bond angle.

Hybridization	$s\%$	Angle
$sp$	50	$180^\circ$
$sp^2$	33.3	$120^\circ$
$sp^3$	25	$109.28^\circ$
$sp^3d^1$	20	$90^\circ$ and $120^\circ$

108. (b)  $IF_7$  molecule show  $sp^3d^3$ -hybridization.

110. (a)  $PCl_3$  contain three bonding and one lone pair electron. Hence shows  $sp^3$ -hybridization.

111. (a) Ammonia and  $(BF_4)^-$  shows  $sp^3$ -hybridization.

112. (b) For square planar geometry hybridization is  $dsp^2$  involving  $s, p_x, p_y$  and  $d_{x^2-y^2}$  orbital.

113. (b) All carbon atoms of benzene consist of alternate single and double bond and show  $sp^2$  hybridization.

116. (c)  $BCl_3$  molecule show  $sp^2$ -hybridization and planar structure.

117. (c)  $BCl_3$  Boron trichloride molecule show  $sp^2$ -hybridization and trigonal planar structure.

118. (b)  $SO_2$  molecule shows  $sp^2$ -hybridization and bent structure.

119. (c) Due to multiple bonding in  $N_2$  molecule.

120. (a) % of  $s$ -character in

$$CH_4 = \frac{100}{4} = 25, \quad C_2H_4 = \frac{100}{3} = 33, \\ (sp^3) \quad (sp^2)$$

$$C_2H_2 = \frac{100}{2} = 50 \\ (sp)$$

121. (a) Acidic character increases when we come down a group, so  $HI$  is the strongest acid.


122. (c)  $SO_2$  has  $sp^2$  hybridization have the  $V$  shape structure ( $<120^\circ$ ) due to 2 lone pair of electron over  $S$  atom.  $CO_2$  and  $N_2O$  have the  $sp$  hybridization.

123. (a) In  $H_2CO_3$  and  $BF_3$  central atom are in  $sp^2$  hybridization but in  $H_2CO_3$  due to the ionic character of  $O-H$  bond it will be polar (High electronegativity of oxygen).

124. (a) Due to  $sp^3$  hybridization and presence of lone pair of electron on  $p$  atom  $PCl_3$  are of pyramidal shape like that of  $NH_3$ .

125. (b) There is  $sp$  hybridization in  $C_2H_2$  so it has the linear structure.

126. (c) In octahedral molecule six hybrid orbitals directed towards the corner of a regular octahedron with a bond angle of  $90^\circ$ .




according to this geometry, the number of  $X-M-X$  bond at  $180^\circ$  must be three.






127. (d)  $sp^3d^2$  hybrid orbital have octahedral shape

128. (c) In the formation of  $d^2sp^3$  hybrid orbitals two  $(n-1)d$  orbitals of  $e_g$  set [*i.e.*,  $(n-1)d_{z^2}$  and  $(n-1)d_{x^2-y^2}$  orbitals] one  $ns$  and three  $np$  [ $np_x, np_y$  and  $np_z$ ] orbitals combine together and form six  $d^2sp^3$  hybrid orbitals.


129. (c) The correct order of bond angle (Smallest first) is




$$92.6^\circ < 107^\circ < 109^\circ 28' < 120^\circ$$




Square planar



Regular tetrahedral




See saw shaped




Square planar

131. (b)



132. (b) In the formation of  $BF_3$  molecule, one  $s$  and  $2p$  orbital hybridise. Therefore it is  $sp^2$  hybridization.


133. (e) In  $NCl_3$  and  $H_2S$  the central atom of both ( $N$  and  $S$ ) are in  $sp^3$  hybridization state



while in  $BF_3$  and  $NCl_3$  central atoms are in  $sp^2$  and  $sp^3$  hybridization respectively. In  $H_2S$  and  $BeCl_2$  central atom are in  $sp^3$  and  $sp^2$  hybridization. In  $BF_3$ ,  $NCl_3$  &  $H_2S$  central atom are in  $sp^2$ ,  $sp^3$  &  $sp^3$  hybridization and in the central atom are in  $sp^3$  and  $sp$  hybridization.

134. (c)  $C_{\text{ground state}} = 2s^2, 2p_x^1 p_y^1$ ;  $C_{\text{excited state}} = 2s^1, 2p_x^1 p_y^1 p_z^1$   
 $O_{\text{ground state}} = 2s^2, 2p_x^2 p_y^1 p_z^1$


In the formation of  $CO_2$  molecule, hybridization of orbitals of carbon occur only to a limited extent involving only one  $s$  and one  $p$  orbitals there is thus  $sp$  hybridisation of valence shell orbitals of the carbon atom resulting in the formation of two  $sp$  hybrid orbitals.




135. (d) In  $NH_3$ ,  $N$  undergoes  $sp^3$  hybridization. Due to the presence of one lone pair, it is pyramidal in shape.

136. (b)  $NO_2$   $SF_4$   $PF_6^-$   
 $sp$   $sp^3d$   $sp^3d^2$

137. (b) The configuration of  ${}_5B = 1s^2, 2s^2, 2p^1$



138. (d) In  $SO_3$  molecule,  $S$  atom remains  $sp^2$  hybrid, hence it has trigonal planar structure




139. (a) In  $PCl_3$  molecule, phosphorous is  $sp^3$  - hybridised but due to presence of lone pair of electron, it has pyramidal structure



140. (a) The electronic configuration of

$I = [Xe] 5s^2, 5p^5$  hence




$IF_7$  shows  $sp^3d^2$  hybridization. So, its structure is pentagonal bipyramidal.

141. (a) Compound containing highly electronegative element ( $F$ ,  $O$ ,  $N$ ) attached to an electropositive element ( $H$ ) show hydrogen bonding. Fluorine ( $F$ ) is highly electronegative and has smaller size. So hydrogen fluoride shows the strongest hydrogen bonding in the liquid phase.

142. (b) In the ammonia molecule  $N$  atom is  $sp^3$  - hybridized but due to the presence of one lone pair of  $e^-$  (i.e. due to greater  $L_p - b_p$  repulsion) it has distorted tetrahedral (or pyramidal) geometry.




143. (a)  ${}_4Be \rightarrow 1s^2, 2s^2, 2p^0$



144. (a) Except  $CO_3$  other choice  $CO_2$ ,  $CS_2$  and  $BeCl_2$  have  $sp$ -hybridization and shows the linear structure while  $CO_3$  have  $sp^3$  hybridization and show the non linear structure because  $sp^3$  generate tetrahedral structure.

145. (a)  $dsp^3$  or  $sp^3d$  hybridization exhibit trigonal bipyramidal geometry e.g.,  $PCl_5$




146. (b) Carbon has only two unpaired electrons by its configuration but hybridization is a concept by which we can explain its valency 4.

147. (c) Hybridization is due to overlapping of orbitals of same energy content.

148. (d)  $MX_3$  show the  $sp^2$  hybridization in which  $3sp^2$  hybridized orbital of  $M$  bonded by  $3X$  from  $\sigma$  bond and having the zero dipole moment.
149. (bcd)  $SnCl_2$  has V-shaped geometry.
150. (a)  $NF_3$  is predominantly covalent in nature and has pyramidal structure (the central atom is  $sp^3$  hybridised) with a lone pair of electrons in the fourth orbital.
151. (ac)  $PCl_3, NH_3 \rightarrow$  Pyramidal.  
 $CH_4, CCl_4 \rightarrow$  Tetrahedral.
152. (a)  $dsp^3$  or  $sp^3d$ : one  $s^+$  three  $p^+$  one  $d(d_{z^2})$ .

### Resonance

1. (d) Choice (a), (b), (c) are the resonance structures of  $CO_2$ .
2. (b) In  $NH_3$  nitrogen has one lone pair of electron.
5. (b) In  $CN^-$  ion formal negative charge is on nitrogen atom due to lone pair of electrons.
7. (a)  $CH_3 - \overset{\cdot\cdot}{\underset{|}{C}} = CH_2$  has  $9\sigma, 1\pi$  and 2 lone pairs.
8. (c) In resonance structure there should be the same number of electron pairs.
9. (b) There are three resonance structure of  $CO_3^{2-}$  ion.




11. (abcd) It has all the characteristics.

### VSEPR Theory

2. (a) The bond angle in  $PH_3$  would be expected to be close to  $90^\circ$ . (The bond angle  $H-P-H$  in  $PH_3$  is  $93^\circ$ )
3. (b) In  $BF_3$  molecule Boron is  $sp^2$  hybridised so its all atoms are co-planar.
4. (c) Due to  $lp-lp$  repulsions, bond angle in  $H_2O$  is lower ( $104^\circ.5'$ ) than that in  $NH_3$  ( $107^\circ$ ) and  $CH_4$  ( $109^\circ 28'$ ).  $BeF_2$  on the other hand, has  $sp$ -hybridization and hence has a bond angle of  $180^\circ$ .
5. (c) Compound is carbontetrachloride because  $CCl_4$  has  $sp^3$  hybridization 4 orbitals giving regular tetrahedron geometry. In others the geometry is little distorted inspite of  $sp^3$  hybridization due to different atoms on the vertices of tetrahedron.
6. (b)  $SO_4^{2-}$  ion is tetrahedral since hybridization of  $S$  is  $sp^3$ .
7. (b)  $NH_3$  molecule has one lone pair of electrons on the central atom *i.e.* Nitrogen.
8. (c)  $C_2H_2$  has linear structure because carbons are  $sp$ -hybridised and lies at  $180^\circ$ .


9. (b)  $XeF_6$  is distorted Octahedral. It has  $sp^3d^3$  hybridisation with lone pair of electron on  $Xe$ , so its shape is distorted.



12. (a)  $CO_2$  has bond angle  $180^\circ$ .
13. (a) As the  $s$ -character of hybridized orbitals decreases the bond angle also decreases  
 In  $sp^3$  hybridisation:  $s$ -character  $1/4$ , bond angle  $109^\circ$   
 In  $sp^2$  hybridisation:  $s$ -character  $1/3$ , bond angle  $120^\circ$   
 In  $sp$  hybridisation:  $s$ -character  $1/2$ , bond angle  $180^\circ$
14. (a)  $XeF_2$  molecule is Linear because  $Xe$  is  $sp$  hybridised.
15. (c)  $SO_4^{2-}$  has 42 electrons;  $CO_3^{2-}$  has 32 electrons;  $NO_3^-$  has 32 electrons.
16. (c) Molecular oxygen contains unpaired electron so it is paramagnetic (according to MOT).
17. (b) Structure of  $H_2O$  is a bent structure due to repulsion of lone pair of oxygen.
18. (d) Bond angle between two hybrid orbitals is  $105^\circ$  it means orbitals are  $sp^3$  hybridised but to lone pair repulsion bond angle get changed from  $109^\circ$  to  $105^\circ$ . So its % of  $s$ -character is between 22-23%.
22. (d) Number of electrons in  $ClO_2^-$   
 $= 7 + 6 + 6 + 1 = 20$   
 Number of electrons in  $CF_2^+$   $= 7+7+7 - 1=20$ .
23. (b) Central atom having four electron pairs will be of tetrahedral shape.

24. (d)
- 
26. (c) It shows  $sp^2$ -hybridization and show trigonal planar structure.
28. (b)  $H_2S$  show bond angle nearly  $90^\circ$ .
31. (a) Bond angle of hydrides is decreases top to bottom in the group.  
 $NH_3 > PH_3 > AsH_3 > SbH_3$
32. (c)
- 
- Three bond pair and one lone pair of electron.
33. (c) Unpaired electrons are present in  $KO_2$  while others have paired electron  
 $NO_2^+ = 22$  electrons ;  $BaO_2 = 72$  electrons  
 $AlO_2 = 30$  electrons ;  $KO_2 = 35$  electrons
34. (a) Bond angle decreases from  $H_2O$  to  $H_2Te$ .

35. (c)  $BF_3$  does not contain lone pair of electron.  
 36. (b)



Bent T-shaped geometry in which both lone pairs occupy the equatorial position of the trigonal bipyramidal here

$$(l_p - l_p) \text{ repulsion} = 0$$

$$(l_p - b_p) \text{ repulsion} = 4 \text{ and}$$


$$(b_p - b_p) \text{ repulsion} = 2$$

37. (b) The overall value of the dipole moment of a polar molecule depends on its geometry and shape *i.e.*, vectorial addition of dipole moment of the constituent bonds water has angular structure with bond angle  $105^\circ$  as it has dipole moment. However  $BeF_2$  is a linear molecule since dipole moment summation of all the bonds present in the molecule cancel each other.




38. (d)  $BCl_3$ ,  $BBr_3$  and  $BF_3$ , all of these have same structure *i.e.* trigonal planar ( $sp^2$  hybridization) Hence bond angle is same for all of them (*i.e.*, equal to  $120^\circ$ )

39. (d) We know that molecule of  $(NH_3)$  has maximum repulsion due to lone pair of electron. Its shape is pyramidal and is  $sp^3$  hybridization.



- $\therefore NH_3$  has largest bond angle.
41. (c) In  $NH_3$ ,  $sp^3$ -hybridization is present but bond angle is  $106^\circ 45'$  because Nitrogen has lone pair of electron according to VSEPR theory due to  $bp-lp$  repulsion bond angle decreases from  $109^\circ 45'$  to  $106^\circ 45'$ .


42. (a) Bond strength decreases as the size of the halogen increases from  $F$  to  $I$ .
43. (b)  $NH_3$  has pyramidal structure, yet nitrogen is  $sp^3$  hybridised. This is due to the presence of lone pair of electron.
44. (c)  $SiF_4$  has symmetrical tetrahedral shape which is due to  $sp^3$  hybridization of the central sulphur atom in its excited state configuration.  $SF_4$  has distorted tetrahedral or See-Saw geometry which arise due to  $sp^3d$  hybridization of central sulphur atom and due to the presence of lone pair of electron in one of the equatorial hybrid orbital.



## Molecular orbital theory

2. (c)  $B.O. = \frac{\text{No. of bonding } e^- - \text{No. of antibonding } e^-}{2}$
- $$= \frac{8 - 3}{2} = \frac{5}{2} = 2.5$$
3. (b) One bonding M.O. and one anti-bonding M.O.
4. (b)  $O_2^{2-}$  is least stable.
5. (c) B.O. of  $O_2$  is 2, B.O. of  $O_2^{-1}$  is 1.5, B.O. of  $O_2^{+1}$  is 2.5 and of  $O_2^{2-}$  is 1.
6. (d) Hydride of boron does not exist in  $BH_3$  form. It is stable as its dimer di borane ( $B_2H_6$ ).
10. (c)  $O_2^-(2 \times 8 + 1 = 17)$  has odd number of electrons and hence it is paramagnetic. All the remaining molecules/ions, *i.e.*,  $CN^-(6 + 7 + 1 = 14)$  diamagnetic  $NO(7 + 8 = 15)$  has odd number of electrons and hence it is paramagnetic.
11. (c)  $B.O. = \frac{\text{No. of } N_b - \text{No. of } N_a}{2} = \frac{5}{2} = 2.5$ .
12. (b) Bond order of  $O_2^+$  is highest so its bond length is smallest.
13. (c) Oxygen is paramagnetic due to the presence of two unpaired electron :
- $$O_2 = \sigma(1s)^2 \sigma^*(1s)^2 \sigma(2s)^2 \sigma^*(2s)^2$$
- $$\sigma(2p_x)^2 \pi(2p_y)^2 \pi(2p_x)^2 \pi^*(2p_y)^1 \pi^*(2p_z)^1$$
17. (d) In  $CH_3CN$  bond order between  $C$  and  $N$  is 3 so its bond length is minimum.
18. (b)
- |                 | $He_2^+$             | $H_2$                | $H_2^+$       | $H_2^-$              |
|-----------------|----------------------|----------------------|---------------|----------------------|
| $\sigma(1s)$    | $\uparrow\downarrow$ | $\uparrow\downarrow$ | $\uparrow$    | $\uparrow\downarrow$ |
| $\sigma^*(1s)$  | $\uparrow$           | —                    | —             | $\uparrow$           |
| B.O.            | $\frac{1}{2}$        | 1                    | $\frac{1}{2}$ | $\frac{1}{2}$        |
| Magnetic nature | P                    | D                    | P             | P                    |
- (P = Paramagnetic, D = Diamagnetic)
19. (c) Due to unpaired  $e^-$   $ClO_2$  is paramagnetic.
20. (c) The Bond order in  $N_2$  molecule is 3,  $N \equiv N$  Here,  $N_b = 2 + 4 + 2 = 8$  and  $N_a = 2$
- $$\therefore B.O. = (8 - 2) / 2 = 3$$
21. (d)  $H_2^+$  has the bond order  $\frac{1}{2}$ , it has only one electron so it will be paramagnetic.
22. (c) When bond forms between two atom then their energy get lower than that of separate atoms because bond formation is an exothermic process.
23. (b) Valency of  $A$  is 3 while that of  $B$  is 2 so according to Criss Cross rule the formula of the compound between these two will be  $A_2B_3$ .
24. (c) Due to resonance bond order of  $C - C$  bonds in benzene is between 1 and 2.


25. (a) Nitrogen does not have vacant 'd'-orbitals so it can't have +5 oxidation state i.e. the reason  $PCl_5$  exists but  $NCl_5$  does not.
26. (d) Molecules having unpaired electrons show paramagnetism.
27. (b)  $NO_2$  has unpaired electrons so it would be paramagnetic.
30. (c) Helium molecule does not exist as bond order of  $He_2 = 0$ .
31. (c) Structure of  $P_4O_{10}$  is



Each phosphorus is attached to 4 oxygen atoms.

33. (c) B.O. of carbon =  $\frac{N_b - N_a}{2} = \frac{8 - 4}{2} = 2$ .
34. (a) B.O. =  $\frac{N_b - N_a}{2} = \frac{10 - 4}{2} = 3$ .
37. (b) B.O. =  $\frac{N_b - N_a}{2} = \frac{8 - 3}{2} = \frac{5}{2} = 2.5$ .
38. (a) Electronic configuration of  $O_2$  is  
 $O_2 = \sigma(1s)^2 \sigma^*(1s)^2 \sigma(2s)^2 \sigma^*(2s)^2 \sigma(2p_x)^2 \pi(2p_y)^2 \pi(2p_z)^2 \pi^*(2p_y)^1 \pi^*(2p_z)^1$   
 The molecule has two unpaired electrons So, it is paramagnetic
40. (c)  $\pi^* 2p_y$  has two nodal planes.
42. (a) Element with atomic number 26 is Fe. It is a ferromagnetic.
43. (b) Correct Sequence of bond order is  
 $O_2^+ > O_2 > O_2^{2-}$   
 B.O. - 2.5    2    1.5
44. (a) Due to small bond length.
45. (a)  $S^{-2}$  have all paired electrons so it is diamagnetic.
46. (c)  $NO$  has 15 electrons.
47. (b) In the conversion of  $O_2$  into  $O_2^-$  bond order decreases.
49. (c)  $O_2^{2-}$  does not have any unpaired electron so it is diamagnetic.
50. (a)  $O_2^{2-}$  consist of four antibonding electron pair [1s and 2s have two antibonding and  $2p_x 2p_y$  have two antibonding electron pair].
51. (c) The electron's distribution in molecular orbitals is  $1s^2, 2s^1$   
 B.O. =  $\frac{2-1}{2} = \frac{1}{2} = 0.5$ .
52. (b)  $ClO_2^-$  has all paired electrons hence it does not show paramagnetism.
53. (a) B.O. =  $\frac{1}{2}[N_b - N_a]$   
 $N_2 = \frac{1}{2}[10 - 4] = \frac{6}{2} = 3$ ;  $O_2^{2+} = \frac{1}{2}[10 - 4] = \frac{6}{2} = 3$ .
54. (a) B.O. for  $N_2^+ = \frac{1}{2}[N_b - N_a] = \frac{1}{2}[9 - 4] = \frac{5}{2} = 2.5$ .

55. (a)  $H_2O_2$  contain bond angle between two  $O-H$  planes about  $90^\circ$ .
56. (c) Nitrogen molecule has highest bond energy due to presence of triple bond.
57. (c)  $Cu^{2+} = [Ar]_{18} 3d^9 4s^0$  it has one unpaired electron so it is paramagnetic.
59. (a)  $CN^- = 14$  electrons ;  $CO = 14$  electrons  
 B.O. =  $\frac{1}{2}[10 - 4] = \frac{6}{2} = 3$ .
60. (a) B.O. =  $\frac{1}{2}[10 - 5] = \frac{5}{2} = 2.5$ , paramagnetic
61. (a)
64. (c) The paramagnetic property in oxygen came through unpaired electron which can be explained by molecular orbital theory.



65. (a) Bond order =  $\frac{\text{Total number of bonds between atoms}}{\text{Total number of resonating structure}}$   
 $= \frac{5}{4} = 1.25$
66. (c) We know that carbonate ion has following resonating structures  

 Bond order =  $\frac{\text{Total number of bonds between atoms}}{\text{Total number of resonating structure}}$   
 $= \frac{1+1+2}{3} = \frac{4}{3} = 1.33$ .
67. (a)  $O_2^+(15e^-) = K : K^*(\sigma 2s)^2 (\sigma^* 2s)^2 (\sigma 2p_x)^2 (\pi 2p_y)^2 (\pi 2p_z)^2 (\pi^* 2p_y)^1 (\pi^* 2p_z)^0$   
 Hence, bond order =  $\frac{1}{2}(10 - 5) = 2.5$   
 $N_2^+(13e^-) = KK^*(\sigma 2s)^2 (\sigma^* 2s)^2 (\sigma 2p_x)^2 (\pi 2p_y)^2 (\pi 2p_z)^1$   
 Hence, bond order =  $\frac{1}{2}(9 - 4) = 2.5$ .
68. (a) Electronic configuration of  $O_2$  is

$$O_2 = (\sigma 1s)^2 (\sigma^* 1s)^2 (\sigma 2s)^2 (\sigma^* 2s)^2 (\sigma 2s)^2 (\sigma 2p_z)^2$$

$$(\pi 2p_x)^2 (\pi 2p_y)^2 (\pi^* 2p_x)^1 (\pi^* 2p_y)^1$$

$$\text{Hence bond order} = \frac{1}{2} [N_b - N_a] = \frac{1}{2} [10 - 6] = 2.$$

69. (c) Nitrogen form triple bond  $N \equiv N$   
In which 6 electron take part.
70. (a) As bond order increase bond length decrease the bond order of species are

$$= \frac{\text{number of bonding electron} - \text{Number of } a.b. \text{ electron}}{2}$$

$$\text{For } O_2 = \frac{10 - 6}{2} = 2 ;$$

$$O_2^+ = \frac{10 - 5}{2} = 2.5$$

$$O_2^- = \frac{10 - 7}{2} = 1.5$$

So, bond order  $O_2^+ > O_2 > O_2^-$  and bond length are  $O_2^+ > O_2 > O_2^-$ .

71. (b)  $O_2 : \sigma 1s^2, \sigma^* 1s^2, \sigma 2s^2, \sigma^* 2s^2, \sigma 2p_x^2 \left\{ \begin{array}{l} \pi 2p_y^2 \\ \pi 2p_z^2 \end{array} \right\} \left\{ \begin{array}{l} \pi^* 2p_y^1 \\ \pi^* 2p_z^1 \end{array} \right\}$

$$\text{Bond order} = \frac{10 - 6}{2} = 2.0$$

(Two unpaired electrons in antibonding molecular orbital)

$$O_2^+ : \sigma 1s^2, \sigma^* 1s^2, \sigma 2s^2, \sigma^* 2s^2, \sigma 2p_x^2 \left\{ \begin{array}{l} \pi 2p_y^2 \\ \pi 2p_z^2 \end{array} \right\} \left\{ \begin{array}{l} \pi^* 2p_y^1 \\ \pi^* 2p_z^0 \end{array} \right\}$$

$$\text{Bond order} = \frac{10 - 5}{2} = 2.5$$

(One unpaired electron in antibonding molecular orbital so it is paramagnetic)

72. (b) Higher the bond order, shorter will be the bond length, thus  $NO^+$  having the higher bond order that is 3 as compared to  $NO$  having bond order 2 so  $NO^+$  has shorter bond length.

73. (d) Oxygen molecule ( $O_2$ ) boron molecule ( $B_2$ ) and  $N_2^+$  ion, all of them have unpaired electron, hence they all are paramagnetic.

74. (c) Bond order of  $NO^+, NO$  and  $NO^-$  are 3, 2.5 and 2 respectively, bond energy  $\propto$  bond order

75. (a) Paramagnetic property arise through unpaired electron.  $B_2$  molecule have the unpaired electron so it show paramagnetism.

$$B_2 \rightarrow \sigma 1s^2 \sigma^* 1s^2, \sigma 2s^2 \sigma^* 2s^2, \pi 2p_x^1 = \pi 2p_y^1$$

(2 unpaired electron)

$$C_2 \rightarrow \sigma 1s^2 \sigma^* 1s^2, \sigma 2s^2 \sigma^* 2s^2, \pi 2p_x^2, \pi 2p_y^2$$

(No unpaired electron)

$$N_2 \rightarrow \sigma 1s^2 \sigma^* 1s^2, \sigma 2s^2 \sigma^* 2s^2, \sigma 2p_x^2, \pi 2p_y^2, \pi 2p_z^2$$

(No unpaired electron)

$$F_2 \rightarrow \sigma 1s^2, \sigma^* 1s^2, \sigma 2s^2, \sigma^* 2s^2, \sigma 2p_x^2, \pi 2p_y^2, \pi 2p_z^2,$$

(No unpaired electron)

$$\pi^* 2p_y^2, \pi^* 2p_z^2$$

So only  $B_2$  exist unpaired electron and show the paramagnetism.

76. (b)  $O_2 \rightarrow \sigma 1s^2, \sigma^* 1s^2, \sigma 2s^2, \sigma^* 2s^2, \pi 2p_x^2 \left\{ \begin{array}{l} \pi 2p_y^2 \\ \pi 2p_z^2 \end{array} \right\} \left\{ \begin{array}{l} \pi^* 2p_y^1 \\ \pi^* 2p_z^1 \end{array} \right\}$

So two unpaired electron found in  $O_2$  at ground stage by which it shows paramagnetism.

77. (b) Due to greater electron affinity  $Cl_2$  has the highest bond energy.

78. (a) Molecular orbital electronic configuration of these species are :

$$O_2^-(17e^-) = \sigma 1s^2 \sigma^* 1s^2, \sigma 2s^2 \sigma^* 2s^2, \sigma 2p_x^2, \pi 2p_y^2,$$

$$\pi 2p_z^2, \pi^* 2p_y^2, \pi^* 2p_z^1$$

$$O_2(16e^-) = \sigma 1s^2 \sigma^* 1s^2, \sigma 2s^2 \sigma^* 2s^2, \sigma 2p_x^2, \pi 2p_y^2,$$

$$\pi 2p_z^2, \pi^* 2p_y^1, \pi^* 2p_z^1$$

$$O_2^{2-}(18e^-) = \sigma 1s^2 \sigma^* 1s^2, \sigma 2s^2 \sigma^* 2s^2, \sigma 2p_x^2, \pi 2p_y^2,$$


$$\pi 2p_z^2, \pi^* 2p_y^2, \pi^* 2p_z^2$$

Hence number of antibonding electrons are 7, 6, and 8 respectively.

79. (c) Species with unpaired electrons is paramagnetic  $O_2$  has 2 unpaired electrons,  $O_2^-$  has one unpaired,  $O_2^{2-}$  has zero unpaired electrons,  $O_2^{2+}$  has one unpaired.

80. (a)  $O_2$  has 2 unpaired electron while  $O_2^+$  and  $O_2^-$  has one each unpaired electrons while  $O_2^{2+}$  does not have any unpaired electron.

81. (c)  $H-O-O-H, O \leftarrow O = O, O = O$



Due to resonance in  $O_3$   $O-O$  bond length will be in b/w  $O=O$  and  $O-O$ .

82. (a) From valency bond theory, bond order in  $CO$ , i.e.  $:\bar{C} \equiv \overset{+}{O}:$  is 3, that of  $O=C=O$  is 2 while that of  $CO_3^{2-}$  ion is 1.33. Since the bond length increases as the bond order decreases, i.e.  $CO < CO_2 < CO_3^{2-}$ .

83. (c)  $N_2 : KK\sigma(2s)^2 \sigma^*(2s)^2 \pi(2p_x)^2 \pi(2p_y)^2 \sigma(2p_z)^2$  (diamagnetic)

$$C_2 : KK\sigma(2s)^2 \sigma^*(2s)^2 \pi(2p_x)^2 \pi(2p_y)^2$$

(diamagnetic)

$$N_2^+ : KK\sigma(2s)^2 \sigma^*(2s)^2 \pi(2p_x)^2 \pi(2p_y)^2 \sigma(2p_z)^2$$

(paramagnetic)

$$O_2^{2-} : KK\sigma(2s)^2 \sigma^*(2s)^2 \sigma(2p_z)^2 \pi(2p_x)^2 \pi(2p_y)^2$$

$$\pi^*(2p_x)^2 \pi^*(2p_y)^2$$

(diamagnetic)

84. (d)  $NH_3 = 107^\circ, PH_3 = 93^\circ, H_2O = 104.5^\circ$

$$H_2Se = 91^\circ, H_2S = 92.5^\circ$$



## Hydrogen bonding


- (d) Hydrogen bonding will be maximum in  $F-H$  bond due to greater electronegativity difference.
- (b) Ice has hydrogen bonding.
- (b)  $H - F$  has highest boiling point because it has hydrogen bonding.
- (d)  $CO_2$  is  $sp$ -hybridised
- (b)  $sp$ -hybridization gives two orbitals at  $180^\circ$  with Linear structure.
- (d) Hydrogen bonding increases the boiling point of compound.
- (c)  $o$ -Nitrophenol has intramolecular hydrogen bonding but  $p$ -Nitrophenol has intermolecular hydrogen bonding so boiling point of  $p$ -Nitrophenol is more than  $o$ -Nitrophenol.
- (c) The strongest hydrogen bond is in hydrogen fluoride because the power of hydrogen bond  $\propto$  electronegativity of atom and

$$\text{electronegativity} \propto \frac{1}{\text{atomic size}}$$

So fluorine has maximum electronegativity and minimum atomic size.

- (d)  $H_2O$  can form hydrogen bonds rest  $CH_4$  and  $CHCl_3$  are organic compound having no oxygen while  $NaCl$  has itself intraionic attraction in the molecule.
- (b)  $PH_3$  has the lowest boiling point because it does not form Hydrogen bond.
- (b) Hydrogen bonding increases heat of vaporisation.
- (d) Only  $NH_3$  forms H-bonds.
- (a) Water molecule has hydrogen bonding so molecules get dissociated so it is liquid.
- (d) In case of water, five water molecules are attached together through four hydrogen bonding.
- (c) Hydrogen bond is strongest in hydrogen fluoride.
- (c) Boiling point of  $H_2O$  is more than that of  $H_2S$  because  $H_2O$  forms hydrogen bonding while  $H_2S$  does not.

- (c) Interamolecular H-bonding.




- (a) Hydrogen bond is formed when hydrogen is attached with the atom which is highly electronegative and having small radius.
- (a) Water is dense than ice because of hydrogen bonding interaction and structure of ice.
- (a) Ethanol have hydrogen bonding so its boiling point is higher than its isomer dimethyl ether.
- (a) A compound having maximum electronegative element will form strong Hydrogen bond.
- (a) Due to electronegativity difference of  $N_2$  and  $H_2$ ,  $NH_3$  form hydrogen bond.
- (b) Intermolecular hydrogen bonding compound contain more b.p. compare to intramolecular hydrogen bonding compound.
- (d) Water molecule contain hydrogen bonding.
- (c) It contain intermolecular hydrogen bonding.
- (b) Ethyl alcohol has a intermolecular hydrogen bond.
- (b)  $HCl$  contain weak covalent bond.
- (c) Due to intermolecular hydrogen bonding water molecules come close to each other and exist in liquid state.
- (b) Due to greater resonance stabilization.


- (d)  $C_2H_5OH$  will dissolve in water because it forms hydrogen bond with water molecule.
- (b) In ice cube all molecules are held by inter molecular hydrogen bond.
- (d) Hydrogen bonding is developed due to inter atomic attraction so it is the weakest.

## Types of bonding and Forces in solid

- (b) In electrovalent crystal has cation and anion are attached by electrostatic forces.
- (d) Mercury has very weak interatomic forces so it remains in liquid state.
- (c) The melting and boiling points of argon is low hence, in solid argon atoms are held together by weak Vander Waal's forces.
- (c)  $NaF$  is the strongest ionic crystal so its melting point would be highest.
- (b) Diamond is the hardest substance it's melting point would be highest.
- (c) Bond is formed by attractive and repulsive forces of both the atoms.
- (a) Generally zero group elements are linked by the Vander Waal's force. Hence these show weakest intermolecular forces.
- (d) Glycerol has a three  $OH$  group hence it is viscous in nature.
- (c) Vander waal's forces is the weakest force of attraction.
- (b)  $NH_4^+$  contain all three types of bond in its structure



- (d) In  $NaOH$  covalent bond is present in  $O - H$  bond while ionic bond is formed between  $OH^-$  and  $Na^+$ .
- (a) Bond formation is an exothermic reaction so there is decrease in energy of product.
- (d) Blue vitriol is  $CuSO_4 \cdot 5H_2O$  and it has all types of bonds.




Ionic bond = 1, Covalent bond = 3  
Co-ordinate bond = 1.

## Critical Thinking Questions

- (d) We know that ionic characters  
 $= 16 [E_A - E_B] + 3.5 \times [E_A - E_B]^2$   
 or ionic characters = 72.24%
- (c) Configuration of  $O_2$  molecule is  
 $[\sigma(1s)^2 \sigma^*(1s)^2 \sigma(2s)^2 \sigma^*(2s)^2 \pi(2p_x)^2 \pi(2p_y)^2 \sigma(2p_z)^2 \pi^*(2p_x)^1 \pi^*(2p_y)^1]$   
 No. of pair are 7 so total no. of paired electrons are 14.
- (a)  $H - \overset{\cdot\cdot}{O} : + H^+ \rightarrow H - \overset{\cdot\cdot}{O} \rightarrow H$   
 $\quad \quad \quad | \quad \quad \quad |$   
 $\quad \quad \quad H \quad \quad \quad H$
- (b) The correct order of increasing dipole moment is

*p*-dichlorobenzene < Toluene < *m*-dichlorobenzene < *o*-dichlorobenzene.

8. (a) The dipole moment of  $CH_4 = 0D$ ,  $NF_3 = 0.2D$ ,  $NH_3 = 1.47D$  and  $H_2O = 1.85D$ . Therefore the correct order of the dipole moment is  $CH_4 < NF_3 < NH_3 < H_2O$ .
10. (d) Ammonia molecule is more basic than nitrogen trifluoride and Boron trifluoride because ammonia molecule easily gives lone pair of electron.
11. (a) Chlorine atom in  $ClO_2^-$  is  $sp^3$  hybridised but its shape is angular.
12. (c)  $[NF_3]$  and  $[H_3O^+]$  are pyramidal while  $[NO_3^-]$  and  $[BF_3]$  are planar. Hence answer (c) is correct.
13. (d)  $CH_2 = \underset{\substack{sp^2 \\ \text{hybridised}}}{CH} - \overset{sp^3}{CH_2} - C \equiv CH$
14. (d) B.O. in  $CO$  i.e.,  $:\overset{-}{C} \equiv \overset{+}{O}:$  is 3, that of  $O = C = O$  is 2 while that of  $CO_3^{2-}$  ion is 1.33. Since the bond length increases as the bond order decreases i.e.  $CO < CO_2 < CO_3^{2-}$ . Thus option (d) is correct.
15. (b) Dichromate dianion has following structure




6,  $Cr-O$  bonds are equivalent.

17. (b)  $ClF_3$  is a  $[AB_3]$  type of molecule because it consist of three bonding pair and two lone pair of electrons hence this compound shows  $sp^3d$  hybridization.
20. (a)  $BeF_3^-$  does not show  $sp^3$ -hybridization because this compound is not formed.
21. (a)  $K_3[Fe(CN)_6]$

$$Fe_{26} = 4s^2 3d^6$$


$$Fe^{3+} = 3d^5 4s^0$$




Unpaired electron

$d^2sp^3$ -hybridization

22. (d)  $N_2^+$  has one unpaired electron so it would be paramagnetic.
23. (a) Each of the species has 14 electron so isoelectronic and shows bond order 3.
- $$B.O. = \frac{1}{2}[N_b - N_a] = \frac{1}{2}[10 - 4] = \frac{6}{2} = 3.$$




27. (c)  $CuSO_4 \cdot 5H_2O$  has electrovalent, covalent and coordinate bonds.




## Assertion & Reason

1. (a) Solubility in water depends on hydration energy and lattice energy.
2. (a) Polarity in covalent bond developed due to shifting of electrons towards one of the bonded atoms.
5. (c)  $SiF_4$  have  $sp^3$  hybridization & shape of regular tetrahedral where the bond angle of  $F-S-F$  are found  $109.5^\circ$  which is greater than  $90^\circ$  but less than  $180^\circ$ . Repulsion sequence are  $Lp-Lp > Lp-Bp > Bp-Bp$  so assertion are true but the reason are false.
9. (c)  $N_2$  molecule is diamagnetic. The diamagnetic character is due to the presence of paired electron  $N_2$  molecule does not contain any unpaired electron. Thus, assertion is correct but the reason is false.
10. (a) It is correct that during formation of ice from water there are vacant spaces between hydrogen bonded molecules of ice. Ice has a cage like structure. Due to this reason ice is less dense than liquid water. hence both assertion & reason are true & reason are the correct explanation of assertion.
11. (b) Water is liquid while  $H_2S$  is gas because oxygen is of small size & more electronegative in comparison to sulphur. Hence water molecules exist as associated molecules to form liquid state due to hydrogen bonding  $H_2S$  does not have hydrogen bonding & can't associated hence it is gas.
12. (d) Iodine is more soluble in  $CCl_4$  than in  $H_2O$  because iodine is non polar & thus it dissolve in  $CCl_4$  because like dissolves like.
13. (a) *o* & *p*-nitrophenols can be separated by steam distillation because *o*-nitrophenol is steam volatile. Here, both assertion & reason are correct & reason is correct explanation of assertion.
14. (e) Fluorine is highly reactive  $F-F$  bond has low bond dissociation energy. Here assertion is false but reason is true.
15. (c) It is true that sigma ( $\sigma$ ) bond is stronger than pi ( $\pi$ ) bond but the reason that there is free rotation of atoms is false.
16. (c) Energy is released in the formation of the crystal lattice. It is qualitative measure of the stability of an ionic compound so assertion is true & reason are false.
17. (c)  $Li, Na$  &  $K$  are alkali metals & not alkaline earth metal so, size of alkali metal increases So. Assertion is true & reason are false.
18. (b) Hess's law states that the enthalpy of a reaction is the same, whether it takes place in a single step or in more than one step. In born haber cycle the formation of an cycle ionic compound may occur either by direct combination of the element or by a stepwise process involving vaporization of elements, conversion of the gaseous atoms into ions & the combination of the gaseous ions to form the ionic solid.




19. (a) With increase in bond order, bond length decreases & hence bond energy increases so both assertion & reason are true & reason are the correct explanation of assertion.
20. (c) Electron affinity is experimentally measurable while electronegativity is a relative number so assertion is true but reason are false.
21. (b) Assertion & reason both are correct but reason is not the correct explanation of assertion sulphur has five electrons pairs whose arrangement should be trigonal bipyramidal according to VSEPR theory. Two structure are possible



(a)


Lone pair in the axial position (three l.p - b.p repulsion at  $90^\circ$ )



(b)

Lone pair in the equatorial position (two L.p - b.p repulsion)

22. (e)  $BF_3$  has zero dipole moment because of its structure.




$H_2S$  has two lone pairs on sulphur atom & hence. It has irregular shape.

Thus it possess dipole moment. So assertion is false but reason are true.

23. (d) Both assertion & reason are false because pairs of electron will have different spins. Electrons are equally shared between them.

24. (d) In  $B_2$ , total number of electrons = 10



Presence of unpaired electron shows the paramagnetic nature.

The highest occupied molecular orbital is of  $\pi$ -type.

25. (a) Both assertion & reason are true & reason is the correct explanation of the assertion because. At any given instant, at room temperature each water molecules forms hydrogen bonds with other water molecules. The  $H_2O$  molecules are in continuous motion. So hydrogen bonds are constantly & rapidly broken & formed. In ice  $H_2O$  molecules are however fixed in the space lattice.

26. (a) Both assertion & reason are true & reason is the correct explanation of assertion, because helium molecule is formed by linking two helium atoms. both have  $1s$  orbitals. These will combine to form two molecular orbitals  $\sigma(1s)$  &  $\sigma^*(1s)$  four available electrons are accommodated as  $\sigma(1s)^2$  &  $\sigma^*(1s)^2$ .

- Nature of the bond formed between two elements depends on the
  - Oxidation potential
  - Electronegativity
  - Ionization potential
  - Electron affinity
- Two elements  $X$  and  $Y$  have following electronic configurations  
 $X = 1s^2, 2s^2 2p^6, 3s^2 3p^6, 4s^2$  and  
 $Y = 1s^2, 2s^2 2p^6, 3s^2 3p^5$ . The expected compound formed by combination of  $X$  and  $Y$  is [BHU 1990]
  - $XY_2$
  - $X_5Y_2$
  - $X_2Y_5$
  - $XY_5$
- Electricity do not pass through ionic compounds
  - In solution
  - In solid state
  - In melted state
  - None of these
- From the following which compound on heating readily sublimates
  - $NaCl$
  - $MgCl_2$
  - $BaCl_2$
  - $AlCl_3$
- Which one in the following contains ionic as well as covalent bond [IIT 1979; CPMT 1983; DPMT 1983]
  - $CH_4$
  - $H_2$
  - $KCN$
  - $KCl$
- The solution of sugar in water contains [NCERT 1972; MP PET 2000]
  - Free atoms
  - Free molecules
  - Free ions
  - Free atoms and free molecules
- In which of the following reactions, there is no change in the valency [NCERT 1974; CPMT 1971, 78]
  - $4KClO_3 \rightarrow 3KClO_4 + KCl$
  - $SO_2 + 2H_2S \rightarrow 2H_2O + 3S$
  - $BaO_2 + H_2SO_4 \rightarrow BaSO_4 + H_2O_2$
  - $2BaO + O_2 \rightarrow 2BaO_2$
- The octet rule is not followed in [BHU 1981]
  - $F_2$
  - $NaF$
  - $CaF_2$
  - $BF_3$
- Sodium chloride is an ionic compound whereas hydrogen chloride is a gas because [KCET 2002]
  - Sodium is reactive
  - Covalent bond is weaker than ionic bond
  - Hydrogen chloride is a gas
  - Covalent bond is stronger than ionic bond
- Which one of the following molecules has a coordinate bond
  - $NH_4Cl$
  - $AlCl_3$
  - $NaCl$
  - $Cl_2$
- Co-ordinate bond is absent in [RPMT 2002]
  - $BH_4^\ominus$
  - $CO_3^{2-}$
  - $H_3O^+$
  - $NH_4^\oplus$
- The dipole moment of chlorobenzene is 1.73 D. The dipole moment of  $p$ -dichlorobenzene is expected to be [CPMT 1991]
  - 3.46 D
  - 0.00 D
  - 1.73 D
  - 1.00 D
- Polarization of electrons in acrolein may be written as [IIT 1988]
  - $\overset{\delta^-}{C}H_2 = CH - \overset{\delta^+}{C}H = O$
  - $\overset{\delta^-}{C}H_2 = CH - CH = \overset{\delta^+}{O}$
  - $\overset{\delta^-}{C}H_2 = \overset{\delta^+}{C}H - CH = O$
  - $\overset{\delta^+}{C}H_2 = CH - CH = \overset{\delta^-}{O}$
- The order of dipole moments of the following molecules is [Roorkee 2000]
  - $CHCl_3 > CH_2Cl_2 > CH_3Cl > CCl_4$
  - $CH_2Cl_2 > CH_3Cl > CHCl_3 > CCl_4$
  - $CH_3Cl > CH_2Cl_2 > CHCl_3 > CCl_4$
  - $CH_2Cl_2 > CHCl_3 > CH_3Cl > CCl_4$
- The electronegativity of  $C, H, O, N$  and  $S$  are 2.5, 2.1, 3.5, 3.0 and 2.5 respectively. Which of the following bond is most polar
  - $O-H$
  - $S-H$
  - $N-H$
  - $C-H$
- Which of the following bond has the most polar character [DPMT 1982; CBSE PMT 1992; CPMT 1999]
  - $C-O$
  - $C-Br$
  - $C-S$
  - $C-F$
- The geometry of  $H_2S$  and its dipole moment are [IIT 1999]
  - Angular and non-zero
  - Angular and zero
  - Linear and non-zero
  - Linear and zero
- How many  $\sigma$  and  $\pi$  bonds are there in the molecule of tetracyanoethylene
 
$$N \equiv C \begin{array}{c} \diagup \\ \diagdown \end{array} C = C \begin{array}{c} \diagdown \\ \diagup \end{array} C \equiv N$$
  - Nine  $\sigma$  and nine  $\pi$
  - Five  $\sigma$  and nine  $\pi$
  - Nine  $\sigma$  and seven  $\pi$
  - Five  $\sigma$  and eight  $\pi$
- The shape of  $H_3O^+$  ion is [EAMCET 1993; CPMT 2001]
  - Linear
  - Angular
  - Trigonal planar
  - Triangular pyramidal
- The hybridization in sulphur dioxide is [IIT 1986; DPMT 1990]
  - $sp$  [CPMT 1988, 94]
  - $sp^3$
  - $sp^2$
  - $dsp^2$
- The number and type of bonds between two carbon atoms in  $CaC_2$  are [IIT 1996]
  - One sigma ( $\sigma$ ) and one pi ( $\pi$ ) bonds


- (b) One sigma ( $\sigma$ ) and two pi ( $\pi$ ) bonds  
 (c) One sigma ( $\sigma$ ) and one and a half pi ( $\pi$ ) bonds  
 (d) One sigma ( $\sigma$ ) bond
22. Which of the following resonating structures of  $N_2O$  is the most contributing  
 [Roorkee Qualifying 1998]  
 (a)  $N \equiv N - O$  (b)  $N - N \equiv O$   
 (c)  $N = N - O$  (d)  $N - N = O$
23. The hybridization of atomic orbitals of nitrogen in  $NO_2^+$ ,  $NO_3^-$ , and  $NH_4^+$  are  
 [IIT Screening 2000]  
 (a)  $sp$ ,  $sp^3$  and  $sp^2$  respectively  
 (b)  $sp$ ,  $sp^2$  and  $sp^3$  respectively  
 (c)  $sp^2$ ,  $sp$  and  $sp^3$  respectively  
 (d)  $sp^2$ ,  $sp^3$  and  $sp$  respectively
24. The molecule having one unpaired electron is  
 [IIT 1985; MP PMT 1989]  
 (a)  $NO$  (b)  $CO$   
 (c)  $CN^-$  (d)  $O_2$
25. The geometry of  $ClO_3^-$ , according to valence shell electron pair repulsion (VSEPR) theory will be  
 [KCET 1996; MP PET 1997]  
 (a) Planar triangle (b) Pyramidal  
 (c) Tetrahedral (d) Square planar
26. Which of the following halogens has the highest bond energy  
 (a)  $F_2$  (b)  $Cl_2$   
 (c)  $Br_2$  (d)  $I_2$
27. What bond order does  $O_2^{2-}$  have  
 [Pb. PMT 2001]  
 (a) 3 (b) 2  
 (c) 1 (d) 1/2
28. In the process,  $O_2^+ \rightarrow O_2^{+2} + e$  the electron lost is from  
 [Orissa JEE 2002]  
 (a) Bonding  $\pi$ -orbital (b) Antibonding  $\pi$ -orbital  
 (c)  $2p_z$  orbital (d)  $2p_x$  orbital
29. The maximum number of hydrogen bonds formed by a water molecule in ice is  
 [MP PET 1993; AFMC 2002; UPSEAT 1999, 2001, 02]  
 (a) 4 (b) 3  
 (c) 2 (d) 1
30. Hydrogen bonding is not present in  
 [AIIMS 1998; MP PET/PMT 1998]  
 (a) Glycerine  
 (b) Water  
 (c) Hydrogen sulphide  
 (d) Hydrogen fluoride
31. The bonds in  $K_4[Fe(CN)_6]$  are  
 [EAMCET 1991]  
 (a) All ionic  
 (b) All covalent  
 (c) Ionic and covalent  
 (d) Ionic, covalent and coordinate covalent
32. In which of the following ionic, covalent and coordinate bonds are present  
 [UPSEAT 2002]  
 (a) Water  
 (b) Ammonia  
 (c) Sodium cyanide  
 (d) Potassium bromide

## AS Answers and Solutions

(SET -3)


1. (b) If the two elements have similar electronegativities, the bond between them will be covalent, while a large difference in electronegativities leads to an ionic bond.
2. (a) From electronic configuration valencies of  $X$  and  $Y$  are +2 and -1 respectively so formula of compound is  $XY_2$ .
3. (b) Ionic compounds can't pass electricity in solid state because they don't have mobile ion in solid state.
4. (d)  $AlCl_3$  sublimes readily on heating.
5. (c) Structure of  $KCN$  is  $[K^+(C \equiv \overset{\cdot\cdot}{N})^-]$ .

6. (b) Sugar is an organic compound which is covalently bonded so in water it remains as free molecules.
7. (c) In the reaction  $BaO_2 + H_2SO_4 \rightarrow BaSO_4 + H_2O$  valency is not changing.
8. (d)  $BF_3$  does not have octet, it has only six electrons so it is electron deficient compound.
9. (b)  $NaCl$  is an ionic compound because it consists of more electronegativity difference compare to  $HCl$ .
10. (a)  $NH_4Cl$  has a coordinate bond besides covalent and ionic



11. (b)  $O=C=O$  has covalent bonds only.
12. (b) Due to symmetry dipole moment of *p*-dichloro benzene is zero.
13. (d)
14. (d)  $CCl_4$  has zero dipole moment because of symmetric tetrahedral structure.  $CH_3Cl$  has slightly higher dipole moment which is equal to 1.86D. Now  $CH_3Cl$  has less electronegativity than  $CH_2Cl_2$ . But  $CH_2Cl_2$  has greater dipole moment than  $CHCl_3$ .

15. (a) More the difference in electronegativity of atoms. Bond between them will be more polar.
16. (d)  $C-F$  bond has the most polar character due to difference of their electronegativity.
17. (a)  $H_2S$  has angular geometry and have some value of dipole moment.



18. (a)  $9\pi$  and  $9\sigma$  bonds.
19. (d)  $H_3O^+$  has  $sp^3$  hybridization and its shape is triangular pyramidal due to lone pair on oxygen.
20. (c)  $SO_2$  molecule has  $sp^2$  hybridisation.

21. (b) In  $\begin{array}{c} C \\ ||| \\ C \\ | \\ C \end{array}$  two carbons are joined with  $1\sigma$  and  $2\pi$  bonds.

\*\*\*

22. (a) In  $N_2O$  molecule  $N \equiv N - O$  structure is most contributed.
23. (b) The shape of  $NO_2^+$ ,  $NO_3^-$  and  $NH_4^+$  are linear trigonal planar and tetrahedral respectively. Thus the hybridization of atomic orbitals of nitrogen in these species are  $sp$ ,  $sp^2$  and  $sp^3$  respectively.
24. (a)  $NO$  has one unpaired electron with Nitrogen.




25. (b)  $\begin{array}{c} \cdot\cdot \\ | \\ O - Cl - O \\ | \\ O \end{array}$

26. (b) Bond energy of  $Cl_2$  is highest among all halogen molecule. Bond energies of  $F_2$ ,  $Cl_2$ ,  $Br_2$ ,  $I_2$  are 37, 58, 46 and 36  $Kcal\ mol^{-1}$  respectively.

27. (c)  $O_2^{2-}$  have bond order one


$$B.O. = \frac{1}{2}[10 - 8] = \frac{2}{2} = 1.$$

28. (b) Electron lost from antibonding  $\pi$  orbital.
29. (a) In ice each water molecule forms four hydrogen bond through which each water molecule is tetrahedrally attached with other water molecule.



30. (c) Hydrogen bonding is present in molecules which have F, O, or N atoms.

31. (d) Structure of  $K_4[Fe(CN)_6]$  is



32. (c) Sodium cyanide contain ionic, covalent and coordinate bond.