

Ordinary Thinking

Objective Questions

General Characteristics

1. The number of unpaired electrons in Cr^{+} will be
(a) 3 (b) 4
(c) 5 (d) 6
2. The highest oxidation state of Cr will be
(a) 2 (b) 3
(c) 4 (d) 6
3. Which statement is true about the transitional elements [MP PMT 1995]
(a) They are highly reactive
(b) They show variable oxidation states
(c) They have low M.P.
(d) They are highly electropositive
4. The transitional metal which form green compound in +3 oxidation state and yellow orange compound in +6 oxidation state is
(a) Fe (b) Ni
(c) Cr (d) Co
5. Highest (+7) oxidation state is shown by [MP PMT 1990, 2001; RPMT 1999; AIIMS 1999; JIPMER 2001; CBSE PMT 1994, 2002; MP PET 1989, 2003]
(a) Co (b) Cr
(c) V (d) Mn
6. Transitional elements are
(a) All metals
(b) Few metals and few non-metals
(c) All solids
(d) All highly reactive
7. Which of the following has highest ionic radii [MP PMT 1990]
(a) Cr^{+3} (b) Mn^{+3}
(c) Fe^{+3} (d) Co^{+3}
8. In a reaction the ferrous (Fe^{++}) iron is oxidised to ferric (Fe^{+++}) ion. The equivalent weight of the ion in the above reaction is equal to [CPMT 1985]
(a) Half of the atomic weight
(b) $1/5$ of the atomic weight
(c) The atomic weight
(d) Twice the atomic weight
9. Which of the following element has maximum density
(a) Hg (b) Au
(c) Os (d) Pb
10. Which is heaviest among the following [CPMT 1986]
(a) Iron (b) Copper
(c) Gold (d) Silver
11. Transitional elements exhibit variable valencies because they release electrons from the following orbits [MP PET/PMT 1988; MP PET 1989; UPSEAT 2001]
(a) ns orbit (b) ns and np orbits
(c) $(n-l)d$ and ns orbits (d) $(n-l)d$ orbit
12. The tendency towards complex formation is maximum in
(a) s -block elements (b) p -block elements
(c) d -block elements (d) f -block elements
13. Which forms coloured salts [CPMT 1984; MP PET 1995]
(a) Metals (b) Non-metals
(c) p -block elements (d) Transitional elements
14. Which element belongs to d -block [CPMT 1984]
(a) Na (b) Ca
(c) Cu (d) Ar
15. Variable valency is shown by [MP PMT 1986; AMU 1999]
(a) Na (b) Cu
(c) Mg (d) Al
16. The element with a atomic number 26 is [CPMT 1972]
(a) A non-metal (b) Krypton
(c) Iron (d) Manganese
17. One of the following metals forms a volatile carbonyl compound and this property is taken advantage of for its extraction. This metal is [NCERT 1984]
(a) Iron (b) Nickel
(c) Cobalt (d) Tungston
18. The coinage metals are
(a) Iron, Cobalt, Nickel (b) Copper and Zinc
(c) Copper, Silver and Gold (d) Gold and Platinum
19. Which of the following structure is that of a coinage metal [CPMT 1973, 86]
(a) 2, 8, 1 (b) 2, 8, 18, 1
(c) 2, 8, 8 (d) 2, 18, 8, 3
20. An elements in +3 oxidation state has the electronic configuration $(Ar)3d^3$. Its atomic number is [JIPMER 2002]
(a) 24 (b) 23
(c) 22 (d) 21
21. The catalytic activity of the transition metals and their compounds is ascribed to their [Kerala (Engg.) 2002]
(a) Chemical reactivity
(b) Magnetic behaviour
(c) Unfilled d -orbitals
(d) Ability to adopt multiple oxidation states and their complexing ability
22. What is the general electronic configuration for 2nd row transition series [Orissa JEE 2002]
(a) $[Ne]3d^{1-10}, 4s^2$ (b) $[Ar]3d^{1-10}, 4s^{1-2}$
(c) $[Kr]4d^{1-10}, 5s^{1-2}$ (d) $[Xe]5d^{1-10}, 5s^{1-2}$
23. Transitional elements are named transition elements because their characters are
(a) In between s and p -block elements
(b) Like that of p and d -block elements
(c) They are members of $I-A$ group
(d) They are like inactive elements
24. Those elements whose two outermost orbitals are incompletely filled with electrons are
(a) p -block elements
(b) s -block elements
(c) Transitional elements
(d) Both s and p -block elements
25. Which ion has maximum magnetic moment [AIIMS 1983; MP PMT 1990]
(a) V^{+3} (b) Mn^{+3}
(c) Fe^{+3} (d) Cu^{+2}
26. Which of the following transition metal is present in misch metal
(a) La (b) Sc

- (c) *Ni* (d) *Cr*
27. Which of the following statements is not true in regard to transition elements [EAMCET 1988, 89]
- They readily form complex compounds
 - They show variable valency
 - All their ions are colourless
 - Their ions contain partially filled *d*-electron levels
28. Which of the following represents the electronic configuration of a transition element [EAMCET 1987]
- $1s^2, 2s^2 p^6 \dots n s^2 p^3$
 - $1s^2, 2s^2 p^6 \dots n s^2 p^6 d^3, (n+1)s^2$
 - $1s^2, 2s^2 p^6 \dots n s^2 p^6 d^{10}, (n+1)s^2 p^1$
 - $1s^2, 2s^2 p^6 \dots n s^2 p^6$
29. The general electronic configuration of transition elements is [CPMT 1984, 90, 2002; CBSE PMT 1991, 96; AIIMS 2001; Pb. CET 2000; MP PMT 2003]
- $(n-1)d^{1-5}$ (b) $(n-1)d^{1-10} n s^1$
 - $(n-1)d^{1-10} n s^{1-2}$ (d) $n s^2 (n-1)d^{10}$
30. Transition elements are coloured [MP PMT 1986; Pb. CET 1989; RPET 1999]
- Due to small size
 - Due to metallic nature
 - Due to unpaired *d*-electrons
 - All of these
31. Which of the following has the maximum number of unpaired *d*-electrons [BIT 1992; CBSE PMT 1999]
- Zn* (b) Fe^{2+}
 - Ni^{3+} (d) Cu^+
32. Which is not amphoteric [CPMT 1991]
- Al^{3+} (b) Cr^{3+}
 - Fe^{3+} (d) Zn^{2+}
33. Which does not form amalgam [AFMC 1988; MP PET 2001]
- Fe* (b) *Co*
 - Ag* (d) *Zn*
34. Transition metals are often paramagnetic owing to [Bihar CEE 1992]
- Their high M.P. and B.P.
 - The presence of vacant orbitals
 - The presence of one or more unpaired electrons in the system
 - Their being less electropositive than the elements of groups I-A and II-A
35. Elements which generally exhibit multiple oxidation states and whose ions are usually coloured are [NCERT 1973; MP PMT 2000]
- Metalloids (b) Transition elements
 - Non-metals (d) Gases
36. Which of the following transition metal cation has maximum unpaired electrons [MP PET/PMT 1988; MP PMT 1991; RPMT 1997]
- Mn^{+2} (b) Fe^{+2}
 - Co^{+2} (d) Ni^{+2}
37. Maximum number of oxidation states of transition metal is derived from the following configuration [MP PET/PMT 1988]
- ns* electron (b) $(n-1)d$ electron
 - $(n+1)d$ electron (d) $ns + (n-1)d$ electron
38. Which of the following statement is correct [MP PET/PMT 1988; MP PMT 1991]
- Iron belongs to 3rd transition series of the periodic table
 - Iron belongs to *f*-block of the periodic table
 - Iron belongs to second transition series of the periodic table
 - Iron belongs to group VIII of the periodic table
39. Zinc does not show variable valency like *d*-block elements because [MP PET 1989]
- It is a soft metal
 - d*-orbital is complete
 - It is low melting
 - Two electrons are present in the outermost orbit
40. Which of the following is a transitional element [MP PMT 1989; DPMT 1982]
- Al* (b) *As*
 - Ni* (d) *Rb*
41. Which is not true for transition elements
- They are all metals
 - They show variable valency
 - They form coloured ions
 - They do not form co-ordinate compounds
42. The main reason for not using a mercury electrolytic cell in *NaOH* manufacture is that [CPMT 1988]
- Hg* is toxic
 - Hg* is a liquid
 - Hg* has a high vapour pressure
 - Hg* is a good conductor of electricity
43. Lanthanum is grouped with *f*-block elements because [AMU 2000]
- It has partially filled *f*-orbitals
 - It is just before *Ce* in the periodic table
 - It has both partially filled *f* and *d*-orbitals
 - The properties of Lanthanum are very similar to the elements of *4f* block
44. The element having electronic configuration belongs to $ns^2(n-1)d^{1-10}(n-2)f^{1-14}$ [UPSEAT 2001]
- s*-block (b) *p*-block
 - d*-block (d) *f*-block
45. Variable valency is shown by [UPSEAT 2001]
- Typical elements (b) Normal elements
 - Transition elements (d) None of these
46. Which ion is not coloured [DPMT 2001]
- Cr^{3+} (b) Co^{2+}
 - Cr^{2+} (d) Cu^+
47. The number of unpaired electrons in ferrous ion is [JIPMER (Med.) 2001]
- 5 (b) 4
 - 3 (d) 2
48. *Fe*, *Co* and *Ni* have valuable catalytic properties in process involving

- (a) Organic compound (b) Oxidation
(c) Hydrogenation (d) Compounds of hydrogen
49. Which of the following statement is not correct [NCERT 1983]
(a) Metals contribute their valency electrons to the common sea of electrons
(b) Metals have high co-ordination number
(c) Metals tend to adopt closely packed structures
(d) Metals have high lattice energy
50. Zinc, cadmium and mercury show the properties of
(a) Typical elements (b) Normal elements
(c) Transitional elements (d) Rare elements
51. Iron is
(a) A normal element (b) A typical element
(c) A transitional element (d) An inert element
52. Platinum, palladium, iridium etc., are called noble metals because [NCERT 1975; CPMT 1976]
(a) Alfred Noble discovered them
(b) They are inert towards many common reagents
(c) They are shining lustrous and pleasing to look at
(d) They are found in active state
53. Which of the following statement is not true about *Mohr's salt* [CPMT 1988]
(a) It decolourises $KMnO_4$
(b) It is a primary standard
(c) It is a double salt
(d) Oxidation state of iron is +3 in this salt
54. Which one of the following statement is true for transition elements [AIIMS 1983; AFMC 1987; JIPMER 1997]
(a) They exhibit diamagnetism
(b) They exhibit inert pair effect
(c) They do not form alloys
(d) They show variable oxidation states
55. The valence shell electronic configuration of Cr^{2+} ion is [Orissa JEE 2005]
(a) $4s^0 3d^4$ (b) $4s^2 3d^2$
(c) $4s^2 3d^0$ (d) $3p^6 4s^2$
56. The hardness of *Cr* is due to and metallic lusture is due to
(a) Covalent bond, metallic bond
(b) Covalent bond, hydrogen bond
(c) Metallic bond, covalent bond
(d) Metallic bond, hydrogen bond
57. In the first transition series, the highest B.P. and M.P. is of
(a) *Cr* (b) *V*
(c) *Ni* (d) *Fe*
58. In the following transition elements, the lowest M.P. and B.P. is exhibited by
(a) *Cr* (b) *Hg*
(c) *Cu* (d) *Au*
59. In the following members of transition elements, the lowest ionization energy is of
(a) *Ti* (b) *Sc*
(c) *V* (d) *Mn*
60. Which of the following has second ionisation potential less than expected
(a) *Cr* (b) *Zn*
(c) *V* (d) *Mo*
61. A transition element *X* has a configuration $[Ar]3d^4$ in its + 3 oxidation state. Its atomic number is [CBSE PMT 1996]
(a) 25 (b) 26
(c) 22 (d) 19
62. The transition element have a characteristic electronic configuration which can be represented as [MP PMT/PET 1988; MP PMT 1989]
(a) $(n-2)s^2 p^6 d^{1-10} (n-1)s^2 p^6 ns^2$
(b) $(n-2)s^2 p^6 d^{1-10} (n-1)s^1 p^6 d^1$ or $d^1 ns^1$
(c) $(n-1)s^1 p^6 d^{10} ns^2 p^6 nd^{1-10}$
(d) $(n-1)s^2 p^6 d^{1-10} ns^1$ or ns^2
63. Number of unpaired electrons in Fe^{+++} ($Z = 26$) is [MP PMT 1995; RPET 2003]
(a) 4 (b) 5
(c) 6 (d) 3
64. Of the ions Zn^{2+} , Ni^{2+} and Cr^{3+} [atomic number of *Zn* = 30, *Ni* = 28, *Cr* = 24] [MP PET 1996]
(a) Only Zn^{2+} is colourless and Ni^{2+} and Cr^{3+} are coloured
(b) All three are colourless
(c) All three are coloured
(d) Only Ni^{2+} is coloured and Zn^{2+} and Cr^{3+} are colourless
65. Common oxidation state of scandium, a transition element is/are [atomic number of *Sc* = 21] [MP PET 1996]
(a) + 4 (b) + 1
(c) + 2 and + 3 (d) + 4 and +1
66. Which of the following is not correct about transition metals [MP PET 1996]
(a) Their melting and boiling points are high
(b) Their compounds are generally coloured
(c) They can form ionic or covalent compounds
(d) They do not exhibit variable valency
67. From +6 to +1 oxidation state is shown by the element of..... group
(a) V-B (b) VI-B
(c) VII-B (d) VIII
68. The electronic configuration of cobalt is
(a) $1s^2, 2s^2 2p^6, 3s^2 3p^6 3d^1, 4s^2$
(b) $1s^2, 2s^2 2p^6, 3s^2 3p^6 3d^7, 4s^2$
(c) $1s^2, 2s^2 2p^6, 3s^2 3p^6 3d^3, 4s^2$
(d) $1s^2, 2s^2 2p^6, 3s^2 3p^6 3d^5, 4s^2$
69. Out of all the known elements the number of transitional elements is
(a) 80 (b) 61
(c) 43 (d) 38
70. Cigarette or gas lighter is made up of
(a) Misch metal (b) Alkali metal
(c) Noble metal (d) None
71. Bullet-proof steel alloy is prepared by using
(a) *Sc* (b) *Ni*
(c) *Zr* (d) *Zn*
72. In making gun-berrel, the steel used is
(a) *Fe - Mn* (b) *Fe - Cr*



- (c) $Fe - W$ (d) $Ni - Mo$
73. Neobium and tantalum metals are used in making surgical instruments because they are
(a) Non-corrosive (b) Hard
(c) Soft (d) All
74. To support tungsten filament in electric bulb, the steel used is
(a) Cr (b) Ni
(c) Mn (d) Mo
75. The elements belonging to the first transition series have atomic number from
(a) 19 to 37 (b) 22 to 32
(c) 24 to 30 (d) 21 to 30
76. Which of the following elements does not belong to the first transition series [BHU 2000; MP PMT 1995]
(a) Fe (b) V
(c) Ag (d) Cu
77. Fe^{2+} shows [RPET 2000]
(a) Ferromagnetism (b) Paramagnetism
(c) Diamagnetism (d) None of these
78. Zinc and mercury do not show variable valency like d -block elements because [RPMT 2000; MP PMT 2000]
(a) They are soft
(b) Their d -shells are complete
(c) They have only two electrons in the outermost subshell
(d) Their d -shells are incomplete
79. Cuprous ion is colourless while cupric ion is coloured because
(a) Both have half filled p and d -orbitals
(b) Cuprous ion has incomplete d -orbital and cupric ion has a complete d -orbital
(c) Both have unpaired electrons in the d -orbitals
(d) Cuprous ion has a complete d -orbital and cupric ion has an incomplete d -orbital
80. Transition metals are related to which block [MP PMT 2003; CPMT 1991]
(a) s -block (b) p -block
(c) d -block (d) None of these
81. The number of unpaired electrons in cobalt atom is (atomic number of $Co = 27$) [MP PMT 2003]
(a) 2 (b) 3
(c) 4 (d) 1
82. Zn is related to which group [MP PMT 2003]
(a) 11B (b) 11A
(c) 1A (d) 1B
83. Which of the following element does not show variable valency
(a) Ni (b) Zn
(c) Cu (d) Mn
84. Which of the following is diamagnetic transitional metal ion
(a) Ni^{+2} (b) Zn^{+2}
(c) Co^{+2} (d) Cu^{+2}
85. Which of the following is not an actinide [DPMT 2005]
(a) Curium (b) Californium
(c) Uranium (d) Terbium
86. The ability to form complex compounds by the transitional metal ion is due to
(a) Small size (b) Vacant ' d ' orbitals
(c) High nuclear charge (d) All of these
87. Which of the following will give green hydrated ion
(a) Fe^{+2} (b) Ni^{+2}
- (c) (a) and (b) both (d) V^{+3}
88. Magnetic moment is expressed in
(a) Faraday (b) Calorie
(c) Bohr Magneton (d) Debye
89. If the colours of salts of transition elements are due to the presence of unpaired electrons in the transition metal ions, which of the following ions will be colourless in aqueous solution
(a) Ti^{3+} (b) Ti^{4+}
(c) Fe^{2+} (d) Fe^{3+}
(The at. no. of Ti and Fe are 22 and 26 respectively)
90. In the periodic table first transition series is placed in
(a) Third period (b) Fourth period
(c) Fifth period (d) None of these
91. The element having general electronic configuration $3d^4 4s^1$ is [BHU 1978; CPMT 1991]
(a) Noble gas (b) Non-metal
(c) Metalloid (d) Transition metal
92. Which of the following general configuration of outermost shell represents chromium element [Cr 's atomic number = 24] [CBSE PMT 1991; MP PMT 1992, 2001; RPET/PMT 1999]
(a) $d^5 s^1$ (b) $d^6 s^0$
(c) $d^4 s^2$ (d) $d^3 s^2$
93. Which element gives maximum balanced oxide [MP PMT 1990]
[KCET 2000]
(a) V (b) Cr
(c) Mn (d) Fe
94. In first transition series, the melting point of Mn is low because [MP PMT/PET 1993]
(a) Due to d^{10} configuration, metallic bonds are strong
(b) Due to d^7 configuration, metallic bonds are weak
(c) Due to d^5 configuration, metallic bonds are weak
(d) None of these
95. Which of the following ions has the least magnetic moment [MP PMT 1993]
(a) Cu^{+2} (b) Ni^{+2}
(c) Co^{+3} (d) Fe^{+2}
96. Among the following outermost configurations of transition metals, which shows the highest oxidation state [MP PMT 1993; MP PET 1995, 2001]
[MP PMT 2003]
(a) $3d^2 4s^2$ (b) $3d^5 4s^1$
(c) $3d^5 4s^2$ (d) $3d^6 4s^2$
97. Which of the following is not true for transition metals [MP PET 1993]
(a) They are malleable and ductile
(b) They have high boiling and melting points
(c) They crystallize with body centred cubic and hexagonal close-packed structures only
(d) They show variable oxidation states although not always
98. The most malleable of all the metals is
(a) Silver (b) Sodium
(c) Gold (d) Platinum
99. Paramagnetism is exhibited by molecules [NCERT 1981; Manipal MEE 1995]
(a) Not attracted in a magnetic field
(b) Containing only paired electrons

- (c) Carrying a positive charge
(d) Carrying unpaired electrons
100. The higher number of unpaired electrons are in [DCE 2001]
(a) Fe (b) Fe^+
(c) Fe^{+2} (d) Fe^{+3}
101. Which one of the following statements concerning lanthanides elements is false [AMU 2001]
(a) Lanthanides are separated from one another by ion exchange methods
(b) The ionic radii of trivalent lanthanides steadily increase with increase in atomic number
(c) All lanthanides are highly dense metals
(d) Most typical oxidation of lanthanides is +3
102. Complex ion is shown by [CPMT 2001]
(a) Ag (b) Au
(c) Cu (d) All of these
103. Which of the following transition metal is used as a catalyst [CPMT 2001]
(a) Nickel (b) Cobalt
(c) Gold (d) Both (a) and (b)
104. For which of the following transition metal ion is shown highest magnetic moment having outer electronic configuration
(a) $3d^1$ (b) $3d^8$
(c) $3d^5$ (d) $3d^7$
105. The number of unpaired electrons is maximum in (Atomic no. : $Ti = 22$; $V = 23$; $Cr = 24$; $Fe = 26$) [MP PET 2003]
(a) Cr (b) Fe
(c) V (d) Sc
106. Europium is [DPMT 2005]
(a) s -block element (b) p -block element
(c) d -block element (d) f -block element
107. Which of the following elements is alloyed with copper to form brass [MP PET 2003; MP PMT 2004]
(a) Lead (b) Silver
(c) Zinc (d) Antimony
108. In which of the following metallic bond is strongest [MP PET 2003]
(a) Fe (b) Sc
(c) V (d) Cr
109. Which of the following is a colourless ion [EAMCET 1992]
(a) Cu^{+2} (b) Fe^{+3}
(c) Ti^{+3} (d) Zn^{+2}
110. The substance used in cancer therapy is [DPMT 2002]
(a) Rn (b) Ni
(c) Fe (d) Co
111. In solution of $AgNO_3$, if Cu is a solution become blue due to [MH CET 2002]
(a) Oxidation of Ag (b) Oxidation of Cu
(c) Reduction of Ag (d) Reduction of Cu
112. Lanthanide for which + II and + III oxidation states are common is
(a) La (b) Nd
(c) Ce (d) Eu
113. The number of unpaired electrons in Zn^{++} is
(a) 2 (b) 3
(c) 4 (d) 0
114. The first transition element is
(a) Chromium (b) Scandium
(c) Nickel (d) Copper
115. The electronic configuration (outermost) of Mn^{+2} ion (atomic no. of $Mn = 25$) in its ground state is [MP PET 1993; MP PMT 1994; AFMC 2002]
(a) $3d^5 4s^0$ (b) $3d^4 4s^1$
(c) $3d^3 4s^2$ (d) $3d^2 4s^2 4p^2$
116. The highest magnetic moment is shown by the transition metal ion with the outer electronic configuration [MP PET 1993; MP PMT 1995; RPMT 1999]
(a) $3d^2$ (b) $3d^5$
(c) $3d^7$ (d) $3d^9$
117. Which has valency two
(a) Al (b) Rb
(c) Cu (d) Ge
118. All the following statements about the transitional elements are true except that [Kerala (Med.) 2003]
(a) All of the transitional elements are predominantly metallic
(b) In [MP PET 2003] solution many of their simple ions are coloured
(c) Most of the transitional elements show pronounced catalytic activity
(d) Most of the transitional elements show only one valence state
119. Which of the following transition metals can have highest oxidation state [RPET 2003]
(a) Cr (b) Co
(c) Ni (d) Cu
120. Which of the following melts in boiling water [MNR 1984]
(a) Gun metal (b) Wood's metal
(c) Monel metal (d) Bell metal
(e) None of these
121. On the basis of position in the electrochemical series, the metal which does not displace hydrogen from water and acid is
(a) Hg (b) Al
(c) Pb (d) Ba
122. The tendency of $3d$ -metal ions to form stable complexes is due to their [MP PMT 1997]
(a) Variable oxidation state
(b) Strong electronegative nature
(c) High charge/size ratio and vacant d -orbitals
(d) Very low ionization energies
123. The $3d$ -metal ions are paramagnetic in nature because [MP PMT 1997]
(a) They are reducing agents
(b) They form coloured salts
(c) They have one or more paired s -electrons
(d) They have one or more unpaired d -electrons
124. Lanthanide contraction occurs because [AIIMS 2003] [AMU 2000; BHU 2003]
(a) f -orbital electrons are easily lost
(b) f -orbital an incompletely filled
(c) f -orbital electron a poor shielders of nuclear charge
(d) f -orbital do not come out on the surface of atom and are buried inside

125. Which is most reactive metal [BHU 1979]
 (a) *Fe* (b) *Pt*
 (c) *Ni* (d) *Co*
126. Least reactive metal is
 (a) *Fe* (b) *Os*
 (c) *Ni* (d) *Pt*
127. Which occludes hydrogen
 (a) *Os* (b) *Pt*
 (c) *Ni* (d) All of these
128. Which has the maximum ferromagnetic character
 (a) *Fe* (b) *Co*
 (c) *Ni* (d) *Pt*
129. Which forms interstitial compounds [BHU 1982; MP PMT 1983]
 (a) *Fe* (b) *Co*
 (c) *Ni* (d) All of these
130. Which occurs in nature in free state
 (a) *Fe* (b) *Co*
 (c) *Ni* (d) *Pt*
131. $3d^{10} 4s^0$ electronic configuration exhibits
 (a) Zn^{++} (b) Cu^{++}
 (c) Cd^{++} (d) Hg^{++}
132. $3d^0 4s^0$ electronic configuration exhibits
 (a) Pd^{+2} (b) Sc^{+2}
 (c) Ti^{+4} (d) Zn^{+2}
133. Rare-earth elements are exhibited by
 (a) At. No. 58 to 71 (b) At. No. 21 to 30
 (c) At. No. 39 to 71 (d) At. No. 81 to 91
134. All those elements belong to *f*-block whose atomic numbers are
 (a) 58 to 71 (b) 90 to 103
 (c) Both (a) and (b) (d) None
135. The correct order of density is
 (a) $Cu > Ni > Zn > Sc$
 (b) $Ni > Cu > Zn > Sc$
 (c) $Zn > Cu > Ni > Sc$
 (d) $Sc > Zn > Ni > Cu$
136. The property exhibited by only transitional elements
 (a) To form paramagnetic compounds
 (b) To form coloured compounds
 (c) To form complex compounds
 (d) To show inert tendency
137. Which of the following will have standard oxidation potential less than *SHE*
 (a) *Zn* (b) *Cu*
 (c) *Fe* (d) *Ni*
138. Hydrated Cu^{+2} ion will be
 (a) Green (b) Violet
 (c) Blue (d) Colourless
139. The placement of *Zn*, *Cd* and *Hg* along with '*d*' block elements is not proper because
 (a) Their '*d*' orbitals are completely filled
 (b) Their '*d*' orbitals are empty
 (c) They do not form complex compounds
 (d) They do not form coloured compounds
140. Which of the following is the weakest reducing agent
 (a) *Zn* (b) *Cu*
 (c) H_2 (d) *Li*
141. The decrease in atomic volume from *Cr* to *Cu* is very negligible because
 (a) Increase in nuclear charge
 (b) Screening effect
 (c) Unpaired electrons of *Cr*
 (d) None
142. The heaviest atom amongst the following is [Kurukshetra CEE 1998]
 (a) Uranium (b) Radium
 (c) Lead (d) Mercury
143. Thallium shows different oxidation states because [Kurukshetra CEE 1998]
 (a) It is a transition metal
 (b) Of inert-pair effect
 (c) Of its high reactivity
 (d) Of its amphoteric character
144. The test of ozone O_3 can be done by [AFMC 1997]
 (a) *Ag* (b) *Hg*
 (c) *Au* (d) *Cu*
145. Which of the following set of elements does not belong to transitional elements set [EAMCET 1998]
 (a) *Fe*, *Co*, *Ni* (b) *Cu*, *Ag*, *Au*
 (c) *Ti*, *Zr*, *Hf* (d) *Ga*, *In*, *Tl*
146. The transition metals mostly are [MP PMT 2000; Kerala (Med.) 2002]
 (a) Diamagnetic
 (b) Paramagnetic
 (c) Neither diamagnetic nor paramagnetic
 (d) Both diamagnetic and paramagnetic
147. The correct statement in respect of *d*-block elements [MP PMT 2000, 02]
 (a) They are all metals
 (b) They show variable valency
 (c) They form coloured ions and complex salts
 (d) All above statements are correct
148. Which one of the following is an example of non-typical transition elements [MP PMT 2002]
 (a) *Li*, *K*, *Na* (b) *Be*, *Al*, *Pb*
 (c) *Zn*, *Cd*, *Hg* (d) *Ba*, *Ca*, *Sr*
149. Which one is wrong in the following statements [Kurukshetra CET 2002]
 (a) Gold is considered to be the king of metals
 (b) Gold is soluble in mercury
 (c) Copper is added to gold to make it hard
 (d) None of these
150. The number of unpaired electrons in Cr^{3+} ion is

- [Kurukshestra CET 2002]
- (a) 3 (b) 5
(c) 4 (d) 1
151. The metal ion which does not form coloured compound is [Kurukshestra CET 2002]
- (a) Chromium (b) Manganese
(c) Zinc (d) Iron
152. Super alloys are usually [Kurukshestra CET 2002]
- (a) Iron based
(b) Nickel based
(c) Cobalt based
(d) Based on all of these
153. The transitional metal which shows oxidation state from +2 to +7 belong to group
- (a) VII B (b) VI B
(c) II B (d) III B
154. Which of the following may be colourless [RPMT 1997; RPET/PMT 1999]
- (a) Cr^{+3} (b) Cu^{+}
(c) Fe^{+3} (d) Cu^{2+}
155. Which of the following ions is paramagnetic
- (a) Cu^{+} (b) Zn^{+2}
(c) Ti^{+3} (d) Ti^{+4}
156. Which of the following metals absorbs hydrogen
- (a) K (b) Al
(c) Zn (d) Pd
157. Which of the following ions is coloured [BHU 1997]
- (a) Cu^{+} (b) Cu^{2+}
(c) Ti^{4+} (d) V^{5+}
158. The metal present in B_{12} is [BHU 1997]
- (a) Magnesium (b) Iron
(c) Cobalt (d) Manganese
159. Which metal does not give the following reaction $M + \text{water or steam} \rightarrow \text{oxide} + H_2$ [Pb. PMT 2001]
- (a) Mercury (b) Iron
(c) Sodium (d) Magnesium
160. Ionisation potential values of d -block elements as compared to ionization potential value of f -block elements are
- (a) Higher (b) Equal
(c) lower (d) All of these
161. Which one of the following properties is not of transition elements [MP PET 1999; CPMT 2002]
- (a) Colour (b) Paramagnetism
(c) Fixed valency (d) None of the above
162. In which of the following, tendency towards formation of coloured ions is maximum [MP PET 1999]
- (a) s -block elements (b) d -block elements
(c) p -block elements (d) f -block elements
163. The electronic configuration $1s^2, 2s^2 p^6, 3s^2 p^6 d^6$ corresponds to [MP PET 1994]
- (a) Mn^{2+} (b) Fe^{2+}
(c) Co (d) Ge
164. Which of the following statements is not correct about the electronic configuration of gaseous chromium atom [MP PET 1994]
- (a) It has 5 electrons in $3d$ and one electron in $4s$ orbitals
(b) The principal quantum numbers of its valence electrons are 3 and 4
(c) It has 6 electrons in $3d$ orbital
(d) Its valance electrons have quantum number ' l ' 0 and 2
165. Zn and Hg belong to the same group, they differ in many of their properties. The property that is shared by both is [Pb. PMT 1998]
- (a) They form oxide readily
(b) They react with steam readily
(c) They react with hot concentrated sulphuric acid
(d) They react with hot sodium hydroxide
166. Which of the following ionic species will impart colour to an aqueous solution [CBSE PMT 1998; BHU 2001]
- (a) Ti^{4+} (b) Cu^{+}
(c) Zn^{2+} (d) Cr^{3+}
167. The number of electrons in the outermost shell of the $3d$ -transition elements generally remains [MP PMT 1997]
- (a) $(n-1)d^n$ (b) nd^n
(c) ns^2 (d) $(n-1)s^2$
168. The $3d$ -elements show variable oxidation states. What is the maximum oxidation state shown by the element Mn [MP PMT 1997; JIPMER 2002]
- (a) +4 (b) +5
(c) +6 (d) +7
169. Which of the following ions gives coloured solution [MP PET 1995]
- (a) Cu^{+} (b) Zn^{++}
(c) Ag^{+} (d) Fe^{++}
170. Which metal represents more than one oxidation state [CPMT 1990]
- (a) Al (b) Na
(c) Mg (d) Fe
171. A reduction in atomic size with increase in atomic number is a characteristic of elements of [AIEEE 2003]
- (a) High atomic masses (b) d -block
(c) f -block (d) Radioactive series
172. Which one of the following characteristics of the transition metals is associated with their catalytic activity [CBSE PMT 2003]
- (a) Variable oxidation states
(b) High enthalpy of atomization
(c) Paramagnetic behaviour
(d) Colour of hydrated ions
173. The number of oxidation states of manganese is
- (a) 4 (b) 6
(c) 3 (d) 8
174. Which of the following transitional metal has lowest density
- (a) Sc (b) Ti
(c) V (d) Cr
175. Which of the following transitional metal has lowest boiling point (B.P.)

- (a) Zn (b) Sc
(c) Ti (d) V
176. Which of the following electronic configuration is that of a transitional element [NCERT 1983; CPMT 1989, 97; MP PET/PMT 1997; AIIMS 2000; MP PMT 2002]
- (a) $1s^2, 2s^2 p^6, 3s^2 p^6 d^{10}, 4s^2 p^6$
(b) $1s^2, 2s^2 p^6, 3s^2 p^6 d^{10}, 4s^2 p^1$
(c) $1s^2, 2s^2 p^6, 3s^2 p^6 d^2, 4s^2$
(d) $1s^2, 2s^2 p^6, 3s^2 p^6, 4s^2$
177. Fe has been placed in the eighth group, the number of electrons in the outermost orbit is
(a) 1 (b) 2
(c) 3 (d) 4
178. Due to covalent bonding, the transitional metals are
(a) Lustrous (b) Conductor
(c) Hard and brittle (d) Ductile
179. The magnetic moment of a metal ion of first transition series is $2.83 BM$. Therefore it will have unpaired electrons
(a) 6 (b) 4
(c) 3 (d) 2
180. Which of the following pair of ions may exhibit same colour
(a) Cr^{+++} and Fe^{++}
(b) Ti^{+++} and V^{++}
(c) Fe^{+++} and Mn^{++}
(d) Cu^+ and Ni^{++}
181. The number of incomplete orbitals in inner transition element is
(a) 2 (b) 3
(c) 4 (d) 1
182. Most common oxidation states of Cs (cesium) are [AIEEE 2002]
(a) +2, +3 (b) +2, +4
(c) +3, +4 (d) +3, +5
183. The $3d$ elements show variable oxidation states because the energies of the following sets of orbitals are almost similar
(a) $ns, (n-1)d$ (b) ns, nd
(c) $(n-1)s, nd$ (d) $np, (n-1)d$
184. Which of the following $3d$ bivalent metal ions has the smallest number of unpaired d electrons
(a) $3d^6$ (b) $3d^7$
(c) $3d^8$ (d) $3d^9$
185. The $3d$ metal ions form coloured compounds because the energy corresponding to the following lies in the visible range of electromagnetic spectrum
(a) Free energy change of complex formation by $3d$ metal ions
(b) $d-d$ transitions of $3d$ electrons
(c) Heat of hydration of $3d$ metal ions
(d) Ionisation energy of $3d$ metal ions
186. The oxidation number of iron in potassium ferrocyanide is
(a) +2 (b) +3
(c) +4 (d) Zero
187. In transition elements, the orbitals partially filled by electrons are [DPMT 1984; AIIMS 1984; MP PMT 1997; AIIMS 2000; MP PMT 2002]
(a) s -orbitals (b) p -orbitals
(c) d -orbitals (d) f -orbitals
188. Number of unpaired electrons in Mn^{2+} is [CPMT 1997; Pb. PET/PMT 1999]
(a) 3 (b) 5
(c) 4 (d) 1
189. Mercury is the only metal which is liquid at $0^\circ C$. This is due to its [CBSE PMT 1997; AIIMS 1998; MP PMT 1999]
(a) Very high ionisation energy and weak metallic bond
(b) Low ionisation potential
(c) High atomic weight
(d) High vapour pressure
190. Essential constituent of an amalgam is [DPMT 1982; CPMT 1973, 77, 78, 89]
(a) Iron (b) An alkali metal
(c) Silver (d) Mercury
191. Mercury is transported in metal containers made of [DPMT 1982; CPMT 1973]
(a) Silver (b) Lead
(c) Iron (d) Aluminium
192. The electroplating of chromium is undertaken because [MP PMT 1994]
(a) Electrolysis of chromium is easier
(b) Chromium can form alloys with other metals
(c) Chromium gives protective and decorative coating to the base metal
(d) Of the high reactivity of metallic chromium [Pb. PMT 2001]
193. An element having the electronic configuration $[Ar]3d^2 4s^2$ belongs to [MP PMT 1993]
(a) s -block elements (b) p -block elements
(c) d -block elements (d) f -block elements
194. Which one of the following is not a transition metal [MP PMT 1999]
(a) Chromium (b) Titanium
(c) Lead (d) Tungsten
195. The atomic number of an element is 22. The highest oxidation state exhibited by it in its compounds is [MP PMT 1996]
(a) 1 (b) 2
(c) 3 (d) 4
196. d -block elements form
(a) Ionic compounds
(b) Covalent compounds
(c) Ionic and covalent compounds
(d) Only complex compounds
197. The transition metals have a less tendency to form ions due to [Bihar CEE 1995]
(a) High ionisation energy
(b) Low heat of hydration of ion
(c) High heat of sublimation
(d) All of these
198. The electronic configuration of Ag atom is [CPMT 1984]
(a) $[Kr]3d^{10} 4s^1$ (b) $[Xe]4f^{14} d^{10} 6s^1$

- (c) $[Kr]4d^{10}5s^1$ (d) $[Kr]4d^95s^2$
199. Most powerful oxidizing property of manganese is shown by which of the following oxidation state [MP PET 2003]
 (a) $Mn(+7)$ (b) $Mn(+2)$
 (c) $Mn(+4)$ (d) $Mn(+5)$
200. Which one of the following ions is colourless [MP PET 1999; RPET/PMT 1999]
 (a) Cu^+ (b) Co^{2+}
 (c) Ni^{2+} (d) Fe^{3+}
201. The atomic radii of the elements are almost same of which series
 (a) $Fe - Co - Ni$ (b) $Na - K - Rb$
 (c) $F - Cl - Br$ (d) $Li - Be - B$
202. In human body if necessary, the plate, screw or wire used for surgery are made up of
 (a) Ni (b) Au
 (c) Pt (d) Ta
203. Manganese is related to which block of periodic table [MP PMT 2003]
 (a) s-block (b) p-block
 (c) d-block (d) f-block
204. A hard and resistant metal (alloy) generally used in tip of nib of fountain pen is [BHU 1982]
 (a) $Os.Ir$ (b) $Pt.Cr$
 (c) $V.Fe$ (d) $Fe.Cr$
205. Chloride of which of the following elements will be coloured [MP PMT 1999]
 (a) Silver (b) Mercury
 (c) Zinc (d) Cobalt
206. Which of the following ions has the highest magnetic moment [JIPMER 1997; AIEEE 2002]
 (a) Ti^{3+} (b) Sc^{3+}
 (c) Mn^{2+} (d) Zn^{2+}
207. Cerium ($Z = 58$) is an important member of the lanthanoids. Which of the following statements about cerium is incorrect [AIEEE 2004] 217. Which of the following is the correct sequence of atomic weights of given elements [Pb. CET 2002]
 (a) The +4 oxidation state of cerium is not known in solutions
 (b) The +3 oxidation state of cerium is more stable than the +4 oxidation state
 (c) The common oxidation states of cerium are +3 and +4
 (d) Cerium (IV) acts as an oxidizing agent
 (a) $Fe > Co > Ni$ (b) $Ni > Co > Fe$
 (c) $Co > Ni > Fe$ (d) $Fe > Ni > Co$
208. Of the following outer electronic configurations of atoms, the highest oxidation state is achieved by which one of them [AIEEE 2004]
 (a) $(n-1)d^3ns^2$ (b) $(n-1)d^5ns^1$
 (c) $(n-1)d^8ns^2$ (d) $(n-1)d^5ns^2$
209. Among the following series of transition metal ions, the one where all metals ions have $3d^2$ electronic configuration is [CBSE PMT 2004]
 (a) $Ti^{4+}, V^{3+}, Cr^{2+}, Mn^{3+}$ (b) $Ti^{2+}, V^{3+}, Cr^{4+}, Mn^{5+}$
 (c) $Ti^{3+}, V^{2+}, Cr^{3+}, Mn^{4+}$ (d) $Ti^+, V^{4+}, Cr^{6+}, Mn^{7+}$
210. Lanthanoids are [CBSE PMT 2004]
 (a) 14 elements in the sixth period (atomic no. = 58 to 71) that are filling 4f sublevel
 (b) 14 elements in the seventh period (atomic no. = 58 to 71) that are filling 4f sublevel
- (c) 14 elements in the sixth period (atomic no. = 90 to 103) that are filling 4f sublevel
 (d) 14 elements in the seventh period (atomic no. = 90 to 103) that are filling 4f sublevel
211. Which of the following metals make the most efficient catalyst [BHU 1995]
 (a) Transition (b) Alkali
 (c) Alkaline earth (d) Coloured metals
212. Lanthanides and actinides resemble in [AFMC 2004]
 (a) Electronic configuration (b) Oxidation state
 (c) Ionization energy (d) Formation of complexes
213. The lanthanide contraction relates to [Kerala PMT 2004]
 (a) Atomic radii
 (b) Atomic as well as M^{3+} radii
 (c) Valence electrons
 (d) Oxidation states
 (e) Ionisation energy
214. Which of the following species is expected to show the highest magnetic moment? (At. Nos.: $Cr=24, Mn=25, Co=27, Ni=28, Cu=29$) [Kerala PMT 2004]
 (a) Cr^{2+} (b) Mn^{2+}
 (c) Cu^{2+} (d) Co^{2+}
 (e) Ni^{2+}
215. Which one belongs to 3d-transition series [MP PMT 2004]
 (a) Copper (b) Gold
 (c) Cobalt (d) Silver
216. Which one of the following organisation's iron and steel plant was built to use charcoal as a source of power, to start with, but later switched over to hydroelectricity [AIIMS 2004]
 (a) The Tata Iron and Steel Company
 (b) The Indian Iron and Steel Company
 (c) Mysore Iron and Steel Limited
 (d) Hindustan Steel Limited
218. Which of the following element has maximum first ionisation potential [Pb. CET 2002]
 (a) V (b) Ti
 (c) Cr (d) Mn
219. A metal M having electronic configuration $M - 1s^2 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^1$ [DCE 2002]
 (a) s-block element (b) d-block element
 (c) p-block element (d) None of these
220. Identify the transition element [MH CET 2003]
 (a) $1s^2, 2s^2 2p^6, 3s^2, 3p^6, 4s^2$
 (b) $1s^2, 2s^2 2p^6, 3s^2, 3p^6 3d^2, 4s^2$
 (c) $1s^2, 2s^2 2p^6, 3s^2, 3p^6 3d^{10}, 4s^2 4p^2$
 (d) $1s^2, 2s^2 2p^6, 3s^2, 3p^6 3d^{10}, 4s^2 4p^1$
221. What is the name of element with atomic number 105 [CPMT 2004]
 (a) Kurchatovium (b) Dubnium
 (c) Nobelium (d) Holmium

222. Electrons in a paramagnetic compound are [UPSEAT 2004]
 (a) Shared (b) Unpaired
 (c) Donated (d) Paired
223. Which of the following pairs involves isoelectronic ions [UPSEAT 2004]
 (a) Mn^{3+} and Fe^{2+} (b) Mn^{2+} and Fe^{3+}
 (c) Cr^{3+} and Mn^{2+} (d) Fe^{2+} and Co^{2+}
224. Which of the following is paramagnetic [Pb. CET 2000]
 (a) Ni^{++} (b) Cu^{+}
 (c) Zn^{++} (d) Sc^{+++}
225. The electronic configuration of chromium is [BHU 2005; Pb. CET 2000]
 (a) $[Ne]3s^2 3p^6 3d^4, 4s^2$ (b) $[Ne]3s^2 3p^6 3d^5, 4s^1$
 (c) $[Ne]3s^2 3p^6 3d^6, 4s^1$ (d) $[Ne]3s^2 3p^5 3d^5, 4s^2$
226. Electronic configuration of $Cu(Z = 29)$ is [Pb. CET 2001]
 (a) $[Ar]3d^9 4s^2$ (b) $[Ar]3d^{10} 4s^1$
 (c) $[Ar]3d^5 4s^2$ (d) $[Ar]3d^6 4s^2$
227. $Ce-58$ is a member of [Pb. CET 2002]
 (a) s -block (b) p -block
 (c) d -block (d) f -block
228. How many unpaired electrons are there in Ni^{2+} [MP PET 2004]
 (a) 2 (b) 4
 (c) 5 (d) 0
229. The main reason for larger number of oxidation states exhibited by the actinoids than the corresponding lanthanoids is [CBSE PMT 2005]
 (a) Lesser energy difference between $5f$ and $6d$ orbitals than between $4f$ and $5d$ orbitals
 (b) Larger atomic size of actinoids than the lanthanoids
 (c) More energy difference between $5f$ and $6d$ orbitals than between $4f$ and $5d$ orbitals
 (d) Greater reactive nature of the actinoids than the lanthanoids
230. Four successive members of the first row transition elements are listed below with their atomic numbers. Which one of them is expected to have the highest third ionization enthalpy [CBSE PMT 2005]
 (a) Vanadium ($Z = 23$) (b) Chromium ($Z = 24$)
 (c) Iron ($Z = 26$) (d) Manganese ($Z = 25$)
231. The aqueous solution containing which one of the following ions will be colourless [CBSE PMT 2000, 05]
 (a) Sc^{3+} (b) Fe^{2+}
 (c) Ti^{3+} (d) Mn^{2+}
 (Atomic number $Sc = 21$, $Fe = 26$, $Ti = 22$, $Mn = 25$)
232. Which of the following trivalent ion has the largest atomic radii in the lanthanide series [BHU 2002]
 (a) La (b) Ce
 (c) Pm (d) Lu
233. Which of the following does not have valence electron in $3d$ -subshell [AIIMS 2002]
 (a) $Fe(III)$ (b) $Mn(II)$
 (c) $Cr(I)$ (d) $P(0)$
234. Among the following pairs of ions, the lower oxidation state in aqueous solution is more stable than the other in [AIIMS 2005]
 (a) Tl^+ , Tl^{3+} (b) Cu^+ , Cu^{2+}
 (c) Cr^{2+} , Cr^{3+} (d) V^{2+} , VO^{2+}
235. The lanthanide contraction is responsible for the fact that [AIEEE 2005]
 (a) Zr and Y have about the same radius
 (b) Zr and Nb have similar oxidation state
 (c) Zr and Hf have about the same radius
 (d) Zr and Zn have the same oxidation state
236. Which of the following factors may be regarded as the main cause of lanthanide contraction [AIEEE 2005]
 (a) Poor shielding of one of $4f$ electron by another in the subshell
 (b) Effective shielding of one of $4f$ electrons by another in the subshell
 (c) Poorer shielding of $5d$ electrons by $4f$ electrons
 (d) Greater shielding of $5d$ electron by $4f$ electrons
237. Which of the following have maximum number of unpaired electrons [BHU 2005]
 (a) Fe (b) Fe
 (c) Co (d) Co
238. Transition metals show paramagnetism [BHU 2005]
 (a) Due to characteristic configuration
 (b) High lattice energy
 (c) Due to variable oxidation states
 (d) Due to unpaired electrons
239. Which of the following pairs of elements cannot form an alloy [KCET 2005]
 (a) Zn, Cu (b) Fe, Hg
 (c) Fe, C (d) Hg, Na
240. Which belongs to the actinides series [J & K 2005]
 (a) Ce (b) Cf
 (c) Ca (d) Cs
241. Effective magnetic moment of Sc^{+3} ion is [Kerala CET 2005]
 (a) 1.73 (b) 0
 (c) 5.92 (d) 2.83
 (e) 3.87



Compounds of Transitional elements

1. Potassium permanganate acts as an oxidant in neutral, alkaline as well as acidic media. The final products obtained from it in the three conditions are, respectively
[MP PMT 1997]
(a) MnO_2, MnO_2, Mn^{2+} (b) $MnO_4^{2-}, Mn^{3+}, Mn^{2+}$
(c) $MnO_2, MnO_4^{2-}, Mn^{3+}$ (d) MnO, MnO_4, Mn^{2+}
 2. In acidic medium one mole of MnO_4^- accepts how many moles of electrons in a redox process ? [MP PET/PMT 1998]
(a) 1 (b) 3
(c) 5 (d) 6
 3. In acidic medium potassium dichromate acts as an oxidant according to the equation,
 $Cr_2O_7^{2-} + 14H^+ + 6e^- \rightarrow 2Cr^{3+} + 7H_2O$. What is the equivalent weight of $K_2Cr_2O_7$? (mol. Wt. = M)
[MP PET/PMT 1998]
(a) M (b) $M/2$
(c) $M/3$ (d) $M/6$
 4. The correct formula of permanganic acid is [MP PET 1999]
(a) $HMnO_4$ (b) $HMnO_5$
(c) H_2MnO_4 (d) H_2MnO_3
 5. Acidified potassium dichromate is treated with hydrogen sulphide. In the reaction, the oxidation number of chromium
[MP PET 1996]
(a) Increases from +3 to +6
(b) Decreases from +6 to +3
(c) Remains unchanged
(d) Decreases from +6 to +2
 6. When $KMnO_4$ reacts with acidified $FeSO_4$ [MP PET 1996]
(a) Only $FeSO_4$ is oxidised
(b) Only $KMnO_4$ is oxidised
(c) $FeSO_4$ is oxidised $KMnO_4$ and is reduced
(d) None of these
 7. When calomel reacts with NH_4OH , we get
[CBSE PMT 1996]
(a) $HgNH_2Cl$ (b) $NH_2 - Hg - Hg - Cl$
(c) Hg_2O (d) HgO
 8. $AgCl$ dissolves in a solution of NH_3 but not in water because [MP PMT 1984, 86]
(a) NH_3 is a better solvent than H_2O
(b) Ag^+ forms a complex ion with NH_3
(c) NH_3 is a stronger base than H_2O
(d) The dipole moment of water is higher than NH_3
 9. In solid $CuSO_4 \cdot 5H_2O$ copper is co-ordinated to
[MP PET 1985, 86]
(a) Five water molecules (b) Four water molecules
(c) One sulphate anion (d) One water molecule
 10. A white powder soluble in NH_4OH but insoluble in water is [AFMC 1987, 74]
(a) $BaSO_4$ (b) $CuSO_4$
(c) $PbSO_4$ (d) $AgCl$
 11. Verdigris is [BHU 1987]
(a) Basic copper acetate (b) Basic lead acetate
(c) Basic lead (d) None of these
 12. Number of moles of $K_2Cr_2O_7$ reduced by one mole of Sn^{2+} ions is [KCET 1996]
(a) 1/3 (b) 3
(c) 1.6 (d) 6
 13. Which one of the following is reduced by hydrogen peroxide in acid medium [EAMCET 1997]
(a) Potassium permanganate
(b) Potassium iodide
(c) Ferrous sulphate
(d) Potassium ferrocyanide
 14. Which oxide of manganese is amphoteric [AFMC 1995]
(a) MnO_2 (b) Mn_2O_3
(c) Mn_2O_7 (d) MnO
 15. Which one of the following oxides is ionic [IIT-JEE 1995]
(a) MnO (b) Mn_2O_7
(c) CrO_3 (d) P_2O_5
 16. Correct formula of calomel is [CPMT 1994; AFMC 1998]
(a) Hg_2Cl_2 (b) $HgCl_2$
(c) $HgCl_2 \cdot H_2O$ (d) $HgSO_4$
 17. One of the important use of ferrous sulphate is in the
(a) Manufacture of blue black ink
(b) Manufacture of chalks
(c) Preparation of hydrogen sulphide
(d) Preparation of anhydrous ferric chloride
 18. Copper sulphate is not used
(a) In electrotyping
(b) In dyeing and calicoprinting
(c) In detecting water
(d) As fertilizer
 19. Blue vitriol is [AFMC 1992]
(a) $CuSO_4$ (b) $CuSO_4 \cdot 5H_2O$
(c) Cu_2SO_4 (d) $CuSO_4 \cdot H_2O$
 20. A solution of copper sulphate may be kept safely in the container made up of
(a) Fe (b) Ag
(c) Zn (d) Al
 21. Silver nitrate produces a black stain on skin due to
(a) Being a strong reducing agent
(b) Its corrosive action
(c) Formation of complex compound
(d) Its reduction to metallic silver
 22. When hypo solution is added to cupric sulphate solution, the blue colour of the latter is discharged, due to formation of
(a) CuS_2O_3 (b) $Na_2S_4O_6$
(c) $NaCuS_2O_3$ (d) Cu_2O
 23. Metal oxides which decomposes on heating is [MNR 1984; UPSEAT 1999]
(a) ZnO (b) Al_2O_3
(c) CuO (d) Na_2O
(e) HgO
- Anhydrous sample of ferric chloride is prepared by heating
(a) $Fe + HCl$ (b) $Fe + Cl_2$
(c) $FeCl_2 + Cl_2$ (d) Hydrated ferric chloride

25. Light green crystals of ferrous sulphate lose water molecule and turn brown on exposure to air. This is due to its oxidation to
 (a) Fe_2O_3 (b) $Fe_2O_3 \cdot H_2O$
 (c) $Fe(OH)SO_4$ (d) $Fe_2O_3 + FeO$
26. In alkaline condition $KMnO_4$, reacts as follows :
 $2KMnO_4 + 2KOH \rightarrow 2K_2MnO_4 + H_2O + O$
 Therefore its equivalent weight will be
 [NCERT 1974; CPMT 1977; DCE 2002]
 (a) 31.5 (b) 52.7
 (c) 72.0 (d) 158.0
27. Equivalent weight of $KMnO_4$ acting as an oxidant in acidic medium is equal to [CPMT 1990; MP PMT 1999]
 (a) Molecular weight of $KMnO_4$
 (b) $\frac{1}{2} \times$ Molecular weight of $KMnO_4$
 (c) $\frac{1}{3} \times$ Molecular weight of $KMnO_4$
 (d) $\frac{1}{5} \times$ Molecular weight of $KMnO_4$
28. In which of the following ionic radii of chromium would be smallest [MP PET 1994]
 (a) K_2CrO_4 (b) CrO_2
 (c) $CrCl_3$ (d) CrF_2
29. $CoO \cdot Al_2O_3$ is called
 (a) Cobalt aluminate (b) Thenard's blue
 (c) Both (a) and (b) (d) None of these
30. $ZnO \cdot CoO$ is called
 (a) Cobalt zincate (b) Rinman's green
 (c) Both (a) and (b) (d) None of these
31. $FeSO_4 \cdot (NH_4)_2SO_4 \cdot 6H_2O$ is called [Bihar CEE 1995]
 (a) Mohr's salt (b) Green salt
 (c) Alum (d) Glauber's salt
32. Molybdenum compounds are used in
 (a) Dye industry (b) For colouring leather
 (c) For colouring rubber (d) All of these
33. When copper turnings and concentrated HCl is heated with copper sulphate the compound formed is [CPMT 1984]
 (a) Cupric chloride (b) Cuprous chloride
 (c) Copper sulphate (d) SO_2
34. The compound of copper which turns green on keeping in air is [CPMT 1984]
 (a) Copper sulphate (b) Copper nitrate
 (c) Cupric chloride (d) Cuprous chloride
35. Cu_2Cl_2 with HCl in presence of oxidising agents gives [CPMT 1984]
 (a) $CuCl_2$ (b) H_2CuCl_2
 (c) Hydrogen gas (d) Chlorine gas
36. $K_2Cr_2O_7$ on heating with aqueous $NaOH$ gives [CBSE PMT 1997]
 (a) CrO_4^{2-} (b) $Cr(OH)_3$
 (c) $Cr_2O_7^{2-}$ (d) $Cr(OH)_2$
37. $KMnO_4$ reacts with oxalic acid according to the equation :
 $2MnO_4^- + 5C_2O_4^{2-} + 16H^+ \rightarrow 2Mn^{2+} + 10CO_2 + 8H_2O$
 Here 20 ml of 0.1 M $KMnO_4$ is equivalent to [CBSE PMT 1996]
 (a) 20ml of 0.5 M $C_2H_2O_4$
 (b) 50ml of 0.1 M $C_2H_2O_4$
 (c) 50ml of 0.5 M $C_2H_2O_4$
 (d) 20ml of 0.1 M $C_2H_2O_4$
38. The equivalent weight of potassium permanganate for acid solution is [MP PET 1999]
 (a) 158 (b) 31.6
 (c) 52.16 (d) 79
39. Which statement is not correct [MP PET 1999]
 (a) Potassium permanganate is a powerful oxidising substance
 (b) Potassium permanganate is a weaker oxidising substance than potassium dichromate
 (c) Potassium permanganate is a stronger oxidising substance than potassium dichromate
 (d) Potassium dichromate oxidises a secondary alcohol into a ketone
40. The formula of corrosive sublimate is [CPMT 1997]
 (a) $HgCl_2$ (b) Hg_2Cl_2
 (c) Hg_2O (d) Hg
41. Which is mild oxidising agent [AFMC 1971]
 (a) Ag_2O (b) $KMnO_4$
 (c) $K_2Cr_2O_7$ (d) Cl_2
42. The equivalent weight of $K_2Cr_2O_7$ in acidic medium
 (a) 294 (b) 298
 (c) 49 (d) 50
43. The transitional element of III-B group is isomorphic with $Al_2(SO_4)_3$. Therefore the compound will be
 (a) Purple (b) Blue
 (c) White (d) Uncertain
44. A copper salt is isomorphic with $ZnSO_4$, the salt will be
 (a) Paramagnetic (b) Diamagnetic
 (c) Ferromagnetic (d) None
45. V_2O_5 is useful as catalyst in
 (a) Manufacture of H_2SO_4
 (b) Manufacture of HNO_3
 (c) Manufacture of Na_2CO_3
 (d) It is not a catalyst
46. $KMnO_4$ in basic medium is reduced to [Orissa JEE 2005]
 (a) K_2MnO_4 (b) MnO_2
 (c) $Mn(OH)_2$ (d) Mn^{2+}
47. When $KMnO_4$ is reduced with oxalic acid in acidic solution, the oxidation number of Mn changes from [CPMT 1989]
 (a) 7 to 4 (b) 6 to 4
 (c) 7 to 2 (d) 4 to 2
48. Nessler's reagent is [MP PET 1991; MP PMT 1993; AFMC 2001]
 (a) K_2HgI_4 (b) $K_2HgI_4 + KOH$
 (c) $K_2HgI_2 + KOH$ (d) $K_2HgI_4 + Hg$
49. When ammonium dichromate is heated, the gas formed is [MP PMT 1993; IIT-JEE 1999]
 (a) N_2 (b) O_2
 (c) H_2 (d) NH_3
50. Acidified potassium dichromate on reacting with a sulphite is reduced to

- (a) CrO_2Cl_2 (b) CrO_4^{2-}
(c) Cr^{3+} (d) Cr^{2+}
51. The product of oxidation of I^- ion by MnO_4^- in alkaline medium is
(a) I_2 (b) IO_3^-
(c) IO_4^- (d) I_3^-
52. Identify the statement which is not correct regarding copper sulphate [UPSEAT 2000, 01]
(a) It reacts with KI to give iodine
(b) It reacts with KCl to give Cu_2Cl_2
(c) It reacts with $NaOH$ and glucose to give Cu_2O
(d) It give CuO on strong heating in air
53. Acidified potassium permanganate solution is decolourised by [MNR 1984]
(a) Bleaching powder (b) White vitriol
(c) Mohr's salt (d) Microcosmic salt
(e) Laughing gas
54. Which of the following oxides is white but becomes yellow on heating [MP PET 1995]
(a) AgO (b) Ag_2O
(c) FeO (d) ZnO
55. Amalgams are
(a) Highly coloured alloys
(b) Always solid
(c) Alloys which contain mercury as one of the contents
(d) Alloys which have great resistance to abrasion
56. In photography sodium thiosulphate is used as [DPMT 2005]
(a) Complexing agent (b) Oxidising agent
(c) Reducing agent (d) None of these
57. The substance that sublimes on heating [EAMCET 1978, 82; MP PMT 1999]
(a) Magnesium chloride (b) Silver chloride
(c) Mercurous chloride (d) Sodium chloride
58. $K_3[Fe(CN)_6]$ is called
(a) Potassium ferricyanide
(b) Red prussiate of potash
(c) Potassium hexacyanoferrate (III)
(d) All of these
59. Which of the following will show increase in weight when kept in magnetic field
(a) TiO_2 (b) $Fe_2(SO_4)_3$
(c) $KMnO_4$ (d) $ScCl_3$
60. Amongst TiF_6^{2-} , CoF_6^{3-} , Cu_2Cl_2 and $NiCl_4^{2-}$ (Atomic number $Ti = 22$, $Co = 27$, $Cu = 29$, $Ni = 28$). The colourless species are [CBSE PMT 1995]
(a) CoF_6^{3-} and $NiCl_4^{2-}$ (b) TiF_6^{2-} and CoF_6^{3-}
(c) Cu_2Cl_2 and $NiCl_4^{2-}$ (d) TiF_6^{2-} and Cu_2Cl_2
61. Which of the following imparts green colour to the glass [CPMT 1993]
(a) Cu_2O (b) CdS
(c) MnO_2 (d) Cr_2O_3
62. On the heating copper nitrate strongly, is finally obtained [CPMT 1971, 74, 78]
(a) Copper (b) Copper oxide
(c) Copper nitrate (d) Copper nitride
63. On adding KI to a solution of copper sulphate [CPMT 1973; NCERT 1977; MP PMT 1989]
(a) Cupric oxide is precipitated
(b) Metallic copper is precipitated
(c) Cuprous iodide is precipitated with liberation of iodine
(d) No change occurs
64. Which of the following statements is correct about equivalent weight of $KMnO_4$ [MP PET 1994]
(a) It is one third of its molecular weight in alkaline medium
(b) It is one fifth of its molecular weight in alkaline medium
(c) It is equal to its molecular weight in acidic medium
(d) It is one third of its molecular weight in acidic medium
65. The reaction of $K_2Cr_2O_7$ with $NaCl$ and conc. H_2SO_4 gives [MP PET 1995]
(a) $CrCl_3$ (b) $CrOCl_2$
(c) CrO_2Cl_2 (d) Cr_2O_3
66. Silver nitrate is supplied in coloured bottles because it is [CPMT 1985]
(a) Oxidised in air
(b) Decomposes in sunlight
(c) Explosive in sunlight
(d) Reactive towards air in sunlight
67. A nitrate when mixed with common salt gives a white precipitate which is soluble in dilute NH_4OH . It is the nitrate of [CPMT 1985]
(a) Copper (b) Mercury
(c) Silver (d) Gold
68. Which one of the following is lunar caustic [CPMT 1984]
(a) $AgNO_3$ (b) Cu_2Cl_2
(c) $CuCl_2$ (d) Hg_2Cl_2
69. Invar, an alloy of Fe and Ni is used in watches and meter scale, its characteristic property is [Kerala (Engg.) 2002]
(a) Small coefficient of expansion
(b) Resistance to corrosion
(c) Hardness and elasticity
(d) Magnetic nature
70. The extraction of nickel involves
(a) The formation of $Ni(CO)_4$
(b) The decomposition of $Ni(CO)_4$
(c) The formation and thermal decomposition of $Ni(CO)_4$
(d) The formation and catalytic decomposition of $Ni(CO)_4$
71. On adding excess of NH_3 solution to $CuSO_4$ solution, the dark blue colour is due to [CPMT 1990; AIIMS 1982; MP PMT 1989, 92; BHU 1996; JIPMER 1997]
(a) $[Cu(NH_3)_4]^{++}$ (b) $[Cu(NH_3)_2]^{++}$
(c) $[Cu(NH_3)]^+$ (d) None of the above
72. If M is the molecular weight of $KMnO_4$, its equivalent weight will be when it is converted into K_2MnO_4 [MP PET 1993]
(a) M (b) $M/3$
(c) $M/5$ (d) $M/7$
73. While writing the formula of ferrous oxide it is written as (FeO) , because it is
(a) Non-stoichiometric (b) Non-existent
(c) Paramagnetic (d) Ferromagnetic
74. Which of the following exhibit maximum oxidation state of vanadium

- (a) $VOCl_3$ (b) VCl_4
(c) VCl_3 (d) VCl_2
75. Prussian blue is due to the formation of [BHU 1980; CBSE PMT 1990; KCET 1992; MP PET 1995]
(a) $Fe_4[Fe(CN)_6]_3$ (b) $Fe_2[Fe(CN)_6]$
(c) $Fe_3[Fe(CN)_6]$ (d) $Fe[Fe(CN)_6]_3$
76. The Nessler's reagent contains [CPMT 1976, 88; NCERT 1987; MP PMT 1985; BHU 1996]
(a) Hg_2^{++} (b) Hg^{++}
(c) HgI_2^{-} (d) HgI_4^{-}
77. Formula of ferric sulphate is [AFMC 2003]
(a) $FeSO_4$ (b) $Fe(SO_4)_2$
(c) Fe_2SO_4 (d) $Fe_2(SO_4)_3$
78. When $CuSO_4$ is hydrated, then it becomes [AFMC 2003]
(a) Acidic (b) basic
(c) Neutral (d) Amphoteric
79. Silvering of mirror is done by [AFMC 2003]
(a) $AgNO_3$ (b) Ag_2O_3
(c) Fe_2O_3 (d) Al_2O_3
80. The colour of $K_2Cr_2O_7$ changes from red orange to lemon yellow on treatment with aqueous KOH because of [MP PMT 1994]
(a) The reduction of Cr^{VI} to Cr^{III}
(b) The formation of chromium hydroxide
(c) The conversion of dichromate to chromate
(d) The oxidation of potassium hydroxide to potassium peroxide
81. On heating pyrolusite with KOH in presence of air we get
(a) $KMnO_4$ (b) K_2MnO_4
(c) $Mn(OH)_2$ (d) Mn_3O_4
82. $Cu(CN)_4^{2-}$ is colourless as it absorbs light in
(a) Visible region (b) Ultraviolet region
(c) Infrared region (d) All above are wrong
83. Acidified solution of chromic acid on treatment with hydrogen peroxide yields [MP PET 1999; AFMC 2000]
(a) $CrO_3 + H_2O + O_2$ (b) $Cr_2O_3 + H_2O + O_2$
(c) $CrO_5 + H_2O$ (d) $H_2Cr_2O_7 + H_2O + O_2$
84. Which of the following metals corrodes readily in moist air [CPMT 1972, 82; CBSE PMT 1989]
(a) Gold (b) Silver
(c) Nickel (d) Iron
85. Which one of the following compounds is not coloured [AIIMS 1997]
(a) Na_2CuCl_4 (b) Na_2CdCl_4
(c) $K_4Fe(CN)_6$ (d) $K_3Fe(CN)_6$
86. Acidified $KMnO_4$ is decolorized by [AMU 1999]
(a) Br_2 (b) O_3
(c) HCl (d) HBr
87. Which of the following compound expected to be colourless
(a) ScO (b) V_2O_3
(c) $CuCN$ (d) $Cr_2(SO_4)_3$
88. Crystals of which pair are isomorphous [CPMT 1990]
(a) $ZnSO_4, SnSO_4$ (b) $MgSO_4, CaSO_4$
(c) $ZnSO_4, MgSO_4$ (d) $PbSO_4, NiSO_4$
89. On heating $Mn(OH)_2$ with PbO_2 and conc. HNO_3 pink colour is obtained due to the formation of [MP PET 1995]
(a) $KMnO_4$ (b) $HMnO_4$
(c) $Pb(MnO_4)_2$ (d) $PbMnO_4$
90. Which of the following is used as white pigment
(a) TiO_2 (b) V_2O_5
(c) CuO (d) HgO
91. Which metal oxide is used to make blue glass
(a) Fe_2O_3 (b) CoO
(c) Cu_2O (d) NiO
92. The chemical formula for certain pyrophosphate is $Ca_2P_2O_7$, the formula of its ferric pyrophosphate will be
(a) $Fe_2(P_2O_7)_3$ (b) $Fe_4(P_4O_{14})$
(c) $Fe_4(P_2O_7)_3$ (d) Fe_3PO_4
93. Which of the following compounds does not dissolve in ammonium hydroxide solution
(a) AgF (b) $AgBr$
(c) $AgCl$ (d) AgI
94. Which of the following is non-stoichiometric
(a) Fe_3O_4 (b) Fe_2O_3
(c) FeO (d) All the above
95. Ferrosilicon is used in steel industry as [Kerala (Med.) 2003]
(a) A flux (b) Scavenger of hydrogen
(c) A reducing agent (d) A cutting tool
(e) Alloying agent
96. In the reaction,
 $2KMnO_4 + 16HCl \rightarrow 5Cl_2 + 2MnCl_2 + 2KCl + 8H_2O$
the reduction product is [Kerala (Med.) 2003]
(a) Cl_2 (b) $MnCl_2$
(c) H_2O (d) KCl
97. Which one of the following statements is correct [AIEEE 2003]
(a) Manganese salts give violet borax bead test in the reducing flame
(b) From a mixed precipitate of $AgCl$ and AgI ammonia solution dissolves only $AgCl$
(c) Ferric ions give a deep green precipitate on adding potassium ferrocyanide solution
(d) On boiling a solution having K^+, Ca^{2+} and HCO_3^- ions we get a precipitate of $K_2Ca(CO_3)_2$
98. Collin's reagent is [RPMT 2002]
(a) MNO_2 / HCl (b) MNO_4 / C_5H_5N
(c) $K_2Cr_2O_7 / H_2SO_4$ (d) $Cr_2O_3 / 2C_5H_5N$
99. Which compound has coloured aqueous solution [RPMT 2002]
(a) $Zn(NO_3)_2$ (b) $LiNO_3$
(c) $Co(NO_3)_2$ (d) $Ba(NO_3)_2$
100. Zinc oxide when treated with $NaOH$ solution gives
(a) $Zn(OH)_2$ (b) $ZnCl_2$
(c) Na_2ZnO_2 (d) $Zn(OH)_3$

101. Which of the following compounds volatilises on heating [BHU 1998]
- (a) $MgCl_2$ (b) $HgCl_2$
(c) $CaCl_2$ (d) $FeCl_3$
102. Which of the following statement is not true
- (a) Colourless compounds of transition elements are paramagnetic
(b) Coloured compounds of transition elements are paramagnetic
(c) Colourless compounds of transition elements are diamagnetic
(d) Transition elements form the complex compounds
103. Calamine is a mineral, which is [MP PMT 2003]
- (a) $ZnCO_3$ (b) ZnS
(c) $ZnSO_4$ (d) ZnO
104. Super conductors are derived from compounds of [Kerala (Engg.) 2002]
- (a) *p*-block elements (b) Lanthanides
(c) Actinides (d) Transition elements
105. Manganese achieves its highest oxidation state in its compound [MP PET 1993, 2001; MP PMT 2004]
- (a) MnO_3 (b) Mn_3O_4
(c) $KMnO_4$ (d) K_2MnO_4
106. Which can be reduced to the metal by heating it in a stream of hydrogen [DPMT 2000]
- (a) Copper (II) oxide (b) Magnesium oxide
(c) Aluminium oxide (d) Calcium oxide
107. Which of the following is coloured
- (a) $ScCl_3$ (b) TiO_2
(c) $MnSO_4$ (d) $ZnSO_4$
108. Chrome green is
- (a) Chromium sulphate (b) Chromium chloride
(c) Chromium nitrate (d) Chromium oxide
109. The colour of $(NH_4)_2SO_4 \cdot Fe(SO_4)_3 \cdot 24H_2O$ is [BHU 1982; CPMT 1989]
- (a) White (b) Green
(c) Violet (d) Blue
110. Correct formula of potassium ferricyanide is [DPMT 1982, 83; CPMT 1974; AFMC 2005]
- (a) $K_4Fe(CN)_6$ (b) $K_3Fe(CN)_6$
(c) $K_3[Fe(CN)_3]$ (d) $K_3[Fe(CN)_4]$
111. The form of iron having the highest carbon content is [DPMT 2005]
- (a) Cast iron (b) Wrought iron
(c) Strain steel (d) Mild steel
112. Aqueous solution of ferric chloride is [MP PMT 1999]
- (a) Acidic (b) Basic
(c) Neutral (d) Amphoteric
113. In the reduction of dichromate by $Fe(II)$ the number of electrons involved per chromium atom is [Pb. PMT 2001]
- (a) 2 (b) 3
(c) 4 (d) 1
114. A group of acidic oxide is [MP PET 2003]
- (a) CrO_3, Mn_2O_7 (b) ZnO, Al_2O_3
(c) CaO, ZnO (d) Na_2O, Al_2O_3
115. Silver nitrate is mainly used [CPMT 1988, 93]
- (a) In photography (b) In model formation
(c) As reducing agent (d) As dehydrating agent
116. The correct order of magnetic moments (spin only values in B.M.) among is [AIEEE 2004]
- (a) $[Fe(CN)_6]^{4-} > [MnCl_4]^{2-} > [CoCl_4]^{2-}$
(b) $[MnCl_4]^{2-} > [Fe(CN)_6]^{4-} > [CoCl_4]^{2-}$
(c) $[MnCl_4]^{2-} > [CoCl_4]^{2-} > [Fe(CN)_6]^{4-}$
(d) $[Fe(CN)_6]^{4-} > [CoCl_4]^{2-} > [MnCl_4]^{2-}$
(Atomic nos. $Mn = 25, Fe = 26, Co = 27$)
117. Hybridization of $[Ni(CO)_4]$ is [Pb. CET 2000]
- (a) sp^3 (b) d^2sp^3
(c) sp^3d (d) sp^2
118. What is the oxidation number of iron in the compound $[Fe(H_2O)_5(NO)]SO_4$ [Pb. CET 2001]
- (a) +2 (b) +3
(c) +4 (d) +4
119. Which of the following metal gives hydrogen gas, when heated with hot concentrated alkali [Pb. CET 2002]
- (a) Cu (b) Ag
(c) Zn (d) Ni
120. When ferric oxide reacts with $NaOH$, the product formed is [Pb. CET 2002]
- (a) NaF (b) $FeCl_3$
(c) $Fe(OH)_3$ (d) $NaFeO_2$
121. The compound insoluble in water is [AIIMS 2004]
- (a) Mercurous nitrate (b) Mercuric nitrate
(c) Mercurous chloride (d) Mercurous perchlorate
122. Which is an amphoteric oxide [JEE Orissa 2004, 05]
- (a) ZnO (b) CaO
(c) BaO (d) SrO
123. What is the magnetic moment of $[FeF_6]^{3-}$ [JEE Orissa 2004]
- (a) 5.92 (b) 5.49
(c) 2.32 (d) 4
124. How H_2S is liberated in laboratory [JEE Orissa 2004]
- (a) $FeSO_4 + H_2SO_4$
(b) $FeS + dil. H_2SO_4$
(c) $FeS + conc. H_2SO_4$
(d) Elementary H_2 + elementary S
125. The spin magnetic moment of cobalt in the compound $Hg[Co(SCN)_4]$ is [IIT JEE Screening 2004]
- (a) $\sqrt{3}$ (b) $\sqrt{8}$
(c) $\sqrt{15}$ (d) $\sqrt{24}$
126. In which of these processes platinum is used as a catalyst [DCE 2004]
- (a) Oxidation of ammonia to form HNO_3
(b) Hardening of oils
(c) Production of synthetic rubber
(d) Synthesis of methanol
127. Iron is dropped in dil. HNO_3 , it gives [DCE 2004]
- (a) Ferric nitrate
(b) Ferric nitrate and NO_2
(c) Ferrous nitrate and ammonium nitrate

- (d) Ferrous nitrate and nitric oxide
128. CrO_3 dissolves in aqueous $NaOH$ to give [J & K 2005]
- (a) CrO_4^{2-} (b) $Cr(OH)_3^-$
(c) $Cr_2O_7^{2-}$ (d) $Cr(OH)_2$
129. KI and $CuSO_4$ solution when mixed, give [CPMT 2004; UPSEAT 2004]
- (a) $CuI_2 + K_2SO_4$ (b) $Cu_2I_2 + K_2SO_4$
(c) $K_2SO_4 + Cu_2I_2 + I_2$ (d) $K_2SO_4 + CuI_2 + I_2$
130. When Cu reacts with $AgNO_3$ solution, the reaction takes place is [CPMT 2004]
- (a) Oxidation of Cu (b) Reduction of Cu
(c) Oxidation of Ag (d) Reduction of NO_3^-
131. By annealing, steel [BHU 2004]
- (a) Becomes soft
(b) Becomes liquid
(c) Becomes hard and brittle
(d) Is covered with a thin film of Fe_3O_4
132. Which of the following is more soluble in ammonia [MH CET 2003]
- (a) $AgCl$ (b) $AgBr$
(c) AgI (d) None of these
133. Potassium permanganate works as oxidising agent both in acidic and basic medium. In both state product obtained by $KMnO_4$ is respectively [Kerala CET 2005]
- (a) MnO_2^- and Mn^{3+} (b) Mn^{3+} and Mn^{2+}
(c) Mn^{2+} and Mn^{3+} (d) MnO_2 and Mn^{2+}
(e) Mn^{2+} and MnO_2
134. Which of the following is the green coloured powder produced when ammonium dichromate is used in fire works [J & K 2005]
- (a) Cr (b) CrO_3
(c) Cr_2O_3 (d) $CrO(O_2)$
135. Which compound does not dissolve in hot dilute HNO_3 [DCE 2002; NCERT 1977]
- (a) HgS (b) CuS
(c) PbS (d) CdS
136. The least stable oxide at room temperature is [DCE 2002]
- (a) ZnO (b) CuO
(c) Sb_2O_3 (d) Ag_2O
137. Which of the following pair of elements cannot form an alloy [KCET 2005]
- (a) $Zn - Cu$ (b) $Fe - Hg$
(c) Fe, C (d) Na, Hg
138. Which of the following shows dimerisation [DCE 2002]
- (a) $HgCl_2$ (b) B_2H_6
(c) $TiCl_4$ (d) SO_2
139. Which of the following is also known as "Fools gold" [DCE 2003]
- (a) Wurtzite (b) Iron pyrites
(c) Chalcocite (d) Silver glance
140. Which one of the following is highest melting halide [DCE 2003]
- (a) $AgCl$ (b) $AgBr$
(c) AgF (d) AgI
141. $4K_2Cr_2O_7 \xrightarrow{\text{heat}} 4K_2CrO_4 + 3O_2 + X$. In the above reaction X is [DCE 2004]
- (a) CrO_3 (b) Cr_2O_7
(c) Cr_2O_3 (d) CrO_5
142. Mond's process is used for [AFMC 2004]
- (a) Ni (b) Al
(c) Fe (d) Cu
143. Stainless steel is an alloy of [AFMC 2004]
- (a) Copper (b) Nickel and chromium
(c) Manganese (d) Zinc
144. Percentage of silver in German silver is [AFMC 2004; CPMT 1985; CBSE PMT 2000; MP PMT 2001]
- (a) 0% (b) 1%
(c) 5% (d) None of these
145. Which process of purification is represented by the following scheme [Kerala PMT 2004]
- $$\text{impure } Ti + 2I_2 \xrightarrow{250^\circ C} TiI_4 \xrightarrow[1400^\circ C]{\text{pure}} Ti + I_2$$
- (a) Cupellation (b) Poling
(c) Electrolytic refining (d) Zone refining
(e) Van-Arkel process
146. Which of the following sulphides when heated strongly in air gives the corresponding metal [Kerala PMT 2004]
- (a) Cu_2S (b) CuS
(c) Fe_2S_3 (d) FeS
(e) HgS
147. Guignet's green is known as [Kerala PMT 2004]
- (a) $Cr_2O_3 \cdot 2H_2O$ (b) $FeO_3 \cdot 2H_2O$
(c) Cu_2O_3 (d) $FeCO_3 \cdot Cr_2O_3$
(e) $FeO \cdot Cr_2O_3$
148. Vanadium (III) oxide is a strong [Kerala PMT 2004]
- (a) Drying agent (b) Oxidising agent
(c) Reducing agent (d) Wetting agent
(e) Precipitating agent
149. Stainless steel does not rust because [KCET 2004]
- (a) Chromium and nickel combine with iron
(b) Chromium forms an oxide layer and protects iron from rusting
(c) Nickel present in it, does not rust
(d) Iron forms a hard chemical compound with chromium present in it.
150. The main product obtained when a solution of sodium carbonate reacts with mercuric chloride is [KCET 2004]
- (a) $Hg(OH)_2$ (b) $HgCO_3 \cdot HgO$
(c) $HgCO_3$ (d) $HgCO_3 \cdot Hg(OH)_2$
151. Which of the following has diamagnetic character [Pb. CET 2003]
- (a) $[NiCl_4]^{2-}$ (b) $[CoF_6]^{3-}$
(c) $[Fe(H_2O)_6]^{2+}$ (d) $[Ni(CN)_4]^{2-}$
152. The solubility of silver bromide in hypo solution due to the formation of [Pb. CET 2003; CPMT 1987]
- (a) $[Ag(S_2O_3)]^{-3}$ (b) Ag_2SO_3
(c) $[Ag(S_2O_3)]^{-}$ (d) $Ag_2S_2O_3$
153. Brass is an alloy of [DPMT 1982, 83; CPMT 1972, 80, 89; MLNR 1985; AFMC 1990; Pb. CET 2004; EAMCET 1993; MP PMT 1996; KCET 2000]
- (a) Zn and Sn (b) Zn and Cu
(c) Cu, Zn and Sn (d) Cu and Sn

154. Iodine is formed when KI reacts with a solution of [Pb. CET 2004]
 (a) $CuSO_4$ (b) $(NH_4)_2SO_4$
 (c) $ZnSO_4$ (d) $FeSO_4$
155. Rust is [Pb. CET 2004]
 (a) $FeO + Fe(OH)_2$ (b) Fe_2O_3
 (c) $Fe_2O_3 + Fe(OH)_2$ (d) Fe_2O_3 and $Fe(OH)_3$
156. $[Sc(H_2O)_6]^{3+}$ ion is [Pb. CET 2004]
 (a) Colourless and diamagnetic
 (b) Coloured and octahedral
 (c) Colourless and paramagnetic
 (d) Coloured and paramagnetic
157. Which of the following is called white vitriol [MP PET 1990; Bihar MEE 1995; BVP 2004]
 (a) $ZnCl_2$ (b) $MgSO_4 \cdot 7H_2O$
 (c) $ZnSO_4 \cdot 7H_2O$ (d) $Al_2(SO_4)_3$
158. $FeSO_4 \cdot 7H_2O$ shows isomorphism with [BVP 2004]
 (a) $ZnSO_4 \cdot 7H_2O$ (b) $MnSO_4 \cdot 4H_2O$
 (c) $CaSO_4 \cdot 5H_2O$ (d) $CaCl_2 \cdot 2H_2O$
159. Which pair of compound is expected to show similar colour in aqueous medium [IIT Screening 2005]
 (a) $FeCl_2$ and $CuCl_2$ (b) $VOCl_2$ and $CuCl_2$
 (c) $VOCl_2$ and $FeCl_2$ (d) $FeCl_2$ and $MnCl_2$
160. Which of the following dissolves in hot conc. $NaOH$ solution [CPMT 2004]
 (a) Fe (b) Zn
 (c) Sn (d) Ag
161. Which of the following sulphides is yellow in colour [CPMT 1983, 88, 2004; NCERT 1976]
 (a) CuS (b) CdS
 (c) ZnS (d) CoS
162. Which of the following is not oxidized by O_3 [IIT Screening 2005]
 (a) KI (b) $FeSO_4$
 (c) $KMnO_4$ (d) K_2MnO_4
163. The number of moles of $KMnO_4$ reduced by one mole of KI in alkaline medium is [CBSE PMT 2005]
 (a) One fifth (b) Five
 (c) One (d) Two
164. Excess of KI reacts with $CuSO_4$ solution and then $Na_2S_2O_3$ solution is added to it. Which of the statements is **incorrect** for this reaction [AIEEE 2004]
 (a) $Na_2S_2O_3$ is oxidised (b) CuI_2 is formed
 (c) Cu_2I_2 is formed (d) Evolved I_2 is reduced
165. The only cations present in a slightly acidic solution are Fe^{3+} , Zn^{2+} and Cu^{2+} . The reagent that when added in excess to this solution would identify and separate Fe^{3+} in one step is [IIT 1997]
 (a) $2M HCl$ (b) $6M NH_3$
 (c) $6M NaOH$ (d) H_2S gas
166. Which element is alloyed with copper to form bronze [CPMT 1972, 80, 93; CPMT 1980, 82]
 (a) Fe (b) Mn
 (c) Sn (d) Zn
167. Emery consists of [AFMC 1999]
 (a) Impure corundum (b) Impure carborundum
 (c) Impure graphite (d) Purest form of iron
168. The metal commonly present in brass and german silver is [EAMCET 1988]
 (a) Mg (b) Zn
 (c) C (d) Al
169. In the equation
 $4M + 8CN^- + 2H_2O + O_2 \longrightarrow 4[M(CN_2)]^- + 4OH^-$
 The metal M is [MP PET 2000]
 (a) Copper (b) Iron
 (c) Gold (d) Zinc
170. The term plating is [Kerala (Med.) 2002]
 (a) Platinum painting
 (b) Flat sheet of platinum
 (c) Platinum manufacturing
 (d) Platinum used as a catalyst
171. Purple of cassius is [BHU 2002]
 (a) Gold solution (b) Silver solution
 (c) Copper solution (d) Platinum solution
172. Match the items under List 1 with the compounds/elements from the List 2. Select the correct answer from the sets (a), (b), (c) and (d).
- | List 1 | List 2 |
|------------------------------------|------------------------------------|
| (i) Explosive | (A) NaN_3 |
| (ii) Artificial gem | (B) Fe_3O_4 |
| (iii) Self reduction | (C) Sn |
| (iv) Magnetic material | (D) Al_2O_3 |
| | (E) $Pb(N_3)_2$ |
| | (F) Fe_2O_3 |
| | (G) Cu |
| | (H) SiC |
| (a) (i) A, (ii) D, (iii) G, (iv) B | (b) (i) A, (ii) D, (iii) G, (iv) F |
| (c) (i) E, (ii) D, (iii) G, (iv) B | (d) (i) E, (ii) H, (iii) C, (iv) F |
173. Blood haemoglobin contains the metal
 (a) Al (b) Mg
 (c) Cu (d) Fe
174. Percentage of carbon in steel is
 (a) 2.5 – 4.5% (b) 0.25 – 0.5%
 (c) 0.2 – 1.5% (d) 3 – 3.5%
175. Steel is manufactured from
 (a) Wrought iron (b) Cast iron
 (c) (a) and (b) both (d) Haematite
176. Modern method for the manufacture of steel is
 (a) Bessemer process
 (b) Seimen-Martin's open hearth process
 (c) Duplex method
 (d) L.D. process
177. **Spiegeleisen** is an alloy of
 (a) Fe , C and Mn (b) Fe , Mg and C
 (c) Fe , Co and Cr (d) Fe , Cu and Ni
178. Stainless steel is an alloy steel of the following metals [MP PET 1990; Pb. PET 1999; KCET 2000]
 (a) Fe Only (b) Cr and Ni
 (c) W and Cr (d) Ni and Be
179. In the manufacture of steel, the Bessemer converter is containing lining of
 (a) SiO_2 (b) CaO
 (c) CaO and MgO (d) Fe_2O_3
180. Which of the following alloys contain only Cu and Zn

- (a) Bronze (b) Brass [DCE 1999]
(c) Gun metal (d) Bell metal
181. Steel becomes soft and pliable by [MP PET 1989]
(a) Annealing (b) Nitriding
(c) Tempering (d) Case hardening
182. Most stable oxidation state of iron is [AFMC 1976; CPMT 1988]
(a) +2 (b) +3
(c) -2 (d) -3
183. Nickel steel contain % of Ni [MP PMT/PET 1988]
(a) 1 - 5% (b) 3 - 5%
(c) 6 - 5% (d) 8 - 5%
184. Permanent magnet is made from [MP PET/PMT 1988; CBSE 1989]
(a) Cast iron (b) Steel
(c) Wrought Iron (d) All of these
185. In nitriding process of steel [MP PET/PMT 1988; CBSE 1989]
(a) Steel is heated in an atmosphere of ammonia
(b) Steel is made red hot and then cooled
(c) Steel is made red hot and then plunged into oil for cooling
(d) None of these
186. Iron on reacting with carbon give
(a) FeC (b) Fe_2C
(c) Fe_3C (d) FeC_2
187. Iron loses magnetic property at [KCET 2002]
(a) Melting point (b) 1000K
(c) Curie point (d) Boiling point
188. Heat treatment alters the properties of steel due to [KCET 2002]
(a) Chemical reaction on heating
(b) Partial rusting
(c) Change in the residual energy
(d) Change in the lattice structure due to differential rate of cooling
189. Pure conc. HNO_3 makes iron passive as the surface is covered with protective layer of [Orissa JEE 2002; EAMCET 1993]
(a) Fe_2O_3 (b) FeO
(c) Fe_3O_4 (d) $Fe(NO_3)_3$
190. Red hot iron absorbs SO_2 giving the product [Orissa JEE 2002]
(a) $FeS + O_2$ (b) $Fe_2O_3 + FeS$
(c) $FeO + FeS$ (d) $FeO + S$
191. If steel is heated to a temperature well below red heat and is then cooled slowly, the process is called [Kerala (Med.) 2002]
(a) Tempering (b) Hardening
(c) Softening (d) Annealing
192. In smelting of iron, which of the following reactions takes place in Blast furnace at $400^\circ C - 600^\circ C$ [MP PET 2002]
(a) $CaO + SiO_2 \rightarrow CaSiO_3$
(b) $2FeS + 3O_2 \rightarrow 2Fe + SO_2$
(c) $FeO + SiO_2 \rightarrow FeSiO_3$
(d) $Fe_2O_3 + 3CO \rightarrow 2Fe + 3CO_2$
193. Soil containing both Al and Fe is called [DPMT 2002]
(a) Laterite (b) Bauxite
(c) Pedalfers (d) Clay
194. German silver is an alloy of [EAMCET 1979; CPMT 1986, 93; MP PET/PMT 1998; UPSEAT 1999; CBSE PMT 2000; KCET 2000; MP PMT 2001]
(a) Copper, zinc and nickel (b) Copper and silver
(c) Copper, zinc and tin (d) Copper, zinc and silver
195. Iron is rendered passive by the action of [IIT 1982; MP PET 1985; MP PMT 1987; DPMT 1984; KCET 1993]
(a) Conc. H_2SO_4 (b) Conc. H_3PO_4
(c) Conc. HCl (d) Conc. HNO_3
196. Iron sheets are galvanized by depositing a coating of or In galvanisation, iron surface is coated with [MP PET 1985, 86, 87, 89, 92, 96; NCERT 1980; Bihar CEE 1995]
(a) Zinc (b) Tin
(c) Chromium (d) Nickel
197. Chemical formula of rust is [BHU 1986; MP PET 1990]
(a) FeO (b) Fe_3O_4
(c) $Fe_2O_3 \cdot xH_2O$ (d) $FeO \cdot xH_2O$
198. Heating steel to bright redness and then cooling suddenly by plunging it into oil or water, makes it [MP PET 1990]
(a) Hard and pliable (b) Soft and pliable
(c) Soft and brittle (d) Hard and brittle
199. Which of the following is found in body [CPMT 1975]
(a) Pb (b) Fe
(c) Cd (d) Al
200. Which of the following pairs of elements might form an alloy [NCERT 1981]
(a) Zinc and lead (b) Iron and mercury
(c) Iron and carbon (d) Mercury and platinum
201. Ferrous sulphate on strong heating gives
(a) SO_2 (b) $Fe_2(SO_4)_3$
(c) $FeO + SO_3$ (d) $Fe_2O_3 + SO_2 + SO_3$
202. Green vitriol is [DPMT 1985; BHU 1997; RPET 1999; JIPMER 2002]
(a) $CuSO_4 \cdot 5H_2O$ (b) $FeSO_4 \cdot 7H_2O$
(c) $CaSO_4 \cdot 2H_2O$ (d) $ZnSO_4 \cdot 7H_2O$
203. When conc. sulphuric acid is added slowly to a solution of ferrous sulphate containing nitrate ion, a brown colour ring is formed. The composition of the ring is [CPMT 1989]
(a) $[Fe(H_2O)_5NO]SO_4$ (b) $FeSO_4 \cdot NO_2$
(c) $Fe[(H_2O)_5](NO_3)_2$ (d) None of these
204. F_2 is the formed by reacting K_2MnF_6 with [AIIMS 2005]
(a) SbF_5 (b) MnF_3
(c) $KSbF_6$ (d) MnF_4
205. Railway wagon axles are made by heating rods of iron embedded in charcoal powder. The process is known as [CPMT 1972; DCE 2000; KCET 2003; UPSEAT 2001]
(a) Case hardening (b) Sheradizing
(c) Annealing (d) Tempering
206. The alloy of steel that is used in making automobile parts and utensils [EAMCET 1979; MP PMT 1992]
(a) Stainless steel (b) Nickel steel
(c) Tungstun steel (d) Chromium steel
207. Which of the following has lowest percentage of carbon

- [DPMT 1984; CPMT 1989, 91, 94; KCET 2000]
- (a) Cast iron (b) Wrought iron
(c) Steel (d) All have same percentage
208. Galvanisation is the [CPMT 1980, 86, 91, 99; MP PET/PMT 1988; Pb. PET 1999]
- (a) Deposition of Zn on Fe
(b) Deposition of Al on Fe
(c) Deposition of Sn on Fe
(d) Deposition of Cu on Fe
209. Tempered steel is
- (a) Soft and pliable
(b) Hard and brittle
(c) Neither so hard nor so brittle
(d) Very soft
210. Best quality of steel is manufactured by [BHU 1996]
- (a) Siemen –Martin's open hearth process
(b) Electrical process
(c) Bessemer process
(d) Blast furnace
211. The presence of Si in steel gives it
- (a) Fibrous structure (b) Silicate type structure
(c) Sheet type structure (d) None of these
212. The presence of Mn in steel produces
- (a) Elasticity
(b) Increases tensile strength
(c) Both (a) and (b)
(d) None of these
213. Presence of Cr in steel makes it
- (a) Resistant to chemical action
(b) Useful for making cutlery
(c) Increases chemical action
(d) (a) and (b) both
214. The addition of metals like Cr, Mn, W and Ni to ordinary steel makes it
- (a) More useful
(b) Alters the properties of ordinary steel
(c) Both (a) and (b)
(d) None of these
215. Stainless steel is non-corrosive. This character is more prominent in
- (a) Mn steel (b) Ordinary steel
(c) Ti steel (d) All of these
216. When little vanadium is mixed with steel, it becomes
- (a) More hard (b) More tensility
(c) Both (a) and (b) (d) No effect
217. To obtain steel entirely free from sulphur and phosphorus, the process used is
- (a) Electrothermal process (b) Bessemer process
(c) Open-hearth process (d) Duplex process
218. Stainless steel contains Cr
- (a) 14% (b) 5%
(c) 50% (d) 2.5%
219. The chief constituents of steel made in India are [MP PMT/PET 1988]
- (a) Mn and Cr (b) Al and Zn
(c) V and Co (d) Ni and Mg
220. Which of the following is used to prepare medical instruments
- (a) Cast iron (b) Wrought iron
(c) Steel (d) Alloy of Cu and Fe
221. A clock spring is heated to redness and then plunged into cold water. This treatment will cause it to become [NCERT 1979]
- (a) Soft and ductile (b) More springy than before
(c) Strongly magnetic (d) Hard and brittle
222. Mark the steel in which carbon % is highest
- (a) Mild steel (b) Hard steel
(c) Alloy steel (d) None of these
223. Mark the variety of iron which has highest melting point
- (a) Pig iron (b) Cast iron
(c) Wrought iron (d) Steel
224. Bessemer converter is used in the manufacture of [CPMT 1991]
- (a) Pig iron (b) Steel
(c) Wrought iron (d) Cast iron
225. Steel contains [MP PMT 1989; KCET 2000]
- (a) $Fe + C + Mn$ (b) $Fe + C + Al$
(c) $Fe + Mn$ (d) $Fe + Mn + Cr$
226. Steel differs from pig iron and wrought iron in that it contains [KCET 1991]
- (a) No carbon at all
(b) Less carbon than either
(c) More carbon than either
(d) An amount of carbon intermediate between two
227. Finely divided iron combines with CO to give [MNR 1994]
- (a) $Fe(CO)_5$ (b) $Fe_2(CO)_9$
(c) $Fe_3(CO)_{12}$ (d) $Fe(CO)_6$
228. Mohr's salt is [KCET 1993; DCE 1999; AIIMS 2000]
- (a) $FeSO_4 \cdot 7H_2O$
(b) $Fe(NH_4)SO_4 \cdot 6H_2O$
(c) $(NH_4)_2SO_4 \cdot FeSO_4 \cdot 6H_2O$
(d) $[Fe(NH_4)_2](SO_4)_2 \cdot 6H_2O$
229. Mohr's salt is [MNR 1986]
- (a) Normal salt (b) Acid salt
(c) Basic salt (d) Double salt
230. An example of double salt is [CPMT 1986; CBSE PMT 1989; Roorkee 1990]
- (a) Bleaching powder (b) $K_4[Fe(CN)_6]$
(c) Hypo (d) Potash alum
231. The passivity of iron in concentrated nitric acid is due to [MP PMT 1994]
- (a) Ferric nitrate coating on the metal
(b) Ammonium nitrate coating on the metal
(c) A thin oxide layer coating on the metal
(d) A hydride coating on the metal
232. The action of steam on heated iron is represented as [MP PMT 1994]
- (a) $3Fe + 4H_2O \rightarrow Fe_3O_4 + 4H_2$
(b) $2Fe + 3H_2O \rightarrow Fe_2O_3 + 3H_2$
(c) $Fe + H_2O \rightarrow FeO + H_2$
(d) $2Fe + H_2O + O_2 \rightarrow Fe_2O_3 + H_2$
233. Which metal is used to make alloy steel for armour plates, safes and helmets [KCET 2003]
- (a) Al (b) Mn
(c) Cr (d) Pb
234. Rusting on iron needs [MP PMT 1995]
- (a) Dry air
(b) Air and water
(c) Distilled water and carbon dioxide
(d) Oxygen and carbon dioxide
235. Iron when treated with concentrated nitric acid [MP PET 1996]



- (a) Readily reacts (b) Slowly reacts
(c) Becomes passive (d) Gives ferrous nitrate
236. An alloy which does not contain copper is [DPMT 1984]
(a) Solder (b) Bronze
(c) Brass (d) Bell metal
237. Which one of the following statements shows the correct percentage of carbon in steel, pig iron and wrought iron
(a) Steel containing less than 0.15% carbon; wrought iron 0.15 to 2.0% carbon; and pig iron over 2% carbon
(b) Pig iron less than 0.15% carbon; wrought iron 0.15 to 2.0% carbon; and steel over 2% carbon
(c) Wrought iron less than 0.15% carbon; steel 0.15 to 2.0% carbon; and pig iron over 2% carbon
(d) Wrought iron less than 0.15% carbon; pig iron 0.15 to 2.0% carbon; and steel over 2.0% carbon
238. In the Bessemer and open-hearth process for the manufacture of steel, which one of the following is used for the removal of carbon in part or whole
- | Bessemer | Open-hearth |
|------------------|--------------|
| (a) Ferric oxide | Air |
| (b) Air | Ferric oxide |
| (c) Oxygen | Scrap iron |
| (d) Air | Scrap iron |
239. About the basic open hearth process, which statement is wrong
(a) Limestone is added to the charge
(b) Phosphorus impurity cannot be removed by this process
(c) Carbon content of the steel can be uniformly controlled over a series of batches
(d) Iron scrap can be utilized
240. Which of the following statements is wrong
(a) Heating to a high temperature and then cooling suddenly, e.g. by dipping in water, makes steel hard and brittle
(b) Steel can be softened by heating it to a high temperature for a prolonged time and then cooling slowly. This is called quenching.
(c) Tempering of hardened steel is done by heating it to just below red heat at controlled temperature and duration
(d) Phosphorus impurity makes steel 'cold short'
241. Bell metal is an alloy of [DPMT 1990, 96; CBSE PMT 1999; Kerala PMT 2002]
(a) Cu, Zn and Sn (b) Cu, Zn and Ni
(c) Cu and Zn (d) Cu and Sn
242. Turnbull's blue is [Bihar CEE 1995]
(a) Ferricyanide (b) Ferrous ferricyanide
(c) Ferrous cyanide (d) Ferri-ferricyanide
243. Addition of high proportions of manganese makes steel useful in making rails of rail-roads because manganese [IIT 1998]
(a) Gives hardness to steel
(b) Helps the formation of oxides of iron
(c) Can remove oxygen and sulphur
(d) Can show highest oxidation state of + 7
244. Copper displaces which of the metal from their salt solutions [CPMT 1988]
(a) $AgNO_3$ (b) $ZnSO_4$
(c) $FeSO_4$ (d) All of these
245. Which of the following statement(s) is(are) correct with reference to the ferrous and ferric ions [IIT 1998]
(a) Fe^{3+} gives brown colour with potassium ferricyanide
(b) Fe^{2+} gives blue precipitate with potassium ferricyanide
(c) Fe^{3+} gives red colour with potassium thiocyanate
(d) Fe^{2+} gives brown colour with ammonium thiocyanate
246. Which of the following element constitutes a major impurity in pig iron [CBSE PMT 1998]
(a) Silicon (b) Oxygen
(c) Sulphur (d) Graphite
247. Annealing is [Pune CET 1998; AFMC 2002]
(a) Heating steel in nitrogen and cooling
(b) Heating steel to bright redness and then cooling slowly
(c) Heating wrought iron with carbon to redness
(d) Heating steel to high temperature and cooling suddenly by plunging in water
248. In electroplating, the metal that is not used for plating is [Pune CET 1998]
(a) Fe (b) Zn
(c) Ni (d) Au
249. Which one of the following is a wrong statement about cast iron [KCET 1998]
(a) It is also called pig iron
(b) It contains about 4.5% carbon
(c) It is corrosion resistant
(d) It contracts on cooling
250. Iron pipes lying under acidic soil are often attached to blocks of magnesium for protection from rusting. Magnesium offers protection to iron against corrosion because it [KCET 1998]
(a) Is more readily converted into positive ions
(b) Is lighter than iron
(c) Forms a corrosion-resistant alloy with iron
(d) Prevents air from reaching the surface of iron
251. FeS_2 is [RPET 1999]
(a) Artificial silver (b) Fool's gold
(c) Mohr's salt (d) Cast iron
252. Stainless steel is an alloy of iron with [DCE 1999]
(a) 8% Cr , 50% Mn (b) 10% Ni , 2% Mn
(c) 2% Cr , 3% C (d) 12% Cr , 1% N
253. The chemical processes in the production of steel from haematite ore involve [IIT-JEE (Screening) 2000]
(a) Reduction
(b) Oxidation
(c) Reduction followed by oxidation
(d) Oxidation followed by reduction
254. The protection of steel by chrome plating is due to [MP PMT 2001]
(a) Cathodic protection
(b) Anodic protection
(c) Covering of steel surface
(d) Formation of alloy with iron
255. The most convenient method to protect bottom of ship made of iron is [CBSE PMT 2001; Kerala (Engg.) 2002]
(a) White tin plating
(b) Coating with red lead oxide
(c) Connecting with 'PB' block
(d) Connecting with 'Mg' block
256. Carbon monoxide reacts with iron to form [KCET (Med.) 2001]
(a) $Fe(CO)_5$ (b) $FeCO_2$
(c) $FeO + C$ (d) $Fe_2O_3 + C$
257. Iron is extracted from magnetite by reduction with

[UPSEAT 2001]

- (a) H_2 (b) C
 (c) Mg (d) Al
258. Malachite is a mineral of
 [MP PMT 1990; MP PET 1992, 98, 2000; MP PMT 1998]
 (a) Zn (b) Fe
 (c) Hg (d) Cu
259. The most important oxidation state of copper is
 [MP PMT 1987]
 (a) +1 (b) +2
 (c) +3 (d) +4
260. Hot and conc. nitric acid when reacts with copper, the gas obtained is
 (a) N_2 (b) Nitrous oxide
 (c) NO (d) NO_2
261. Which of the following property is not expected to be shown by copper
 [MP PET/PMT 1988; NCERT 1975; MP PET 1989]
 (a) High thermal conductivity
 (b) Low electrical conductivity
 (c) Ductility
 (d) Malleability
262. Which of the following metal gives more than one chloride
 (a) Cu (b) Al
 (c) Ag (d) Na
263. The metal which is the best conductor of electricity is
 [CPMT 1996]
 (a) Iron (b) Copper
 (c) Silver (d) Aluminium
264. Paris green is
 (a) Double salt of copper carbonate and copper nitrate
 (b) Double salt of copper acetate and copper arsenite
 (c) Double salt of copper acetate and copper sulphate
 (d) Double salt of copper and silver nitrate
265. Reaction between the following pairs will produce H_2 except [CPMT 1973; CBSE PMT 1998]
 (a) Na + ethyl alcohol (b) Fe + steam
 (c) $Fe + H_2SO_4 (aq.)$ (d) $Cu + HCl (aq.)$
266. Which of the following is correct [BHU 1995]
 (a) Gun metal : $Cu + Zn + Sn$
 (b) Duralumin : $Al + Cu + Mg + Ag$
 (c) German silver : $Cu + Zn + C$
 (d) Solder : $Pb + Al$
267. Solder is an alloy of [IIT 1995; MP PET 1995; AFMC 2005]
 (a) 70% lead, 30% tin (b) 30% lead, 70% tin
 (c) 80% lead, 20% tin (d) 90% Cu , 10% tin
268. Zinc forms two important alloys, (i) Brass and (ii) German silver. Metals present in them mainly are
 (a) In (i) zinc and tin; and in (ii) zinc, silver and nickel
 (b) In (i) zinc and iron; and in (ii) zinc, nickel and cobalt
 (c) In (i) zinc and copper; and in (ii) zinc, copper and nickel
 (d) In (i) zinc and aluminium; and in (ii) zinc, nickel and aluminium
269. One of the constituent of german silver is
 [IIT 1980; Kurukshetra CEE 1998; DCE 1999]
 (a) Ag (b) Cu
 (c) Mg (d) Al
270. Gun metal is an alloy of [MP PMT 1990; CPMT 1997]
 (a) Cu and Al (b) Cu, Sn and Zn

(c) Cu, Zn and Ni (d) Cu and Sn

271. Besides Zn and Cu , german silver contains the metal
 [MP PET 1997]
 (a) Sn (b) Ag
 (c) Ni (d) Mg
272. Which metal is present in brass, bronze and German silver
 [CPMT 1997; AFMC 1998; AIIMS 1999; J & K 2005]
 (a) Zn (b) Mg
 (c) Cu (d) Al
273. Which of the following is wrongly matched [KCET (Med.) 1999]
 (a) German silver $Cu + Zn + Ni$
 (b) Alnico $Fe + Al + Ni + CO$
 (c) Monel metal $Cu + Zn + Sn$
 (d) Duralumin $Al + Cu + Mg + Mn$
274. An extremely hot copper wire reacts with steam to give
 [CPMT 1988]
 (a) CuO (b) Cu_2O
 (c) Cu_2O_2 (d) CuO_2
275. From a solution of $CuSO_4$, the metal used to recover copper is [MP PET 1992; B]
 (a) Sodium (b) Iron
 (c) Silver (d) Hg
276. Copper sulphate is commercially made from copper scraps by
 [CPMT 1973]
 (a) Dissolving in hot conc. H_2SO_4
 (b) The action of dil. H_2SO_4 and air
 (c) Heating with sodium sulphate
 (d) Heating with sulphur
277. Cuprous ion is colourless, while cupric ion is coloured because [EAMCET 1992; B]
 (a) Both have unpaired electrons in d -orbital
 (b) Cuprous ion has a completed d -orbital and cupric ion has an incomplete d -orbital
 (c) Both have half-filled p and d -orbitals
 (d) Cuprous ion has incomplete d -orbital and cupric ion has a completed d -orbital
278. A blue colouration is not obtained when [CBSE PMT 1989]
 (a) Ammonium hydroxide dissolves in copper sulphate
 (b) Copper sulphate solution reacts with $K_4[Fe(CN)_6]$
 (c) Ferric chloride reacts with sodium ferrocyanide
 (d) Anhydrous $CuSO_4$ is dissolved in water
279. Identify the statement which is not correct regarding $CuSO_4$ [MNR 1992; Pb. P]
 (a) It reacts with KI to give iodine
 (b) It reacts with KCl to give Cu_2Cl_2
 (c) It reacts with $NaOH$ and glucose to give Cu_2O
 (d) It gives CuO on strong heating in air
280. Copper sulphate solution reacts with KCN to give [MP PMT 1992; MNR 1994; IIT 1996; AIIMS 1999; CBSE PMT 2002]
 (a) $Cu(CN)_2$ (b) $CuCN$
 (c) $K_2[Cu(CN)_4]$ (d) $K_3[Cu(CN)_4]$

281. If excess of NH_4OH is added to $CuSO_4$ solution, it forms blue coloured complex which is
[MP PMT 1971, 79; Bihar CEE 1995; RPET 1999; AFMC 2001]
- (a) $Cu(NH_3)_4SO_4$ (b) $Cu(NH_3)_2SO_4$
(c) $Cu(NH_4)_4SO_4$ (d) $Cu(NH_4)_2SO_4$
282. Which of the following metals displaces SO_2 gas from concentrated sulphuric acid
(a) Mg (b) Zn
(c) Cu (d) None of these
283. The method of zone refining of metals is based on the principle of
(a) Greater solubility of the impurity in the molten state than in the solid
(b) Greater mobility of the pure metal than that of the impurity
(c) Higher melting point of the impurity than that of the pure metal
(d) Greater noble character of the solid metal than that of the impurity
284. A metal when left exposed to the atmosphere for some time becomes coated with green basic carbonate. The metal in question is [NDA 1999]
(a) Copper (b) Nickel
(c) Silver (d) Zinc
285. When $CuSO_4$ solution is added to $K_4[Fe(CN)_6]$, the formula of the product formed is [Bihar CEE 1995]
(a) $Cu_2Fe(CN)_6$ (b) KCN
(c) $Cu(CN)_3$ (d) $Cu(CN)_2$
286. MnO_4^- on reduction in acidic medium forms [MP PMT 1995]
(a) MnO_2 (b) Mn^{++}
(c) MnO_4^{--} (d) Mn
287. Which of the following metals will not react with a solution of $CuSO_4$ [CPMT 1996]
(a) Fe (b) Zn
(c) Mg (d) Hg
288. Which one of the following metals will not reduce H_2O [EAMCET 1997]
(a) Ca (b) Fe
(c) Cu (d) Li
289. The reaction, which forms nitric oxide, is [KCET (Med.) 2001]
(a) C and N_2O (b) Cu and N_2O
(c) Na and NH_3 (d) Cu and HNO_3
290. A cuprous ore among the following is [KCET 2002]
(a) Cuprite (b) Malachite
(c) Chalcopyrites (d) Azurite
291. When metallic copper comes in contact with moisture, a green powdery/ pasty coating can be seen over it. This is chemically known as [AFMC 2002]
(a) Copper sulphide - Copper carbonate
(b) Copper carbonate - Copper sulphate
(c) Copper carbonate - Copper hydroxide
(d) Copper Sulphate - Copper sulphide
292. Orford process is used in extraction of
(a) Fe (b) Co
(c) Pt (d) Ni
293. Horn silver is
(a) $AgCl$ (b) Ag
(c) $AgBr$ (d) CH_3COOAg
294. Which of the following is used in photography [CPMT 1980]
(a) $AgCl$ (b) $AgBr$
(c) AgI (d) Ag_2O
295. Silver halides are used in photography because [CBSE PMT 2003] [MP PMT 1989]
(a) They are photosensitive
(b) Soluble in hypo
(c) Soluble in NH_4OH
(d) Soluble in acids
296. $AgCl$ when heated with Na_2CO_3 gives [CPMT 1980; MP PET 1989; MP PMT 1982, 84]
(a) Ag_2O (b) Ag
(c) Ag_2CO_3 (d) $NaAgCO_3$
297. $AgNO_3$ gives a red ppt. with [NCERT 1972; BHU 1978; MP PMT 1995]
(a) KI (b) $NaBr$
(c) $NaNO_3$ (d) K_2CrO_4
298. Silver nitrate is prepared by [CPMT 1984]
(a) The action of only conc. HNO_3 on silver
(b) Heating silver oxide with NO_2
(c) The action of hot dil. HNO_3 on silver
(d) Dissolve Ag in aqua-regia
299. $AgCl$ is soluble in [EAMCET 1992]
(a) Aqua-regia (b) H_2SO_4
(c) HCl (d) NH_3 (aq)
300. Which of the following is least soluble in water [NCERT 1974, 78; MNR 1984, 89]
(a) AgI (b) $AgCl$
(c) $AgBr$ (d) Ag_2S
301. Photographic films and plates have an essential ingredient of [CPMT 1980; CBSE PMT 1989]
(a) Silver nitrate (b) Silver bromide
(c) Sodium chloride (d) Oleic acid
302. Which of the following does not react with $AgCl$ [AIIMS 1997]
(a) $NaNO_3$ (b) Na_2CO_3
(c) $Na_2S_2O_3$ (d) NH_4OH
303. Which one of the following is known as lunar caustic when in fused state [MP PMT 1999; JIPMER 2002]
(a) Silver nitrate (b) Silver sulphate
(c) Silver chloride (d) Sodium sulphate
304. Which silver halide is used in medicine [DPMT 1996]
(a) $AgNO_3$ (b) $AgCl$
(c) $AgBr$ (d) AgF

305. When silver nitrate is heated to red hot, what is formed
[CPMT 1996; NCERT 1970]
- (a) Ag (b) Ag_2O
(c) Ag_2O_3 (d) AgO_2
306. For making Ag from $AgNO_3$, which of the following is used [AFMC 1998]
- (a) PH_3 (b) AsH_3
(c) Na_2CO_3 (d) NH_3
307. Which one of the following reacts with concentrated sulphuric acid [EAMCET 1998]
- (a) Au (b) Ag
(c) Pt (d) Pb
308. The nitrate of which metal left globule on heating strongly
[JIPMER 2001]
- (a) $Pb(NO_3)_2$ (b) $NaNO_3$
(c) $AgNO_3$ (d) $Cu(NO_3)_2$
309. During extraction of silver, which of the following is formed
[MP PET 2002]
- (a) $Na[Ag(CN)_2]$ (b) $Na_2[Ag(CN)_2]$
(c) $Na_4[Ag(CN)_2]$ (d) None of these
310. Colourless solutions of the following four salts are placed separately in four different test tubes and a strip of copper is dipped in each one of these. Which solution will turn Blue
[MP PET 2002]
- (a) KNO_3 (b) $AgNO_3$
(c) $Zn(NO_3)_2$ (d) $ZnSO_4$
311. Zinc when reacted with excess of $NaOH$ gives
[CPMT 1974, 78, 94; MP PMT 1999]
- (a) Zinc hydroxide (b) Zinc oxide
(c) Di sodium zincate (d) Sodium zincate
312. Pair of metals which dissolves in $NaOH$ solution
- (a) Al, Cu (b) Zn, Hg
(c) Zn, Cu (d) Zn, Al
313. Lucas reagent is
[CPMT 1980; AIIMS 1980, 82; DPMT 1983; MP PET 1995; MP PMT 1997, 98]
- (a) Anhydrous $ZnCl_2 + conc. HCl$
(b) Hydrated $ZnCl_2 + dil. HCl$
(c) Conc. $HNO_3 + anhydrous ZnCl_2$
(d) Conc. $HNO_3 + anhydrous MgCl_2$
314. What is the effect of shaking dil. H_2SO_4 with small quantity of anhydrous $CuSO_4$
[NCERT 1975; CPMT 1975, 88]
- (a) The white solid dissolves to form a colourless solution
(b) The white solid dissolves to form a green solution
(c) The white solid turns blue but does not dissolve
(d) The white solid dissolves to form a blue solution
315. Which metal is electro-deposited on iron surface to prevent rusting [MP PET 1990]
- (a) Cu (b) Zn
(c) Mg (d) Pb
316. To prevent corrosion, iron pipes carrying drinking water are covered with zinc. The process involved is
[CPMT 1986; MP PMT 1993; MP PET 1999]
- (a) Photoelectrolysis (b) Electroplating
(c) Galvanization (d) Cathodic protection
317. From aqueous solution of $ZnSO_4$, normal zinc carbonate may be precipitated by [CPMT 1973]
- (a) Boiling with $CaCO_3$ (b) Adding Na_2CO_3
(c) Adding $NaHCO_3$ (d) Passing CO_2
318. Which one of the following dissolve in hot concentrated $NaOH$ solution [IIT 1980]
- (a) Fe (b) Zn
(c) Cu (d) Ag
319. Which of the following metal forms an amphoteric oxide
[CPMT 1976]
- (a) Ca (b) Fe
(c) Cu (d) Zn
320. Reaction of zinc with cold and very dilute nitric acid yields
[MP PET 1985, 92, 97; BHU 1995, 2000; NCERT 1974; MP PMT 1995]
- (a) $Zn(NO_3)_2 + N_2O$ (b) $Zn(NO_3)_2 + NO$
(c) $Zn(NO_3)_2 + NH_4NO_3$ (d) $Zn(NO_3)_2 + NO_2$
321. The number of unpaired electrons in Zn^{2+} is
- (a) 2 (b) 3
(c) 4 (d) 0
322. The trace metal present in insulin is [KCET 1991]
- (a) Iron (b) Cobalt
(c) Zinc (d) Manganese
323. The chemical name of borax is [CPMT 1994]
- (a) Sodium orthoborate
(b) Sodium metaborate
(c) Sodium tetraborate
(d) Sodium tetraborate decahydrate
324. Hydrogen is not obtained when zinc reacts with
[CPMT 1994]
- (a) Cold water (b) Dilute H_2SO_4
(c) Dilute HCl (d) Hot 20% $NaOH$
325. The metal which gives hydrogen on treatment with acid as well as sodium hydroxide is [MP PET 1996]
- (a) Iron (b) Zinc
(c) Copper (d) None of the above
326. To protect iron against corrosion, the most durable metal plating on it, is [CBSE PMT 1994]
- (a) Nickel plating (b) Tin plating
(c) Copper plating (d) Zinc plating
327. The compound $ZnFe_2O_4$ is [Kerala (Engg.) 2002]
- (a) A normal spinel compound
(b) Interstitial compound
(c) Covalent compound
(d) Co-ordination compound
328. ZnO when heated with BaO at $1100^\circ C$ gives a compound. Identify the compound [AFMC 2002]
- (a) $BaZnO_2$ (b) $BaO_2 + Zn$
(c) $BaCdO_2$ (d) $Ba + ZnO_2$
329. Zn gives hydrogen gas with H_2SO_4 and HCl but not with HNO_3 because [CBSE PMT 2002]
- (a) NO_2 is reduced in preference to H_3O^+

- (b) HNO_3 is weaker acid than H_2SO_4 and HCl
 (c) Zn acts as oxidising agent when reacts with HNO_3
 (d) In electrochemical series Zn is placed above the hydrogen
- 330.** The metal used for making radiation shield is [Kerala (Med.) 2002]
 (a) Aluminium (b) Iron
 (c) Zinc (d) Lead
- 331.** Which of the following metal is obtained by leaching out process using a solution of $NaCN$ and then precipitating the metal by addition of zinc dust [NCERT 1984; AIIMS 1983; CBSE PMT 1989]
 (a) Copper (b) Silver
 (c) Nickel (d) Iron
- 332.** While extracting an element from its ore, the ore is ground and leached with dilute KCN solution to form the soluble product potassium argentocyanide. The element is [CBSE PMT 1989]
 (a) Lead (b) Chromium
 (c) Manganese (d) Silver
- 333.** In Mc Arthur Forest method, silver is extracted from the solution of $Na[Ag(CN)_2]$ by the use of: [CPMT 2004]
 (a) Fe (b) Mg
 (c) Cu (d) Zn
- 334.** Iron obtained from blast furnace is known as [DPMT 1981; CPMT 1988; MP PET 2000]
 (a) Wrought iron (b) Cast iron
 (c) Pig iron (d) Steel
- 335.** Extraction of silver from commercial lead is possible by [BHU 1979]
 (a) Mond's process (b) Park's process
 (c) Haber's process (d) Clark's process
- 336.** Impurities of lead in silver are removed by [AIIMS 1987]
 (a) Park process (b) Solvey process
 (c) Cyanide process (d) Amalgamation process
- 337.** Park's process is used in the extraction of [BHU 1977; CBSE PMT 1992; MP PMT 1996; Kurukshetra CEE 1998]
 (a) Iron (b) Zinc
 (c) Silver (d) Lead
- 338.** From argentite (Ag_2S) ore the method used in obtaining metallic silver is [MP PMT 1989]
 (a) Fused mixture of Ag_2S and KCl is electrolysed
 (b) Ag_2S is reduced with CO
 (c) Ag_2S is roasted to Ag_2O which is reduced with carbon
 (d) Treating argentite with $NaCN$ solution followed by metal displacement with zinc
- 339.** In the extraction of zinc which gas is burnt in the jackets surrounding the retorts
 (a) Water gas (b) Producer gas
 (c) Oil gas (d) Coal gas
- 340.** MacArther process is used for [BHU 1995]
 (a) Hg (b) Fe
 (c) Cl (d) O_2
- 341.** In the metallurgy of zinc, the zinc dust obtained from roasting and reduction of zinc sulphide contains some ZnO . How is this removed [MP PET 1993; AFMC 2002]
 (a) Absorbance of ultraviolet light and re-emission of white light is employed
 (b) Shock cooling by contact with a shower of molten lead is done
 (c) X-ray method is used
 (d) Smelting is employed
- 342.** In the metallurgy of copper, metallic copper is finally formed in the furnace by the reactions
 (a) $Cu_2S + O_2 \rightarrow 2Cu + SO_2$
 (b) $2CuS + 3O_2 \rightarrow 2CuO + 2SO_2$
 $2CuO + CuS \rightarrow 3Cu + SO_2$
 (c) $2Cu_2S + 3O_2 \rightarrow 2Cu_2O + 2SO_2$
 $Cu_2S + 2Cu_2O \rightarrow 6Cu + SO_2$
 (d) $CuS + O_2 \rightarrow Cu + SO_2$
- 343.** In the smelting of roasted copper pyrites ore, melting occurs so that the first reaction is
 (a) All the sulphur preferentially combines with iron to form FeS and CuO is formed
 (b) All the sulphur preferentially combines with copper to form CuS and FeO is formed
 (c) All the sulphur preferentially combines with iron to form FeS and Cu_2O is formed
 (d) All the sulphur preferentially combines with copper to form Cu_2S and FeO is formed
- 344.** In the oxidation of Cu , the reaction which takes place in bessemer converter is [CPMT 1999]
 (a) $2CuFeS_2 + O_2 \rightarrow Cu_2S + FeS + SO_2$
 (b) $2Cu_2S + 3O_2 \rightarrow 2Cu_2O + 2SO_2$
 (c) $2Cu_2O + Cu_2S \rightarrow 6Cu + SO_2$
 (d) $2FeS + 3O_2 \rightarrow 2FeO + 2SO_2$
- 345.** Silica is added to roasted copper ore during smelting in order to remove [KCET 1998]
 (a) Cuprous sulphide (b) Cuprous oxide
 (c) Ferrous oxide (d) Ferrous sulphide
- 346.** Parke's process is used to extract [MP PMT 2002]
 (a) Silver using $NaCN$
 (b) Copper using $CuFeS_2$
 (c) Silver from argentiferrous lead
 (d) Silver by forming amalgam
- 347.** Identify the reaction that doesn't take place during the smelting process of copper extraction [KCET 2003]
 (a) $2FeS + 3O_2 \rightarrow 2FeO + 2SO_2 \uparrow$
 (b) $Cu_2O + FeS \rightarrow Cu_2S + FeO$
 (c) $2Cu_2S + 3O_2 \rightarrow 2Cu_2O + 2SO_2 \uparrow$
 (d) $FeO + SiO_2 \rightarrow FeSiO_3$
- 348.** The extraction of which of the following metals involves bessemerisation [DCE 2004]
 (a) Fe (b) Ag
 (c) Al (d) Cu
- 349.** Bessemer converter is used for [AFMC 2004]
 (a) Steel (b) Wrought iron
 (c) Pig iron (d) Cast iron
- 350.** In the cyanide process for the extraction of silver, sodium cyanide is used to [MP PMT 1994]
 (a) Convert silver into a soluble silver complex
 (b) Reduce silver

- (c) Precipitate silver
(d) Oxidise silver
351. Parke's process of desilverization of lead depends upon
- (a) Partition coefficient of silver between molten zinc/molten lead having a high value
(b) Partition coefficient of silver between molten zinc/molten lead having a low value
(c) Crystallizing out of pure lead while the silver-lead eutectic which has a lower melting point is left behind in liquid form
(d) Chemical combination of zinc and silver which precipitates out easily
352. Which is impure form of iron
- (a) Cast iron (b) Wrought iron
(c) Steel iron (d) None
353. Blister copper is [CPMT 1976, 85, 2002; DPMT 1982; MP PET 1995; Bihar CEE 1995]
- (a) Pure copper (b) Ore of copper
(c) Alloy of copper (d) 1% impure copper
354. In the extraction of copper when molten copper is cooled slowly, blister copper is obtained due to evolution of the following gas
- (a) Water vapour (b) Sulphur dioxide
(c) Carbon dioxide (d) Carbon monoxide
355. In the electrolytic purification of copper some gold is found in the [CPMT 1972; AFMC (1995); MP PET 2003]
- (a) Cathode (b) Cathode mud
(c) Anode mud (d) Electrolyte
356. Purest form of iron is [CPMT 1975, 80, 84, 87, 89; DPMT 1982, 83; MP PMT 1987, 90, 91; MP PET 1995; BHU 1999; MH CET 2003]
- (a) Cast iron (b) Wrought iron
(c) Hot steel (d) Stainless steel
357. *Spelter* is [CPMT 1988]
- (a) Impure *Cu* (b) Impure *Zn*
(c) *ZnO* (d) *CuO*
358. A copper coin is completely covered with a gold film and is placed in dilute HNO_3 . This will result in formation of [CPMT 1981]
- (a) Gold nitrate (b) Copper nitrate
(c) None of these (d) Purple of cassins
359. When zinc is added to $CuSO_4$ copper gets ppt. due to [CPMT 1979]
- (a) Reduction of copper ions
(b) Oxidation of copper ions
(c) Hydrolysis of copper sulphate
(d) Complex formation
360. Addition of iron filings to $CuSO_4$ solution caused precipitation of *Cu* owing to the [CPMT 1990]
- (a) Reduction of Cu^{++}
(b) Oxidation of Cu^{++}
(c) Reduction of *Fe*
(d) Reduction of Fe^{+++}
361. Oxygen gas can be prepared from solid $KMnO_4$ by [DPMT 2001]
- (a) Strongly heating the solid

- (b) Treating the solid with H_2 gas
(c) Dissolving the solid in dil. H_2SO_4
(d) Dissolving the solid in dil. *HCl*

Critical Thinking

Objective Questions

1. Transition metal with low oxidation number will act as [DCE 2001]
- (a) A base
(b) An acid
(c) An oxidising agent
(d) None of these
2. Which of the following pair will have effective magnetic moment equal
- (a) Cr^{+3} and Mn^{+2} (b) Cr^{+2} and Fe^{+2}
(c) V^{+2} and Sc^{+3} (d) Ti^{+2} and V^{+2}
3. Which is least soluble in water [MNR 1984, 89]
- (a) *AgCl* (b) *AgBr*
(c) *AgI* (d) *Ag_2S*
(e) *AgF*
4. Which one of the following has the maximum number of unpaired electrons [UPSEAT 2001]
- (a) Mg^{2+} (b) Ti^{3+}
(c) V^{3+} (d) Fe^{2+}
5. Which of the following ions form most stable complex compound [MP PMT 1995]
- (a) Cu^{++} (b) Ni^{++}
(c) Fe^{++} (d) Mn^{++}
6. Mn^{++} can be converted into Mn^{7+} by reacting with [UPSEAT 2003]
- (a) SO_2 (b) Cl_2
(c) PbO_2 (d) $SnCl_2$
7. General configuration of outermost and penultimate shell is $(n-1)s^2(n-1)p^6(n-1)d^xns^2$. If $n = 4$ and $x = 5$ then no. of proton in the nucleus will be [MP PET 2003]
- (a) > 25 (b) < 24
(c) 25 (d) 30
8. Which transition metal reduces steam to evolve hydrogen [MP PMT 2003; DCE 2002]
- (a) *Mg* (b) *Fe*
(c) *Sc* (d) *Pt*
9. Chloride of which of the following element is coloured [MP PMT 1990]
- (a) *Ag* (b) *Hg*
(c) *Zn* (d) *Co*
10. Arrange Ce^{+3} , La^{+3} , Pm^{+3} and Yb^{+3} in increasing order of their ionic radii [AIEEE 2002]
- (a) $Yb^{+3} < Pm^{+3} < Ce^{+3} < La^{+3}$
(b) $Ce^{+3} < Yb^{+3} < Pm^{+3} < La^{+3}$
(c) $Yb^{+3} < Pm^{+3} < La^{+3} < Ce^{+3}$
(d) $Pm^{+3} < La^{+3} < Ce^{+3} < Yb^{+3}$

11. $KMnO_4$ reacts with ferrous sulphate according to the equation
 $MnO_4^- + 5Fe^{2+} + 8H^+ \rightarrow Mn^{2+} + 5Fe^{3+} + 4H_2O$ Here
 10ml of 0.1M $KMnO_4$ is equivalent to [CPMT 1999]
- (a) 20ml of 0.1M $FeSO_4$
 (b) 30ml of 0.1M $FeSO_4$
 (c) 40ml of 0.1M $FeSO_4$
 (d) 50ml of 0.1M $FeSO_4$
12. Which of the following is/are soluble in ethanol [Roorkee Qualifying 1998]
- (a) HgF_2 (b) $HgCl_2$
 (c) $HgBr_2$ (d) HgI_2
13. Out of all the known elements, the percentage of transitional elements is approximately
- (a) 30% (b) 50%
 (c) 60% (d) 75%
14. Atomic number of Cr and Fe are 24 and 26 respectively. Which of the following is paramagnetic with the spin of electron [CBSE PMT 2002]
- (a) $[Cr(NH_3)_6]^{+3}$ (b) $[Fe(CO)_5]$
 (c) $[Fe(CN)_6]^{-4}$ (d) $[Cr(CO)_6]$
15. Which of the following is not an element [DCE 2001]
- (a) Graphite (b) Diamond
 (c) 22-carat gold (d) Rhombic sulphur
16. Which of the following is more paramagnetic [DCE 2001]
- (a) Fe^{+2} (b) Fe^{+3}
 (c) Cr^{+3} (d) Mn^{+3}
17. The number of d-electrons in Fe^{2+} (at no. of $Fe = 26$) is not equal to that of the [MNR 1994]
- (a) p - electrons in Ne (at. no. = 10)
 (b) s - electrons in Mg (at. no. = 12)
 (c) d - electrons in Fe
 (d) p - electrons in Cl^- (at. no. of $Cl = 17$)
18. The basic character of the transition metal monoxides follows the order [CBSE PMT 2003]
- (a) $TiO > VO > CrO > FeO$
 (b) $VO > CrO > TiO > FeO$
 (c) $CrO > VO > FeO > TiO$
 (d) $TiO > FeO > VO > CrO$
 (Atomic no. $Ti = 22, V = 23, Cr = 24, Fe = 26$)
19. Amongst following the lowest degree of paramagnetism per mole of the compound at 298 K will be shown by
- (a) $MnSO_4 \cdot 4H_2O$ (b) $CuSO_4 \cdot 5H_2O$
 (c) $FeSO_4 \cdot 6H_2O$ (d) $FeSO_4 \cdot 5H_2O$
20. In nitroprusside ion, the iron and NO exist as Fe^{II} and NO^+ rather than Fe^{III} and NO . These forms can be differentiated by [IIT-JEE 1998]
- (a) Estimating the concentration of iron
 (b) Measuring the concentration of CN^-
 (c) Measuring the solid state magnetic moment
 (d) Thermally decomposing the compound
21. Among the following, the compound that is both paramagnetic and coloured is [IIT-JEE 1997]
- (a) $K_2Cr_2O_7$ (b) $(NH_4)_2(TiCl_6)$
 (c) VO_2 (d) $K_3[Cu(CN)_4]$
22. The number of moles of $KMnO_4$ that will be needed to react completely with one mole of ferrous oxalate $Fe(C_2O_4)$ in acidic solution is [IIT-JEE 1997; KCET 1996]
- (a) 3/5 (b) 2/5
 (c) 4/5 (d) 1
23. In following reaction
 $yMnO_4^- + xH^+ + C_2O_4^{2-} \rightarrow yMn^{2+} + 2CO_2 + \frac{x}{2}H_2O$,
 x and y are [CPMT 1997]
- (a) 2 and 16 (b) 16 and 2
 (c) 8 and 16 (d) 5 and 2
24. Which of the following weighs less when weighed in magnetic field
- (a) VCl_3 (b) $ScCl_3$
 (c) $TiCl_3$ (d) $FeCl_3$
25. An element is in M^{3+} form. Its electronic configuration is $[Ar]3d^1$ the ion is [JIPMER 2002]
- (a) Ti^{3+} (b) Ti^{4+}
 (c) Ca^{2+} (d) Sc^{3+}
26. The atomic number of vanadium (V), chromium (Cr), manganese (Mn) and iron (Fe) are respectively 23, 24, 25 and 26 which one of these may be expected to have the highest second ionization enthalpy [AIEEE 2003]
- (a) V (b) Cr
 (c) Mn (d) Fe

Assertion & Reason

For AIIMS Aspirants

Read the assertion and reason carefully to mark the correct option out of the options given below :

- (a) If both assertion and reason are true and the reason is the correct explanation of the assertion.
 (b) If both assertion and reason are true but reason is not the correct explanation of the assertion.
 (c) If assertion is true but reason is false.
 (d) If the assertion and reason both are false.
 (e) If assertion is false but reason is true.

1. Assertion : Cuprous ion (Cu^+) has unpaired electrons while cupric ion (Cu^{++}) does not
 Reason : Cuprous ion (Cu^+) is colourless whereas cupric ion (Cu^{++}) is blue in the aqueous solution [AIIMS 2002]
2. Assertion : Zn^{2+} is diamagnetic
 Reason : The electrons are lost from 4 s orbital to form Zn^{2+} [IIT-JEE 1998]
- Assertion : Transition metals show variable valence.
 Reason : Due to a large energy difference between the ns^2 and $(n-1)d$ electrons. [AIIMS 1996]
4. Assertion : The aqueous solution of $FeCl_3$ is basic in nature.
 Reason : $FeCl_3$ hydrolyses in water. [AIIMS 1998]
5. Assertion : $AgCl$ dissolves in NH_4OH solution.
 Reason : Due to formation of a complex.

[AIIMS 1998]

6. Assertion : Pure iron is not used for making tools and machines.

Reason : Pure iron is hard. [AIIMS 1998]

7. Assertion : Solution of Na_2CrO_4 in water is intensely coloured.

Reason : Oxidation state of Cr in Na_2CrO_4 is +VI.

[AIIMS 2003]

8. Assertion : Copper metal gets readily corroded in an acidic aqueous solution.

Reason : Free energy change for this process is positive. [AIIMS 2004]

9. Assertion : The free gaseous Cr atom has six unpaired electrons.

Reason : Half filled 's' orbital has greater stability.

[AIIMS 2004]

10. Assertion : Fe^{2+} is paramagnetic.

Reason : Fe^{2+} contains four unpaired electrons.

11. Assertion : Transition metals are good catalysts.

Reason : V_2O_5 or Pt is used in the preparation of H_2SO_4 by contact process.

12. Assertion : Rusting of an iron is an example of corrosion.

Reason : Rusting of iron is decreased by acids and electrolytes.

13. Assertion : $AgBr$ is used in photography.

Reason : $AgBr$ undergoes photochemical reaction.

14. Assertion : Tungsten filament is used in electric bulbs.

Reason : Tungsten is a metal of high melting point.

15. Assertion : $Na_2Cr_2O_7$ is not a primary standard in volumetric analysis.

Reason : $Na_2Cr_2O_7$ is hygroscopic.

16. Assertion : Promethium is a man made element.

Reason : It is radioactive and has been prepared by artificial means.

17. Assertion : Magnetic moment values of actinides are lesser than the theoretically predicted values.

Reason : Actinide elements are strongly paramagnetic.

18. Assertion : The degree of complex formation in actinides decreases in the order $M^{4+} > MO_2^{2+} > M^{3+} > MO_2^+$.

Reason : Actinides form complexes with π -bonding ligands such as alkyl phosphines and thioethers.



19. Assertion : In transition elements ns orbital is filled up first and $(n-1)d$ afterwards, during ionization ns electrons are lost prior to $(n-1)d$ electrons.
- Reason : The effective nuclear charge felt by $(n-1)d$ electrons is higher as compared to that by ns electrons.
20. Assertion : Extraction of iron metal from iron oxide ore is carried out by heating with coke.
- Reason : The reaction $Fe_2O_3(s) \rightarrow Fe(s) + \frac{3}{2}O_2(g)$ is a spontaneous process. [AIIMS 2005]

Answers

General Characteristics

1	c	2	d	3	b	4	c	5	d
6	a	7	a	8	c	9	c	10	c
11	c	12	c	13	d	14	c	15	b
16	c	17	b	18	c	19	b	20	a
21	d	22	c	23	a	24	c	25	c
26	a	27	c	28	b	29	c	30	c
31	b	32	c	33	a	34	c	35	b
36	a	37	d	38	d	39	b	40	c
41	d	42	d	43	d	44	d	45	c
46	d	47	b	48	c	49	a	50	c
51	c	52	b	53	d	54	d	55	a
56	a	57	a	58	b	59	b	60	b
61	a	62	d	63	b	64	a	65	c
66	d	67	b	68	b	69	b	70	a
71	c	72	d	73	a	74	d	75	d
76	c	77	b	78	b	79	d	80	c
81	b	82	a	83	b	84	b	85	d
86	d	87	c	88	c	89	b	90	b
91	d	92	a	93	c	94	c	95	a
96	c	97	c	98	c	99	d	100	d
101	b	102	d	103	d	104	c	105	a
106	d	107	c	108	d	109	d	110	d
111	b	112	a	113	d	114	b	115	a
116	b	117	c	118	d	119	c	120	b
121	a	122	c	123	d	124	d	125	a
126	d	127	d	128	a	129	d	130	d
131	a	132	c	133	a	134	c	135	b
136	c	137	b	138	c	139	a	140	b
141	b	142	a	143	b	144	b	145	d

146	b	147	d	148	c	149	d	150	a
151	c	152	d	153	a	154	b	155	c
156	d	157	b	158	c	159	a	160	a
161	c	162	b	163	b	164	c	165	a
166	d	167	c	168	d	169	d	170	d
171	c	172	a	173	b	174	a	175	a
176	c	177	b	178	c	179	d	180	a
181	b	182	c	183	a	184	d	185	b
186	a	187	c	188	b	189	a	190	d
191	c	192	c	193	c	194	c	195	d
196	c	197	d	198	c	199	a	200	a
201	a	202	d	203	c	204	b	205	d
206	c	207	a	208	d	209	b	210	a
211	a	212	b	213	b	214	b	215	c
216	a	217	c	218	d	219	b	220	b
221	b	222	b	223	b	224	a	225	b
226	b	227	d	228	a	229	a	230	d
231	a	232	a	233	d	234	a	235	c
236	a	237	a	238	d	239	b	240	b
241	b								

Compounds of Transitional elements

1	a	2	c	3	d	4	a	5	b
6	c	7	a	8	b	9	b	10	d
11	a	12	a	13	a	14	a	15	a
16	a	17	a	18	d	19	b	20	b
21	d	22	c	23	e	24	b	25	c
26	d	27	d	28	a	29	c	30	c
31	a	32	d	33	b	34	d	35	a
36	a	37	b	38	b	39	b	40	a
41	a	42	c	43	c	44	a	45	a
46	b	47	c	48	b	49	a	50	c
51	b	52	b	53	c	54	d	55	c
56	a	57	c	58	d	59	b	60	d
61	d	62	b	63	c	64	a	65	c
66	b	67	c	68	a	69	a	70	c
71	a	72	a	73	a	74	a	75	a
76	d	77	d	78	d	79	a	80	c
81	b	82	d	83	c	84	d	85	b
86	a	87	c	88	c	89	b	90	a
91	b	92	c	93	d	94	c	95	e
96	b	97	b	98	d	99	c	100	c
101	b	102	a	103	a	104	d	105	c

106	a	107	c	108	d	109	b	110	b
111	a	112	a	113	b	114	a	115	a
116	c	117	a	118	a	119	c	120	c
121	c	122	a	123	a	124	b	125	c
126	a	127	c	128	a	129	c	130	a
131	a	132	a	133	d	134	c	135	a
136	d	137	b	138	a	139	b	140	c
141	c	142	a	143	b	144	d	145	e
146	e	147	a	148	c	149	b	150	b
151	d	152	a	153	b	154	a	155	c
156	a	157	c	158	a	159	b	160	b
161	b	162	c	163	d	164	b	165	d
166	c	167	a	168	b	169	c	170	d
171	a	172	c	173	d	174	c	175	c
176	d	177	a	178	b	179	c	180	b
181	a	182	b	183	b	184	b	185	a
186	c	187	c	188	d	189	c	190	c
191	a	192	d	193	c	194	a	195	d
196	a	197	c	198	d	199	b	200	c
201	d	202	b	203	a	204	a	205	a
206	a	207	b	208	a	209	c	210	a
211	a	212	c	213	a	214	c	215	c
216	c	217	a	218	a	219	a	220	c
221	d	222	b	223	c	224	b	225	a
226	d	227	a	228	c	229	d	230	d
231	c	232	a	233	b	234	b	235	c
236	a	237	c	238	d	239	a	240	d
241	d	242	b	243	ab	244	a	245	bc
246	d	247	b	248	a	249	d	250	a
251	b	252	a	253	d	254	a	255	a
256	d	257	b	258	d	259	b	260	d
261	b	262	a	263	c	264	b	265	d
266	a	267	b	268	c	269	b	270	b
271	c	272	c	273	c	274	b	275	b
276	b	277	b	278	b	279	b	280	d
281	a	282	c	283	a	284	a	285	a
286	b	287	d	288	c	289	d	290	a
291	c	292	d	293	a	294	b	295	a
296	b	297	d	298	c	299	d	300	d
301	b	302	a	303	a	304	a	305	a
306	a	307	b	308	c	309	a	310	b
311	d	312	d	313	a	314	d	315	b
316	c	317	c	318	b	319	d	320	c
321	d	322	c	323	d	324	a	325	b

326	d	327	a	328	a	329	b	330	d
331	b	332	d	333	d	334	c	335	b
336	a	337	c	338	d	339	b	340	a
341	d	342	c	343	a	344	c	345	c
346	c	347	c	348	d	349	c	350	a
351	a	352	a	353	d	354	b	355	c
356	b	357	b	358	c	359	a	360	a
361	a								

Critical Thinking Questions

1	c	2	b	3	d	4	d	5	a
6	c	7	b	8	b	9	d	10	a
11	d	12	abc	13	c	14	a	15	c
16	b	17	d	18	a	19	b	20	c
21	c	22	a	23	b	24	b	25	a
26	b								

Assertion & Reason

1	e	2	b	3	c	4	e	5	a
6	c	7	a	8	d	9	c	10	a
11	b	12	c	13	b	14	a	15	a
16	a	17	b	18	b	19	a	20	d

AS Answers and Solutions

General characteristics

- (c)

	$3d^5$					$4s^1$
Cr	↑	↑	↑	↑	↑	↑
Cr ⁺	↑	↑	↑	↑	↑	□
- (d) There are 6 electrons in its ultimate and penultimate shell.
- (b) They show variable oxidation state due to participation of ns and $(n-1)d$ electrons.
- (c) $(Cr^{+6}K_2Cr_2O_7 - \text{yellow } Cr^{+3}Cr_2(SO_4)_3 - \text{green})$
- (a) Ionic radii $\propto \frac{1}{\text{Atomic No.}}$ Ionic radius decreases from left to right in a period.
- (c) The atomic weight;
$$\text{Equivalent weight} = \frac{\text{Atomic weight}}{\text{No. of } e^- \text{ lost or gained}}$$

$$Fe^{2+} \rightarrow Fe^{3+} + e^-$$

∴ Equivalent weight = Atomic weight

10. (c) Gold; $[Xe]5d^{10}6s^1$.
11. (c) $(n-1)d$ and ns orbits.
12. (c) d -block elements; because
(i) Small atomic size
(ii) High nuclear charge
(iii) Presence of vacant d -orbitals
13. (d) Transitional elements form coloured salts due to the presence of unpaired electrons in d -orbital.
14. (c) Cu ; because last electron enters d -orbital $(3d^{10}4s^1)$.
15. (b) Cu due to the presence of vacant d -orbital.
17. (b) Nickel; $Ni + 4CO \rightarrow [Ni(CO)_4]$ (volatile)
18. (c) Copper, silver and gold; all the three were used for making coins.
19. (b) 2, 8, 18, 1 = Cu
23. (a) In between s and p -block elements.
25. (c) Fe^{+3}


S. No.	Outer configuration	No. of unpaired e	Colour of ion	Magnetic moment
V^{+3}	$3d^2$	2	Green	2.76
Mn^{+3}	$3d^4$	4	Violet	1.9
Fe^{+3}	$3d^5$	5	Yellow	5.96
Cu^{+2}	$3d^9$	1	Blue	1.9

26. (a) Misch metal is an alloy of rare earth metals with composition :
Rare earth metals – 94.95%
Iron (Fe) – 5%
 S, C, Ca, Al, \dots – Traces
27. (c) "All their ions are colourless" this sentence is false because they are 90% coloured and only few are colourless.
28. (b) $1s^2, 2s^2 2p^6, \dots, ns^2 p^6 d^3, (n+1)s^2$ as last electron enters d -subshell.
30. (c) Due to unpaired d -electrons.
31. (b) $Fe^{+2} - 3d^6 4s^0 - 4$ unpaired e^- .
32. (c) All the oxides of $Fe(FeO, Fe_2O_3$ and $Fe_3O_4)$ are basic in nature.
34. (c) The presence of one or more unpaired electrons in the system.
35. (b) They show multiple oxidation state due to availability of vacant d -orbitals.
They are coloured due to $d-d$ transition.
36. (a) $Mn^{2+} - 5$ unpaired electrons.
38. (d) Iron belongs to group VIII B of the periodic table.
39. (b) d -orbital is complete; $Zn - 3d^{10} 4s^2$
41. (d) Transition elements form co-ordinate compounds because of
(i) High nuclear charge
(ii) Small size
(iii) Vacant d -orbital
42. (d) Hg is a good conductor of electricity.
45. (c) Transition metals show variable valency.
46. (d) Cu^+ do not have any unpaired electron.
47. (b) Fe^{2+} ion have 4 unpaired electrons
 $Fe^{2+} = 26 - 2 = 24 = 1s^2 2s^2 2p^6 3s^2 3p^6 3d^6$

48. (c) Hydrogenation – because they have tendency to occlude hydrogen on free surface.
49. (a) Metals contribute their valency electrons to the common sea of electrons.
52. (b) They are inert towards many common reagents.
53. (d) Oxidation state of iron in mohl's salt
 $FeSO_4(NH_4)_2SO_4.6H_2O$ is + 2.
54. (d) They show variable oxidation states due to participation of $(n-1) d$ -orbitals electrons with ns orbital electrons.
55. (a) Electronic configuration of chromium
 $Cr \rightarrow [Ar]3d^5 4s^1$
 $Cr^{2+} \rightarrow [Ar]3d^4 4s^0$.
56. (a) Covalent bond is constituted by electrons of d -orbitals and lusture is due to free electrons of s -orbital in metallic bond.
57. (a) Cr has highest M.P. and B.P. due to maximum no. of unpaired electrons.
58. (b) Hg as there is no unpaired electron so M.P. and B.P are low. Hg is therefore liquid at room temperature with 234K.
60. (b) Zn due to increased shielding effect the attraction of electrons towards nucleus decreases.
61. (a) Number of electrons in excited state
 $X^{+3} = 18 + 4 = 22$
Number of electrons in ground state
 $X = 22 + 3 = 25$.
62. (d) $(n-1)s^2 p^6 d^{1-10} ns^1$ or ns^2
64. (a) Ni^{2+} and Cr^{2+} are coloured. But Zn^{2+} is colourless because of absence of unpaired e^- .
66. (d) They show variable valency due to presence of vacant d -orbitals.
67. (b) Maximum oxidation state = 6
Maximum no. of e^- in last shell = 6
∴ Group is VI-B.
76. (c) Ag belongs to second ($4d$) transition series remaining all are in first transition series.
77. (b) Fe^{+2} ion have 4 unpaired electron so it is paramagnetic.
78. (b) $_{30}Zn$ and $_{80}Hg$ have their d orbitals completely filled so they do not show any variable valency.
80. (c) d -block elements are known as transition elements. These show variable valency due to their incomplete d -subshell.
81. (b) Electronic configuration of $_{27}Co$ -
 $1s^2 2s^2 2p^6 3s^2 3p^6 3d^7 4s^2$,
hence unpaired $e^- = 3$
82. (a) $_{30}Zn$ has been placed in II B group of d -block in the long form of periodic table.
83. (b) The electronic configuration of Zn is $(Ar)3d^{10} 4s^2$. Hence due to complete d -subshell, it does not show variable valency.
84. (b) $Zn^{+2} - 3d^{10}$
no unpaired electrons. Hence, diamagnetic in nature.
85. (d) Terbium is lanthanide as it belongs to $4f$ -series having configuration $[Xe]4s^9 6s^2$. However the remaining members belong to $5f$ -series (actinides).
87. (c) Fe^{+2} and Ni^{+2} both.
89. (b) $Ti^{+4} \rightarrow 3d^0 4s^0$ ∴ no unpaired e^- .
91. (d) Transition metal as its last electron enters d -orbital.

92. (a) $d^5 s^1$
- | | |
|---|---|
| d | s |
| | |
- This configuration is more stable because of symmetrical distribution of e and exchange energy.

93. (c) Among the transition metals Mn forms maximum no. of oxides.



94. (c) Due to d^5 configuration, metallic bonds are weak. d^5 orbital is half filled as a result $3d$ electrons are more tightly held by the nucleus and this reduces the de-localization of electrons resulting in weaker metallic bonding.

95. (a) Cu^{+2}

S.No	Ion	Electronic configuration	No. of unpaired electrons
(i)	Cu^{+2}	d^9	1
(ii)	Ni^{+2}	d^8	2
(iii)	Co^{+2}	d^7	3
(iv)	Fe^{+2}	d^6	4

$$\mu = \sqrt{n(n+2)} \text{ (magnetic moment)}$$

$$\mu \propto \sqrt{n}$$

Cu^{+2} there is only 1 unpaired electron so its magnetic moment is least.

96. (c) In the first transition series $Mn(3d^5 4s^2)$ shows the maximum oxidation state of +7.

97. (c) They crystallize with body centered cubic and hexagonal close packed structure.

99. (d) Carrying unpaired electrons.



102. (d) All are transition elements and form complex ion.

103. (d) Ni and Co are used as catalyst.

104. (c) Magnetic moment depend upon the no. of unpaired electrons.

105. (a) Cr has 6 unpaired electrons.



106. (d) Europium is a f -block elements as it follows the general electronic configuration of the f -block elements



107. (c) 70% Cu and 30% Zn are mixed to form brass which is used in making utensils, artificial jewelry.

108. (d) Strength of metallic bond depends upon number of unpaired electrons. As number of unpaired electrons increases, the bond strength increases. So Cr, Mo, W show stronger bonding due to maximum number of unpaired electron.

109. (d) Zn^{+2} as there are no unpaired electrons.



110. (d) Cobalt is used in cancer therapy.

111. (b) Cu is oxidised which turns the solution blue.

113. (d) $Zn^{+2} - 3d^{10} 4s^0$ so there are no unpaired electrons.

114. (b) $Sc - 21 \rightarrow 3d^1 4s^2$


116. (b) $3d^5$ as this configuration corresponds to maximum number of unpaired electrons.

121. (a) Hg because it lies below H_2 in electrochemical series and thus cannot reduce it.

122. (c) High charge/size - ratio and vacant d -orbitals.


123. (d) They have one or more unpaired d - electron.

125. (a) Fe because it easily gets oxidised in moist air.



126. (d) Pt because it is a noble metal and does not react with air, water or acid at room temperature.

128. (a) Iron:




Therefore, Fe is most ferromagnetic as it has maximum number of unpaired electrons.


129. (d) The transition metals form a large number of interstitial compounds in which small atoms like hydrogen, carbon, boron and nitrogen occupy interstitial sites in their lattices.

130. (d) Because Pt is a noble metal.

131. (a) $Zn - 3d^{10} 4s^2$



132. (c) $Ti \rightarrow 3d^2 4s^2$



133. (a) Atomic no. 58 to 71 are rare earth metals.

Lanthanides

134. (c) 58 to 71 and 90 to 103 (Lanthanides & actinides).

136. (c) To form complex compounds.

137. (b) Cu as it comes after H in electrochemical series.

139. (a) Their d -orbitals are completely filled.

140. (b) Cu as it comes after H in electrochemical series. It has positive standard reduction potential thus does not provide electrons for reduction.

144. (b) $6Hg + O_3 \rightarrow 3Hg_2O$

Mercurous oxide

During this reaction, mercury loses its meniscus and starts sticking glass.

145. (d) Ga, In, Tl ; they belong to p -block.


148. (c) Zn, Cd and Hg are non typical transition elements because they have complete d -orbitals.

150. (a) In Cr^{3+} number of unpaired $e^- = 3$. A electronic configuration of $Cr^{3+} = 1s^2 2s^2 2p^6 3s^2 3p^6 3d^3$.


151. (c) Zinc does not have any unpaired electron pair so it forms only colourless compound.

153. (a) VII B groups as the metal must contain 7 electrons in ultimate shell in order to show +7 oxidation state.

154. (b)



155. (c) Ti^{+3} is paramagnetic due to the presence of an unpaired electron.



156. (d) At 350K and 1 atm pressure 1 unit volume of Pd absorbs 900 unit volume of H_2 .

157. (b) $Cu^{2+} \rightarrow 3d^9 4s^0$

1 unpaired e^- .

159. (a) Mercury does not give H_2 on reacting with water because its ionisation energy is so much high.

160. (a) Removal of electron is easier in f -block elements due to more shielding.

161. (c) Transition metal show variable valency due to presence of vacant d -orbitals.

163. (b) $Fe^{2+} - 1s^2 2s^2 p^6 3s^2 p^6 d^6$.

164. (c) It has 6 electrons in $3d$ orbital.

165. (a) They form oxide readily.

166. (d) Cr^{+3} due to presence of unpaired electrons.

169. (d) Fe^{+2} due to presence of 4 unpaired electrons.

170. (d) Fe because it belongs to transition series.

171. (c) Lanthanide contraction takes place.

172. (a) Variable oxidation states give free valencies.

173. (b) $Mn = +2, +3, +4, +5, +6, +7$

175. (a) Zn due to no unpaired electron in d -orbital.

176. (c) $1s^2, 2s^2 p^6, 3s^2 p^6 d^2, 4s^2$.

179. (d) $\mu = \sqrt{n(n+2)}$ (μ = magnetic moment)

(n = no. of unpaired electron).

$$2.83 = \sqrt{n(n+2)}$$

$$n(n+2) = 8$$

$$n^2 + 2n - 8 = 0$$

$$n = 2.$$

180. (a) $Cr^{+++} \rightarrow$ green

$Fe^{+++} \rightarrow$ Pale-green

181. (b) Inner transition elements means f -block element, they have three incomplete outer orbitals.

184. (d) $3d$
- $$\boxed{\uparrow\downarrow} \quad \boxed{\uparrow\downarrow} \quad \boxed{\uparrow\downarrow} \quad \boxed{\uparrow\downarrow} \quad \boxed{\uparrow} \quad 1 \text{ unpaired } e^- \text{ only.}$$

185. (b) $d-d$ transition of $3d$ electrons.

186. (a) $K_4[Fe(CN)_6]$

$$4 + x - 6 = 0$$

$$x = 6 - 4$$

$$x = +2.$$

188. (b) $Mn^{2+} - 3d^5 \rightarrow$ unpaired e^- .

189. (a) Very high ionisation energy and weak metallic bond.

191. (c) Iron because mercury does not form amalgam with iron.

192. (c) Chromium gives protective and decorative coating to the base metal.

193. (c) d -block elements as last electron enters in d -orbital.

195. (d) Highest oxidation state \rightarrow no. of $s-e^- +$ no. of $d-e^-$
 $\frac{2}{2} + \frac{2}{2} = 4$

196. (c) Ionic and covalent compounds.

198. (c) $[Kr]4d^{10}5s^1$ (atomic no. = 47)

199. (a) Magnese is stronger oxidising agent in + 7 oxidising state. e.g. $KMnO_4$


200. (a) $Cu^+ - 3d^{10}4s^0$; no unpaired e^- .

201. (a) $Fe-CO-Ni$. With the increase in the d -electrons, screening effect increases, this counter balances the increased nuclear charge due to increase in atomic number. As a result atomic radii remain practically same after chromium.

202. (d) Ta because it is non-corrosive.

205. (d) Cobalt due to presence of unpaired e^- .

206. (c) $Ti^{3+} \rightarrow 3d^1 4s^0$; $Sc^{3+} \rightarrow 3d^0$



In Mn^{2+} number of unpaired $d e^- = 5$. So it has maximum magnetic moment according to the formula. $\mu = \sqrt{n(n+2)}$

207. (a) + 4 oxidation state of cerium is also known in solution.

208. (d) $(n-1)d$ ns
- $$\boxed{\uparrow} \quad \boxed{\uparrow} \quad \boxed{\uparrow} \quad \boxed{\uparrow} \quad \boxed{\uparrow} \quad \boxed{\uparrow\downarrow}$$
- $(n-1)d^5 ns^2$ can achieve the maximum oxidation state of + 7.

209. (b) $Ti_{22} = 3d^2 4s^2$; $Ti^{2+} = 3d^2$

$$V_{23} = 3d^3 4s^2; V^{2+} = 3d^2$$

$$Cr_{24} = 3d^4 4s^2; Cr^{4+} = 3d^2$$

$$Mn_{25} = 3d^5 4s^2; Mn^{5+} = 3d^2$$

210. (a) As sixth period can accommodate only 18 element in the table, 14 member of HF series (atomic number 58 to 71) are separately accommodated in a horizontal row below the periodic table. These are called as lanthanides.

212. (b) The oxidation state in both (lanthanide and actinide) is +3. The property of actinide are very similar to those of lanthanide when both are in +3 state.

213. (b) The lanthanide contraction relates to atomic as well as M^{3+} radii in which the regular decrease in the size of lanthanoid ion from La^{3+} to Lu^{3+} are found.



214. (b) Highest magnetic moment depends upon number of unpaired electron since

$$Cr^{2+} = 3d^4 4s^0, Mn^{2+} = 3d^5 4s^0$$

$$Cu^{2+} = 3d^9 4s^0, Co^{2+} = 3d^7 4s^0, Ni^{2+} = 3d^6 4s^0$$


So Mn^{2+} contain maximum number of unpaired electron i.e. 5.

215. (c) Cobalt 27 belong to $3d$ transition series having in complete $3d$ orbitals i.e., $3d^7$.
216. (a) It is the Tata iron and steel company.
217. (c) The atomic weight of Co, Ni and Fe are 58.90, 58.60, 55.85 respectively. Therefore $Co > Ni > Fe$ is the correct sequence of atomic weight.
218. (d) The first ionization energies of Ti, V, Cr and Mn are 656, 650, 652 and 717 $kJ/mole$ respectively. I.E. increase in a period from $L \rightarrow R$ hence, manganese has maximum first ionisation potential.
219. (b) Metal M belongs to d -block. Its electronic configuration can be arranged as, $1s^2 2s^2 2p^6 3s^2 3p^6 4s^1 3d^{10}$.
220. (b) We know that transition element are those element which have partially filled d -subshell in their elementary form. Therefore, the general electronic configuration of d -block element is $(n-1)d^{1-10} ns^{1-2}$.
221. (b) The element with atomic no. 105 is dubnium. In IUPAC nomenclature, it is known as un-nil pentin.
222. (b) The compound which have the unpaired electron show the paramagnetic property.
223. (b) Among the given choice Mn^{2+} and Fe^{3+} involve isoelectronic ions.
224. (a) Elements or ions containing unpaired electrons are paramagnetic.



Because Ni^{2+} have 2 unpaired electrons in $3d$ subshell therefore it is paramagnetic.

225. (b) $Cr(Z = 24)$ has electronic configuration $1s^2 2s^2 2p^6 3s^2 3p^6 3d^5 4s^1$ (\because half filled d orbital is more stable than incompletely filled d orbital)
226. (b) The expected electronic configuration of Cu (29) is $[Ar] 3d^9 4s^2$ but actually it is found to be $[Ar] 3d^{10} 4s^1$. This is because fully filled d orbitals are more stable than incompletely filled d -orbitals. So there is a migration of one e^- , from $4s$ orbital to $3d$ orbital to give a more stable configuration.
227. (d) $Ce - 58$ have configuration $1s^2, 2s^2 2p^6, 3s^2 3p^6 3d^{10}, 4s^2 4p^6 4d^{10} 4f^2, 5s^2 5p^6 5d^0, 6s^2$
Since, its last electron enter in f sub-shell, therefore it is a member of f -block.
228. (a) $Ni \rightarrow 1s^2 2s^2 2p^6 3s^2 3p^6 3d^8 4s^2$
 $Ni^{2+} \rightarrow 1s^2 2s^2 2p^6 3s^2 3p^6 3d^8$
 $3d^8 = 10 - 2 = 2$ unpaired electron.
230. (d) ${}_{25}Mn = 3d^5 4s^2$
After losing two electron electronic configuration will be like this (${}_{25}Mn^{++} 3d$) and this is most stable configuration due to half filled orbitals hence third ionization enthalpy will be highest in this case.
231. (a) ${}_{21}Sc \rightarrow 1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^1$






Due to presence of paired electron it will be colourless.

232. (a) Lanthanum is the first element of Lanthanide series so size decreases with increase in atomic number so La has the largest atomic radii.
234. (a) Tl ions are more stable than Tl and thus, Tl ions change to Tl ions thereby acting as oxidising agents.
 Tl compounds + $2e \rightarrow Tl$ compounds
(less stable oxidising agent) (more stable oxidising agent)
237. (a) $Fe \rightarrow [Ar] 3d 4s$, number of unpaired electrons = 5
 $Fe \rightarrow [Ar] 3d 4s$, number of unpaired electrons = 4
 $Co \rightarrow [Ar] 3d 4s$, number of unpaired electrons = 3
 $Co \rightarrow [Ar] 3d 4s$, number of unpaired electrons = 4
238. (d) Paramagnetic character is actually due to presence of unpaired electrons.
239. (b) Mercury has the property of dissolving nearly all metals, forming liquid or solid solutions called amalgams. It amalgamates well with gold, silver and tin, but does not dissolve iron or platinum. Presence of these may result in sickening or flouring of the mercury.
240. (b) Ce -lanthanide, Cs -alkali metal, Cf -actinide, Ca -alkaline earth metal.
241. (b) Where $n =$ number of unpaired electron
For $Sc^{3+} = 3d^0, n = 0, \therefore \mu = 0$

Compounds of Transitional elements

1. (a) MnO_2, MnO_2, Mn^{2+}
In neutral medium :
 $2KMnO_4 + 3MnSO_4 + 2H_2O \rightarrow K_2SO_4 + 2H_2SO_4 + 5MnO_2$
In alkaline medium :
 $2KMnO_4 + H_2O \rightarrow 2MnO_2 + 2KOH + 3O$
In acidic medium :
 $2KMnO_4 + 3H_2SO_4 \rightarrow K_2SO_4 + 2MnSO_4^{+2} + 5H_2O + 5O$
2. (c) Oxidation state of Mn changes from +7 to +2 in acidic medium i.e. one mole of it accepts 5 mole of electrons.
3. (d) Since it accept $6e^-$ its
Equivalent weight = $\frac{M}{6}$.
5. (b) Decreases from + 6 to + 3.
 $K_2Cr_2O_7 + 4H_2SO_4 \rightarrow K_2SO_4 + Cr_2(SO_4)_3 + 4H_2O + 3O$
 $[H_2S + [O] \rightarrow S + H_2O] \times 3$
 $K_2Cr_2O_7 + 4H_2SO_4 + 3H_2S \rightarrow K_2SO_4 + Cr_2(SO_4)_3 + 7H_2O + 3S$
6. (c) $FeSO_4$ is oxidised and $KMnO_4$ is reduced.
 $2KMnO_4 + 3H_2SO_4 \rightarrow K_2SO_4 + 2MnSO_4$
 $[2FeSO_4 + H_2SO_4 \rightarrow Fe_2(SO_4)_3 + 2H] \times 5$




In this reaction oxidation state of *Mn* is changing from +7 to +2 while oxidation state of *Fe* is changing from +2 to +3.

7. (a) $Hg_2Cl_2 + 2NH_4OH \rightarrow NH_4Cl + 2H_2O + Hg + HgNH_2Cl$
Calomel
8. (b) Ag^+ forms a complex ion with NH_3
 $AgCl + 2NH_3 \rightarrow [Ag(NH_3)_2]Cl$
9. (b) Four water molecules.
 $CuSO_4 \cdot 5H_2O$ is a crystalline salt. Four H_2O molecules attach to copper forming a square planar symmetry and two oxygen atoms from SO_4^{2-} ion complete the distorted octahedron. The fifth water molecule is attached through hydrogen bonding between one of the co-ordinated water molecule and one of the sulphate ion.
10. (d) $AgCl$ is a covalent compound hence it is insoluble in water also it form complex with NH_4OH , thus is soluble in NH_4OH .
 $AgCl + 2NH_4OH \rightarrow [Ag(NH_3)_2]Cl + H_2O$
11. (a) Basic copper acetate.
(Verdigris - $(CH_3COO)_2Cu \cdot Cu(OH)_2$) is a blue green powder used in green pigment and in dyes. Also in manufacture of insecticides and fungicides.
13. (a) H_2O_2 reduces acidified $KMnO_4$ solution. As a result. The pink colour of $KMnO_4$ is changed.
14. (a) MnO_2 forms amphoteric oxide due to intermediate oxidation state.
15. (a) MnO is ionic due to lower oxidation state.
17. (a) Manufacture of blue black ink.
18. (d) As fertilizer because it is not required by plants.
20. (b) Since *Ag* is less reactive than *Cu* therefore it does not displace *Cu* from $CuSO_4$ while other metals are more reactive, they displace *Cu* from $CuSO_4$.
21. (d) Its reduction to metallic silver.
 $2AgNO_3 \rightarrow 2Ag + N_2 + 3O_2$
black stain
22. (c) $Na_2S_2O_3 + CuSO_4 \rightarrow NaCuS_2O_3$
23. (e) $2HgO \xrightarrow{\Delta} 2Hg + O_2$
24. (b) $2Fe + 3Cl_2 \rightarrow 2FeCl_3$
25. (c) $Fe(OH)SO_4$
 $FeSO_4 \xrightarrow{H_2O} Fe(OH)SO_4$
27. (d) $\frac{1}{5} \times$ molecular weight of $KMnO_4$
as transfer of $5e^-$ takes place when $KMnO_4$ acts as oxidant in acidic medium.
 $2KMnO_4 + 3H_2SO_4 \rightarrow K_2SO_4 + 2MnSO_4 + 3H_2O + 5O$
28. (a) Oxidation number of *Cr* in options *a*, *b*, *c* and *d* are +6, +4, +3, +3 respectively.

In given options (a) has high oxidation number therefore its radii will be low. Atomic radii decreases with increase in oxidation no.

33. (b) Cuprous chloride.
 $Cu(s) + 2HCl + CuSO_4 \rightarrow 2CuCl + H_2SO_4$
Cuprous chloride
34. (d) Cuprous chloride slowly oxidises to green basic cupric chloride.
 $CuCl \xrightarrow{\text{air}} 3CuO \cdot CuCl_2 \cdot 3H_2O$
green coloured
35. (a) $2CuCl + 2HCl + [O] \rightarrow 2CuCl_2 + H_2O$
38. (b) Equivalent weight of $KMnO_4$ in acidic medium is $M/5$
 \therefore Equivalent weight = $\frac{158}{5} = 31.6$
39. (b) In acidic medium, $KMnO_4$ gives 5 oxygen while acidic $K_2Cr_2O_7$ gives 3 oxygen.
41. (a) Ag_2O is mild oxidising agent as greater the oxidation number of metal stronger oxidising agent.
42. (c) $K_2Cr_2O_7 + 3H_2SO_4 \rightarrow K_2SO_4 + Cr_2(SO_4)_3 + 3(O) + 3H_2O$
No. of electrons lossed = $12 - 6 = 6$
 \therefore Equivalent weight = $\frac{M}{6} = \frac{294}{6} = 49$.
44. (a) $ZnSO_4 \rightarrow Zn^{++} + SO_4^{--}$
 $Zn^{++} \approx Cu^{++}$
 $Cu^{+2} \rightarrow 3d^9 - 1 \text{ unpaired } e^-$
 \therefore paramagnetic in nature.
46. (b) $KMnO_4$ is first reduced to manganate and then to insoluble manganese dioxide. Colour changes first from purple to green and finally becomes colourless.
 $2KMnO_7 + 2KOH \rightarrow 2K_2MnO_4 + H_2O + O$
 $2K_2MnO_4 + 2H_2O \rightarrow 2MnO_2 + 4KOH + 2O$
 $2KMnO_2 + H_2O \xrightarrow{\text{alkaline}} 2MnO_2 + 2KOH + 3[O]$
47. (c) $2KMnO_4 + 3H_2SO_4 + 5C_2H_2O_4 \rightarrow$
 $2Mn^{2+} + 8H_2O + 10CO_2$
48. (b) $2KI + HgI_2 \rightarrow \underbrace{K_2HgI_4}_{\text{Nessler's reagent}} + 2KI$
50. (c) $Cr_2O_7^{2-} + 8H^+ + 2SO_3^{2-} \rightarrow 2Cr^{+3} + 3SO_4^{2-} + 4H_2O$
51. (b) $KI + MnO_4^- \rightarrow K^+ IO_3^- + Mn^{+2}$
52. (b) Among all the reactions given $CuSO_4$ does not react with KCl to give Cu_2Cl_2 .
53. (c) Mohr's salt.
 $2KMnO_4 + 8H_2SO_4 + 10FeSO_4 \rightarrow$
 $K_2SO_4 + 2MnSO_4 + 5Fe_2(SO_4)_3 + 8H_2O$
Mohr's salt decolourises $KMnO_4$ by reducing Mn^{+7} ions to Mn^{+2} ions.
55. (c) Amalgams are alloys which contain mercury as one of the contents.
56. (a) In order to make the image permanent, it is necessary to remove the unreduced silver bromide from the surface of the developed film. This operation is called fixing of image. Fixing is

done by dipping the developed film or plate in sodium thiosulphate (hypo) solution. the hypo solution dissolves the unreduced silver bromide by forming a complex.




Thus sodium thiosulphate acts as a complexing agent.

58. (d) All of these.
59. (b) $Fe^{+3} - 3d^5 - 5$ electrons unpaired.
 Fe will be attracted in magnetic field so will show increase in weight.
60. (d) TiF_6^{2-} and Cu_2Cl_2 due to absence of unpaired electrons.
62. (b) Copper oxide; $2Cu(NO_3)_2 \rightarrow 2CuO + 4NO_2 + O_2$
63. (c) Cuprous iodide is precipitated with liberation of iodine.
 $2KI + CuSO_4 \rightarrow CuI_2 + K_2SO_4 + I_2$
64. (a) It is one third of its molecular weight in alkaline medium because it gives 3 nascent oxygen in alkaline medium.
 $2KMnO_4 + H_2O \xrightarrow{+3e^-} 2KOH + 2MnO_2 + 3[O]$
 \therefore Equivalent weight = $\frac{M}{3}$
65. (c) $NaCl + H_2SO_4 + K_2Cr_2O_7 \rightarrow$
 $CrO_2Cl_2 + K_2SO_4 + Na_2SO_4$
Chromyl chloride
66. (b) Decomposes in sunlight.
 $2AgNO_3 \xrightarrow{\Delta} 2Ag + 2NO_2 + O_2$
67. (c) Silver; $AgNO_3 + NaCl \rightarrow AgCl + NaNO_3$
White ppt.
70. (c) The formation and thermal decomposition of $Ni(CO)_4$.
71. (a) $CuSO_4 + 4NH_3 \rightarrow [Cu(NH_3)_4]^{++} SO_4^{--}$
72. (a) Equivalent wt. = $\frac{\text{molecular wt.}}{\text{total no. of } e^- \text{ gained or lost}} = \frac{M}{1} = M$
76. (d) $HgI_2 + 2KI \rightarrow K_2[HgI_4] \rightleftharpoons 2K^+ + [HgI_4]^{--}$
80. (c) The conversion of dichromate to chromate.
 $K_2Cr_2O_7 + 2KOH \rightarrow \underbrace{2K_2CrO_4}_{\text{yellow}} + H_2O$
81. (b) $MnO_2 + KOH \rightarrow K_2MnO_4$
pyrolusite
83. (c) $K_2Cr_2O_7 + H_2SO_4 + 4H_2O_2 \rightarrow K_2SO_4 + 2CrO_5 + 5H_2O$
84. (d) Iron; $Fe + H_2O/H^+ \rightarrow Fe_2O_3 \cdot xH_2O$
85. (b) Na_2CdCl_4 - no unpaired electrons.
87. (c) $CuCN$; $3d^{10}$ $4s^0$
 Cu^+

↑↓	↑↓	↑↓	↑↓	↑↓	
----	----	----	----	----	--

as all the electrons are paired. It is expected to be colourless.
88. (c) $ZnSO_4, MgSO_4$ are isomorphous i.e. having same structure.
90. (a) TiO_2 because of 3d state i.e. no unpaired electrons.
91. (b) $CoO \rightarrow Co^{+2}$ is blue colour.
92. (c) $Ca_2P_2O_7 \rightarrow 2Ca^{++} + (P_2O_7)^{4-}$ $Fe^{+3} + (P_2O_7)^{4-} \rightarrow Fe_4(P_2O_7)_3$
93. (d) AgI because of high covalent character along with strong Vander Waal's attraction and electrostatic attraction between silver and iodide ions.

100. (c) $ZnO + 2NaOH \rightarrow Na_2ZnO_2 + H_2O$
Sodium zincate
101. (b) $HgCl_2$ compound is easily volatile. They are insoluble in water and soluble in acids.
102. (a) Colourless compounds are those which have no unpaired electrons and paramagnetic substance do have unpaired electrons. Therefore paramagnetic substance possess colour.
107. (c) $MnSO_4 \rightarrow Mn^{2+} + SO_4^{2-}$



due to presence of unpaired electrons it will form coloured salt.

111. (a) Cast iron : iron -93-95%
Carbon - 2.5-5%, impurities about 2%.
112. (a) $FeCl_3$ is a salt of strong acid and weak base. It gives $Fe(OH)_3$ and HCl on hydrolysis. $Fe(OH)_3$ is a weak base and HCl is strong acid.

So the aqueous solution of $FeCl_3$ will be acidic in nature

113. (b) $Cr_2^{6+} + 6e^- \rightarrow 2Cr^{3+}$; $Fe^{2+} \rightarrow Fe^{3+} + e^-$
114. (a) CrO_3 and Mn_2O_7 are acidic oxide since they react with water to form acid.
 $CrO_3 + H_2O \rightarrow H_2CrO_4$; $Mn_2O_7 + H_2O \rightarrow 2HMnO_4$
Chromic acid Permanganic acid

115. (a) In photography as it is sensitive towards light.
116. (c) $[MnCl_4]^{2-} \rightarrow$

↑	↑	↑	↑	↑
---	---	---	---	---

 $4s$ $4p$

↑↓	↑↓	↑↓
----	----	----

Number of unpaired electron = 5
 $[CoCl_4]^{2-} \rightarrow$

↑↓	↑↓	↑	↑	↑
----	----	---	---	---

 $4s$ $4p$

↑↓	↑↓	↑↓
----	----	----

Number of unpaired electron = 3
 $[Fe(CN)_6]^{4-} \rightarrow$

↑↓	↑↓	↑↓	↑↓	↑↓
----	----	----	----	----

 $4s$ $4p$

↑↓	↑↓	↑↓
----	----	----

Number of unpaired electron = 0
Magnetic moment = $\sqrt{n(n+2)}$ dsp

Where, n = number of unpaired electron i.e., greater the number of unpaired electron greater will be the paramagnetic character.

117. (a) In the compound $[Ni(CO)_4]$
Oxidation number of $Ni = 0$
Co-ordination number of $Ni = 4$
(Co (carbonyl) is a strong ligand so it cause pairing of electron)
 $Ni[Ar]$

↑↓	↑↓	↑↓	↑	↑
----	----	----	---	---

 $4s$ $4p$

↑↓	↑↓	↑↓
----	----	----

 Ni in $[Ni(CO)_4]$

↑↓	↑↓	↑↓	↑↓	↑↓
----	----	----	----	----

 $4s$ $4p$

--	--	--	--	--

 $[Ni(CO)_4]$

↑↓	↑↓	↑↓	↑↓	↑↓
----	----	----	----	----

 $4s$ $4p$

↑↓	↑↓	↑↓	↑↓
----	----	----	----

 $[Ar]$

↑↓	↑↓	↑↓	↑↓
----	----	----	----

 CO CO CO CO
 sp^3 Hybridization

118. (a) Applying the oxidation number rule in $[Fe(H_2O)_5(NO)]SO_4$
($\therefore H_2O$ and NO are neutral)

$$[x + (0) \times 5 + 0] + (-2) = 0$$

$$x + 0 + 0 - 2 = 0 \Rightarrow x = +2$$

Hence, oxidation number of Fe is +2.



119. (c) $Zn + NaOH \rightarrow Na_2ZnO_2 + H_2$
Conc. Sodium zincate Hydrogen

Thus in this reaction hydrogen gas is produced when zinc is treated with concentrated $NaOH$.

120. (c) $Fe_2O_3 + 6NaOH \rightarrow 2Fe(OH)_3 \downarrow + 3Na_2O$
Brown (insoluble in $NaOH$)

121. (c) Mercurous chloride are insoluble in water while rest are soluble in water.

122. (a) ZnO is an amphoteric oxide,






123. (a) Fe^{3+} ion has $[Ar] 3d^5$ configuration hence number of unpaired electron is 5.

124. (b) $FeS + H_2SO_4 \rightarrow FeSO_4 + H_2S$

125. (c) In this complex CO^{2+} ion have 3 unpaired electron so spin only magnetic moment will be $\sqrt{3(3+2)}$ i.e., $\sqrt{15}$ B.M.

126. (a) Platinum acts as catalyst in the oxidation of ammonia to form nitric oxide. This reaction is used in the Ostwald's method of nitric acid preparation




127. (c) Iron is oxidised to ferrous nitrate and nitric acid is changed to ammonium nitrate




128. (a) $CrO_3 + 2NaOH \rightarrow Na_2CrO_4 + H_2O$
Yellow solution

129. (c) $2KI + CuSO_4 \rightarrow CuI_2 + K_2SO_4$
Unstable



Hence, solution contains Cu_2I_2 , I_2 and K_2SO_4 .


130. (a) Cu is placed above Ag in electrochemical series, hence it can replace Ag from its salts solution. Therefore the reaction occurs as follows,




131. (a) When the quenched steel is heated to temperature below red hot and then allowed to cool slowly, it becomes soft. This process is known as annealing.

132. (a) We know that ammonia the order of solubility is $AgCl > AgBr > AgI$. Therefore, $AgCl$ is more soluble in ammonia.

133. (d) In alkaline medium, $KMnO_4$ first reduced in manganate & then in insoluble manganese dioxide.



In acidic medium, Manganous sulphate formed



134. (c) $(NH_4)_2Cr_2O_7 \xrightarrow{\Delta} 2K_2CrO_4 + Cr_2O_3 + \frac{3}{2}O_2$

135. (a) HgS is soluble in aqua-regia and it is insoluble in hot dil. HNO_3 .

136. (d) $Ag_2O \xrightarrow{\Delta} 2Ag + \frac{1}{2}O_2$

138. (a) $HgCl_2$ show dimerisation. It found in dimer form.

139. (b) Iron pyrites (FeS_2) is also called 'fools gold'.

140. (c) According to Fajan, small anion is polarised to lesser extent than the larger anion. Hence AgF will be the most ionic and has high melting point.

141. (c) Potassium dichromate, on heating gives oxygen and chromic oxide (Cr_2O_3).



142. (a) Nickel is purified by Mond's process



143. (b) Stainless steel contains 11.5% Cr and 2.0% Ni with Fe .

144. (d) German silver is an alloy of copper not silver containing $Cu = 56.0\%$, $Zn = 24.0\%$ and $Ni = 20.0\%$.

145. (e) This scheme take place in Van Arkel process by this process ultrapure metal is prepared, the impure metal is first converted into a volatile stable compound generally iodide (leaving behind the impurities which is then decomposed at a higher temperature to give the pure metal. Metal like titanium, zirconium are purified by this method.


146. (e) HgS on strong heating gives Hg .

147. (a) $Cr_2O_3 \cdot 2H_2O$ is known as cruet green.

148. (c) Vanadium (III) oxide is a strong reducing agent because vanadium is electropositive metal and have high reduction potential. It has low heat of sublimation, low ionisation potential.


149. (b) Stainless steel does not rust because chromium forms an oxide layer and protect iron from rusting.

150. (b) $HgCl_2 + Na_2CO_3 \rightarrow HgCO_3 + 2NaCl$



151. (d) Ni in presence of CN^-

$[Ni^{2+}$ in presence of $CN^- = [Ar]$




As $[Ni(CN)_4]^{2-}$ has no unpaired electron. It is diamagnetic.

152. (a) The solubility of silver bromide in hyposolution due to the formation of $[Ag(S_2O_3)_2]^{3-}$.

153. (b) Brass is an alloy of Zn and Cu .

154. (a) Iodine being a strong reducing agent reduce Cu^{2+} ions to Cu^+ ions and itself gets oxidised to iodine.




155. (c) Rust is Fe_2O_3 and $Fe(OH)_3$

156. (a) ${}_{21}\text{Sc} = [\text{Ar}] 3d^1 4s^2$
 $\text{Sc}^{3+} = [\text{Ar}] 3d^0 4s^0$ no unpaired electrons in d sub shell, so it is diamagnetic and colourless.

157. (c) Zinc sulphate ($\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$) is called white vitriol. It, when heated with barium sulphide, forms a white pigment lithopone.


158. (a) Isomorphous compound are those compounds which forms same type of crystals *i.e.*, have similar structure. $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ is isomorphous with $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$.

159. (b) Colour of transition metal ion salt is due to $d-d$ transition of unpaired electrons of d -orbital. Metal ion salt having similar number of similar number of unpaired electron in d -orbitals shows similar colour in aqueous medium



Number of unpaired electrons = 1




160. (b) Zn dissolve in Conc. NaOH due to the formation of Sodium Zincate.



161. (b) CdS — Yellow
 CuS — Black
 ZnS — White
 CoS — Black

162. (c) KMnO_4 will not be oxidized further by ozone as manganese is already present in its highest possible oxidation state *i.e.* +7


163. (d) In alkaline medium



166. (c) Bronze contain $\text{Cu} = 75 - 90\%$, $\text{Sn} = 10 - 25\%$

168. (b) Brass ($\text{Cu} + \text{Zn}$)

German silver ($\text{Cu} + \text{Zn} + \text{Ni}$)



170. (d) Platforming is a process for manufacturing platinum.

171. (a) We know by reducing auric chloride by stannous chloride, the colloidal solution of gold is obtained. It is known as purple of cassium.

172. (c) List 1

List 2


- | | |
|------------------------|---------------------------|
| (i) Explosive | $\text{Pb}(\text{N}_3)_2$ |
| (ii) Artificial gem | Al_2O_3 |
| (iii) Self reduction | Cu |
| (iv) Magnetic material | Fe_3O_4 |

178. (b) $\text{Fe} + \text{Cr} + \text{Ni} = \text{Stainless steel}$
82% 11% 7%

179. (c) CaO and MgO are refractory materials. They have very high melting point.

180. (b) Brass contain $\text{Cu} = 60\%$ and $\text{Zn} = 40\%$ in its composition.

181. (a) Annealing is a process of heating steel to redness followed by slow cooling



182. (b) ${}_{26}\text{Fe}^{2+}$



+3 state is most stable because of half filled d sub-shell


183. (b) Fe and Ni .
95-97% 3-5%

185. (a) The process of producing a hard coating of iron nitride on steel is called nitriding.

187. (c) Iron loses magnetic property at curie point.


188. (d) Heat treatment alters the properties of steel due to change in the lattice structure due to differential rate of cooling.

189. (c) The passivity of iron is due to the formation of a thin insoluble and invisible iron film on surface which prevents its further reactions. The film is due to the formation of Fe_3O_4 .







191. (a) The steel obtained by this process retains its hardness but is not brittle.

192. (d) In blast furnace at 400-600°C for the smelting of iron, following takes place:-



195. (d) Iron is rendered passive by conc. HNO_3 and other oxidising agents like $\text{K}_2\text{Cr}_2\text{O}_7$, KMnO_4 , Chloric acid, chromic acid, silver nitrate etc. A specimen of passive iron can be rendered active by scratching the film mechanically or chemically.

199. (b) Fe present in Haemoglobin



In this reaction, the stronger Lewis acid SbF_5 displaces the weaker one, MnF_6 from its salt. MnF_6 is unstable and readily decomposes to give MnF_5 and fluorine.

205. (a) Case hardening : The process of hardening the surface of wrought iron by depositing a surface layer of steel on it is called case-hardening it is done by heating wrought iron in contact with potassium ferrocyanide.

Alternatively, case hardening can also be done by heating wrought iron with charcoal and then plunging it a suitable oil.


206. (a) Stainless steel is not corroded by air, moisture and small change in pH . Made up of Fe , N and Cr .
82% 7% 11%

207. (b) Variety of irons % of Carbon


Cast or Pig iron	2.5 - 4%
Wrought iron	0.12 - 0.25%
Steel	0.2 - 0.5%

208. (a)
 209. (c) Tempering : If the quenched or hardened steel is reheated to a temperature between 503 to 573 K and then allowed to cool slowly, the process is called tempering.

226. (d) % of carbon in wrought iron is 0.2 - 0.5%, in steel 0.5 - 1.5% and in Pig iron 2.5 - 4%.



230. (d) Potash alum $K_2SO_4 \cdot Al_2(SO_4)_3 \cdot 24H_2O$
231. (c) A thin layer of Fe_3O_4 is formed on the Fe metal
232. (a) $3Fe + 4H_2O \rightarrow Fe_3O_4 + 4H_2$
233. (b) Mn used to make alloy steel for armour plates, safes and helmets.
236. (a) Solder (Pb + Sn), Bronze (Cu + Sn), Brass (Cu + Zn), Bell metal (Cu + Sn).
241. (d) $Cu + Sn \rightarrow$ Bell metal
80% 20%
242. (b) Turnbull's blue Ferrous ferri cyanide $Fe_3[Fe(CN)_6]_2$
244. (a) $Cu + 2AgNO_3 \rightarrow 2Ag + Cu(NO_3)_2$
 $Cu + ZnSO_4 \rightarrow$ no reaction
 $Cu + FeSO_4 \rightarrow$ no reaction
- $$\left. \begin{aligned} E^0 Zn^{+2}/Zn &= -0.76 V \\ E^0 Cu^{2+}/Cu &= +0.34 V \\ E^0 Fe^{2+}/Fe &= -0.40 V \\ E^0 Ag^+/Ag &= +0.80 V \end{aligned} \right\}$$
- As it is clear that reduction potential of copper is more than Zn and Fe. Hence it is unable to displace them from their salts.
245. (b,c) Ferrous salts react with potassium ferricyanide to give blue colouration due to the formation of Turnbull's blue in this reaction, first ferrous salt is oxidised to ferric salt by the ferricyanide ion which itself is reduced to ferrocyanide.
- $$Fe^{+2} + [Fe(CN)_6]^{3-} \rightarrow Fe^{+3} + [Fe(CN)_6]^{4-}$$
- Ferricyanide Ferrocyanide
- $$Fe^{+3} + [Fe(CN)_6]^{4-} \rightarrow \{Fe[Fe(CN)_6]\}^-$$
- $$Fe^{+3} + [Fe(CN)_6]^{4-} + K^+ \rightarrow K\{Fe[Fe(CN)_6]\}$$
- Pot. ferricferrocyanide
or Turnbull's blue
- Ferric ions react with potassium thiocyanate to give blood red colouration due to the formation of ferric thiocyanate
- $$FeCl_3 + 3KCNS \rightarrow Fe(CNS)_3 + 3KCl$$
- Ferric thiocyanate
(Blood red)
250. (a) Electric protection : In this Mg acts as anode while iron pipe as cathode. Mg loses electrons prior to iron.
251. (b) Pyrite (FeS_2) known as fool's gold
252. (a) Stainless steel contains mainly Iron, Carbon, Nickel alongwith Chromium and Manganese.
253. (d) Firstly, carbon which is added along which crushed haematite ore is oxidised to CO (and CO_2) and second the produced CO acts as chief reducing agent for the reduction of haematite to steel.
255. (a) By white tin plating iron can be protect by water.
256. (d) $2Fe + 3CO \xrightarrow{\text{Heat}} Fe_2O_3 + 3C$
(cast iron)
257. (b) Magnetite is reduced by carbon.
258. (d) Malachite ($Cu(OH)_2 \cdot CuCO_3$) Basic copper carbonate
259. (b) + 2 is most important oxidation state
 $Cu^{+1} + e^- \rightarrow Cu; E^0 = +0.52 V$
 $Cu^{+2} + 2e^- \rightarrow Cu; E^0 = +0.34 V$
260. (d) $Cu + 4HNO_3 \rightarrow Cu(NO_3)_2 + 2NO_2 + 2H_2O$
262. (a) Metal Chloride in + 1 + 2 oxidation states
Cu CuCl Cu₂Cl₂



265. (d) $Cu + HCl \rightarrow$ no reaction
Copper is less reactive than hydrogen. Therefore it is unable to displace hydrogen from acid.
 $E_{Cu}^0 = +0.34$ and $E_H^0 = 0.00$
266. (a) Gun metal contains Cu (88%), Zn (2%), Sn(10%), Pb (0.5%)
267. (b) Solder – Sn 67% and Pb 33%.
268. (c) Brass contains Cu = 80% , Zn = 20%
German silver contains Cu = 60% , Zn = 20% , Ni = 20%
269. (b) German silver contain Cu = 60, Zn = 20, Ni = 20%
270. (b) Cu = 88%, Sn = 10%, Zn = 2%, Pb = 0.5% → Gun Metal
274. (b) $4Cu + 2H_2O \rightarrow 2Cu_2O + 2H_2$
276. (b) $2Cu + 2H_2SO_4 + O_2 \rightarrow 2CuSO_4 + 2H_2O$
277. (b) Cuprous ion (Cu^+) $3d^{10}$ Completely filled d sub shell
- $$3d^{10} \quad \begin{array}{|c|c|c|c|c|} \hline \uparrow\downarrow & \uparrow\downarrow & \uparrow\downarrow & \uparrow\downarrow & \uparrow\downarrow \\ \hline \end{array}$$
- Cupric ion Cu^{+2}
- $$3d^9 \quad \begin{array}{|c|c|c|c|c|} \hline \uparrow\downarrow & \uparrow\downarrow & \uparrow\downarrow & \uparrow\downarrow & \uparrow \\ \hline \end{array}$$
- One unpaired electron
278. (b) $CuSO_4 + K_4[Fe(CN)_6] \rightarrow$ no reaction
 $4NH_4OH + CuSO_4 \rightarrow [Cu(NH_3)_4]SO_4 + 4H_2O$
Deep blue
 $CuSO_4 + 5H_2O \rightarrow CuSO_4 \cdot 5H_2O$
Anhydrous Blue
 $4FeCl_3 + 3Na_4[Fe(CN)_6] \rightarrow Fe_4[Fe(CN)_6] + 12NaCl$
Ferric ferrocyanide
(Prussian blue)
280. (d) $CuSO_4 + 2KCN \rightarrow Cu(CN)_2 + K_2SO_4$
 $2Cu(CN)_2 \rightarrow Cu_2(CN)_2 + (CN)_2$
 $Cu_2(CN)_2 + 6KCN \rightarrow 2K_3[Cu(CN)_4]$
281. (a) $CuSO_4 + 4NH_4OH \rightarrow [Cu(NH_3)_4]SO_4 + 4H_2O$
Blue
282. (c) $Cu + 2H_2SO_4 \rightarrow CuSO_4 + SO_2 + 2H_2O$
284. (a) $2Cu + CO_2 + H_2O + O_2 \rightarrow CuCO_3 \cdot Cu(OH)_2$
Basic copper carbonate
285. (a) $2CuSO_4 + K_4(Fe(CN)_6) \rightarrow Cu_2[Fe(CN)_6] + 2K_2SO_4$
287. (d) $CuSO_4 + Hg \rightarrow$ No reaction
Hg is less reactive than Cu it comes below Cu in the reactivity series
288. (c) $Cu + H_2O \rightarrow$ No reaction
 $E_{Li^+/Li}^0 = -3.04 V$ $E_{H^+/H_2}^0 = 0.00 V$
 $E_{Ca^{+2}/Ca}^0 = -2.87 V$ $E_{Cu^+/Cu}^0 = +0.34 V$
Cu comes below H in the electrochemical series. Hence, unable to displace hydrogen from water.
289. (d) $3Cu + 8HNO_3 \rightarrow 3Cu(NO_3)_2 + 2NO + 4H_2O$
291. (c) $Cu + O_2 + CO_2 + H_2O \rightarrow Cu(OH)_2 \cdot CuCO_3$

292. (d) Orford process → During electrolytic refining of Ni from orford bottoms the Pt metal together with gold and silver collect as anode mud in concentrated form which is later processed to separate the metals.
294. (b) AgBr is used in photography because it is light sensitive.
296. (b) $4AgCl + 2Na_2CO_3 \rightarrow 4Ag + 4NaCl + 2CO_2 + O_2$
297. (d) $2AgNO_3 + K_2CrO_4 \rightarrow Ag_2CrO_4 + 2KNO_3$
(Red)
298. (c) $3Ag + 4HNO_3 \xrightarrow[\text{dilute}]{\text{heat}} 3AgNO_3 + NO + 2H_2O$
299. (d) $AgCl + 2NH_3 \rightarrow [Ag(NH_3)_2]Cl$ (soluble complex)
301. (b) AgBr is most sensitive to light and undergoes photochemical reduction
 $2AgBr \xrightarrow{\text{Light}} 2Ag + Br_2$
302. (a) NaNO₃ is purely ionic while AgCl is covalent other compounds reacts with AgCl
 $2AgCl + Na_2CO_3 \rightarrow 2Ag + 2NaCl + CO_2 + \frac{1}{2}O_2$
 $AgCl + 2Na_2CO_3 \rightarrow Na_2[Ag(S_2O_3)_2] + NaCl$
 $AgCl + 2NH_4OH \rightarrow [Ag(NH_3)_2Cl] + 2H_2O$
304. (a) A very dilute solution is used in cauterisation of eyes and dental antiseptic
305. (a) $2AgNO_3 \rightarrow 2Ag + 2NO_2 + O_2$
306. (a) $2AgNO_3 \xrightarrow{PH_3} 2Ag + 2NO_2 + O_2$
307. (b) $2Ag + 2H_2SO_4 \rightarrow Ag_2SO_4 + SO_2 + 2H_2O$
308. (c) Ag salts on strong heating from Ag
309. (a) Silver metal is extracted from the argentite ore (Ag₂S) by the cyanide process, in which ore is treated with sodium cyanide sold. Dicyanoargentate(I) $[2Na\{Ag(CN)_2\}]$ is formed.
310. (b) When a strip of copper is dipped in the solution of silver nitrate, the solution becomes blue. Cu is placed above Ag in electrochemical series.
 $2AgNO_3 + Cu \rightarrow Cu(NO_3)_2 + 2Ag$
311. (d) $ZnO + 2NaOH \rightarrow Na_2ZnO_2 + H_2O$
312. (d) $Zn + 2NaOH \rightarrow Na_2ZnO_2 + H_2$
 $2Al + 2NaOH \rightarrow 2NaAlO_2$
314. (d) The white solid dissolves to form a blue solution.
 $CuSO_4 + \text{dil } H_2SO_4 \rightarrow CuSO_4 \cdot 5H_2O$
blue vitriol
317. (c) $ZnSO_4 + 2NaHCO_3 \rightarrow ZnCO_3 + Na_2SO_4 + H_2O + CO_2$
318. (b) $Zn + 2NaOH \xrightarrow{\text{heat}} Na_2ZnO_2 + H_2$
319. (d) $ZnO + 2HCl \rightarrow ZnCl_2 + H_2O$
 $ZnO + 2NaOH \rightarrow Na_2ZnO_2 + H_2O$
320. (c) Very dil HNO₃ → Ammonium nitrate is formed
 $4Zn + 10HNO_3 \rightarrow 4Zn(NO_3)_2 + NH_4NO_3 + 3H_2O$
321. (d) $Zn_{30} \rightarrow 3d^{10}, 4s^2$
 $Zn^{+2} \rightarrow 3d^{10}$
- | | | | | |
|----|----|----|----|----|
| ↑↓ | ↑↓ | ↑↓ | ↑↓ | ↑↓ |
|----|----|----|----|----|
- No unpaired electrons
323. (d) Sodium tetraborate decahydrate (Na₂B₄O₇·10H₂O)
324. (a) Zn does not react with cold water. However it reacts with hot water and yield H₂
 $Zn + H_2O \xrightarrow{\text{Boil}} ZnO + H_2$
 $Zn + H_2SO_4(\text{dil}) \rightarrow ZnSO_4 + H_2$
 $Zn + 2HCl(\text{dil}) \rightarrow ZnCl_2 + H_2$
 $Zn + 2NaOH \xrightarrow[20\%]{\text{heat}} Na_2ZnO_2 + H_2$
325. (b) $Zn + 2HCl \rightarrow ZnCl_2 + H_2$
 $Zn + 2NaOH \rightarrow Na_2ZnO_2 + H_2$
327. (a) The compound ZnFe₂O₄ is a normal spinel compound.
328. (a) $ZnO + BaO \xrightarrow{1100^\circ C} BaZnO_2$
329. (b) $4Zn + 10HNO_3 \rightarrow 4Z(NO_3)_2 + NH_4NO_3 + 3H_2O$
330. (d) Lead is used for making radiation shield.
331. (b) $Ag_2S + 4NaCN \rightarrow 2Na[Ag(CN)_2] + Na_2S$
 $2Na[Ag(CN)_2] + Zn \rightarrow Na_2[Zn(CN)_4] + 2Ag$
333. (d) In Mc Arthur Forest method, Silver is extracted from solution of sodium argenticyanide by using Zn
 $2Na[Ag(CN)_2] + Zn \rightarrow Na_2[Zn(CN)_4] + 2Ag \downarrow$
338. (d) $Ag_2S + 4NaCN \rightarrow 2Na[Ag(CN)_2] + Na_2S$
 $2Na[Ag(CN)_2] + Zn \rightarrow Na_2[Zn(CN)_4] + 2Ag$
342. (c) Self reduction
 $2Cu_2S + 3O_2 \rightarrow 2Cu_2O + 2SO_2$
 $Cu_2S + 2Cu_2O \rightarrow 6Cu + SO_2$
345. (c) $FeO + SiO_2 \rightarrow FeSiO_3$
Impurity Flux Slag
346. (c) Parke's process is used to extract silver from argentiferous lead.
347. (c) $2Cu_2S + 3O_2 \rightarrow 2Cu_2O + 2SO_2 \uparrow$
348. (d) Copper metallurgy involves bessemerisation. In bessemerisation converter, the impurities of Ferric Oxide forms slag with Silica and Copper Oxide gets reduced to give blister copper.
 $FeO + SiO_2 \rightarrow FeSiO_3$
Slag
 $Cu_2S + Cu_2O \rightarrow 6Cu \downarrow + SO_2 \uparrow$
349. (c) Bessemer converter is used to purify Pig Iron by passing compressed air over pig iron in Bessemer converter to produce slag.
 $2Mn + O_2 \rightarrow 2MnO; Si + O_2 \rightarrow SiO_2$
 $2C + O_2 \rightarrow 2CO; MnO + SiO_2 \rightarrow MnSiO_3$
Slag
353. (d) 1% Impure copper
358. (c) None of the above. Since, gold is a noble metal and common acids do not attack on it if used singly.
359. (a) Due to reduction of copper
 $Zn + CuSO_4 \rightarrow Cu + ZnSO_4$





360. (a) Reduction of Cu^{++} .
 $Fe + CuSO_4 \rightarrow FeSO_4 + Cu$


Critical Thinking Questions

1. (c) Transition metal which have low oxidation number show the oxidising nature because of great tendency to lose the electron.





2. (b) Cr^{+2} and Fe^{+2}




3. (d) The solubility order is :



4. (d) No of unpaired electron in different ion are as under









5. (a) The magnitude of stability constants for some divalent metal ions of the first transition series with oxygen or nitrogen donor ligands increases in the order.




6. (c) Strong oxidising agents such as PbO_2 or sodium bismuthate ($NaBiO_3$) oxidise Mn^{2+} to MnO_4^- or Mn^{7+} .

7. (c) Given $n = 4$ $x = 5$




8. (b) Iron decomposes steam into hydrogen when it is passed over red hot iron




9. (d) $CoCl_3 - Co^{+3} - 3d^6 4s^0$

4 unpaired electrons. So it will be coloured.

10. (a) Due to Lanthanoid contraction order will be



11. (d) In this reaction



5 time quantity of Fe^{2+} consumed.

So 5 time of $FeSO_4$ will be equivalent to 50 ml

12. (abc) Due to less capacity of hydrogen bonding of I_2 with water HgI_2 is less soluble in water.

13. (c) $\frac{\text{Transition element} + \text{Inner transition element}}{\text{Total element}} \times 100$



$$\frac{33 + 28}{105} \times 100 = 58.09 \approx 60\%$$

14. (a) All metal carbonyls are diamagnetic cyanide complexes are also diamagnetic.

15. (c) 22 carat gold is alloy of copper and gold.

16. (b) Fe^{3+} have highest no. of unpaired electron so it will be more paramagnetic.

17. (d) p -electrons in Cl^- (atomic no. of $Cl = 17$)



In Fe^{2+} total number of $de^- = 6$ which is not equal to pe^- in $Cl^- = 12$.

18. (a) Basic character of oxide decreases from left to right in a period of periodic table.


19. (b) $CuSO_4 \cdot 5H_2O$ because it has only one unpaired electrons.

20. (c) The existence of Fe^{2+} and NO^+ in Nitroprusside ion $[Fe(CN)_5NO]^{2-}$ can be established by measuring the magnetic moment of the solid compound which should correspond to ($Fe^{2+} = 3d^6$) four unpaired electrons.

21. (c) $V^{+4} \rightarrow 3d^1 4s^0$

1 unpaired electrons. Hence, it is paramagnetic and coloured compound.


22. (a) $3MnO_4^- + 5(Fe^{2+} + C_2O_4^{2-}) + 24H^+ \rightarrow$



Thus 5M of FeC_2O_4 is oxidised by 3M of $KMnO_4$ then 1M of FeC_2O_4 is oxidised by 3/5 mole of $KMnO_4$.

23. (b) $2MnO_4^- + 16H^+ + C_2O_4^{2-} \rightarrow 2Mn^{+2} + 2CO_2 + 8H_2O$


24. (b) $ScCl_3 \rightarrow Sc^{+3} + 3Cl^-$





No unpaired electron so will show diamagnetic character and will be repelled, so will weigh less.

25. (a) $(Ar) 3s^1 + 3 = Ti$, it means M^{3+} form Ti^{3+} ion.

26. (b) $V = 1s^2 2s^2 2p^6 3s^2 3p^6 3d^3 4s^2$






In second ionization enthalpy Cr^+ has exact half filled d -sub shell.

Assertion and Reason


2. (b) Zn^{2+} is diamagnetic because it has no unpaired electrons
3. (c) The assertion is correct but the reason is false. Actually transition metal show variable valency due to very small difference between the ns^2 and $(n-1)d$ electrons, Therefore, assertion is correct but reason is false.

4. (e) The aqueous solution of $FeCl_3$ is acidic in nature because $FeCl_3$ hydrolyse in water to produce acid ion.



Therefore, assertion is false but reason is true.

5. (a) $AgCl$ on adding to a solution of NH_4OH solution dissolves to form a complex diamine silver chloride.




Therefore, both assertion and reason are true and reason is a correct explanation of assertion.

6. (c) Pure iron is not used for making tools and machines as it is soft. Therefore, cannot be used for this purpose. Assertion is true but reason is false.

7. (a) A solution of Na_2CrO_4 in water is intensely coloured due to oxidation state of chromium in Na_2CrO_4 is +6. Here both assertion and reason are correct.

8. (d) Copper corrods at negligible rates in unpolluted air, water and deaerated non-oxidizing acids. Pure copper and the high copper alloys can be considered to exhibit similar resistance to most corrosive environments. They possess excellent resistance to atmospheric environments. Corrosion is a spontaneous process for which free energy change must be negative.

9. (c) ${}_{24}Cr \rightarrow [Ar] 3d^4 4s^2$




Full-filled s -orbital has greater stability.

10. (a) Fe has atomic no. 26.

So its electronic configuration is $[Ar] 3d^6 4s^2$.

Fe^{2+} has electronic configuration $[Ar] 3d^6$.




It has four unpaired electrons and it is paramagnetic.

11. (b) Due to larger surface area and variable valencies to form intermediate absorbed complex easily, transition metals are used as good catalysts.

12. (c) Rusting involves reduction of absorbed oxygen to OH^- ions and oxidation of iron to Fe^{2+} ions. The two ions combine to yield $Fe(OH)_2$ which gets oxidized to give $Fe_2O_3 \cdot nH_2O$ (rust). The presence of acid helps dissolution of pure iron to ferrous ions while electrolytes increase the conductivity and assist the cell action.

13. (b) $AgBr$ is the most sensitive silver halide to photo reduction. Hence it is used as the light sensitive material in photographic films. The unchanged $AgBr$ is dissolved in hypo solution to cast an image on photographic plate.



14. (a) Tungsten is a metal of high melting point and its filament gives brilliant light on passing electric current.

17. (b) The magnetic moments are lesser than the fact that $5f$ electrons of actinides are less effectively shielded which results in quenching of orbital contribution.


18. (b) The higher the charge on the metal ion, smaller is the ionic size and more is the complex forming decreases in the order $M^{4+} > MO_2^{2+} > M^{3+} > MO_2^+$. The higher tendency of complex formation of MO_2^{2+} of charge on metal atom M in MO_2^{2+}

20. (d) Extraction of iron metal from iron oxide ore is carried out by heating with coke and flux (calcium carbonate). Flux is a slag forming substance. It converts infusible impurities into fusible slag.

The reaction : $FeO_{(s)} \rightarrow Fe_{(s)} + 3/2 O_{2(g)}$ is not a spontaneous process. $FeO_{(s)}$ is converted to FeO at about $400^\circ C$. FeO is converted to Fe at about $800^\circ C - 1000^\circ C$.

- The equivalent weight of $MnSO_4$ is half of its molecular weight when it is converted to
 - Mn_2O_3
 - MnO_2
 - MnO_4^-
 - MnO_4^{2-}
- In which compound chromium has +6 oxidation state [CPMT 2003]
 - $K_2Cr_2O_7$
 - $CrCl_3$
 - $Cr(SO_4)_3$
 - None of these
- Which of the following metal does not show variable valency [RPET 2000]
 - Fe
 - Hg
 - Zn
 - Cu
- Which of the following metals will not react with solution of $CuSO_4$ [CPMT 1974, 80; MH CET 2004]
 - Fe
 - Zn
 - Mg
 - Ag
- Which metal among following has strongest tendency to undergo oxidation [CPMT 1989]
 - Zn
 - Cu
 - Mg
 - Al
- Which of the following has highest paramagnetic character
 - Mn (II)
 - Fe (II)
 - Co (II)
 - Ni (II)
- Ammonia is a Lewis base. It forms complexes with cations. Which one of the following cations does not form complex with ammonia
 - Ag^+
 - Cu^{++}
 - Cd^{++}
 - Pb^{++}
- Which of the following is expected to form colourless complex [AMU 2000]
 - Ni^{2+}
 - Cu^+
 - Ti^{3+}
 - Fe^{3+}
- Which of the following is ferromagnetic
 - Cr
 - Mn
 - W
 - Co
- The most stable oxidation state of Mn is
 - +2
 - +4
 - +5
 - +7
- The number of unpaired electrons in Mn^{+3} is
 - 4
 - 3
 - 2
 - Zero
- The correct order of ionic radii of Y^{3-} , La^{3+} , Eu^{3+} and Lu^{3+} is [CBSE PMT 2000]
 - $La^{3+} < Eu^{3+} < Lu^{3+} < Y^{3+}$
 - $Y^{3-} < La^{3+} < Eu^{3+} < Lu^{3+}$
 - $Lu^{3+} < Y^{3+} < Eu^{3+} < La^{3+}$
 - $Lu^{3+} < Eu^{3+} < La^{3+} < Y^{3+}$
 (Atomic No. $Y = 39$, $La = 57$, $Eu = 63$, $Lu = 71$)
- One mole of potassium dichromate completely oxidises the following number of moles of ferrous sulphate in acidic medium [MP PET 1997]
 - 1
 - 3
 - 5
 - 6
- The atomic radii from Cr to Cu is almost identical because of
 - Increasing nuclear charge from Cr to Cu
 - Repulsion among increased electrons
 - Increased screening effect to nullify increased nuclear charge
 - All the above
- Oxidation number of Mn in K_2MnO_4 and in $KMnO_4$ are respectively [MP PET 1991, 2001]
 - +6 and +7
 - +6 and +6
 - +7 and +7
 - +7 and +6
- When phosphine is passed through aqueous solution of copper sulphate, the product produced is
 - $Cu(OH)_2$
 - Cu_3P_2
 - $[Cu(PH_3)_4]^{2+}$
 - $[Cu(PH_3)_2]^{2+}$
- Hydroxide soluble in ammonia is [NCERT 1973, 77; MNR 1984; KCET 1992]
 - $Al(OH)_3$
 - $Fe(OH)_3$
 - $Cr(OH)_3$
 - $Cu(OH)_2$
- Which of the following pair of transitional elements exhibit highest and lowest density
 - Os and Sc
 - Os and Pt
 - Hg and Sc
 - Os and Ir
- When an acidified solution of ferrous ammonium sulphate is treated with potassium permanganate solution, the ion which is oxidised is [BHU 1979]
 - MnO_4^-
 - NH_4^+
 - Fe^{++}
 - SO_4^{2-}

1. (b) $MnSO_4 \xrightarrow{-2e^-} MnO_2$
 Equivalent wt. = $\frac{\text{molecular wt.}}{\text{total no. of } e^- \text{ gained or lost}} = \frac{M}{2}$
2. (a) In $K_2Cr_2O_7$, Cr has + 6 oxidation state.
3. (c) Zn shows only + 2 valency.
4. (d) Because Ag comes below in the electromotive series also standard electrode potential of Cu and Ag are:
 $Cu^{2+}(aq) + 2e^- \rightarrow Cu(s); [SEP - E_{298}^0 (\text{volt}) = +0.18]$
 $Ag^+(aq) + e^- \rightarrow Ag(s); [SEP - E_{298}^0 (\text{volt}) = +0.80]$
5. (c) Mg; because of its high hydration energy.
6. (a) $Mn^{+2} - 3d^5$
 5 unpaired element in d-subshell so it has highest paramagnetic.
7. (d) Pb^{+2} because it does not have vacant d-orbitals nor high nuclear charge and it does not belong to transition series.
8. (b) In Cu^{+1} (cuprous ion) d orbitals are completely filled so it will form colourless complex.
9. (d) The substances which are strongly attracted by magnetic field and show permanent magnetism even in absence of magnetic field are ferromagnetic e.g., Co, Fe, Ni
10. (a)



As half filled orbitals are more stable than partial filled ones. Therefore, + 2 is most stable oxidation state.

11. (a) Mn^{+3}

$3d^5$	$4s^0$
↑	↑
12. (c) Lanthanide contraction results in small size of Lu^{3+} , so
 $Lu^{3+} < Y^{3+} < Cu^{3+} < La^{3+}$
13. (d) Oxidation number of chromium in potassium dichromate is +6 so it oxidise 6 moles of ferrous sulphate in acidic medium.
14. (c) Increased screening effect to nullify increased nuclear charge.
15. (a) O.N. of K_2MnO_4

$$2 + x - 8 = 0$$

$$x = 6$$

O.N. of $KMnO_4$

$$1 + x - 8 = 0$$

$$x = 7$$

16. (b) $3CuSO_4 + 2PH_3 \rightarrow Cu_3P_2 + 3H_2SO_4$
17. (d) Due to formation of complex
 $Cu(OH)_2 + 4NH_3 \rightarrow [Cu(NH_3)_4](OH)_2$
18. (a) Os and Sc
 $Os = 22.60 \text{ gm/cm}^3$
 $Sc = 3.01 \text{ gm/cm}^3$
19. (c) $Fe^{2+} \xrightarrow{\text{oxidises}} Fe^{3+}$