

DPP - Daily Practice Problems

Chapter-wise Sheets

Date : _____

Start Time : _____

End Time : _____

CHEMISTRY (CC06)

SYLLABUS : Thermodynamics

Max. Marks : 180

Marking Scheme : + 4 for correct & (-1) for incorrect

Time : 60 min.

INSTRUCTIONS : This Daily Practice Problem Sheet contains 45 MCQ's. For each question only one option is correct. Darken the correct circle/ bubble in the Response Grid provided on each page.

1. In a closed insulated container a liquid is stirred with a paddle to increase the temperature. Which of the following is true?
(a) $\Delta E = W \neq 0$ (b) $\Delta E = W, q = 0$
(c) $\Delta E = 0, W = q \neq 0$ (d) $W = 0, \Delta W = q \neq 0$
2. One mole of a non-ideal gas undergoes a change of state ($2.0\text{ atm}, 3.0\text{ L}, 95\text{ K}$) \rightarrow ($4.0\text{ atm}, 5.0\text{ L}, 245\text{ K}$) with a change in internal energy, $\Delta U = 30.0\text{ Latm}$. The change in enthalpy ΔH of the process in L atm is.
(a) 40.0 (b) 42.3
(c) 44.0 (d) Not defined because pressure is not constant
3. Assuming that water vapour is an ideal gas, the internal energy change (ΔU) when 1 mol of water is vapourised at 1 bar pressure and 100°C , (given : molar enthalpy of vapourisation of water at 1 bar and $373\text{ K} = 41\text{ kJ mol}^{-1}$ and $R = 8.3\text{ J mol}^{-1}\text{ K}^{-1}$) will be
(a) 41.00 kJ mol^{-1} (b) 4.100 kJ mol^{-1}
(c) $3.7904\text{ kJ mol}^{-1}$ (d) $37.904\text{ kJ mol}^{-1}$
4. A piston filled with 0.04 mole of an ideal gas expands reversibly from 50.0 mL to 375 mL at a constant temperature of 37.0°C . As it does so, it absorbs 208 J of heat. The values of q and w for the process will be:
($R = 8.314\text{ J/molK}$) ($\ln 7.5 = 2.01$)
(a) $q = +208\text{ J}, w = -208\text{ J}$ (b) $q = -208\text{ J}, w = -208\text{ J}$
(c) $q = -208\text{ J}, w = +208\text{ J}$ (d) $q = +208\text{ J}, w = +208\text{ J}$
5. Among the following, the intensive property is (properties are)
(a) molar conductivity (b) electromotive force
(c) resistance (d) heat capacity
6. The variation of heat of reaction with temperature is given by:
(a) Van't Hoff equation
(b) Clausius- Clapeyron equation
(c) Nernst equation
(d) Kirchoff's equation
7. The difference between heats of reaction at constant pressure and constant volume for the reaction :
 $2\text{ C}_6\text{H}_6(\text{l}) + 15\text{ O}_2(\text{g}) \rightarrow 12\text{ CO}_2(\text{g}) + 6\text{ H}_2\text{O}(\text{l})$ at 25°C in kJ is
(a) -7.43 (b) +3.72
(c) -3.72 (d) +7.43

RESPONSE
GRID



- | | | | | |
|--|--|--|--|--|
| 1. <input type="radio"/> a <input type="radio"/> b <input type="radio"/> c <input type="radio"/> d | 2. <input type="radio"/> a <input type="radio"/> b <input type="radio"/> c <input type="radio"/> d | 3. <input type="radio"/> a <input type="radio"/> b <input type="radio"/> c <input type="radio"/> d | 4. <input type="radio"/> a <input type="radio"/> b <input type="radio"/> c <input type="radio"/> d | 5. <input type="radio"/> a <input type="radio"/> b <input type="radio"/> c <input type="radio"/> d |
| 6. <input type="radio"/> a <input type="radio"/> b <input type="radio"/> c <input type="radio"/> d | 7. <input type="radio"/> a <input type="radio"/> b <input type="radio"/> c <input type="radio"/> d | | | |

Space for Rough Work

C-22

DPP/ CC06

8. The enthalpy of vapourisation of water from the following two equations is.



- (a) 6.02 kJ (b) 40.5 kJ (c) 62.3 kJ (d) 1.25 kJ


9. For complete combustion of ethanol,



the amount of heat produced as measured in bomb calorimeter, is $1364.47 \text{ kJ mol}^{-1}$ at 25°C . Assuming ideality the enthalpy of combustion, $\Delta_c H$, for the reaction will be: ($R = 8.314 \text{ kJ mol}^{-1}$)

- (a) $-1366.95 \text{ kJ mol}^{-1}$ (b) $-1361.95 \text{ kJ mol}^{-1}$
 (c) $-1460.95 \text{ kJ mol}^{-1}$ (d) $-1350.50 \text{ kJ mol}^{-1}$

10. The direct conversion of A to B is difficult, hence it is carried out by the following shown path :



Given

$\Delta S_{(A \rightarrow C)} = 50 \text{ e.u.}$, $\Delta S_{(C \rightarrow D)} = 30 \text{ e.u.}$, $\Delta S_{(B \rightarrow D)} = 20 \text{ e.u.}$, where e.u. is the entropy unit, then $\Delta S_{(A \rightarrow B)}$ is

- (a) $+60 \text{ e.u.}$ (b) $+100 \text{ e.u.}$ (c) -60 e.u. (d) -100 e.u.

11. The heat of combustion of $\text{CH}_4(\text{g})$, C (graphite), $\text{H}_2(\text{g})$ are -20 kcal , -40 kcal , -10 kcal respectively. The heat of formation of methane is

- (a) -40 kcal (b) $+40 \text{ kcal}$
 (c) -80.0 kcal (d) $+80 \text{ kcal}$

12. From the following bond energies:


H – H bond energy: $431.37 \text{ kJ mol}^{-1}$

C = C bond energy: $606.10 \text{ kJ mol}^{-1}$

C – C bond energy: $336.49 \text{ kJ mol}^{-1}$

C – H bond energy: $410.50 \text{ kJ mol}^{-1}$

Enthalpy for the reaction,



will be:

- (a) $-243.6 \text{ kJ mol}^{-1}$ (b) $-120.0 \text{ kJ mol}^{-1}$
 (c) $553.0 \text{ kJ mol}^{-1}$ (d) $1523.6 \text{ kJ mol}^{-1}$

RESPONSE GRID


- | | | | | |
|------------------|------------------|------------------|------------------|------------------|
| 8. (a)(b)(c)(d) | 9. (a)(b)(c)(d) | 10. (a)(b)(c)(d) | 11. (a)(b)(c)(d) | 12. (a)(b)(c)(d) |
| 13. (a)(b)(c)(d) | 14. (a)(b)(c)(d) | 15. (a)(b)(c)(d) | 16. (a)(b)(c)(d) | 17. (a)(b)(c)(d) |
| 18. (a)(b)(c)(d) | 19. (a)(b)(c)(d) | 20. (a)(b)(c)(d) | | |

Space for Rough Work

21. The following two reactions are known :



The value of ΔH for the following reaction



- (a) +6.2 kJ (b) +10.3 kJ (c) -43.3 kJ (d) -10.3 kJ


22. For a particular reversible reaction at temperature T , ΔH and ΔS were found to be both +ve. If T_e is the temperature at equilibrium, the reaction would be spontaneous when

- (a) $T_e > T$ (b) $T > T_e$
 (c) T_e is 5 times T (d) $T = T_e$

23. The standard enthalpy of formation of NH_3 is $-46.0 \text{ kJ mol}^{-1}$. If the enthalpy of formation of H_2 from its atoms is -436 kJ mol^{-1} and that of N_2 is -712 kJ mol^{-1} , the average bond enthalpy of N – H bond in NH_3 is
 (a) -964 kJ mol^{-1} (b) $+352 \text{ kJ mol}^{-1}$

- (c) +1056 kJ mol⁻¹ (d) -1102 kJ mol⁻¹

24. Standard entropy of X_2 , T_2 and $X-T_3$ are 60, 40 and $50 \text{ J K}^{-1} \text{ mol}^{-1}$, respectively. For the reaction,



- temperature will be

25. A heat engine absorbs heat Q_1 at temperature T_1 and heat Q_2 at temperature T_2 . Work done by the engine is J ($Q_1 + Q_2$). This data



- (a) violates 1st law of thermodynamics

- (a) violates 1st law of thermodynamics
 (b) violates 1st law of thermodynamics if $Q_1 = -vc$
 (c) violates 1st law of thermodynamics if $Q_2 = -vc$
 (d) does not violate 1st law of thermodynamics.

26. The standard enthalpies of formation of $\text{CO}_2(\text{g})$, $\text{H}_2\text{O}(\text{l})$ and glucose(s) at 25°C are -400 kJ/mol , -300 kJ/mol and -1300 kJ/mol , respectively. The standard enthalpy of combustion per gram of glucose at 25°C is

- (a) +2900 kJ (b) -2900 kJ (c) -16.11 kJ (d) +16.11 kJ
 Using the data provided, calculate the multiple bond energy (kJ mol^{-1}) of a $\text{C} \equiv \text{C}$ bond in C_2H_2 . That energy is (take the answer to 3 significant figures).

- $$2\text{C}(s) + \text{H}_2(g) \longrightarrow \text{HC}\equiv\text{CH}(g); \quad \Delta H = 225 \text{ kJ mol}^{-1}$$



- The enthalpy of neutralisation of NH_3OH and CH_3COOH

28. The enthalpy of neutralisation of NH_4OH and CH_3COOH is $-10.5 \text{ kcal mol}^{-1}$ and enthalpy of neutralisation of

RESPONSE GRID

- 21.** a b c d **22.** a b c d **23.** a b c d **24.** a b c d **25.** a b c d
26. a b c d **27.** a b c d **28.** a b c d **29.** a b c d **30.** a b c d
31. a b c d **32.** a b c d **33.** a b c d

Space for Rough Work

34. The ΔH at 358 K for the reaction





given that $\Delta H_{298} = -33.29 \text{ kJ mol}^{-1}$ and C_p for $\text{Fe}_2\text{O}_3(\text{s})$, $\text{Fe}(\text{s})$, $\text{H}_2\text{O}(\text{l})$ and $\text{H}_2(\text{g})$ are 103.8, 25.1, 75.3 and 28.8 $\text{J K}^{-1} \text{ mol}^{-1}$
 (a) $-28.136 \text{ kJ mol}^{-1}$ (b) $-38.3 \text{ kJ mol}^{-1}$
 (c) 42.5 kJ mol^{-1} (d) 56.2 kJ mol^{-1}

35. The enthalpy of neutralisation of a weak acid in 1 M solution with a strong base is $-56.1 \text{ kcal mol}^{-1}$. If the enthalpy of ionisation of acid is $1.5 \text{ kcal mol}^{-1}$ and enthalpy of neutralisation of the strong acid with a strong base is -57.3 kJ eq^{-1} . What is the % ionisation of the weak acid in molar solution (assume the acid is monobasic)
 (a) 25 (b) 20 (c) 15 (d) 10

36. ΔH_f° of NF_3 is -113 kJ mol^{-1} and N–F bond energy is $273.0 \text{ kJ mol}^{-1}$. If N≡N and F–F bond energies are in the ratios 6 : 1, their magnitudes will be
 (a) $780.0 \text{ kJ mol}^{-1}, 130.0 \text{ kJ mol}^{-1}$
 (b) $840 \text{ kJ mol}^{-1}, 140 \text{ kJ mol}^{-1}$
 (c) $950.0 \text{ kJ mol}^{-1}, 158.3 \text{ kJ mol}^{-1}$
 (d) $941.3 \text{ kJ mol}^{-1}, 156.9 \text{ kJ mol}^{-1}$

37. Consider the following processes :
 $\Delta H (\text{kJ/mol})$



For $\text{B} + \text{D} \rightarrow \text{E} + 2\text{C}$, ΔH will be :

- (a) 525 kJ/mol (b) -175 kJ/mol
 (c) -325 kJ/mol (d) 325 kJ/mol

38. For an isothermal reversible expansion process, the value of q can be calculated by the expression

$$(a) q = 2.303nRT \log \frac{V_2}{V_1} \quad (b) q = -2.303nRT \log \frac{V_2}{V_1}$$

$$(c) q = -P_{\text{exp}} nRT \log \frac{V_1}{V_2} \quad (d) \text{None of these}$$

39. Match the columns

Column-I

- A. $\text{C}_4\text{H}_{10} + \frac{13}{2}\text{O}_2 \rightarrow 4\text{CO}_2 + 5\text{H}_2\text{O}; \Delta H = -w$
 B. $\text{CH}_4 \rightarrow \text{C} + 4\text{H}; \Delta H = x$
 C. $\text{H}_2 + \text{Br}_2 \rightarrow 2\text{HBr}; \Delta H = y$
 D. $\text{Na}^-(\text{s}) \rightarrow \text{Na}(\text{g}); \Delta H = z$

Column-II

- I. Enthalpy of atomisation
 II. Enthalpy of formation
 III. Enthalpy of combustion
 IV. Enthalpy of sublimation


- (a) A – IV; B – I; C – II; D – III

- (b) A – II; B – III; C – I; D – IV

- (c) A – III; B – I; C – II; D – IV

- (d) A – I; B – II; C – IV; D – III

40. From the following data ΔH of the following reactions



Calculate the mole composition of the mixture of steam and oxygen on being passed over coke at 1273 K, keeping the temperature constant.

- (a) 1 : 0.6 (b) 0.6 : 1 (c) 2 : 3 (d) 3 : 2

41. The heats of neutralisation of CH_3COOH , HCOOH , HCN and H_2S are -13.2 , -13.4 , -2.9 and -3.8 kJ/mol respectively. Arrange the acids in increasing order of strength

- (a) $\text{HCOOH} > \text{CH}_3\text{COOH} > \text{H}_2\text{S} > \text{HCN}$

- (b) $\text{CH}_3\text{COOH} > \text{HCOOH} > \text{H}_2\text{S} > \text{HCN}$

- (c) $\text{H}_2\text{S} > \text{HCOOH} > \text{CH}_3\text{COOH} > \text{HCN}$

- (d) $\text{HCOOH} > \text{H}_2\text{S} > \text{CH}_3\text{COOH} > \text{HCN}$

42. Which of the following salts should cause maximum cooling when 1 mole of it is dissolved in the same volume of water?

- (a) $\text{NaCl}; \Delta H^\circ = 5.35 \text{ kJ mol}^{-1}$

- (b) $\text{KNO}_3; \Delta H^\circ = 53.5 \text{ kJ mol}^{-1}$

- (c) $\text{KOH}; \Delta H^\circ = -56.0 \text{ kJ mol}^{-1}$

- (d) $\text{HBr}; \Delta H^\circ = -83.3 \text{ kJ mol}^{-1}$

43. Read the following statements carefully and choose the correct option

- (i) Internal energy U of the system is a state function.

- (ii) $-w$ shows, that work is done on the system.

- (iii) $+w$ shows, that work is done by the system

- (a) (i) and (ii) are correct (b) (ii) and (iii) are correct

- (c) (i) and (iii) are correct (d) Only (i) is correct

44. Consider the following reaction occurring in an automobile






the sign of ΔH , ΔS and ΔG would be

- (a) $+, -, +$ (b) $- , +, -$

- (c) $- , +, +$ (d) $+ , +, -$

45. Choose the reaction with negative ΔS value.



RESPONSE GRID

34. (a) (b) (c) (d)
 39. (a) (b) (c) (d)
 44. (a) (b) (c) (d)

35. (a) (b) (c) (d)
 40. (a) (b) (c) (d)
 45. (a) (b) (c) (d)

36. (a) (b) (c) (d)

41. (a) (b) (c) (d)

37. (a) (b) (c) (d)

42. (a) (b) (c) (d)

38. (a) (b) (c) (d)

43. (a) (b) (c) (d)

Space for Rough Work

**DAILY PRACTICE
PROBLEMS**
**CHEMISTRY
SOLUTIONS**
DPP/CC06

1. (b) In insulated system no heat or matter is exchanged, $q = 0 \therefore \Delta E = W$. The work done on the system increases internal energy.

$$\begin{aligned} 2. (c) \quad \Delta H &= H_2 - H_1 = (E_2 + P_2 V_2) - (E_1 + P_1 V_1) \\ &= (E_2 - E_1) + (P_2 V_2 - P_1 V_1) \\ &= 30 + 4 \times 5 - 2 \times 3 = 44 \text{ J atm} \end{aligned}$$

$$\begin{aligned} 3. (d) \quad \text{Given } \Delta H &= 41 \text{ kJ mol}^{-1} = 41000 \text{ J mol}^{-1} \\ T &= 100^\circ\text{C} = 273 + 100 = 373 \text{ K} \\ n &= 1 \\ \Delta U &= \Delta H - \Delta n RT = 41000 - (1 \times 8.314 \times 373) \\ &= 37898.88 \text{ J mol}^{-1} \approx 37.9 \text{ kJ mol}^{-1} \end{aligned}$$

4. (a) Process is isothermal reversible expansion, hence $\Delta U = 0$, therefore $q = -W$. Since $q = +208 \text{ J}$, $W = -208 \text{ J}$


5. (b) An intensive property is a physical quantity whose value does not depend on the amount of the substance for which it is measured.

$$6. (d) \quad \text{Kirchoff's equation, } \frac{\Delta H_2 - \Delta H_1}{T_2 - T_1} = \Delta C_p$$

$$\begin{aligned} 7. (a) \quad \text{Heat capacity at constant volume } (q_v) &= \Delta E \\ \text{Heat capacity of constant pressure } (q_p) &= \Delta H \\ \Delta H &= \Delta E + \Delta n RT \text{ or } \Delta H - \Delta E = \Delta n RT \\ \Delta n &= \text{no. of moles of gaseous products} \\ &\quad - \text{no. of moles of gaseous reactants} \\ &= 12 - 15 = -3 \\ \Delta H - \Delta E &= -3 \times 8.314 \times 298 \text{ J} = -7.43 \text{ kJ.} \end{aligned}$$

8. (b) Desired equation is $\text{H}_2\text{O(l)} \rightarrow \text{H}_2\text{O(g)}$, $\Delta H = ?$
(Equation II - Equation I)

$$\Delta H = -245.5 \text{ kJ} - (-286 \text{ kJ}) = 40.5 \text{ kJ}$$



Bomb calorimeter gives ΔU of the reaction



$$\text{Given, } \Delta U = -1364.47 \text{ kJ mol}^{-1}$$

$$\Delta n_g = -1$$

$$\begin{aligned} \Delta H &= \Delta U + \Delta n_g RT = -1364.47 - \frac{1 \times 8.314 \times 298}{1000} \\ &= -1366.93 \text{ kJ mol}^{-1} \end{aligned}$$

$$10. (a) \quad \Delta S_{(A \rightarrow B)} = \Delta S_{(A \rightarrow C)} + \Delta S_{(C \rightarrow D)} - \Delta S_{(B \rightarrow D)} \\ = 50 + 30 - 20 = 60 \text{ c.u.}$$

$$11. (a) \quad \text{CH}_4 + 2\text{O}_2 \rightarrow \text{CO}_2 + 2\text{H}_2\text{O}; \Delta H = 20 \text{ kcal} \dots \text{(1)} \\ \text{C} + \text{O}_2 \rightarrow \text{CO}_2; \Delta H = -40 \text{ kcal} \dots \text{(2)}$$



$$\text{or } \Delta H = -20 + (-40) - (-20) = -40 \text{ kcal}$$

$$12. (b) \quad \text{Enthalpy of reaction} \\ = \text{B.E}_{(\text{Reactant})} - \text{B.E}_{(\text{Product})}$$

$$= [\text{B.E}_{(\text{C=C})} + 4\text{B.E}_{(\text{C-H})} + \text{B.E}_{(\text{H-H})}]$$

$$- [\text{B.E}_{(\text{C-C})} + 6\text{B.E}_{(\text{C-H})}]$$

$$= [606.1 + (4 \times 410.5) + 431.37] - [336.49 + (6 \times 410.5)] \\ = -120.0 \text{ kJ mol}^{-1}$$

$$13. (a) \quad \Delta G = \Delta H - T\Delta S, \Delta H + vc, \Delta S \text{ is +vc; } T\Delta S > \Delta H \text{ for spontaneous process. It will make } \Delta G - vc$$

$$14. (d) \quad \Delta H = \Delta E + \Delta n RT \text{ For } \Delta H \neq \Delta E, \Delta n \neq 0 \\ \text{Where } \Delta n = \text{no. of moles of gaseous products} - \text{no. of moles of gaseous reactants}$$

$$(a) \Delta n = 2 - 2 = 0$$

$$(b) \Delta n = 0 \quad (\because \text{they are either in solid or liquid state})$$

$$(c) \Delta n = 1 - 1 = 0 \quad (\because \text{C is in solid state})$$

$$(d) \Delta n = 2 - 4 = -2$$

$\therefore (d)$ is correct answer

$$15. (b) \quad \text{The species in its elemental form has zero standard molar enthalpy of formation at 298 K. At 298 K, Cl}_2 \text{ is gas while Br}_2 \text{ is liquid.}$$

$$16. (b) \quad \text{High pressure reduces volume, decreases entropy, hence } \Delta S \text{ negative.}$$

$$17. (d) \quad \Delta S^\circ = 2S^\circ_{\text{HCl}} - (S^\circ_{\text{H}_2} + S^\circ_{\text{Cl}_2})$$

$$= 2 \times 186.7 - (130.6 + 223.0)$$

$$= 19.8 \text{ JK}^{-1} \text{ mol}^{-1}$$

$$18. (a) \quad \Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$$

For a spontaneous reaction $\Delta G^\circ < 0$

$$\text{or } \Delta H^\circ - T\Delta S^\circ < 0 \Rightarrow T > \frac{\Delta H^\circ}{\Delta S^\circ}$$

$$\Rightarrow T > \frac{179.1 \times 10^3}{160.2} > 1117.9 \text{ K} \approx 1118 \text{ K}$$

$$19. (d) \quad \text{For the reaction, } \text{C}_{(\text{s})} + \frac{1}{2}\text{O}_{2(\text{g})} \longrightarrow \text{CO}$$

$$\Delta H = \Delta U + \Delta n RT \text{ or } \Delta H - \Delta U = \Delta n RT$$


$$\Delta n = 1 - \frac{1}{2} = \frac{1}{2}; \Delta H - \Delta U = \frac{1}{2} \times 8.314 \times 298$$

$$= 1238.78 \text{ J mol}^{-1}$$


$$20. (c) \quad \Delta G^\bullet = \Delta H^\bullet - T\Delta S^\circ; -RT \ln K = \Delta H^\bullet - T\Delta S^\bullet$$



$$\ln K = -\frac{\Delta H^\circ - T\Delta S^\circ}{RT}$$



eq. (i) - 2 × eq. (ii), we get



$$\Delta H = -26.8 + 33.0 = +6.2 \text{ kJ}$$

22. (b) At equilibrium $\Delta G = 0$
Hence, $\Delta G = \Delta H - T_e \Delta S = 0$
 $\therefore \Delta H = T_e \Delta S$ or $T_e = \frac{\Delta H}{\Delta S}$

For a spontaneous reaction

ΔG must be negative which is possible only if $\Delta H - T\Delta S < 0$

$$\therefore \Delta H < T\Delta S \text{ or } T > \frac{\Delta H}{\Delta S}; T_e > T$$



Let x be the bond enthalpy of N-H bond then
[Note : Enthalpy of formation or bond formation enthalpy is given which is negative but the given reaction involves bond breaking hence values should be taken as positive.]

$\Delta H = \Sigma \text{Bond energies of reactants} - \Sigma \text{Bond energies of products}$

$$2 \times -46 = 712 + 3 \times (436) - 6x$$

$$-92 = 2020 - 6x$$


$$6x = 2020 + 92$$

$$6x = 2112$$

$$x = +352 \text{ kJ/mol}$$

24. (c) For a reaction to be at equilibrium $\Delta G = 0$. Since $\Delta G = \Delta H - T\Delta S$ so at equilibrium $\Delta H - T\Delta S = 0$ or $\Delta H = T\Delta S$

For the reaction



Calculating ΔS for the above reaction, we get

$$\begin{aligned} \Delta S &= 50 - \left[\frac{1}{2} \times 60 + \frac{3}{2} \times 40 \right] \text{ J K}^{-1} \\ &= 50 - (30 + 60) \text{ J K}^{-1} = -40 \text{ J K}^{-1} \end{aligned}$$

At equilibrium, $T\Delta S = \Delta H \quad [\because \Delta G = 0]$


$$\therefore T \times (-40) = -30 \times 1000 \quad [\because 1 \text{ kJ} = 1000 \text{ J}]$$

$$\text{or } T = \frac{-30 \times 1000}{-40} \text{ or } 750 \text{ K}$$

25. (a) According to first law of thermodynamics energy can neither be created nor destroyed although it can be converted from one form to another.

NOTE : Carnot cycle is based upon this principle but during the conversion of heat into work some mechanical energy is always converted to other form of energy hence this data violates 1st law of thermodynamics.

26. (c) The standard enthalpy of the combustion of glucose can be calculated by the eqn.




$$\Delta H_C = 6 \times \Delta H_f(\text{CO}_2) + 6 \times \Delta H_f(\text{H}_2\text{O}) - \Delta H_f[\text{C}_6\text{H}_{12}\text{O}_6]$$

$$\Delta H^\circ = 6(-400) + 6(-300) - (-1300)$$

$$\Delta H^\circ = -2900 \text{ kJ/mol}$$

For one gram of glucose, enthalpy of combustion

$$\Delta H^\circ = -\frac{2900}{180} = -16.11 \text{ kJ/g}$$



From equation (i) :

$$225 = [2 \times \Delta H_{\text{C}(s) \rightarrow \text{c}(g)} + 1 \times \text{BE}_{\text{H}-\text{H}}]$$

$$-[2 \times \text{BE}_{\text{C}-\text{H}} + 1 \times \text{BE}_{\text{C} \equiv \text{C}}]$$

$$225 = [1410 + 1 \times 330] - [2 \times 350 + 1 \times \text{BE}_{\text{C} \equiv \text{C}}]$$

$$225 = 1740 - 700 - \text{BE}_{\text{C} \equiv \text{C}}$$

$$\text{BE}_{\text{C} \equiv \text{C}} = 815 \text{ kJ mol}^{-1}$$

28. (b) ΔH_N for strong base and strong acid

$$= -13.7 \text{ kca l eq}^{-1}$$



$$= -12.5 - (-13.7) = 1.2 \text{ kcal mol}^{-1}$$




$$= -10.5 - (-13.7) - \Delta H_{\text{ion}}(\text{CH}_3\text{COOH})$$


$$= 13.7 - 10.5 - 1.2$$

$$= 2 \text{ kcal mol}^{-1}$$


29. (b) Write the proper chemical equations.




$$\Delta H = -393 \text{ kJ} \quad \dots(i)$$



$$\Delta H = -286 \text{ kJ} \quad \dots(ii)$$



From $(2 \times i + 3 \times ii) - (iii)$ we get



$$\therefore 2(-393 \text{ kJ}) + 3(-286 \text{ kJ}) - (-1360 \text{ kJ}) = -284 \text{ kJ}$$

30. (a) $\Delta H_{\text{hyd.}} = \Delta H_{\text{sol.}} - \Delta H_{\text{lattice}}$

$$= 1 - 180 = -179 \text{ kcal mol}^{-1}$$

$$\text{Then } \Delta H_{\text{hyd}}(\text{Na}^+) + \Delta H_{\text{hyd}}(\text{Cl}^-) = -179$$

$$\text{or } \Delta H_{\text{hyd}}(\text{Na}^+) + \frac{2}{3} \Delta H_{\text{hyd}}(\text{Na}^+) = -179$$

$$\text{or } \Delta H_{\text{hyd}}(\text{Na}^+) = -107.4 \text{ kcal/mol}^{-1}$$

31. (a) For non spontaneous reaction

$$\Delta G = + \text{ve}$$

$$\Delta G = \Delta H - T \Delta S \text{ and}$$

$$\Delta S = 121 \text{ J K}^{-1}$$

For $\Delta G = + \text{ve}$

ΔH has to be positive. Hence the reaction is endothermic.

The minimum value of ΔH can be obtained by putting

$$\Delta G = 0$$

$$\Delta H = T \Delta S = 298 \times 121 \text{ J} \\ = 36.06 \text{ kJ}$$

32. (c) The standard enthalpy of reaction is the enthalpy change for a reaction when all the participating substances are in their standard states. The standard state of a substance at a specified temperature is its pure form at 1 bar. For example, the standard state of liquid ethanol at 298 K is pure liquid ethanol at 1 bar. Standard state of solid iron at 500 K is pure iron at 1 bar.

33. (d)

$$34. (a) \Delta C_p = 2 \times 25.1 + 3 \times 75.3 - [103.8 + 3 \times 28.8]$$

$$= 85.9 \text{ J K}^{-1} \text{ mol}^{-1}$$

$$\frac{\Delta H_2 - \Delta H_1}{T_2 - T_1} = \Delta C_p;$$

$$\frac{\Delta H_2 - (-33290)}{358 - 298} = 85.9$$

$$\Delta H = -28136 \text{ J mol}^{-1}$$

$$= -28.136 \text{ kJ mol}^{-1}$$

35. (b) The enthalpy of ionisation of weak acid is given by

$$\Delta H_{\text{ion(HA)}}$$


$$= \Delta H_{\text{N(weak acid/strong base)}} - \Delta H_{\text{N(strong acid/strong base)}}$$

$$= -56.1 - (-57.3) = 1.2 \text{ kJ mol}^{-1}$$

$$\Delta H_{\text{(ionisation)}} = 1.5 \text{ kJ mol}^{-1}$$

Hence % ionisation in 1 M solution

$$= \frac{(1.5 - 1.2)}{1.5} \times 100 = 20$$



$$\text{or } \Delta H_{\text{N=N}} + \frac{3}{2} \Delta H_{\text{F-F}} - 3 \Delta H_{\text{N-F}} = -113 \text{ kJ}$$

Let $x \text{ kJ mol}^{-1}$ be the bond energy of F-F bond then bond energy of N≡N bond = $6x$

$$\therefore \frac{1}{2} \times 6x + \frac{3}{2} \times x - 3 \times 273 = -113 \text{ kJ}$$

On solving,

$$x = 156.9 \text{ kJ mol}^{-1} \text{ and}$$

$$\text{N≡N bond energy} = 6 \times 156.9 = 941.4 \text{ kJ mol}^{-1}$$

37. (b) Given ΔH



To calculate ΔH operate

$$2 \times \text{eq. (1)} + \text{eq. (2)} - \text{eq. (3)}$$

$$\Delta H = 300 - 125 - 350 = -175$$

$$38. (a) q = -W = 2.303nR T \log \frac{V_2}{V_1}$$

39. (c)

40. (a) The first reaction is exothermic and the second reaction is endothermic. On passing the mixture of O₂ and H₂O (steam) over coke while keeping the temperature constant ΔH of both the reactions must be same. Moles of O₂ needed to evolve 132 kJ

$$= \frac{0.5 \times 132}{110} = 0.6$$

Hence steam : O₂ ratio must be 1 : 0.6

41. (a) The greater the (negative value) of heat of neutralisation, the more is the strength of the acid. Hence, HCOOH > CH₃COOH > H₂S > HCN

42. (b) Dissolution of KNO₃ is endothermic, hence heat is absorbed and cooling is observed.

43. (d) The positive sign expresses when work is done on the system. Similarly, negative sign expresses when work is done by the system.

44. (b) This is combustion reaction, which is always exothermic hence

$$\Delta H = - \text{ve}$$

As the no. of gaseous molecules are increasing hence entropy increases so $\Delta S = + \text{ve}$.

$$\text{now } \Delta G = \Delta H - T \Delta S$$


For a spontaneous reaction

$$\Delta G = - \text{ve}$$

which is possible in this case as $\Delta H = - \text{ve}$ and $\Delta S = + \text{ve}$.

45. (c) ΔS has negative value if number of gaseous moles decreases during a reaction, $\Delta n_g = - \text{ve}$

For the reaction



$$\Delta n_g = 2 - 3 = -1$$

