

# Biomolecules

## Question1

Which of the following is a nucleotide?

[NEET 2024 Re]

Options:

A.

Uridine

B.

Adenylic acid

C.

Guanine

D.

Guanosine

**Answer: B**

**Solution:**

Uridine is a nucleoside. Thus, option (1) is incorrect.

Adenylic acid is a nucleotide. Thus, option (2) is correct.

Guanine is a nitrogenous base. Thus, option (3) is incorrect.

Guanosine is a nucleoside. Thus, option (4) is incorrect.

---

## Question2

**Ligases is a class of enzymes responsible for catalysing the linking together of two compounds.**

Which of the following bonds is not catalysed by it?

[NEET 2024 Re]

Options:

A.

C – C



B.

P – O

C.

C – O

D.

C – N

**Answer: A**

**Solution:**

Option (1) is the correct answer because, ligases are the enzymes that catalyse the linking together of 2 compounds, e.g., enzymes which catalyse joining of C – O, C – S, C – N, P – O etc. bonds.

⇒ Options (2), (3) and (4) are wrong as ligase catalyse the joining of P – O, C – O & C – N bonds.

---

## Question3

**Which of the following are not fatty acids?**

**A. Glutamic acid**

**B. Arachidonic acid**

**C. Palmitic acid**

**D. Lecithin**

**E. Aspartic acid**

**Choose the correct answer from the options given below :**

**[NEET 2024 Re]**

**Options:**

A.

C, D and E only

B.

A and B only

C.

A, D and E only

D.

B and C only

**Answer: C**

**Solution:**



Option (3) is the correct answer because glutamic acid and aspartic acid are amino acids while lecithin is a phospholipid. Palmitic acid and arachidonic acid are fatty acids.


## Question4

Which of the following graphs depicts the effect of substrate concentration on velocity of enzyme catalysed reaction?


[NEET 2024 Re]

Options:


A.




B.



C.




D.



**Answer: A**

**Solution:**

Option (1) is the correct answer because with the increase in substrate concentration, the velocity of enzymatic reaction rises at first. The reaction ultimately reaches maximum velocity ( $V_{max}$ ) which is not exceeded by any further rise in concentration of the substrate. This is because the enzyme molecules are fewer than substrate molecules and after saturation of these molecules, there are no free enzyme molecules to bind the additional substrate molecules.



Option (2) is incorrect as velocity of reaction is continuously increasing in the given graph.

In option (3) after reaching at  $V_{max}$ , velocity declines while in option (4) velocity of reaction declines from high on increasing substrate concentration. So, option (3) and (4) are incorrect.

## Question5

**Enzymes that catalyse the removal of groups from substrates by mechanisms other than hydrolysis leaving double bonds, are known as :**

**[NEET 2024 Re]**

**Options:**

A.

Transferases

B.

Oxidoreductases

C.

Dehydrogenases

D.

Lyases

**Answer: D**

**Solution:**

The correct answer is option (4) because lyases are group of enzymes that catalyse the removal of groups from substrates by mechanisms other than hydrolysis leaving double bonds.

Option (1) is incorrect because transferases are enzymes catalysing a transfer of a group G (other than hydrogen) between a pair of substrate S and S' .

Option (2) is incorrect because oxidoreductases are enzymes which catalyse oxidation-reduction between two substrates S and S'.

Option (3) is incorrect because dehydrogenases are also known as oxidoreductases.

## Question6

**Match List-I with List-II.**

	List-I		List-II
A.	Primary structure of protein	I	Human haemoglobin
B.	Secondary structure of protein	II	Disulphide bonds
C.	Tertiary structure of protein	III	Polypeptide chain
D.	Quaternary structure of protein	IV	Alpha helix and $\beta$ sheet

**Choose the correct answer from the options given below :**

**[NEET 2024 Re]**

**Options:**

A.

A-III, B-IV, C-II, D-I

B.

A-III, B-II, C-I, D-IV

C.

A-I, B-III, C-II, D-IV

D.

A-IV, B-III, C-II, D-I

**Answer: A**

**Solution:**

The correct answer is option (1) because

	List-I		List-II
A.	Primary structure of protein	III	Polypeptide chain with positional information of aminoacids
B.	Secondary structure of protein	IV	Alpha helix and $\beta$ sheet structure
C.	Tertiary structure of protein	II	Hollow woolen ball like structure with hydrogen and disulphide bonds
D.	Quaternary structure of protein	I	Assembly of more than one polypeptides, seen in adult human haemoglobin

Hence, A-III, B-IV, C-II, D-I is the correct match.

## Question7

**The cofactor of the enzyme carboxypeptidase is:**

**[NEET 2024]**

**Options:**

A. Zinc

B. Niacin

C. Flavin

D. Haem

**Answer: A**

**Solution:**

The correct answer is option (1) as the cofactor of the enzyme carboxypeptidase is zinc. Niacin is associated with coenzyme NAD and NADP. Option (4) is incorrect as haem is the prosthetic group in peroxidase and catalase.

---

## Question8

**Inhibition of Succinic dehydrogenase enzyme by malonate is a classical example of:  
[NEET 2024]**

**Options:**

- A. Cofactor inhibition
- B. Feedback inhibition
- C. Competitive inhibition
- D. Enzyme activation

**Answer: C**

**Solution:**

Correct answer is option (3) because malonate shows close structural similarity with the substrate and it competes with the substrate for the substrate binding site of the enzyme succinic dehydrogenase. Option (1), (2) and (4) are incorrect as enzyme activation, co-factor inhibition are not showing structural similarity with substrate.

---

## Question9

**Match List-I with List-II**

List-I		List-II	
A.	GLUT-4	I.	Hormone
B.	Insulin	II.	Enzyme
C.	Trypsin	III.	Intercellular ground substance
D.	Collagen	IV.	Enables glucose transport into cells

**Choose the correct answer from the options given below  
[NEET 2024]**

**Options:**

- A. A-IV, B-I, C-II, D-III
- B. A-I, B-II, C-III, D-IV
- C. A-II, B-III, C-IV, D-I
- D. A-III, B-IV, C-I, D-II

**Answer: A**

**Solution:**

Correct answer is option (1)

List-I		List-II	
A.	GLUT-4	IV.	Enables glucose transport into cells
B.	Insulin	I.	Hormone
C.	Trypsin	II.	Enzyme
D.	Collagen	III.	Intercellular ground substance

## Question10

**Match List I with List II :**

List-I		List-II	
A.	Cocaine	I.	Effective sedative in surgery
B.	Heroin	II.	Cannabis sativa
C.	Morphine	III.	Erythroxyllum
D.	Marijuana	IV.	Papaver somniferum

**Choose the correct answer from the options given below:  
[NEET 2024]**

**Options:**

- A. A-IV, B-III, C-I, D-II
- B. A-I, B-III, C-II, D-IV
- C. A-II, B-I, C-III, D-IV
- D. A-III, B-IV, C-I, D-II

**Answer: D**

**Solution:**

The correct option is (4) as

A. Cocaine - Obtained from plant *Erythroxylum coca*, stimulating action on CNS.

B. Heroin - Formed by the acetylation of morphine which is obtained from plant *Papaver somniferum*.

C. Morphine - Obtained from *Papaver somniferum*, is an effective sedative in surgery.

D. Marijuana - Obtained from *Cannabis sativa*, produces hallucinogenic effect and affects cardiovascular system of the body.

## Question 11

Match List I with List II :

List-I		List-II	
A.	Lipase	I.	Peptide bond
B.	Nuclease	II.	Ester bond
C.	Protease	III.	Glycosidic bond
D.	Amylase	IV.	Phosphodiester bond

Choose the correct answer from the options given below :  
[NEET 2024]

Options:

A. A-IV, B-II, C-III, D-I

B. A-III, B-II, C-I, D-IV

C. A-II, B-IV, C-I, D-III

D. A-IV, B-I, C-III, D-II

Answer: C

Solution:

The correct answer is option (3) as

List-I		List-II	
A.	Lipase	-	Digests ester bond found in lipids.
B.	Nuclease	-	Helps in digestion of phosphodiester bonds found in nucleic acids.
C.	Protease	-	Helps in digestion of peptide bond found in proteins.
D.	Amylase	-	Digests/breaks the glycosidic bonds found in carbohydrates i.e., digest starch into smaller molecules, ultimately yielding maltose, which in turn is cleaved into two glucose molecules by maltase.

## Question 12

Regarding catalytic cycle of an enzyme action, select the correct sequential steps :





- A. Substrate enzyme complex formation.**
  - B. Free enzyme ready to bind with another substrate.**
  - C. Release of products.**
  - D. Chemical bonds of the substrate broken.**
  - E. Substrate binding to active site.**
- Choose the correct answer from the options given below :**  
**[NEET 2024]**

**Options:**

- A. E, A, D, C, B
- B. A, E, B, D, C
- C. B, A, C, D, E
- D. E, D, C, B, A

**Answer: A**

**Solution:**

The correct answer is option (1) which is E, A, D, C, B.

The catalytic cycle of an enzyme action can be described in the following steps.

- (1) First, the substrate binds to the active site of the enzyme, fitting into the active site.
- (2) The binding of the substrate induces the enzyme to alter its shape, fitting more tightly around the substrate.
- (3) The active site of the enzyme, now in close proximity of the substrate breaks the chemical bonds of the substrate and the new enzyme-product complex is formed.
- (4) The enzyme releases the products of the reaction and the free enzyme is ready to bind to another molecule of the substrate and run through the catalytic cycle once again.

Options (2), (3) and (4) are incorrect as the steps mentioned are in the wrong sequence.

## Question 13

**Match List-I with List-II**

List - I		List - II	
(A)	Protein	(I)	C=C double bonds
(B)	Unsaturated fatty acid	(II)	Phosphodiester bond
(C)	Nucleic acid	(III)	Glycosidic bonds
(D)	Polysaccharide	(IV)	Peptide bonds

**Choose the correct answer from the options given below :**  
**[NEET 2023 mpr]**

**Options:**

- A. (A)-(II), (B)-(I), (C)-(IV), (D)-(III)
- B. (A)-(IV), (B)-(III), (C)-(I), (D)-(II)
- C. (A)-(IV), (B)-(I), (C)-(II), (D)-(III)

D. (A)-(I), (B)-(IV), (C)-(III), (D)-(II)

**Answer: C**

**Solution:**

The correct matchings are :

(A) Protein - This is a polymer made up of amino acids linked by peptide bonds. So, (A) matches with (IV) Peptide bonds.

(B) Unsaturated fatty acid - These have one or more C=C double bonds in their hydrocarbon chain. So, (B) matches with (I) C=C double bonds.

(C) Nucleic acid - These are polymers made up of nucleotides linked by phosphodiester bonds. So, (C) matches with (II) Phosphodiester bond.

(D) Polysaccharide - These are polymers made up of monosaccharides linked by glycosidic bonds. So, (D) matches with (III) Glycosidic bonds.

So, the correct answer is Option C : (A)-(IV), (B)-(I), (C)-(II), (D)-(III).

---

## Question14

**Inulin is a polymer of :  
[NEET 2023 mpr]**

**Options:**

A. Fructose

B. Galactose

C. Amino acids

D. Glucose

**Answer: A**

**Solution:**

Inulin is a polymer of fructose. Therefore, the correct answer is Option A : Fructose.

---

## Question15

**Which of the following is not a secondary metabolite?  
[NEET 2023 mpr]**

**Options:**

A. Curcumin

B. Morphine

C. Anthocyanin



D. Lecithin

**Answer: D**

**Solution:**

Secondary metabolites are organic compounds produced by organisms that are not directly involved in the normal growth, development, or reproduction of the organism. Examples include antibiotics, pigments, and toxins.

Curcumin, morphine, and anthocyanin are all examples of secondary metabolites.

Lecithin, however, is a type of phospholipid, which is a primary metabolite as it is directly involved in the normal growth and development of cells by being a key component of cell membranes.

---

## Question16

**Cellulose does not form blue colour with iodine because  
[NEET 2023]**

**Options:**

- A. It is a helical molecule
- B. It does not contain complex helices and hence cannot hold iodine molecules
- C. It breaks down when iodine reacts with it
- D. It is a disaccharide

**Answer: B**

**Solution:**

**Solution:**

Option (2) is the correct answer because cellulose does not contain complex helices and hence cannot hold iodine molecules.

Option (1), (3) and (4) are not correct as cellulose is a polysaccharide.

---

## Question17

**Melonate inhibits the growth of pathogenic bacteria by inhibiting the  
activity of  
[NEET 2023]**

**Options:**

- A. Amylase
- B. Lipase
- C. Dinitrogenase
- D. Succinic dehydrogenase



**Answer: D**

### **Solution:**

Option (4) is correct answer of this question because malonate is a competitive inhibitor of enzyme succinate dehydrogenase.

Inhibition of succinic dehydrogenase by malonate occurs due to close resemblance of malonate with substrate succinate in structure. Competitive inhibitors are often used in the control of bacterial pathogens.

---

## **Question18**

**Given below are two statements :**

**Statement I : Low temperature preserves the enzyme in a temporarily inactive state whereas high temperature destroys enzymatic activity because proteins are denatured by heat.**

**Statement II : When the inhibitor closely resembles the substrate in its molecular structure and inhibits the activity of the enzyme, it is known as competitive inhibitor.**

**In the light of the above statements, choose the correct answer from the options given below :**

**[NEET 2023]**

**Options:**

- A. Both Statement I and Statement II are false.
- B. Statement I is true but Statement II is false.
- C. Statement I is false but Statement II is true.
- D. Both Statement I and Statement II are true.

**Answer: D**

### **Solution:**

**Solution**

The correct answer is option (4) as low temperature preserves the enzyme in a temporarily inactive state whereas high temperature destroys enzymatic activity because proteins are denatured by heat.

- Competitive inhibitor due to its close structural similarity with the substrate, competes with the substrate for the substrate-binding site of the enzyme.
- 

## **Question19**

**Given below are two statements:**

**Statement I: A protein is imagined as a line, the left end represented by first amino acid (C-terminal) and the right end represented by last amino acid ( N-terminal).**

**Statement II: Adult human haemoglobin, consists of 4 subunits (two**



subunits of  $\alpha$  type and two subunits of  $\beta$  type.)

In the light of the above statements, choose the correct answer from the options given below

[NEET 2023]

**Options:**

- A. Both Statement I and Statement II are false.
- B. Statement I is true but Statement II is false.
- C. Statement I is false but Statement II is true.
- D. Both Statement I and Statement II are true

**Answer: C**

**Solution:**

The correct answer is option (3) as a protein is imagined as a line, the left end represented by the first amino acid and the right end is represented by the last amino acid. The first amino acid is also called N-terminal amino acid. The last amino acid is called the C-terminal amino acid.

## Question20

List-I	List-II
(a) Adenine	(i) Pigment
(b) Anthocyanin	(ii) Polysaccharide
(c) Chitin	(iii) Alkaloid
(d) Codeine	(iv) Purine

Choose the correct answer from the options given below .  
[NEET Re-2022]

**Options:**

- A. (a) - (i), (b) - (iv), (c) - (iii), (d) - (ii)
- B. (a) - (iv), (b) - (i), (c) - (ii), (d) - (iii)
- C. (a) - (iv), (b) - (iii), (c) - (ii), (d) - (i)
- D. (a) - (iii), (b) - (i), (c) - (iv), (d) - (ii)

**Answer: B**

**Solution:**

Adenine: Purine (double ringed structure) found in both DNA and RNA)



Anthocyanin Pigment (secondary metabolite)

Chitin Homopolysaccharide of  $\text{N-acetylglucosamine}$  found in fungal cell wall and exoskeleton of Arthropods

Codeine Alkaloid (secondary metabolite)

---

## Question21

**Primary proteins are also called as polypeptides because:  
[NEET Re-2022]**

**Options:**

- A. They can assume many conformations
- B. They are linear chains
- C. They are polymers of peptide monomers
- D. Successive amino acids are joined by peptide bonds

**Answer: D**

**Solution:**

**Solution:**

Primary proteins are the linear chains of amino acids, joined by peptide bonds.

---

## Question22

**Given below are two statements : Statement I : Amino acids have a property of ionizable nature of  $-\text{NH}_2$  and  $-\text{COOH}$  groups, hence have different structures at different pH.**

**Statement II : Amino acids can exist as Zwitterionic form at acidic and basic pH. In the Light of the above statements, choose the most appropriate answer from the options given below :**

**[NEET Re-2022]**

**Options:**

- A. Statement I is incorrect but Statement II is correct
- B. Both Statement I and Statement II are correct
- C. Both Statement I and Statement II are incorrect
- D. Statement I is correct but Statement II is incorrect

**Answer: B**

**Solution:**



Statement I :Correct

Statement II: Correct, Zwitterion formation takes place at isoelectric point that can be at acidic or basic pH.

---

## Question23

**In the enzyme which catalyses the breakdown of:**



**the prosthetic group is:**

**[NEET Re-2022]**

**Options:**

- A. Niacin
- B. Nicotinamide adenine dinucleotide
- C. Haem
- D. Zinc

**Answer: C**

**Solution:**

**Solution:**

Peroxidase and catalase enzymes catalyze the breakdown of Hydrogen peroxide to water and oxygen, Haem is the prosthetic group.

---

## Question24

**Read the following statements on lipids and find out correct set of statements:**

- (a) Lecithin found in the plasma membrane is a glycolipid
- (b) Saturated fatty acids possess one or more  $\text{C} = \text{C}$  bonds
- (c) Gingly oil has lower melting point, hence remains as oil in winter
- (d) Lipids are generally insoluble in water but soluble in some organic solvents
- (e) When fatty acid is esterified with glycerol, monoglycerides are formed

**Choose the correct answer from the option given below:**

**[NEET-2022]**

**Options:**

- A. (a), (b) and (c) only
- B. (a), (d) and (e) only
- C. (c), (d) and (e) only
- D. (a), (b) and (d) only



**Answer: C**

**Solution:**

Option (3) is the correct answer because statements (c), (d) and (e) are correct as oils have lower melting point and hence remain oil in winters. Lipids are generally insoluble in water but soluble in some organic solvents.





Option (1), (2) and (4) are incorrect because statements (a) and (b) are incorrect. Lecithin is a type of phospholipid found in plasma membrane. Saturated fatty acids are without double bond.

## Question25

**A dehydration reaction links two glucose molecules to product maltose. If the formula for glucose is  $C_6H_{12}O_6$  then what is the formula for maltose?**

**[NEET-2022]**

**Options:**




**Answer: C**

**Solution:**

**Solution:**

Option (3) is correct because maltose is a disaccharide formed by dehydration process i.e., synthesis by elimination of one water molecule to form a glycosidic bond in between two glucose molecules. So, its molecular formula is.



## Question26

**Which of the following are not secondary metabolites in plants?**  
**[NEET 2021]**

**Options:**

A. Morphine, codeine

B. Amino acids, glucose

C. Vinblastin, curcumin

D. Rubber, gums





**Answer: B**

### Solution:

The correct option is (2)

- Amino acids and glucose are included under the category of primary metabolites as they have identifiable functions and play known roles in normal physiological processes.
- Rubber, gums, morphine, codeine, vinblastine and curcumin are included under the category of secondary metabolites as their role or functions in host organisms is not known yet. However, many of them are useful to human welfare.

## Question27

### Match List-I with List-II.

	List-I		List-II
(a)	Protein	(i)	C = C double bonds
(b)	Unsaturated fatty acid	(ii)	Phosphodiester bonds
(c)	Nucleic acid	(iii)	Glycosidic bonds
(d)	Polysaccharide	(iv)	Peptide bonds

**Choose the correct answer from the options given below.  
[NEET 2021]**

### Options:

- A. (a)-(iv) (b)-(i) (c)-(ii) (d)-(iii)
- B. (a)-(i) (b)-(iv) (c)-(iii) (d)-(ii)
- C. (a)-(ii) (b)-(i) (c)-(iv) (d)-(iii)
- D. (a)-(iv) (b)-(iii) (c)-(i) (d)-(ii)

**Answer: A**

### Solution:

#### Solution:

- In a polypeptide or a protein, amino acids are linked by a peptide bond which is formed when the carboxyl (-COOH) group of one amino acid reacts with amino (-NH<sub>2</sub>) group of the next amino acid with the elimination of a water moiety.
- Unsaturated fatty acids are with one or more C = C double bonds.
- In nucleic acids, a phosphate moiety links the 3'-carbon of one sugar of one nucleotide to the 5'-carbon of the sugar of the succeeding nucleotide. The bond between the phosphate and hydroxyl group is an ester bond. As there is one such ester bond on either side, it is called phosphodiester bond.
- In a polysaccharide, the individual monosaccharides are linked by a glycosidic bond.

## Question28

**Following are the statements with reference to 'lipids'.**

**(a) Lipids having only single bonds are called unsaturated fatty acids**

**(b) Lecithin is a phospholipid.**

**(c) Trihydroxy propane is glycerol.**

**(d) Palmitic acid has 20 carbon atoms including carboxyl carbon.**

**(e) Arachidonic acid has 16 carbon atoms.**

**Choose the correct answer from the options given below.**

**[NEET 2021]**

**Options:**

A. (a) and (b) only

B. (c) and (d) only

C. (b) and (c) only

D. (b) and (e) only

**Answer: C**

**Solution:**

**Solution:**

● The correct option is (3) because lipids having only single bonds are called saturated fatty acids and lipids having one or more C = C double bonds are called unsaturated fatty acids.

● Palmitic acid has 16 carbon atoms including carboxyl carbon.

● Arachidonic acid has 20 carbon atoms including the carboxyl carbon.

● Lecithin is a phospholipid found in cell membrane.

● Glycerol has 3 carbons, each bearing a hydroxyl (-OH) group.

.....

## Question29

**Identify the substances having glycosidic bond and peptide bond, respectively in their structure**

**[NEET-2020]**

**Options:**

A. Glycerol, trypsin

B. Cellulose, lecithin

C. Inulin, insulin

D. Chitin, cholesterol

**Answer: C**

**Solution:**



Inulin is a fructan (polysaccharide of fructose). Adjacent fructose units are linked through glycosidic bond.

Insulin is a protein composed of 51 amino acids. Adjacent amino acids are attached through peptide bond.

---

## Question30

### Match the following

(a) Inhibitor of catalytic activity	(i) Ricin
(b) Possess peptide bonds	(ii) Malonate
(c) Cell wall material in fungi	(iii) Chitin
(d) Secondary metabolite	(iv) Collagen

Choose the correct option from the following

	(a)	(b)	(c)	(d)
(1)	(iii)	(i)	(iv)	(ii)
(2)	(iii)	(iv)	(i)	(ii)
(3)	(ii)	(iii)	(i)	(iv)
(4)	(ii)	(iv)	(iii)	(i)

### [NEET-2020]

#### Options:

- A. a
- B. b
- C. c
- D. d

**Answer: D**

#### Solution:

##### Solution:

Option (4) is the correct answer because Malonate is the competitive inhibitor of catalytic activity of succinic dehydrogenase, so (a) matches with (ii) in column II.

Collagen is proteinaceous in nature and possesses peptide bonds, so (b) matches with (iv) in column II.

Chitin is a homopolymer present in the cell wall of fungi and exoskeleton of arthropods, so, (c) matches with (iii) in column II.

Abrin and Ricin are toxins, secondary metabolites, so (d) in column I matches with (i) in column II.

---

## Question31

**"Ramachandran plot" is used to confirm the structure of :-**

**[NEET OD 2019]**



**Options:**

- A. RNA
- B. Proteins
- C. Triacylglycerides
- D. DNA

**Answer: B**

---

## Question32

**Purines found both in DNA and RNA are [NEET 2019]**

**Options:**

- A. Adenine and guanine
- B. Guanine and cytosine
- C. Cytosine and thymine
- D. Adenine and thymine

**Answer: A**

**Solution:**

**Solution:**

Purines found both in DNA and RNA are Adenine and guanine

---

## Question33

**Prosthetic groups differ from co-enzymes in that :- [NEET OD 2019]**

**Options:**

- A. they require metal ions for their activity.
- B. they (prosthetic groups) are tightly bound to apoenzymes.
- C. their association with apoenzymes is transient.
- D. they can serve as co-factors in a number of enzyme-catalyzed reactions.

**Answer: B**

---

## Question34

## Which of the following glucose transporters is insulin-dependent? [NEET 2019]

**Options:**

- A. GLUT II
- B. GLUT III
- C. GLUT IV
- D. GLUT I

**Answer: C**

**Solution:**

**Solution:**

GLUT-IV is insulin dependent and is responsible for majority of glucose transport into muscle and adipose cells in anabolic conditions. Whereas GLUT-I is insulin independent and is widely distributed in different

---

## Question35

**Consider the following statement :**

**(A) Coenzyme or metal ion that is tightly bound to enzyme protein is called prosthetic group.**

**(B) A complete catalytic active enzyme with its bound prosthetic group is called apoenzyme. Select the correct option.**

**[NEET 2019]**

**Options:**

- A. (A) is true but (B) is false.
- B. Both (A) and (B) are false.
- C. (A) is false but (B) is true.
- D. Both (A) and (B) are true.

**Answer: B**

**Solution:**

**Solution:**

“Coenzymes, metal ions and prosthetic groups that is bound to enzyme protein is called co- factor. Prosthetic groups are tightly bound to enzyme where as metal ions and coenzymes are not tightly bound to enzyme.”

---

## Question36

**The two functional groups characteristic of surgars are  
[NEET 2018]**

**Options:**

- A. hydroxyl and methyl
- B. carbonyl and methyl
- C. carbonyl and hydroxyl
- D. carbonyl and phosphate

**Answer: C**

**Solution:****Solution:**

Sugar is a common term used to denote carbohydrate.

Carbohydrates are polyhydroxy aldehyde, ketone or their derivatives, which means they have carbonyl and hydroxyl groups.

---

## Question37

**Which one of the following statements is correct,with reference to enzymes ?**

**[NEET 2017]**

**Options:**

- A. Holoenzyme = Apoenzyme + Coenzyme
- B. Coenzyme = Apoenzyme + Holoenzyme
- C. Holoenzyme = Coenzyme + Co-factor
- D. Apoenzyme = Holoenzyme + Coenzyme

**Answer: A**

**Solution:****Solution:**

Holoenzyme is conjugated enzyme in which protein part is apoenzyme while non-protein is cofactor.

Coenzyme are also organic compounds but their association with apoenzyme is only transient and serve as cofactors.

---

## Question38

**Which of the following are not polymeric ?**

**[NEET 2017]**

**Options:**

- A. Proteins
- B. Polysaccharides



C. Lipids

D. Nucleic acids

**Answer: A**

---

## Question39

**A non-proteinaceous enzyme is  
[NEET 2016 P2]**

**Options:**

A. deoxyribonuclease

B. lysozyme

C. Ribozyme

D. ligase

**Answer: B**

**Solution:**

**Solution:**

Ribozyme is non proteinaceous enzyme as it is 23 rRNA acts as a catalyst during protein synthesis.

---

## Question40

**Which of the following is the least likely to be involved in stabilizing the  
three-dimensional folding of most proteins?  
[NEET 2016 P2]**

**Options:**

A. Ester bonds

B. Hydrogen bonds

C. Electrostatic interaction

D. Hydrophobic interaction

**Answer: C**

**Solution:**


**Solution:**

Ester bonds are formed in nucleic acids and lipids, but not proteins.

---

## Question41

Which of the following describes the given graph correctly?



[NEET 2016 P2]

Options:

- A. Exothermic reaction with energy A in absence of enzyme and B in presence of enzyme
- B. Endothermic reaction with energy A in presence of enzyme and B in absence of enzyme
- C. Exothermic reaction with energy A in presence of enzyme and B in absence of enzyme
- D. Endothermic reaction with energy A in absence of enzyme and B in presence of enzyme.

Answer: B

---

## Question42

A typical fat molecule is made up of :

[NEET 2016 P1]

Options:

- A. Three glycerol and three fatty acid molecules
- B. Three glycerol molecules and one fatty acid molecule
- C. One glycerol and three fatty acid molecules
- D. One glycerol and one fatty acid molecule

Answer: B

Solution:

Solution:

fat is a triglyceride which is made up of 3 molecules of fatty acids and one molecule of glycerol

---

## Question43

Which one of the following statements is wrong?

[NEET 2016 P1]

Options:



- A. Glycine is a sulphur containing amino acid.
- B. Sucrose is a disaccharide.
- C. Cellulose is a polysaccharide.
- D. Uracil is a pyrimidine.

**Answer: A**

**Solution:**

**Solution:**

Glycine is the simplest amino acid which is devoid of sulphur content

-----

## Question44

**The chitinous exoskeleton of arthropods is formed by the polymerisation of**  
**[NEET 2015]**

**Options:**

- A. N-acetyl glucosamine
- B. lipoglycans
- C. keratin sulphate and chondrotin sulphate
- D. D - glucosamine.

**Answer: A**

**Solution:**

**Solution:**

Chitin is a structural polysaccharide that constitutes the exoskeleton of arthropods. It is a complex carbohydrate in which N-acetyl glucosamine monomers are joined together by (1, 4) $\beta$ -linkages. Chitinous exoskeleton provides strength and elasticity to arthropods.

-----

## Question45

**Which of the following biomolecules does have a phosphodiester bond?**  
**[NEET 2015]**

**Options:**

- A. Amino acids in a polypeptide
- B. Nucleic acids in a nucleotide
- C. Fatty acids in a diglyceride
- D. Monosaccharides in a polysaccharide



**Answer: B**

**Solution:**

Nucleic acids have phosphodiester bond in a nucleotide.

---

## Question46

**Which of the following statements is incorrect ?  
[NEET 2015 C]**

**Options:**

- A. In competitive inhibition, the inhibitor molecule is not chemically changed by the enzyme.
- B. The competitive inhibitor does not affect the rate of break down of the enzyme-substrate complex
- C. The presence of the competitive inhibitor decreases the  $K_m$  of the enzyme for the substrate.
- D. A competitive inhibitor reacts reversibly with the enzyme to form an enzyme-inhibitor complex

**Answer: C**

---

## Question47

**Select the option which is not correct with respect to enzyme action:  
[NEET 2014]**

**Options:**

- A. Substrate binds with enzyme at its active site.
- B. Addition of lot of succinate does not reverse the inhibition of succinic dehydrogenase by malonate
- C. A non-competitive inhibitor binds the enzyme at a site distinct from that which binds the substrate
- D. Malonate is a competitive inhibitor of succinic dehydrogenase

**Answer: B**

**Solution:**

Inhibition of succinic dehydrogenase by malonate is an example of competitive inhibition. This is a reversible reaction. On increasing the substrate (succinate) concentration the effect of inhibitor is removed and  $V_{max}$  remains the same.

---

## Question48

**Which one of the following is a non-reducing carbohydrate?  
[NEET 2014]**

**Options:**

- A. Maltose
- B. Sucrose
- C. Lactose
- D. Ribose 5-phosphate

**Answer: B**

**Solution:**

**Solution:**

Lactose, Maltose, Ribose 5-phosphate all are reducing sugars.

---

## Question49

**A phosphoglyceride is always made up of  
(NEET 2013)**

**Options:**

- A. a saturated or unsaturated fatty acid esterified to a glycerol molecule to which a phosphate group is also attached.
- B. a saturated or unsaturated fatty acid esterified to a phosphate group which is also attached to a glycerol molecule.
- C. only a saturated fatty acid esterified to a glycerol molecule to which a phosphate group is also attached.
- D. only an unsaturated fatty acid esterified to a glycerol molecule to which a phosphate group is also attached.

**Answer: A**

**Solution:**

**Solution:**

(a) : Phosphoglycerides are the triesters of fatty acids (either saturated or unsaturated) and glycerol to which a phosphate group is also attached.

---



## Question50

**Macromolecule chitin is (NEET 2013)**

**Options:**

- A. sulphur containing polysaccharide
- B. simple polysaccharide
- C. nitrogen containing polysaccharide
- D. phosphorous containing polysaccharide.

**Answer: C**

**Solution:**

(c) : Chitin is a structural polysaccharide that constitutes the exoskeleton of arthropods. It is a complex carbohydrate in which N -acetyl glucosamine monomers are joined together by (1,4)  $\beta$ -linkages. Chitinous exoskeleton provides strength and elasticity to arthropods.

---

## Question51

**Transition state structure of the substrate formed during an enzymatic reaction is (NEET 2013)**

**Options:**

- A. transient and unstable
- B. permanent and stable
- C. transient but stable
- D. permanent but unstable.

**Answer: A**

**Solution:**

**Solution:**

(a) : Transition state is formation of unstable intermediate structural state. During this, substrate bonds are broken and new bonds are established that transform the substrate molecules into products. This state is transient and highly unstable.

---

## Question52

**The essential chemical components of many coenzymes are (NEET 2013)**



**Options:**

- A. carbohydrates
- B. vitamins
- C. proteins
- D. nucleic acids.

**Answer: B****Solution:**

(b) : Coenzyme is the non protein organic group which gets attached to the apoenzyme to form holoenzyme or conjugate enzyme. It helps in removing a product of chemical reaction besides bringing contact between the substrate and the enzyme. Most of the coenzymes are made of water soluble vitamins B and C, e. g., thiamine, riboflavin, nicotinamide, pyridoxine.

## Question53

**Which of the following statements about enzymes is wrong?  
(KN NEET 2013)**

**Options:**


- A. Enzymes are denatured at high temperatures.
- B. Enzymes are mostly proteins but some are lipids also.
- C. Enzymes are highly specific.
- D. Enzymes require optimum pH and temperature for maximum activity.

**Answer: B****Solution:****Solution:**

(b) : Enzymes are mostly proteins but some are RNA (ribozymes). No lipid working as enzymes are known.

## Question54

**The figure shows a hypothetical tetrapeptide portion of a protein with parts labelled A -D. Which one of the following options is correct?**

**(KN NEET 2013)**

**Options:**

- A. D is the acidic amino acid-glutamic acid.
- B. C is an aromatic amino acid-tryptophan.
- C. A is the C-terminal amino acid and D is N-terminal amino acid.
- D. A is a sulphur containing amino acid methionine.

**Answer: A**

**Solution:**

**Solution:**

(a) Glutamic acid and aspartic acid are acidic amino acids with two carboxylic groups and one amino group.

---

## Question55

**Uridine, present only in RNA is a  
(KN NEET 2013)**

**Options:**

- A. nucleoside
- B. nucleotide
- C. purine
- D. pyrimidine.

**Answer: A**

**Solution:**

**Solution:**

(a) : The combination of pentose sugar with nitrogenous bases (purines or pyrimidines) is called nucleoside. Examples are adenosine, guanosine, cytidine, thymidine and uridine.

---

## Question56

**Which one out of A – D given below correctly represents the structural formula of the basic amino acid?**

A	B	C	D
$  \begin{array}{c}  \text{NH}_2 \\    \\  \text{H}-\text{C}-\text{COOH} \\    \\  \text{CH}_2 \\    \\  \text{CH}_2 \\    \\  \text{C} \\  // \quad \backslash \\  \text{O} \quad \text{OH}  \end{array}  $	$  \begin{array}{c}  \text{NH}_2 \\    \\  \text{H}-\text{C}-\text{COOH} \\    \\  \text{CH}_2 \\    \\  \text{OH}  \end{array}  $	$  \begin{array}{c}  \text{CH}_2\text{OH} \\    \\  \text{CH}_2 \\    \\  \text{CH}_2 \\    \\  \text{NH}_2  \end{array}  $	$  \begin{array}{c}  \text{NH}_2 \\    \\  \text{H}-\text{C}-\text{COOH} \\    \\  \text{CH}_2 \\    \\  \text{CH}_2 \\    \\  \text{CH}_2 \\    \\  \text{CH}_2 \\    \\  \text{CH}_2 \\    \\  \text{NH}_2  \end{array}  $

**(2012)**

**Options:**

- A. C
- B. D
- C. A
- D. B

**Answer: B**

**Solution:**

(b) : Basic amino acids have an additional amino group without forming amides thus they are diamino monocarboxylic acids e.g., arginine, lysine, etc.

## Question57

**Which one is the most abundant protein in the animal world?  
(2012)**

**Options:**

- A. Trypsin
- B. Haemoglobin
- C. Collagen
- D. Insulin

**Answer: C**


**Solution:**

(c) : Collagen is an insoluble fibrous protein found extensively in the connective tissue of skin, tendons and bone. Collagen accounts for over 30% of the total body proteins of mammals and it is the most abundant animal protein.

## Question58

The given diagrammatic representation shows one of the categories of small molecular weight organic compounds in the living tissues.

Identify the category shown and the one blank component "X" in it.



	Category	Component
(a)	Cholesterol	Guanine
(b)	Amino acid	$NH_2$
(c)	Nucleotide	Adenine
(d)	Nucleoside	Uracil

(2012)

Options:

- A. (a)
- B. (b)
- C. (c)
- D. (d)

Answer: D

Solution:

Solution:

(d) : The given structure corresponds with the structure of ribose sugar. As it lacks a phosphoric acid hence it can be a nucleoside not a nucleotide.

## Question59

Which one of the following is wrong statement?

(2012)

Options:

- A. Anabaena and Nostoc are capable of fixing nitrogen in free-living state also.
- B. Root nodule forming nitrogen fixers live as aerobes under free-living conditions.
- C. Phosphorus is a constituent of cell membranes, certain nucleic acids and all proteins.
- D. Nitrosomonas and Nitrobacter are chemoautotrophs



**Answer: C**

**Solution:**

(c) : Phosphorus is present in plasma membrane in the form of phospholipid bilayer. It is an essential component of all nucleic acids not 'certain' nucleic acids. Moreover, phosphorus is never found in proteins.

---

## Question60

**Which one of the following biomolecules is correctly characterized? (Mains 2012)**

**Options:**

- A. Lecithin - a phosphorylated glyceride found in cell membrane.
- B. Palmitic acid-an unsaturated fatty acid with 18 carbon atoms.
- C. Adenylic acid - adenosine with a glucose phosphate molecule.
- D. Alanine amino acid - contains an amino group and an acidic group anywhere in the molecule.

**Answer: A**

**Solution:**


**Solution:**

(a) Palmitic acid is one of the most common, saturated fatty acids found in animals and plants. It has 16 carbons including the carboxyl carbon. Adenylic acid is a nucleotide consisting of adenine, ribose or deoxyribose, and a phosphate group. It is a constituent of DNA or RNA. It is also called adenosine monophosphate. Amino acids are organic acids (with carboxylic group COOH) having amino group ( $-NH_2$ ) generally attached to Carbon or carbon next to carboxylic group. The carbon also bears a variable alkyl group (R) or hydrogen or hydrocarbon. In alanine (R) is represented by methyl group.

---

## Question61

**The curve given below shows enzymatic activity in relation to three conditions ( pH , temperature and substrate concentration). What do the two axes ( x and y ) represent?**



	x -axis	y -axis
(a)	enzymatic activity	pH
(b)	temperature	enzyme activity
(c)	substrate concentration	enzymatic activity
(d)	enzymatic activity	temperature

**(2011)**

**Options:**

- A. (a)
- B. (b)
- C. (c)
- D. (d)

**Answer: B**


**Solution:**

**Solution:**

(b) : Enzymes generally function in a narrow range of temperature and pH. Each enzyme shows its highest activity at a particular temperature and pH called the optimum temperature and optimum pH. Activity declines both below and above the optimum values. X-axis always represents temperature or pH and Y axis represents enzyme activity.

## Question62

**Which one of the following structural formulae of two organic compounds is correctly identified along with its related function?**



**(2011)**

**Options:**


- A. B : Adenine -A nucleotide that makes up nucleic acids
- B. A : Triglyceride - Major source of energy
- C. B : Uracil - A component of DNA

D. A : Lecithin- A component of cell membrane

**Answer: D**

**Solution:**

(d) : 'A' is a structural formula of lecithin. It is probably the most common phospholipid. Phospholipids are major components in the lipid bilayers of cell membrane.



---

## Question63

**Three of the following statements about enzymes are correct and one is wrong. Which one is wrong? (Mains 2010)**

**Options:**

- A. Enzymes require optimum pH for maximal activity.
- B. Enzymes are denatured at high temperature but in certain exceptional organisms they are effective even at temperatures  $80^{\text{circ}} - 90^{\text{circ}} \text{mathrm C}$
- C. Enzymes are highly specific.
- D. Most enzymes are proteins but some are lipids.

**Answer: D**

**Solution:**


**Solution:**

(d) : Enzymes are mostly proteins but some are RNA (ribozymes). No lipid working as enzymes are known.

---

## Question64

**The figure given below shows the conversion of a substrate into product by an enzyme. In which one of the four options (a-d) the components of reaction labelled as A, B, C and D are identified correctly?**



	A	B	C	D
(a)	Potential energy	Transition state	Activation energy with enzyme	Activation energy without enzyme
(b)	Transition state	potential energy	Activation energy without enzyme	Activation energy with enzyme
(c)	Potential energy	Transition state	Activation energy with enzyme	Activation energy without enzyme
(d)	Activation energy wit enzyme	Transition state	Activation energy without enzyme	Potential energy

**(Mains 2010)**

**Options:**

- A. (a)
- B. (b)
- C. (c)
- D. (d)

**Answer: B**

## Question65

**Carbohydrates are commonly found as starch in plants storage organs. Which of the following five properties of starch (1-5) make it useful as a storage material?**

- (1) Easily translocated**
- (2) Chemically non-reactive**
- (3) Easily digested by animals**
- (4) Osmotically inactive**
- (5) Synthesized during photosynthesis**

**The useful properties are (2009)**

**Options:**



- A. (1),(3) and (5)
- B. (1) and (5)
- C. (2) and (3)
- D. (2) and (4)

**Answer: D**

**Solution:**

**Solution:**

(d) : Starch is the major storage carbohydrate of plants. In most plant species it is accumulated in the chloroplast of leaves, whereas in storage organ it accumulates in amyloplast as reserve starch. It is the osmotically inactive form of photosynthetic product and is a hexosan polysaccharide made of large number of glucose unit so, chemically non reactive.

---

## Question66

**Cellulose is the major component of cell walls of (2008)**

**Options:**

- A. Pseudomonas
- B. Saccharomyces
- C. Pythium
- D. Xanthomonas.

**Answer: C**

**Solution:**

**Solution:**

(c) : The cell wall of most fungi consist of chitin or cellulose. In Pythium, the hyphal wall contains cellulose whereas, in yeast the cell wall is thin and is composed of chitin in combination with other compounds (carbohydrates, glucan and mannan). The bacterial cell wall contains N -acetyl glucosamine and N-acetyl muramic acid.

---

## Question67

**A competitive inhibitor of succinic dehydrogenase is (2008)**

**Options:**

- A. m - ketoglutarate
- B. malate
- C. malonate



D. oxaloacetate.

**Answer: 0**

**Solution:**

**Solution:**

(c, d) Malonate or oxaloacetate, which resemble succinate in structure and inhibit the activity of succinate dehydrogenase. Such competitive inhibitors are often used in control of bacterial pathogen.

---

## Question68

**Modern detergents contain enzyme preparations of (2008)**

**Options:**

- A. thermoacidophiles
- B. thermophiles
- C. acidophiles
- D. alkaliphiles.

**Answer: D**

**Solution:**

**Solution:**

(d) : Modern detergents contain enzyme preparation of alkaline protease which are called alkaliphiles, for removing protein stain.

---

## Question69

**About 98 percent of the mass of every living organism is composed of just six elements including carbon, hydrogen, nitrogen, oxygen and (2007)**

**Options:**

- A. sulphur and magnesium
- B. magnesium and sodium
- C. calcium and phosphorus
- D. phosphorus and sulphur

**Answer: D**

**Solution:**



(d) : Living organisms requires 6 elements in relatively large amounts. C, H, O, N, P, S. These elements contribute to the structural organization of living organisms.

---

## Question70

**An organic substance bound to an enzyme and essential for its activity is called (2006)**

**Options:**

- A. isoenzyme
- B. coenzyme
- C. holoenzyme
- D. apoenzyme.

**Answer: B**

**Solution:**

**Solution:**

(b) : Enzymes are simple if they are made of only proteins ( e.g., pepsin, amylase etc.) while conjugate enzymes have an additional non-protein cofactor which may be organic or inorganic. Loosely attached organic cofactor is coenzyme. It plays an accessory role in enzyme catalyzed processes often by acting as a donor or acceptor of a substance involved in the reaction. ATP and NAD are common coenzymes.

---

## Question71

**The catalytic efficiency of two different enzymes can be compared by the (2006)**

**Options:**

- A. formation of the product
- B. pH of optimum value
- C.  $K_m$  value
- D. molecular size of the enzyme.

**Answer: C**

**Solution:**

(c) :  $K_m$  value or Michaelis constant is defined as the substrate concentration at which half of the enzyme molecules are forming (ES) complex or concentration of the substrate when the velocity of the enzyme reaction is half the maximal possible. The  $K_m$  varies from enzyme to enzyme and is used in characterizing the different enzymes. A smaller  $K_m$  value indicates greater affinity of the enzyme for its substrate, hence, shows a quicker reaction.  $K_m$  value is a constant characteristic of an enzyme for its conversion of a substrate.



## Question72

**Which one of the following statements regarding enzyme inhibition is correct?  
(2005)**

**Options:**

- A. Competitive inhibition is seen when a substrate competes with an enzyme for binding to an inhibitor protein.
- B. Competitive inhibition is seen when the substrate and the inhibitor compete for the active site on the enzyme.
- C. Non-competitive inhibition of an enzyme can be overcome by adding large amount of substrate.
- D. Non-competitive inhibitors often bind to the enzyme irreversibly.

**Answer: B**

**Solution:**

**Solution:**

(b): Competitive inhibition is a reversible inhibition where inhibitor competes with the normal substrate for the active site of enzyme. A competitive inhibitor is usually chemically similar to the normal substrate and therefore, fits into the active site of an enzyme and binds with it. The inhibition is thus due to substrate analogue. The enzyme, now cannot act upon the substrate and reaction products are not formed. E.g., the activity of succinate dehydrogenase is inhibited by malonate.  $K_m$  value or Michaelis constant is defined as the substrate concentration at which half of the enzyme molecules are forming enzyme substrate (ES) complex, or concentration of the substrate when the velocity of the enzyme reaction is half the maximal possible. A smaller  $K_m$  value indicates greater affinity of the enzyme for its substrate, hence, shows a quicker reaction. The competitive inhibitor decreases the affinity of enzyme for substrate, thus increases the  $K_m$  value.

---

## Question73

**Enzymes, vitamins and hormones can be classified into a single category of biological chemicals, because all of these  
(2005)**

**Options:**

- A. help in regulating metabolism
- B. are exclusively synthesized in the body of a living organism as at present
- C. are conjugated proteins
- D. enhance oxidative metabolism.

**Answer: A**

**Solution:**



(a) : Enzymes control all the life processes. They increase the rate of a biological reaction. The magnitude of increase may be greater than those affected by other catalysts. Vitamins are accessory indispensable food factor, organic in nature (organic acid, amino acid esters, alcohols, steroids etc.) required by an organism in small amounts to maintain normal growth and regulate the metabolic processes. Hormones are biologically active organic substance that are produced in minute quantities by some specialized organs and exert physiological effects at sites remote from their origin.

---

## Question74

**Which of the following is the simplest amino acid?  
(2005)**

**Options:**


- A. Alanine
- B. Asparagine
- C. Glycine
- D. Tyrosine

**Answer: C**

**Solution:**

**Solution:**

(c) : Glycine is considered as the simplest amino acid as it has one amino group, one carboxylic group and no substituent functional group.



---

## Question75

**Nucleotides are building blocks of nucleic acids. Each nucleotide is a composite molecule formed by  
(2005)**

**Options:**

- A. base-sugar-phosphate
- B. base-sugar-OH
- C. (base-sugar-phosphate)  $n_n$
- D. sugar-phosphate.

**Answer: A**

**Solution:**



(a) : Each nucleotide consists of three distinct units - a phosphate group derived from phosphoric acid, a pentose sugar and a ring shaped nitrogenous base.

Nucleoside + Phosphoric acid + Nucleotide + H<sub>2</sub>O

---

## Question76

**Carbohydrates, the most abundant biomolecule on earth, are produced by (2005)**

**Options:**

- A. some bacteria, algae and green plant cells
- B. fungi, algae and green plant cells
- C. all bacteria, fungi and algae
- D. viruses, fungi and bacteria.

**Answer: A**

**Solution:**

**Solution:**

(a) : Carbohydrates are organic compounds synthesized in the chlorophyll containing cells of some bacteria, algae and green plant cells, during photosynthesis. Certain photo-autotrophic bacteria e.g., Green sulphur bacteria and purple sulphur bacteria contain pigments like chlorobium chlorophyll and bacteriochlorophyll respectively that helps them in photosynthesis.

---

## Question77

**In which one of the following enzymes, is copper necessarily associated as an activator? (2004)**

**Options:**

- A. Carbonic anhydrase
- B. Tryptophanase
- C. Lactic dehydrogenase
- D. Tyrosinase

**Answer: D**

**Solution:**

(d) : Copper is associated as an activator with tyrosinase. It is widely distributed in plants, animals and man. It is also known as polyphenol oxidase or catecholase. It oxidizes tyrosine to melanin in mammals and causes the cut surfaces of many fruits and vegetable to darken.

---



## Question78

**The major role of minor elements inside living organisms is to act as (2003)**

**Options:**

- A. co-factors of enzymes
- B. building blocks of important amino acids
- C. constituent of hormones
- D. binder of cell structure.

**Answer: A**

**Solution:**

**Solution:**

(a) : Minor element are those which are required in quantity of less than milligram/gram of dry matter but they are essential for proper growth and development of an organism e . g., Cl , M n, B, Z n, Cu Mo etc. These elements work as non-protein cofactor in enzymes e . g., Z n, Cu etc. They also take part in oxidation reduction reactions e . g., Cu, with variable valency. Chloride ion enhances activity of salivary amylase. Zinc is required for activity of carbonic anhydrase and alcohol dehydrogenase, etc.

---

## Question79

**Lipids are insoluble in water because lipid molecules are (2002)**

**Options:**

- A. hydrophilic
- B. hydrophobic
- C. neutral
- D. zwitter ions.

**Answer: B**

**Solution:**

**Solution:**

(b) : Lipid molecules are insoluble or sparingly soluble in water but are ~~freely soluble in organic solvents like ether,~~ alcohol and benzene. Insolubility of lipids in water is due to the fact that the polar groups they contain are much smaller than their nonpolar portions. The nonpolar chains are long complex hydrophobic hydrocarbon chains. If shaken in water lipids often form small droplets or micelles. The complex formed is called emulsions. These non polar proteins give them water repellent or hydrophobic property.

---



## Question80

**Which of the following is a reducing sugar?  
(2002)**

**Options:**

- A. Galactose
- B. Gluconic acid
- C.  $\beta$  -methyl galactoside
- D. Sucrose

**Answer: A**

**Solution:**

**Solution:**

(a) : All those sugars which have free aldehyde or ketone group are called reducing sugars. These are able to reduce cupric ions ( $\text{Cu}^{+2}$ ) into cuprous ions ( $\text{Cu}^{+}$ ). Sucrose, starch are non-reducing sugars.

---

## Question81

**Enzyme first used for nitrogen fixation  
(2001)**

**Options:**

- A. nitrogenase
- B. nitroreductase
- C. transferase
- D. transaminase.

**Answer: A**

**Solution:**

**Solution:**

(a) : Nitrogen fixation involves conversion of atmospheric nitrogen to ammonia. It is done with the help of nitrogenase enzyme which occurs inside thick walled heterocysts of the blue green algae. These provide suitable anaerobic environment for nitrogenase activity even in aerobic conditions.

---

## Question82

**Role of enzyme in reactions is to/as  
(2000)**

**Options:**




- A. decrease activation energy
- B. increase activation energy
- C. inorganic catalyst
- D. none of the above.

**Answer: A**

**Solution:**

(a) : All molecules require certain amount of energy for activation (to overcome energy barrier) before they can react. This energy is called activation energy. This energy is recovered when products are formed. The essence of an enzyme is its ability to speed up (catalyze) a reaction by making or breaking specific covalent bonds (bonds in which atoms are held together by sharing of electrons). Enzymes act by somehow lowering the temperature at which a given bond is unstable i.e., they speed up a reaction by lowering the activation energy. It is the magnitude of the activation energy which determines how fast the reaction will proceed.



**Graph showing energy requirement of catalysed and uncatalysed reactions.**

## Question83

**Which factor is responsible for inhibition of enzymatic process during feedback? (2000)**

**Options:**


- A. Substrate
- B. Enzymes
- C. End product
- D. Temperature

**Answer: C**

**Solution:**

**Solution:**

(c) : Feedback inhibition or end product inhibition is the inhibition of the activity of an enzyme catalysing some early reactions of the series by the end product of the metabolic pathway. For example a substrate A is converted into a product F through B, C, D and E intermediate products. As the concentration of end product F increases, it diffuses to allosteric enzyme (E) causing a reduced synthesis of the product B which in turn lowers the rate of enzymatic reactions in rest of the pathway.



**Feedback inhibition :** Product F inhibits the action of enzyme E<sub>1</sub>.

## Question84

**Enzymes are not found in (2000)**

**Options:**

- A. fungi
- B. algae
- C. virus
- D. cyanobacteria.

**Answer: C**

**Solution:**

**Solution:**

(c) : Viruses do not have enzymes so they cannot synthesize proteins. They multiply only inside the living host cell and for multiplication and metabolism they take over the machinery of the host cell. They lack their own cellular machinery and enzymes.

## Question85

**ATP is (2000)**

**Options:**

- A. nucleotide
- B. nucleoside
- C. nucleic acid
- D. vitamin.

**Answer: A**

**Solution:**

(a) : ATP is a nucleotide as it is composed of adenine, ribose sugar and phosphoric acid. There are two additional phosphate groups attached to the phosphate group of AMP. The last two phosphate molecules are connected by high energy bonds.

---

## Question86

**Which of the following have carbohydrate as prosthetic group?  
(2000)**

**Options:**

- A. Glycoprotein
- B. Chromoprotein
- C. Lipoprotein
- D. Nucleoprotein

**Answer: A**

**Solution:**

**Solution:**

(a) : Glycoproteins are proteins that contain sugars like carbohydrates as prosthetic group. In most glycoproteins, the linkage is between asparagine and N-acetyl-D-glucosamine. Some glycoproteins are immunoglobulins, membrane proteins and muscle proteins. Lipoproteins are protein complexed with lipids like triglycerides, phospholipids etc. Nucleoproteins are proteins associated with nucleic acids and chromoproteins are proteins associated with pigments e.g., cytochrome, phytochrome.

---

## Question87

**Cellulose, the most important constituent of plant cell wall is made up of  
(1998)**

**Options:**

- A. branched chain of glucose molecules linked by  $\beta - 1, 4$  glycosidic bond in straight chain and  $\alpha - 1, 6$  glycosidic bond at the site of branching
- B. unbranched chain of glucose molecules linked by  $\alpha - 1, 4$  glycosidic bond
- C. branched chain of glucose molecules linked by  $\alpha - 1, 6$  glycosidic bond at the site of branching
- D. unbranched chain of glucose molecules linked by  $\beta - 1, 4$  glycosidic bond.

**Answer: D**

**Solution:**

Cellulose ( $C_6H_{10}O_5)_n$  is the most abundant organic polymer. It is a polysaccharide and consists of long unbranched



chains of glucose residues linked by ( $\beta$ , 1 – 4 glycosidic bonds. In plants, cellulose is formed from sugar. It serves as building material in the formation of cell wall.

---

## Question88

**Lactose is composed of (1998)**

**Options:**

- A. glucose + galactose
- B. fructose + galactose
- C. glucose + fructose
- D. glucose + glucose.

**Answer: A**

**Solution:**

(a) : Lactose is popularly known as milk sugar. It is a disaccharide composed of one molecule of glucose and one molecule of galactose. The covalent bond that joins these two monosaccharide units is called glycosidic bond or glycosidic linkage. It is a reducing sugar.

---

## Question89

**Co-factor (prosthetic group) is a part of holoenzyme. It is (1997)**

**Options:**

- A. loosely attached organic part
- B. loosely attached inorganic part
- C. accessory non-protein substance attached firmly
- D. none of these.

**Answer: C**

---

## Question90

**Which is a typical example of 'feedback inhibition'? (1996)**




**Options:**

- A. Cyanide and cytochrome reaction
- B. Sulpha drugs and folic acid synthesizer bacteria
- C. Allosteric inhibition of hexokinase by glucose 6 -phosphate
- D. Reaction between succinic dehydro-genase and succinic acid

**Answer: C****Solution:**

(c) : Feedback inhibition or end product inhibition occurs when the end product of a metabolic pathway inhibits the activity of an enzyme catalysing some early reactions of the series. The end product is the inhibitor and the enzyme inactivated is called allosteric enzyme. The enzyme is regulated by modulators that bind noncovalently at site other than the active site. An example of feedback inhibition is the inhibition of the activity of the enzyme hexokinase by glucose 6 -phosphate in glycolysis. This enzyme catalyses conversion of glucose into glucose 6 -phosphate but as the reaction proceeds, increase in concentration of glucose 6-phosphate inhibits the activity of hexokinase.



---

## Question91

**In which of the following groups are all polysaccharides?  
(1996)**

**Options:**

- A. Sucrose, glucose and fructose
- B. Maltose, lactose and fructose
- C. Glycogen, sucrose and maltose
- D. Glycogen, cellulose and starch

**Answer: D****Solution:**

(d) : Polysaccharides are complex long chain carbohydrates which are formed by dehydrate synthesis or polymerisation of more than 10 but generally very large number of units called monosaccharides. Starch, glycogen and cellulose are all polysaccharides. Starch is a glucosan homopolysaccharide which is the main reserve food of plants. Glycogen is also a glucosan homopoly-saccharide which is the major reserve food of fungi, animals and some bacteria. It is also called animal starch. Cellulose is the structural polysaccharide of plant cell walls, some fungi, protists. It is a fibrous glucosan homopolysaccharide of high tensile strength.



## Question92

**What are the most diverse molecules in the cell?  
(1996)**

**Options:**

- A. Lipids
- B. Mineral salts
- C. Proteins
- D. Carbohydrates

**Answer: C**

**Solution:**

(c) : Proteins show enormous diversity because of different proportions and sequences of twenty amino acid within the protein molecule. A large number of permutations and combinations of these amino acids are responsible for the unlimited variety of proteins. Proteins are the most abundant and most varied of the macromolecules having one or more polypeptides (chains of amino acids). The proteins constitute almost 50 % of the total dry weight of the cell. Proteins may be simple or conjugated. Among conjugated, proteins may be phosphoprotein, glycoprotein, nucleoprotein, chromoprotein, lipoprotein, flavoprotein, metallo protein etc. Functionally proteins may be structural protein, enzymes, hormones, respiratory pigment etc.

---

## Question93

**Which purine base is found in RNA?  
(1996)**

**Options:**

- A. Thymine
- B. Uracil
- C. Cytosine
- D. Guanine

**Answer: D**

**Solution:**

**Solution:**

(d) : The bases are of two types-purines and pyrimidines. The purine derivatives adenine (A) and guanine (G) are double ring structures whereas pyrimidine derivatives thymine, cytosine and uracil are single ring structures. Thymine (T) and cytosine (C) are found in DNA and cytosine (C) and uracil (U) is found in RNA.

---

## Question94



**Which of the following nucleotide sequences contains 4 pyrimidine bases?  
(1994)**

**Options:**

- A. GATCAATGC
- B. GCUAGACAA
- C. UAGCGGUAA
- D. Both (b) and (c)

**Answer: A**

**Solution:**

**Solution:**

(a) : In the given question there are 4 pyrimidines as 2 cytosine and 2 thymine in option 'a'

---

## Question95

**The four elements that make up 99% of all elements found in a living system are  
(1994)**

**Options:**

- A. C, H, O and P
- B. C, N, O and P
- C. H, O, C and N
- D. C, H, O and S

**Answer: C**

**Solution:**

**Solution:**

(c) : Carbon, hydrogen, oxygen and nitrogen are called four big elements of living body they make up about 99% of the mass of most cells. As C, H, O and N are lightest elements so the bonds they form are the strongest covalent bonds. So that the compounds formed are stable, varied in size and shapes. Carbon constitutes more than 50% of the dry matter. It has been observed that human body contains 0.5% hydrogen, 18.5% carbon, 65% oxygen and 3.3% nitrogen. Other elements are present in very lesser amount.

---

## Question96

**Which is wrong about nucleic acids?  
(1993)**



**Options:**

- A. DNA is single stranded in some viruses.
- B. RNA is double stranded occasionally.
- C. Length of one helix is  $45\text{\AA}$  in B-DNA.
- D. One turn of Z-DNA has 12 bases.

**Answer: C**

**Solution:**

**Solution:**

(c) : One complete turn of DNA is  $34\text{\AA}$  long and has 10 base pairs.

---

## Question97

### Glycogen is a polymer of (1993)

**Options:**

- A. galactose
- B. glucose
- C. fructose
- D. sucrose.

**Answer: B**

**Solution:**

**Solution:**

(b) : Glycogen (animal starch) is a polysaccharide consisting of a highly branched polymer of glucose occurring in animal tissues, especially in liver and muscle cells. It is the major store of carbohydrate energy in animal cells.

---

## Question98

### In RNA, thymine is replaced by (1992)

**Options:**

- A. adenine
- B. guanine
- C. cytosine
- D. uracil.



**Answer: D**

**Solution:**

**Solution:**

(d) : The bases are of two types-purines and pyrimidines. The purine derivatives adenine (A) and guanine (G) are double ring structures whereas pyrimidine derivatives thymine, cytosine and uracil are single ring structures. Thymine (T) and cytosine (C) are found in DNA and cytosine (C) and uracil (U) is found in RNA.

---

## Question99

**Amino acids are mostly synthesised from (1992)**

**Options:**

- A. mineral salts
- B. fatty acids
- C. volatile acids
- D.  $\alpha$  -ketoglutaric acid.

**Answer: D**

**Solution:**

**Solution:**

(d) : Amino acids are mostly synthesized from  $\alpha$  -ketoglutaric acid. These are the precursors of amino acids. A five carbon compound formed during Krebs' cycle is a  $\alpha$  -ketoglutaric acid which is the first dicarboxylic acid formed. Pyruvic acid converted into alanine,  $\alpha$  -ketoglutaric acid into glutamic acid, OAA into aspartic acid, polymerization of such amino acids results into formation of proteins.

---

## Question100

**Which is distributed more widely in a cell? (1992)**

**Options:**

- A. DNA
- B. RNA
- C. Chloroplasts
- D. Sphaerosomes

**Answer: B**

**Solution:**

(b) : RNA occurs in the nucleus as well as in the cytoplasm of the eukaryotic cells and in prokaryotic cell, it is found in the cytoplasm. DNA is found in the nucleus, mitochondria and chloroplast. Chloroplast and sphaerosomes are found only in cytoplasm.

---

## Question101

**Living cell contains 60-95% water. Water present in human body is (1992)**

**Options:**

- A. 60-65 %
- B. 50-55 %
- C. 75-80 %
- D. 65-70 %

**Answer: D**

**Solution:**

**Solution:**

(d) : Water is the most abundant substance of living beings. The water content of actively living cells varies between 60-95 %. In human beings maximum water content is found in the embryo 90-95 %. Water content decreases thereafter in adult and the aged where it is 65 -70 %

---

## Question102

**Adenine is (1992)**

**Options:**

- A. purine
- B. pyrimidine
- C. nucleoside
- D. nucleotide

**Answer: A**

**Solution:**

**Solution:**

(a) : The bases are of two types-purines and pyrimidines. The purine derivatives adenine (A) and guanine (G) are double ring structures whereas pyrimidine derivatives thymine, cytosine and uracil are single ring structures. Thymine (T) and cytosine (C) are found in DNA and cytosine (C) and uracil (U) is found in RNA.

---



## Question103

**Enzymes having slightly different molecular structure but performing identical activity are (1991)**

**Options:**

- A. holoenzymes
- B. isoenzymes
- C. apoenzymes
- D. coenzymes

**Answer: B**

**Solution:**

**Solution:**

(b) There are certain enzymes which have slightly different molecular structure but have similar catalytic function such enzymes are known as isoenzymes. Holoenzyme is the active compound formed by combination of a coenzyme and an apoenzyme. Apoenzyme is the protein component of an enzyme, to which the coenzyme attaches to form an active enzyme. Coenzymes are organic non-protein molecules that bind with the protein molecule (apoenzyme) to form the active enzyme (holoenzyme).

---

## Question104

**A nucleotide is formed of (1991)**

**Options:**

- A. purine, pyrimidine and phosphate
- B. purine, sugar and phosphate
- C. nitrogen base, sugar and phosphate
- D. pyrimidine, sugar and phosphate

**Answer: C**

**Solution:**

**Solution:**

(c) : Nucleotide is an organic compound consisting of a nitrogen-containing purine or pyrimidine base linked to a sugar (ribose or deoxyribose) and a phosphate group.

---

## Question105

**DNA is composed of repeating units of**

**(1991)**

**Options:**

- A. ribonucleosides
- B. deoxyribonucleosides
- C. ribonucleotides
- D. deoxyribonucleotides

**Answer: D**

**Solution:**

**Solution:**

(d) : DNA is the largest macromolecule in the organisms. It is a long, double chain of deoxyribonucleotide, or deoxyribotide units. The two deoxyribonucleotide chains are twisted around a common axis to form a right-handed double helix (spiral) that encloses a cylindrical space in it. Each deoxyribonucleotide unit, in turn, consists of three different molecules : phosphate, ( $\text{PO}_4^{3-}$ ), a 5 -carbon deoxyribose sugar ( $\text{C}_5\text{H}_{10}\text{O}_4$ ) and a nitrogenous base.

---

## Question106

**A segment of DNA has 120 adenine and 120 cytosine bases. The total number of nucleotides present in the segment is (1991)**

**Options:**

- A. 120
- B. 240
- C. 60
- D. 480

**Answer: D**

**Solution:**

**Solution:**

(d) : According to Chargaff's rules, the amount of adenine is always equal to that of thymine, and the amount of guanine is always equal to that of cytosine i.e.,  $A = T$  (120) and  $G = C$  (120), therefore, the total no. of nucleotides would be  $120 \times 4 = 480$ .

---

## Question107

**The basic unit of nucleic acid is (1991)**

**Options:**





- A. pentose sugar
- B. nucleoid
- C. nucleoside
- D. nucleotide.

**Answer: D**

**Solution:**

(d) : The nucleic acids (DNA and RNA) are the molecules having complex structure and very high molecular weights. The nucleic acid is composed of a large number of nucleotide molecules joined into a linear, unbranched chain. Nucleotide is an organic compound consisting of a nitrogen-containing purine or pyrimidine base linked to a sugar (ribose or deoxyribose) and a phosphate group.

---

## Question108

**Mineral associated with cytochrome is (1991)**

**Options:**

- A. Cu
- B. Mg
- C. Cu and Mg
- D. Fe

**Answer: D**

**Solution:**

**Solution:**

(d) : Cytochromes are generally membranebound hemoproteins that contain heme groups and carry out electron transport. The heme group is a highly conjugated ring system (which means its electrons are very mobile) surrounding a metal ion, which readily interconverts between the oxidation states. For many cytochromes the metal ion present is that of iron, which interconverts between  $Fe^{2+}$  (reduced) and  $Fe^{3+}$  (oxidized) states (electron-transfer processes) or between  $Fe^{2+}$  (reduced) and  $Fe^{3+}$  (formal, oxidized) states (oxidative processes). Cytochromes are thus capable of performing oxidation and reduction.

---

## Question109

**Which is not consistent with double helical structure of DNA? (1990)**

**Options:**

- A.  $A = T, C = G$
- B. Density of DNA decreases on heating.



C.  $A + T/C + G$  is not constant.

D. Both (a) and (b)

**Answer: C**

**Solution:**

(c) : The density of DNA decreases on heating as hydrogen bonds breakdown. According to Chargaff's rules, the amount of adenine is always equal to that of thymine, and the amount of guanine is always equal to that of cytosine i.e.,  $A = T$  and  $G = C$ . The base ratio  $A + T/G + C$  may vary from species to species, but is constant for a given species.

---

## Question110

**RNA does not possess (1988)**

**Options:**

A. uracil

B. thymine

C. adenine

D. cytosine.

**Answer: B**

**Solution:**

**Solution:**

(b) : The bases are of two types-purines and pyrimidines. The purine derivatives adenine (A) and guanine (G) are double ring structures whereas pyrimidine derivatives thymine, cytosine and uracil are single ring structures. Thymine (T) and cytosine (C) are found in DNA and cytosine (C) and uracil (U) is found in RNA.

---

## Question111

**In double helix of DNA, the two DNA strands are (1988)**

**Options:**

A. coiled around a common axis

B. coiled around each other

C. coiled differently

D. coiled over protein sheath.

**Answer: A**

## **Solution:**

(a) : According to Watson-Crick model, the DNA molecule consists of two long, parallel chains which are joined together by short crossbars at regular intervals. The two chains are spirally coiled around a common axis in a regular manner to form a right handed double helix.

---

# Chemistry In Everyday Life

## Question1

Some tranquilizers are listed below. Which one from the following belongs to barbiturates?

[NEET 2023]

**Options:**

A.

Meprobamate

B.

Valium

C.

Veronal

D.

Chlordiazepoxide

**Answer: C**

**Solution:**

**Solution:**

Veronal is the derivative of Barbituric acid and considered as barbiturate.

Meprobamate, valium and chlordiazepoxide are other tranquilizers.

-----

## Question2

Cheilosis occurs due to deficiency of \_\_\_\_\_.

[NEET 2023 mpr]

**Options:**

A.

thiamine

B.

nicotinamide

C.



pyridoxamine

D.

riboflavin

**Answer: D**

**Solution:**

Cheilosis (Fissuring at corners of mouth and lips) occurs due to deficiency of vitamin B<sub>2</sub> (Riboflavin)

---

### Question3

**Which amongst the following is used in controlling depression and hypertension?**

**[NEET 2023 mpr]**

**Options:**

A.

Seldane

B.

Valium

C.

Equanil

D.

Prontosil

**Answer: C**

**Solution:**

Equanil is used in controlling depression and hyper tension.

---

### Question4

**Match List-I with List-II.**

List - I	List - II
(a) Sodium laurylsulphate	(i) Toilet soap
(b) Cetyltrimethyl ammonium chloride	(ii) Non-ionic detergent
(c) Sodium stearate	(iii) Anionic detergent
(d) Polyethyleneglycyl stearate	(iv) Cationic detergent

**Choose the correct answer from the options given below:  
[NEET-2022]**

**Options:**

- A. (a) - (iii), (b) - (ii), (c) - (iv), (d) - (i)  
 B. (a) - (iii), (b) - (iv), (c) - (ii), (d) - (i)  
 C. (a) - (i), (b) - (iv), (c) - (ii), (d) - (iii)  
 D. (a) - (iv), (b) - (iii), (c) - (i), (d) - (ii)

**Answer: B**

**Solution:**

**Solution:**

- Cimetidine is an antacid
- Seldane is an antihistamine
- Morphine is an analgesic
- Salvarsan is an antimicrobial drug

## Question5

**The incorrect statement regarding enzymes is  
[NEET-2022]**

**Options:**

- A. Enzymes are biocatalysts.  
 B. Like chemical catalysts enzymes reduce the activation energy of bio processes.  
 C. Enzymes are polysaccharides.  
 D. Enzymes are very specific for a particular reaction and substrate.

**Answer: C**

**Solution:**

**Solution:**

Enzymes are complex nitrogenous organic compounds which are produced by living plants and animals. They are



protein molecules of high molecular mass. They are not polysaccharides.

---

## Question6

**Match List - I with List - II :**

List - I	List - II
(a) Sodium laurylsulphate	(i) Toilet soap
(b) Cetyltrimethyl ammonium chloride	(ii) Non-ionic detergent
(c) Sodium stearate	(iii) Anionic detergent
(d) Polyethyleneglycyl stearate	(iv) Cationic detergent

**[NEET Re-2022]**

**Options:**

- A. (a) – (iii), (b) – (i), (c) – (ii), (d) – (iv)
- B. (a) - (iv), (b) - (iii), (c) - (i), (d) - (ii)
- C. (a) - (i), (b) - (iv), (c) - (ii), (d) - (iii)
- D. (a) - (iii), (b) - (iv), (c) - (i), (d) - (ii)

**Answer: D**

**Solution:**

**Solution:**

- (a) Sodium laurylsulphate → Anionic detergent
  - (b) Cetyltrimethylammonium chloride → cationic detergent
  - (c) Sodium stearate → Toilet soap
  - (d) Polyethyleneglycyl stearate → Non-ionic detergent
- 

## Question7

**Given below are two statements :**

**Statement I :**

**Aspirin and Paracetamol belong to the class of narcotic analgesics.**

**Statement II :**

**Morphine and Heroin are non-narcotic analgesics. In the light of the above statements, choose the correct answer from the options given below.**

**[NEET 2021]**

**Options:**

©

- A. Both Statement I and Statement II are true
- B. Both Statement I and Statement II are false
- C. Statement I is correct but Statement II is false
- D. Statement I is incorrect but Statement II is true.

**Answer: B**

**Solution:**

**Solution:**

- Aspirin and paracetamol belong to the class of non-narcotic analgesics
  - Morphine and Heroin are Narcotic analgesics
- ∴ Both statement I and statement II are false

---

## Question8

**The RBC deficiency is deficiency disease of :  
[NEET 2021]**

**Options:**

- A. Vitamin B<sub>12</sub>
- B. Vitamin B<sub>6</sub>
- C. Vitamin B<sub>1</sub>
- D. Vitamin B<sub>2</sub>

**Answer: A**

**Solution:**

**Solution:**

- Deficiency of vitamin B<sub>2</sub> (Riboflavin) causes cheilosis, digestive disorders and burning sensation of the skin.
- Deficiency of vitamin B<sub>12</sub> causes Pernicious anaemia which is RBC deficiency in haemoglobin.
- Deficiency of vitamin B<sub>6</sub> (Pyridoxine) causes Convulsions.
- Deficiency of vitamin B<sub>1</sub> (Thiamine) causes Beri-Beri (loss of appetite and retarded growth).

---

## Question9

**Which of the following is a cationic detergent?  
[2020]**

**Options:**

- A. Sodium stearate

©






- B. Cetyltrimethyl ammonium bromide
- C. Sodium dodecylbenzene sulphonate
- D. Sodium lauryl sulphate

**Answer: B**

**Solution:**



## Question10

**Among the following, the narrow spectrum antibiotic is (NEET 2019)**

**Options:**

- A. chloramphenicol
- B. penicillin G
- C. ampicillin
- D. amoxycillin.

**Answer: B**

---

## Question11

**The artificial sweetner stable at cooking temperature and does not provide calories is (Odisha NEET 2019)**

**Options:**

- A. saccharin
- B. aspartame
- C. sucralose

©



D. alitame.

**Answer: C**

---

## Question12

**The liquified gas that is used in dry cleaning along with a suitable detergent is (Odisha NEET 2019)**

**Options:**

- A. water gas
- B. petroleum gas
- C. NO<sub>2</sub>
- D. CO<sub>2</sub>

**Answer: D**

---

## Question13

**Mixture of chloroxylenol and terpineol acts as (NEET 2017)**

**Options:**

- A. antiseptic
- B. antipyretic
- C. antibiotic
- D. analgesic.

**Answer: A**

**Solution:**

Dettol which is a well known antiseptic is a mixture of chloroxylenol and  $\alpha$ -terpineol in a suitable solvent.

---



## Question14

**Which of the following is an analgesic?  
(NEET-I 2016)**

**Options:**

- A. Streptomycin
- B. Chloromycetin
- C. Novalgin
- D. Penicillin

**Answer: C**

**Solution:**

**Solution:**

Streptomycin, chloromycetin and penicillin are antibiotics while novalgin is an analgesic.

---

## Question15

**Bithional is generally added to the soaps as an additive to function as  
a/an  
(2015 Cancelled)**

**Options:**

- A. buffering agent
- B. antiseptic
- C. softener
- D. dryer.

**Answer: B**



## Question16

**Artificial sweetner which is stable under cold conditions only is (2014)**

**Options:**

- A. saccharine
- B. sucralose
- C. aspartame
- D. alitame

**Answer: C**

**Solution:**

**Solution:**

Aspartame is stable under cold conditions and unstable at cooking temperature.

---

## Question17

**Antiseptics and disinfectants either kill or prevent growth of microorganisms. Identify which of the following statements is not true. (2013 NEET)**

**Options:**

- A. Dilute solutions of boric acid and hydrogen peroxide are strong antiseptics.
- B. Disinfectants harm the living tissues
- C. A 0.2% solution of phenol is an antiseptic while 1% solution acts as a disinfectant.
- D. Chlorine and iodine are used as strong disinfectants.

**Answer: A**

**Solution:**

**Solution:**

Dilute solutions of boric acid and hydrogen peroxide are weak antiseptics.

---



## Question18

**Dettol is the mixture of  
(Karnataka NEET 2013)**

**Options:**

- A. chloroxylenol and bithionol
- B. chloroxylenol and terpineol
- C. phenol and iodine
- D. terpineol and bithionol.

**Answer: B**

**Solution:**

**Solution:**

Dettol is the mixture of chloroxylenol and  $\alpha$  -terpineol.

---

## Question19

**Chloroamphenicol is an  
(2012 Mains)**

**Options:**

- A. antifertility drug
- B. antihistaminic
- C. antiseptic and disinfectant
- D. antibiotic-broad spectrum

**Answer: D**

---

## Question20

**Which one of the following is employed as Antihistamine?  
(2011)**



**Options:**

- A. Chloramphenicol
- B. Diphenyl hydramine
- C. Norethindrone
- D. Omeprazole

**Answer: B**

**Solution:**

**Solution:**

Diphenylhydramine is employed as antihistamine drug.

---

## Question21

**Which one of the following is employed as a tranquilizer drug? (2010)**

**Options:**

- A. Promethazine
- B. Valium
- C. Naproxen
- D. Mifepristone

**Answer: B**

**Solution:**

**Solution:**

Valium is a tranquilizer.

---

## Question22

**Which one of the following is employed as a tranquilizer? (2009)**

**Options:**

- A. Naproxen

- B. Tetracycline
- C. Chlorpheniramine
- D. Equanil.

**Answer: D**

**Solution:**

**Solution:**

Equanil is used for the treatment of stress, mild and severe mental diseases i.e., as a tranquilizer.

---

## Question23

**Chloropicrin is obtained by the reaction of (2004)**


**Options:**

- A. steam on carbon tetrachloride
- B. nitric acid on chlorobenzene
- C. chlorine on picric acid
- D. nitric acid on chloroform.

**Answer: D**

**Solution:**

When chloroform is treated with concentrated nitric acid, its hydrogen is replaced by nitro group.



## Question24

**Which of the following forms cationic micelles above certain concentration? (2004)**

**Options:**

- A. Sodium dodecyl sulphate
- B. Sodium acetate
- C. Urea



D. Cetyltrimethylammonium bromide

**Answer: B**

**Solution:**

Sodium acetate forms cationic micelles. In the molecule of soaps and detergents the negative ions aggregate to form a micelle of colloidal size. The negative ion has a long hydrocarbon chain and a polar group ( $\text{COO}^-$ ) at one end.

---

## Question25

**Gammexane is (1999)**


**Options:**

- A. bromobenzene
- B. benzyl chloride
- C. chlorobenzene
- D. benzenehexachloride.

**Answer: D**

**Solution:**

Gammexane is an isomeric form of benzene hexachloride (BHC).



## Question26

**The decomposition of organic compounds, in the presence of oxygen and without the development of odoriferous substances, is called (1999)**

**Options:**

- A. nitrification

©



B.  $N_2$  -fixation

C. decay

D. denitrification.

**Answer: C**

**Solution:**

**Solution:**

The other three type of reactions takes place only in the presence of bacteria, in which the nitrogen compound is decomposed either to nitrogen or nitrates or nitrites. While in decay reaction an organic compound is decomposed in presence of oxygen.

## Question27

**Aspirin is an acetylation product of (1998)**

**Options:**

A. m -Hydroxybenzoic acid

B. o -Dihydroxybenzene


C. o -Hydroxybenzoic acid

D. p -Dihydroxybenzene.

**Answer: C**

**Solution:**

Aspirin is acetyl salicylic acid, which is formed by acetylation of o - hydroxybenzoic acid.



## Question28

**Which of the following can possibly be used as analgesic without causing addiction and mood modification? (1997)**

**Options:**

©



- A. Diazepam
- B. Tetrahydrocannabinol
- C. Morphine
- D. N -Acetyl-para-aminophenol.

**Answer: D**

**Solution:**

**Solution:**

N-Acetyl-para-aminophenol (or paracetamol) is an antipyretic which can also be used as an analgesic to relieve pains.

---

## Question29

**Diazo coupling is useful to prepare some (1994)**

**Options:**

- A. pesticides
- B. dyes
- C. proteins
- D. vitamins.

**Answer: B**

**Solution:**

**Solution:**

Azo dyes are derived by coupling of a phenol adsorbed on the surface of a fabric with a diazonium salt. Dyes can be prepared by diazo coupling. For example,

---

## Question30

**Which one of the following statements is not true? (1994)**

**Options:**

- A. Ampicillin is a natural antibiotic.
- B. Aspirin is both analgesic and antipyretic.
- C. Sulphadiazine is a synthetic antibacterial drug.



D. Some disinfectants can be used as antiseptics.

**Answer: A**

**Solution:**

Ampicillin is a modification of penicillin and thus is not a natural antibiotic.

---