

Some Basic Concepts of Chemistry

Question1


Mass of methane required to produce 22g of CO₂ after complete combustion is g.

(Given Molar mass in g mol⁻¹ C = 12.0 H = 1.0 O = 16.0)

[27-Jan-2024 Shift 1]

Answer: 8

Solution:



$$\text{Moles of CO}_2 = \frac{22}{44} = 0.5$$

So, required moles of CH₄ = 0.5

$$\text{Mass} = 0.5 \times 16 = 8 \text{ gm}$$

Question2



9.3g of aniline is subjected to reaction with excess of acetic anhydride to prepare acetanilide. The mass of acetanilide produced if the reaction is 100% completed is ___ × 10⁻¹g. (Given molar mass in gmol⁻¹ N : 14, O : 16, C : 12, H : 1)

[27-Jan-2024 Shift 2]

Answer: 13.5

Solution:





$$n_{\text{Acetanilide}} = n_{\text{Aniline}}$$

$$\Rightarrow \frac{m}{135} = \frac{9.3}{93}$$

$$\Rightarrow m = 13.5\text{g}$$


Question3

1 mole of PbS is oxidised by "X" moles of O₃ to get "Y" moles of O₂. X + Y =

[27-Jan-2024 Shift 2]

Answer: 8

Solution:



$$x = 4, y = 4$$

Question4

0.05cm thick coating of silver is deposited on a plate of 0.05m² area. The number of silver atoms deposited on plate are _____ × 10²³. (At mass Ag = 108, d = 7.9gcm⁻³)

[30-Jan-2024 Shift 1]

Answer: 11

Solution:

$$\text{Volume of silver coating} = 0.05 \times 0.05 \times 10000$$

$$= 25\text{cm}^3$$

$$\text{Mass of silver deposited} = 25 \times 7.9\text{g}$$

$$\text{Moles of silver atoms} = \frac{25 \times 7.9}{108}$$

$$\text{Number of silver atoms} = \frac{25 \times 7.9}{108} \times 6.023 \times 10^{23}$$

$$= 11.01 \times 10^{23}$$

Ans. 11


Question5

Number of moles of methane required to produce 22gCO₂(g) after combustion is $x \times 10^{-2}$ moles. The value of x is

[31-Jan-2024 Shift 1]

Answer: 50

Solution:



$$n_{\text{CO}_2} = \frac{22}{44} = 0.5 \text{ moles}$$

So moles of CH₄ required = 0.5 moles i.e. 50×10^{-2} mole

$$x = 50$$

Question6

A sample of CaCO₃ and MgCO₃ weighed 2.21g is ignited to constant weight of 1.152g. The composition of mixture is :

(Given molar mass in gmol⁻¹

CaCO₃ : 100, MgCO₃ : 84)


[31-Jan-2024 Shift 2]

Options:


A.

$$1.187\text{gCaCO}_3 + 1.023\text{gMgCO}_3$$


B.



C.



D.



Answer: A

Solution:



Let the weight of CaCO_3 be x gm

$$\therefore \text{weight of MgCO}_3 = (2.21 - x) \text{ gm}$$

Moles of CaCO_3 decomposed = moles of CaO formed

$$\frac{x}{100} = \text{moles of CaO formed}$$

$$\therefore \text{weight of CaO formed} = \frac{x}{100} \times 56$$

Moles of MgCO_3 decomposed = moles of MgO formed

$$\frac{(2.21 - x)}{84} = \text{moles of MgO formed}$$

$$\therefore \text{weight of MgO formed} = \frac{2.21 - x}{84} \times 40$$

$$\Rightarrow \frac{2.21 - x}{84} \times 40 + \frac{x}{100} \times 56 = 1.152$$

$$\therefore x = 1.1886\text{g} = \text{weight of CaCO}_3$$

$$\& \text{ weight of MgCO}_3 = 1.0214\text{g}$$

Question7

The molarity of 1L orthophosphoric acid (H_3PO_4) having 70% purity by weight (specific gravity 1.54gcm^{-3}) is _____ M.

(Molar mass of $\text{H}_3\text{PO}_4 = 98\text{gmol}^{-1}$)

[31-Jan-2024 Shift 2]

Answer: 11

Solution:

Specific gravity (density) = 1.54g/cc.

Volume = 1L = 1000ml

Mass of solution = 1.54 × 1000

= 1540g

% purity of H₂SO₄ is 70%


So weight of H₃PO₄ = 0.7 × 1540 = 1078g

Mole of H₃PO₄ = $\frac{1078}{98} = 11$

Molarity = $\frac{11}{1L} = 11$

Question8

Consider the following reaction:



If 72mmol² of PbCl₂ is mixed with 50mmol of (NH₄)₃PO₄, then amount of Pb₃(PO₄)₂ formed is _____mmol. (nearest integer)

[1-Feb-2024 Shift 1]

Answer: 24

Solution:

Limiting Reagent is PbCl₂ mmol of Pb₃(PO₄)₂ formed = $\frac{\text{mmol of PbCl}_2 \text{ reacted}}{3} = 24 \text{ mmol}$



Question9

10mL of gaseous hydrocarbon on combustion gives 40mL of CO₂(g) and 50mL of water vapour. Total number of carbon and hydrogen atoms in the hydrocarbon is_____

[1-Feb-2024 Shift 2]

Answer: 14

Solution:



$$10x = 40$$

$$x = 4$$

$$5y = 50$$

$$y = 10$$



Question 10

When $\text{Fe}_{0.93}\text{O}$ is heated in presence of oxygen, it converts to Fe_2O_3 . The number of correct statement/s from the following is ___

A. The equivalent weight of $\text{Fe}_{0.93}\text{O}$ is $\frac{\text{Molecular weight}}{0.79}$.

B. The number of moles of Fe^{2+} and Fe^{3+} in 1 mole of $\text{Fe}_{0.93}\text{O}$ is 0.79 and 0.14 respectively.


C. $\text{Fe}_{0.93}\text{O}$ is metal deficient with lattice comprising of cubic closed packed arrangement of O^{2-} ions.

D. The % composition of Fe^{2+} and Fe^{3+} in $\text{Fe}_{0.93}\text{O}$ is 85% and 15% respectively.

[24-Jan-2023 Shift 1]

Answer: 4

Solution:



$$nf = \left(3 - \frac{200}{93}\right) \times 0.93$$

$$nf = 0.79$$

$$\text{B : } 2x + (0.93 - x) \times 3 = 2$$

$$x = 0.79$$

$$\text{Fe}^{2+} = 0.79, \text{Fe}^{3+} = 0.21$$

$$\text{C : Fact}^D : \% \text{Fe}^{2+} = \frac{0.79}{0.93} \times 100 = 85\%; \text{Fe}^{3+} = 15\%$$

Question 11



The number of units, which are used to express concentration of solutions from the following is ____

Mass percent, Mole, Mole fraction, Molarity, ppm, Molality.

[24-Jan-2023 Shift 2]

Answer: 5

Solution:

Mass percent, mole fraction, molarity, ppm, molality are used for measuring concentration terms.

Question12

What is the mass ratio of ethylene glycol ($C_2H_6O_2$, molar mass = 62g / mol) required for making 500g of 0.25 molal aqueous solution and 250 mL of 0.25 molar aqueous solution ?

[25-Jan-2023 Shift 2]

Options:

A. 1 : 1

B. 3 : 1

C. 2 : 1

D. 1 : 2

Answer: C

Solution:

Assume : Mass of solvent \approx Mass of solution

Case I :-

$$0.25 = \frac{W_1}{62} \times \frac{1000}{500}$$

Case II :-

$$0.25 = \frac{W_2}{62} \times \frac{1000}{250}$$

$$\frac{W_1}{W_2} = \frac{2}{1}$$

Question13

Number of hydrogen atoms per molecule of a hydrocarbon A having 85.8% carbon is _____



(Given : Molar mass of A = 84gmol^{-1})
[25-Jan-2023 Shift 2]

Answer: 12

Solution:

Element	Percentage	Mole	Mole ratio
C	85.8	$\frac{85.8}{12} = 7.15$	1
H	14.2	$\frac{14.2}{1} = 14.2$	2

Empirical formula (CH_2)

$$14 \times n = 84$$

$$n = 6$$





\therefore Molecular formula C_6H_{12}

Question14

When a hydrocarbon A undergoes combustion in the presence of air, it requires 9.5 equivalents of oxygen and produces 3 equivalents of water. What is the molecular formula of A ?


[29-Jan-2023 Shift 2]

Options:



Answer: A

Solution:



$$x + \frac{y}{4} = 9.5$$

$$\frac{y}{2} = 3$$

$$\Rightarrow x = 8, y = 6$$



Question15

When 0.01 mol of an organic compound containing 60% carbon was burnt completely, 4.4g of CO_2 was produced. The molar mass of compound is _____ gmol^{-1} (Nearest integer)
[29-Jan-2023 Shift 2]

Answer: 200

Solution:

Let M is the molar mass of the compound (g / mol)

mass of compound = 0.01 Mgm

mass of carbon = $0.01M \times \frac{60}{100}$

moles of carbon = $\frac{0.01M}{12} \times \frac{60}{100}$

moles of CO_2 from combustion = $\frac{4.4}{44}$ = moles of carbon

$$\frac{0.01M}{12} \times \frac{60}{100} = \frac{4.4}{44}$$

$$M = \frac{4.4}{44} \times \frac{100}{60} \times \frac{12}{0.01} = 200 \text{ gm / mol}$$

Question16

Some amount of dichloromethane (CH_2Cl_2) is added to 671.141 mL of chloroform (CHCl_3) to prepare $2.6 \times 10^{-3}\text{M}$ solution of CH_2Cl_2 (DCM). The concentration of DCM is ppm _____ (by mass).
Given: Atomic mass: C = 12; H : 1; Cl = 35.5 density of $\text{CHCl}_3 = 1.49\text{gcm}^{-3}$
[30-Jan-2023 Shift 1]

Answer: 148

Solution:

$$\text{Molarity} = \frac{\text{mole}}{\text{volume}}$$

$$2.6 \times 10^{-3} = \frac{x / 85}{0.67141}$$

$$x = 0.148\text{g}$$



$$\text{conc. of DCM in ppm} = \frac{0.148}{1.49 \times 671.141} \times 10^6$$
$$= 148 \text{ ppm}$$

Question 17

Match List I with List II:

List I (Mixture)	List II (Separation Technique)
(A) $\text{CHCl}_3 + \text{C}_6\text{H}_5\text{NH}_2$	I. Steam distillation
(B) $\text{C}_6\text{H}_{14} + \text{C}_5\text{H}_{12}$	II. Differential extraction
(C) $\text{C}_6\text{H}_5\text{NH}_2 + \text{H}_2\text{O}$	III. Distillation
(D) Organic compound in H_2O	IV. Fractional distillation

[30-Jan-2023 Shift 2]

Options:

- A. A-IV, B-I, C-III, D-II
- B. A-III, B-IV, C-I, D-II
- C. A-II, B-I, C-III, D-IV
- D. A-III, B-I, C-IV, D-II

Answer: B

Solution:

Solution:



List I (Mixture)	List II (Separation Technique)
$\text{CHCl}_3 + \text{C}_6\text{H}_5\text{NH}_2$	Distillation
$\text{C}_6\text{H}_{14} + \text{C}_5\text{H}_{12}$	Fractional distillation
$\text{C}_6\text{H}_5\text{NH}_2 + \text{H}_2\text{O}$	Steam distillation
Organic compound in H_2O	Differential extraction

Question 18

On complete combustion, 0.492g of an organic compound gave 0.792g of CO_2 .

The % of carbon in the organic compound is _____
(Nearest integer)

[31-Jan-2023 Shift 1]

Answer: 44

Solution:

weight of C in 0.792 gm CO_2

$$= \frac{12}{44} \times 0.792 = 0.216$$

$$\% \text{ of C in compound} = \frac{0.216}{0.492} \times 100$$

$$= 43.90\%$$

Ans: 44

Question 19

Zinc reacts with hydrochloric acid to give hydrogen and zinc chloride.


The volume of hydrogen gas produced at STP from the reaction of 11.5g of zinc with excess HCl is _____ L (Nearest integer)

(Given : Molar mass of Zn is 65.4 gmol^{-1} and Molar volume of H_2 at STP = 22.7L)

[31-Jan-2023 Shift 1]

Answer: 4

Solution:



$$\text{Moles of Zn used} = \frac{11.5}{65.4} = \text{Moles of H}_2 \text{ evolved}$$





$$\text{Volume of H}_2 = \frac{11.5}{65.4} \times 22.7\text{L} = 3.99\text{L}$$

Question20

When a hydrocarbon A undergoes complete combustion it requires 11 equivalents of oxygen and produces 4 equivalents of water. What is the molecular formula of A ?


[31-Jan-2023 Shift 2]

Options:



Answer: A

Solution:



$$\frac{y}{2} = 4 \therefore y = 8$$


$$x + \frac{8}{4} = 11$$

$$\therefore x = 9$$

$$\therefore \text{Hydrocarbon will be } = \text{C}_9\text{H}_8$$

Question21

Assume carbon burns according to following equation :



When 12g carbon is burnt in 48g of oxygen, the volume of carbon monoxide produced is _____ $\times 10^{-1}$ L at STP [nearest integer]


[Given : Assume CO as ideal gas, Mass of C is 12gmol^{-1} , Mass of O is 16gmol^{-1} and molar volume of an ideal gas at STP is 22.7Lmol^{-1}]

[31-Jan-2023 Shift 2]

Answer: 227

Solution:

Solution:



1 mol 1.5 mol

Limiting reagent is carbon. One mole carbon produces one mole CO. Hence, volume at STP is 22.7×10^{-1} litre

Question22

The density of 3M solution of NaCl is 1.0gmL^{-1} .

Molality of the solution is _____ $\times 10^{-2}\text{m}$

(Nearest integer).

Given: Molar mass of Na and Cl is 23 and 35.5g mol^{-1} respectively.
[1-Feb-2023 Shift 1]

Answer: 364

Solution:

$$\begin{aligned} m &= \frac{1000 \times M}{1000 \times d - M \times M.W \text{ of solute}} \\ &= \frac{1000 \times 3}{1000 \times 1 - (3 \times 58.5)} = 3.64 \\ &= 364 \times 10^{-2} \end{aligned}$$

Question23

The molality of a 10%(v / v) solution of di-bromine solution in CCl_4 (carbon tetrachloride) is 'x'. $x = \text{_____} \times 10^{-2}\text{M}$. (Nearest integer)

[Given : molar mass of $\text{Br}_2 = 160\text{gmol}^{-1}$

atomic mass of C = 12gmol^{-1}

atomic mass of Cl = 35.5gmol^{-1}

density of dibromine = 3.2gcm^{-3}

density of $\text{CCl}_4 = 1.6\text{gcm}^{-3}$]

[1-Feb-2023 Shift 2]



Answer: 139

Solution:

(10 ml solute in 90 ml solvent
mass of solute = $10 \times 3.2 = 32\text{g}$
mass of solvent = 90×1.6
 $m = \frac{32 \times 1000}{160 \times 90 \times 1.6} = 1.388$
 $m = 138.8 \times 10^{-2} = 139$

Question24


If 5 moles of BaCl_2 is mixed with 2 moles of Na_3PO_4 , the maximum number of moles of $\text{Ba}_3(\text{PO}_4)_2$ formed is

(Nearest integer)

[6-Apr-2023 shift 1]

Answer: 1

Solution:



Question25

0.5g of an organic compound (X) with 60% carbon will produce _____ $\times 10^{-1}\text{g}$ of CO_2 on complete combustion.

[8-Apr-2023 shift 1]

Answer: 11

Solution:

$$\text{Moles of carbon} = \frac{0.5 \times 0.6}{12}$$
$$\text{Moles of CO}_2 = \frac{0.5 \times 0.6}{12}$$
$$\text{Mass of CO}_2 = \frac{0.5 \times 0.6}{12} \times 44 = 11 \times 10^{-1} \text{ gram}$$



Question26

Which of the following have same number of significant figures?

A. 0.00253

B. 1. 0003

C. 15.0

D. 163

Choose the correct answer from the options given below

[8-Apr-2023 shift 2]

Options:

A. B and C only

B. A, B and C only

C. A, C and D only

D. C and D only

Answer: C

Solution:

Solution:

0.00253, 15.0, 163

All have three significant figures.

Question27

The number of molecules and moles in 2.8375 litres of O₂ at STP are respectively

[10-Apr-2023 shift 1]

Options:

A. 7.527×10^{22} and 0.125 mol

B. 1.505×10^{23} and 0.250 mol

C. 7.527×10^{23} and 0.125 mol

D. 7.527×10^{22} and 0.250 mol

Answer: A

Solution:

$$\text{Moles of O}_2(n_{\text{O}_2}) = \frac{\text{Volume of O}_2}{22.7} = 0.125 \text{ moles}$$

$$\text{Molecules of O}_2 = \text{moles} \times N_A$$

$$= 0.125 \times 6.022 \times 10^{23}$$

$$= 7.527 \times 10^{22} \text{ molecules}$$

Ans (1) 7.527×10^{22} and 0.125 mole

Question28

Given below are two statements: one is labelled as Assertion A and the other is labelled as Reason R

Assertion A : 3.1500g of hydrated oxalic acid dissolved in water to make 250.0 mL solution will result in 0.1M oxalic acid solution.

Reason R : Molar mass of hydrated oxalic acid is 126gmol^{-1} In the light of the above statements, choose the correct answer from the options given below:

[10-Apr-2023 shift 2]

Options:

A. A is false but R is true

B. A is true but R is false


C. Both A and R are true but R is NOT the correct explanation of A

D. Both A and R are true and R is the correct explanation of A

Answer: D

Solution:

Assertion is correct.



$$M = \frac{3.15 \times 1000}{126 \times 250}$$

$$= \frac{12.6}{126} = 0.1$$

Reason is correct. It is used as a fact in explanation of assertion.

Question29

Match List I with List II



LIST I	LIST II
A 16g of CH ₄ (g)	I. Weight 28g
B 1g of H ₂ (g)	II 60.2 × 10 ²³ electrons
C 1 mole of N ₂ (g)	III Weight 32g
D 0.5 mol of SO ₂ (g)	IV Occupies 11.4L volume at STP

**Choose the correct answer from the options given below:
[10-Apr-2023 shift 2]**

Options:

- A. A-II, B-IV, C-I, D-III
- B. A-II, B-IV, C-III, D-I
- C. A-II, B-III, C-IV, D-I
- D. A-I, B-III, C-II, D-IV

Answer: A

Solution:

Solution:

16gCH₄ = mole = 1
 e⁻ = 60.0 × 10²³
 19 Hz = 0.5 mole = 11.4(L) STP
 1 mole N₂ = 2 rg
 0.5 mol SO₂ = weights 32g.

Question30

**25 mL of silver nitrate solution (1M) is added dropwise to 25 mL of potassium iodide (1.05M) solution. The ion(s) present in very small quantity in the solution is/are
[11-Apr-2023 shift 1]**

Options:

- A. NO₃⁻ only
- B. Ag⁺ and I⁻ both



C. K^+ only

D. I only

Answer: B

Solution:

Solution:

On adding $AgNO_3$ into KI, AgI will form and solubility of AgI is very low.

So, $[Ag^+]$ and $[I^-]$ will be present in very small quantity.

Question31

A solution of sugar is obtained by mixing 200g of its 25% solution and 500g of its 40% solution (both by mass). The mass percentage of the resulting sugar solution is _____ (Nearest integer)

[11-Apr-2023 shift 1]

Answer: 36

Solution:

$$\text{Solution (I)} \rightarrow \text{Mass of sugar} = 200 \times \frac{25}{100} = 50 \text{ gm}$$

$$\text{Mass of solution} = 200 \text{ gm}$$

$$\text{Solution (II)} \rightarrow \text{Mass of solution} = 500 \text{ gm}$$

$$\text{Mass of sugar} = \frac{40}{100} \times 500 = 200 \text{ gm}$$

$$\text{Final \%w / w} = \frac{\text{Total mass of sugar}}{\text{Total mass of solution}} \times 100$$

$$= \frac{50 + 200}{200 + 500} \times 100 = \frac{250}{7}$$

$$= 35.71\% \approx 36$$

Question32

A solution is prepared by adding 2g of " X " to 1 mole of water. Mass percent of " X " in the solution is

[11-Apr-2023 shift 2]

Options:

A. 5%

B. 20%

C. 2%



D. 10%

Answer: D

Solution:

Solute (X) = 2g

Solvent (H₂O) = 1 mole = 18g

Total mass = 2 + 18 = 20g

% mass of X = $\frac{2}{20} \times 100 = 10\%$

Question33


The volume of hydrogen liberated at STP by treating 2.4g of magnesium with excess of hydrochloric acid is _____ $\times 10^{-2}$ L.

Given: Molar volume of gas is 22.4L at STP. Molar mass of magnesium is 24gmol⁻¹

[11-Apr-2023 shift 2]

Answer: 224

Solution:



w = 2.4g

N = $\frac{2.4}{24} = 0.1$ mole

1 mole of gas at STP \Rightarrow 22.4 lit.

\therefore 0.1 mole of gas = 0.1 \times 22.4

= 2.24 lit. = 224 $\times 10^{-2}$ litre

Question34

A metal chloride contains 55.0% of chlorine by weight . 100 mL vapours of the metal chloride at STP weigh 0.57g. The molecular formula of the metal chloride is (Given: Atomic mass of chlorine is 35.5u)

[12-Apr-2023 shift 1]

Options:

A. MCl

B. MCl₃

C. MCl₂

D. MCl_4

Answer: C

Solution:

Solution:

Molecular weight of metal chloride

$$= \frac{0.57}{100} \times 22700$$

$$= 129.39$$

$$\text{weight of Cl} = 129.39 \times 0.55$$

$$= 71.1645$$

$$\therefore \text{Mole of } MCl = \frac{71.1645}{35.5} \cong 2$$

Hence MCl_2

Question35

An organic compound gives 0.220g of CO_2 and 0.126g of H_2O on complete combustion. If the % of carbon is 24 then the % of hydrogen is _____ $\times 10^{-1}$. (Nearest integer)

[13-Apr-2023 shift 1]

Answer: 56

Solution:

Solution:

$$\% \text{ of carbon} = \frac{\frac{0.220}{44} \times 12}{x} \times 100$$

(x = mass of organic compound)

$$24 = \frac{6}{x}$$

$$x = 0.25 \text{ gm}$$

$$\% \text{ of H} = \frac{\frac{0.126}{18} \times 2 \times 1}{0.25} \times 100$$

$$= 5.6 = 56 \times 10^{-1}$$

Question36

1g of a carbonate (M_2CO_3) on treatment with excess HCl produces 0.01 mol of CO_2 . The molar mass of M_2CO_3 is _____ $gmol^{-1}$. (Nearest integer)


[13-Apr-2023 shift 2]



Answer: 100

Solution:

Solution:



From principle of atomic conservation of carbon atom, Mole of $\text{M}_2\text{CO}_3 \times 1 = \text{Mole of CO}_2 \times 1$

$$\frac{1\text{ gm}}{\text{molar mass of M}_2\text{CO}_3} = 0.01 \times 1$$

$$\therefore \text{Molar mass of M}_2\text{CO}_3 = 100\text{ gm / mole}$$

Question37

Compound A contains 8.7% Hydrogen, 74% Carbon and 17.3% Nitrogen.





The molecular formula of the compound is,

Given : Atomic masses of C, H and N are 12,1 and 14 amu respectively.

The molar mass of the compound A is 162gmol^{-1} .

[28-Jun-2022-Shift-2]

Options:



Answer: D

Solution:

Solution:

Mole ratio of H, C and N

$$= \frac{8.7}{1} : \frac{74}{12} : \frac{17.3}{14}$$

$$= 8.7 : 6.167 : 1.23$$

$$= \frac{8.7}{1.23} : \frac{6.167}{1.23} : \frac{1.23}{1.23}$$

$$= 7 : 5 : 1$$

$$\therefore \text{Emperical formula} = \text{C}_5\text{H}_7\text{N}$$

$$\therefore \text{Molecular formula} = (\text{C}_5\text{H}_7\text{N})_n$$

$$\text{Given molecular mass} = 162$$

$$\text{Molecular mass of } (\text{C}_5\text{H}_7\text{N})_n$$

$$= (5 \times 12 + 7 \times 1 + 14) \times n$$

$$= (81) \times n$$

$$\therefore 81 \times n = 162$$

$$\Rightarrow n = 2$$

$$\therefore \text{Molecular formula} = \text{C}_{10}\text{H}_{14}\text{N}_2$$



Question38

The complete combustion of 0.492g of an organic compound containing ' C ', ' H ' and ' O ' gives 0.793g of CO₂ and 0.442g of H₂O. The percentage of oxygen composition in the organic compound is _____ (nearest integer)
[28-Jun-2022-Shift-2]

Answer: 46

Solution:

Solution:

Total organic compound = 0.492 gm

Produced CO₂ = 0.793 gm

$$\therefore \text{Moles of CO}_2 = \frac{0.793}{44}$$

$$\therefore \text{Moles of C atoms} = \frac{0.793}{44}$$

$$\therefore \text{Weight of C atoms} = \frac{0.793}{44} \times 12 = 0.216\text{g}$$

Produced H₂O = 0.442 gm

$$\therefore \text{Moles of H}_2\text{O} = \frac{0.442}{18}$$

$$\therefore \text{Moles of H atoms} = \frac{0.442}{18} \times 2$$

$$\therefore \text{Weight of H atoms} = \frac{0.442}{18} \times 2 = 0.05\text{g}$$

\therefore Weight of O atoms

$$= 0.492 - (0.216 + 0.05)$$


$$= 0.226\text{ gm}$$

% by mass of oxygen in compound

$$= \frac{0.226}{0.492} \times 100 = 46\%$$

Question39

Production of iron in blast furnace follows the following equation



when 4.640 kg of Fe₃O₄ and 2.520 kg of CO are allowed to react then the amount of iron (in g) produced is:

[29-Jun-2022-Shift-1]

Options:

A. 1400

B. 2200

C. 3360

D. 4200

Answer: C

Solution:

Given,

Mass of $\text{Fe}_3\text{O}_4 = 4.640 \text{ kg} = 4640 \text{ gm}$

Molar mass of $\text{Fe}_3\text{O}_4 = 56 \times 3 + 16 \times 4 = 232 \text{ g}$


$$\therefore \text{Moles of } \text{Fe}_3\text{O}_4 = \frac{4640}{232} = 20$$

Also, given

Mass of $\text{CO} = 2.520 \text{ kg} = 2520 \text{ gm}$

Molar mass of $\text{CO} = 12 + 16 = 28 \text{ gm}$

$$\therefore \text{Molar of } \text{CO} = \frac{2520}{28} = 90$$



Here Fe_3O_4 is limiting reagent as to find limiting reagent, divide the given moles of reactants with their respective stoichiometric coefficient and reactant for which this ratio is minimum will be limiting reagent

$$\text{For } \text{Fe}_3\text{O}_4, \frac{\text{moles}}{\text{stoichiometric coefficient}} = \frac{20}{1}$$

$$\text{For } \text{CO}, \frac{\text{moles}}{\text{stoichiometric coefficient}} = \frac{90}{4} = 22.5$$

$\therefore \text{Fe}_3\text{O}_4$ is limiting reagent.

Now produced $\text{Fe} = 20 \times 3 = 60 \text{ mol}$

\therefore Weight of $\text{Fe} = 60 \times 56 = 3360 \text{ g}$

Question40

Geraniol, a volatile organic compound, is a component of rose oil. The density of the vapour is 0.46 gL^{-1} at 257°C and 100 mm Hg . The molar mass of geraniol is _____ gmol^{-1} . (Nearest Integer)

[. Given : $R = 0.082 \text{ L atm K}^{-1} \text{ mol}^{-1}$]

[29-Jun-2022-Shift-1]

Answer: 152

Solution:

From ideal gas equation we know

$$PV = nRT$$

$$\Rightarrow PV = \frac{W}{M}RT$$

$$\Rightarrow P = \frac{W}{V} \cdot \frac{RT}{M}$$

$$\Rightarrow P = d \cdot \frac{RT}{M} \left[\because d = \frac{W}{V} \right]$$

We know, 760 mm of Hg = 1 atm

$$\therefore 100 \text{ mm of Hg} = \frac{100}{760} \text{ atm}$$

$$\therefore \text{Pressure (P)} = \frac{100}{760} \text{ atm}$$

Density (d) = 0.46

$$R = 0.082 \text{ L atm K}^{-1} \text{ mol}^{-1}$$

$$T = (257 + 273) \text{ K} = 530 \text{ K}$$

Putting the values in above equation, we get

$$\frac{100}{760} = \frac{0.46 \times 0.082 \times 530}{M}$$

$$\Rightarrow M = 152$$

Question41

Using the rules for significant figures, the correct answer for the expression $\frac{0.02858 \times 0.112}{0.5702}$ will be

[29-Jun-2022-Shift-2]

Options:

A. 0.005613

B. 0.00561

C. 0.0056

D. 0.006

Answer: B

Solution:

Solution:

$$\frac{0.02858 \times 0.112}{0.5702} = 0.00561$$

Reported answer should not be more precise than least precise term (0.112 is the least precise term with three significant figures) in calculations, so there should be three significant figures in reported answer.

Question42

If a rocket runs on a fuel ($C_{15}H_{30}$) and liquid oxygen, the weight of oxygen required and CO_2 released for every litre of fuel respectively are

:

(Given : density of the fuel is 0.756g / mL)

[24-Jun-2022-Shift-1]

Options:

A. 1188g and 1296g




B. 2376g and 2592g

C. 2592g and 2376g

D. 3429g and 3142g

Answer: C

Solution:



Given, volume of fuel = 1L = 1000ml

And density of fuel = 0.756g / ml

We know,

$$d = \frac{w}{v}$$

$$\Rightarrow 0.756 = \frac{w}{1000}$$

$$\Rightarrow w = 756\text{gm}$$

\therefore weight of fuel = 756gm

Molar mass of $\text{C}_{15}\text{H}_{30} = 15 \times 12 + 30 = 210$

$$\therefore \text{Moles of } \text{C}_{15}\text{H}_{30} = \frac{756}{210}$$

From equation you can see,

1 mole of $\text{C}_{15}\text{H}_{30}$ react with $\frac{45}{2}$ mole of O_2

$\therefore \frac{756}{210}$ moles of $\text{C}_{15}\text{H}_{30}$ react with $\frac{45}{2} \times \frac{756}{210}$ moles of O_2

$$\therefore \text{Moles of } \text{O}_2 \text{ required} = \frac{45}{2} \times \frac{756}{210}$$

$$\therefore \text{Mass of } \text{O}_2 \text{ required} = \frac{45}{2} \times \frac{756}{210} \times 32 = 2592\text{g}$$

Also,

From 1 mole of $\text{C}_{15}\text{H}_{30}$ 15 moles of CO_2 formed

\therefore From $\frac{756}{210}$ moles of $\text{C}_{15}\text{H}_{30}$ $15 \times \frac{756}{210}$ moles of CO_2 formed

$$\therefore \text{Moles of } \text{CO}_2 \text{ formed} = 15 \times \frac{756}{210}$$

$$\therefore \text{Mass of } \text{CO}_2 \text{ formed} = 15 \times \frac{756}{210} \times 44 =$$

2376g

Question43

120g of an organic compound that contains only carbon and hydrogen gives 330g of CO_2 and 270g of water on complete combustion. The percentage of carbon and hydrogen, respectively are [24-Jun-2022-Shift-2]

Options:

A. 25 and 75

B. 40 and 60


C. 60 and 40

D. 75 and 25

Answer: D

Solution:

Solution:



From the reaction,

Produced $CO_2 = x$ mol

and produced $H_2O = \frac{y}{2}$ mol

Given produced $CO_2 = 330$ g

$$\therefore \text{moles of } CO_2 = \frac{330}{44} = \frac{30}{4} = x$$

Also given produced $H_2O = 270$ gm

$$\therefore \text{Moles of } H_2O = \frac{270}{18} = 15 = \frac{y}{2}$$

$$\Rightarrow y = 30$$

$$\therefore x : y = \frac{30}{4} : 30 = 1 : 4$$

Formula of the compound = $(CH_4)_n$

\therefore Weight of C in $(CH_4)_n = 12n$

Weight of H in $(CH_4)_n = 4n$

\therefore Weight ratio of C and H

$$= 12n : 4n$$

$$= 3 : 1$$

$$\therefore \% \text{ of C} = \frac{3}{4} \times 100 = 75$$

$$\text{and \% of H} = \frac{1}{4} \times 100 = 25$$

Question44

The number of N atoms in 681g of $C_7H_5N_3O_6$ is $x \times 10^{21}$. The value of x is ($N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$)

(Nearest Integer)

[25-Jun-2022-Shift-1]

Answer: 5418

Solution:

Molar mass of $C_7H_5N_3O_6$

$$= 12 \times 7 + 5 + 14 \times 3 + 16 \times 6$$

$$= 227 \text{ gm}$$

Given mass of $C_7H_5N_3O_6 = 681$ gm

$$\therefore \text{Moles of } C_7H_5N_3O_6 = \frac{681}{227} = 3$$

In one molecule of $C_7H_5N_3O_6$, 3N atoms present.

\therefore In 1 mole $C_7H_5N_3O_6$, 3 moles of N atoms presents.

\therefore In 3 moles of $C_7H_5N_3O_6$, $3 \times 3 = 9$ moles of N atoms presents.

We know, 1 mole of N atoms = 6.02×10^{23} N atoms.

$$\therefore 9 \text{ moles of N atoms} = 9 \times 6.02 \times 10^{23} = 54.18 \times 10^{23} = 5418 \times 10^{21}$$

Question45

A protein ' A ' contains 0.30% of glycine (molecular weight 75). The minimum molar mass of the protein ' A ' is $___ \times 10^3 \text{ gmol}^{-1}$ [nearest integer]

[25-Jun-2022-Shift-2]

Answer: 25

Solution:

Solution:

Let, molar mass of protein A = x
Protein A contains 0.30% glycine

$$\therefore \frac{x \times 0.3}{100} = 75$$

$$\Rightarrow x = 25000 = 25 \times 10^3$$

Question46

A commercially sold conc. HCl is 35% HCl by mass. If the density of this commercial acid is 1.46g / mL, the molarity of this solution is:
(Atomic mass : Cl = 35.5 amu, H = 1 amu)

[26-Jun-2022-Shift-1]

Options:

A. 10.2M

B. 12.5M

C. 14.0M

D. 18.2M

Answer: C

Solution:

Solution:

35% HCl by mass means in 100 gm HCl solution 35 gm HCl present.

Now, volume of 100 gm HCl solution

$$= \frac{100}{1.46} \text{ ml}$$

$$= \frac{100}{1.46} \text{ l}$$

$$\text{Moles of HCl} = \frac{35}{36.5}$$



$$\text{Now, molarity} = \frac{\text{moles of solute}}{\text{volume of solution (in L)}}$$


$$= \frac{\frac{35}{36.5}}{\frac{100}{1.46}} = 14$$

Question47

On complete combustion 0.30g of an organic compound gave 0.20g of carbon dioxide and 0.10g of water. The percentage of carbon in the given organic compound is ____ (Nearest integer)
[26-Jun-2022-Shift-1]

Answer: 18

Solution:



Given organic compound $C_xH_y = 0.3 \text{ gm}$

Produced carbon dioxide (CO_2) = 0.2 gm

Produced water (H_2O) = 0.1 gm

$$\text{Moles of } CO_2 = \frac{0.2}{44}$$

$$\therefore \text{Moles of C atom} = \frac{0.2}{44}$$

$$\therefore \text{Mass of C atom} = \frac{0.2}{44} \times 12 = 0.0545$$

$$\text{Moles of } H_2O = \frac{0.1}{18}$$

$$\therefore \text{Moles of H atoms} = \frac{0.1}{18} \times 2$$

$$\therefore \text{Mass of H atoms} = \frac{0.1 \times 2}{18} \times 1 = 0.0111$$

$$\therefore \% \text{ of C atom} = \frac{0.0545}{0.3} \times 100 = 18\%$$


Question48

The moles of methane required to produce 81g of water after complete combustion is ____ $\times 10^{-2}$ mol. [nearest integer]
[26-Jun-2022-Shift-2]

Answer: 225

Solution:





POAC on H atom

$$n_{\text{CH}_4} \times 4 = n_{\text{H}_2\text{O}} \times 2$$

$$n_{\text{CH}_4} = \frac{81}{18} \times 2 \times \frac{1}{4} = \frac{81}{36}$$

$$n_{\text{CH}_4} = 2.25$$

$$= 225 \times 10^{-2}$$

$$\text{Nearest Integers} = 225$$

Question49

116g of a substance upon dissociation reaction, yields 7.5g of hydrogen, 60g of oxygen and 48.5g of carbon. Given that the atomic masses of H, O and C are 1, 16 and 12, respectively. The data agrees with how many formulae of the following?

- A. CH_3COOH ,
- B. HCHO ,
- C. CH_3OOCH_3 ,
- D. CH_3CHO

[27-Jun-2022-Shift-2]

Answer: 2

Solution:

$$\% \text{H} = \frac{7.5}{116} \times 100 = 6.5$$

$$\% \text{O} = \frac{60}{116} \times 100 = 51.7$$

$$\% \text{C} = \frac{48.5}{116} \times 100 = 41.8$$

Relative atomicities = H \Rightarrow 6.5

$$\text{O} \Rightarrow \frac{51.7}{16} = 3.25$$

$$\text{C} \Rightarrow \frac{41.8}{12} = 3.5$$

Empirically formula is approx.. CH_2O

(A) $\text{C}_2\text{H}_4\text{O}_2$ (B) CH_2O relate to this formula.

Question50


56.0 L of nitrogen gas is mixed with excess hydrogen gas and it is found that 20L of ammonia gas is produced. The volume of unused nitrogen gas is found to be _____ L.

[25-Jul-2022-Shift-2]

Answer: 46

Solution:

Solution:



Since H_2 is in excess and 20L of ammonia gas is produced.

Hence, 2 moles $\text{NH}_3 \equiv 1$ mole N_2 ($v \propto n$)

$$20\text{LNH}_3 \equiv 10\text{LN}_2$$

$$\text{Volume of N}_2 \text{ left} = 56 - 10$$

$$= 46\text{L}$$

Question51

Chlorophyll extracted from the crushed green leaves was dissolved in water to make 2L solution of Mg of concentration 48 ppm. The number of atoms of Mg in this solution is $x \times 10^{20}$ atoms. The value of x is _____ . (Nearest Integer)

(Given : Atomic mass of Mg is 24gmol^{-1} ; $N_A = 6.02 \times 10^{23}\text{mol}^{-1}$)

[26-Jul-2022-Shift-1]

Answer: 24

Solution:

Solution:

In 2L \rightarrow 96 mg of Mg

$$\text{Number of atoms of Mg} = \frac{96 \times 10^{-3}}{24} \times N_A$$

$$= 4 \times 10^{-3} \times 6 \times 10^{23}$$

$$= 24 \times 10^{20}$$

$$x = 24$$

Question52

When 800 mL of 0.5M nitric acid is heated in a beaker, its volume is reduced to half and 11.5g of nitric acid is evaporated. The molarity of the remaining nitric acid solution is $x \times 10^{-2}\text{M}$. (Nearest integer)

(Molar mass of nitric acid is 63gmol^{-1})

[26-Jul-2022-Shift-1]



Answer: 54

Solution:

Solution:

$$m \text{ moles of HNO}_3 = 800 \times 0.5$$

$$\text{Moles of HNO}_3 = 400 \times 10^{-3} = 0.4 \text{ moles}$$

$$\text{Weight of HNO}_3 = 0.4 \times 63\text{g} = 25.2\text{g}$$

$$\text{Remaining acid} = 25.2 - 11.5 = 13.7\text{g}$$

$$M = \frac{13.7 \times 1000}{400 \times 63}$$

$$= \frac{137}{252} = 0.54$$

$$= 54 \times 10^{-2}$$

Question53

Hemoglobin contains 0.34% of iron by mass. The number of Fe atoms in 3.3g of hemoglobin is (Given: Atomic mass of Fe is 56u, $N_A = 6.022 \times 10^{23} \text{mol}^{-1}$.)

[26-Jul-2022-Shift-2]

Options:

A. 1.21×10^5

B. 12.0×10^{16}

C. 1.21×10^{20}

D. 3.4×10^{22}

Answer: C

Solution:

Solution:

According to the question,

100g of hemoglobin contains 0.34g of iron

3.3g of hemoglobin contains $\frac{0.34}{100} \times 3.3\text{g}$ of iron

$$\text{moles of Fe} = \frac{0.34 \times 3.3}{100 \times 56} = \frac{N}{N_A}$$

$$N = \frac{0.34 \times 3.3 \times 6.022 \times 10^{23}}{100 \times 56}$$
$$= 1.21 \times 10^{20}$$

Question54

250g solution of D-glucose in water contains 10.8% of carbon by weight. The molality of the solution is nearest to



(Given: Atomic Weights are, H, 1u; C, 12u; O, 16u)
[27-Jul-2022-Shift-1]

Options:

- A. 1.03
- B. 2.06
- C. 3.09
- D. 5.40

Answer: B

Solution:

Solution:

$C_6H_{12}O_6 \rightarrow$ Glucose

$$\text{We know: } \frac{\text{mass of C}}{\text{mass of glucose}} = \frac{72}{180}$$

$$\text{Given: } \%C = 10.8 = \frac{\text{mass of C}}{\text{mass of solution}} \times 100$$

$$\frac{10.8 \times 250}{100} = \text{mass of C} \Rightarrow \text{Mass of C} = 27 \text{ gm}$$

$$\therefore \text{mass of glucose} = 67.5 \text{ gm}$$

$$\therefore \text{moles of glucose} = 0.375 \text{ moles}$$

$$\text{Mass of solvent} = 250 - 67.5 \text{ gm} = 182.5 \text{ gm}$$

$$\therefore \text{Molality} = \frac{0.375}{0.1825} = 2.055 \approx 2.06$$

Question55

The normality of H_2SO_4 in the solution obtained on mixing 100 mL of 0.1M H_2SO_4 with 50 mL of 0.1M NaOH _____ is $\times 10^{-1}N$. (Nearest Integer)

[27-Jul-2022-Shift-2]

Answer: 1

Solution:

Solution:

$$\text{No. of equivalents of } H_2SO_4 = 100 \times 0.1 \times 2 = 20$$

$$\text{No. of equivalents of NaOH} = 50 \times 0.1 = 5$$


$$\text{No. of equivalents of } H_2SO_4 \text{ left} = 20 - 5 = 15$$

$$\Rightarrow 150 \times x = 15$$

$$x = \frac{1}{10} = 0.1N = 1 \times 10^{-1}N$$



Question56



In the above reaction, 5g of toluene is converted into benzaldehyde with 92% yield. The amount of benzaldehyde produced is _____ $\times 10^{-2}$ g. (Nearest integer)
[27-Jul-2022-Shift-2]

Answer: 530

Solution:



$$\text{Moles} = \frac{5}{92}$$

$$\text{Moles of benzaldehyde produced} = \frac{5}{92} \times 0.92 = 0.05$$

$$\begin{aligned} \therefore \text{Mass of benzaldehyde formed} &= 0.05 \times 106 \\ &= 5.3\text{g} \\ &= 530 \times 10^{-2} \end{aligned}$$

Question57

In the given reaction,




if one mole of each of X and Y with 0.05 mol of Z gives compound XYZ_3 - (Given : Atomic masses of X, Y and Z are 10, 20 and 30 amu, respectively.) The yield of XYZ_3 is _____g. (Nearest integer)

[28-Jul-2022-Shift-1]

Answer: 2

Solution:





$$\text{Limiting reagent is } Z = \frac{0.05}{3} = .016$$

3 moles of Z → 1 mole of XYZ₃

0.05 mole of Z → $\frac{1}{3} \times 0.05$ mole of XYZ₃

$$\begin{aligned} \text{M.wt. of XYZ}_3 &= 10 + 20 + 90 \\ &= 120 \text{ amu} \end{aligned}$$

$$\begin{aligned} \text{Wt. of XYZ}_3 &= \frac{.05}{3} \times 120 \\ &= 2\text{g} \end{aligned}$$

Question58

**On complete combustion of 0.492g of an organic compound containing C, H and O, 0.7938g of CO₂ and 0.4428g of H₂O was produced. The % composition of oxygen in the compound is _____.
[28-Jul-2022-Shift-1]**

Answer: 46

Solution:

$$\% \text{ of H} = \frac{2}{18} \times \frac{\text{wt. of H}_2\text{O}}{\text{wt. of organic compound}} \times 100$$

$$= \frac{2}{18} \times \frac{0.4428}{0.492} \times 100$$

$$= 0.11 \times 0.9 \times 100$$

$$= 0.099 \times 100 = 9.9$$

$$\% \text{ of C} = \frac{12}{44} \times \frac{0.7938}{0.492} \times 100$$


$$= 0.27 \times 1.61 \times 100$$

$$= 43.47$$

$$\% \text{ Oxygen} = 100 - (43.47 + 9.9)$$

$$= 100 - 53.37 \approx 46$$

Question59



Consider the above reaction, the limiting reagent of the reaction and number of moles of NH₃ formed respectively are :

[29-Jul-2022-Shift-1]

Options:

A. H₂, 1.42 moles

B. H₂, 0.71 moles




C. N₂, 1.42 moles

D. N₂, 0.71 moles

Answer: C

Solution:

Solution:



Ideally 28gN₂ reacts with 6gH₂ limiting reagent is N₂


∴ Amount of NH₃ formed on reacting 20gN is,

$$= \frac{34 \times 20}{28} = 24.28g$$

$$= 1.42 \text{ moles}$$

Question60

Consider the reaction



The amount of HNO₃ required to produce 110.0g of KNO₃ is

(Given: Atomic masses of H, O, N and K are 1, 16, 14 and 39 , respectively.)

[29-Jul-2022-Shift-2]

Options:

A. 32.2g

B. 69.4g


C. 91.5g

D. 162.5g

Answer: C

Solution:

Solution:



$$\therefore 110g \text{ of } KNO_3 \Rightarrow \text{moles of } KNO_3 = \frac{110}{101}$$

$$= 1.089 \text{ mol}$$


As, 4 mole of HNO₃ produces 3 mol of KNO₃.

Hence, the moles of HNO₃ required to produce

$$1.089 \text{ moles of } KNO_3 = \frac{4}{3} \times 1.089 = 1.452 \text{ mol}$$

Hence, mass of HNO₃ required is $1.452 \times 63 = 91.5g$

Question61



When coal of purity 60% is allowed to burn in presence of insufficient oxygen, 60% of carbon is converted into 'CO' and the remaining is converted into 'CO₂'. The heat generated when 0.6 kg of coal is burnt is _____.

[29-Jul-2022-Shift-2]

Options:

A. 1600 kJ

B. 3200 kJ

C. 4400 kJ

D. 6600 kJ

Answer: D

Solution:

Solution:

Weight of coal = 0.6 kg = 600 gm

∴ 60% of it is carbon

So weight of carbon = $600 \times \frac{60}{100} = 360 \text{ g}$

∴ moles of carbon = $\frac{360}{12} = 30 \text{ moles}$

∴ Heat generated = $12 \times 400 + 18 \times 100 = 6600 \text{ kJ}$

Question62

A 1.84 mg sample of polyhydric alcoholic compound 'X' of molar mass 92.0g / mol gave 1.344 mL of H₂ gas at STP. The number of alcoholic hydrogens present in compound 'X' is _____.

[29-Jul-2022-Shift-2]

Answer: 3

Solution:

Solution:

Volume of H₂ gas = 1.344 mL

Mole of H₂ gas = $\frac{1.344}{22400} = 6 \times 10^{-5}$

No of H atoms per molecule of H₂ = 2.

No. of moles of organic compound = $\frac{1.84 \times 10^{-3}}{92} = 2 \times 10^{-5}$



$$\text{No. of } -\text{OH (hydroxyl group in one molecule)} = \frac{6 \times 10^{-5}}{2 \times 10^{-5}} = 3$$

Question63

The number of significant figures in 50000.020×10^{-3} is
[26 Feb 2021 Shift 1]

Answer: 7

Solution:

Solution:

Non-zero digits are always significant. Any zeros between two significant digits are significant.
 \therefore Zero's between 5 and 2 are all significant.
(Number of significant figures = 7)

Question64

The NaNO_3 weighed out to make 50mL of an aqueous solution containing 70.0mg Na^+ per mL is g. (Rounded off to the nearest integer) [Given : Atomic weight in gmol^{-1} , -Na : 23; N : 14; O : 16].
[26 Feb 2021 Shift 2]

Answer: 13

Solution:

Solution:

70mg Na^+ is present in 1mL of NaNO_3 solution.

\therefore 50mL of NaNO_3 will contain = $70 \times 50\text{mg}$

$$= \frac{70 \times 50}{1000} = 3.5\text{gm}$$

Moles of Na^+ in solution = Moles of NaNO_3 in solution

[$\therefore \text{NaNO}_3 \rightarrow \text{Na}^+ + \text{NO}_3^-$]

$$= \frac{3.5}{23} \text{mol} \quad [\therefore \text{Molar mass of } \text{Na}^+ = 23\text{gmol}^{-1}]$$

Mass of NaNO_3 = mole \times molar mass

[\therefore molar mass of $\text{NaNO}_3 = 85\text{gmol}^{-1}$]

$$= \frac{3.5}{23} \times 85 = 12.934\text{g} \sim \text{eq}13\text{g}$$



Question65

Complete combustion of 1.80g of an oxygen containing compound ($C_xH_yO_z$) gave 2.64g of CO_2 and 1.08g of H_2O . The percentage of oxygen in the organic compound is [25 Feb 2021 Shift 1]


Options:

- A. 50.33
- B. 53.33
- C. 63.53
- D. 51.63

Answer: B

Solution:

Solution:



$$n_C = \frac{CO}{(Moles)_2} = \frac{2.64 \text{ (Given mass)}}{44 \text{ (Molecular mass)}} = 0.06$$

$$n_H = 2 \times n_{H_2O} = \frac{1.08}{18} \times 2 = 0.12$$

Weight of oxygen in $C_xH_yO_z$

$$= 1.80 - 12 \times \frac{2.64}{44} - \frac{1.08}{18} \times 2$$

$$= 1.80 - 0.72 - 0.12 = 0.96g$$

$$\% \text{ of oxygen by weight} = \frac{0.96}{1.80} \times 100 = 53.33\%$$


Question66

The formula of a gaseous hydrocarbon, which requires 6 times of its own volume of O_2 for complete oxidation and produces 4 times its own volume of CO_2 is C_xH_y . The value of y is [24 Feb 2021 Shift 2]

Answer: 8

Solution:

Combustion reaction:



Suppose, volume of C_xH_y is V and volume of O_2 is 6 times greater than $C_xH_y = 6V$

then volume of $x\text{CO}_2 \Rightarrow V_x = 4V$

$$x = 4$$

Since, $V_{\text{O}_2} = 6 \times V_{\text{C}_x\text{H}_y}$

$$V \left(x + \frac{y}{4} \right) = 6V$$

$$\left(x + \frac{y}{4} \right) = 6$$

Put value of $x = 4$ in Eq. (i) We get,

$$4 + \frac{y}{4} = 6 \Rightarrow y = 8$$


Question67

1.86g of aniline completely reacts to form acetanilide. 10% of the product is lost during purification. Amount of acetanilide obtained after purification (in g) is $\times 10^{-2}$.

[24 Feb 2021 Shift 2]

Answer: 243

Solution:



Given, weight = 1.86g

Here, 1 mole of aniline gives 1 mole of acetanilide

\therefore mole of aniline = mole of acetanilide

$$\Rightarrow \frac{1.86}{93} = \frac{W_{\text{Acetanilide}}}{135}$$

$$W_{\text{Acetanilide}} = \frac{1.86 \times 135}{93} \text{g} = 2.70 \text{g}$$

But efficiency of reaction is 90% only.

Hence, mass of acetanilide produced

$$= 2.70 \times \frac{90}{100} \text{g} = 2.43 \text{g} = 243 \times 10^2 \text{g}$$

$$x = 243$$

Question68


15mL of aqueous solution of Fe^{2+} in acidic medium completely reacted with 20mL of 0.03M aqueous $\text{Cr}_2\text{O}_7^{2-}$. The molarity of the Fe^{2+} solution is $\times 10^{-2}\text{M}$ (Round off to the nearest integer).

[17 Mar 2021 Shift 1]



Answer: 24

Solution:



Dichromate ($\text{Cr}_2\text{O}_7^{2-}$) oxidised F e^{2+} to F e^{3+} and itself get reduced to Cr^{3+} .

Valency factor of $\text{Cr}_2\text{O}_7^{2-} = 6$ as Cr gets reduced from +6 to +3

From law of equivalence,

Milliequivalent of $\text{Cr}_2\text{O}_7^{2-} = \text{Milliequivalent of F e}^{2+}$

$$M_1 V_1 n_1 = M_2 V_2 n_2$$

$$(M_1 \times V_1) \times 6 = (M_2 \times V_2) \times 1$$

$$(0.03 \times 20) \times 6 = (M_2 \times 15) \times 1$$

$$\Rightarrow M_2 = 0.24 \text{ molar}$$

$$= 24 \times 10^{-2} \text{ molar}$$

Question69

**The mole fraction of a solute in a 100 molal aqueous solution
 $\times 10^{-2}$ (Round off to the nearest integer).**

[Given, atomic masses H : 1.0u, O : 16.0u]

[17 Mar 2021 Shift 1]

Answer: 64

Solution:

Solution:

Given, molality = 100

$\Rightarrow 100$ moles of solute in 1kg of solvent

$$\text{Mole fraction of solute } (X_{\text{solute}}) = \frac{\text{Moles of solute}}{\text{Total moles}}$$

$$= \frac{n_{\text{Solute}}}{n_{\text{Solute}} + n_{\text{Solvent}}}$$

$$n_{\text{solvent}} = \frac{1000}{18} = 55.5$$

$$X_{\text{solute}} = \frac{100}{100 + 55.5} = \frac{100}{155.5} = 0.643$$

$$X_{\text{solute}} = 64.3 \times 10^{-2} \Rightarrow 64 \times 10^{-2}$$

Question70

10.0mL of $\text{N a}_2\text{C O}_3$ solution is titrated against 0.2M H Cl solution. The following titre values were obtained in 5 readings. 4.8mL, 4.9mL, 5.0mL, 5.0mL and 5.0mL based on these readings and convention of




titrimetric estimation of concentration of Na_2CO_3 solution is mM
(Round off to the nearest integer).
[18 Mar 2021 Shift 2]

Answer: 50

Solution:

10.0mL of Na_2CO_3 solution is titrated against 0.2M HCl

Volume of Na_2CO_3 solution (mL)	Volume of 0.2M HCl solution (mL)	Mean volume of HCl (ml)
10	4.8	
10	4.9	
10	5.0	5.0
10	5.0	
10	5.0	



10mL 0.2M_{HCl}

$M_{\text{Na}_2\text{CO}_3} = ?$ 5mL

M_{eq} of $\text{Na}_2\text{CO}_3 = M_{\text{eq}}$ of HCl

$M_{\text{Na}_2\text{CO}_3} \times 10 \times 2 = 0.2 \times 5 \times 1$

$M_{\text{Na}_2\text{CO}_3} = 5 \times 10^{-2}\text{M} = 50 \times 10^{-3}\text{M} = 50\text{mM}$


Question71

Complete combustion of 750g of an organic compound provides 420g of CO_2 and 210g of H_2O . The percentage composition of carbon and hydrogen in organic compound is 15.3 and respectively (Round off to the nearest integer).
[16 Mar 2021 Shift 1]

Answer: 3

Solution:





Any hydrocarbon, on combustion gives CO_2 and H_2O . This is Liebig's method for estimation of 'C' and 'H' percentage.

Mass of water formed = 210g

18g of H_2O contains = 2g of hydrogen

210g of H_2O contains = $\left(\frac{2}{18} \times 210 = \frac{70}{3}\right)$ g of hydrogen.

Given, mass of organic compound = 750g

Percentage of hydrogen

$$= \frac{\text{Mass of hydrogen}}{\text{Mass of organic compound}} \times 100$$

$$= \frac{70}{3 \times 750} \times 100$$

$$= 3.11\%$$

Nearest integer = 3

Question72

10.0 mL of 0.05 M $KMnO_4$ solution was consumed in a titration with 10.0 mL of given oxalic acid dihydrate solution. The strength of given oxalic acid solution is $\times 10^{-2}$ g/L.

(Round off to the nearest integer)

[27 Jul 2021 Shift 2]

Answer: 1575

Solution:

$$n_{eq}KMnO_4 = n_{eq}H_2C_2O_4 \cdot 2H_2O$$

$$\text{or, } \frac{10 \times 0.05}{1000} \times 5 = \frac{10 \times M}{1000} \times 2$$

$$\therefore \text{Conc. of oxalic acid solution} = 0.125M$$

$$= 0.125 \times 126g / L = 15.75g / L$$

$$= 1575 \times 10^{-2}g / L$$

Question73

When 10mL of an aqueous solution of Fe^{2+} ions was titrated in the presence of dil H_2SO_4 using diphenylamine indicator, 15mL of 0.02M solution of $K_2Cr_2O_7$ was required to get the end point. The molarity of the solution containing Fe^{2+} ions is $x \times 10^{-2}M$. The value of x is _____. (Nearest integer)

[25 Jul 2021 Shift 1]

Answer: 18



Solution:

$$\begin{aligned} \text{milli-equivalents of } \text{Fe}^{2+} &= \text{milli-equivalents of } \text{K}_2\text{Cr}_2\text{O}_7 \\ M \times 10 \times 1 &= 0.02 \times 15 \times 6 \\ M &= 0.18 = 18 \times 10^{-2} \text{M} \end{aligned}$$

Question74

If the concentration of glucose ($\text{C}_6\text{H}_{12}\text{O}_6$) in blood is 0.72gL^{-1} , the molarity of glucose in blood is _____ $\times 10^{-3}\text{M}$. (Nearest integer)
[Given: Atomic mass of C = 12, H = 1, O = 16u]
[22 Jul 2021 Shift 2]

Answer: 4

Solution:

$$[\text{Glucose}] = \frac{C(\text{gm/l})}{M(\text{gm/mol})} = \frac{0.72}{180} = 4 \times 10^{-3}\text{M}$$

Question75

The number of significant figures in 0.00340 is _____ .
[25 Jul 2021 Shift 2]

Answer: 3

Solution:

Number of significant figures = 3

Question76

4g equimolar mixture of NaOH and Na_2CO_3 contains xg of NaOH and y g of Na_2CO_3 . The value of x is _____ g. (Nearest integer)
[20 Jul 2021 Shift 2]

Answer: 1

Solution:

Total mass = 4g

Now

NaOH : a mol $W_{\text{NaOH}} + W_{\text{Na}_2\text{CO}_3} = 4$

Na₂CO₃ : 'a' mol $\Rightarrow 40a + 106a = 4$

$$\Rightarrow a = \frac{4}{146} \text{ mol}$$

\Rightarrow therefore mass of NaOH is : $\frac{4}{146} \times 40\text{g}$

$$= 1.095 \approx 1$$

Question77

250mL of 0.5M NaOH was added to 500mL of 1M HCl. The number of unreacted HCl molecules in the solution after complete reaction is _____ $\times 10^{21}$.

(Nearest integer)

($N_A = 6.022 \times 10^{23}$)

[20 Jul 2021 Shift 1]

Answer: 226


Solution:

We know that no. of moles = $V_{\text{litre}} \times \text{Molarity}$ & No. of millimoles = $V_{\text{ml}} \times \text{Molarity}$

so millimoles of NaOH = $250 \times 0.5 = 125$

Millimoles of HCl = $500 \times 1 = 500$

Now reaction is



t = 0 125 500 0 0

t = t 0 375 125 125

so millimoles of HCl left = 375

Moles of HCl = 375×10^{-3}

No. of HCl molecules = $6.022 \times 10^{23} \times 375 \times 10^{-3}$

$$= 225.8 \times 10^{21}$$

$$\approx 226 \times 10^{21} = 226$$

Question78


Sodium oxide reacts with water to produce sodium hydroxide. 20.0g of

**sodium oxide is dissolved in 500 mL of water. Neglecting the change in volume, the concentration of the resulting NaOH solution is $\times 10^{-1}$ M.(Nearest integer)
 [Atomic mass: Na = 23.0, O = 16.0, H = 1.0]
 [31 Aug 2021 Shift 2]**

Answer: 13

Solution:

For the reaction,



$$\text{Moles of NaOH formed} \Rightarrow x = \frac{20}{62} \times 2$$

$$\text{Concentration of NaOH} = \frac{\text{Moles of NaOH}}{\text{Volume of solution(in litre)}}$$

$$= \frac{\frac{20}{62} \times 2}{\frac{500}{1000}} = 1.29\text{M} = 13 \times 10^{-1}\text{M}$$


Question79

**100g of propane is completely reacted with 1000g of oxygen. The mole fraction of carbon dioxide in the resulting mixture is $x \times 10^{-2}$. The value of x is (Nearest integer)
 [Atomic weight : H = 1.008, C = 12.00, O = 16.00]
 [27 Aug 2021 Shift 2]**

Answer: 19

Solution:

For the reaction,



1 mole of propane reacts completely with 5 moles of oxygen to form 3 moles of carbon dioxide and 4 moles of steam.

44 g of propane = 1 mole of propane


$$100\text{g of propane} = \frac{1}{44} \times 100 = 2.27 \text{ mol}$$

32g of O_2 = 1 mole of O_2

$$1000\text{g of } \text{O}_2 = \frac{1}{32} \times 1000 = 31.25 \text{ moles}$$

\therefore 2.27 moles of propane requires $5 \times 2.27 = 11.35$ moles of O_2

moles of CO_2 formed = $3 \times 2.7 = 6.681$ mol of CO_2



When reaction is completed 19.90 moles of O₂,
6.81 moles of CO₂ and 9.08 moles of steam are left in the flask.

$$\text{Mole fraction of CO}_2 = \frac{\text{Moles of CO}_2(\text{g})}{\text{Total number of moles}}$$

$$= \frac{6.81}{19.90 + 6.81 + 9.08} = 0.19$$

$$x \times 10^{-2} = 0.19$$

$$x = 19$$

Question80

A chloro compound A, (i) Forms aldehydes on ozonolysis followed by the hydrolysis.

(ii) When vaporised completely, 1.53g of A gives 448 mL of vapour at STP.

The number of carbon atoms in a molecule of compound A is
[26 Aug 2021 Shift 2]

Answer: 3

Solution:

Solution:

Given, 448 mL of A gives 1.53 g of vapours.

$$\therefore 22400 \text{ mL of A gives } = \frac{1.53}{448} \times 22400 = 76.50 \text{ g of A}$$

\therefore Molecular mass is 76.5

\therefore The possible compound is CH₃ – CH = CH – Cl

On ozonolysis followed by hydrolysis, it gives aldehyde as follows.

The compound (A) CH₃ – CH = CH – Cl (chloropropene) has 3 carbon atoms.

Question81

The number of atoms in 8g of sodium is $x \times 10^{23}$. The value of x is
(Nearest integer) [Given : N_A = 6.02 × 10²³ mol⁻¹ Atomic mass of

Na = 23.0u]

[1 Sep 2021 Shift 2]

Answer: 2

Solution:

Given, mass of Na = 8g

Molar mass of Na = 23g mol^{-1}

$$\frac{\text{Weight of sodium atom}}{\text{Molecular mass of sodium atom}} = \frac{\text{Number of atoms}}{\text{Avogadro's number}}$$

$$\frac{8\text{g}}{23\text{g}} = \frac{\text{Number of atoms}}{6.022 \times 10^{23}}$$

$$\text{Number of atoms} = \frac{8 \times 6.022}{23} \times 10^{23}$$

$$\text{Number of atoms} = 2.09 \times 10^{23}$$

$x \approx 2$

Hence, answer is 2.

Question82

4.5g of compound A(M W = 90) was used to make 250mL of its aqueous solution. The molarity of the solution is M is $x \times 10^{-1}$. The value of x is (Rounded off to the nearest integer) [2020]

Answer: 2

Solution:

Solution:

$$M = \frac{4.5 / 90}{250 / 1000} = 0.2 = 2 \times 10^{-1}$$

Question83

The molarity of HNO_3 in a sample which has density 1.4 g / mL and mass percentage of 63% is _____. (Molecular Weight of $\text{HNO}_3 = 63$) [NV, Jan. 09, 2020(I)]

Answer: 14

Solution:

Mass percent of $\text{HNO}_3 = 63$

Thus, 100g of nitric acid solution contains 63g of nitric acid by mass.

$$\text{No. of moles} = \frac{63\text{g}}{63\text{g mol}^{-1}} = 1$$

$$\text{Volume of 100g of nitric acid solution} = \frac{\text{Mass}}{\text{Density}} = \frac{100\text{g}}{1.4\text{g / mL}} = 71.4\text{mL}$$

$$\text{Molarity} = \frac{\text{No. of moles}}{\text{volume(mL)}} \times 1000 = \frac{1}{71.4} \times 1000 = 14\text{M}$$

Question84

Amongst the following statements, that which was not proposed by Dalton was:

[Jan. 07,2020 (I)]

Options:

- A. Chemical reactions involve reorganization of atoms. These are neither created nor destroyed in a chemical reaction.
- B. All the atoms of a given element have identical properties including identical mass. Atoms of different elements differ in mass.
- C. When gases combine or reproduced in a chemical reaction they do so in a simple ratio by volume provided all gases are at the same T & P.
- D. Matter consists of indivisible atoms.

Answer: C

Solution:

Solution:

Except (3) all postulates was given by the Dalton.

Question85

10.30mg of O₂ is dissolved into a liter of sea water of density 1.03g / mL.

The concentration of O₂ in ppm is _____.

[NV, Jan. 09, 2020 (II)]

Answer: 10

Solution:

Solution:

$$\text{ppm} = \frac{10.3 \times 10^{-3}}{1.03 \times 1000} \times 10^6 = 10$$

Question86



Ferrous sulphate heptahydrate is used to fortify foods with iron. The amount (in grams) of the salt required to achieve 10 ppm of iron in 100kg of wheat is _____ .

Atomic weight: Fe = 55.85; S = 32.00; O = 16.00

[NV, Jan. 08 , 2020 (I)]

Answer: 4.96

Solution:

Solution:

$$10\text{ppm} = \frac{\text{Mass of Fe (in g)}}{100 \times 1000} \times 10^6$$

$$\Rightarrow \text{Mass of Fe} = 1\text{g}$$

$$\text{Molar mass of FeSO}_4 \cdot 7\text{H}_2\text{O} = 278$$

$$56\text{g of iron present in 1mole of FeSO}_4 \cdot 7\text{H}_2\text{O}$$

$$1\text{g of Fe present } \frac{278}{56}\text{g in of salt} = 4.96\text{g}$$

Question87

NaClO₃ is used, even in spacecrafts, to produce O₂. The daily consumption of pure O₂ by a person is 492L at 1 atm, 300K . How much amount of NaClO₃, in grams, is required to produce O₂ for the daily consumption of a person at 1atm, 300K ? _____.




$$R = 0.082 \text{ Latm mol}^{-1}\text{K}^{-1}$$

[NV, Jan. 08,2020 (II)]

Answer: 2130

Solution:

Solution:



$$\text{Moles of NaClO}_3 = \text{Moles of O}_2$$

$$\text{Moles of O}_2 = \frac{PV}{RT} = \frac{1 \times 492}{0.082 \times 300} = 20\text{mol}$$

$$\text{Mass of NaClO}_3 = 20 \times 106.5 = 2130\text{g}$$

Question88



The ammonia (NH_3) released on quantitative reaction of 0.6g urea (NH_2CONH_2) with sodium hydroxide (NaOH) can be neutralized by:
[Jan. 07, 2020 (II)]

Options:

- A. 200mL of 0.4N HCl
- B. 200mL of 0.2N HCl
- C. 100mL of 0.2N HCl
- D. 100mL of 0.1N HCl

Answer: C

Solution:

Solution:

1 mol of urea = 2 mol of NH_3

60g of urea = 2mol of NH_3

0.6g of urea = $\frac{2}{60} \times 0.6\text{mol} = 0.02\text{mol}$ of NH_3

mol of NH_3 = mol of HCl

\therefore mol of HCl = 0.02mol

\Rightarrow Normality of HCl = 0.2N

Volume of HCl = 100mL

Question89

A solution of two components containing n_1 moles of the 1st component and n_2 moles of the 2nd component is prepared. M_1 and M_2 are the molecular weights of component 1 and 2 respectively. If d is the density of the solution in gmL^{-1} , C_2 is the molarity and x_2 is the mole fraction of the 2nd component, then C_2 can be expressed as:

[Sep. 06,2020(I)]

Options:

A. $C_2 = \frac{1000x_2}{M_1 + x_2(M_2 - M_1)}$

B. $C_2 = \frac{d x_2}{M_2 + x_2(M_2 - M_1)}$

C. $C_2 = \frac{1000d x_2}{M_1 + x_2(M_2 - M_1)}$

D. $C_2 = \frac{d x_1}{M_2 + x_2(M_2 - M_1)}$

Answer: C

Solution:



	1st component	2nd component
mole	n_1	n_2
m.w	M_1	M_2
mass	n_1M_1	n_2M_2

[∴ mass = mole × m.w.]

$$\text{Mass of solution} = n_1M_1 + n_2M_2$$

$$\text{Mole fraction of the 2nd component (x₂)} = \frac{n_2}{n_1 + n_2}$$

$$\Rightarrow n_1 = \frac{n_2(1 - x_2)}{x_2}$$

$$\text{Mass of solution} = n_1M_1 + n_2M_2$$

$$= \frac{n_2M_1(1 - x_2)}{x_2} + n_2M_2$$

$$= \frac{n_2}{x_2}[M_2x_2 - x_2M_1 + M_1]$$

$$\text{Volume of solution} = \frac{n_2[M_2x_2 - x_2M_1 + M_1]}{1000d x_2}$$

$$C_2 = \frac{1000n_2d x_2}{n_2[M_2x_2 - x_2M_1 + M_1]}$$

$$\Rightarrow C_2 = \frac{1000d x_2}{M_1 + x_2(M_2 - M_1)}$$

Question90

The average molar mass of chlorine is 35.5gmol^{-1} . The ratio of ^{35}Cl to ^{37}Cl in naturally occurring chlorine is close to:

[Sep. 06, 2020 (II)]

Options:

A. 4 : 1

B. 3 : 1


C. 2 : 1

D. 1 : 1

Answer: B

Solution:

Solution:



$$M_{\text{avg}} = 35 \times x + 37(1 - x) = 35.5$$

$$= 35x + 37(1 - x) = 35.5$$

$$\Rightarrow 2x = 1.5$$

$$x = \frac{3}{4}$$

So, ratio of $^{35}\text{Cl} : ^{37}\text{Cl} = \frac{3}{4} / \frac{1}{4} = 3 : 1$.

Question91

The ratio of the mass percentages of 'C&H' and 'C&O' of a saturated acyclic organic compound 'X' are 4 : 1 and 3 : 4 respectively. Then, the moles of oxygen gas required for complete combustion of two moles of organic compound 'X' is _____.

[NV, Sep. 02, 2020 (II)]

Answer: 5

Solution:

Solution:

Mass ratio of C : H is 4 : 1 \Rightarrow 12 : 3 and


C : O is 3 : 4 \Rightarrow 12 : 16

So,

	mass	mole	mole ratio
C	12	1	1
H	3	3	3
O	16	1	1

Empirical formula $\Rightarrow \text{CH}_3\text{O}$

As compound is saturated acyclic, so molecular formula is $\text{C}_2\text{H}_6\text{O}_2$



\therefore Number of moles of O_2 required to oxidise 2 moles of (X) = 5.

Question92

The minimum number of moles of O_2 required for complete combustion of 1 mole of propane and 2 moles of butane is _____.


[NV, Sep. 05, 2020(I)]

Answer: 18



Solution:






Complete combustion of hydrocarbons can be represented by the following reaction.



For propane combustion reaction is



Similarly, for butane is



Question93



The volume, in mL, of 0.02M $K_2Cr_2O_7$ solution required to react with 0.288g of ferrous oxalate in acidic medium is _____. (Molar mass of Fe = 56gmol⁻¹)

[NV, Sep. 05,2020(II)]

Answer: 50

Solution:

Solution:



$$V \times 0.02 \times 6 = \frac{0.288}{144} \times 3 \times 1000$$

$$V = 50\text{mL}$$


Question94

A 20.0mL solution containing 0.2 g impure H_2O_2 reacts completely with 0.316g of $KMnO_4$ in acid solution. The purity of H_2O_2 (in %) is _____. (mol. wt. of H_2O_2 = 34; mol. wt. of $KMnO_4$ = 158)

[NV, Sep. 04,2020(I)]

Answer: 85

Solution:



$$\text{Moles of KMnO}_4 = \frac{0.316}{158} = 2 \times 10^{-3}$$

$$\text{Equivalents of H}_2\text{O}_2 = \text{Equivalents of KMnO}_4$$

$$\text{Equivalents of KMnO}_4 = 2 \times 10^{-3} \times 5 = 0.01$$

$$\text{Moles of H}_2\text{O}_2 = \frac{0.01}{2} = 0.005$$

$$\text{Mass of pure H}_2\text{O}_2 = 0.005 \times 34 = 0.170\text{g}$$

$$\text{Percentage purity} = \frac{0.17}{0.2} \times 100 = 85\%$$

Question95

The mass of ammonia in grams produced when 2.8kg of dinitrogen quantitatively reacts with 1kg of dihydrogen is _____.


[NV, Sep. 04,2020(I)]

Answer: 3400

Solution:

Solution:

$$\text{Mole of N}_2 = \frac{2800}{28} = 100 \text{ and Mole of H}_2 = \frac{1000}{2} = 500$$



$$\text{Mass of NH}_3 \text{ formed} = 200 \times 17 = 3400\text{g}$$

Question96

A 100mL solution was made by adding 1.43g of $\text{Na}_2\text{CO}_3 \cdot x\text{H}_2\text{O}$. The normality of the solution is 0.1N.

The value of x is _____.

(The atomic mass of Na is 23g / mol)

[NV, Sep. 04, 2020 (II)]

Answer: 10

Solution:

Solution:

$$\text{Normality} = \frac{\text{No. of equivalents of solute}}{\text{Volume of solution (in L)}}$$



$$0.1 = \frac{1.43}{\frac{(106 + 18x)}{2} \times 0.1} \Rightarrow \frac{106 + 18x}{2} = 143$$

$$\Rightarrow 18x = 286 - 106 = 180 \Rightarrow x = 10.$$

Question97

The mole fraction of glucose ($C_6H_{12}O_6$) in an aqueous binary solution is 0.1. The mass percentage of water in it, to the nearest integer, is _____.

[NV, Sep. 03, 2020 (I)]

Answer: 47

Solution:

Solution:

Let total mole of solution = 1

So, mole of glucose = 0.1

Mole of H_2O = 0.9

$$\%(w/w) \text{ of } H_2O = \left[\frac{0.9 \times 18}{0.9 \times 18 + 0.1 \times 180} \right] \times 100$$

$$= 47.368 = 47.37.$$

Question98

6.023×10^{22} molecules are present in 10g of a substance 'x'. The molarity of a solution containing 5g of substance 'x' in 2 L solution is _____ $\times 10^{-3}$

[NV, Sep. 03, 2020 (II)]

Answer: 25

Solution:

$$\text{Number of mole of } x = \frac{6.022 \times 10^{22}}{6.022 \times 10^{23}} = \frac{10}{\text{Molar mass of } x}$$

So molar mass of x = 100g

$$\text{Molarity} = \frac{5}{100 \times 2} = 0.025M.$$



Question99


The volume (in mL) of 0.1N N aOH required to neutralise 10mL of 0.1N phosphinic acid is _____.

[NV, Sep. 03, 2020 (II)]

Answer: 10

Solution:

Phosphinic acid is hypophosphorous acid (H_3PO_2).



For neutrization,

$$(N_1 V_1)_{\text{add}} = (N_2 V_2)_{\text{base}}$$

$$0.1 \times 10 = 0.1 \times (V_{\text{mL}})_{\text{NaOH}}$$

$$V_{\text{NaOH}} = 10\text{mL}$$

Question100

8g of NaOH is dissolved in 18g of H_2O . Mole fraction of NaOH in solution and molality (in mol kg^{-1}) of the solution respectively are :
[Jan. 12, 2019 (II)]

Options:

A. 0.2,22.20

B. 0.2,11.11

C. 0.167,11.11

D. 0.167,22.20

Answer: C

Solution:

$$\text{No. of moles of } \text{H}_2\text{O}(n_1) = \frac{18}{18} = 1$$

$$\text{No. of moles of NaOH}(n_2) = \frac{8}{40} = \frac{1}{5}$$

$$\text{Mole fraction of NaOH} = \frac{n_2}{n_2 + n_1} = \frac{\frac{1}{5}}{\frac{1}{5} + 1} = 0.167$$

$$\text{Molality} = \frac{\text{No. of moles of solute}}{\text{Mass of solvent (kg)}} = \frac{1}{5} \times \frac{1000}{18} = 11.11\text{m}$$



Question101

The amount of sugar ($C_{12}H_{22}O_{11}$) required to prepare 2 L of its 0.1M aqueous solution is:
[Jan. 10 2019(II)]

Options:

- A. 136.8g
- B. 17.1g
- C. 68.4g
- D. 34.2g

Answer: C

Solution:

Solution:

As we know,

$$\text{Molarity} = \frac{\text{No. of moles of sugar}}{\text{Volume of solution (in L)}}$$

$$\Rightarrow 0.1 = \frac{\text{No. of moles of sugar}}{2L}$$

So, no. of moles of sugar = 0.2 mole

$$\therefore \text{Mass of sugar} = \text{No. of moles of sugar} \times \text{Molar mass of sugar} = 0.2 \times 342 = 68.4g$$

Question102

50mL of 0.5M oxalic acid is needed to neutralize 25mL of sodium hydroxide solution. The amount of NaOH in 50mL of the given sodium hydroxide solution is:
[Jan. 12,2019 (I)]

Options:

- A. 40g
- B. 10g
- C. 20g
- D. 80g
- E. None of Above

Answer: E

Solution:

Solution:

Oxalic acid Sodium hydroxide

$$N_1V_1 = N_2V_2$$

$$(2 \times M_1)V_1 = M_2V_2$$



$$2 \times 0.5 \times 50 = M_2 \times 25$$

$$M_2 = 2$$

$$\text{Molarity} = \frac{\text{No. of moles}}{\text{Vol. (L)}}$$

$$2 = \frac{\text{No. of moles}}{50 / 1000}$$

$$\text{No. of moles} = \frac{1}{10} = 0.1$$

$$\text{No. of moles} = \frac{W}{\text{Molar mass}}$$

$$W = 0.1 \times 40 = 4\text{g}$$

No option is correct.

Question103

A 10mg effervescent tablet containing sodium bicarbonate and oxalic acid releases 0.25mL of CO₂ at T = 298.15K and P = 1 bar. If molar volume of CO₂ is 25.0L under such condition, what is the percentage of sodium bicarbonate in each tablet?

[Molar mass of NaHCO₃ = 84gmol⁻¹]

[Jan. 11, 2019 (I)]

Options:

A. 0.84

B. 33.6


C. 16.8

D. 8.4

Answer: D

Solution:

Solution:



$$\text{Moles of CO}_2 \text{ evolved} = \frac{0.25}{25 \times 10^3} = 10^{-5}$$

$$\therefore \text{Moles of NaHCO}_3 = 10^{-5}$$

$$\therefore \text{Mass of NaHCO}_3 = 84 \times 10^{-5}\text{g}$$

$$= 0.84 \times 10^{-3}\text{g} = 0.84\text{mg}$$

$$\therefore \% \text{ by weight} = \times 100 = 8.4\%$$

Question104

25mL of the given HCl solution requires 30mL of 0.1M sodium carbonate solution. What is the volume of this HCl solution required to titrate 30mL of 0.2M aqueous NaOH solution?

[Jan.11,2019 (II)]

Options:



- A. 25mL
- B. 75mL
- C. 50mL
- D. 12.5mL

Answer: A

Solution:

Solution:

25mL of HCl solution requires 30mL of 0.1M Na_2CO_3 solution.

$$\therefore N_1 V_1 = N_2 V_2$$

$$\therefore 25 \times N_1 = 30 \times 0.2 \quad (0.1M \text{Na}_2\text{CO}_3 = 0.2N \text{Na}_2\text{CO}_3)$$

$$N_1 = \frac{6}{25} = 0.24N$$


Now, HCl solution is titrated with NaOH solution.

$$M_1 V_1 = M_2 V_2; 0.24N \text{HCl} = 0.24M \text{HCl}$$

$$\therefore V \times 0.24 \times 1 = 30 \times 0.2 \times 1 \Rightarrow V = 25\text{mL}$$

Question105

For the following reaction the mass of water produced from 445g of $\text{C}_{57}\text{H}_{110}\text{O}_6$ is:



[Jan. 9, 2019 (II)]

Options:


- A. 490g
- B. 445g
- C. 495g
- D. 890g

Answer: C

Solution:

Solution:

For the given reaction:



$$\text{Moles of } \text{C}_{57}\text{H}_{110}\text{O}_6 = \frac{445}{890} = 0.5$$

$$\text{Now, moles of water} = \frac{110}{2} \times 0.5 = 27.5$$

$$\therefore \text{mass} = 27.5 \times 18 = 495\text{g}$$

Question106



**The percentage composition of carbon by mole in methane is :
[April 8, 2019(II)]**

Options:

- A. 75%
- B. 80%
- C. 25%
- D. 20%

Answer: D

Solution:

Solution:

CH_4 has one atom of carbon among 5 atoms (1C + 4H)

$$\therefore \text{Mole \% of C} = \frac{1}{5} \times 100 = 20\%$$

Question107

**5 moles of AB_2 weigh $125 \times 10^{-3}\text{kg}$ and 10 moles of A_2B_2 weigh $300 \times 10^{-3}\text{kg}$. The molar mass of A(M_A) and molar mass of B(M_B) in kgmol^{-1} are:
[April 12, 2019(I)]**

Options:

- A. $M_A = 10 \times 10^{-3}$ and $M_B = 5 \times 10^{-3}$
- B. $M_A = 50 \times 10^{-3}$ and $M_B = 25 \times 10^3$
- C. $M_A = 25 \times 10^{-3}$ and $M_B = 50 \times 10^{-3}$
- D. $M_A = 5 \times 10^{-3}$ and $M_B = 10 \times 10^{-3}$

Answer: D

Solution:

Solution:

5mol AB_2 weighs 125g

$$\therefore \text{AB}_2 = 25\text{g} / \text{mol}$$

10mol A_2B_2 weighs 300g

$$\therefore \text{A}_2\text{B}_2 = 30\text{g} / \text{mol}$$

$$\therefore \text{Molar mass of A}(M_A) = 5\text{g} \text{ or } 5 \times 10^{-3}\text{kg}$$

$$\text{Molar mass of B}(M_B) = 10\text{g} \text{ or } 10 \times 10^{-3}\text{kg}$$



Question108

The minimum amount of $O_2(g)$ consumed per gram of reactant is for the reaction : (Given atomic mass: Fe = 56, O = 16, Mg = 24, P = 31, C = 12, H = 1)
[April 10, 2019 (II)]


Options:

- A. $4Fe(s) + 3O_2(g) \rightarrow 2Fe_2O_3(s)$
B. $P_4(s) + 5O_2(g) \rightarrow P_4O_{10}(s)$
C. $C_3H_8(g) + 5O_2(g) \rightarrow 3CO_2(g) + 4H_2O(l)$
D. $2Mg(s) + O_2(g) \rightarrow 2MgO(s)$


Answer: A

Solution:


Solution:




1g Fe requires $= \frac{3 \times 32}{4 \times 56} = 0.43g$ of oxygen



1g of P requires $= \frac{5 \times 32}{31 \times 4} = 1.3g$ of oxygen



1g of C_3H_8 requires $= \frac{5 \times 32}{44} = 3.6g$ of O_2



1 g Mg requires $= \frac{32}{2 \times 24}g = 0.66g$ of O_2

Question109

For a reaction, $N_2(g) + 3H_2(g) \rightarrow 2NH_3(g)$; identify dihydrogen (H_2) as a limiting reagent in the following reaction mixtures.
[April 9, 2019 (I)]

Options:

- A. 56g of N_2 + 10g of H_2
B. 35g of N_2 + 8g of H_2
C. 28g of N_2 + 6g of H_2
D. 14g of N_2 + 4g of H_2


Answer: A

Solution:



Solution:

According to the stoichiometry of balanced equation, 28gN_2 reacts with 6gH_2



$\frac{1 \text{ mole } 28 \text{ g}}{3 \text{ mole } 6 \text{ g}}$

\therefore For 56g of N_2 , 12g of H_2 is required.

Question110

0.27g of a long chain fatty acid was dissolved in 100cm^3 of hexane. 10mL of this solution was added dropwise to the surface of water in a round watch glass. Hexane evaporates and a monolayer is formed. The distance from edge to centre of the watch glass is 10cm. What is the height of the monolayer?

[Density of fatty acid = 0.9gcm^{-3} ; $\pi = 3$]

[April 8, 2019 (II)]

Options:

A.

10^{-8} m

B.

10^{-6} m

C.

10^{-4} m

D.

10^{-2} m

Answer: B

Solution:**Solution:**

Given: 0.27g is present in 100cm^3 of hexane

\therefore 10mL of aqueous solution contains only 0.027 g acid.

$$\text{Volume of } 0.027\text{g acid} = \frac{0.027}{0.9} \text{ mL}$$

$$\therefore \pi r^2 h = \frac{0.027}{0.9} \quad (\text{given } r = 10\text{cm}, \pi = 3)$$

$$\therefore h = 10^{-4} \text{ cm} = 10^{-6} \text{ m}$$

Question111

An unknown chlorohydrocarbon has 3.55% of chlorine. If each molecule of the hydrocarbon has one chlorine atom only, chlorine atoms present in 1g of chlorohydrocarbon are:

(Atomic wt. of Cl = 35.5u ; Avogadro constant = $6.023 \times 10^{23} \text{ mol}^{-1}$)



[Online April 16,2018]

Options:

- A. 6.023×10^9
- B. 6.023×10^{23}
- C. 6.023×10^{21}
- D. 6.023×10^{20}

Answer: D

Solution:

Solution:

Given percentage of chlorine in an hydrocarbon = 3.55%

i.e.,

100g of chlorohydrocarbon has 3.55g of chlorine.

1 g of chlorohydrocarbon will have $\frac{3.55}{100} = 0.0355$ g of chlorine.

Atomic wt. of Cl = 35.5g / mol

Number of moles of Cl = $\frac{0.0355\text{g}}{35.5\text{g / mol}} = 0.001\text{mol}$

Number of atoms of Cl = $0.001\text{mol} \times 6.023 \times 10^{23}\text{mol}^{-1}$
= 6.023×10^{20}

Question112

The ratio of mass percent of C and H of an organic compound ($C_xH_yO_z$) is 6 : 1. If one molecule of the above compound ($C_xH_yO_2$) contains half as much oxygen as required to burn one molecule of compound C_xH_y completely to CO_2 and H_2O . The empirical formula of compound $C_xH_yO_z$ is:

[2018]

Options:

- A. $C_3H_6O_3$
- B. C_2H_4O
- C. $C_3H_4O_2$
- D. $C_2H_4O_3$


Answer: D

Solution:

Solution:

So, $x = 1, y = 2$

Equation for combustion of C_xH_y



$$\text{Oxygen atoms required} = 2\left(x + \frac{y}{4}\right)$$



As mentioned,

$$2\left(x + \frac{y}{4}\right) = 2z; \left(x + \frac{y}{4}\right) = z$$

Now putting the values of x and y

$$\Rightarrow \left(1 + \frac{2}{4}\right) = z \Rightarrow z = 1.5$$

∴ Molecule ($C_xH_yO_2$) can be written as



Question 113

A sample of $NaClO_3$ is converted by heat to $NaCl$ with a loss of 0.16 g of oxygen. The residue is dissolved in water and precipitated as $AgCl$. The mass of $AgCl$ (in g) obtained will be: (Given: Molar mass of $AgCl = 143.5 \text{ gmol}^{-1}$)
[Online April 15, 2018 (I)]

Options:

- A. 0.35
- B. 0.54
- C. 0.41
- D. 0.48


Answer: D

Solution:

Solution:

No. of moles of oxygen in 0.16g of oxygen molecule

$$= \frac{0.16\text{g}}{32\text{g/mol}} = 0.005\text{mol}$$



According to the reaction,

3 moles of $O_2 = 2$ moles of $NaCl = 2$ moles of $AgCl$

Molar mass of $AgCl = 143.5 \text{ g/mol}$

$$0.005 \text{ moles of } O_2 \text{ will ppt.} = 0.005 \times \frac{2}{3} \text{ moles } AgCl$$

$$= 0.0033 \text{ moles of } AgCl$$

∴ Mass of $AgCl$ (in g) obtained will be

$$= 143.5 \text{ g/mol} \times 0.0033 \text{ moles} = 0.48 \text{g.}$$

Question 114

Excess of $NaOH$ (aq) was added to 100mL of $FeCl_3$ (aq) resulting into 2.14g of $Fe(OH)_3$. The molarity of $FeCl_3$ (aq) is:



(Given molar mass of Fe = 56g mol^{-1} and molar mass of Cl = 35.5g mol^{-1})

[Online April 8, 2017]


Options:

- A. 0.2M
- B. 0.3M
- C. 0.6M
- D. 1.8M

Answer: A

Solution:

Solution:



$$\text{Moles of Fe}(\text{OH})_3 = \frac{\text{weight in g}}{\text{Mol. mass of Fe}(\text{OH})_3}$$

$$= \frac{2.14\text{g}}{107\text{g/mol}} = 0.02\text{mol}$$

1.0mole of Fe(OH)₃ is obtained from = 1.0mole of FeCl₃

$$0.02 \text{ mole of Fe}(\text{OH})_3 \text{ will be obtained from } = 0.02 \text{ mole FeCl}_3 \text{ Molarity} = \frac{\text{No. of moles}}{\text{Volume in L}} = \frac{0.02\text{mole}}{0.1\text{L}} = 0.2\text{M}$$

Question 115

The most abundant elements by mass in the body of a healthy human adult are: [2017] Oxygen (61.4%); Carbon (22.9%), Hydrogen (10.0%); and Nitrogen (2.6%). The weight which a 75kg person would gain if all ¹H atoms are replaced by ²H atoms is [2017]

Options:

- A. 15kg
- B. 37.5kg
- C. 7.5kg
- D. 10kg

Answer: C

Solution:

Solution:

Percentage (by mass) of elements given in the body of a healthy human adult is :

Oxygen = 61.4%, Carbon = 22.9%

Hydrogen = 10.0% and Nitrogen = 2.6%

∴ Total weight of person = 75kg



\therefore Mass due to ^1H is $= 75 \times \frac{10}{100} = 7.5\text{kg}$

If ^1H atoms are replaced by ^2H atoms.

Mass gain by person would be $= 7.5\text{kg}$

Question116

What quantity (in mL) of a 45% acid solution of a monoprotic strong acid must be mixed with a 20% solution of the same acid to produce 800mL of a 29.875% acid solution?

[Online April 9,2017]

Options:

A. 320

B. 325

C. 316

D. 330

Answer: C

Solution:

Solution:

$$\frac{V \times 45}{100} + \frac{(800 - V) \times 20}{100} = \frac{800 \times 29.875}{100}$$
$$\Rightarrow \frac{9V}{20} + 160 - \frac{V}{5} = 239 \Rightarrow V = 316$$

Question117

5L of an alkane requires 25L of oxygen for its complete combustion. If all volumes are measured at constant temperature and pressure, the alkane is:

[Online April 9, 2016]

Options:

A. Isobutane

B. Ethane

C. Butane


D. Propane

Answer: D

Solution:



Since the compound undergoing combustion is an alkane. Hence the combustion reaction can be written as



Since volumes are measured at constant T & P, hence according to Avogadro's law

Volume \propto mole

1 Lalkane requires $\frac{3n+1}{2}$ Lof O_2

5 Lalkane requires 25L of O_2

$$\frac{1}{5} = \frac{\frac{3n+1}{2}}{25}$$





$$\therefore n = 3$$

Hence alkane is propane (C_3H_8)

Question118

At 300K and 1atm, 15mL of a gaseous hydrocarbon requires 375mL air containing 20% O_2 by volume for complete combustion. After combustion the gases occupy 330mL. Assuming that the water formed is in liquid form and the volumes were measured at the same temperature and pressure, the formula of the hydrocarbon is: [2016]

Options:




E. None of above

Answer: E

Solution:

Solution:



$$\text{Volume of } O_2 \text{ used} = \frac{20}{100} \times 375 = 75\text{mL}$$

$$\text{Volume of air} = 375 - 75 = 300\text{mL}$$

$$\text{Total volume of gases after combustion} \\ = \text{vol. of } CO_2 + \text{vol. of air} = 330\text{mL}$$

$$\text{Volume of } CO_2 = 330 - 300 = 30\text{mL}$$

$$15\text{mL } C_xH_y \text{ gives} = 30\text{mL } CO_2$$

$$1\text{mL } C_xH_y \text{ gives} = \frac{30}{15} = 2\text{mL } CO_2$$

At constant T and P; Volume \propto mole

$$\therefore 1\text{mol } C_xH_y = 2\text{mol } CO_2$$

$$x = 2$$

$$\left(\frac{4x+y}{4} \right) = \frac{75}{15}$$

$$4x + y = 20$$

$$y = 20 - 4 \times 2 = 12$$



Hence, formula of the hydrocarbon is C_2H_{12} .

Question119

The amount of arsenic pentasulphide that can be obtained when 35.5g arsenic acid is treated with excess H_2S in the presence of conc. HCl

(assuming 100% conversion) is:

[Online April 9, 2016]


Options:

- A. 0.25mol
- B. 0.50mol
- C. 0.333mol
- D. 0.125mol

Answer: D

Solution:

Solution:



\therefore The molar mass of H_3AsO_4 is $3 \times 1 + 79 + 4 \times 16 = 142g / mol$

\therefore Number of moles of $H_3AsO_4 = \frac{35.5}{142} = 0.25mol$

\therefore Number of moles of $As_2S_5 = \frac{0.25}{2} = 0.125mol$.

Question120

The volume of 0.1N dibasic acid sufficient to neutralize 1 g of a base that furnishes 0.04 mole of OH^- in aqueous solution is :

[Online April 10,2016]

Options:

- A. 400mL
- B. 600mL
- C. 200mL
- D. 800mL

Answer: A

Solution:



Applying law of equivalence
Equivalence of acid = Equivalence of base
Equivalent of acid = Normality \times volume = $0.1 \times V$
Another formula of equivalence = n factor \times number of moles
 \therefore Equivalent of base = n factor of $\text{OH}^- \times$ moles of OH^-
= 1×0.04
 $0.1 \times V = 1 \times 0.04$
 $V = 0.4\text{L} = 0.4 \times 1000 = 400\text{mL}$.

Question121

The molecular formula of a commercial resin used for exchanging ions in water softening is $\text{C}_8\text{H}_7\text{SO}_3 - \text{Na}^+$ (Mol. wt. 206). What would be the maximum uptake of Ca^{2+} ions by the resin when expressed in mole per gram resin?

[2015]

Options:

A. $\frac{2}{309}$

B. $\frac{1}{412}$

C. $\frac{1}{103}$

D. $\frac{1}{206}$

Answer: B

Solution:

Solution:

2 mole of water softner require 1 mole of Ca^{2+} ion

So, 1 mole of water softner require $\frac{1}{2}$ mole of Ca^{2+} ion




Thus, $\frac{1}{2 \times 206} = \frac{1}{412}$ mol / g will be maximum uptake.

Question122

Choose the incorrect formula out of the four compounds for an element X below:

[Online April 11, 2015]

Options:



D. XPO_4

Answer: C

Solution:

Solution:

Mass of substance = 250mg = 0.250g

Mass of AgBr = 141mg = 0.141g

1 mole of AgBr = 1g atom of Br

188g of AgBr = 80g of Br

\therefore fof AgBr contain bromine = 80g

0.141g of AgBr contain bromine

$$= \frac{80}{188} \times 0.141 = 0.06g$$

0.06g of bromine is present in 0.250g of organic compound

$$\therefore \% \text{ of bromine} = \frac{0.06}{0.250} \times 100 = 24\%$$

Question123

3 g of activated charcoal was added to 50mL of acetic acid solution (0.06N) in a flask. After an hour it was filtered and the strength of the filtrate was found to be 0.042N . The amount of acetic acid adsorbed (per gram of charcoal) is: [2015]

Options:

A. 42mg

B. 54mg

C. 18mg

D. 36mg

Answer: C

Solution:

Solution:

Let the weight of acetic acid initially be w_1 in 50mL of 0.060N solution.

$$N = \frac{w_1 \times 1000}{M \cdot wt. \times 50} \quad (\text{Normality} = 0.06N)$$

$$0.06 = \frac{w_1 \times 1000}{60 \times 50}$$

$$\Rightarrow w_1 = \frac{0.06 \times 60 \times 50}{1000} = 0.18g = 180mg$$

After an hour, the strength of acetic acid = 0.042N

so, let the weight of acetic acid be w_2

$$N = \frac{w_2 \times 1000}{60 \times 50}$$

$$0.042 = \frac{W_2 \times 1000}{3000}$$

$$\Rightarrow w_2 = 0.126g = 126mg$$

So amount of acetic acid adsorbed per 3g

$$= 180 - 126mg = 54mg$$



∴ amount of acetic acid absorbed per g = $54 / 3 = 18\text{mg}$

Question124





A sample of a hydrate of barium chloride weighing 61g was heated until all the water of hydration is removed. The dried sample weighed 52g.

The formula of the hydrated salt is : (atomic mass,

Ba = 137amu, Cl = 35.5 amu)

[Online April 10, 2015]

Options:



Answer: D

Solution:

Solution:

Weight of hydrated $\text{BaCl}_2 = 61\text{g}$

Weight of anhydrous $\text{BaCl}_2 = 52\text{g}$

Loss in mass = 9g

Assuming $\text{BaCl}_2 \cdot x\text{H}_2\text{O}$ as hydrate

Mass of $\text{H}_2\text{O} = 9\text{g}$

$$\text{Moles of } \text{H}_2\text{O} = \frac{9}{18} = 0.5\text{mol}$$

Gross molecular wt. of $\text{BaCl}_2 = 208\text{g} / \text{mol}$.

$$\% \text{ of } \text{H}_2\text{O} \text{ in this hydrated } \text{BaCl}_2 = \frac{9}{61} \times 100 = 14.75\%$$

$$\% \text{ of } \text{H}_2\text{O} \text{ in } \text{BaCl}_2 \cdot x\text{H}_2\text{O} = \frac{18x}{208 + 18x} \times 100$$

$$\text{Thus, } \frac{18x}{208 + 18x} \times 100 = 14.75$$

On solving $x = 2$

Hence, the formula of hydrated salt is $\text{BaCl}_2 \cdot 2\text{H}_2\text{O}$

Question125

$\text{A} + 2\text{B} + 3\text{C} \rightleftharpoons \text{AB}_2\text{C}_3$ Reaction of 6.0g of A, 6.0×10^{23} atoms of B, and 0.036 mol of C yields 4.8g of compound AB_2C_3 . If the atomic mass of A

and C are 60 and 80amu, respectively, the atomic mass of B is (

Avogadro no. = 6×10^{23}) :

[Online April 11, 2015]

Options:



B. 60amu


C. 70amu

D. 40amu

Answer: A

Solution:

Solution:



$$\text{No. of moles of A} = \frac{6.0\text{g}}{60\text{g/mol}} = 0.1\text{mol}$$

$$\text{No. of moles of B} = \frac{6.00 \times 10^{23}}{6.000 \times 10^{23}} = 1\text{mol}$$

$$\text{No. of moles of C} = 0.036$$

Therefore, C is the limiting reagent,

The number of moles of product formed

$$= \frac{0.036}{3} = 0.012\text{mol}$$

The expression for the molar mass is Molar mass

$$= \frac{\text{Given mass}}{\text{mole of product}}$$

$$60 + (2 \times x) + (3 \times 80) = \frac{4.8}{0.012}$$

$$x = 50\text{amu}$$

Hence, atomic mass of B is 50 amu

Question126

In Carius method of estimation of halogens, 250mg of an organic compound gave 141mg of AgBr. The percentage of bromine in the compound is :

(at. mass Ag = 108; Br = 80)

[2015]

Options:

A. 48

B. 60

C. 24

D. 36

Answer: C

Solution:

Solution:

$$\text{Mass of substance} = 250\text{mg} = 0.250\text{g}$$

$$\text{Mass of AgBr} = 141\text{mg} = 0.141\text{g}$$

$$1 \text{ mole of AgBr} = 1\text{g atom of Br}$$

$$188\text{g of AgBr} = 80\text{g of Br}$$

$$\therefore 188\text{g of AgBr contain bromine} = 80\text{g}$$

$$0.141 \text{ g of AgBr contain bromine}$$



$$= \frac{80}{188} \times 0.141 = 0.06\text{g}$$

0.06g of bromine is present in 0.250g of organic compound

$$\therefore \% \text{ of bromine} = \frac{0.06}{0.250} \times 100 = 24\%$$

Question127

Dissolving 120g of a compound of (mol. wt. 60) in 1000g of water gave a solution of density 1.12g / mL. The molarity of the solution is: [Online April 9,2014]

Options:

- A. 1.00M
- B. 2.00M
- C. 2.50M
- D. 4.00M

Answer: B

Solution:

Solution:

Given

mass of solute (w) = 120g

mass of solvent (w) = 1000g

Mol. mass of solute = 60g

density of solution = 1.12g / mL

From the given data,

Mass of solution = 1000 + 120 = 1120g

$$\therefore d = \frac{\text{Mol. mass}}{V} \text{ or } V = \frac{\text{Mol. mass}}{d}$$

$$\text{Volume of solution } V = \frac{1120}{1.12} = 1000\text{mL or } = 1 \text{ litre}$$

$$\text{Now molarity (M)} = \frac{w}{\text{Mol. mass} \times V(\text{L})} = \frac{120}{60 \times 1} = 2\text{M}$$

Question128

The amount of oxygen in 3.6 moles of water is: [Online April 9, 2014]

Options:

- A. 115.2g
- B. 57.6g
- C. 28.8g
- D. 18.4g

Answer: B



Solution:

1 mole of water contains = 16 g of oxygen
∴ 3.6 mole of water contains
= $16 \times 3.6 = 57.6\text{g}$

Question 129

A gaseous compound of nitrogen and hydrogen contains 12.5% (by mass) of hydrogen. The density of the compound relative to hydrogen is 16. The molecular formula of the compound is:

[Online April 11, 2014]

Options:

A. NH_2

B. N_3H





C. NH_3

D. N_2H_4

E. None of above.

Answer: D

Solution:



Since Cl^{2-} does not exist. So, X_2Cl_3 is incorrect.

The correct formula should be XCl_3

Question 130

A gaseous compound of nitrogen and hydrogen contains 12.5% (by mass) of hydrogen. The density of the compound relative to hydrogen is 16. The molecular formula of the compound is:


[Online April 11, 2014]

Options:

A. NH_2

B. N_3H



**Answer: D****Solution:****Solution:**


In an unknown compounds containing N and H given % of H = 12.5%

 \therefore % of N = $100 - 12.5 = 87.5\%$

Element	Percentage	Atomic ratio	Simple ratio
H	12.5%	$\frac{12.5}{1} = 12.5$	$\frac{12.5}{6.25} = 2$
N	87.5	$\frac{87.5}{14} = 6.25$	$\frac{6.25}{6.25} = 1$

Empirical formula = NH_2 Mol. wt = $2 \times$ vapour density = $16 \times 2 = 32$.Molecular formula = $n \times$ empirical formula mass

$$n = \frac{32}{16} = 2$$

 \therefore Molecular formula of the compound will be = $(\text{NH}_2)_2$ 

Question131


The amount of BaSO_4 formed upon mixing 100mL of 20.8% BaCl_2 solution with 50mL of 9.8% H_2SO_4 solution will be:**(Ba = 137, Cl = 35.5, S = 32, H = 1 and O = 16)****[Online April 12, 2014]****Options:**

A. 23.3g

B. 11.65g

C. 30.6g

D. 33.2g

Answer: B**Solution:****Solution:**

$$\text{Mass of BaCl}_2 \text{ in solution} = 100 \times \frac{20.8}{100} = 20.8\text{g}$$

$$\text{Mass of H}_2\text{SO}_4 \text{ in solution} = 50 \times \frac{9.8}{100} \times 4.9 = 4.9\text{g}$$

98g of H_2SO_4 reacts with 208g BaCl_2

$4.9\text{gH}_2\text{SO}_4$ will react with $\frac{208}{98} \times 4.9 = 10.4\text{gBaCl}_2$

H_2SO_4 reacts as a limiting reagent because BaCl_2 is given in excess

$98\text{gH}_2\text{SO}_4$ produces 233gBaSO_4

$4.9\text{gH}_2\text{SO}_4$ will produce $\frac{233}{98} \times 4.9 = 11.65\text{gBaSO}_4$

Question132

The density of 3M solution of sodium chloride is 1.252g mL^{-1} . The molality of the solution will be:

(molar mass, $\text{NaCl} = 58.5\text{gmol}^{-1}$)

[Online April 22, 2013]

Options:

- A. 260m
- B. 2.18m
- C. 2.79m
- D. 3.00m

Answer: C

Solution:

Solution:

The relation between molarity (M) and molality (m) is $d = M \left(\frac{1}{m} + \frac{M_2}{1000} \right)$, $M_2 =$ Mol. mass of solute

On putting value $1.252 = 3 \left(\frac{1}{m} + \frac{58.5}{1000} \right)$

On solving $m = 2.79$

Question133

Number of atoms in the following samples of substances is the largest in:

[Online April 23, 2013]

Options:

- A. 4.0g of hydrogen
- B. 71.0g of chlorine
- C. 127.0g of iodine
- D. 48.0g of magnesium

Answer: A

Solution:



4g of hydrogen = 4 mole of hydroge

$$n = 4 \times 6.023 \times 10^{23} \text{ atoms}$$

71.0g of chlorine = $\frac{71.0}{71.0} = 1$ moles of chlorine

$$= 6.023 \times 10^{23} \text{ atoms}$$

127g of iodine = $\frac{127}{254}$ mol

$$= 6.023 \times 10^{23} \times \frac{1}{2} \text{ atoms}$$

48.0g of magnesium = $\frac{48.0}{24.0}$ mol

$$= 2 \times 6.023 \times 10^{23} \text{ atoms}$$

\therefore 4.0gH₂ has largest number of atoms.

Question134

10mL of 2(M)NaOH solution is added to 200mL of 0.5 (M) of NaOH solution. What is the final concentration ?

[Online April 25, 2013]

Options:

A. 0.57(M)

B. 5.7(M)

C. 11.4(M)

D. 1.14(M)

Answer: A

Solution:

Solution:

From molarity equation

$$M_1V_1 + M_2V_2 = MV_{(\text{total})}$$

$$M = \frac{M_1V_1 + M_2V_2}{\text{Total}} = \frac{2 \times 10 + 0.5 \times 200}{210}$$

$$M = \frac{120}{210} = 0.57M$$

Question135

6 litres of an alkene require 27 litres of oxygen at constant temperature and pressure for complete combustion. The alkene is:

[Online April 25, 2013]

Options:

A. Ethene

B. Propene



C. 1 -Butene

D. 2 -Butene



Answer: B

Solution:

Solution:

General combustion reaction for hydrocarbons is $C_xH_y + \left(x + \frac{y}{4}\right)O_2 \rightarrow xCO_2 + \frac{y}{2}H_2O$

For alkane, $x = n$ and $y = 2n$



\therefore 1 mole alkene reacts with $\frac{3n}{2}$ mole of O_2

moles \propto volume (at constant temp & spress.)

1 L alkene requires $\frac{3n}{2}$ L of O_2

6L alkene requires 27L of O_2

$$\frac{1}{6} = \frac{\frac{3n}{2}}{27}$$

$$3n = \frac{54}{6}$$

$$n = 3^6$$

Hence alkene is propene (C_3H_6)

Question136

A gaseous hydrocarbon gives upon combustion 0.72g of water and 3.08g. of CO_2 . The empirical formula of the hydrocarbon is :

[2013]

Options:

A. C_2H_4

B. C_3H_4

C. C_6H_5

D. C_7H_8

Answer: D

Solution:

Solution:

\therefore 18g, H_2O contains = 2gH

\therefore 0.72g H_2O contains

$$= \frac{2}{18} \times 0.72g = 0.08gH$$

\therefore 44g CO_2 contains = 12gC

$$\therefore 3.08gCO_2 \text{ contains } = \frac{12}{44} \times 3.08 = 0.84gC$$



$$\therefore \text{C} : \text{H} = \frac{0.84}{12} : \frac{0.08}{1}; 0.07 : 0.08 = 7 : 8$$

\therefore Empirical formula = C_7H_8

Question137

**1 gram of a carbonate (M_2CO_3) on treatment with excess HCl produces 0.01186mol e of CO_2 . The molar mass of M_2CO_3 in gmol^{-1} is:
[2013]**

Options:


- A. 1186
- B. 84.3
- C. 118.6
- D. 11.86

Answer: B

Solution:

Solution:

Given chemical eqⁿ



0.01186mol From the above chemical eqⁿ.

$$n\text{M}_2\text{CO}_3 = n\text{CO}_2$$

$$\frac{1}{\text{Molar mass of } \text{M}_2\text{CO}_3} = 0.01186$$

$$\therefore \text{Molar mass of } \text{M}_2\text{CO}_3 = \frac{1}{0.01186}$$

$$\text{Molar mass} = 84.3\text{g / mol}$$

Question138

**The concentrated sulphuric acid that is peddled commercial is 95% H_2SO_4 by weight. If the density of this commercial acid is 1.834 gcm^{-3} , the molarity of this solution is
[Online May 7, 2012]**

Options:

- A. 17.8M
- B. 12.0M
- C. 10.5M
- D. 15.7M

Answer: A



Solution:

95%H₂SO₄ by weight means 100gH₂SO₄ solution contains 95gH₂SO₄ by mass.

Molar mass of H₂SO₄ = 98gmol⁻¹

Moles in 95gH₂SO₄ = $\frac{95}{98} = 0.969\text{mol}$

Volume of 100gH₂SO₄ solution

$$= \frac{\text{mass}}{\text{density}} = \frac{100\text{g}}{1.834\text{gcm}^{-3}}$$

$$= 54.52\text{cm}^3 = 54.52 \times 10^{-3}\text{L}$$

$$\text{Molarity} = \frac{\text{Moles of solute}}{\text{Volume of solution in L}}$$

$$= \frac{0.969}{54.52 \times 10^{-3}} = 17.8\text{M}$$

Question139

The ratio of number of oxygen atoms (O) in 16.0 g ozone (O₃), 28.0g carbon monoxide (CO) and 16.0 oxygen (O₂) is (Atomic mass: C = 12, O = 16 and Avogadro's constant N_A = 6.0 × 10²³ mol⁻¹)
[Online May 7, 2012]

Options:

A. 3 : 1 : 2

B. 1 : 1 : 2

C. 3 : 1 : 1

D. 1 : 1 : 1

Answer: D

Solution:

Solution:

O₃ molecular weight = 16 + 16 + 16 = 48g / mol

It means weight of 1mol of O₃ is 48g and in 1mol of O₃ we have 3 atoms of Oxygen

In 48g of O₃, number of atoms of oxygen = 3

so in 16g of O₃, number of atoms of oxygen = $(3 / 48) \times 16 = 1$

CO molecular weight = 12 + 16 = 28g / mol

It means weight of 1 mol of CO is 28g and in 1 mol of CO we have 1 atom of Oxygen

so in 28g of CO, number of atoms of oxygen = 1

O₂ molecular weight = 12 + 16 = 32g / mol

It means weight of 1 mol of O₂ is 32g and in 1mol of O₂ we have 2 atoms of Oxygen

In 32g of O₂, number of atoms of oxygen = 2

so in 16g of O₂, number of atoms of oxygen = $(2 / 32) \times 16 = 1$

So answer is 1:1:1

Question140



The ppm level of F^- in a 500g sample of a tooth paste containing $0.2gF^-$ is
[Online May 12, 2012]

Options:

- A. 400
- B. 1000
- C. 250
- D. 200

Answer: A

Solution:

$$\begin{aligned} \text{ppm} &= \frac{\text{mass of solute (g)}}{\text{mass of solution (g)}} \times 10^6 \\ &= \frac{0.2}{500} \times 10^6 = 400 \text{ ppm} \end{aligned}$$

Question141

5g of benzene on nitration gave 6.6g of nitrobenzene. The theoretical yield of the nitrobenzene will be
[Online May 12, 2012]


Options:

- A. 4.5g
- B. 5.6g
- C. 8, 09g
- D. 6.6g

Answer: C

Solution:

Solution:



Now since 78g of benzene on nitration give = 123g nitrobenzene
hence 5g of benzene on nitration give

$$= \frac{123}{78} \times 5 = 7.88g$$


The nearest answer is (c) i.e. theoretical yield = 7.88g

Question142



A transition metal M forms a volatile chloride which has a vapour density of 94.8. If it contains 74.75% of chlorine the formula of the metal chloride will be
[Online May 26, 2012]

Options:



Answer: C

Solution:

Solution:

74.75% of chlorine means 74.75g chlorine is present in 100g of metal chloride.

$$\begin{aligned}\text{Weight of metal} &= 100 - 74.75 \\ &= 25.25\text{g}\end{aligned}$$

$$\text{Equivalent weight} = \frac{\text{weight of metal}}{\text{weight of chlorine}} \times 35.5$$

$$= \frac{25.25}{74.75} \times 35.5 = 12$$

Valency of metal

$$= \frac{2 \times V \cdot D}{\text{Equivalent wt. of metal} + 35.5}$$

$$= \frac{2 \times 94.8}{12 + 35.5} = 4$$

∴ Formula of metal chloride is MCl_4

Alternate method:

Mol. wt = 2 × vapour density

$$= 2 \times 94.8 = 189.6\text{g}$$

Since 74.75% is chlorine therefore,

189.6 metal chloride contains

$$= \frac{74.75}{100} \times 189.6 = 141.72\text{g chloride}$$

$$\text{Number of atoms of chloride} = \frac{141.72}{35.5} = 3.99 \approx 4$$

Hence, formula of metal chloride is MCl_4

Question143

An aqueous solution of oxalic acid dihydrate contains its 6.3 g in 250mL. The volume of 0.1N NaOH required to completely neutralize 10mL of this solution
[Online May 12, 2012]

Options:

A. 4mL

B. 20mL

C. 2mL

D. 40mL

Answer: D

Solution:

Normality of oxalic acid solution

$$= \frac{6.3 \times 1000}{63 \times 250} = 0.4N$$

Now from

$$N_1 V_1 = N_2 V_2$$

$$0.4 \times 10 = 0.1 \times V_2$$

$$V_2 = 40\text{mL}$$

Question144

The molality of a urea solution in which 0.0100g of urea, $[(\text{NH}_2)_2\text{CO}]$ is added to 0.3000d m^3 of water at STP is:
[2011RS]

Options:

A. $5.55 \times 10^{-4}\text{m}$

B. 33.3 m

C. $3.33 \times 10^{-2}\text{m}$

D. 0.555m

Answer: A

Solution:

Solution:

Molality = Moles of solute/ Mass of solvent in kg

$$\begin{aligned} \text{Molality} &= \frac{0.01 / 60}{0.3} = \frac{0.01}{60 \times 0.3} \\ &= 5.55 \times 10^{-4}\text{m} \end{aligned}$$

Question145

The density (in gmL^{-1}) of a 3.60M sulphuric acid solution that is 29% H_2SO_4 (molar mass = 98gmol^{-1}) by mass will be
[2007]

Options:

A. 1.45

B. 1.64



C. 1.88

D. 1.22

Answer: D

Solution:

Solution:

Since molarity of solution is 3.60M . It means 3.6 moles of H_2SO_4 is present in its 1 litre solution.

Mass of 3.6 moles of H_2SO_4

= Moles \times Molecular mass

= $3.6 \times 98g = 352.8g$

\therefore 1000mL solution has 352.8g of H_2SO_4

Given that 29g of H_2SO_4 is present in

= 100g of solution

\therefore 352.8g of H_2SO_4 is present in

= $\frac{100}{29} \times 352.8g$ of solution


= 1216g of solution

Density = $\frac{\text{Mass}}{\text{Volume}} = \frac{1216}{1000}$

= 1.216g / mL = 1.22g / mL

Question 146

In the reaction, [2007]



[2007]

Options:

A. 11.2L $H_2(g)$ at STP is produced for every mole of $HCl(aq)$ consumed

B. 6L $HCl(aq)$ is consumed for every 3L of $H_2(g)$ produced


C. 33.6L $H_2(g)$ is produced regardless of temperature and pressure for every mole of Al that reacts

D. 67.2L $H_2(g)$ at STP is produced for every mole of Al that reacts.

Answer: A

Solution:

Solution:



\therefore 6 moles of HCl produces = 3 moles of H_2

= $3 \times 22.4L$ of H_2 at S.T.P

\therefore 1 mole of HCl produces

= $\frac{3 \times 22.4}{6}L$ of H_2 at S.T.P

= 11.2L of H_2 at STP



Question147

How many moles of magnesium phosphate, $Mg_3(PO_4)_2$ will contain 0.25 mole of oxygen atoms?
[2006]

Options:

- A. 1.25×10^{-2}
- B. 2.5×10^{-2}
- C. 0.02
- D. 3.125×10^{-2}

Answer: D

Solution:

Solution:

1 Mole of $Mg_3(PO_4)_2$ contains 8 moles of oxygen atoms

\therefore 8 mole of oxygen atoms \equiv 1 mole of $Mg_3(PO_4)_2$

0.25 mole of oxygen atom $\equiv \frac{1}{8} \times 0.25$ mole of $Mg_3(PO_4)_2$

$= 3.125 \times 10^{-2}$ mole of $Mg_3(PO_4)_2$

Question148

Density of a 2.05M solution of acetic acid in water is 1.02 g / mL. The molality of the solution is
[2006]

Options:

- A. 2.28mol kg^{-1}
- B. 0.44mol kg^{-1}
- C. 1.14mol kg^{-1}
- D. 3.28mol kg^{-1}

Answer: A

Solution:

Solution:

Apply the formula $d = M \left(\frac{1}{m} + \frac{M_2}{1000} \right)$

$\therefore 1.02 = 2.05 \left(\frac{1}{m} + \frac{60}{1000} \right)$

On solving we get, $m = 2.288 \text{mol / kg}$



Question149

Two solutions of a substance (non electrolyte) are mixed in the following manner: 480mL of 1.5M first solution + 520mL of 1.2M second solution. What is the molarity of the final mixture? [2005]

Options:

- A. 2.70M
- B. 1.344M
- C. 1.50M
- D. 1.20M

Answer: B

Solution:

Solution:

From the molarity equation

$$M_1V_1 + M_2V_2 = MV$$

Let M be the molarity of final mixture,

$$M = \frac{M_1V_1 + M_2V_2}{V} \text{ where } V = V_1 + V_2$$

$$M = \frac{480 \times 1.5 + 520 \times 1.2}{480 + 520} = 1.344M$$

Question150

If we consider that 1 / 6, in place of 1 / 12, mass of carbon atom as the relative atomic mass unit, the mass of one mole of the substance will [2005]

Options:

- A. be a function of the molecular mass of the substance
- B. remain unchanged
- C. increase two fold
- D. decrease twice

Answer: D

Solution:

Solution:

$$\text{Relative atomic mass} = \frac{\text{Mass of one atom of the element}}{1 / 12^{\text{th}} \text{ part of the mass of one atom of carbon} - 12}$$

$$\text{or } \frac{\text{Mass of one atom of the element}}{\text{Mass of one atom of the C - 12}} \times 12$$



Now if we use $\frac{1}{6}$ in place of $\frac{1}{12}$ the formula becomes Relative atomic mass

$$= \frac{\text{Mass of one atom of element}}{\text{Mass of one atom of carbon}} \times 6$$

∴ Relative atomic mass decrease twice.

Question151

6.02×10^{20} molecules of urea are present in 100ml of its solution. The concentration of urea solution is

(Avogadro constant, $N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$)

[2004]

Options:

A. 0.02M

B. 0.01M

C. 0.001M

D. 0.1M

Answer: B

Solution:

Solution:

Moles of urea present in 100mL of sol.

$$= \frac{6.02 \times 10^{20}}{6.02 \times 10^{23}} \text{ mol}$$

$$\therefore M = \frac{6.02 \times 10^{20} \times 1000}{6.02 \times 10^{23} \times 100} = 0.01M$$

[∴ M = Moles of solute present in 1L of solution]

Question152

To neutralise completely 20mL of 0.1M aqueous solution of phosphorous acid (H_3PO_3), the value of 0.1M aqueous KOH solution required is

[2004]

Options:

A. 40mL

B. 20mL

C. 10mL

D. 60mL

Answer: A



Solution:

Solution:

The neutralization reaction is $\text{H}_3\text{PO}_3 + 2\text{KOH} \rightarrow \text{K}_2\text{HPO}_3 + 2\text{H}_2\text{O}$.

Phosphorous acid is diprotic acid as it has two ionizable hydrogens. Thus, 1 mole of phosphorous acid will neutralize 2 moles of KOH.

The number of moles of phosphorous acid present in 20 mL of 0.1M aqueous solution is $0.1 \times 20 \times \frac{1}{1000} = 0.002$ mol.

They will neutralize 2×0.002 mol = 0.004 moles of KOH.

The molarity of KOH solution is 0.1M.

The volume of KOH solution required will be $\frac{0.004}{0.1} = 0.04\text{L} = 40$ ml.

Question153


25mL of a solution of barium hydroxide on titration with a 0.1 molar solution of hydrochloric acid gave a titre value of 35mL. The molarity of barium hydroxide solution was [2003]

Options:

- A. 0.14
- B. 0.28
- C. 0.35
- D. 0.07

Answer: D

Solution:



$$N_1 V_1 = N_2 V_2$$

$$N_1 \times 25 = 0.1 \times 35$$

$$N_1 = 0.14$$

Since, Ba(OH)_2 is diacid base

$$\text{Hence } N = M \times 2 \text{ or } M = \frac{N}{2}$$

$$M = 0.07\text{M}$$

Question154

What volume of hydrogen gas, at 273K and 1atm. pressure will be consumed in obtaining 21.6g of elemental boron (atomic mass = 10.8) from the reduction of boron trichloride by hydrogen ? [2003]

Options:

- A. 67.2L



B. 44.8L

C. 22.4L

D. 89.6 L

Answer: A

Solution:



Now, since 10.8g boron requires hydrogen

$$= \frac{3}{2} \times 22.4\text{L at S.T.P.}$$

Hence 21.6 boron requires hydrogen

$$= \frac{3}{2} \times \frac{22.4}{10.8} \times 21.6 = 67.2\text{L at S.T.P.}$$

Question155

**With increase of temperature, which of these changes?
[2002]**

Options:

A. molality

B. weight fraction of solute

C. molarity

D. mole fraction.

Answer: C

Solution:

Solution:

Among all the given options, molarity changes with temperature because the term molarity involves volume which increases on increasing temperature.

Question156

**Number of atoms in 558.5 gram Fe (at. wt. of Fe = 55.85gmol⁻¹) is
[2002]**

Options:

A. twice that in 60g carbon

B. 6.023×10^{22}



C. half that in 8g He

D. $558.5 \times 6.023 \times 10^{23}$

Answer: A

Solution:

Solution:

$$Fe(\text{No. of moles}) = \frac{558.5}{55.85} = 10\text{mol}$$





$$C(\text{No. of moles}) \text{ in } 60\text{g of } C = 60 / 12 = 5\text{mol.}$$

Question157

In a compound C, H and N atoms are present in 9 : 1 : 3.5 by weight. Molecular weight of compound is 108 . Molecular formula of compound is

[2002]

Options:



Answer: C

Solution:

Element	%	Relative no. of atoms	Simple ratio of atoms
C	9	$\frac{9}{12} = \frac{3}{4}$	3
H	1	$\frac{1}{1} = 1$	4
N	3.5	$\frac{3.5}{14} = \frac{1}{4}$	1

Empirical formula = C_3H_4N



$$(C_3H_4N)_n = 108$$

$$(12 \times 3 + 4 \times 1 + 14)_n = 108$$

$$(54)_n = 108 \quad n = \frac{108}{54} = 2$$

$$\therefore \text{Molecular formula} = C_6H_8N_2$$

