

Continuity and Differentiability



TOPIC 1 Continuity



1. Let $f(x) = x \cdot \left[\frac{x}{2} \right]$, for $-10 < x < 10$, where $[t]$ denotes the greatest integer function. Then the number of points of discontinuity of f is equal to _____.

[NA Sep. 05, 2020 (I)]

2. If a function $f(x)$ defined by

$$f(x) = \begin{cases} ae^x + be^{-x}, & -1 \leq x < 1 \\ cx^2, & 1 \leq x \leq 3 \\ ax^2 + 2cx, & 3 < x \leq 4 \end{cases} \text{ be continuous for some}$$

$a, b, c \in \mathbf{R}$ and $f'(0) + f'(2) = e$, then the value of a is:

[Sep. 02, 2020 (I)]

- (a) $\frac{1}{e^2 - 3e + 13}$ (b) $\frac{e}{e^2 - 3e - 13}$
 (c) $\frac{e}{e^2 + 3e + 13}$ (d) $\frac{e}{e^2 - 3e + 13}$

3. Let $[t]$ denote the greatest integer $\leq t$ and $\lim_{x \rightarrow 0} x \left[\frac{4}{x} \right] = A$.

Then the function, $f(x) = [x^2] \sin(\pi x)$ is discontinuous, when x is equal to:

[Jan. 9, 2020 (II)]

- (a) $\sqrt{A+1}$ (b) $\sqrt{A+5}$
 (c) $\sqrt{A+21}$ (d) \sqrt{A}

4. If the function f defined on $\left(-\frac{1}{3}, \frac{1}{3}\right)$ by

$$f(x) = \begin{cases} \frac{1}{x} \log_e \left(\frac{1+3x}{1-2x} \right), & \text{when } x \neq 0 \\ k, & \text{when } x = 0 \end{cases} \text{ is continuous, then } k$$

is equal to _____.

[NA Jan. 7, 2020 (II)]

5. If the function f defined on $\left(\frac{\pi}{6}, \frac{\pi}{3}\right)$ by

$$f(x) = \begin{cases} \frac{\sqrt{2} \cos x - 1}{\cot x - 1}, & x \neq \frac{\pi}{4} \\ k, & x = \frac{\pi}{4} \end{cases}$$

is continuous, then k is equal to: [April 09, 2019 (I)]

- (a) 2 (b) $\frac{1}{2}$ (c) 1 (d) $\frac{1}{\sqrt{2}}$

6. If $f(x) = [x] - \left[\frac{x}{4} \right]$, $x \in \mathbf{R}$, where $[x]$ denotes the greatest integer function, then:

[April 09, 2019 (II)]

- (a) f is continuous at $x = 4$.
 (b) $\lim_{x \rightarrow 4^+} f(x)$ exists but $\lim_{x \rightarrow 4^-} f(x)$ does not exist.
 (c) Both $\lim_{x \rightarrow 4^-} f(x)$ and $\lim_{x \rightarrow 4^+} f(x)$ exist but are not equal.
 (d) $\lim_{x \rightarrow 4^-} f(x)$ exists but $\lim_{x \rightarrow 4^+} f(x)$ does not exist.

7. If the function

$$f(x) = \begin{cases} a|\pi - x| + 1, & x \leq 5 \\ b|x - \pi| + 3, & x > 5 \end{cases}$$

is continuous at $x = 5$, then the value of $a - b$ is:

[April 09, 2019 (II)]

- (a) $\frac{2}{\pi+5}$ (b) $\frac{-2}{\pi+5}$ (c) $\frac{2}{\pi-5}$ (d) $\frac{2}{5-\pi}$

8. Let $f: [-1, 3] \rightarrow \mathbf{R}$ be defined as

$$f(x) = \begin{cases} |x| + [x], & -1 \leq x < 1 \\ x + |x|, & 1 \leq x < 2 \\ x + [x], & 2 \leq x \leq 3, \end{cases}$$

where $[t]$ denotes the greatest integer less than or equal to t . Then, f is discontinuous at: [April 08, 2019 (II)]

- (a) only one point (b) only two points
 (c) only three points (d) four or more points



9. Let $f: \mathbf{R} \rightarrow \mathbf{R}$ be a function defined as

$$f(x) = \begin{cases} 5, & \text{if } x \leq 1 \\ a + bx, & \text{if } 1 < x < 3 \\ b + 5x, & \text{if } 3 \leq x < 5 \\ 30, & \text{if } x \geq 5 \end{cases}$$

Then, f is :

[Jan 09, 2019 (I)]

- (a) continuous if $a = 5$ and $b = 5$
 (b) continuous if $a = -5$ and $b = 10$
 (c) continuous if $a = 0$ and $b = 5$
 (d) not continuous for any values of a and b

10. If the function f defined as

$$f(x) = \frac{1}{x} - \frac{k-1}{e^{2x}-1}$$

$x \neq 0$, is continuous at $x = 0$,
 then the ordered pair $(k, f(0))$ is equal to?

[Online April 16, 2018]

- (a) (3, 1) (b) (3, 2) (c) $(\frac{1}{3}, 2)$ (d) (2, 1)

11. Let $f(x) = \begin{cases} (x-1)^{\frac{1}{2-x}}, & x > 1, x \neq 2 \\ k, & x = 2 \end{cases}$

The value of k for which f is continuous at $x = 2$ is

[Online April 15, 2018]

- (a) e^{-2} (b) e (c) e^{-1} (d) 1

12. The value of k for which the function

$$f(x) = \begin{cases} \left(\frac{4}{5}\right)^{\frac{\tan 4x}{\tan 5x}}, & 0 < x < \frac{\pi}{2} \\ k + \frac{2}{5}, & x = \frac{\pi}{2} \end{cases} \text{ is continuous at } x = \frac{\pi}{2}, \text{ is :}$$

[Online April 9, 2017]

- (a) $\frac{17}{20}$ (b) $\frac{2}{5}$ (c) $\frac{3}{5}$ (d) $-\frac{2}{5}$

13. Let $a, b \in \mathbf{R}, (a \neq 0)$. If the function f defined as

$$f(x) = \begin{cases} \frac{2x^2}{a}, & 0 \leq x < 1 \\ a, & 1 \leq x < \sqrt{2} \\ \frac{2b^2 - 4b}{x^3}, & \sqrt{2} \leq x < \infty \end{cases}$$

is continuous in the interval $[0, \infty)$, then an ordered pair

(a, b) is : [Online April 10, 2016]

- (a) $(-\sqrt{2}, 1 - \sqrt{3})$ (b) $(\sqrt{2}, -1 + \sqrt{3})$
 (c) $(\sqrt{2}, 1 - \sqrt{3})$ (d) $(-\sqrt{2}, 1 + \sqrt{3})$

14. Let k be a non-zero real number.

[Online April 11, 2015]

$$\text{If } f(x) = \begin{cases} \frac{(e^x - 1)}{\sin\left(\frac{x}{k}\right) \log\left(1 + \frac{x}{4}\right)}, & x \neq 0 \\ 12, & x = 0 \end{cases}$$

is a continuous function then the value of k is:

- (a) 4 (b) 1 (c) 3 (d) 2

15. If the function

$$f(x) = \begin{cases} \frac{\sqrt{2 + \cos x} - 1}{(\pi - x)^2}, & x \neq \pi \\ k, & x = \pi \end{cases}$$

is continuous at $x = \pi$, then k equals:

[Online April 19, 2014]

- (a) 0 (b) $\frac{1}{2}$ (c) 2 (d) $\frac{1}{4}$

16. If $f(x)$ is continuous and $f\left(\frac{9}{2}\right) = \frac{2}{9}$, then

$$\lim_{x \rightarrow 0} f\left(\frac{1 - \cos 3x}{x^2}\right) \text{ is equal to:}$$

[Online April 9, 2014]

- (a) $\frac{9}{2}$ (b) $\frac{2}{9}$ (c) 0 (d) $\frac{8}{9}$

17. Consider the function :

$f(x) = [x] + |1 - x|$, $-1 \leq x \leq 3$ where $[x]$ is the greatest integer function.

Statement 1 : f is not continuous at $x = 0, 1, 2$ and 3 .

$$\text{Statement 2 : } f(x) = \begin{cases} -x, & -1 \leq x < 0 \\ 1 - x, & 0 \leq x < 1 \\ 1 + x, & 1 \leq x < 2 \\ 2 + x, & 2 \leq x \leq 3 \end{cases}$$

[Online April 25, 2013]

- (a) Statement 1 is true ; Statement 2 is false,
 (b) Statement 1 is true; Statement 2 is true; Statement 2 is not correct explanation for Statement 1.
 (c) Statement 1 is true; Statement 2 is true; Statement It is a correct explanation for Statement 1.
 (d) Statement 1 is false; Statement 2 is true.

18. Let f be a composite function of x defined by

$$f(u) = \frac{1}{u^2 + u - 2}, u(x) = \frac{1}{x - 1}$$

Then the number of points x where f is discontinuous is :

[Online April 23, 2013]

- (a) 4 (b) 3 (c) 2 (d) 1

19. Let $f(x) = -1 + |x - 2|$, and $g(x) = 1 - |x|$; then the set of all points where $f \circ g$ is discontinuous is :

[Online April 22, 2013]

- (a) $\{0, 2\}$ (b) $\{0, 1, 2\}$
- (c) $\{0\}$ (d) an empty set

20. If $f : R \rightarrow R$ is a function defined by $f(x) = [x] \cos\left(\frac{2x-1}{2}\right)\pi$, where $[x]$ denotes the greatest integer function, then f is . [2012]

- (a) continuous for every real x .
- (b) discontinuous only at $x = 0$
- (c) discontinuous only at non-zero integral values of x .
- (d) continuous only at $x = 0$.

21. Let $f : [1, 3] \rightarrow R$ be a function satisfying

$$\frac{x}{[x]} \leq f(x) \leq \sqrt{6-x}, \text{ for all } x \neq 2 \text{ and } f(2) = 1,$$

where R is the set of all real numbers and $[x]$ denotes the largest integer less than or equal to x .

Statement 1: $\lim_{x \rightarrow 2^-} f(x)$ exists. [Online May 19, 2012]

Statement 2: f is continuous at $x = 2$.

- (a) Statement 1 is true, Statement 2 is true, Statement 2 is a correct explanation for Statement 1.
- (b) Statement 1 is false, Statement 2 is true.
- (c) Statement 1 is true, Statement 2 is true, Statement 2 is not a correct explanation for Statement 1.
- (d) Statement 1 is true, Statement 2 is false.

22. **Statement 1:** A function $f: R \rightarrow R$ is continuous at x_0 if and only if $\lim_{x \rightarrow x_0} f(x)$ exists and $\lim_{x \rightarrow x_0} f(x) = f(x_0)$

Statement 2: A function $f: R \rightarrow R$ is discontinuous at x_0 if and only if, $\lim_{x \rightarrow x_0} f(x)$ exists and $\lim_{x \rightarrow x_0} f(x) \neq f(x_0)$.

[Online May 12, 2012]

- (a) Statement 1 is true, Statement 2 is true, Statement 2 is not a correct explanation of Statement 1.
- (b) Statement 1 is false, Statement 2 is true.
- (c) Statement 1 is true, Statement 2 is true, Statement 2 is a correct explanation of Statement 1.
- (d) Statement 1 is true, Statement 2 is false.

23. Define $f(x)$ as the product of two real functions [2011RS]

$$f_1(x) = x, x \in R, \text{ and } f_2(x) = \begin{cases} \sin \frac{1}{x}, & \text{if } x \neq 0 \\ 0, & \text{if } x = 0 \end{cases}$$

as follows :

$$f(x) = \begin{cases} f_1(x) \cdot f_2(x), & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$$

Statement - 1 : $f(x)$ is continuous on R .

Statement - 2 : $f_1(x)$ and $f_2(x)$ are continuous on R .

- (a) Statement -1 is true, Statement-2 is true; Statement-2 is a correct explanation for Statement-1.
- (b) Statement-1 is true, Statement-2 is true; Statement-2 is NOT a correct explanation for Statement-1
- (c) Statement-1 is true, Statement-2 is false
- (d) Statement-1 is false, Statement-2 is true

24. The values of p and q for which the function [2011]

$$f(x) = \begin{cases} \frac{\sin(p+1)x + \sin x}{x}, & x < 0 \\ q, & x = 0 \\ \frac{\sqrt{x+x^2} - \sqrt{x}}{x^{3/2}}, & x > 0 \end{cases}$$

are

- (a) $p = \frac{5}{2}, q = \frac{1}{2}$ (b) $p = -\frac{3}{2}, q = \frac{1}{2}$
- (c) $p = \frac{1}{2}, q = \frac{3}{2}$ (d) $p = \frac{1}{2}, q = -\frac{3}{2}$

25. The function $f : R/\{0\} \rightarrow R$ given by [2007]

$$f(x) = \frac{1}{x} - \frac{2}{e^{2x} - 1}$$

can be made continuous at $x = 0$ by defining $f(0)$ as

- (a) 0 (b) 1
- (c) 2 (d) -1

26. Let $f(x) = \frac{1 - \tan x}{4x - \pi}, x \neq \frac{\pi}{4}, x \in \left[0, \frac{\pi}{2}\right]$.

If $f(x)$ is continuous in $\left[0, \frac{\pi}{2}\right]$, then $f\left(\frac{\pi}{4}\right)$ is [2004]

- (a) -1 (b) $\frac{1}{2}$
- (c) $-\frac{1}{2}$ (d) 1

27. f is defined in $[-5, 5]$ as [2002]

$$f(x) = \begin{cases} x & \text{if } x \text{ is rational} \\ -x & \text{if } x \text{ is irrational.} \end{cases}$$

- (a) $f(x)$ is continuous at every x , except $x = 0$
- (b) $f(x)$ is discontinuous at every x , except $x = 0$
- (c) $f(x)$ is continuous everywhere
- (d) $f(x)$ is discontinuous everywhere

TOPIC 2 Differentiability


28. Let $f: \mathbf{R} \rightarrow \mathbf{R}$ be a function defined by $f(x) = \max\{x, x^2\}$. Let S denote the set of all points in \mathbf{R} , where f is not differentiable. Then: **[Sep. 06, 2020 (II)]**

(a) $\{0, 1\}$ (b) $\{0\}$
 (c) ϕ (an empty set) (d) $\{1\}$

29. If the function $f(x) = \begin{cases} k_1(x-\pi)^2 - 1, & x \leq \pi \\ k_2 \cos x, & x > \pi \end{cases}$ is twice differentiable, then the ordered pair (k_1, k_2) is equal to: **[Sep. 05, 2020 (I)]**

(a) $\left(\frac{1}{2}, 1\right)$ (b) $(1, 0)$
 (c) $\left(\frac{1}{2}, -1\right)$ (d) $(1, 1)$

30. Let f be a twice differentiable function on $(1, 6)$. If $f(2) = 8$, $f'(2) = 5$, $f'(x) \geq 1$ and $f''(x) \geq 4$, for all $x \in (1, 6)$, then: **[Sep. 04, 2020 (I)]**

(a) $f(5) + f'(5) \leq 26$ (b) $f(5) + f'(5) \geq 28$
 (c) $f'(5) + f''(5) \leq 20$ (d) $f(5) \leq 10$

31. Suppose a differentiable function $f(x)$ satisfies the identity $f(x+y) = f(x) + f(y) + xy^2 + x^2y$, for all real x and y . If

$$\lim_{x \rightarrow 0} \frac{f(x)}{x} = 1, \text{ then } f'(3) \text{ is equal to } \underline{\hspace{2cm}}.$$

[NA Sep. 04, 2020 (I)]

32. The function $f(x) = \begin{cases} \frac{\pi}{4} + \tan^{-1} x, & |x| \leq 1 \\ \frac{1}{2}(|x| - 1), & |x| > 1 \end{cases}$ is: **[Sep. 04, 2020 (II)]**

(a) continuous on $\mathbf{R} - \{1\}$ and differentiable on $\mathbf{R} - \{-1, 1\}$.
 (b) both continuous and differentiable on $\mathbf{R} - \{1\}$.
 (c) continuous on $\mathbf{R} - \{-1\}$ and differentiable on $\mathbf{R} - \{-1, 1\}$.
 (d) both continuous and differentiable on $\mathbf{R} - \{-1\}$.

33. If $f(x) = \begin{cases} \frac{\sin(a+2)x + \sin x}{x} & ; x < 0 \\ b & ; x = 0 \\ \frac{(x+3x^2)^{1/3} - x^{1/3}}{x^{4/3}} & ; x > 0 \end{cases}$

is continuous at $x = 0$, then $a + 2b$ is equal to:

[Jan. 9, 2020 (I)]

(a) 1 (b) -1 (c) 0 (d) -2

34. Let f and g be differentiable functions on \mathbf{R} such that fg is the identity function. If for some $a, b \in \mathbf{R}$, $g'(a) = 5$ and $g(a) = b$, then $f'(b)$ is equal to: **[Jan. 9, 2020 (II)]**

(a) $\frac{1}{5}$ (b) 1 (c) 5 (d) $\frac{2}{5}$

35. Let S be the set of all functions $f: [0, 1] \rightarrow \mathbf{R}$, which are continuous on $[0, 1]$ and differentiable on $(0, 1)$. Then for every f in S , there exists a $c \in (0, 1)$, depending on f , such that: **[Jan. 8, 2020 (II)]**

(a) $|f(c) - f(1)| < (1-c)f'(c)$
 (b) $\frac{f(1) - f(c)}{1-c} = f'(c)$
 (c) $|f(c) + f(1)| < (1+c)|f'(c)|$
 (d) $|f(c) - f(1)| < |f'(c)|$

36. Let the function, $f: [-7, 0] \rightarrow \mathbf{R}$ be continuous on $[-7, 0]$ and differentiable on $(-7, 0)$. If $f(-7) = -3$ and $f'(x) \geq 2$, for all $x \in (-7, 0)$, then for all such functions, $f'(-1) + f(0)$ lies in the interval: **[Jan. 7, 2020 (I)]**

(a) $(-\infty, 20]$ (b) $[-3, 11]$
 (c) $(-\infty, 11]$ (d) $[-6, 20]$

37. Let S be the set of points where the function, $f(x) = |2 - |x - 3||$, $x \in \mathbf{R}$, is not differentiable.

Then $\sum_{x \in S} f(f(x))$ is equal to ———. **[NA Jan. 7, 2020 (I)]**

38. If $f(x) = \begin{cases} \frac{\sin(p+1)x + \sin x}{x}, & x < 0 \\ q, & x = 0 \\ \frac{\sqrt{x+x^2} - \sqrt{x}}{x^{3/2}}, & x > 0 \end{cases}$

is continuous at $x = 0$, then the ordered pair (p, q) is equal to: **[April 10, 2019 (I)]**

(a) $\left(-\frac{3}{2}, -\frac{1}{2}\right)$ (b) $\left(-\frac{1}{2}, \frac{3}{2}\right)$
 (c) $\left(-\frac{3}{2}, \frac{1}{2}\right)$ (d) $\left(\frac{5}{2}, \frac{1}{2}\right)$

39. Let $f(x) = \log_e(\sin x)$, $(0 < x < \pi)$ and $g(x) = \sin^{-1}(e^{-x})$, $(x \geq 0)$. If α is a positive real number such that $a = (f \circ g)'(\alpha)$ and $b = (f \circ g)(\alpha)$, then: **[April 10, 2019 (II)]**

(a) $a\alpha^2 + b\alpha + a = 0$ (b) $a\alpha^2 - b\alpha - a = 1$
 (c) $a\alpha^2 - b\alpha - a = 0$ (d) $a\alpha^2 + b\alpha - a = -2a^2$

40. Let $f: \mathbf{R} \rightarrow \mathbf{R}$ be differentiable at $c \in \mathbf{R}$ and $f(c) = 0$. If $g(x) = |f(x)|$, then at $x = c$, g is : **[April 10, 2019 (I)]**
 (a) not differentiable if $f'(c) = 0$
 (b) differentiable if $f''(c) \neq 0$
 (c) differentiable if $f'(c) = 0$
 (d) not differentiable
41. Let $f(x) = 15 - |x - 10|$; $x \in \mathbf{R}$. Then the set of all values of x , at which the function, $g(x) = f(f(x))$ is not differentiable, is: **[April 09, 2019 (I)]**
 (a) $\{5, 10, 15\}$ (b) $\{10, 15\}$
 (c) $\{5, 10, 15, 20\}$ (d) $\{10\}$
42. If $f(1) = 1, f'(1) = 3$, then the derivative of $f(f(f(x))) + (f(x))^2$ at $x = 1$ is : **[April 08, 2019 (II)]**
 (a) 33 (b) 12 (c) 15 (d) 9
43. Let f be a differentiable function such that $f(1) = 2$ and $f'(x) = f(x)$ for all $x \in \mathbf{R}$. If $h(x) = f(f(x))$, then $h'(1)$ is equal to : **[Jan. 12, 2019 (II)]**
 (a) $2e^2$ (b) $4e$ (c) $2e$ (d) $4e^2$
44. Let $f(x) = \begin{cases} -1, & -2 \leq x < 0 \\ x^2 - 1, & 0 \leq x \leq 2 \end{cases}$ and $g(x) = |f(x)| + f(|x|)$. Then, in the interval $(-2, 2)$, g is : **[Jan. 11, 2019 (I)]**
 (a) differentiable at all points
 (b) not continuous
 (c) not differentiable at two points
 (d) not differentiable at one point
45. If $x \log_e (\log_e x) - x^2 + y^2 = 4 (y > 0)$, then $\frac{dy}{dx}$ at $x = e$ is equal to : **[Jan. 11, 2019 (I)]**
 (a) $\frac{(1+2e)}{2\sqrt{4+e^2}}$ (b) $\frac{(2e-1)}{2\sqrt{4+e^2}}$
 (c) $\frac{(1+2e)}{\sqrt{4+e^2}}$ (d) $\frac{e}{\sqrt{4+e^2}}$
46. Let K be the set of all real values of x where the function $f(x) = \sin |x| - |x| + 2(x - \pi) \cos |x|$ is not differentiable. Then the set K is equal to : **[Jan. 11, 2019 (II)]**
 (a) \emptyset (an empty set) (b) $\{\pi\}$
 (c) $\{0\}$ (d) $\{0, \pi\}$
47. Let $f(x) = \begin{cases} \max\{|x|, x^2\} & |x| \leq 2 \\ 8 - 2|x| & 2 < |x| \leq 4 \end{cases}$
 Let S be the set of points in the interval $(-4, 4)$ at which f is not differentiable. Then S : **[Jan 10, 2019 (I)]**
 (a) is an empty set
 (b) equals $\{-2, -1, 0, 1, 2\}$
 (c) equals $\{-2, -1, 1, 2\}$
 (d) equals $\{-2, 2\}$
48. Let $f: (-1, 1) \rightarrow \mathbf{R}$ be a function defined by $f(x) = \max\{-|x|, -\sqrt{1-x^2}\}$. If K be the set of all points at which f is not differentiable, then K has exactly: **[Jan. 10, 2019 (II)]**
 (a) five elements (b) one element
 (c) three elements (d) two elements
49. Let $S = \{t \in \mathbf{R} : f(x) = |x - \pi| (e^{|x|} - 1) \sin |x| \text{ is not differentiable at } t\}$. Then the set S is equal to : **[2018]**
 (a) $\{0\}$ (b) $\{\pi\}$
 (c) $\{0, \pi\}$ (d) \emptyset (an empty set)
50. Let $S = \{(\lambda, \mu) \in \mathbf{R} \times \mathbf{R} : f(t) = (\lambda|e^t| - \mu) \cdot \sin(2|t|), t \in \mathbf{R}$, is a differentiable function}. Then S is a subset of? **[Online April 15, 2018]**
 (a) $\mathbf{R} \times [0, \infty)$ (b) $(-\infty, 0) \times \mathbf{R}$
 (c) $[0, \infty) \times \mathbf{R}$ (d) $\mathbf{R} \times (-\infty, 0)$
51. If the function $f(x) = \begin{cases} -x, & x < 1 \\ a + \cos^{-1}(x+b), & 1 \leq x \leq 2 \end{cases}$ is differentiable at $x = 1$, then $\frac{a}{b}$ is equal to : **[Online April 9, 2016]**
 (a) $\frac{\pi+2}{2}$ (b) $\frac{\pi-2}{2}$
 (c) $\frac{-\pi-2}{2}$ (d) $-1 - \cos^{-1}(2)$
52. If the function $g(x) = \begin{cases} k\sqrt{x+1}, & 0 \leq x \leq 3 \\ mx+2, & 3 < x \leq 5 \end{cases}$ is differentiable, then the value of $k+m$ is : **[2015]**
 (a) $\frac{10}{3}$ (b) 4
 (c) 2 (d) $\frac{16}{5}$
53. Let $f: \mathbf{R} \rightarrow \mathbf{R}$ be a function such that $|f'(x)| \leq x^2$, for all $x \in \mathbf{R}$. Then, at $x = 0$, f is: **[Online April 19, 2014]**
 (a) continuous but not differentiable.
 (b) continuous as well as differentiable.
 (c) neither continuous nor differentiable.
 (d) differentiable but not continuous.

54. Let $f, g: \mathbb{R} \rightarrow \mathbb{R}$ be two functions defined by

$$f(x) = \begin{cases} x \sin\left(\frac{1}{x}\right), & x \neq 0 \\ 0, & x = 0 \end{cases}, \text{ and } g(x) = x f(x)$$

Statement I: f is a continuous function at $x = 0$.

Statement II: g is a differentiable function at $x = 0$.

[Online April 12, 2014]

- (a) Both statement I and II are false.
 (b) Both statement I and II are true.
 (c) Statement I is true, statement II is false.
 (d) Statement I is false, statement II is true.
55. Consider the function, $f(x) = |x-2| + |x-5|, x \in \mathbb{R}$.
Statement-1: $f'(4) = 0$
Statement-2: f is continuous in $[2, 5]$, differentiable in $(2, 5)$ and $f(2) = f(5)$. [2012]
- (a) Statement-1 is false, Statement-2 is true.
 (b) Statement-1 is true, statement-2 is true; statement-2 is a correct explanation for Statement-1.
 (c) Statement-1 is true, statement-2 is true; statement-2 is not a correct explanation for Statement-1.
 (d) Statement-1 is true, statement-2 is false.
56. If $f(x) = a |\sin x| + b e^{|x|} + c|x|^3$, where $a, b, c \in \mathbb{R}$, is differentiable at $x = 0$, then [Online May 26, 2012]
- (a) $a = 0, b$ and c are any real numbers
 (b) $c = 0, a = 0, b$ is any real number
 (c) $b = 0, c = 0, a$ is any real number
 (d) $a = 0, b = 0, c$ is any real number
57. If $x + |y| = 2y$, then y as a function of x , at $x = 0$ is [Online May 7, 2012]
- (a) differentiable but not continuous
 (b) continuous but not differentiable
 (c) continuous as well as differentiable
 (d) neither continuous nor differentiable
58. If function $f(x)$ is differentiable at $x = a$,

then $\lim_{x \rightarrow a} \frac{x^2 f(a) - a^2 f(x)}{x - a}$ is : [2011RS]

- (a) $-a^2 f'(a)$ (b) $a f(a) - a^2 f'(a)$
 (c) $2af(a) - a^2 f'(a)$ (d) $2af(a) + a^2 f'(a)$
59. Let $f(x) = \begin{cases} (x-1) \sin \frac{1}{x-1} & \text{if } x \neq 1 \\ 0 & \text{if } x = 1 \end{cases}$ [2008]

Then which one of the following is true?

- (a) f is neither differentiable at $x = 0$ nor at $x = 1$
 (b) f is differentiable at $x = 0$ and at $x = 1$
 (c) f is differentiable at $x = 0$ but not at $x = 1$
 (d) f is differentiable at $x = 1$ but not at $x = 0$

60. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a function defined by

$$f(x) = \min \{x+1, |x|+1\}, \text{ Then which of the following is true?}$$

- (a) $f(x)$ is differentiable everywhere [2007]
 (b) $f(x)$ is not differentiable at $x = 0$
 (c) $f(x) \geq 1$ for all $x \in \mathbb{R}$
 (d) $f(x)$ is not differentiable at $x = 1$

61. The set of points where $f(x) = \frac{x}{1+|x|}$ is differentiable is [2006]

- (a) $(-\infty, 0) \cup (0, \infty)$ (b) $(-\infty, -1) \cup (-1, \infty)$
 (c) $(-\infty, \infty)$ (d) $(0, \infty)$

62. If f is a real valued differentiable function satisfying $|f(x) - f(y)| \leq (x - y)^2, x, y \in \mathbb{R}$ and $f(0) = 0$, then $f(1)$ equals [2005]

- (a) -1 (b) 0
 (c) 2 (d) 1

63. Suppose $f(x)$ is differentiable at $x = 1$ and

$$\lim_{h \rightarrow 0} \frac{1}{h} f(1+h) = 5, \text{ then } f'(1) \text{ equals [2005]}$$

- (a) 3 (b) 4 (c) 5 (d) 6

64. If $f(x) = \begin{cases} x e^{-\left(\frac{1}{|x|} + \frac{1}{x}\right)}, & x \neq 0 \\ 0, & x = 0 \end{cases}$ then $f(x)$ is

- (a) discontinuous every where [2003]
 (b) continuous as well as differentiable for all x
 (c) continuous for all x but not differentiable at $x = 0$
 (d) neither differentiable nor continuous at $x = 0$

TOPIC 3

Chain Rule of Differentiation, Differentiation of Explicit & Implicit Functions, Parametric & Composite Functions, Logarithmic & Exponential Functions, Inverse Functions, Differentiation by Trigonometric Substitution



65. The derivative of $\tan^{-1} \left(\frac{\sqrt{1+x^2}-1}{x} \right)$ with respect to

$$\tan^{-1} \left(\frac{2x\sqrt{1-x^2}}{1-2x^2} \right) \text{ at } x = \frac{1}{2} \text{ is : [Sep. 05, 2020 (II)]}$$

- (a) $\frac{2\sqrt{3}}{5}$ (b) $\frac{\sqrt{3}}{12}$ (c) $\frac{2\sqrt{3}}{3}$ (d) $\frac{\sqrt{3}}{10}$

66. If $(a + \sqrt{2}b \cos x)(a - \sqrt{2}b \cos y) = a^2 - b^2$, where $a > b > 0$, then $\frac{dx}{dy}$ at $\left(\frac{\pi}{4}, \frac{\pi}{4}\right)$ is: [Sep. 04, 2020 (I)]

- (a) $\frac{a-2b}{a+2b}$ (b) $\frac{a-b}{a+b}$ (c) $\frac{a+b}{a-b}$ (d) $\frac{2a+b}{2a-b}$

67. If $y = \sum_{k=1}^6 k \cos^{-1} \left\{ \frac{3}{5} \cos kx - \frac{4}{5} \sin kx \right\}$, then $\frac{dy}{dx}$ at $x=0$ is _____ . [NA Sep. 02, 2020 (II)]

68. If $x = 2\sin\theta - \sin 2\theta$ and $y = 2\cos\theta - \cos 2\theta$, $\theta \in [0, 2\pi]$, then $\frac{d^2y}{dx^2}$ at $\theta = \pi$ is : [Jan. 9, 2020 (II)]

- (a) $\frac{3}{4}$ (b) $-\frac{3}{8}$ (c) $\frac{3}{2}$ (d) $-\frac{3}{4}$

69. If $y(\alpha) = \sqrt{2 \left(\frac{\tan \alpha + \cot \alpha}{1 + \tan^2 \alpha} \right) + \frac{1}{\sin^2 \alpha}}$, $\alpha \in \left(\frac{3\pi}{4}, \pi\right)$, then $\frac{dy}{d\alpha}$ at $\alpha = \frac{5\pi}{6}$ is: [Jan. 7, 2020 (I)]

- (a) 4 (b) $\frac{4}{3}$ (c) -4 (d) $-\frac{1}{4}$

70. Let $y = y(x)$ be a function of x satisfying $y\sqrt{1-x^2} = k - x\sqrt{1-y^2}$ where k is a constant and $y\left(\frac{1}{2}\right) = -\frac{1}{4}$. Then $\frac{dy}{dx}$ at $x = \frac{1}{2}$, is equal to: [Jan. 7, 2020 (II)]

- (a) $-\frac{\sqrt{5}}{4}$ (b) $-\frac{\sqrt{5}}{2}$ (c) $\frac{2}{\sqrt{5}}$ (d) $\frac{\sqrt{5}}{2}$

71. If $e^y + xy = e$, the ordered pair $\left(\frac{dy}{dx}, \frac{d^2y}{dx^2}\right)$ at $x=0$ is equal to : [April 12, 2019 (I)]

- (a) $\left(\frac{1}{e}, -\frac{1}{e^2}\right)$ (b) $\left(-\frac{1}{e}, \frac{1}{e^2}\right)$
 (c) $\left(\frac{1}{e}, \frac{1}{e^2}\right)$ (d) $\left(-\frac{1}{e}, -\frac{1}{e^2}\right)$

72. The derivative of $\tan^{-1} \left(\frac{\sin x - \cos x}{\sin x + \cos x} \right)$, with respect to $\frac{x}{2}$, where $x \in \left(0, \frac{\pi}{2}\right)$ is : [April 12, 2019 (II)]

- (a) 1 (b) $\frac{2}{3}$ (c) $\frac{1}{2}$ (d) 2

73. If $2y = \left(\cot^{-1} \left(\frac{\sqrt{3} \cos x + \sin x}{\cos x - \sqrt{3} \sin x} \right) \right)^2$, $x \in \left(0, \frac{\pi}{2}\right)$ then $\frac{dy}{dx}$ is equal to : [April 08, 2019 (I)]

- (a) $\frac{\pi}{6} - x$ (b) $x - \frac{\pi}{6}$ (c) $\frac{\pi}{3} - x$ (d) $2x - \frac{\pi}{3}$

74. Let S be the set of all points in $(-\pi, \pi)$ at which the function $f(x) = \min \{ \sin x, \cos x \}$ is not differentiable. Then S is a subset of which of the following? [Jan. 12, 2019 (I)]

- (a) $\left\{ -\frac{\pi}{4}, 0, \frac{\pi}{4} \right\}$ (b) $\left\{ -\frac{3\pi}{4}, -\frac{\pi}{4}, \frac{3\pi}{4}, \frac{\pi}{4} \right\}$
 (c) $\left\{ -\frac{\pi}{2}, -\frac{\pi}{4}, \frac{\pi}{4}, \frac{\pi}{2} \right\}$ (d) $\left\{ -\frac{3\pi}{4}, -\frac{\pi}{2}, \frac{\pi}{2}, \frac{3\pi}{4} \right\}$

75. For $x > 1$, if $(2x)^{2y} = 4e^{2x-2y}$, then $(1 + \log_e 2x)^2 \frac{dy}{dx}$ is equal to : [Jan. 12, 2019 (I)]

- (a) $\frac{x \log_e 2x - \log_e 2}{x}$ (b) $\log_e 2x$
 (c) $\frac{x \log_e 2x + \log_e 2}{x}$ (d) $x \log_e 2x$

76. Let $f: \mathbf{R} \rightarrow \mathbf{R}$ be a function such that $f(x) = x^3 + x^2 f'(1) + x f''(2) + f'''(3)$, $x \in \mathbf{R}$. Then $f(2)$ equals: [Jan 10, 2019 (I)]

- (a) -4 (b) 30 (c) -2 (d) 8

77. If $x = 3 \tan t$ and $y = 3 \sec t$, then the value of $\frac{d^2y}{dx^2}$ at $t = \frac{\pi}{4}$, is: [Jan. 09, 2019 (II)]

- (a) $\frac{1}{3\sqrt{2}}$ (b) $\frac{1}{6\sqrt{2}}$ (c) $\frac{3}{2\sqrt{2}}$ (d) $\frac{1}{6}$

78. If $x = \sqrt{2^{\csc^{-1} t}}$ and $y = \sqrt{2^{\sec^{-1} t}}$ ($|t| \geq 1$), then $\frac{dy}{dx}$ is equal to. [Online April 16, 2018]

- (a) $\frac{y}{x}$ (b) $-\frac{y}{x}$ (c) $-\frac{x}{y}$ (d) $\frac{x}{y}$

79. If $f(x) = \begin{vmatrix} \cos x & x & 1 \\ 2 \sin x & x^2 & 2x \\ \tan x & x & 1 \end{vmatrix}$, then $\lim_{x \rightarrow 0} \frac{f'(x)}{x}$ [Online April 15, 2018]

- (a) Exists and is equal to -2
 (b) Does not exist
 (c) Exist and is equal to 0
 (d) Exists and is equal to 2

80. If $f(x) = \sin^{-1}\left(\frac{2 \times 3^x}{1+9^x}\right)$, then $f'\left(-\frac{1}{2}\right)$ equals.
[Online April 15, 2018]
 (a) $\sqrt{3} \log_e \sqrt{3}$ (b) $-\sqrt{3} \log_e \sqrt{3}$
 (c) $-\sqrt{3} \log_e 3$ (d) $\sqrt{3} \log_e 3$
81. If $x^2 + y^2 + \sin y = 4$, then the value of $\frac{d^2 y}{dx^2}$ at the point $(-2, 0)$ is
[Online April 15, 2018]
 (a) -34 (b) -32 (c) -2 (d) 4
82. If for $x \in \left(0, \frac{1}{4}\right)$, the derivative of $\tan^{-1}\left(\frac{6x\sqrt{x}}{1-9x^3}\right)$ is $\sqrt{x} \cdot g(x)$, then $g(x)$ equals :
[2017]
 (a) $\frac{3}{1+9x^3}$ (b) $\frac{9}{1+9x^3}$
 (c) $\frac{3x\sqrt{x}}{1-9x^3}$ (d) $\frac{3x}{1-9x^3}$
83. For $x \in \mathbb{R}$, $f(x) = |\log 2 - \sin x|$ and $g(x) = f(f(x))$, then :
[2016]
 (a) $g'(0) = -\cos(\log 2)$
 (b) g is differentiable at $x = 0$ and $g'(0) = -\sin(\log 2)$
 (c) g is not differentiable at $x = 0$
 (d) $g'(0) = \cos(\log 2)$
84. If $f(x) = x^2 - x + 5$, $x > \frac{1}{2}$, and $g(x)$ is its inverse function, then $g'(7)$ equals:
[Online April 12, 2014]
 (a) $-\frac{1}{3}$ (b) $\frac{1}{13}$ (c) $\frac{1}{3}$ (d) $-\frac{1}{13}$
85. If $y = \sec(\tan^{-1} x)$, then $\frac{dy}{dx}$ at $x = 1$ is equal to : **[2013]**
 (a) $\frac{1}{\sqrt{2}}$ (b) $\frac{1}{2}$ (c) 1 (d) $\sqrt{2}$
86. If the curves $\frac{x^2}{\alpha} + \frac{y^2}{4} = 1$ and $y^3 = 16x$ intersect at right angles, then a value of α is : **[Online April 23, 2013]**
 (a) 2 (b) $\frac{4}{3}$ (c) $\frac{1}{2}$ (d) $\frac{3}{4}$
87. For $a > 0$, $t \in \left(0, \frac{\pi}{2}\right)$, let $x = \sqrt{a^{\sin^{-1} t}}$ and $y = \sqrt{a^{\cos^{-1} t}}$,
 Then, $1 + \left(\frac{dy}{dx}\right)^2$ equals : **[Online April 22, 2013]**
 (a) $\frac{x^2}{y^2}$ (b) $\frac{y^2}{x^2}$ (c) $\frac{x^2 + y^2}{y^2}$ (d) $\frac{x^2 + y^2}{x^2}$
88. Let $f(x) = \frac{x^2 - x}{x^2 + 2x}$, $x \neq 0, -2$. Then $\frac{d}{dx}[f^{-1}(x)]$ (wherever it is defined) is equal to : **[Online April 9, 2013]**
 (a) $\frac{-1}{(1-x)^2}$ (b) $\frac{3}{(1-x)^2}$
 (c) $\frac{1}{(1-x)^2}$ (d) $\frac{-3}{(1-x)^2}$
89. If $f'(x) = \sin(\log x)$ and $y = f\left(\frac{2x+3}{3-2x}\right)$, then $\frac{dy}{dx}$ equals
[Online May 12, 2012]
 (a) $\sin\left[\log\left(\frac{2x+3}{3-2x}\right)\right]$
 (b) $\frac{12}{(3-2x)^2}$
 (c) $\frac{12}{(3-2x)^2} \sin\left[\log\left(\frac{2x+3}{3-2x}\right)\right]$
 (d) $\frac{12}{(3-2x)^2} \cos\left[\log\left(\frac{2x+3}{3-2x}\right)\right]$
90. Let $f: (-1, 1) \rightarrow \mathbb{R}$ be a differentiable function with $f(0) = -1$ and $f'(0) = 1$. Let $g(x) = [f(2f(x) + 2)]^2$. Then $g'(0) =$
[2010]
 (a) -4 (b) 0 (c) -2 (d) 4
91. Let y be an implicit function of x defined by $x^{2x} - 2x^x \cot y - 1 = 0$. Then $y'(1)$ equals
[2009]
 (a) 1 (b) $\log 2$ (c) $-\log 2$ (d) -1
92. If $x^m \cdot y^n = (x+y)^{m+n}$, then $\frac{dy}{dx}$ is
[2006]
 (a) $\frac{y}{x}$ (b) $\frac{x+y}{xy}$ (c) xy (d) $\frac{x}{y}$
93. If $x = e^{y+e^{y+\dots+\infty}}$, $x > 0$, then $\frac{dy}{dx}$ is
[2004]
 (a) $\frac{1+x}{x}$ (b) $\frac{1}{x}$ (c) $\frac{1-x}{x}$ (d) $\frac{x}{1+x}$
94. Let $f(x)$ be a polynomial function of second degree. If $f(1) = f(-1)$ and a, b, c are in A.P., then $f'(a), f'(b), f'(c)$ are in
[2003]
 (a) Arithmetic - Geometric Progression
 (b) A.P.
 (c) G.P.
 (d) H.P.
95. If $f(x+y) = f(x) \cdot f(y) \forall x, y$ and $f(5) = 2$,
 $f'(0) = 3$, then $f'(5)$ is
[2002]
 (a) 0 (b) 1 (c) 6 (d) 2



TOPIC 4

Differentiation of Infinite Series, Successive Differentiation, nth Derivative of Some Standard Functions, Leibnitz's Theorem, Rolle's Theorem, Lagrange's Mean Value Theorem



96. For all twice differentiable functions $f : \mathbb{R} \rightarrow \mathbb{R}$, with $f(0) = f(1) = f'(0) = 0$ [Sep. 06, 2020 (II)]
- (a) $f''(x) \neq 0$ at every point $x \in (0, 1)$
 - (b) $f''(x) = 0$, for some $x \in (0, 1)$
 - (c) $f''(0) = 0$
 - (d) $f''(x) = 0$, at every point $x \in (0, 1)$

97. If $y^2 + \log_e(\cos^2 x) = y$, $x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$, then : [Sep. 03, 2020 (I)]
- (a) $y''(0) = 0$
 - (b) $|y'(0)| + |y''(0)| = 1$
 - (c) $|y''(0)| = 2$
 - (d) $|y'(0)| + |y''(0)| = 3$

98. If c is a point at which Rolle's theorem holds for the function, $f(x) = \log_e\left(\frac{x^2 + a}{7x}\right)$ in the interval $[3, 4]$, where $\alpha \in \mathbb{R}$, then $f'''(c)$ is equal to: [Jan. 8, 2020 (I)]
- (a) $-\frac{1}{12}$
 - (b) $\frac{1}{12}$
 - (c) $-\frac{1}{24}$
 - (d) $\frac{\sqrt{3}}{7}$

99. Let $x^k + y^k = a^k$, ($a, k > 0$) and $\frac{dy}{dx} + \left(\frac{y}{x}\right)^{\frac{1}{3}} = 0$, then k is: [Jan. 7, 2020 (I)]
- (a) $\frac{3}{2}$
 - (b) $\frac{4}{3}$
 - (c) $\frac{2}{3}$
 - (d) $\frac{1}{3}$

100. The value of c in the Lagrange's mean value theorem for the function $f(x) = x^3 - 4x^2 + 8x + 11$, when $x \in [0, 1]$ is: [Jan. 7, 2020 (II)]
- (a) $\frac{4 - \sqrt{5}}{3}$
 - (b) $\frac{4 - \sqrt{7}}{3}$
 - (c) $\frac{2}{3}$
 - (d) $\frac{\sqrt{7} - 2}{3}$

101. If $2x = y^{\frac{1}{5}} + y^{-\frac{1}{5}}$ and $(x^2 - 1) \frac{d^2y}{dx^2} + \lambda x \frac{dy}{dx} + ky = 0$, then $\lambda + k$ is equal to: [Online April 9, 2017]
- (a) -23
 - (b) -24
 - (c) 26
 - (d) -26

102. Let f be a polynomial function such that $f(3x) = f'(x)$, $f''(x)$, for all $x \in \mathbb{R}$. Then : [Online April 9, 2017]
- (a) $f(b) + f'(b) = 28$
 - (b) $f''(b) - f'(b) = 0$
 - (c) $f''(b) - f'(b) = 4$
 - (d) $f(b) - f'(b) + f''(b) = 10$

103. If $y = \left[x + \sqrt{x^2 - 1}\right]^{15} + \left[x - \sqrt{x^2 - 1}\right]^{15}$, then $(x^2 - 1) \frac{d^2y}{dx^2} + x \frac{dy}{dx}$ is equal to [Online April 8, 2017]
- (a) $12y$
 - (b) $224y^2$
 - (c) $225y^2$
 - (d) $225y$

104. If Rolle's theorem holds for the function $f(x) = 2x^3 + bx^2 + cx$, $x \in [-1, 1]$, at the point $x = \frac{1}{2}$, then $2b + c$ equals : [Online April 10, 2015]
- (a) -3
 - (b) -1
 - (c) 2
 - (d) 1

105. If f and g are differentiable functions in $[0, 1]$ satisfying $f(0) = 2 = g(1)$, $g(0) = 0$ and $f(1) = 6$, then for some $c \in]0, 1[$ [2014]
- (a) $f'(c) = g'(c)$
 - (b) $f'(c) = 2g'(c)$
 - (c) $2f'(c) = g'(c)$
 - (d) $2f'(c) = 3g'(c)$

106. Let $f(x) = x|x|$, $g(x) = \sin x$ and $h(x) = (\text{gof})(x)$. Then [Online April 11, 2014]
- (a) $h(x)$ is not differentiable at $x = 0$.
 - (b) $h(x)$ is differentiable at $x = 0$, but $h'(x)$ is not continuous at $x = 0$
 - (c) $h'(x)$ is continuous at $x = 0$ but it is not differentiable at $x = 0$
 - (d) $h'(x)$ is differentiable at $x = 0$

107. Let for $i = 1, 2, 3$, $p_i(x)$ be a polynomial of degree 2 in x , $p_i'(x)$ and $p_i''(x)$ be the first and second order derivatives of $p_i(x)$ respectively. Let,

$$A(x) = \begin{bmatrix} p_1(x) & p_1'(x) & p_1''(x) \\ p_2(x) & p_2'(x) & p_2''(x) \\ p_3(x) & p_3'(x) & p_3''(x) \end{bmatrix}$$

- and $B(x) = [A(x)]^T A(x)$. Then determinant of $B(x)$: [Online April 11, 2014]
- (a) is a polynomial of degree 6 in x .
 - (b) is a polynomial of degree 3 in x .
 - (c) is a polynomial of degree 2 in x .
 - (d) does not depend on x .

108. If the Rolle's theorem holds for the function $f(x) = 2x^3 + ax^2 + bx$ in the interval $[-1, 1]$ for the point $c = \frac{1}{2}$, then the value of $2a + b$ is: [Online April 9, 2014]
- (a) 1
 - (b) -1
 - (c) 2
 - (d) -2

109. If $f(x) = \sin(\sin x)$ and $f''(x) + \tan x f'(x) + g(x) = 0$, then $g(x)$ is :
[Online April 23, 2013]
 (a) $\cos^2 x \cos(\sin x)$ (b) $\sin^2 x \cos(\cos x)$
 (c) $\sin^2 x \sin(\cos x)$ (d) $\cos^2 x \sin(\sin x)$
110. Consider a quadratic equation $ax^2 + bx + c = 0$, where $2a + 3b + 6c = 0$ and let $g(x) = a\frac{x^3}{3} + b\frac{x^2}{2} + cx$.
[Online May 19, 2012]
Statement 1: The quadratic equation has at least one root in the interval $(0, 1)$.
Statement 2: The Rolle's theorem is applicable to function $g(x)$ on the interval $[0, 1]$.
 (a) Statement 1 is false, Statement 2 is true.
 (b) Statement 1 is true, Statement 2 is false.
 (c) Statement 1 is true, Statement 2 is true, Statement 2 is not a correct explanation for Statement 1.
 (d) Statement 1 is true, Statement 2 is true, Statement 2 is a correct explanation for Statement 1.
111. $\frac{d^2x}{dy^2}$ equals : **[2011]**
 (a) $-\left(\frac{d^2y}{dx^2}\right)^{-1} \left(\frac{dy}{dx}\right)^{-3}$ (b) $\left(\frac{d^2y}{dx^2}\right) \left(\frac{dy}{dx}\right)^{-2}$
 (c) $-\left(\frac{d^2y}{dx^2}\right) \left(\frac{dy}{dx}\right)^{-3}$ (d) $\left(\frac{d^2y}{dx^2}\right)^{-1}$
112. Let $f(x) = x|x|$ and $g(x) = \sin x$.
Statement-1 : $g \circ f$ is differentiable at $x = 0$ and its derivative is continuous at that point.
Statement-2 : $g \circ f$ is twice differentiable at $x = 0$. **[2009]**
 (a) Statement-1 is true, Statement-2 is true; Statement-2 is not a correct explanation for Statement-1.
 (b) Statement-1 is true, Statement-2 is false.
 (c) Statement-1 is false, Statement-2 is true.
 (d) Statement-1 is true, Statement-2 is true; Statement-2 is a correct explanation for Statement-1.
113. A value of c for which conclusion of Mean Value Theorem holds for the function $f(x) = \log_e x$ on the interval $[1, 3]$ is **[2007]**
 (a) $\log_3 e$ (b) $\log_e 3$ (c) $2 \log_3 e$ (d) $\frac{1}{2} \log_3 e$
114. Let f be differentiable for all x . If $f(1) = -2$ and $f'(x) \geq 2$ for $x \in [1, 6]$, then **[2005]**
 (a) $f(6) \geq 8$ (b) $f(6) < 8$ (c) $f(6) < 5$ (d) $f(6) = 5$
115. If the equation $a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x = 0$, $a_1 \neq 0$, $n \geq 2$, has a positive root $x = \alpha$, then the equation $na_n x^{n-1} + (n-1)a_{n-1} x^{n-2} + \dots + a_1 = 0$ has a positive root, which is **[2005]**
 (a) greater than α
 (b) smaller than α
 (c) greater than or equal to α
 (d) equal to α
116. If $2a + 3b + 6c = 0$, then at least one root of the equation $ax^2 + bx + c = 0$ lies in the interval **[2004]**
 (a) $(1, 3)$ (b) $(1, 2)$ (c) $(2, 3)$ (d) $(0, 1)$
117. If $f(x) = x^n$, then the value of **[2003]**
 $f(1) - \frac{f'(1)}{1!} + \frac{f''(1)}{2!} - \frac{f'''(1)}{3!} + \dots + \frac{(-1)^n f^n(1)}{n!}$ is
 (a) 1 (b) 2^n (c) $2^n - 1$ (d) 0.
118. Let $f(a) = g(a) = k$ and their n th derivatives $f^n(a), g^n(a)$ exist and are not equal for some n . Further if
 $\lim_{x \rightarrow a} \frac{f(a)g(x) - f(x)g(a)}{g(x) - f(x)} = 4$
 then the value of k is **[2003]**
 (a) 0 (b) 4 (c) 2 (d) 1
119. If $y = (x + \sqrt{1+x^2})^n$, then $(1+x^2) \frac{d^2y}{dx^2} + x \frac{dy}{dx}$ is **[2002]**
 (a) $n^2 y$ (b) $-n^2 y$ (c) $-y$ (d) $2x^2 y$
120. If $2a + 3b + 6c = 0$, ($a, b, c \in R$) then the quadratic equation $ax^2 + bx + c = 0$ has **[2002]**
 (a) at least one root in $[0, 1]$
 (b) at least one root in $[2, 3]$
 (c) at least one root in $[4, 5]$
 (d) None of these



Hints & Solutions



1. (8) We know $[x]$ discontinuous for $x \in \mathbb{Z}$

$f(x) = x \left[\frac{x}{2} \right]$ may be discontinuous where $\frac{x}{2}$ is an

integer.

So, points of discontinuity are,

$$x = \pm 2, \pm 4, \pm 6, \pm 8 \text{ and } 0$$

but at $x = 0$

$$\lim_{x \rightarrow 0^+} f(x) = 0 = f(0) = \lim_{x \rightarrow 0^-} f(x)$$

So, $f(x)$ will be discontinuous at $x = \pm 2, \pm 4, \pm 6$ and ± 8 .

2. (d) Since, function $f(x)$ is continuous at $x = 1, 3$

$$\therefore f(1) = f(1^+)$$

$$\Rightarrow ae + be^{-1} = c \quad \dots(i)$$

$$f(3) = f(3^+)$$

$$\Rightarrow 9c = 9a + 6c \Rightarrow c = 3a \quad \dots(ii)$$

From (i) and (ii),

$$b = ae(3 - e) \quad \dots(iii)$$

$$f'(x) = \begin{cases} ae^x - be^{-x} & -1 < x < 1 \\ 2cx & 1 < x < 3 \\ 2ax + 2c & 3 < x < 4 \end{cases}$$

$$f'(0) = a - b, f'(2) = 4c$$

$$\text{Given, } f'(0) + f'(2) = e$$

$$a - b + 4c = e \quad \dots(iv)$$

From eqs. (i), (ii), (iii) and (iv),

$$a - 3ae + ae^2 + 12a = e$$

$$\Rightarrow 13a - 3ae + ae^2 = e$$

$$\Rightarrow a = \frac{e}{e^2 - 3e + 13}$$

3. (a) $\lim_{x \rightarrow 0} x \left[\frac{4}{x} \right] = A \Rightarrow \lim_{x \rightarrow 0} x \left[\frac{4}{x} - \left\{ \frac{4}{x} \right\} \right] = A$

$$\Rightarrow \lim_{x \rightarrow 0} 4 - x \left\{ \frac{4}{x} \right\} = A \Rightarrow 4 - 0 = A$$

As, $f(x) = [x^2] \sin(\pi x)$ will be discontinuous at non-integers

$$\text{And, when } x = \sqrt{A+1} \Rightarrow x = \sqrt{5},$$

which is not an integer.

Hence, $f(x)$ is discontinuous when x is equal to $\sqrt{A+1}$

4. (5) $\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \left(\frac{1}{x} \ln \left(\frac{1+3x}{1-2x} \right) \right)$

$$= \lim_{x \rightarrow 0} \left(\frac{\ln(1+3x)}{x} - \frac{\ln(1-2x)}{x} \right)$$

$$= \lim_{x \rightarrow 0} \left(\frac{3 \ln(1+3x)}{3x} - \frac{2 \ln(1-2x)}{-2x} \right)$$

$$= 3 + 2 = 5$$

$\therefore f(x)$ will be continuous

$$\therefore k = f(0) = \lim_{x \rightarrow 0} f(x) = 5$$

5. (b) Since, $f(x)$ is continuous, then

$$\lim_{x \rightarrow \frac{\pi}{4}} f(x) = f\left(\frac{\pi}{4}\right)$$

$$\lim_{x \rightarrow \frac{\pi}{4}} \frac{\sqrt{2} \cos x - 1}{\cot x - 1} = k$$

Now by L- hospital's rule

$$\lim_{x \rightarrow \frac{\pi}{4}} \frac{\sqrt{2} \sin x}{x \operatorname{cosec}^2 x} = k \Rightarrow \frac{\sqrt{2} \left(\frac{1}{\sqrt{2}} \right)}{(\sqrt{2})^2} = k \Rightarrow k = \frac{1}{2}$$

6. (a) L.H.L. $\lim_{x \rightarrow 4^-} \left[[x] - \left[\frac{x}{4} \right] \right] = 3 - 0 = 3$

$$\text{R.H.L. } \lim_{x \rightarrow 4^+} \left[[x] - \left[\frac{x}{4} \right] \right] = 4 - 1 = 3$$

$$f(4) = \left[[4] - \left[\frac{4}{4} \right] \right] = 4 - 1 = 3$$

$\therefore \text{LHL} = f(4) = \text{RHL}$

$\therefore f(x)$ is continuous at $x = 4$

7. (d) R.H.L. $\lim_{x \rightarrow 5^+} b|(x - \pi)| + 3 = (5 - \pi)b + 3$

$$f(5) = \text{L.H.L. } \lim_{x \rightarrow 5^-} a|(\pi - x)| + 1 = a(5 - \pi) + 1$$

\therefore function is continuous at $x = 5$

$\therefore \text{LHL} = \text{RHL}$

$$(5 - \pi)b + 3 = (5 - \pi)a + 1$$

$$\Rightarrow 2 = (a - b)(5 - \pi) \Rightarrow a - b = \frac{2}{5 - \pi}$$

8. (c) Given function is,

$$f(x) = \begin{cases} |x| + [x], & -1 \leq x < 1 \\ x + |x|, & 1 \leq x < 2 \\ x + [x], & 2 \leq x \leq 3 \end{cases}$$

$$= \begin{cases} -x-1, & -1 \leq x < 0 \\ x, & 0 \leq x < 1 \\ 2x, & 1 \leq x < 2 \\ x+2, & 2 \leq x < 3 \\ 6, & x = 3 \end{cases}$$

$$\Rightarrow f(-1) = 0, f(-1^+) = 0;$$

$$f(0^-) = -1, f(0) = 0, f(0^+) = 0;$$

$$f(1^-) = 1, f(1) = 2, f(1^+) = 2;$$

$$f(2^-) = 4, f(2) = 4, f(2^+) = 4;$$

$$f(3^-) = 5, f(3) = 6$$

$f(x)$ is discontinuous at $x = \{0, 1, 3\}$

Hence, $f(x)$ is discontinuous at only three points.

9. (d) Let $f(x)$ is continuous at $x = 1$, then

$$f(1^-) = f(1) = f(1^+)$$

$$\Rightarrow 5 = a + b \quad \dots(1)$$

Let $f(x)$ is continuous at $x = 3$, then

$$f(3^-) = f(3) = f(3^+)$$

$$\Rightarrow a + 3b = b + 15 \quad \dots(2)$$

Let $f(x)$ is continuous at $x = 5$, then

$$f(5^-) = f(5) = f(5^+)$$

$$\Rightarrow b + 25 = 30$$

$$\Rightarrow b = 30 - 25 = 5$$

From (1), $a = 0$

But $a = 0, b = 5$ do not satisfy equation (2)

Hence, $f(x)$ is not continuous for any values of a and b

10. (a) If the function is continuous at $x = 0$, then

$$\lim_{x \rightarrow 0} f(x) \text{ will exist and } f(0) = \lim_{x \rightarrow 0} f(x)$$

$$\text{Now, } \lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \left(\frac{1}{x} - \frac{k-1}{e^{2x}-1} \right)$$

$$= \lim_{x \rightarrow 0} \left(\frac{e^{2x}-1-kx+x}{(x)(e^{2x}-1)} \right)$$

$$= \lim_{x \rightarrow 0} \left[\frac{\left(1 + 2x + \frac{(2x)^2}{2!} + \frac{(2x)^3}{3!} + \dots \right) - 1 - kx + x}{(x) \left(\left(1 + 2x + \frac{(2x)^2}{2!} + \frac{(2x)^3}{3!} + \dots \right) - 1 \right)} \right]$$

$$= \lim_{x \rightarrow 0} \left[\frac{(3-k)x + \frac{4x^2}{2!} + \frac{8x^3}{3!} + \dots}{\left(2x^2 + \frac{4x^3}{2!} + \frac{8x^3}{3!} + \dots \right)} \right]$$

For the limit to exist, power of x in the numerator should be greater than or equal to the power of x in the denominator.

Therefore, coefficient of x in numerator is equal to zero

$$\Rightarrow 3 - k = 0$$

$$\Rightarrow k = 3$$

So the limit reduces to

$$\lim_{x \rightarrow 0} \frac{(x^2) \left(\frac{4}{2!} + \frac{8x}{3!} + \dots \right)}{(x^2) \left(2 + \frac{4x}{2!} + \frac{8x^2}{3!} + \dots \right)}$$

$$= \lim_{x \rightarrow 0} \frac{\frac{4}{2!} + \frac{8x}{3!} + \dots}{2 + \frac{4x}{2!} + \frac{8x^2}{3!} + \dots} = 1$$

Hence, $f(0) = 1$

11. (c) Since $f(x)$ is continuous at $x = 2$.

$$\therefore \lim_{x \rightarrow 2} f(x) = f(2)$$

$$\Rightarrow \lim_{x \rightarrow 2} (x-1)^{2-x} = k \quad (1^\infty \text{ form})$$

$$\therefore e^l = k$$

$$\text{where } l = \lim_{x \rightarrow 2} (x-1-1) \times \frac{1}{2-x} = \lim_{x \rightarrow 2} \frac{x-2}{2-x}$$

$$= \lim_{x \rightarrow 2} \left(\frac{x-2}{x-2} \right)$$

$$\Rightarrow k = e^{-1}$$

12. (c) $\lim_{x \rightarrow \pi/2} f(x) = f(\pi/2)$

$$\Rightarrow k + 2/5 = 1 \Rightarrow k = 1 - \frac{2}{5} \Rightarrow k = \frac{3}{5}$$

$$13. (c) \frac{\frac{2x^2}{a} \quad \frac{2b^2-4b}{a} \quad \frac{1}{x^3}}{0 \quad 1 \quad \sqrt{2}}$$

Continuity at $x = 1$

$$\frac{2}{a} = a \Rightarrow a = \pm\sqrt{2}$$

Continuity at $x = \sqrt{2}$ $a = \sqrt{2}$

$$a = \frac{2b^2-4b}{2\sqrt{2}}$$

Put $a = \sqrt{2}$

$$2 = b^2 - 2b \Rightarrow b^2 - 2b - 2 = 0$$

$$b = \frac{2 \pm \sqrt{4+4 \cdot 2}}{2} = 1 \pm \sqrt{3}$$

So, $(a, b) = (\sqrt{2}, 1 - \sqrt{3})$

14. (c) Since $f(x)$ is a continuous function therefore limit of $f(x)$ at $x \rightarrow 0 =$ value of $f(x)$ at 0.

$$\begin{aligned} \therefore \lim_{x \rightarrow 0} f(x) &= \lim_{x \rightarrow 0} \frac{(e^x - 1)^2}{\sin\left(\frac{x}{k}\right) \log\left(1 + \frac{x}{4}\right)} \\ &= \lim_{x \rightarrow 0} \frac{x^2 \left(\frac{e^x - 1}{x}\right)^2}{\frac{x}{R} \left[\frac{\sin\left(\frac{x}{R}\right)}{\frac{x}{R}}\right] \cdot \frac{\log\left(1 + \frac{x}{4}\right)}{\left(\frac{x}{4}\right)}} \times \left(\frac{x}{4}\right) \\ &= \lim_{x \rightarrow 0} \frac{x^2 \left(\frac{e^x - 1}{x}\right)^2 \cdot 4k}{\frac{x}{k} \cdot \frac{x}{4}} \end{aligned}$$

on applying limit we get
 $4k = 12 \Rightarrow k = 3$

15. (d) Since $f(x) = \frac{\sqrt{2 + \cos x} - 1}{(\pi - x)^2}$ is
 Continuous at $x = \pi$
 \therefore L.H.L = R.H.L = $f(\pi)$
 Let $(\pi - x) = \theta, \theta \rightarrow 0$ when $x \rightarrow \pi$

$$\begin{aligned} \therefore \lim_{\theta \rightarrow 0} \frac{\sqrt{2 - \cos \theta} - 1}{\theta^2} \\ &= \lim_{\theta \rightarrow 0} \frac{(2 - \cos \theta) - 1}{\theta^2} \times \frac{1}{\sqrt{2 - \cos \theta} + 1} \\ &= \lim_{\theta \rightarrow 0} \frac{1 - \cos \theta}{\theta^2} \cdot \frac{1}{2} \quad (\because \cos 0 = 1) \\ &= \frac{1}{2} \lim_{\theta \rightarrow 0} \frac{2 \sin^2 \theta / 2}{\theta^2} = \frac{2}{2} \lim_{\theta \rightarrow 0} \frac{\sin^2 \theta / 2}{\frac{\theta^2}{4}} \\ &= \frac{1}{4} \left(\because \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \right) \end{aligned}$$

16. (b) Given that $f\left(\frac{9}{2}\right) = \frac{2}{9}$

$$\begin{aligned} \lim_{x \rightarrow 0} f\left(\frac{1 - \cos 3x}{x^2}\right) &= \lim_{x \rightarrow 0} \left(\frac{x^2}{1 - \cos 3x}\right) \\ &= \lim_{x \rightarrow 0} \left(\frac{x^2}{2 \sin^2 \frac{3x}{2}}\right) = \frac{1}{2} \lim_{x \rightarrow 0} \left(\frac{\frac{9}{4} \cdot x^2 \cdot \frac{4}{9}}{\sin^2 \frac{3x}{2}}\right) \end{aligned}$$

$$\begin{aligned} &= \frac{4}{9 \times 2} \lim_{x \rightarrow 0} \left(\frac{1}{\frac{\sin^2 \frac{3x}{2}}{\left(\frac{3x}{2}\right)^2}}\right) \\ &= \frac{2}{9} \left[\frac{\lim_{x \rightarrow 0} 1}{\lim_{x \rightarrow 0} \frac{\sin^2 \frac{3x}{2}}{\left(\frac{3x}{2}\right)^2}} \right] \left\{ \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \right\} \\ &= \frac{2}{9} \cdot \left[\frac{1}{1} \right] = \frac{2}{9} \end{aligned}$$

17. (a) Let $f(x) = [x] + |1 - x|, -1 \leq x \leq 3$
 where $[x] =$ greatest integer function.
 f is not continuous at $x = 0, 1, 2, 3$
 But in statement-2 $f(x)$ is continuous at $x = 3$.
 Hence, statement-1 is true and 2 is false.

18. (b) $\mu(x) = \frac{1}{x-1}$, which is discontinuous at $x = 1$

$$f(u) = \frac{1}{u^2 + u - 2} = \frac{1}{(u+2)(u-1)},$$

which is discontinuous at $u = -2, 1$

when $u = -2$, then $\frac{1}{x-1} = -2 \Rightarrow x = \frac{1}{2}$

when $u = 1$, then $\frac{1}{x-1} = 1 \Rightarrow x = 2$

Hence given composite function is discontinuous at three points, $x = 1, \frac{1}{2}$ and 2.

19. (d) $f \circ g = f(g(x)) = f(1 - |x|)$
 $= -1 + |1 - |x|| - 2|$
 $= -1 + |-|x| - 1| = -1 + ||x| + 1|$

Let $f \circ g = y$

$$\therefore y = -1 + ||x| + 1|$$

$$\Rightarrow y = \begin{cases} -1 + x + 1, & x \geq 0 \\ -1 - x + 1, & x < 0 \end{cases}$$

$$\Rightarrow y = \begin{cases} x, & x \geq 0 \\ -x, & x < 0 \end{cases}$$

LHL at $(x=0) = \lim_{x \rightarrow 0} (-x) = 0$

RHL at $(x=0) = \lim_{x \rightarrow 0} (x) = 0$

When $x = 0$, then $y = 0$

Hence, LHL at $(x=0) =$ RHL at $(x=0)$
 $=$ value of y at $(x=0)$

Hence y is continuous at $x = 0$.

Clearly at all other point y continuous. Therefore, the set of all points where $f \circ g$ is discontinuous is an empty set.

20. (a) Let $f(x) = [x] \cos\left(\frac{2x-1}{2}\right)$

We know that $[x]$ is discontinuous at all integral points and $\cos x$ is continuous at $x \in \mathbb{R}$.

So, check at $x = n, n \in I$

$$\begin{aligned} \text{L.H.L} &= \lim_{x \rightarrow n^-} [x] \cos\left(\frac{2x-1}{2}\right) \pi \\ &= (n-1) \cos\left(\frac{2n-1}{2}\right) \pi = 0 \end{aligned}$$

($\because [x]$ is the greatest integer function)

$$\begin{aligned} \text{R.H.L} &= \lim_{x \rightarrow n^+} [x] \cos\left(\frac{2x-1}{2}\right) \pi \\ &= n \cos\left(\frac{2n-1}{2}\right) \pi = 0 \end{aligned}$$

Now, value of the function at $x = n$ is

$$f(n) = 0$$

Since, L.H.L = R.H.L = $f(n)$

$\therefore f(x) = [x] \cos\left(\frac{2x-1}{2}\right)$ is continuous for every real x .

21. (d) Consider $\frac{x}{[x]} \leq f(x) \leq \sqrt{6-x}$

$$\Rightarrow \lim_{x \rightarrow 2^-} \frac{x}{[x]} = \frac{2}{1} = 2$$

$$\Rightarrow \lim_{x \rightarrow 2^-} \sqrt{6-x} = 2$$

$$\therefore \lim_{x \rightarrow 2^-} f(x) = 2 \quad [\text{By Sandwich theorem}]$$

$$\text{Now } \lim_{x \rightarrow 2^+} \frac{x}{[x]} = 1, \quad \lim_{x \rightarrow 2^+} \sqrt{6-x} = 2$$

Hence by Sandwich theorem $\lim_{x \rightarrow 2^+} f(x)$ does not exist.

Therefore f is not continuous at $x = 2$. Thus statement-1 is true but statement-2 is not true

22. (d) Statement - 1 is true.

It is the definition of continuity.

Statement - 2 is false.

23. (c) Given that $f(x) = \begin{cases} x \sin(1/x), & x \neq 0 \\ 0, & x = 0 \end{cases}$

At $x = 0$

$$\begin{aligned} \text{LHL} &= \lim_{h \rightarrow 0^-} \left\{ -h \sin\left(-\frac{1}{h}\right) \right\} \\ &= 0 \times \text{a finite quantity between } -1 \\ &\text{and } 1 = 0 \end{aligned}$$

$$\text{RHL} = \lim_{h \rightarrow 0^+} h \sin \frac{1}{h} = 0$$

Also, $f(0) = 0$

Thus LHL = RHL = $f(0)$

$\therefore f(x)$ is continuous on \mathbb{R} .

but $f_2(x)$ is not continuous at $x = 0$

24. (b) $L.H.L = \lim_{x \rightarrow 0^-} f(x)$

$$= \lim_{h \rightarrow 0} \frac{\sin\{(p+1)(-h)\} - \sinh}{-h} = p+1+1 = p+2$$

$$R.H.L = \lim_{x \rightarrow 0^+} f(x)$$

$$= \lim_{h \rightarrow 0} \frac{\sqrt{x+x^2} - \sqrt{x}}{x^{3/2}} \times \frac{\sqrt{x+x^2} + \sqrt{x}}{\sqrt{x+x^2} + \sqrt{x}} = \frac{1}{1+1} = \frac{1}{2}$$

$$f(0) = 2$$

Given that $f(x)$ is continuous at $x = 0$

$$\therefore p+2 = q = \frac{1}{2}$$

$$\Rightarrow p = -\frac{3}{2}, q = \frac{1}{2}$$

25. (b) Given, $f(x) = \frac{1}{x} - \frac{2}{e^{2x}-1}$ is continuous at $x = 0$

$$\Rightarrow f(0) = \lim_{x \rightarrow 0} \frac{1}{x} - \frac{2}{e^{2x}-1}$$

$$= \lim_{x \rightarrow 0} \frac{(e^{2x}-1) - 2x}{x(e^{2x}-1)}; \left[\frac{0}{0} \text{ form} \right]$$

\therefore Applying, L'Hospital rule

Differentiate two times, we get

$$f(0) = \lim_{x \rightarrow 0} \frac{4e^{2x}}{2(xe^{2x} + e^{2x} \cdot 1) + e^{2x} \cdot 2}$$

$$= \lim_{x \rightarrow 0} \frac{4e^{2x}}{4xe^{2x} + 2e^{2x} + 2e^{2x}} \left[\frac{0}{0} \text{ form} \right]$$

$$= \lim_{x \rightarrow 0} \frac{4e^{2x}}{4(xe^{2x} + e^{2x})} = \frac{4 \cdot e^0}{4(0 + e^0)} = 1$$

26. (c) Given that $f(x) = \frac{1 - \tan x}{4x - \pi}$ is continuous in $\left[0, \frac{\pi}{2}\right]$

$$\therefore f\left(\frac{\pi}{4}\right) = \lim_{x \rightarrow \frac{\pi}{4}^-} f(x) = \lim_{x \rightarrow \frac{\pi}{4}^+} f(x)$$

$$\lim_{x \rightarrow \frac{\pi}{4}^-} f(x) = \lim_{h \rightarrow 0} f\left(\frac{\pi}{4} + h\right)$$

$$= \lim_{h \rightarrow 0} \frac{1 - \tan\left(\frac{\pi}{4} + h\right)}{4\left(\frac{\pi}{4} + h\right) - \pi}, h > 0 = \lim_{h \rightarrow 0} \frac{1 - \frac{1 + \tan h}{1 - \tan h}}{4h}$$

$$= \lim_{h \rightarrow 0} \frac{-2}{1 - \tan h} \cdot \frac{\tan h}{4h} = \frac{-2}{4} = -\frac{1}{2} \left[\because \lim_{\theta \rightarrow 0} \frac{\tan \theta}{\theta} = 1 \right]$$

27. (b) Let a is a rational number other than 0, in $[-5, 5]$,

then $f(a) = a$ and $\lim_{x \rightarrow a} f(x) = -a$

$\therefore x \rightarrow a^-$ and $x \rightarrow a^+$ is tends to irrational number

$\therefore f(x)$ is discontinuous at any rational number

If a is irrational number, then


$$f(a) = -a \text{ and } \lim_{x \rightarrow a} f(x) = a$$

$\therefore f(x)$ is not continuous at any irrational number. For $x=0$,

$$\lim_{x \rightarrow 0} f(x) = f(0) = 0$$

$\therefore f(x)$ is continuous at $x=0$

28. (a)



$$f(x) = \max \{x, x^2\}$$

$$\Rightarrow f(x) = \begin{cases} x^2, & x < 0 \\ x, & 0 \leq x < 1 \\ x^2, & x \geq 1 \end{cases}$$

$\therefore f(x)$ is not differentiable at $x=0, 1$

29. (a) $f(x)$ is differentiable then, $f(x)$ is also continuous.

$$\therefore \lim_{x \rightarrow \pi^+} f(x) = \lim_{x \rightarrow \pi^-} f(x) = f(\pi)$$

$$\Rightarrow -1 = -K_2 \Rightarrow K_2 = 1$$

$$\therefore f'(x) = \begin{cases} 2K_1(x - \pi) & ; x \leq \pi \\ -K_2 \sin x & ; x > \pi \end{cases}$$

$$\text{Then, } \lim_{x \rightarrow \pi^+} f(x) = \lim_{x \rightarrow \pi^-} f(x) = 0$$

$$f''(x) = \begin{cases} 2K_1 & ; x \leq \pi \\ -K_2 \cos x & ; x > \pi \end{cases}$$

$$\text{Then, } \lim_{x \rightarrow \pi^+} f(x) = \lim_{x \rightarrow \pi^-} f(x)$$

$$\Rightarrow 2K_1 = K_2 \Rightarrow K_1 = \frac{1}{2}$$

$$\text{So, } (K_1, K_2) = \left(\frac{1}{2}, 1\right)$$

30. (b) Let f be twice differentiable function

$$\therefore f'(x) \geq 1$$

$$\Rightarrow \frac{f(5) - f(2)}{3} \geq 1$$

$$\Rightarrow f(5) \geq 3 + f(2)$$

$$\Rightarrow f(5) \geq 3 + 8 \Rightarrow f(5) \geq 11$$

and also $f''(x) \geq 4$

$$\Rightarrow \frac{f'(5) - f'(2)}{5 - 2} \geq 4 \Rightarrow f'(5) \geq 12 + f'(2)$$

$$\Rightarrow f'(5) \geq 17$$

Hence, $f(5) + f'(5) \geq 28$

31. (10.00)

$$f(x+y) = f(x) + f(y) + xy^2 + x^2y$$

Differentiate w.r.t. x :

$$f'(x+y) = f'(x) + 0 + y^2 + 2xy$$

Put $y = -x$

$$f'(0) = f'(x) + x^2 - 2x^2 \quad \dots(i)$$


$$\therefore \lim_{x \rightarrow 0} \frac{f(x)}{x} = 1 \Rightarrow f(0) = 0$$

$$\therefore f'(0) = 1 \quad \dots(ii)$$

From equations (i) and (ii),

$$f'(x) = (x^2 + 1) \Rightarrow f'(3) = 10.$$

32. (a) $f(x) = \begin{cases} \frac{-x-1}{2}, & x < -1 \\ \frac{\pi}{4} + \tan^{-1} x, & -1 \leq x \leq 1 \\ \frac{1}{2}(x-1), & x > 1 \end{cases}$



It is clear from above graph that,

$f(x)$ is discontinuous at $x = 1$.

i.e. continuous on $R - \{1\}$

$f(x)$ is non-differentiable at $x = -1, 1$

i.e. differentiable on $R - \{-1, 1\}$.

33. (c) LHL = $\lim_{x \rightarrow 0} \frac{\sin(a+2)x + \sin x}{x}$

$$= \lim_{x \rightarrow 0} \left(\frac{\sin(a+2)x}{(a+2)x} \right) (a+2) + \lim_{x \rightarrow 0} \frac{\sin x}{x} = a + 3$$

$$f(0) = b$$

$$\text{RHL} = \lim_{h \rightarrow 0^+} \left[\frac{(1+3h)^{\frac{1}{3}} - 1}{h} \right] = 1$$

\therefore Function $f(x)$ is continuous

$$\therefore \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x) = f(0)$$

$$\therefore a + 3 = 1 \Rightarrow a = -2$$

and $b = 1$

$$\text{Hence, } a + 2b = 0$$

34. (a) It is given that functions f and g are differentiable and $f \circ g$ is identity function.

$$\therefore (f \circ g)(x) = x \Rightarrow f(g(x)) = x$$

Differentiating both sides, we get

$$f'(g(x)) \cdot g'(x) = 1$$

Now, put $x = a$, then

$$f'(g(a)) \cdot g'(a) = 1$$

$$f'(b) \cdot 5 = 1$$

$$f'(b) = \frac{1}{5}$$

35. (Bonus) For a constant function $f(x)$, option (1), (3) and (4) doesn't hold and by LMVT theorem, option (2) is incorrect.

36. (a) From, LMVT for $x \in [-7, -1]$

$$\frac{f(-1) - f(-7)}{(-1) - (-7)} \leq 2 \Rightarrow \frac{f(-1) + 3}{6} \leq 2 \Rightarrow f(-1) \leq 9$$

From, LMVT for $x \in [-7, 0]$

$$\frac{f(0) - f(-7)}{(0) - (-7)} \leq 2$$

$$\frac{f(0) + 3}{7} \leq 2 \Rightarrow f(0) \leq 11$$

$$\therefore f(0) + f(-1) \leq 20$$

37. (c) $\therefore f(x)$ is non differentiable at $x = 1, 3, 5$

[$\because |x-3|$ is not differentiable at $x = 3$]

$$\begin{aligned} \Sigma f(x) &= f(1) + f(3) + f(5) \\ &= 1 + 1 + 1 = 3 \end{aligned}$$

$$38. \text{ (c) } f(x) = \begin{cases} \frac{\sin(p+1)x + \sin x}{x} & x < 0 \\ q & x = 0 \text{ is continuous at } x = 0 \\ \frac{\sqrt{x^2 + x} - \sqrt{x}}{\frac{3}{x^2}} & x > 0 \end{cases}$$

Therefore, $f(0^-) = f(0) = f(0^+) \dots(1)$

$$f(0^-) = \lim_{h \rightarrow 0^-} f(0-h) = \lim_{h \rightarrow 0^-} \frac{\sin(p+1)(-h) + \sin(-h)}{-h}$$

$$= \lim_{h \rightarrow 0^-} \left[\frac{-\sin(p+1)h}{-h} + \frac{\sin h}{h} \right]$$

$$= \lim_{h \rightarrow 0^-} \frac{\sin(p+1)h}{h(p+1)} \times (p+1) + \lim_{h \rightarrow 0^-} \frac{\sin h}{h}$$

$$= (p+1) + 1 = p+2 \quad \dots(2)$$

$$\text{And } f(0^+) = \lim_{h \rightarrow 0^+} f(0+h) = \frac{\sqrt{h^2+h} - \sqrt{h}}{h^{3/2}}$$

$$= \lim_{h \rightarrow 0^+} \frac{(h)^{\frac{1}{2}} \left[\sqrt{h+1} - 1 \right]}{h \left(\frac{1}{h^2} \right)}$$

$$= \lim_{h \rightarrow 0^+} \frac{\sqrt{h+1} - 1}{h} \times \frac{\sqrt{h+1} + 1}{\sqrt{h+1} + 1} = \lim_{h \rightarrow 0^+} \frac{h+1-1}{h(\sqrt{h+1}+1)}$$

$$= \lim_{h \rightarrow 0^+} \frac{1}{\sqrt{h+1}+1} = \frac{1}{1+1} = \frac{1}{2} \quad \dots(3)$$

Now, from equation (1),

$$f(0^-) = f(0) = f(0^+) \Rightarrow p+2 = q = \frac{1}{2}$$

$$\Rightarrow q = \frac{1}{2} \text{ and } p = \frac{-3}{2} \quad \therefore (p, q) = \left(-\frac{3}{2}, \frac{1}{2} \right)$$

39. (b) $f(x) = \ln(\sin x)$, $g(x) = \sin^{-1}(e^{-x})$

$$\Rightarrow f(g(x)) = \ln(\sin(\sin^{-1}(e^{-x}))) = -x$$

$$\Rightarrow f'(g(x)) = -\alpha$$

But given that $(f \circ g)(\alpha) = b$

$$\therefore -\alpha = b \text{ and } f'(g(\alpha)) = a, \text{ i.e., } a = -1$$

$$\therefore a\alpha^2 - b\alpha - a = -\alpha^2 + \alpha^2 - (-1)$$

$$\Rightarrow a\alpha^2 - b\alpha - a = 1.$$

40. (c) $g'(c) = \lim_{x \rightarrow c} \frac{g(x) - g(c)}{x - c}$

$$\Rightarrow g'(c) = \lim_{x \rightarrow c} \frac{|f(x)| - |f(c)|}{x - c}$$

Since, $f(c) = 0$

$$\text{Then, } g'(c) = \lim_{x \rightarrow c} \frac{|f(x)|}{x - c}$$

$$\Rightarrow g'(c) = \lim_{x \rightarrow c} \frac{f(x)}{x - c}; \text{ if } f(x) > 0$$

$$\text{and } g'(c) = \lim_{x \rightarrow c} \frac{-f(x)}{x - c}; \text{ if } f(x) < 0$$

$$\Rightarrow g'(c) = f'(c) = -f'(c)$$

$$\Rightarrow 2f'(c) = 0 \Rightarrow f'(c) = 0$$

Hence, $g(x)$ is differentiable if $f'(c) = 0$

41. (a) Since, $f(x) = 15 - |(10 - x)|$
 $\therefore g(x) = f(f(x)) = 15 - |10 - [15 - |10 - x||]$
 $= 15 - ||10 - x| - 5|$
 \therefore Then, the points where function $g(x)$ is Non-differentiable are
 $10 - x = 0$ and $|10 - x| = 5$
 $\Rightarrow x = 10$ and $x - 10 = \pm 5$
 $\Rightarrow x = 10$ and $x = 15, 5$

42. (a) Let $g(x) = f(f(f(x))) + (f(x))^2$
 Differentiating both sides w.r.t. x , we get
 $g'(x) = f'(f(f(x)))f'(f(x))f'(x) + 2f(x)f'(x)$
 $g'(1) = f'(f(f(1)))f'(f(1))f'(1) + 2f(1)f'(1)$
 $= f'(f(1))f'(1)f'(1) + 2f(1)f'(1)$
 $= 3 \times 3 \times 3 + 2 \times 1 \times 3 = 27 + 6 = 33$

43. (b) Since, $f'(x) = f(x)$

Then, $\frac{f'(x)}{f(x)} = 1$

$\Rightarrow \frac{f'(x)}{f(x)} = dx \Rightarrow \frac{f'(x)}{f(x)} dx = [dx$

$\Rightarrow \ln |f(x)| = x + c$
 $f(x) = \pm e^{x+c} \dots(1)$

Since, the given condition

$f(1) = 2$

From eqⁿ (1) $f(x) = e^{x+c} = e^c e^x$

Then, $f(1) = e^c \cdot e^1$

$\Rightarrow 2 = e^c \cdot e$

$\Rightarrow \frac{2}{e} = e^c$

Then, from eqⁿ (1)

$f(x) = \frac{2}{e} e^x$

$\Rightarrow f'(x) = \frac{2}{e} e^x$

Now $h(x) = f(f(x))$

$\Rightarrow h'(x) = f'(f(x)) \cdot f'(x)$

$h'(1) = f'(2) \cdot f'(1) = \frac{2}{e} e^2 \cdot \frac{2}{e} \cdot e = 4e$

44. (d) $f(x) = \begin{cases} -1, & -2 \leq x < 0 \\ x^2 - 1, & 0 \leq x \leq 2 \end{cases}$

Then, $f(|x|) = \begin{cases} -1, & -2 \leq |x| < 0 \\ |x|^2 - 1, & 0 \leq |x| \leq 2 \end{cases}$

$\Rightarrow f(|x|) = x^2 - 1, -2 \leq x \leq 2$

$\Rightarrow g(x) = \begin{cases} 1 + x^2 - 1, & -2 \leq x < 0 \\ (x^2 - 1) + |x^2 - 1|, & 0 \leq x \leq 2 \end{cases}$

$= \begin{cases} x^2, & -2 \leq x < 0 \\ 0, & 0 \leq x < 1 \\ 2(x^2 - 1), & 1 \leq x \leq 2 \end{cases}$

$g'(0^-) = 0, g'(0^+) = 0, g'(1^-) = 0, g'(1^+) = 4$

$\Rightarrow g(x)$ is non-differentiable at $x = 1$

$\Rightarrow g(x)$ is not differentiable at one point.

45. (b) Consider the equation,

$x \log_e (\log_e x) - x^2 + y^2 = 4$

Differentiate both sides w.r.t. x ,

$\log_e (\log_e x) + x \cdot \frac{1}{x \cdot \log_e x} - 2x + 2y \frac{dy}{dx} = 0$

$\log_e (\log_e x) + \frac{1}{\log_e x} - 2x + 2y \frac{dy}{dx} = 0 \dots(1)$

When $x = e, y = \sqrt{4 + e^2}$. Put these values in (1),

$0 + 1 - 2e + 2\sqrt{4 + e^2} \frac{dy}{dx} = 0$

$\frac{dy}{dx} = \frac{2e - 1}{2\sqrt{4 + e^2}}$

46. (a) $f(x) = \sin |x| - |x| + 2(x - \pi) \cos |x|$

There are two cases,

Case (1), $x > 0$

$f(x) = \sin x - x + 2(x - \pi) \cos x$

$f'(x) = \cos x - 1 + 2(1 - 0) \cos x - 2 \sin (x - \pi)$

$f'(x) = 3 \cos x - 2(x - \pi) \sin x - 1$

Then, function $f(x)$ is differentiable for all $x > 0$

Case (2) $x < 0$

$f(x) = -\sin x + x + 2(x - \pi) \cos x$

$f'(x) = -\cos x + 1 - 2(x - \pi) \sin x + 2 \cos x$

$f'(x) = \cos x + 1 - 2(x - \pi) \sin x$

Then, function $f(x)$ is differentiable for all $x < 0$

Now check for $x = 0$

$f'(0^+) \text{ R.H.D.} = 3 - 1 = 2$


$f'(0^-) \text{ L.H.D.} = 1 + 1 = 2$

L.H.D. = R.H.D.

Then, function $f(x)$ is differentiable for $x = 0$. So it is differentiable everywhere

Hence, $k = \phi$

47. (b) Given $f(x) = \begin{cases} \max\{|x|, x^2\} & |x| \leq 2 \\ 8 - 2|x| & 2 < |x| \leq 4 \end{cases}$




∴ $f(x)$ is not differentiable at $-2, -1, 0, 1$ and 2 .

∴ $S = \{-2, -1, 0, 1, 2\}$

48. (c) Consider the function

$f(x) = \max\{-|x|, -\sqrt{1-x^2}\}$

Now, the graph of the function



From the graph, it is clear that $f(x)$ is not differentiable at $x = \pm \frac{1}{\sqrt{2}}$

$= 0, -\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}$

Then, $K = \left\{-\frac{1}{\sqrt{2}}, 0, \frac{1}{\sqrt{2}}\right\}$

Hence, K has exactly three elements.

49. (d) $f(x) = |x - \pi|(e^{|x|} - 1)\sin|x|$

Check differentiability of $f(x)$ at $x = \pi$ and $x = 0$

at $x = \pi$:

R.H.D. = $\lim_{h \rightarrow 0} \frac{|\pi + h - \pi|(e^{|\pi+h|} - 1)\sin|\pi + h| - 0}{h}$

L.H.D. = $\lim_{h \rightarrow 0} \frac{|\pi - h - \pi|(e^{|\pi-h|} - 1)\sin|\pi - h| - 0}{-h} = 0$

∴ RHD = LHD

Therefore, function is differentiable at $x = \pi$

at $x = 0$:

R.H.D. = $\lim_{h \rightarrow 0} \frac{|h - \pi|(e^{|h|} - 1)\sin|h| - 0}{h} = 0$

L.H.D. = $\lim_{h \rightarrow 0} \frac{|-h - \pi|(e^{-|h|} - 1)\sin|-h| - 0}{-h} = 0$

∴ RHD = LHD

Therefore, function is differentiable.

at $x = 0$.

Since, the function $f(x)$ is differentiable at all the points including π and 0 .

i.e., $f(x)$ is every where differentiable.

Therefore, there is no element in the set S .

⇒ $S = \phi$ (an empty set)

50. (a) $S = \{(\lambda, \mu) \in \mathbb{R} \times \mathbb{R} : f(t) = (|\lambda|e^{|t|} - \mu)\sin(2|t|), t \in \mathbb{R}$
 $f(t) = (|\lambda|e^{|t|} - \mu)\sin(2|t|)$

$= \begin{cases} (|\lambda|e^t - \mu)\sin 2t, & t > 0 \\ (|\lambda|e^{-t} - \mu)(-\sin 2t), & t < 0 \end{cases}$

$f'(t) = \begin{cases} (|\lambda|e^t)\sin 2t + (|\lambda|e^t - \mu)(2\cos 2t), & t > 0 \\ | \lambda | e^{-t} \sin 2t + (|\lambda|e^{-t} - \mu)(-2\cos 2t), & t < 0 \end{cases}$

As, $f(t)$ is differentiable

∴ LHD = RHD at $t = 0$

⇒ $|\lambda| \cdot \sin 2(0) + (|\lambda|e^0 - \mu)2\cos(0)$
 $= |\lambda|e^0 \cdot \sin 2(0) - 2\cos(0)(|\lambda|e^0 - \mu)$

⇒ $0 + (|\lambda| - \mu)2 = 0 - 2(|\lambda| - \mu)$

⇒ $4(|\lambda| - \mu) = 0$

⇒ $|\lambda| = \mu$

So, $S = \{(\lambda, \mu) = \{\lambda \in \mathbb{R} \ \& \ \mu \in [0, \infty)\}$

Therefore set S is subset of $\mathbb{R} \times [0, \infty)$

51. (a) $f(x) = \begin{cases} -x & x < 1 \\ a + \cos^{-1}(x+b) & 1 \leq x \leq 2 \end{cases}$

$f(x)$ is continuous

⇒ $\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} a + \cos^{-1}(x+b) = f(x)$

⇒ $-1 = a + \cos^{-1}(1+b)$

$\cos^{-1}(1+b) = -1 - a$ (a)

$f(x)$ is differentiable

⇒ LHD = RHD

⇒ $-1 = \frac{-1}{\sqrt{1-(1+b)^2}}$

⇒ $1 - (1+b)^2 = 1 \Rightarrow b = -1$ (b)

From (a) ⇒ $\cos^{-1}(0) = -1 - a$

∴ $-1 - a = \frac{\pi}{2}$

$a = -1 - \frac{\pi}{2} \Rightarrow a = \frac{-\pi - 2}{2}$ (c)

∴ $\frac{a}{b} = \frac{\pi + 2}{2}$

52. (c) Since $g(x)$ is differentiable, it will be continuous at $x=3$

$$\therefore \lim_{x \rightarrow 3^-} g(x) = \lim_{x \rightarrow 3^+} g(x)$$

$$2k = 3m + 2 \quad \dots(1)$$

Also $g(x)$ is differentiable at $x = 0$

$$\therefore \lim_{x \rightarrow 3^-} g'(x) = \lim_{x \rightarrow 3^+} g'(x)$$

$$\frac{k}{2\sqrt{3+1}} = m$$

$$k = 4m \quad \dots(2)$$

Solving (1) and (2), we get

$$m = \frac{2}{5}, \quad k = \frac{8}{5}$$

$$k + m = 2$$

53. (b) Let $|f(x)| \leq x^2, \forall x \in R$

Now, at $x = 0, |f(0)| \leq 0$

$$\Rightarrow f(0) = 0$$

$$\therefore f'(0) = \lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h - 0} = \lim_{h \rightarrow 0} \frac{f(h)}{h} \quad \dots(1)$$

$$\text{Now, } \left| \frac{f(h)}{h} \right| \leq |h| \quad (\because |f(x)| \leq x^2)$$

$$\Rightarrow -|h| \leq \frac{f(h)}{h} \leq |h|$$

$$\Rightarrow \lim_{h \rightarrow 0} \frac{f(h)}{h} \rightarrow 0 \quad \dots(2)$$

(using sandwich Theorem)

\therefore from (1) and (2), we get $f'(0) = 0$,

i.e. $-f(x)$ is differentiable, at $x = 0$

Since, differentiability \Rightarrow Continuity

$\therefore |f(x)| \leq x^2$, for all $x \in R$ is continuous as well as differentiable at $x = 0$.

$$54. (b) f(x) = \begin{cases} x \sin\left(\frac{1}{x}\right), & x \neq 0 \\ 0, & x = 0 \end{cases}$$

and $g(x) = x f(x)$

For $f(x)$

$$\text{LHL} = \lim_{h \rightarrow 0^-} \left\{ -h \sin\left(-\frac{1}{h}\right) \right\}$$

$= 0 \times$ a finite quantity between -1 and $1 = 0$

$$\text{RHL} = \lim_{h \rightarrow 0^+} h \sin\left(\frac{1}{h}\right) = 0$$

Also, $f(0) = 0$

Thus $\text{LHL} = \text{RHL} = f(0)$

$\therefore f(x)$ is continuous at $x = 0$

$$g(x) = \begin{cases} x^2 \sin \frac{1}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$$

For $g(x)$

$$\text{LHL} = \lim_{h \rightarrow 0^-} \left\{ -h^2 \sin\left(\frac{1}{h}\right) \right\}$$

$= 0^2 \times$ a finite quantity between -1 and $1 = 0$

$$\text{RHL} = \lim_{h \rightarrow 0^+} h^2 \sin\left(\frac{1}{h}\right) = 0$$

Also $g(0) = 0$

$\therefore g(x)$ is continuous at $x = 0$

$$55. (c) f(x) = |x-2| = \begin{cases} x-2, & x-2 \geq 0 \\ 2-x, & x-2 \leq 0 \end{cases}$$

$$= \begin{cases} x-2, & x \geq 2 \\ 2-x, & x \leq 2 \end{cases}$$

Similarly,

$$f(x) = |x-5| = \begin{cases} x-5, & x \geq 5 \\ 5-x, & x \leq 5 \end{cases}$$

$$\therefore f(x) = |x-2| + |x-5|$$


$$= \{x-2+5-x=3, 2 \leq x \leq 5\}$$

Thus $f(x) = 3, 2 \leq x \leq 5$

$$f'(x) = 0, 2 < x < 5$$

$$f'(4) = 0$$

\therefore Statement-1 is true



Since $f(x) = 3, 2 \leq x \leq 5$ is constant function.

So, it is continuous in $2, 5$ and differentiable in $(2, 5)$

$$\therefore f(2) = 0 + |2-5| = 3$$

and $f(5) = |5-2| + 0 = 3$ statement-2 is also true.

56. (d) $|\sin x|$ and $e^{|x|}$ are not differentiable at $x = 0$ and $|x|^3$ is differentiable at $x = 0$.

\therefore for $f(x)$ to be differentiable at $x = 0$, we must have $a = 0, b = 0$ and c is any real number.

57. (b) Given $x + |y| = 2y$

$$\Rightarrow x + y = 2y \text{ or } x - y = 2y$$

$$\Rightarrow x = y \text{ or } x = 3y$$

This represent a straight line which passes through origin.

Hence, $x + |y| = 2y$ is continuous at $x = 0$.

Now, we check differentiability at $x = 0$

$$x + |y| = 2y \Rightarrow x + y = 2y, y \geq 0$$

$$x - y = 2y, y < 0$$

$$\text{Thus, } f(x) = \begin{cases} x, & y < 0 \\ x/3, & y \geq 0 \end{cases}$$

$$\begin{aligned} \text{Now, L.H.D.} &= \lim_{h \rightarrow 0^-} \frac{f(x+h) - f(x)}{-h} \\ &= \lim_{h \rightarrow 0^-} \frac{x+h-x}{-h} = -1 \end{aligned}$$

$$\begin{aligned} \text{R.H.D.} &= \lim_{h \rightarrow 0^+} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0^+} \frac{\frac{x+h}{3} - \frac{x}{3}}{h} = \lim_{h \rightarrow 0^+} \frac{1}{3} = \frac{1}{3} \end{aligned}$$

Since, L.H.D. \neq R.H.D. at $x = 0$

\therefore given function is not differentiable at $x = 0$

$$58. \text{ (c) } \lim_{x \rightarrow a} \frac{x^2 f(a) - a^2 f(x)}{x - a}$$

Applying L-Hospital rule

$$= \lim_{x \rightarrow a} \frac{2xf(a) - a^2 f'(x)}{1} = 2af(a) - a^2 f'(a)$$

$$59. \text{ (c) } \text{ Given that, } f(x) = \begin{cases} (x-1)\sin\left(\frac{1}{x-1}\right), & \text{if } x \neq 1 \\ 0, & \text{if } x = 1 \end{cases}$$

At $x = 1$

$$\begin{aligned} \text{R.H.D.} &= \lim_{h \rightarrow 0} \frac{f(1+h) - f(1)}{h} \\ &= \lim_{h \rightarrow 0} \frac{h \sin \frac{1}{h} - 0}{h} = \lim_{h \rightarrow 0} \sin \frac{1}{h} = \text{a finite number} \end{aligned}$$

Let this finite number be l

$$\begin{aligned} \text{L.H.D.} &= \lim_{h \rightarrow 0} \frac{f(1-h) - f(1)}{-h} \\ &= \lim_{h \rightarrow 0} \frac{-h \sin\left(\frac{1}{-h}\right)}{-h} = \lim_{h \rightarrow 0} \sin\left(\frac{1}{-h}\right) \\ &= -\lim_{h \rightarrow 0} \sin\left(\frac{1}{h}\right) = -(\text{a finite number}) = -l \end{aligned}$$


Thus R.H.D. \neq L.H.D.

\therefore f is not differentiable at $x = 1$

$$\begin{aligned} \text{At } x=0 \quad f'(0) &= \sin \frac{1}{(x-1)} - \frac{x-1}{(x-1)^2} \cos\left(\frac{1}{x-1}\right) \Bigg|_{x=0} \\ &= -\sin 1 + \cos 1 \end{aligned}$$

$\therefore f$ is differentiable at $x = 0$

$$60. \text{ (a) } f(x) = \min \{x+1, |x|+1\} \\ \Rightarrow f(x) = x+1 \quad \forall x \in R$$



Since $f(x) = x+1$ is polynomial function

Hence, $f(x)$ is differentiable everywhere for all $x \in R$.

$$61. \text{ (c) } f(x) = \begin{cases} \frac{x}{1-x}, & x < 0 \\ \frac{x}{1+x}, & x \geq 0 \end{cases}$$

$f(x) = \frac{x}{1-x}$ is not define at $x \neq 1$ but here $x < 0$ and $f(x)$

$= \frac{x}{1+x}$ is not define at $x = -1$ but here $x > 0$. So, $f(x)$ is continuous for $x \in R$.

$$\text{and } f'(x) = \begin{cases} \frac{x}{(1-x)^2}, & x < 0 \\ \frac{x}{(1+x)^2}, & x \geq 0 \end{cases}$$

$\therefore f'(x)$ exist at everywhere.

$$62. \text{ (b) } \text{ Given that } |f(x) - f(y)| \leq (x-y)^2, x, y \in R \dots (i) \text{ and } f(0) = 0$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$|f'(x)| = \lim_{h \rightarrow 0} \left| \frac{f(x+h) - f(x)}{h} \right| \leq \lim_{h \rightarrow 0} \left| \frac{(h)^2}{h} \right|$$

$$\Rightarrow |f'(x)| \leq 0 \Rightarrow f'(x) = 0$$

$$\Rightarrow f(x) = \text{constant}$$

$$\text{As } f(0) = 0$$

$$\Rightarrow f(1) = 0.$$

$$63. \text{ (c) } f'(1) = \lim_{h \rightarrow 0} \frac{f(1+h) - f(1)}{h};$$

Given that function is differentiable so it is continuous also

and $\lim_{h \rightarrow 0} \frac{f(1+h)}{h} = 5$ and hence $f(1) = 0$

Hence, $f'(1) = \lim_{h \rightarrow 0} \frac{f(1+h)}{h} = 5$

64. (c) Given that $f(0) = 0$; $f(x) = xe^{-\left(\frac{1}{|x|} + \frac{1}{x}\right)}$

R.H.L. = $\lim_{h \rightarrow 0} (0+h)e^{-2/h} = \lim_{h \rightarrow 0} \frac{h}{e^{2/h}} = 0$

L.H.L. = $\lim_{h \rightarrow 0} (0-h)e^{-\left(\frac{1}{h} + \frac{1}{-h}\right)} = 0$

therefore, $f(x)$ is continuous at $x = 0$.

Now, R.H.D = $\lim_{h \rightarrow 0} \frac{(0+h)e^{-\left(\frac{1}{h} + \frac{1}{h}\right)} - 0}{h} = 0$

L.H.D. = $\lim_{h \rightarrow 0} \frac{(0-h)e^{-\left(\frac{1}{h} - \frac{1}{h}\right)} - 0}{-h} = 1$

therefore, L.H.D. \neq R.H.D.

$f(x)$ is not differentiable at $x = 0$.

65. (d) Let $u = \tan^{-1}\left(\frac{\sqrt{1+x^2}-1}{x}\right)$

Put $x = \tan \theta \Rightarrow \theta = \tan^{-1} x$

$\therefore u = \tan^{-1}\left(\frac{\sec \theta - 1}{\tan \theta}\right) = \tan^{-1}\left(\tan \frac{\theta}{2}\right)$

$= \frac{\theta}{2} = \frac{1}{2} \tan^{-1} x$

$\therefore \frac{du}{dx} = \frac{1}{2} \times \frac{1}{(1+x^2)}$

Let $v = \tan^{-1}\left(\frac{2x\sqrt{1-x^2}}{1-2x^2}\right)$

Put $x = \sin \phi \Rightarrow \phi = \sin^{-1} x$

$v = \tan^{-1}\left(\frac{2 \sin \phi \cos \phi}{\cos 2\phi}\right) = \tan^{-1}(\tan 2\phi)$

$= 2\phi = 2 \sin^{-1} x$

$\frac{dv}{dx} = 2 \frac{1}{\sqrt{1-x^2}}$

$\frac{du}{dv} = \frac{du/dx}{dv/dx} = \frac{\sqrt{1-x^2}}{4(1+x^2)}$

$\therefore \left(\frac{du}{dv}\right)_{\left(x=\frac{1}{2}\right)} = \frac{\sqrt{3}}{10}$

66. (c) $(a + \sqrt{2b} \cos x)(a - \sqrt{2b} \cos y) = a^2 - b^2$

Differentiating both sides,

$(-\sqrt{2b} \sin x)(a - \sqrt{2b} \cos y) + (a + \sqrt{2b} \cos x)$

$(\sqrt{2b} \sin y) \frac{dy}{dx} = 0$

$\Rightarrow \frac{dy}{dx} = \frac{(\sqrt{2b} \sin x)(a - \sqrt{2b} \cos y)}{(a + \sqrt{2b} \cos x)(\sqrt{2b} \sin y)}$

$\therefore \left[\frac{dy}{dx}\right]_{\left(\frac{\pi}{4}, \frac{\pi}{4}\right)} = \frac{a-b}{a+b} \Rightarrow \frac{dx}{dy} = \frac{a+b}{a-b}$

67. (91) $y = \sum_{k=1}^6 k \cos^{-1} \left\{ \frac{3}{5} \cos kx - \frac{4}{5} \sin kx \right\}$

Let $\cos a = \frac{3}{5}$ and $\sin a = \frac{4}{5}$

$\therefore y = \sum_{k=1}^6 k \cos^{-1} \{ \cos a \cos kx - \sin a \sin kx \}$

$= \sum_{k=1}^6 k \cos^{-1}(\cos(kx + a))$

$= \sum_{k=1}^6 k(kx + a) = \sum_{k=1}^6 (k^2 x + ak)$

$\therefore \frac{dy}{dx} = \sum_{k=1}^6 k^2 = \frac{6(7)(13)}{6} = 91.$

68. (Bonus) It is given that

$x = 2\sin\theta - \sin 2\theta$... (i)

$y = 2\cos\theta - \cos 2\theta$... (ii)

Differentiating (i) w.r.t. θ , we get

$\frac{dx}{d\theta} = 2\cos\theta - 2\cos 2\theta$

Differentiating (ii) w.r.t. θ ; we get

$\frac{dy}{d\theta} = -2\sin\theta + 2\sin 2\theta$

From (ii) \div (i), we get

$\therefore \frac{dy}{dx} = \frac{\sin 2\theta - \sin \theta}{\cos \theta - \cos 2\theta}$

$= \frac{2\sin \frac{\theta}{2} \cdot \cos \frac{3\theta}{2}}{2\sin \frac{\theta}{2} \cdot \sin \frac{3\theta}{2}} = \cot \frac{3\theta}{2}$

... (iii)

Again, differentiating eqn. (iii), we get

$\frac{d^2y}{dx^2} = \frac{-3}{2} \operatorname{cosec}^2 \frac{3\theta}{2} \cdot \frac{d\theta}{dx}$

$$\frac{d^2y}{dx^2} = \frac{-3 \operatorname{cosec}^2 3\theta}{2(\cos\theta - \cos 2\theta)}$$

$$\frac{d^2y}{dx^2}(\theta = \pi) = -\frac{3}{4(-1-1)} = \frac{3}{8}$$

$$69. \text{ (a) } y(\alpha) = \sqrt{\frac{2 \sin \alpha + \cos \alpha}{\sec^2 \alpha}} = \sqrt{\frac{2 \cos^2 \alpha}{\sin \alpha \cos \alpha} + \frac{1}{\sin^2 \alpha}}$$

$$= \sqrt{2 \cot \alpha + \operatorname{cosec}^2 \alpha} = \sqrt{2 \cot \alpha + 1 + \cot^2 \alpha}$$

$$= |1 + \cot \alpha| = -1 - \cot \alpha \quad \left[\because \alpha \in \left(\frac{3\pi}{4}, \pi \right) \right]$$

$$\frac{dy}{d\alpha} = \operatorname{cosec}^2 \alpha \Rightarrow \left(\frac{dy}{d\alpha} \right)_{\alpha = \frac{5\pi}{6}} = 4$$

$$70. \text{ (b) } \text{ Given, } x = \frac{1}{2}, y = \frac{-1}{4} \Rightarrow xy = \frac{-1}{8}$$

$$y \cdot \frac{1 \cdot (-2x)}{2\sqrt{1-x^2}} + y' \sqrt{1-x^2}$$

$$= - \left\{ 1 \cdot \sqrt{1-y^2} + \frac{x \cdot (-2y)}{2\sqrt{1-y^2}} y' \right\}$$

$$\Rightarrow -\frac{xy}{\sqrt{1-x^2}} + y' \sqrt{1-x^2} = -\sqrt{1-y^2} + \frac{xy \cdot y'}{\sqrt{1-y^2}}$$

$$\Rightarrow y' \left(\sqrt{1-x^2} - \frac{xy}{\sqrt{1-y^2}} \right) = \frac{xy}{\sqrt{1-x^2}} - \sqrt{1-y^2}$$

$$\Rightarrow y' \left(\frac{\sqrt{3}}{2} + \frac{1}{8 \cdot \frac{\sqrt{15}}{4}} \right) = \frac{-1}{8 \cdot \frac{\sqrt{3}}{2}} - \frac{\sqrt{15}}{4}$$

$$\Rightarrow y' \left(\frac{\sqrt{45}+1}{2\sqrt{15}} \right) = -\frac{(1+\sqrt{45})}{4\sqrt{3}}$$

$$\therefore y' = -\frac{\sqrt{5}}{2}$$

$$71. \text{ (b) } \text{ Given, } e^y + xy' = e \quad \dots(i)$$

Putting $x = 0$ in (i), $\Rightarrow e^y = e \Rightarrow y = 1$

On differentiating (i) w. r. to x

$$e^y \frac{dy}{dx} + x \frac{dy}{dx} + y = 0 \quad \dots(ii)$$

Putting $y = 1$ and $x = 0$ in (ii),

$$e \frac{dy}{dx} + 0 + 1 = 0 \Rightarrow \frac{dy}{dx} = -\frac{1}{e}$$

On differentiating (ii) w. r. to x ,

$$e^y \frac{d^2y}{dx^2} + \frac{dy}{dx} \cdot e^y \cdot \frac{dy}{dx} + x \frac{d^2y}{dx^2} + \frac{dy}{dx} + \frac{dy}{dx} = 0 \quad \dots(iii)$$

Putting $y = 1, x = 0$ and $\frac{dy}{dx} = -\frac{1}{e}$ in (iii),

$$e \frac{d^2y}{dx^2} + \frac{1}{e} - \frac{2}{e} = 0 \Rightarrow \frac{d^2y}{dx^2} = \frac{1}{e^2}$$

$$\text{Hence, } \left(\frac{dy}{dx}, \frac{d^2y}{dx^2} \right) = \left(-\frac{1}{e}, \frac{1}{e^2} \right)$$

$$72. \text{ (d) } f(x) = \tan^{-1} \left(\frac{\tan x - 1}{\tan x + 1} \right)$$

$$= -\tan^{-1} \left(\tan \left(\frac{\pi}{4} - x \right) \right) \quad \left[\because \frac{\pi}{4} - x \in \left(-\frac{\pi}{4}, \frac{\pi}{4} \right) \right]$$

$$\text{So, } f(x) = -\left(\frac{\pi}{4} - x \right) = x - \frac{\pi}{4}$$

$$\text{Let } y = \Rightarrow f(y) = 2y - \frac{\pi}{4}$$

Now, differentiate w.r.t. y , $\frac{df(y)}{dy} = 2$.

$$73. \text{ (none) } 2y = \left[\cot^{-1} \left(\frac{\frac{\sqrt{3}}{2} \cos x + \frac{1}{2} \sin x}{\frac{1}{2} \cos x - \frac{\sqrt{3}}{2} \sin x} \right) \right]^2$$


$$\Rightarrow 2y = \left[\cot^{-1} \left(\frac{\cos \left(\frac{\pi}{6} - x \right)}{\sin \left(\frac{\pi}{6} - x \right)} \right) \right]^2$$

$$\Rightarrow 2y = \left[\cot^{-1} \left(\cot \left(\frac{\pi}{6} - x \right) \right) \right]^2 \quad \because \frac{\pi}{6} - x \in \left(-\frac{\pi}{3}, \frac{\pi}{6} \right)$$

$$\Rightarrow 2y = \begin{cases} \left(\frac{7\pi}{6} - x \right)^2, & \text{if } \frac{\pi}{6} - x \in \left(-\frac{\pi}{3}, 0 \right) \\ \left(\frac{\pi}{6} - x \right)^2, & \text{if } \frac{\pi}{6} - x \in \left(0, \frac{\pi}{6} \right) \end{cases}$$

$$\Rightarrow \frac{dy}{dx} = \begin{cases} x - \frac{7\pi}{6} & \text{if } x \in \left(\frac{\pi}{6}, \frac{\pi}{2} \right) \\ x - \frac{\pi}{6} & \text{if } x \in \left(0, \frac{\pi}{6} \right) \end{cases}$$

74. (b) $f(x) = \min \{\sin x, \cos x\}$



$\therefore f(x)$ is not differentiable at $x = -\frac{3\pi}{4}, \frac{\pi}{4}$

$$\therefore S = \left\{ -\frac{3\pi}{4}, \frac{\pi}{4} \right\}$$

$$\Rightarrow S \subseteq \left\{ -\frac{3\pi}{4}, -\frac{\pi}{4}, \frac{3\pi}{4}, \frac{\pi}{4} \right\}$$

75. (a) Consider the equation,

$$(2x)^{2y} = 4e^{2x-2y}$$

Taking log on both sides

$$2y \ln(2x) = \ln 4 + (2x - 2y) \quad \dots(1)$$

Differentiating both sides w.r.t. x ,

$$2y \frac{1}{2x} + 2 \ln(2x) \frac{dy}{dx} = 0 + 2 - 2 \frac{dy}{dx}$$

$$2 \frac{dy}{dx} (1 + \ln(2x)) = 2 - \frac{2y}{x} = \frac{2x - 2y}{x} \quad \dots(2)$$

From (1) and (2),

$$\frac{dy}{dx} (1 + \ln 2x) = 1 - \frac{1}{x} \left(\frac{\ln 2 + x}{1 + \ln 2x} \right)$$

$$\begin{aligned} \Rightarrow (1 + \ln 2x)^2 \frac{dy}{dx} &= 1 + \ln(2x) - \left(\frac{x + \ln 2}{x} \right) \\ &= \frac{x \ln(2x) - \ln 2}{x} \end{aligned}$$

76. (c) Let $f(x) = x^3 + ax^2 + bx + c$

$$f'(x) = 3x^2 + 2ax + b \Rightarrow f'(1) = 3 + 2a + b$$

$$f''(x) = 6x + 2a \Rightarrow f''(2) = 12 + 2a$$

$$f'''(x) = 6 \Rightarrow f'''(3) = 6$$

$$\therefore f(x) = x^3 + f'(1)x^2 + f''(2)x + f'''(3)$$

$$\therefore f'(1) = a \Rightarrow 3 + 2a + b = a \Rightarrow a + b = -3 \quad \dots(1)$$

$$\text{also } f''(2) = b \Rightarrow 12 + 2a = b \Rightarrow 2a - b = -12 \quad \dots(2)$$

$$\text{and } f'''(3) = c \Rightarrow c = 6$$

Add (1) and (2)

$$3a = -15 \Rightarrow a = -5 \Rightarrow b = 2$$

$$\Rightarrow f(x) = x^3 - 5x^2 + 2x + 6$$

$$\Rightarrow f(2) = 8 - 20 + 4 + 6 = -2$$

77. (b) $\therefore x = 3 \tan t \Rightarrow \frac{dx}{dt} = 3 \sec^2 t$

$$\text{and } y = 3 \sec t \Rightarrow \frac{dy}{dt} = 3 \sec t \cdot \tan t$$

$$\therefore \frac{dy}{dx} = \frac{dy/dt}{dx/dt} \therefore \frac{dy}{dx} = \frac{\tan t}{\sec t} = \sin t$$

$$\begin{aligned} \therefore \frac{d^2 y}{dx^2} &= \frac{d}{dt} (\sin t) \cdot \frac{dt}{dx} \\ &= \cos t \cdot \frac{1}{3 \sec^2 t} \end{aligned}$$

$$\begin{aligned} \therefore \frac{d^2 y}{dx^2} \left(\text{at } t = \frac{\pi}{4} \right) &= \frac{1}{3} \cdot \left(\frac{1}{\sqrt{2}} \right)^3 \\ &= \frac{1}{6\sqrt{2}} \end{aligned}$$

78. (b) Here, $\frac{dx}{dt} = \frac{1}{2\sqrt{2^{\csc^{-1}t}}} 2^{\csc^{-1}t} \log 2 \cdot \frac{-1}{x\sqrt{x^2-1}}$

$$\frac{dy}{dt} = \frac{1}{2\sqrt{2^{\sec^{-1}t}}} 2^{\sec^{-1}t} \log 2 \cdot \frac{1}{x\sqrt{x^2-1}}$$

$$\therefore \frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{-\sqrt{2^{\csc^{-1}t}}}{\sqrt{2^{\sec^{-1}t}}} \cdot \frac{2^{\sec^{-1}t}}{2^{\csc^{-1}t}}$$

$$\frac{dy}{dx} = \frac{dy}{dt} = -\sqrt{\frac{2^{\sec^{-1}t}}{2^{\csc^{-1}t}}} = \frac{-y}{x}$$

$$\begin{aligned}
 79. \quad (a) \quad f(x) &= \begin{vmatrix} \cos x & x & 1 \\ 2 \sin x & x^2 & 2x \\ \tan x & x & 1 \end{vmatrix} \\
 &= \cos x (x^2 - 2x^2) - x (2 \sin x - 2x \tan x) \\
 &\quad + 1(2x \sin x - x^2 \tan x) \\
 &= -x^2 \cos x - 2x \sin x + 2x^2 \tan x + 2x \sin x - x^2 \tan x \\
 &= x^2 \tan x - x^2 \cos x = x^2 (\tan x - \cos x) \\
 &\Rightarrow f'(x) = 2x (\tan x - \cos x) + x^2 (\sec^2 x + \sin x) \\
 \therefore \lim_{x \rightarrow 0} \frac{f'(x)}{x} &= \\
 \lim_{x \rightarrow 0} \frac{2x (\tan x - \cos x) + x^2 (\sec^2 x + \sin x)}{x} &= \\
 = \lim_{x \rightarrow 0} (\tan x - \cos x) + x (\sec^2 x + \sin x) &= \\
 = 2(0 - 1) + 0 = -2 &= \\
 \text{So, } \lim_{x \rightarrow 0} \frac{f'(x)}{x} &= -2
 \end{aligned}$$

$$\begin{aligned}
 80. \quad (a) \quad \text{Since } f(x) &= \sin^{-1} \left(\frac{2 \times 3^x}{1 + 9^x} \right) \\
 \text{Suppose } 3^x &= \tan t \\
 \Rightarrow f(x) &= \sin^{-1} \left(\frac{2 \tan t}{1 + \tan^2 t} \right) = \sin^{-1} (\sin 2t) = 2t \\
 &= 2 \tan^{-1} (3x) \\
 \text{So, } f'(x) &= \frac{2}{1 + (3^x)^2} \times 3^x \cdot \log_e 3 \\
 \therefore f' \left(-\frac{1}{2} \right) &= \frac{2}{1 + \left(3^{-\frac{1}{2}} \right)^2} \times 3^{-\frac{1}{2}} \cdot \log_e 3 \\
 &= \frac{1}{2} \times \sqrt{3} \times \log_e 3 = \sqrt{3} \times \log_e \sqrt{3}
 \end{aligned}$$

$$\begin{aligned}
 81. \quad (a) \quad \text{Given, } x^2 + y^2 + \sin y &= 4 \\
 \text{After differentiating the above equation w. r. t. } x &\text{ we get} \\
 2x + 2y \frac{dy}{dx} + \cos y \frac{dy}{dx} &= 0 \quad \dots (1) \\
 \Rightarrow 2x + (2y + \cos y) \frac{dy}{dx} &= 0 \\
 \Rightarrow \frac{dy}{dx} &= \frac{-2x}{2y + \cos y} \\
 \text{At } (-2, 0), \left(\frac{dy}{dx} \right)_{(-2,0)} &= \frac{-2 \times -2}{2 \times 0 + \cos 0} \\
 \Rightarrow \left(\frac{dy}{dx} \right)_{(-2,0)} &= \frac{4}{0+1}
 \end{aligned}$$

$$\Rightarrow \left(\frac{dy}{dx} \right)_{(-2,0)} = 4 \quad \dots (2)$$

Again differentiating equation (1) w. r. t. x , we get

$$2 + 2 \left(\frac{dy}{dx} \right)^2 + 2y \frac{d^2y}{dx^2} - \sin y \left(\frac{dy}{dx} \right)^2 + \cos y \frac{d^2y}{dx^2} = 0$$

$$\Rightarrow 2 + (2 - \sin y) \left(\frac{dy}{dx} \right)^2 + (2y + \cos y) \frac{d^2y}{dx^2} = 0$$

$$\Rightarrow (2y + \cos y) \frac{d^2y}{dx^2} = -2 - (2 - \sin y) \left(\frac{dy}{dx} \right)^2$$

$$\Rightarrow \frac{d^2y}{dx^2} = \frac{-2 - (2 - \sin y) \left(\frac{dy}{dx} \right)^2}{2y + \cos y}$$

So, at $(-2, 0)$,

$$\frac{d^2y}{dx^2} = \frac{-2 - (2 - 0) \times 4^2}{2 \times 0 + 1}$$

$$\Rightarrow \frac{d^2y}{dx^2} = \frac{-2 - 2 \times 16}{1}$$

$$\Rightarrow \frac{d^2y}{dx^2} = -34$$

$$82. \quad (b) \quad \text{Let } F(x) = \tan^{-1} \left(\frac{6x\sqrt{x}}{1 - 9x^3} \right) \text{ where } x \in \left(0, \frac{1}{4} \right).$$

$$= \tan^{-1} \left(\frac{2 \cdot (3x^{3/2})}{1 - (3x^{3/2})^2} \right) = 2 \tan^{-1} (3x^{3/2})$$

$$\text{As } 3x^{3/2} \in \left(0, \frac{3}{8} \right)$$

$$\left[\because 0 < x < \frac{1}{4} \Rightarrow 0 < x^{3/2} < \frac{1}{8} \Rightarrow 0 < 3x^{3/2} < \frac{3}{8} \right]$$

$$\text{So } \frac{dF(x)}{dx} = 2 \times \frac{1}{1 + 9x^3} \times 3 \times \frac{3}{2} \times x^{1/2} = \frac{9}{1 + 9x^3} \sqrt{x}$$

On comparing

$$\therefore g(x) = \frac{9}{1 + 9x^3}$$

$$83. \quad (d) \quad g(x) = f(f(x))$$

In the neighbourhood of $x = 0$,

$$f(x) = |\log 2 - \sin x| = (\log 2 - \sin x)$$

$$\therefore g(x) = |\log 2 - \sin| \log 2 - \sin x || \\ = (\log 2 - \sin(\log 2 - \sin x))$$

$\therefore g(x)$ is differentiable

$$\text{and } g'(x) = -\cos(\log 2 - \sin x) (-\cos x)$$

$$\Rightarrow g'(0) = \cos(\log 2)$$

84. (c) $f(x) = y = x^2 - x + 5$

$$x^2 - x + \frac{1}{4} - \frac{1}{4} + 5 = y$$

$$\left(x - \frac{1}{2}\right)^2 + \frac{19}{4} = y$$

$$\left(x - \frac{1}{2}\right)^2 = y - \frac{19}{4}$$

$$x - \frac{1}{2} = \pm \sqrt{y - \frac{19}{4}}$$

$$x = \frac{1}{2} \pm \sqrt{y - \frac{19}{4}}$$

As $x > \frac{1}{2}$

$$x = \frac{1}{2} + \sqrt{y - \frac{19}{4}}$$

$$g(x) = \frac{1}{2} + \sqrt{x - \frac{19}{4}}$$

$$g'(x) = \frac{1}{2\sqrt{x - \frac{19}{4}}}$$

$$g'(7) = \frac{1}{2\sqrt{7 - \frac{19}{4}}} = \frac{1}{2\frac{\sqrt{28-19}}{2}} = \frac{1}{3}$$

85. (a) Let $y = \sec(\tan^{-1} x) = \sec(\sec^{-1} \sqrt{1+x^2})$

$$\Rightarrow y = \sqrt{1+x^2}$$

$$\Rightarrow \frac{dy}{dx} = \frac{1}{2\sqrt{1+x^2}} \cdot 2x$$

At $x=1$,

$$\frac{dy}{dx} = \frac{1}{\sqrt{2}}$$

86. (b) $\frac{x^2}{\alpha} + \frac{y^2}{4} = 1 \Rightarrow \frac{2x}{\alpha} + \frac{2y}{4} \cdot \frac{dy}{dx} = 0$

$$\Rightarrow \frac{dy}{dx} = \frac{-4x}{\alpha y} \quad \dots(i)$$

$$y^3 = 16x \Rightarrow 3y^2 \cdot \frac{dy}{dx} = 16 \Rightarrow \frac{dy}{dx} = \frac{16}{3y^2} \quad \dots(ii)$$

Since curves intersect at right angles

$$\therefore \frac{-4x}{\alpha y} \times \frac{16}{3y^2} = -1 \Rightarrow 3\alpha y^3 = 64x$$

$$\Rightarrow \alpha = \frac{64x}{3 \times 16x} = \frac{4}{3}$$

87. (d) Let $x = \sqrt{a^{\sin^{-1} t}}$

$$\Rightarrow x^2 = a^{\sin^{-1} t}$$

$$\Rightarrow 2 \log x = \sin^{-1} t \cdot \log a$$

$$\Rightarrow \frac{2}{x} = \frac{\log a}{\sqrt{1-t^2}} \cdot \frac{dt}{dx}$$

$$\Rightarrow \frac{2\sqrt{1-t^2}}{x \log a} = \frac{dt}{dx} \quad \dots(1)$$

Now, let $y = \sqrt{a^{\cos^{-1} t}}$

$$\Rightarrow 2 \log y = \cos^{-1} t \cdot \log a$$

$$\Rightarrow \frac{2}{y} \cdot \frac{dy}{dx} = \frac{-\log a}{\sqrt{1-t^2}} \cdot \frac{dt}{dx}$$

$$\Rightarrow \frac{2}{y} \cdot \frac{dy}{dx} = \frac{-\log a}{\sqrt{1-t^2}} \times \frac{2\sqrt{1-t^2}}{x \log a} \quad \text{(from (1))}$$

$$\Rightarrow \frac{dy}{dx} = -\frac{y}{x}$$

Hence, $1 + \left(\frac{dy}{dx}\right)^2 = 1 + \left(\frac{-y}{x}\right)^2 = \frac{x^2 + y^2}{x^2}$

88. (b) Let $y = \frac{x^2 - x}{x^2 + 2x}$

$$\Rightarrow (x^2 + 2x)y = x^2 - x$$

$$\Rightarrow x(x+2)y = x(x-1)$$

$$\Rightarrow x[(x+2)y - (x-1)] = 0$$

$$\because x \neq 0, \therefore (x+2)y - (x-1) = 0$$

$$\Rightarrow xy + 2y - x + 1 = 0$$

$$\Rightarrow x(y-1) = -(2y+1)$$

$$\therefore x = \frac{2y+1}{1-y} \Rightarrow f^{-1}(x) = \frac{2x+1}{1-x}$$

$$\frac{d}{dx}(f^{-1}(x)) = \frac{2(1-x) - (2x+1)(-1)}{(1-x)^2}$$

$$= \frac{2 - 2x + 2x + 1}{(1-x)^2} = \frac{3}{(1-x)^2}$$

89. (c) Let $f'(x) = \sin[\log x]$ and $y = f\left(\frac{2x+3}{3-2x}\right)$

Now, $\frac{dy}{dx} = f'\left(\frac{2x+3}{3-2x}\right) \cdot \frac{d}{dx}\left(\frac{2x+3}{3-2x}\right)$

$$= \sin\left[\log\left(\frac{2x+3}{3-2x}\right)\right] \cdot \frac{[(6-4x) - (-4x-6)]}{(3-2x)^2}$$

$$= \frac{12}{(3-2x)^2} \cdot \sin\left[\log\left(\frac{2x+3}{3-2x}\right)\right]$$

90. (a) Given that $g(x) = [f(2f(x)) + 2]^2$

$$\therefore g'(x) = 2(f(2f(x)) + 2) \left(\frac{d}{dx}(f(2f(x)) + 2) \right)$$

$$= 2f(2f(x)) + 2 \cdot f'(2f(x)) \cdot (2f'(x))$$

$$\Rightarrow g'(0) = 2f(2f(0)) + 2 \cdot f'(2f(0)) \cdot 2$$

$$= 2f(0) + 4f'(0) = 4(-1) + 4(1) = 0$$

91. (d) $x^{2x} - 2x^x \cot y - 1 = 0$
 $\Rightarrow 2 \cot y = x^x - x^{-x}$
 Let $u = x^x$

$$\Rightarrow 2 \cot y = u - \frac{1}{u}$$

Differentiating both sides with respect to x , we get

$$-2 \operatorname{cosec}^2 y \frac{dy}{dx} = \left(1 + \frac{1}{u^2} \right) \frac{du}{dx}$$

Now $u = x^x$ Taking log both sides

$$\Rightarrow \log u = x \log x$$

$$\Rightarrow \frac{1}{u} \frac{du}{dx} = 1 + \log x$$

$$\Rightarrow \frac{du}{dx} = x^x (1 + \log x)$$

\therefore We get

$$-2 \operatorname{cosec}^2 y \frac{dy}{dx} = (1 + x^{-2x}) \cdot x^x (1 + \log x)$$

$$\Rightarrow \frac{dy}{dx} = \frac{(x^x + x^{-x})(1 + \log x)}{-2(1 + \cot^2 y)} \quad \dots(i)$$

Put $n = 1$ in eqn. $x^{2x} - 2x^x \cot y - 1 = 0$, gives

$$1 - 2 \cot y - 1 = 0$$

$$\Rightarrow \cot y = 0$$

\therefore Putting $x = 1$ and $\cot y = 0$ in eqn. (i), we get

$$y'(1) = \frac{(1+1)(1+0)}{-2(1+0)} = -1$$

92. (a) $x^m \cdot y^n = (x+y)^{m+n}$
 taking log both sides
 $\Rightarrow m \ln x + n \ln y = (m+n) \ln(x+y)$
 Differentiating both sides, we get

$$\therefore \frac{m}{x} + \frac{n}{y} \frac{dy}{dx} = \frac{m+n}{x+y} \left(1 + \frac{dy}{dx} \right)$$

$$\Rightarrow \left(\frac{m}{x} - \frac{m+n}{x+y} \right) = \left(\frac{m+n}{x+y} - \frac{n}{y} \right) \frac{dy}{dx}$$

$$\Rightarrow \frac{my - nx}{x(x+y)} = \left(\frac{my - nx}{y(x+y)} \right) \frac{dy}{dx}$$

$$\Rightarrow \frac{dy}{dx} = \frac{y}{x}$$

93. (c) Given that $x = e^{y+e^{y+\dots\infty}} \Rightarrow x = e^{y+x}$.

Taking log both sides.

$$\log x = y + x \text{ differentiating both side } \Rightarrow \frac{1}{x} = \frac{dy}{dx} + 1$$

$$\therefore \frac{dy}{dx} = \frac{1}{x} - 1 = \frac{1-x}{x}$$

94. (b) $f(x) = ax^2 + bx + c$

$$f(1) = f(-1)$$

$$\Rightarrow a + b + c = a - b + c \text{ or } b = 0$$

$$\therefore f(x) = ax^2 + c \text{ or } f'(x) = 2ax$$

Now $f'(a); f'(b)$ and $f'(c)$

$$\text{are } 2a(a); 2a(b); 2a(c)$$

i.e. $2a^2, 2ab, 2ac$.

\Rightarrow If a, b, c are in A.P. then $f'(a); f'(b)$ and $f'(c)$ are also in A.P.

95. (c) Given that $f(x+y) = f(x) \times f(y)$

Differentiate with respect to x , treating y as constant

$$f'(x+y) = f'(x)f(y)$$

Putting $x = 0$ and $y = x$, we get $f'(x) = f'(0)f(x)$;

$$\Rightarrow f'(5) = 3f(5) = 3 \times 2 = 6.$$

96. (b) Let $f: \mathbb{R} \rightarrow \mathbb{R}$, with $f(0) = f(1) = 0$ and $f'(0) = 0$

$\therefore f(x)$ is differentiable and continuous and

$$f(0) = f(1) = 0.$$

Then by Rolle's theorem, $f'(c) = 0$, $c \in (0, 1)$

Now again

$$\therefore f'(c) = 0, f'(0) = 0$$

Then, again by Rolle's theorem,

$$f''(x) = 0 \text{ for some } x \in (0, 1)$$

97. (c) $y^2 + 2 \log_e(\cos x) = y$...(i)

$$\Rightarrow 2yy' - 2 \tan x = y' \quad \dots(ii)$$

From (i), $y(0) = 0$ or 1

$$\therefore y'(0) = 0$$

Again differentiating (ii) we get,

$$2(y')^2 + 2yy'' - 2 \sec^2 x = y''$$

Put $x = 0$, $y(0) = 0, 1$ and $y'(0) = 0$,

we get, $|y''(0)| = 2$.

98. (b) Since, Rolle's theorem is applicable

$$\therefore f(a) = f(b)$$

$$f(3) = f(4) \Rightarrow \alpha = 12$$

$$f'(x) = \frac{x^2 - 12}{x(x^2 + 12)}$$

As $f'(c) = 0$ (by Rolle's theorem)

$$x = \pm\sqrt{12}, \therefore c = \sqrt{12}, \therefore f''(c) = \frac{1}{12}$$

99. (c) $k \cdot x^{k-1} + k \cdot y^{k-1} \frac{dy}{dx} = 0$

$$\Rightarrow \frac{dy}{dx} = -\left(\frac{x}{y}\right)^{k-1}$$

$$\Rightarrow \frac{dy}{dx} + \left(\frac{x}{y}\right)^{k-1} = 0$$

$$\Rightarrow k-1 = -\frac{1}{3}$$

$$\Rightarrow k = 1 - \frac{1}{3} = \frac{2}{3}$$

100. (b) Since, $f(x)$ is a polynomial function.
 \therefore It is continuous and differentiable in $[0, 1]$
 Here, $f(0) = 11, f(1) = 1 - 4 + 8 + 11 = 16$
 $f'(x) = 3x^2 - 8x + 8$

$$\therefore f'(c) = \frac{f(1) - f(0)}{1 - 0} = \frac{16 - 11}{1}$$

$$= 3c^2 - 8c + 8$$

$$\Rightarrow 3c^2 - 8c + 3 = 0$$

$$\Rightarrow c = \frac{8 \pm 2\sqrt{7}}{6} = \frac{4 \pm \sqrt{7}}{3}$$

$$\therefore c = \frac{4 - \sqrt{7}}{3} \in (0, 1)$$

101. (b) $y^{1/5} + y^{-1/5} = 2x$

$$\Rightarrow \left(\frac{1}{5}y^{-4/5} - \frac{1}{5}y^{-6/5}\right) \frac{dy}{dx} = 2$$

$$\Rightarrow y'(y^{1/5} - y^{-1/5}) = 10y$$

$$\Rightarrow y^{1/5} + y^{-1/5} = 2x$$

$$\Rightarrow y^{1/5} - y^{-1/5} = \sqrt{4x^2 - 4}$$

$$\Rightarrow y'(2\sqrt{x^2 - 1}) = 10y$$

$$\Rightarrow y''(2\sqrt{x^2 - 1}) + y' \cdot \frac{2x}{2\sqrt{x^2 - 1}} = 10y'$$

$$\Rightarrow y''(x^2 - 1) + xy' = 5\sqrt{x^2 - 1}(y')$$

$$\Rightarrow \boxed{y''(x^2 - 1) + xy' - 25y = 0}$$

$$\lambda = 1, k = -25$$

102. (b) Let $f(x) = ax^3 + bx^2 + cx + d$

$$\Rightarrow f(3x) = 27ax^3 + 9bx^2 + 3cx + d$$

$$\Rightarrow f'(x) = 3ax^2 + 2bx + c$$

$$\Rightarrow f''(x) = 6ax + 2b$$

$$\Rightarrow f(3x) = f'(x)f''(x)$$

$$\Rightarrow 27a = 18a^2$$

$$\Rightarrow a = \frac{3}{2}, b = 0, c = 0, d = 0$$

$$\Rightarrow f(x) = \frac{3}{2}x^3,$$

$$f'(x) = \frac{9}{2}x^2, f''(x) = 9x$$

$$\Rightarrow f'(2) = 18$$

$$\text{and } f''(2) = 18$$

$$\Rightarrow f''(b) - f'(b) = 0$$

103. (d) $y = \left\{x + \sqrt{x^2 - 1}\right\}^{15} + \left\{x - \sqrt{x^2 - 1}\right\}^{15}$

Differentiate w.r.t. 'x'

$$\frac{dy}{dx} = 15\left(x + \sqrt{x^2 - 1}\right)^{14} \left[1 + \frac{x}{\sqrt{x^2 - 1}}\right]$$

$$+ 15\left(x - \sqrt{x^2 - 1}\right)^{14} \left(1 - \frac{x}{\sqrt{x^2 - 1}}\right)$$

$$\Rightarrow \frac{dy}{dx} = \frac{15}{\sqrt{x^2 - 1}} \cdot y \quad \dots(i)$$

$$\Rightarrow \sqrt{x^2 - 1} \cdot \frac{dy}{dx} = 15y$$

Again differentiating both sides w.r.t. x

$$\frac{x}{\sqrt{x^2 - 1}} \cdot \frac{dy}{dx} + \sqrt{x^2 - 1} \cdot \frac{d^2y}{dx^2} = 15 \frac{dy}{dx}$$

$$\Rightarrow x \frac{dy}{dx} + (x^2 - 1) \frac{d^2y}{dx^2}$$

$$= 15\sqrt{x^2 - 1} \cdot \frac{15}{\sqrt{x^2 - 1}} \cdot y = 225y$$

104. (b) Conduction for Rolle's theorem

$$f(1) = f(-1)$$

$$\text{and } f'\left(\frac{1}{2}\right) = 0$$

$$c = -2 \text{ and } b = \frac{1}{2}$$

$$2b + c = -1$$

105. (b) Since, f and g both are continuous function on $[0, 1]$ and differentiable on $(0, 1)$ then $\exists c \in (0, 1)$ such that

$$f'(c) = \frac{f(1) - f(0)}{1 - 0} = \frac{6 - 2}{1} = 4$$

$$\text{and } g'(c) = \frac{g(1) - g(0)}{1 - 1} = \frac{2 - 0}{1} = 2$$

Thus, we get $f'(c) = 2g'(c)$

- 106. (c)** Let $f(x) = x|x| = x|x|$, $g(x) = \sin x$
and $h(x) = g \circ f(x) = g[f(x)]$

$$\therefore h(x) = \begin{cases} \sin x^2, & x \geq 0 \\ -\sin x^2, & x < 0 \end{cases}$$

$$\text{Now, } h'(x) = \begin{cases} 2x \cos x^2, & x \geq 0 \\ -2x \cos x^2, & x < 0 \end{cases}$$

Since, L.H.L and R.H.L at $x = 0$ of $h'(x)$ is equal to 0
therefore $h'(x)$ is continuous at $x = 0$

Now, suppose $h'(x)$ is differentiable

$$\therefore h''(x) = \begin{cases} 2(\cos x^2 + 2x^2(-\sin x^2)), & x \geq 0 \\ 2(-\cos x^2 + 2x^2 \sin x^2), & x < 0 \end{cases}$$

Since, L.H.L and R.H.L at $x = 0$ of $h''(x)$ are different
therefore $h''(x)$ is not continuous.

$\Rightarrow h''(x)$ is not differentiable

\Rightarrow our assumption is wrong

Hence $h'(x)$ is not differentiable at $x = 0$.

- 107. (a)** Let $p_1(x) = a_1x^2 + b_1x + c_1$
 $p_2(x) = a_2x^2 + b_2x + c_2$
and $p_3(x) = a_3x^2 + b_3x + c_3$
where $a_1, a_2, a_3, b_1, b_2, b_3, c_1, c_2, c_3$ are real numbers.

$$\therefore A(x) = \begin{bmatrix} a_1x^2 + b_1x + c_1 & 2a_1x + b_1 & 2a_1 \\ a_2x^2 + b_2x + c_2 & 2a_2x + b_2 & 2a_2 \\ a_3x^2 + b_3x + c_3 & 2a_3x + b_3 & 2a_3 \end{bmatrix}$$

$$B(x) = \begin{bmatrix} a_1x^2 + b_1x + c_1 & 2a_1x + b_1 & 2a_1 \\ 2a_1x + b_1 & 2a_2x + b_2 & 2a_3x + b_2 \\ 2a_1 & 2a_2 & 2a_3 \end{bmatrix}$$

$$\times \begin{bmatrix} a_1x^2 + b_1x + c_1 & 2a_1x + b_1 & 2a_1 \\ a_2x^2 + b_2x + c_2 & 2a_2x + b_2 & 2a_2 \\ a_3x^2 + b_3x + c_3 & 2a_3x + b_3 & 2a_3 \end{bmatrix}$$

It is clear from the above multiplication, the degree of determinant of $B(x)$ can not be less than 4.

- 108. (b)** $f(x) = 2x^3 + ax^2 + bx$

let, $a = -1$, $b = 1$

Given that $f(x)$ satisfy Rolle's theorem in interval $[-1, 1]$
 $f(x)$ must satisfy two conditions.

(1) $f(a) = f(b)$

(2) $f'(c) = 0$ (c should be between a and b)

$$f(a) = f(1) = 2(1)^3 + a(1)^2 + b(1) = 2 + a + b$$

$$f(b) = f(-1) = 2(-1)^3 + a(-1)^2 + b(-1) \\ = -2 + a - b$$

$$f(a) = f(b)$$

$$2 + a + b = -2 + a - b$$

$$2b = -4$$

$$b = -2$$

$$\left(\text{given that } c = \frac{1}{2}\right)$$

$$f'(x) = 6x^2 + 2ax + b$$

$$\text{at } x = \frac{1}{2}, f'(x) = 0$$

$$0 = 6\left(\frac{1}{2}\right)^2 + 2a\left(\frac{1}{2}\right) + b$$

$$\frac{3}{2} + a + b = 0$$

$$\frac{3}{2} + a - 2 = 0$$

$$a = 2 - \frac{3}{2} = \frac{1}{2}$$

$$2a + b = 2 \times \frac{1}{2} - 2 = 1 - 2 = -1$$

- 109. (d)** $f(x) = \sin(\sin x)$

$$\Rightarrow f'(x) = \cos(\sin x) \cdot \cos x$$

$$\Rightarrow f''(x) = -\sin(\sin x) \cdot \cos^2 x + \cos(\sin x) \cdot (-\sin x) \\ = -\cos^2 x \cdot \sin(\sin x) - \sin x \cdot \cos(\sin x)$$

$$\text{Now } f'''(x) + \tan x \cdot f'(x) + g(x) = 0$$

$$\Rightarrow g(x) = \cos^2 x \cdot \sin(\sin x) + \sin x \cdot \cos(\sin x)$$

$$- \tan x \cdot \cos x \cdot \cos(\sin x)$$

$$\Rightarrow g(x) = \cos^2 x \cdot \sin(\sin x).$$

- 110. (d)** Let $g(x) = \frac{ax^3}{3} + b \cdot \frac{x^2}{2} + cx$

$$g'(x) = ax^2 + bx + c$$

$$\text{Given: } ax^2 + bx + c = 0 \text{ and } 2a + 3b + 6c = 0$$

Statement-2:

$$(i) \quad g(0) = 0 \text{ and } g(1) = \frac{a}{3} + \frac{b}{2} + c = \frac{2a + 3b + 6c}{6}$$

$$= \frac{0}{6} = 0$$

$$\Rightarrow g(0) = g(1)$$

(ii) g is continuous on $[0, 1]$ and differentiable on $(0, 1)$

\therefore By Rolle's theorem $\exists k \in (0, 1)$ such that $g'(k) = 0$

This holds the statement 2. Also, from statement-2, we can say $ax^2 + bx + c = 0$ has at least one root in $(0, 1)$.

Thus statement-1 and 2 both are true and statement-2 is a correct explanation for statement-1.

$$\begin{aligned} 111. (c) \quad \frac{d^2x}{dy^2} &= \frac{d}{dy} \left(\frac{dx}{dy} \right) = \frac{d}{dx} \left(\frac{dx}{dy} \right) \frac{dx}{dy} = \frac{d}{dx} \left(\frac{1}{dy/dx} \right) \frac{dx}{dy} \\ &= -\frac{1}{\left(\frac{dy}{dx}\right)^2} \cdot \frac{d^2y}{dx^2} \cdot \frac{1}{\frac{dy}{dx}} \left[\because \frac{d}{dx} \left(\frac{1}{x} \right) = -\frac{1}{x^2} \right] \\ &= -\frac{1}{\left(\frac{dy}{dx}\right)^3} \frac{d^2y}{dx^2} \end{aligned}$$

112. (b) Given that $f(x) = x|x|$ and $g(x) = \sin x$
 So that
 $g \circ f(x) = g(f(x)) = g(x|x|) = \sin x|x|$

$$= \begin{cases} \sin(-x^2), & \text{if } x < 0 \\ \sin(x^2), & \text{if } x \geq 0 \end{cases} = \begin{cases} -\sin x^2, & \text{if } x < 0 \\ \sin x^2, & \text{if } x \geq 0 \end{cases}$$

$$\therefore (g \circ f)'(x) = \begin{cases} -2x \cos x^2, & \text{if } x < 0 \\ 2x \cos x^2, & \text{if } x \geq 0 \end{cases}$$

Here we observe

$$L(g \circ f)'(0) = 0 = R(g \circ f)'(0)$$

$\Rightarrow g \circ f$ is differentiable at $x = 0$
 and $(g \circ f)'$ is continuous at $x = 0$

$$\text{Now } (g \circ f)''(x) = \begin{cases} -2 \cos x^2 + 4x^2 \sin x^2, & x < 0 \\ 2 \cos x^2 - 4x^2 \sin x^2, & x \geq 0 \end{cases}$$

Here

$$L(g \circ f)''(0) = -2 \text{ and } R(g \circ f)''(0) = 2$$

$$\therefore L(g \circ f)''(0) \neq R(g \circ f)''(0)$$

$\Rightarrow g \circ f(x)$ is not twice differentiable at $x = 0$.

\therefore Statement - 1 is true but statement -2 is false.

113. (c) Using Lagrange's Mean Value Theorem
 Let $f(x)$ be a function defined on $[a, b]$

$$\text{then, } f'(c) = \frac{f(b) - f(a)}{b - a} \quad \dots(i)$$

$$c \in [a, b]$$

$$\therefore \text{ Given } f(x) = \log_e x \quad \therefore f'(x) = \frac{1}{x}$$

\therefore equation (i) become

$$\frac{1}{c} = \frac{f(3) - f(1)}{3 - 1}$$

$$\Rightarrow \frac{1}{c} = \frac{\log_e 3 - \log_e 1}{2} = \frac{\log_e e^3}{2}$$

$$\Rightarrow c = \frac{2}{\log_e 3} \Rightarrow c = 2 \log_3 e$$

114. (a) As $f(1) = -2$ & $f'(x) \geq 2 \forall x \in [1, 6]$
 Applying Lagrange's mean value theorem

$$\frac{f(6) - f(1)}{5} = f'(c) \geq 2$$

$$\Rightarrow f(6) \geq 10 + f(1)$$

$$\Rightarrow f(6) \geq 10 - 2 \Rightarrow f(6) \geq 8.$$

115. (b) Let $f(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + 0$

The other given equation,

$$n a_n x^{n-1} + (n-1) a_{n-1} x^{n-2} + \dots + a_1 = 0 = f'(x)$$

$$\text{Given } a_1 \neq 0 \Rightarrow f(0) = 0$$

Again $f(x)$ has root α , $\Rightarrow f(\alpha) = 0$

$$\therefore f(0) = f(\alpha)$$

\therefore By Rolle's theorem $f'(x) = 0$ has root between $(0, \alpha)$

Hence $f'(x)$ has a positive root smaller than α .

116. (d) Let us define a function

$$f(x) = \frac{ax^3}{3} + \frac{bx^2}{2} + cx$$

Being polynomial, it is continuous and differentiable, also,

$$f(0) = 0 \text{ and } f(1) = \frac{a}{3} + \frac{b}{2} + c$$

$$\Rightarrow f(1) = \frac{2a + 3b + 6c}{6} = 0 \text{ (given)}$$

$$\therefore f(0) = f(1)$$

$\therefore f(x)$ satisfies all conditions of Rolle's theorem therefore $f'(x) = 0$ has a root in $(0, 1)$

i.e. $ax^2 + bx + c = 0$ has at least one root in $(0, 1)$

117. (d) Given that $f(x) = x^n \Rightarrow f(1) = 1$

$$f'(x) = nx^{n-1} \Rightarrow f'(1) = n$$

$$f''(x) = n(n-1)x^{n-2} \Rightarrow f''(1) = n(n-1)$$

$$f^n(x) = n! \Rightarrow f^n(1) = n!$$

$$f(1) - \frac{f'(1)}{1!} + \frac{f''(1)}{2!} - \frac{f'''(1)}{3!} + \dots + \frac{(-1)^n f^n(1)}{n!}$$

$$= 1 - \frac{n}{1!} + \frac{n(n-1)}{2!} - \frac{n(n-1)(n-2)}{3!} + \dots + (-1)^n \frac{n!}{n!}$$

$$= {}^n C_0 - {}^n C_1 + {}^n C_2 - {}^n C_3 + \dots + (-1)^n {}^n C_n$$

$$= (1 - 1)^n = 0$$

$$118. \text{ (b) } \lim_{x \rightarrow a} \frac{f(a)g'(x) - g(a)f'(x)}{g'(x) - f'(x)} = 4$$

(By Applying L' Hospital rule)

$$\lim_{x \rightarrow a} \frac{k g'(x) - k f'(x)}{g'(x) - f'(x)} = 4 \quad \therefore k = 4.$$

$$119. \text{ (a) } \text{ Given that } y = (x + \sqrt{1+x^2})^n \quad \dots(i)$$

Differentiating both sides w.r. to x

$$\frac{dy}{dx} = n(x + \sqrt{1+x^2})^{n-1} \left(1 + \frac{1}{2}(1+x^2)^{-1/2} \cdot 2x \right)$$

$$\frac{dy}{dx} = n(x + \sqrt{1+x^2})^{n-1} \frac{(\sqrt{1+x^2} + x)}{\sqrt{1+x^2}}$$

$$= \frac{n(\sqrt{1+x^2} + x)^n}{\sqrt{1+x^2}}$$

$$\text{or } \sqrt{1+x^2} \frac{dy}{dx} = ny \quad \text{[from (i)]}$$

$$\Rightarrow \sqrt{1+x^2} y_1 = ny \quad (\because y_1 = \frac{dy}{dx}) \text{ Squaring both sides,}$$

$$\text{we get } (1+x^2)y_1^2 = n^2 y^2$$

Differentiating it w.r. to x ,

$$(1+x^2)2y_1 y_2 + y_1^2 \cdot 2x = n^2 \cdot 2y y_1$$

$$\Rightarrow (1+x^2)y_2 + xy_1 = n^2 y$$

$$120. \text{ (a) } \text{ Let } f(x) = \frac{ax^3}{3} + \frac{bx^2}{2} + cx$$

$$\Rightarrow f(0) = 0 \text{ and}$$

$$f(1) = \frac{a}{3} + \frac{b}{2} + c = \frac{2a+3b+6c}{6} = 0$$

Also $f(x)$ is continuous and differentiable in $[0, 1]$ and $[0, 1[$. So by Rolle's theorem, $f'(x) = 0$.

i.e. $ax^2 + bx + c = 0$ has at least one root in $[0, 1]$.

