

# Chapter

# The p-Block Elements

## (Group 15,16,17,18)



### Topic-1: Group-15 Elements (Nitrogen Family)




#### 1 MCQs with One Correct Answer


- The species formed on fluorination of phosphorus pentachloride in a polar organic solvent are  
(a)  $[\text{PF}_4]^+[\text{PF}_6]^-$  and  $[\text{PCl}_4]^+[\text{PF}_6]^-$  [Adv. 2024]  
(b)  $[\text{PCl}_4]^+[\text{PCl}_4\text{F}_2]^-$  and  $[\text{PCl}_4]^+[\text{PF}_6]^-$   
(c)  $\text{PF}_3$  and  $\text{PCl}_3$   
(d)  $\text{PF}_5$  and  $\text{PCl}_3$
- At room temperature, disproportionation of an aqueous solution of *in situ* generated nitrous acid ( $\text{HNO}_2$ ) gives the species [Adv. 2024]  
(a)  $\text{H}_3\text{O}^+$ ,  $\text{NO}_3^-$  and  $\text{NO}$  (b)  $\text{H}_3\text{O}^+$ ,  $\text{NO}_3^-$  and  $\text{NO}_2$   
(c)  $\text{H}_3\text{O}^+$ ,  $\text{NO}^-$  and  $\text{NO}_2$  (d)  $\text{H}_3\text{O}^+$ ,  $\text{NO}_3^-$  and  $\text{N}_2\text{O}$
- The order of the oxidation state of the phosphorus atom in  $\text{H}_3\text{PO}_2$ ,  $\text{H}_3\text{PO}_3$ ,  $\text{H}_3\text{PO}_4$  and  $\text{H}_4\text{P}_2\text{O}_6$  is [Adv. 2017]  
(a)  $\text{H}_3\text{PO}_3 > \text{H}_3\text{PO}_2 > \text{H}_3\text{PO}_4 > \text{H}_4\text{P}_2\text{O}_6$   
(b)  $\text{H}_3\text{PO}_4 > \text{H}_3\text{PO}_2 > \text{H}_3\text{PO}_3 > \text{H}_4\text{P}_2\text{O}_6$   
(c)  $\text{H}_3\text{PO}_4 > \text{H}_4\text{P}_2\text{O}_6 > \text{H}_3\text{PO}_3 > \text{H}_3\text{PO}_2$   
(d)  $\text{H}_3\text{PO}_2 > \text{H}_3\text{PO}_3 > \text{H}_4\text{P}_2\text{O}_6 > \text{H}_3\text{PO}_4$
- The product formed in the reaction of  $\text{SOCl}_2$  with white phosphorous is [Adv. 2014]  
(a)  $\text{PCl}_3$  (b)  $\text{SO}_2\text{Cl}_2$  (c)  $\text{SCl}_2$  (d)  $\text{POCl}_3$
- Concentrated nitric acid, upon long standing, turns yellow brown due to the formation of [Adv. 2013]  
(a)  $\text{NO}$  (b)  $\text{NO}_2$  (c)  $\text{N}_2\text{O}$  (d)  $\text{N}_2\text{O}_4$
- The reaction of white phosphorus with aqueous  $\text{NaOH}$  gives phosphine along with another phosphorus containing compound. The reaction type; the oxidation states of phosphorus in phosphine and the other product are respectively [2012]  
(a) redox reaction; -3 and -5  
(b) redox reaction; +3 and +5  
(c) disproportionation reaction; -3 and +5  
(d) disproportionation reaction; -3 and +3
- Which ordering of compounds is according to the decreasing order of the oxidation state of nitrogen? [2012]  
(a)  $\text{HNO}_3, \text{NO}, \text{NH}_4\text{Cl}, \text{N}_2$   
(b)  $\text{HNO}_3, \text{NO}, \text{N}_2, \text{NH}_4\text{Cl}$   
(c)  $\text{HNO}_3, \text{NH}_4\text{Cl}, \text{NO}, \text{N}_2$   
(d)  $\text{NO}, \text{HNO}_3, \text{NH}_4\text{Cl}, \text{N}_2$
- Extra pure  $\text{N}_2$  can be obtained by heating [2011]  
(a)  $\text{NH}_3$  with  $\text{CuO}$  (b)  $\text{NH}_4\text{NO}_3$   
(c)  $(\text{NH}_4)_2\text{Cr}_2\text{O}_7$  (d)  $\text{Ba}(\text{N}_3)_2$
- The reaction of  $\text{P}_4$  with  $X$  leads selectively to  $\text{P}_4\text{O}_6$ . The  $X$  is [2009]  
(a) Dry  $\text{O}_2$   
(b) A mixture of  $\text{O}_2$  and  $\text{N}_2$   
(c) Moist  $\text{O}_2$   
(d)  $\text{O}_2$  in the presence of aqueous  $\text{NaOH}$
- The percentage of  $\pi$ -character in the orbitals forming  $\text{P}-\text{P}$  bonds in  $\text{P}_4$  is [2007]  
(a) 25 (b) 33 (c) 50 (d) 75
- When  $\text{PbO}_2$  reacts with conc.  $\text{HNO}_3$ , the gas evolved is [2005S]  
(a)  $\text{NO}_2$  (b)  $\text{O}_2$  (c)  $\text{N}_2$  (d)  $\text{N}_2\text{O}$
- Blue liquid which is obtained on reacting equimolar amounts of two gases at  $-30^\circ\text{C}$  is? [2005S]  
(a)  $\text{N}_2\text{O}$  (b)  $\text{N}_2\text{O}_3$  (c)  $\text{N}_2\text{O}_4$  (d)  $\text{N}_2\text{O}_5$
- Which is the most thermodynamically stable allotropic form of phosphorus? [2005S]  
(a) red (b) white (c) black (d) yellow
- For  $\text{H}_3\text{PO}_3$  and  $\text{H}_3\text{PO}_4$ , the correct choice is: [2003S]  
(a)  $\text{H}_3\text{PO}_3$  is dibasic and reducing  
(b)  $\text{H}_3\text{PO}_3$  is dibasic and non-reducing  
(c)  $\text{H}_3\text{PO}_4$  is tribasic and reducing  
(d)  $\text{H}_3\text{PO}_3$  is tribasic and non-reducing
- Ammonia can be dried by [2000S]  
(a) conc.  $\text{H}_2\text{SO}_4$  (b)  $\text{P}_4\text{O}_{10}$   
(c)  $\text{CaO}$  (d) anhydrous  $\text{CaCl}_2$
- The number of  $\text{P}-\text{O}-\text{P}$  bonds in cyclic metaphosphoric acid is [2000S]  
(a) zero (b) two (c) three (d) four
- In compounds of type  $\text{ECl}_3$ , where  $\text{E} = \text{B}, \text{P}, \text{As}$  or  $\text{Bi}$ , the angles  $\text{Cl}-\text{E}-\text{Cl}$  for different  $\text{E}$  are in the order [1999 - 2 Marks]  
(a)  $\text{B} > \text{P} = \text{As} = \text{Bi}$  (b)  $\text{B} > \text{P} > \text{As} > \text{Bi}$   
(c)  $\text{B} < \text{P} = \text{As} = \text{Bi}$  (d)  $\text{B} < \text{P} < \text{As} < \text{Bi}$
- On heating ammonium dichromate, the gas evolved is [1999 - 2 Marks]  
(a) oxygen (b) ammonia  
(c) nitrous oxide (d) nitrogen



19. One mole of calcium phosphide on reaction with excess water gives [1999 - 2 Marks]  
 (a) one mole of phosphine  
 (b) two moles of phosphoric acid  
 (c) two moles of phosphine  
 (d) one mole of phosphorus pentoxide
20. Amongst the trihalides of nitrogen which one is least basic? [1987 - 1 Mark]  
 (a)  $\text{NF}_3$  (b)  $\text{NCl}_3$   
 (c)  $\text{NBr}_3$  (d)  $\text{NI}_3$
21. Which of the following oxides of nitrogen is a coloured gas? [1987 - 1 Mark]  
 (a)  $\text{N}_2\text{O}$  (b)  $\text{NO}$  (c)  $\text{N}_2\text{O}_5$  (d)  $\text{NO}_2$
22. The bonds present in  $\text{N}_2\text{O}_5$  are : (1986 - 1 Mark)  
 (a) only ionic (b) covalent and coordinate  
 (c) only covalent (d) covalent and ionic
23. Nitrogen dioxide cannot be obtained by heating : [1985 - 1 Mark]  
 (a)  $\text{KNO}_3$  (b)  $\text{Pb}(\text{NO}_3)_2$  (c)  $\text{Cu}(\text{NO}_3)_2$  (d)  $\text{AgNO}_3$
24. Ammonia can be dried by [1980]  
 (a) Conc.  $\text{H}_2\text{SO}_4$  (b)  $\text{P}_2\text{O}_5$   
 (c) Anhydrous  $\text{CuSO}_4$  (d) none
25. The reddish brown coloured gas formed when nitric oxide is oxidised by air is [1979]  
 (a)  $\text{N}_2\text{O}_5$  (b)  $\text{N}_2\text{O}_4$  (c)  $\text{NO}_2$  (d)  $\text{N}_2\text{O}_3$

 2 Integer Value Answer


26. The total number of lone pairs of electrons in  $\text{N}_2\text{O}_3$  is [Adv. 2015]
27. Among the following, the number of compounds than can react with  $\text{PCl}_5$  to give  $\text{POCl}_3$  is [2011]  
 $\text{O}_2, \text{CO}_2, \text{SO}_2, \text{H}_2\text{O}, \text{H}_2\text{SO}_4, \text{P}_4\text{O}_{10}$

 3 Numeric / New Stem Based Questions


28. Dissolving 1.24 g of white phosphorous in boiling  $\text{NaOH}$  solution in an inert atmosphere gives a gas Q. The amount of  $\text{CuSO}_4$  (in g) required to completely consume the gas Q is [Adv. 2022]  
 [Given: Atomic mass of H = 1, O = 16, Na = 23, P = 31, S = 32, Cu = 63]


 4 Fill in the Blanks


29. In  $\text{P}_4\text{O}_{10}$ , the number of oxygen atoms bonded to each phosphorus atom is ..... [1992 - 1 Mark]
30. The basicity of phosphorous acid ( $\text{H}_3\text{PO}_3$ ) is ..... [1990 - 1 Mark]
31. .... phosphorus is reactive because of its highly strained tetrahedral structure. [1987 - 1 Mark]
32. The lowest possible oxidation state of nitrogen is [1980]

 5 True / False

33. Nitric oxide, though an odd electron molecule, is diamagnetic in liquid state. [1991 - 1 Mark]
34. The H-N-H bond angle in  $\text{NH}_3$  is greater than the H-As-H bond angle in  $\text{AsH}_3$ . [1984 - 1 Mark]
35. Red phosphorus is less volatile than white phosphorus because the former has a tetrahedral structure. [1982 - 1 Mark]

 6 MCQs with One or More than One Correct Answer

36. The correct statement(s) related to oxoacids of phosphorous is(are) [Adv. 2021]  
 (a) Upon heating,  $\text{H}_3\text{PO}_3$  undergoes disproportionation reaction to produce  $\text{H}_3\text{PO}_4$  and  $\text{PH}_3$ .  
 (b) While  $\text{H}_3\text{PO}_3$  can act as reducing agent,  $\text{H}_3\text{PO}_4$  cannot.  
 (c)  $\text{H}_3\text{PO}_3$  is a monobasic acid.  
 (d) The H atom of P-H bond in  $\text{H}_3\text{PO}_3$  is not ionizable in water.
37. Based on the compounds of group 15 elements, the correct statement(s) is (are) [Adv. 2018]  
 (a)  $\text{Bi}_2\text{O}_5$  is more basic than  $\text{N}_2\text{O}_5$   
 (b)  $\text{NF}_3$  is more covalent than  $\text{BiF}_3$   
 (c)  $\text{PH}_3$  boils at lower temperature than  $\text{NH}_3$   
 (d) The N-N single bond is stronger than the P-P single bond
38. The compound(s) which generate(s)  $\text{N}_2$  gas upon thermal decomposition below  $300^\circ\text{C}$  is (are) [Adv. 2018]  
 (a)  $\text{NH}_4\text{NO}_3$  (b)  $(\text{NH}_4)_2\text{Cr}_2\text{O}_7$   
 (c)  $\text{Ba}(\text{N}_3)_2$  (d)  $\text{Mg}_3\text{N}_2$
39. The nitrogen containing compound produced in the reaction of  $\text{HNO}_3$  with  $\text{P}_4\text{O}_{10}$  [Adv. 2016]  
 (a) can also be prepared by reaction of  $\text{P}_4$  and  $\text{HNO}_3$   
 (b) is diamagnetic  
 (c) contains one N-N bond  
 (d) reacts with Na metal producing a brown gas
40. The nitrogen oxide(s) that contain(s) N-N bond(s) is(are) [2009]  
 (a)  $\text{N}_2\text{O}$  (b)  $\text{N}_2\text{O}_3$  (c)  $\text{N}_2\text{O}_4$  (d)  $\text{N}_2\text{O}_5$
41. A solution of colourless salt H on boiling with excess  $\text{NaOH}$  produces a non-flammable gas. The gas evolution ceases after sometime. Upon addition of Zn dust to the same solution, the gas evolution restarts. The colourless salt (s) H is (are) [2008]  
 (a)  $\text{NH}_4\text{NO}_3$  (b)  $\text{NH}_4\text{NO}_2$  (c)  $\text{NH}_4\text{Cl}$  (d)  $(\text{NH}_4)_2\text{SO}_4$
42. Ammonia, on reaction with hypochlorite anion, can form [1999 - 3 Marks]  
 (a)  $\text{NO}$  (b)  $\text{NH}_4\text{Cl}$  (c)  $\text{N}_2\text{H}_4$  (d)  $\text{HNO}_2$
43. White phosphorus ( $\text{P}_4$ ) has [1998 - 2 Marks]  
 (a) six P-P single bonds  
 (b) four P-P single bonds  
 (c) four lone pairs of electrons  
 (d) PPP angle of  $60^\circ$
44. Sodium nitrate decomposes above  $800^\circ\text{C}$  to give [1998 - 2 Marks]  
 (a)  $\text{N}_2$  (b)  $\text{O}_2$   
 (c)  $\text{NO}_2$  (d)  $\text{Na}_2\text{O}$
45. Nitrogen(I) oxide is produced by : [1989 - 1 Mark]  
 (a) thermal decomposition of ammonium nitrate  
 (b) disproportionation of  $\text{N}_2\text{O}_4$   
 (c) thermal decomposition of ammonium nitrite  
 (d) interaction of hydroxylamine and nitrous acid.
-  7 Match the Following
46. Match the reactions (in the given stoichiometry of the reactants) in List-I with one of their products given in List-II and choose the correct option. [Adv. 2023]
- | List-I  | List-II   |
|---|---|
| (P) $\text{P}_2\text{O}_3 + 3\text{H}_2\text{O} \rightarrow$      | (1) $\text{P}(\text{O})(\text{OCH}_3)\text{Cl}_2$ |
| (Q) $\text{P}_4 + 3\text{NaOH} + 3\text{H}_2\text{O} \rightarrow$ | (2) $\text{H}_3\text{PO}_3$                       |



- (a)  $\text{P} \rightarrow 2; \text{Q} \rightarrow 3; \text{R} \rightarrow 1; \text{S} \rightarrow 5$   
 (b)  $\text{P} \rightarrow 3; \text{Q} \rightarrow 5; \text{R} \rightarrow 4; \text{S} \rightarrow 2$   
 (c)  $\text{P} \rightarrow 5; \text{Q} \rightarrow 2; \text{R} \rightarrow 1; \text{S} \rightarrow 3$   
 (d)  $\text{P} \rightarrow 2; \text{Q} \rightarrow 3; \text{R} \rightarrow 4; \text{S} \rightarrow 5$

47. Match the following, choosing one item from column X and the appropriate item from column Y. [1983 - 2 Marks]

X	Y
(i) Haber	(a) Activation energy
(ii) Graham	(b) Diffusion of gases
(iii) Arrhenius	(c) Octet rule
(iv) Lewis	(d) Ammonia synthesis

**8 Comprehension/Passage Based Questions**

There are some deposits of nitrates and phosphates in earth's crust. Nitrates are more soluble in water. Nitrates are difficult to reduce under the laboratory conditions but microbes do it easily. Ammonia forms large number of complexes with transition metal ions. Hybridization easily explains the ease of sigma donation capability of  $\text{NH}_3$  and  $\text{PH}_3$ . Phosphine is a flammable gas and is prepared from white phosphorous. [2008]

48. Among the following, the correct statement is  
 (a) Phosphates have no biological significance in humans  
 (b) Between nitrates and phosphates, phosphates are less abundant in earth's crust  
 (c) Between nitrates and phosphates, nitrates are less abundant in earth's crust  
 (d) Oxidation of nitrates is possible in soil

49. Among the following, the correct statement is  
 (a) Between  $\text{NH}_3$  and  $\text{PH}_3$ ,  $\text{NH}_3$  is a better electron donor because the lone pair of electrons occupies spherical  $s$ -orbital and is less directional  
 (b) Between  $\text{NH}_3$  and  $\text{PH}_3$ ,  $\text{PH}_3$  is a better electron donor because the lone pair of electrons occupies  $sp^3$  orbital and is more directional  
 (c) Between  $\text{NH}_3$  and  $\text{PH}_3$ ,  $\text{NH}_3$  is a better electron donor because the lone pair of electrons occupies  $sp^3$  orbital and is more directional  
 (d) Between  $\text{NH}_3$  and  $\text{PH}_3$ ,  $\text{PH}_3$  is a better electron donor because the lone pair of electrons occupies spherical  $s$ -orbital and is less directional

50. White phosphorus on reaction with  $\text{NaOH}$  gives  $\text{PH}_3$  as one of the products. This is a  
 (a) dimerization reaction  
 (b) disproportionation reaction  
 (c) condensation reaction  
 (d) precipitation reaction

**9 Assertion and Reason/Statement Type Questions**

Each question contains **STATEMENT-1 (Assertion)** and **STATEMENT-2 (Reason)**. Each question has 4 choices (a), (b), (c) and (d) out of which **ONLY ONE** is correct. Mark your answer as


- (a) If both Statement -1 and Statement -2 are correct, and Statement -2 is the correct explanation of the Statement -2.  
 (b) If both Statement -1 and Statement -2 are correct, but Statement -2 is not the correct explanation of the Statement -1.

- (c) If Statement -1 is correct but Statement -2 is incorrect.  
 (d) If Statement -1 is incorrect but Statement -2 is correct.

51. **Statement-1** : Although  $\text{PF}_5$ ,  $\text{PCl}_5$  and  $\text{PBr}_5$  are known, the pentahalides of nitrogen have not been observed  
**Statement-2** : Phosphorus has lower electronegativity than nitrogen. [1994 - 2 Marks]

**10 Subjective Problems**

52. Write the structures of  $(\text{CH}_3)_3\text{N}$  and  $(\text{Me}_3\text{Si})_3\text{N}$ . Are they isostructural? Justify your answer. [2005 - 2 Marks]  
 53. How many grams of  $\text{CaO}$  are required to neutralize 852 g of  $\text{P}_4\text{O}_{10}$ ? Draw structure of  $\text{P}_4\text{O}_{10}$  molecule. [2005 - 2 Marks]  
 54. Give reason(s) why elemental nitrogen exists as a diatomic molecule whereas elemental phosphorus as a tetraatomic molecule. [2000 - 2 Marks]  
 55. The Haber process can be represented by the following scheme;



Identify A, B, C, D and E. [1999 - 5 Marks]

56. Reaction of phosphoric acid with  $\text{Ca}_3(\text{PO}_4)_2\text{F}$  yields a fertilizer "triple superphosphate". Represent the same through balanced chemical equation. [1998 - 2 Marks]  
 57. A soluble compound of a poisonous element M, when heated with  $\text{Zn}/\text{H}_2\text{SO}_4$ , gives a colourless and extremely poisonous gaseous compound N, which on passing through a heated tube gives a silvery mirror of element M. Identify M and N. [1997 - 2 Marks]  
 58. Draw the structure of  $\text{P}_4\text{O}_{10}$  and identify the number of single and double P—O bonds. [1996 - 3 Marks]  
 59. Complete and balance the following chemical reactions :  
 (i) Red phosphorus is reacted with iodine in presence of water. [1992 - 1 Mark]  
 $\text{P} + \text{I}_2 + \text{H}_2\text{O} \rightarrow \dots + \dots$   
 (ii) Anhydrous potassium nitrate is heated with excess of metallic potassium. [1992 - 1 Mark]  
 $\text{KNO}_3(\text{s}) + \text{K}(\text{s}) \rightarrow \dots + \dots$   
 60. Arrange the following in : [1991 - 1 Mark]  
 Increasing order of extent of hydrolysis :  
 $\text{CCl}_4, \text{MgCl}_2, \text{AlCl}_3, \text{PCl}_5, \text{SiCl}_4$

61. Write down the balanced equations for the reactions when:  
 (i) a mixture of potassium chlorate, oxalic acid and sulphuric acid is heated; [1985 - 1 Mark]  
 (ii) ammonium sulphate is heated with a mixture of nitric oxide and nitrogen dioxide. [1985 - 1 Mark]
62. Write down the resonance structures of nitrous oxide. [1985 - 2 Marks]

OR

Write the two resonance structures of  $N_2O$  that satisfy the octet rule. [1990 - 1 Mark]

63. State with balanced equations what happens when :  
 (i)  $P_4O_{10} + PCl_5 \rightarrow$  [1998 - 1 Mark]  
 (ii) Phosphorus is treated with concentrated nitric acid. [1997 - 1 Mark]

OR

Manufacture of phosphoric acid from phosphorus. [1997 - 1 Mark]

- (iii) Elemental phosphorus reacts with conc.  $HNO_3$  to give phosphoric acid. [1991 - 1 Mark]  
 (iv) Nitrogen is obtained in the reaction of aqueous ammonia with potassium permanganate. [1991 - 1 Mark]  
 (v) Sodium nitrite is produced by absorbing the oxides of nitrogen in aqueous solution of washing soda. [1991 - 1 Mark]  
 (vi) Aqueous solution of sodium nitrate is heated with zinc dust and caustic soda solution. [1990 - 1 Mark]  
 (vii) Write balanced equations for the preparation of ammonium sulphate from gypsum, ammonia and carbon dioxide. [1990 - 1 Mark]

- (viii) Write balanced equations for the preparation of phosphine from CaO and white phosphorus. [1990 - 2 Marks]  
 (ix) Hypophosphorous acid is heated. [1989 - 1 Mark]  
 (x) Phosphorus reacts with nitric acid to give equimolar ratio of nitric oxide and nitrogen dioxide. [1988 - 1 Mark]  
 (xi) Dilute nitric acid is slowly reacted with metallic tin. [1987 - 1 Mark]  
 (xii) White phosphorous ( $P_4$ ) is boiled with a strong solution of sodium hydroxide in an inert atmosphere. [1982/87 - 1 Mark]

64. Give reasons for the following :  
 (i) The experimentally determined N-F bond length in  $NF_3$  is greater than the sum of the single covalent bond radii of N and F. [1995 - 2 Marks]  
 (ii) Ammonium chloride is acidic in liquid ammonia solvent. [1991 - 1 Mark]  
 (iii) Phosphine has lower boiling point than ammonia. [1989 - 1 Mark]  
 (iv)  $H_3PO_3$  is a dibasic acid. [1989 - 1 Mark]  
 (v) Orthophosphorus acid is not tribasic acid. [1987 - 1 Mark]  
 (vi) A bottle of liquor ammonia should be cooled before opening the stopper. [1983 - 1 Mark]  
 (vii) Orthophosphoric acid,  $H_3PO_4$ , is tribasic, but phosphorous acid,  $H_3PO_3$ , is dibasic. [1982 - 1 Mark]
65. Give structural formula for the following :  
 (i) Phosphorous acid,  $H_3PO_3$  [1981 - 1 Mark]  
 (ii) Pyrophosphoric acid,  $H_4P_2O_7$  [1981 - 1 Mark]




## Topic-2: Group-16 Elements (Oxygen Family)



### 1 MCQs with One Correct Answer

1. In the following reaction sequence in aqueous solution, the species X, Y and Z, respectively, are [Adv.2016]



- (a)  $[Ag(S_2O_3)_2]^{3-}$ ,  $Ag_2S_2O_3$ ,  $Ag_2S$   
 (b)  $[Ag(S_2O_3)_3]^{5-}$ ,  $Ag_2SO_3$ ,  $Ag_2S$   
 (c)  $[Ag(SO_3)_2]^{3-}$ ,  $Ag_2S_2O_3$ ,  $Ag$   
 (d)  $[Ag(SO_3)_3]^{3-}$ ,  $Ag_2SO_4$ ,  $Ag$
2. Aqueous solution of  $Na_2S_2O_3$  on reaction with  $Cl_2$  gives –  
 (a)  $Na_2S_4O_6$  (b)  $NaHSO_4$  [2008]  
 (c)  $NaCl$  (d)  $NaOH$
3. Which of the following is not oxidized by  $O_3$ ? [2005S]  
 (a) KI (b)  $FeSO_4$  (c)  $KMnO_4$  (d)  $K_2MnO_4$
4. The acid having O-O bond is [2004S]  
 (a)  $H_2S_2O_3$  (b)  $H_2S_2O_6$  (c)  $H_2S_2O_8$  (d)  $H_2S_4O_6$
5. The number of S-S bonds in sulphur trioxide trimer ( $S_3O_9$ ) is [2001S]  
 (a) three (b) two (c) one (d) zero
6. Sodium thiosulphate is prepared by [1996 - 1 Mark]  
 (a) reducing  $Na_2SO_4$  solution with  $H_2S$   
 (b) boiling  $Na_2SO_3$  solution with S in alkaline medium  
 (c) neutralising  $H_2S_2O_3$  solution with NaOH  
 (d) boiling  $Na_2SO_3$  solution with S in acidic medium
7. Hydrolysis of one mole of peroxodisulphuric acid produces  
 (a) two moles of sulphuric acid [1996 - 1 Mark]  
 (b) two moles of peroxomonosulphuric acid  
 (c) one mole of sulphuric acid and one mole of peroxomonosulphuric acid  
 (d) one mole of sulphuric acid, one mole of peroxomonosulphuric acid and one mole of hydrogen peroxide.
8.  $H_2SO_4$  cannot be used to prepare HBr from NaBr as it : [1995S]  
 (a) reacts slowly with NaBr (b) oxidises HBr  
 (c) reduces HBr (d) disproportionates HBr
9. The compound which gives off oxygen on moderate heating is : [1986 - 1 Mark]  
 (a) cupric oxide (b) mercuric oxide  
 (c) zinc oxide (d) aluminium oxide
10. A gas that cannot be collected over water is :  
 (a)  $N_2$  (b)  $O_2$  [1985 - 1 Mark]  
 (c)  $SO_2$  (d)  $PH_3$



11. Which of the following is coloured [1980]  
 (a) NO (b) N<sub>2</sub>O (c) SO<sub>3</sub> (d) None

2 Integer Value Answer

12. The total number of compounds having at least one bridging oxo group among the molecules given below is \_\_\_\_\_. N<sub>2</sub>O<sub>3</sub>, N<sub>2</sub>O<sub>5</sub>, P<sub>4</sub>O<sub>6</sub>, P<sub>4</sub>O<sub>7</sub>, H<sub>4</sub>P<sub>2</sub>O<sub>5</sub>, H<sub>5</sub>P<sub>3</sub>O<sub>10</sub>, H<sub>2</sub>S<sub>2</sub>O<sub>3</sub>, H<sub>2</sub>S<sub>2</sub>O<sub>5</sub> [Adv. 2018]

3 Numeric / New Stem Based Questions

13. The amount of water produced (in g) in the oxidation of 1 mole of rhombic sulphur by conc. HNO<sub>3</sub> to a compound with the highest oxidation state of sulphur is \_\_\_\_ (Given data : Molar mass of water = 18g mol<sup>-1</sup>) [Adv. 2019]

4 Fill in the Blanks

14. The lead chamber process involves oxidation of SO<sub>2</sub> by atomic oxygen under the influence of ..... as catalyst. [1992 - 1 Mark]

6 MCQs with One or More than One Correct Answer

15. The compound(s) having peroxide linkage is(are) [Adv. 2024]  
 (a) H<sub>2</sub>S<sub>2</sub>O<sub>7</sub> (b) H<sub>2</sub>S<sub>2</sub>O<sub>8</sub>  
 (c) H<sub>2</sub>S<sub>2</sub>O<sub>5</sub> (d) H<sub>2</sub>SO<sub>5</sub>
16. The correct statement(s) about O<sub>3</sub> is(are) [Adv. 2013-II]  
 (a) O—O bond lengths are equal  
 (b) Thermal decomposition of O<sub>3</sub> is endothermic  
 (c) O<sub>3</sub> is diamagnetic in nature  
 (d) O<sub>3</sub> has a bent structure
17. Which of the following halides react(s) with AgNO<sub>3</sub>(aq) to give a precipitate that dissolves in Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>(aq)? [2012]  
 (a) HCl (b) HF  
 (c) HBr (d) HI

8 Comprehension Passage Based Questions

Passage-1

Upon heating KClO<sub>3</sub> in the presence of catalytic amount of MnO<sub>2</sub>, a gas *W* is formed. Excess amount of *W* reacts with white phosphorus to give *X*. The reaction of *X* with pure HNO<sub>3</sub> gives *Y* and *Z*. [Adv. 2017]




18. *W* and *X* are, respectively  
 (a) O<sub>3</sub> and P<sub>4</sub>O<sub>6</sub> (b) O<sub>2</sub> and P<sub>4</sub>O<sub>6</sub>  
 (c) O<sub>2</sub> and P<sub>4</sub>O<sub>10</sub> (d) O<sub>3</sub> and P<sub>4</sub>O<sub>10</sub>
19. *Y* and *Z* are, respectively  
 (a) N<sub>2</sub>O<sub>3</sub> and H<sub>3</sub>PO<sub>4</sub> (b) N<sub>2</sub>O<sub>5</sub> and HPO<sub>3</sub>  
 (c) N<sub>2</sub>O<sub>4</sub> and HPO<sub>3</sub> (d) N<sub>2</sub>O<sub>4</sub> and H<sub>3</sub>PO<sub>3</sub>

Passage-2

The reactions of Cl<sub>2</sub> gas with cold-dilute and hot-concentrated NaOH in water give sodium salts of two (different) oxoacids of chlorine, *P* and *Q*, respectively. The Cl<sub>2</sub> gas reacts with SO<sub>2</sub> gas, in presence of charcoal, to give a product *R*. *R* reacts with white phosphorus to give a compound *S*. On hydrolysis, *S* gives an oxoacid of phosphorus, *T*. [Adv. 2013]


20. *P* and *Q*, respectively, are the sodium salts of  
 (a) Hypochlorous and chloric acids  
 (b) Hypochlorous and chlorous acids  
 (c) Chloric and perchloric acids  
 (d) Chloric and hypochlorous acids
21. *R*, *S* and *T* respectively, are  
 (a) SO<sub>2</sub>Cl<sub>2</sub>, PCl<sub>5</sub> and H<sub>3</sub>PO<sub>4</sub> (b) SO<sub>2</sub>Cl<sub>2</sub>, PCl<sub>3</sub> and H<sub>3</sub>PO<sub>3</sub>  
 (c) SOCl<sub>2</sub>, PCl<sub>3</sub> and H<sub>3</sub>PO<sub>2</sub> (d) SOCl<sub>2</sub>, PCl<sub>5</sub> and H<sub>3</sub>PO<sub>4</sub>

10 Subjective Problems



Identify the missing compounds. Give the equation from *A* to *B* and *A* to *C*. [2005 - 4 Marks]

23. Identify the following: [2003 - 4 Marks]



- Also mention the oxidation state of S in all the compounds.
24. In the contact process for industrial manufacture of sulphuric acid, some amount of sulphuric acid is used as a starting material. Explain briefly. What is the catalyst used in the oxidation of SO<sub>2</sub>? [1999 - 4 Marks]
25. In the following equation, [1999 - 6 Marks]  
 $A + 2B + \text{H}_2\text{O} \rightarrow C + 2D$   
 ( $A = \text{HNO}_2$ ,  $B = \text{H}_2\text{SO}_3$ ,  $C = \text{NH}_2\text{OH}$ ). Identify *D*. Draw the structures of *A*, *B*, *C* and *D*.

26. Thionyl chloride can be synthesized by chlorinating SO<sub>2</sub> using PCl<sub>5</sub>. Thionyl chloride is used to prepare anhydrous ferric chloride starting from its hexahydrated salt. Alternatively, the anhydrous ferric chloride can also be prepared from its hexahydrated salt by treating with 2, 2 - dimethoxypropane. Discuss all this using balanced chemical equations. [1998 - 6 Marks]

27.  $\text{PbS} \xrightarrow[\text{air}]{\text{heat in}}$  A + PbS  $\xrightarrow{\text{B}}$  Pb + SO<sub>2</sub>; Identify *A* and *B*. [1991 - 2 Marks]

28. Write the two resonance structures of ozone which satisfy the octet rule. [1991 - 1 Mark]

29. Mention the products formed in the following : Sulphur dioxide gas, water vapour and air are passed over heated sodium chloride. [1986 - 1 Mark]

30. Arrange the following in : CO<sub>2</sub>, N<sub>2</sub>O<sub>5</sub>, SiO<sub>2</sub>, SO<sub>3</sub> in the order of increasing acidic character. [1988 - 1 Mark]

31. What happens when :  
 (i) hydrogen sulphide is bubbled through an aqueous solution of sulphur dioxide. [1985 - 1 Mark]  
 (ii) tin is treated with concentrated nitric acid.

[1985 - 1 Mark]

- (iii) Pb<sub>3</sub>O<sub>4</sub> is treated with nitric acid. [1985 - 1 Mark]
32. Show with equations, how the following compound is prepared (equations need not be balanced) : sodium thiosulphate from sodium sulphite. [1982 - 1 Mark]

33. State with balanced equations what happens when : Sulphur is precipitated in the reaction of hydrogen sulphide with sodium bisulphite solution. [1991 - 1 Mark]

34. Give reasons for the following :  
 (i) Sulphur dioxide is a more powerful reducing agent in an alkaline medium than in acidic medium. [1992 - 1 Mark]

(ii) Valency of oxygen is generally two whereas sulphur shows valency of two, four and six. [1988 - 1 Mark]

(iii) Sulphur melts to a clear mobile liquid at 119°C, but on further heating above 160°C, it becomes viscous. [1981 - 1 Mark]



### Topic-3: Group-17 Elements (Halogen Family)



#### 1 MCQs with One Correct Answer

- The reaction of  $\text{HClO}_3$  with  $\text{HCl}$  gives a paramagnetic gas, which upon reaction with  $\text{O}_3$  produces [Adv. 2022]  
(a)  $\text{Cl}_2\text{O}$  (b)  $\text{ClO}_2$  (c)  $\text{Cl}_2\text{O}_6$  (d)  $\text{Cl}_2\text{O}_7$
- A colorless aqueous solution contains nitrates of two metals, X and Y. When it was added to an aqueous solution of  $\text{NaCl}$ , a white precipitate was formed. This precipitate was found to be partly soluble in hot water to give a residue P and a solution Q. The residue P was soluble in aq.  $\text{NH}_3$  and also in excess sodium thiosulfate. The hot solution Q gave a yellow precipitate with  $\text{KI}$ . The metals X and Y, respectively, are [Adv. 2020]  
(a) Ag and Pb (b) Ag and Cd  
(c) Cd and Pb (d) Cd and Zn
- The products obtained when chlorine gas reacts with cold and dilute aqueous  $\text{NaOH}$  are : [2017]  
(a)  $\text{ClO}^-$  and  $\text{ClO}_3^-$  (b)  $\text{ClO}_2^-$  and  $\text{ClO}_3^-$   
(c)  $\text{Cl}^-$  and  $\text{ClO}^-$  (d)  $\text{Cl}^-$  and  $\text{ClO}_2^-$
- Which one of the following species is not a pseudohalide? [1997 - 1 Mark]  
(a)  $\text{CNO}^-$  (b)  $\text{RCOO}^-$  (c)  $\text{OCN}^-$  (d)  $\text{NNN}^-$
- $\text{KF}$  combines with  $\text{HF}$  to form  $\text{KHF}_2$ . The compound contains the species. [1996 - 1 Mark]  
(a)  $\text{K}^+$ ,  $\text{F}^-$  and  $\text{H}^+$  (b)  $\text{K}^+$ ,  $\text{F}^-$  and  $\text{HF}$   
(c)  $\text{K}^+$  and  $[\text{HF}_2]^-$  (d)  $[\text{KHF}]^+$  and  $\text{F}^-$
- Which of the following statements is correct for  $\text{CsBr}_3$ ? [1996 - 1 Mark]  
(a) It is a covalent compound.  
(b) It contains  $\text{Cs}^{3+}$  and  $\text{Br}^-$  ions.  
(c) It contains  $\text{Cs}^+$  and  $\text{Br}_3^-$  ions  
(d) It contains  $\text{Cs}^+$ , and  $\text{Br}^-$  and lattice  $\text{Br}_2$  molecule
- Bromine can be liberated from potassium bromide solution by the action of [1987 - 1 Mark]  
(a) Iodine solution (b) Chlorine water  
(c) Sodium chloride (d) Potassium iodide
- Chlorine acts as a bleaching agent only in presence of [1983 - 1 Mark]  
(a) dry air (b) moisture  
(c) sunlight (d) pure oxygen
- $\text{HBr}$  and  $\text{HI}$  reduce sulphuric acid,  $\text{HCl}$  can reduce  $\text{KMnO}_4$  and  $\text{HF}$  can reduce [1981 - 1 Mark]  
(a)  $\text{H}_2\text{SO}_4$  (b)  $\text{KMnO}_4$   
(c)  $\text{K}_2\text{Cr}_2\text{O}_7$  (d) none of the above
- A solution of  $\text{KBr}$  is treated with each of the following. Which one would liberate bromine [1980]  
(a)  $\text{Cl}_2$  (b)  $\text{HI}$  (c)  $\text{I}_2$  (d)  $\text{SO}_2$
- Which of the following is most stable to heat [1980]  
(a)  $\text{HCl}$  (b)  $\text{HOCl}$  (c)  $\text{HBr}$  (d)  $\text{HI}$



#### 2 Integer Value Answer

- Ozonolysis of  $\text{ClO}_2$  produces an oxide of chlorine. The average oxidation state of chlorine in this oxide is \_\_\_\_\_. [Adv. 2021]
- Reaction of  $\text{Br}_2$  with  $\text{Na}_2\text{CO}_3$  in aqueous solution gives sodium bromide and sodium bromate with evolution of

$\text{CO}_2$  gas. The number of sodium bromide molecules involved in the balanced chemical equation is [2011]



#### 4 Fill in the Blanks

- Solubility of iodine in water is greatly increased by the addition of iodide ions because of the formation of ..... [1994 - 1 Mark]
- ..... acid gives hypo ..... ion. [1988 - 1 Mark]  
(hydrobromic, hypobromous, perbromic, bromide, bromite, perbromate)
- The increase in the solubility of iodine in an aqueous solution of potassium iodide is due to the formation of ..... [1982 - 1 Mark]
- ..... is a weak acid. ( $\text{HF}$ ,  $\text{HCl}$ ,  $\text{HI}$ ) [1981 - 1 Mark]
- Iodine reacts with hot  $\text{NaOH}$  solution. The products are  $\text{NaI}$  and ..... [1980]



#### 5 True / False

- $\text{HBr}$  is a stronger acid than  $\text{HI}$  because of hydrogen bonding. [1993 - 1 Mark]
- In aqueous solution, chlorine is a stronger oxidizing agent than fluorine. [1984 - 1 Mark]
- Dil.  $\text{HCl}$  oxidizes metallic  $\text{Fe}$  to  $\text{Fe}^{2+}$ . [1983 - 1 Mark]



#### 6 MCQs with One or More than One Correct Answer

- With respect to hypochlorite, chlorate and perchlorate ions, choose the correct statement(s).  
(a) The hypochlorite ion is the strongest conjugate base.  
(b) The molecular shape of only chlorate ion is influenced by the lone pair of electrons of  $\text{Cl}$ .  
(c) The hypochlorite and chlorate ions disproportionate to give rise to identical set of ions.  
(d) The hypochlorite ion oxidizes the sulfite ion.
- The colour of the  $X_2$  molecules of group 17 elements changes gradually from yellow to violet down the group. This is due to [Adv. 2017]  
(a) The physical state of  $X_2$  at room temperature changes from gas to solid down the group  
(b) Decrease in ionization energy down the group  
(c) Decrease in  $\pi^* - \sigma^*$  gap down the group  
(d) Decrease in HOMO-LUMO gap down the group
- The correct statement(s) about the oxoacids,  $\text{HClO}_4$  and  $\text{HClO}$ , is (are) [Adv. 2017]  
(a) The central atom in both  $\text{HClO}_4$  and  $\text{HClO}$  is  $sp^3$  hybridized  
(b)  $\text{HClO}_4$  is more acidic than  $\text{HClO}$  because of the resonance stabilization of its anion  
(c)  $\text{HClO}_4$  is formed in the reaction between  $\text{Cl}_2$  and  $\text{H}_2\text{O}$   
(d) The conjugate base of  $\text{HClO}_4$  is weaker base than  $\text{H}_2\text{O}$
- The compounds used as refrigerant are [1990 - 1 Mark]  
(a)  $\text{NH}_3$  (b)  $\text{CCl}_4$  (c)  $\text{CF}_4$  (d)  $\text{CF}_2\text{Cl}_2$   
(e)  $\text{CH}_2\text{F}_2$

### 8 Comprehension/Passage-Based Questions

Bleaching powder and bleach solution are produced on a large scale and used in several household products. The effectiveness of bleach solution is often measured by iodometry. [2012 - II]

26. Bleaching powder contains a salt of an oxoacid as one of its components. The anhydride of that oxoacid is  
(a)  $\text{Cl}_2\text{O}$  (b)  $\text{Cl}_2\text{O}_7$  (c)  $\text{ClO}_2$  (d)  $\text{Cl}_2\text{O}_6$
27. 25 mL of household solution was mixed with 30 mL of 0.50 M KI and 10 mL of 4N acetic acid. In the titration of the liberated iodine, 48 mL of 0.25 N  $\text{Na}_2\text{S}_2\text{O}_3$  was used to reach the end point. The molarity of the household bleach solution is  
(a) 0.48 M (b) 0.96 M (c) 0.24 M (d) 0.024 M

### 9 Assertion and Reason Statement Type Questions

Each question contains STATEMENT-1 (Assertion) and STATEMENT-2 (Reason). Each question has 4 choices (a), (b), (c) and (d) out of which ONLY ONE is correct. Mark your answer as

- (a) If both Statement -1 and Statement -2 are correct, and Statement -2 is the correct explanation of the Statement -2.  
(b) If both Statement -1 and Statement -2 are correct, but Statement -2 is not the correct explanation of the Statement -1.  
(c) If Statement -1 is correct but Statement -2 is incorrect.  
(d) If Statement -1 is incorrect but Statement -2 is correct.
28. **Statement-1:** F atom has less electron affinity than Cl atom.  
**Statement-2:** Additional electrons are repelled more effectively by 3p electrons in Cl atom than by 2p electrons in F atom. [1998 - 2 Marks]

### 10 Subjective Problems

29. Give an example of oxidation of one halide by another halogen. Explain the feasibility of the reaction. [2000 - 2 Marks]
30. Complete the following chemical equations:  
(a)  $\text{KI} + \text{Cl}_2 \rightarrow$  (b)  $\text{KClO}_3 + \text{I}_2 \rightarrow$   
Justify the formation of the products in the above reactions. [1996 - 2 Marks]
31. Gradual addition of KI solution to  $\text{Bi}(\text{NO}_3)_3$  solution initially produces a dark brown precipitate which dissolves in excess of KI to give a clear yellow solution. Write chemical equations for the above reactions. [1996 - 2 Marks]
32. Complete and balance the following chemical reactions:  
(iii)  $\text{NH}_3 + \text{NaOCl} \rightarrow \dots + \dots$  [1993 - 1 Mark]


33. Mention the products formed in the following:  
(i) Chlorine gas is bubbled through a solution of ferrous bromide. [1986 - 1 Mark]  
(ii) Iodine is added to a solution of stannous chloride. [1986 - 1 Mark]
34. Arrange the following in:  
(i)  $\text{HOCl}$ ,  $\text{HOClO}_2$ ,  $\text{HOClO}_3$ ,  $\text{HOClO}$  in increasing order of thermal stability. [1988 - 1 Mark]  
(ii) increasing bond strength  
 $\text{HCl}$ ,  $\text{HBr}$ ,  $\text{HF}$ ,  $\text{HI}$  [1986 - 1 Mark]
35. State with balanced equations what happens when:  
(i) Sodium iodate is added to a solution of sodium bisulphite. [1990 - 1 Marks]  
(ii) Sodium chlorate reacts with sulphur dioxide in dilute sulphuric acid medium. [1989 - 1 Mark]  
(iii) Sodium bromate reacts with fluorine in presence of alkali. [1989 - 1 Mark]  
(iv) Iodate ion reacts with bisulphite ion to liberate iodine. [1988 - 1 Mark]  
(v) Sodium iodate is treated with sodium bisulphite solution. [1982 - 1 Mark]
36. Give reasons for the following:  
(i) Bond dissociation energy of  $\text{F}_2$  is less than that of  $\text{Cl}_2$ . [1992 - 1 Mark]  
(ii) Fluorine cannot be prepared from fluorides by chemical oxidation. [1985 - 1 Mark]  
(iii) Anhydrous  $\text{HCl}$  is a bad conductor of electricity but aqueous  $\text{HCl}$  is a good conductor; [1985 - 1 Mark]  
(iv) In the preparation of hydrogen iodide from alkali iodides, phosphoric acid is preferred to sulphuric acid [1982 - 1 Mark]
37. Write balanced equation involved in the preparation of  
(i) Anhydrous aluminium chloride from alumina.  
(ii) Bleaching powder from slaked lime.  
(iii) Tin metal from cassiterite  
(iv) Chlorine from sodium chloride. [1979]
38. Account for the following. Limit your answer to two sentences  
(i) Hydrogen bromide cannot be prepared by action of concentrated sulphuric acid or sodium bromide.  
(ii) When a blue litmus paper is dipped into a solution of hypochlorous acid, it first turns red and then later gets decolourised. [1979]



## Topic-4: Group-18 Elements (Noble Gases)

### 1 MCQs with One Correct Answer

1. Under ambient conditions, the total number of gases released as products in the final step of the reaction scheme shown below is [Adv. 2014]



2. The shape of  $\text{XeO}_2\text{F}_2$  molecule is [2012]  
(a) trigonal bipyramidal (b) square planar  
(c) tetrahedral (d) see-saw
3. Total number of lone pair of electrons with central atom in  $\text{XeOF}_4$  is [2004S]  
(a) 0 (b) 1 (c) 2 (d) 3

### 3 Numeric / New Stem Based Questions

4. At 143 K, the reaction of  $\text{XeF}_4$  with  $\text{O}_2\text{F}_2$ , produces a xenon compound Y. The total number of lone pair(s) of electron present on the whole molecule of Y is [Adv. 2019]



7 Match the Following

5. The unbalanced chemical reactions given in List-I show missing reagent or condition (?) which are provided in List-II. Match List-I with List-II and select the correct answer using the code given below the lists

List-I	[Adv. 2013-II] List-II
P. $\text{PbO}_2 + \text{H}_2\text{SO}_4 \xrightarrow{?}$ $\text{PbSO}_4 + \text{O}_2 + \text{other product}$	1. NO
Q. $\text{Na}_2\text{S}_2\text{O}_3 + \text{H}_2\text{O} \xrightarrow{?}$ $\text{NaHSO}_4 + \text{other product}$	2. $\text{I}_2$
R. $\text{N}_2\text{H}_4 \xrightarrow{?}$ $\text{N}_2 + \text{other product}$	3. Warm
S. $\text{XeF}_2 \xrightarrow{?}$ $\text{Xe} + \text{other product}$	4. $\text{Cl}_2$

Codes :

	P	Q	R	S
(a)	4	2	3	1
(b)	3	2	1	4
(c)	1	4	2	3
(d)	3	4	2	1

6. All the compounds listed in Column I react with water. Match the result of the respective reactions with the appropriate options listed in Column II. [2010]

Column-I	Column-II
(A) $(\text{CH}_3)_2\text{SiCl}_2$	(p) Hydrogen halide formation
(B) $\text{XeF}_4$	(q) Redox reaction
(C) $\text{Cl}_2$	(r) Reacts with glass
(D) $\text{VCl}_5$	(s) Polymerization
	(t) $\text{O}_2$ formation



8 Comprehension/Passage Based Questions

The noble gases have closed-shell electronic configuration and are monoatomic gases under normal conditions. The low boiling points of the lighter noble gases are due to weak dispersion forces between the atoms and the absence of other interatomic interactions.

The direct reaction of xenon with fluorine leads to a series of compounds with oxidation numbers +2, +4 and +6.  $\text{XeF}_4$  reacts violently with water to give  $\text{XeO}_3$ . The compounds of xenon exhibit rich stereochemistry and their geometries can be deduced considering the total number of electron pairs in the valence shell.

7. Argon is used in arc welding because of its [2007]  
 (a) low reactivity with metal  
 (b) ability to lower the melting point of metal  
 (c) flammability  
 (d) high calorific value
8. The structure of  $\text{XeO}_3$  is [2007]  
 (a) linear (b) planar  
 (c) pyramidal (d) T-shaped
9.  $\text{XeF}_4$  and  $\text{XeF}_6$  are expected to be [2007]  
 (a) oxidizing (b) reducing  
 (c) unreactive (d) strongly basic



10 Subjective Problems

10. Write balanced equations for the reactions of the following compounds with water : [2002 - 5 Marks]  
 (i)  $\text{Al}_4\text{C}_3$  (ii)  $\text{CaNCN}$   
 (iii)  $\text{BF}_3$  (iv)  $\text{NCl}_3$   
 (v)  $\text{XeF}_4$
11. Draw the molecular structures of  $\text{XeF}_2$ ,  $\text{XeF}_4$  and  $\text{XeO}_2\text{F}_2$  indicating the location of lone pair(s) of electrons. [2000 - 3 Marks]



Answer Key

Topic-1 : Group-15 Elements (Nitrogen Family)

1. (b) 2. (a) 3. (c) 4. (a) 5. (b) 6. (c) 7. (b) 8. (d) 9. (b) 10. (d)  
 11. (b) 12. (b) 13. (c) 14. (a) 15. (c) 16. (c) 17. (b) 18. (d) 19. (c) 20. (a)  
 21. (d) 22. (b) 23. (a) 24. (d) 25. (c) 26. (8) 27. (4) 28. (2,38) 29. (four) 30. (two)  
 31. (white/yellow) 32. (-3) 33. True 34. True 35. False 36. (a,b,d) 37. (a,b,c) 38. (b,c) 39. (b,d)  
 40. (a,b,c) 41. (a,b) 42. (c) 43. (a,c,d) 44. (a,b,d) 45. (a,d) 46. (d) 47. (i) (d) (ii) (b) (iii) (a) (iv) (c)  
 48. (c) 49. (c) 50. (b) 51. (b)

Topic-2 : Group-16 Elements (Oxygen Family)

1. (a) 2. (b) 3. (c) 4. (c) 5. (d) 6. (b) 7. (c) 8. (b) 9. (b) 10. (c)  
 11. (d) 12. (6) 13. (288) 14. Nitric oxide, [NO] 15. (b,d) 16. (a,c,d) 17. (a,c,d) 18. (c) 19. (b)  
 20. (a) 21. (a)

Topic-3 : Group-17 Elements (Halogen Family)

1. (c) 2. (a) 3. (c) 4. (b) 5. (c) 6. (c) 7. (b) 8. (b) 9. (d) 10. (a)  
 11. (a) 12. (6) 13. (5) 14.  $\text{I}_3^-$  complex ion 15. Hypobromous; bromite 16.  $\text{KI}_3$  17. HF  
 18.  $\text{NaIO}_3$  19. False 20. False 21. True 22. (a,b,d) 23. (c,d) 24. (a,b,d) 25. (a,d) 26. (a) 27. (c)  
 28. (c)

Topic-4 : Group-18 Elements (Noble Gases)

1. (c) 2. (d) 3. (b) 4. (19) 5. (d)  
 6. (A)-(p), (s); (B)-(p), (q), (r), (t); (C)-(p), (q); (D)-(p), (q) 7. (a) 8. (c) 9. (a)




# Hints & Solutions



## Topic-1: Group-15 Elements (Nitrogen Family)

1. (b) If  $\text{PCl}_5$  is fluorinated in a polar solvent, ionic isomers are formed e.g.:-  
 $[\text{PCl}_4]^+[\text{PCl}_4\text{F}_2]^-$  (colourless crystals)  
 and  $[\text{PCl}_4]^+[\text{PF}_6]^-$  (white crystals)
2. (a) Disproportionation of  $\text{HNO}_2$  in aqueous solution takes place as:



Hence, the products formed are  $\text{H}_3\text{O}^+$ ,  $\text{NO}_3^-$  and  $\text{NO}$ .

3. (c) Let oxidation states of phosphorus in  $\text{H}_3\text{PO}_2$ ,  $\text{H}_3\text{PO}_4$ ,  $\text{H}_3\text{PO}_3$  and  $\text{H}_4\text{P}_2\text{O}_6$  be  $w$ ,  $x$ ,  $y$  and  $z$  respectively.

Thus, in  $\text{H}_3\text{PO}_2$ :

$$3 \times (+1) + w + 2 \times (-2) = 0 \quad \therefore w = +1$$

In  $\text{H}_3\text{PO}_4$ :

$$3 \times (+1) + x + 4 \times (-2) = 0 \quad \therefore x = +5$$


In  $\text{H}_3\text{PO}_3$ :

$$3 \times (+1) + y + 3 \times (-2) = 0 \quad \therefore y = +3$$


In  $\text{H}_4\text{P}_2\text{O}_6$ :

$$4 \times (+1) + 2z + 6 \times (-2) = 0 \quad \therefore z = +4$$

Thus, the order of oxidation state is :



4. (a)  $\text{P}_4 + 8\text{SOCl}_2 \rightarrow 4\text{PCl}_3 + 4\text{SO}_2 + 2\text{S}_2\text{Cl}_2$   
 white phosphorus
5. (b) The slow decomposition of  $\text{HNO}_3$  is represented by the eqn.



6. (c)
- oxidation

$$0 \text{P}_4 + 3\text{NaOH} + 3\text{H}_2\text{O} \xrightarrow[\text{inert atmosphere}]{\Delta} \overset{-3}{\text{PH}_3} + 3\overset{+1}{\text{NaH}_2\text{PO}_2}$$

reduction


(sodium hypophosphite)

7. (b)  $\text{HNO}_3$ ,  $\text{NO}$ ,  $\text{N}_2$ ,  $\text{NH}_4\text{Cl}$

8. (d)  $\text{Ba}(\text{N}_3)_2 \rightarrow \text{Ba} + 3\text{N}_2$

9. (b)  $\text{P}_4 + 3\text{O}_2 \xrightarrow{\text{in presence of N}_2} \text{P}_4\text{O}_6$

Here  $\text{N}_2$  acts as a diluent and thus retards further oxidation. Reaction of  $\text{P}_4$  under other three conditions.




- (c) In moist air,  $\text{P}_4\text{O}_6$  is hydrolysed to form  $\text{H}_3\text{PO}_3$   
 $\text{P}_4\text{O}_6 + 6\text{H}_2\text{O} \rightarrow 4\text{H}_3\text{PO}_3$

- (d) In presence of  $\text{NaOH}$ ,  
 $\text{P}_4 + 3\text{OH}^- + 3\text{H}_2\text{O} \rightarrow \text{PH}_3 + 3\text{H}_2\text{PO}_2^-$

10. (d) In  $\text{P}_4$ , the P-P linkage is formed by  $sp^3-sp^3$  hybridised orbital overlapping. So, the percentage of  $\pi$ -character will be 75%.


11. (b)  $\text{PbO}_2$  is a powerful oxidizing agent and liberate  $\text{O}_2$  when treated with acids.






12. (b)  $\text{NO}(\text{g}) + \text{NO}_2(\text{g}) \xrightarrow{-30^\circ\text{C}} \text{N}_2\text{O}_3$   
 (blue liquid)

13. (c) The ignition temperature of black phosphorus is highest among all its allotropes, hence is most stable.


14. (a) The structure of  $\text{H}_3\text{PO}_3$  is as follows: There are only two -OH groups and hence **dibasic**. The oxidation number of P in this acid is +3. Whereas P can have +5 oxidation state also. Therefore,  $\text{H}_3\text{PO}_3$  can be oxidised which means  $\text{H}_3\text{PO}_3$  is a **reducing agent**.



15. (c)  $\text{NH}_3$  does not react with basic oxide,  $\text{CaO}$ , while other reacts with  $\text{NH}_3$




16. (c) In cyclic metaphosphoric acid number of P-O-P bonds is three.



17. (b) In  $\text{BCl}_3$ ,  $H = \frac{1}{2}(3+3+0-0) = 3$ ;  $sp^2$  hybridization



(bond angle =  $120^\circ$ ). Similarly  $\text{PCl}_3$ ,  $\text{AsCl}_3$  and  $\text{BiCl}_3$  are found to have  $sp^3$  hybridized central atom with one lone pair of electrons on the central atom. The bond angle  $\leq 109^\circ 28'$ , since the central atoms belong to the same group. The bond

- angle of the chlorides decreases as we go down the group. Thus, the order of bond angle is,  $\text{BCl}_3 > \text{PCl}_3 > \text{AsCl}_3 > \text{BiCl}_3$ .
18. (d)  $(\text{NH}_4)_2\text{Cr}_2\text{O}_7 \rightarrow \text{N}_2 + \text{Cr}_2\text{O}_3 + 4\text{H}_2\text{O}$
19. (c)  $\text{Ca}_3\text{P}_2 + 6\text{H}_2\text{O} \rightarrow 3\text{Ca}(\text{OH})_2 + 2\text{PH}_3$ ; i.e. 2 moles of phosphine are produced.
20. (a) Least basic trihalogen of nitrogen is  $\text{NF}_3$  because of the highest electronegativity of fluorine.
21. (d)  $\text{NO}_2$  is reddish brown coloured gas. Rest of the oxides are colourless.





The structure clearly shows the presence of covalent and co-ordinate bonds.

23. (a) Only nitrates of heavy metals and lithium decompose on heating to produce  $\text{NO}_2$ .
24. (d) None; it reacts with all given compounds. It forms addition compounds with them because it is basic in nature and given compounds are acidic in nature. It can be dried over any metal oxide.







Number of lone pairs = 8



When equimolar amounts of  $\text{PCl}_5$  and  $\text{H}_2\text{O}$  are used, the reaction is gentle:  $\text{PCl}_5 + \text{H}_2\text{O} \rightarrow \text{POCl}_3 + 2\text{HCl}$

[Note:  $\text{PCl}_5$  reacts violently with  $\text{H}_2\text{O}$  on complete hydrolysis to produce phosphoric acid.



$\text{PH}_3$ : a non-inflammable gas in its pure form; slightly soluble in water.

When  $\text{PH}_3$  is absorbed in  $\text{CuSO}_4$  solution cupric phosphide forms:  $2\text{PH}_3 + 3\text{CuSO}_4 \rightarrow \text{CuP}_2 + 3\text{H}_2\text{SO}_4$

1 mol of  $\text{P}_4 = 31 \times 4 = 124 \text{ g}$

$\therefore 1.24 \text{ g}$  of white phosphorous = 0.01 mol

$\therefore 0.01 \text{ mol}$  of  $\text{P}_4$  forms 0.01 mol of  $\text{PH}_3$

No. of moles of  $\text{CuSO}_4$  is required for complete

consumption of 0.01 mol =  $0.01 \times \frac{3}{2} = 15 \times 10^{-3}$



M.W. of  $\text{CuSO}_4 = 159 \text{ g/mol}$

$\therefore$  Amount of  $\text{CuSO}_4$  required =  $15 \times 10^{-3} \times 159 = 2.38 \text{ g}$

29. four.

Each 'P' atom is linked to 4 'O' atoms as confirmed by its structure. It is linked to three 'O' atoms by single bond and one 'O' atom by double bond.

Each P atom is linked to 4 oxygen atoms



31. white/ yellow; : In white phosphorus, each phosphorus atom is linked to the other three atoms by covalent bonds. PPP bond angle is  $60^\circ$ , due to which the molecule remains under strain and hence is active in nature.

32. -3; Nitrogen has 5 electrons in its valence shell. Thus, it can accept maximum 3 electrons.

33. True : The molecule of NO has eleven valence electrons (5 due to N and 6 due to O). It is impossible for all of them to be paired, hence the nitric oxide molecule contains an odd electron which makes gaseous nitric oxide paramagnetic.



In the liquid and solid states, nitric oxide is polymerised to a dimer which is diamagnetic.


34. True :

The central element in the metal hydrides of group 15 elements is although in  $sp^3$  hybrid state, the H - M - H bond angle is less than the normal tetrahedral bond angle of  $109^\circ 28'$ ; e.g. the bond angle, H - N - H in  $\text{NH}_3$  is  $106^\circ 45'$ . This is due to greater repulsion between a lone pair and a bond pair of electrons than between the two bond pairs of electrons.


The decrease in bond angle from  $107.8^\circ$  in ammonia to about  $90^\circ$  in  $\text{AsH}_3$  can be explained by the fact that in the latter case  $sp^3$  hybridisation becomes less and less distinct with the increasing size of their electron clouds, i.e., pure p orbitals (instead of  $sp^3$  hybrid orbitals) are used for M - H bonding and the lone pair of electrons is present in spherical s-orbital.

35. False : Red phosphorus is polymeric substance. It exists as chains of  $\text{P}_4$  tetrahedra linked together. Therefore, it is less volatile than white phosphorus.


36. (a, b, d)



(b)  $\text{H}_3\text{PO}_4$  has "P" in its highest oxidation state, hence cannot act as a reducing agent (correct)







- (d) The hydrogen which is directly attached to phosphorous does not ionized in water.



37. (a, b, c)



- (a) Basic character of oxides increases on moving down the group therefore,  $\text{Bi}_2\text{O}_3$  is more basic than  $\text{N}_2\text{O}_5$ .  
 (b) Covalent nature depends on electronegativity difference between bonded atoms. In  $\text{NF}_3$ , N and F are non-metals but in  $\text{BiF}_3$ , Bi is metal while F is non metal therefore,  $\text{NF}_3$  is more covalent than  $\text{BiF}_3$ .  
 (c) In  $\text{PH}_3$ , hydrogen bonding is absent but in  $\text{NH}_3$ , hydrogen bonding is present, therefore  $\text{PH}_3$  boils at lower temperature than  $\text{NH}_3$ .  
 (d) Due to small size in N-N single bond, l.p. - l.p. repulsion is more than P-P single bond therefore, N-N single bond is weaker than the P-P single bond.

38. (b, c)




Hence, only  $(\text{NH}_4)_2\text{Cr}_2\text{O}_7$  and  $\text{Ba}(\text{N}_3)_2$  can provide  $\text{N}_2$  gas on heating below  $300^\circ\text{C}$


39. (b, d)






(b)  $\text{N}_2\text{O}_5$  is diamagnetic in nature




$\text{N}_2\text{O}_5$  contains one N-O-N bond but not N-N bond.



40. (a, b, c)  $\text{N} \equiv \text{N} \rightarrow \text{O}$





41. (a, b) When ammonium salt  $\text{NH}_4\text{NO}_3$  or  $\text{NH}_4\text{NO}_2$  (ammonium salts are colourless) is boiled with excess of  $\text{NaOH}$ , ammonia ( $\text{NH}_3$ ) gas is evolved as follows:



The  $\text{NH}_3$  gas evolved is non-flammable gas.

When the gas evolution ceases we are left with  $\text{NaNO}_2$  or  $\text{NaNO}_3$  in solution.

These salts get reduced when Zn is added to this solution containing salt ( $\text{NaNO}_2$  or  $\text{NaNO}_3$ ). Again  $\text{NH}_3$  gas evolves.




Thus, the colourless salt [H] is either  $\text{NH}_4\text{NO}_2$  or  $\text{NH}_4\text{NO}_3$ . Thus, (a) and (b) are correct answers.


42. (c)  $2\text{NH}_3 + \text{OCI}^- \rightarrow \text{NH}_2\text{NH}_2 + \text{H}_2\text{O} + \text{Cl}^-$

43. (a, c, d) The four atoms in a  $\text{P}_4$  molecule are situated at the corners of a tetrahedron. There are six P - P single bonds with PPP bond angle equal to  $60^\circ$ . Each phosphorus has a lone pair of electrons.

44. (a, b, d) Sodium nitrate on decomposition upto  $500^\circ\text{C}$  gives  $\text{NaNO}_2$  and oxygen.






While at higher temperature (*i.e.* above  $800^\circ\text{C}$ ),  $\text{NaNO}_2$  further decomposes into  $\text{Na}_2\text{O}$ ,  $\text{N}_2$  and  $\text{O}_2$ .



45. (a, d)  $\text{NH}_4\text{NO}_3 \longrightarrow \text{N}_2\text{O} + 2\text{H}_2\text{O}$   
 $\text{NH}_2\text{OH} + \text{HNO}_2 \longrightarrow \text{N}_2\text{O} + 2\text{H}_2\text{O}$

46. (d) (P)  $\text{P}_2\text{O}_3 + 3\text{H}_2\text{O} \rightarrow 2\text{H}_3\text{PO}_3$



47. (i) (d) (ii) (b) (iii) (a) (iv) (c)

48. (c) We know that phosphates have a biological significance in human, therefore statement (a) is not correct. Since nitrates are more soluble in water so they are less abundant in earth's crust where as phosphates are less soluble in water and so they are more abundant in earth's crust. Thus, statement (b) is False and statement (c) is correct.

In nitrates ( $\text{NO}_3^-$ ) nitrogen is in +5 oxidation state which is the highest oxidation state exhibited by nitrogen. Because of this, nitrates can not be oxidized (oxidation means increase in oxidation state). Hence, statement (d) is not correct.


The correct answer is (c).

49. (c) In case of group 15 (nitrogen group), on moving down the group, there occurs a decrease in bond angle of metal hydrides. This decrease in bond angle of metal hydrides of this group may be attributed to the increased  $p$ -character in the bond pair which results in more  $s$ -character in lone pair orbital.

The directional character is more for  $sp^3$  hybrid orbital than on  $s$ -orbital.

Thus, the correct answer is (c).


50. (b) The reaction between NaOH and white phosphorus ( $P_4$ ) can be represented as follows:



In this reaction, Phosphorus is oxidised as well as reduced, so it is a disproportionation reaction.

∴ The correct answer is (b).

51. (b) Nitrogen cannot form pentahalides because it cannot expand its octet due to non-availability of  $d$ -orbitals. So E is not correct explanation of S.
52.  $(CH_3)_3N$  and  $(Me_3Si)_3N$  are not isostructural, the former is pyramidal while the latter is trigonal planar. Silicon has vacant  $d$  orbitals which can accommodate lone pair of electrons from N (back bonding) leading to planar shape.



53.  $6CaO + P_4O_{10} \longrightarrow 2Ca_3(PO_4)_2$






$$\text{Moles of } P_4O_{10} = \frac{852}{284} = 3$$

Moles of CaO =  $3 \times 6 = 18$ ; wt. of CaO =  $18 \times 56 = 1008$  g  
For structure of  $P_4O_{10}$ : [See previous question].

54. Elemental nitrogen exists as a diatomic molecule because nitrogen can form  $p\pi - p\pi$  multiple bonds which is not possible in case of phosphorus due to repulsion between non-bonded electrons of the inner core. There is no such repulsion in case of smaller nitrogen atoms as they have only  $1s^2$  electrons in their inner core.



55. In such a case:

$A = Ca(OH)_2$ ,  $B = NH_4HCO_3$ ,  $C = Na_2CO_3$ ,  $D = NH_4Cl$  and  $E = CaCl_2$




56.  $7H_3PO_4 + Ca_5(PO_4)_3F \rightarrow 5Ca(H_2PO_4)_2 + HF$   
Triple superphosphate

57. The poisonous element  $M$  may be As. So, on the basis of the given facts,



Hence,  $M = As$ ;  $N = AsH_3$


- 58.



No. of P - O single bonds = 12


No. of P - O double bonds = 4

59. (i)  $2P + 3I_2 + 6H_2O \rightarrow 2H_3PO_4 + 6HI$




60. Since carbon has no  $d$ -orbital, it cannot extend its coordination number beyond four, its halides are not attacked (hydrolysed) by water. On the other hand, silicon have vacant  $d$ -orbitals to which water molecules can coordinate and hence their halides are hydrolysed by water. As the charge on central atom increases, the tendency of attack of a nucleophile ( $OH^-$ ) increases.


Increasing order of extent of hydrolysis







61. (i)  $KClO_3 + 2H_2C_2O_4 + H_2SO_4 \rightarrow KHSO_4 + HCl + 6CO_2 + 3H_2O$












62.  $N_2O$  has two principal resonance structures:




63. (i)  $P_4O_{10} + 6PCl_5 \rightarrow 10POCl_3$





(i) Phosphine gas ( $PH_3$ ) is evolved when white phosphorous is boiled with aqueous NaOH or alcoholic solution of potassium hydroxide.



64. (i) Nitrogen and fluorine both are small and have high electron density, they repel the bonded pair of electrons leading to larger bond length than expected.

(ii) It is due to self ionization of  $NH_3$ , the reaction is



Thus, on addition of  $NH_4Cl$  the concentration of  $NH_4^+$  radical increases and therefore,  $NH_4Cl$  acts as an acid in liquid  $NH_3$ .



(iii) As compared to P, N atom has higher electronegativity and small size and shows H-bonding.

Thus, ammonia molecule show association where as phosphine does not.

(iv)  $H_3PO_3$  is a dibasic acid because it contains two OH groups in its molecule.


In the two P-OH bonds, the hydrogen is ionisable.

(v) Orthophosphorus acid is a dibasic acid as it has 2-OH groups in its formula : whereas orthophosphoric acid has 3-OH groups.




(vi) Liquor ammonia possesses high vapour pressure at room temperature and thus before opening a bottle of liquor ammonia, it should be cooled to lower the pressure of  $NH_3$  inside the bottle, otherwise  $NH_3$  will bump out of the bottle.





(vii) In  $H_3PO_4$  and  $H_3PO_3$ , the P atom is attached to 3 and 2 OH groups respectively. The H atom of these P-OH bonds are ionisable. This clearly shows that  $H_3PO_4$  is tribasic and  $H_3PO_3$  is dibasic.



**Topic-2: Group-16 Elements (Oxygen Family)**









- (b) The following reaction occurs  
 $Na_2S_2O_3 + 4Cl_2 + 5H_2O \rightarrow 2NaHSO_4 + 8HCl$
- (c) In  $KMnO_4$ , manganese is already present in its highest possible oxidation state i.e. +7. So, no further oxidation is possible.
- (c) Among oxyacids of sulphur, only Caro's acid ( $H_2SO_5$ ) and Marshall's acid ( $H_2S_2O_8$ ) have the O-O linkage.
- (d) In sulphur trioxide trimer  $S_3O_9$  (also called  $\gamma$ -sulphur trioxide), two sulphur atoms are linked to each other via O atoms, hence there is no S-S bond.









(2 moles of sulphuric acid and 1 mole of  $H_2O_2$ )

But this is not one of the options.






[B]  
Brown fumes  
and pungent smell




Oxidation states of 'S' are: +4 in (A), (+4) in B and +2 in (C), +2.5 in (D)

24. Sulphur trioxide produced in the contact process is passed into 98% sulphuric acid forming  $\text{H}_2\text{S}_2\text{O}_7$ . It is not dissolved in water as the reaction is violent and it gives a dense chemical mist of sulphuric acid particles.





The catalyst used in the contact process is vanadium pentoxide.







The structures of A, B, C and D are as follows.






Thionyl chloride





28. The two resonating structures of ozone are:




Among oxides of the non-metals, the acidic strength increases with oxidation state. Hence  $\text{SO}_3$  (O.S. of S = +6) is most acidic followed by  $\text{N}_2\text{O}_5$  (O.S. of N = +5) and  $\text{CO}_2$  and  $\text{SiO}_2$  (O.S. of C and Si = +4). Further  $\text{CO}_2$  is more acidic than  $\text{SiO}_2$  because of small size of C-atom.



32. By boiling  $\text{Na}_2\text{SO}_3$  solution with powder of sulphur in absence of air, sodium thiosulphate is prepared. Unreacted S is removed, filtrate is evaporated to give crystals of sod. thiosulphate.



34. (i) The reducing nature of  $\text{SO}_2$  is represented as



Hence, with the increase of  $\text{OH}^-$  (alkalinity) the forward reaction is favoured.

(ii) Oxygen is the 2nd most electronegative element after the fluorine and thus invariably show negative oxidation state. Furthermore, it has  $2s^2 2p^4$  configuration and thus requires only two electrons to complete its octet to show -2 oxidation state. Although sulphur also possess  $3s^2 3p^4$  configuration but due to availability of d-orbitals in their outermost shell -2, +2, +4, +6 oxidation state are also shown. Oxygen, however, cannot exceed -2 oxidation state due to non-availability of d-orbitals in its outermost shell.

(iii) Sulphur consists of  $\text{S}_8$  rings held together by weak van der Waal's forces. As sulphur melts at  $119^\circ\text{C}$ , these van der Waal's forces are overcome and  $\text{S}_8$  rings slip and roll over one another giving rise to a clear mobile liquid. Above  $160^\circ\text{C}$ , the  $\text{S}_8$  rings begin to open up and form long chains which gets tangled with each other, thereby gradually increasing the viscosity.



### Topic-3: Group-17 Elements (Halogen Family)

- (c)  $2\text{HClO}_3 + 2\text{HCl} \longrightarrow 2\text{ClO}_2 + \text{Cl}_2 + 2\text{H}_2\text{O}$   
(paramagnetic)

$2\text{ClO}_2 + 2\text{O}_3 \longrightarrow \text{Cl}_2\text{O}_6 + 2\text{O}_2$
- (a) X : Ag, Y : Pb

$\text{AgNO}_3 \xrightarrow{\text{NaCl}} \text{AgCl(s)}; \text{PbNO}_3 \xrightarrow{\text{NaCl}} \text{PbCl}_2(\text{s})$

$\text{AgCl(s)} + \text{NH}_3(\text{aq}) \longrightarrow [\text{Ag}(\text{NH}_3)_2]^+$   
(P) soluble

$\text{PbCl}_2(\text{s}) + \text{KI} \longrightarrow \text{PbI}_2$   
(Q) yellow ppt.

$\text{AgCl(s)} + \text{Na}_2\text{S}_2\text{O}_3 \longrightarrow \text{Na}_3[\text{Ag}(\text{S}_2\text{O}_3)_2] + \text{NaCl}$
- (c)  $\text{Cl}_2 + \text{NaOH} \longrightarrow \text{NaCl} + \text{NaClO} + \text{H}_2\text{O}$   
[cold and dilute]
- (b) The ions consisting of two or more atoms of which at least one is 'N' atom is known as pseudohalide ions. These are univalent, have properties similar to those of halide ions. They combine to form dimers comparable with halogen molecules,  $\text{X}_2$ . These dimers are known as pseudohalogens.
- (c)  $\text{KF} + \text{HF} \rightarrow \text{KHF}_2 \rightleftharpoons \text{K}^+ + (\text{HF}_2)^-$
- (c)  $\text{CsBr}_3$  may be represented as  $\text{Cs}^+\text{Br}_3^-$
- (b) Chlorine is stronger oxidising agent than bromine therefore, chlorine water will liberate bromine from KBr solution.  
 $2\text{KBr} + \text{Cl}_2 \rightarrow 2\text{KCl} + \text{Br}_2$
- (b) Bleaching action of chlorine is only in presence of moisture where nascent oxygen is displaced from  $\text{H}_2\text{O}$ .  
 $\text{Cl}_2 + \text{H}_2\text{O} \rightarrow \text{HCl} + \text{HClO}(\text{unstable})$   
 $\text{HClO} \rightarrow \text{HCl} + \text{O}$
- (d) HI and HBr (in that order) are the strongest reducing hydric acids and hence they reduce  $\text{H}_2\text{SO}_4$ . HCl is quite stable and hence is oxidised by strong oxidising agent like  $\text{KMnO}_4$ . HF is not a reducing agent. In the smallest  $\text{F}^-$  ion, the electron which is to be removed during oxidation is closest to the nucleus and therefore most difficult to be removed. Therefore, HF is a poor reducing agent.
- (a)  $\therefore \text{Cl}_2$  is more reactive than bromine.
- (a) Due to highest bond dissociation energy.
- (6)  $2\text{ClO}_2 + 2\text{O}_3 \rightarrow \text{Cl}_2\text{O}_6 + 2\text{O}_2$   
 $\text{Cl}_2\text{O}_6 \Rightarrow 2x + 6(-2) = 0$   
 $x = +6$   
Average oxidation state of Cl in  $\text{Cl}_2\text{O}_6$  is (+6).
- (5)  $3\text{Br}_2 + 3\text{Na}_2\text{CO}_3 \longrightarrow 5\text{NaBr} + \text{NaBrO}_3 + 3\text{CO}_2$
- $\text{I}_3^-$  complex ion;  $\text{I}_2 + \text{I}^- \longrightarrow \text{I}_3^-$
- Hypobromous; bromite.  $\text{HBrO} \rightleftharpoons \text{H}^+ + \text{BrO}^-$
- $\text{KI}_3$ ; complexes are more soluble in water as compared to normal salts.  $[\text{KI} + \text{I}_2 \longrightarrow \text{KI}_3]$
- HF**; HF is the weakest of the three, because the ionisation (*i.e.* acidic character) of  $\text{HX}$  is a multistep process and when its  $\Delta H$ , heat of ionisation, is calculated it comes out to be the minimum. This is due to the strong H-F bond, large heat of hydration (because of H-bonding) and low value of electron affinity of F-atom.
- NaIO<sub>3</sub>**  
 $3\text{I}_2 + 6\text{NaOH} \xrightarrow{\text{(hot)}} 5\text{NaI} + \text{NaIO}_3 + 3\text{H}_2\text{O}$
- False** : None amongst HBr and HI, exhibit hydrogen bonding. HI is a stronger acid than HBr. Although the electronegativity and electron affinity of Br is greater than that of I, the bond dissociation energy of H-I (+295 kJ/mol) is lower than that of H-Br (+363 kJ/mol). Also the enthalpy of hydration of  $\text{I}^\ominus$  is -394 kJ/mol, whereas for  $\text{Br}^\ominus$  is -339 kJ/mol, which indicates the better stability of  $\text{I}^\ominus$  in aqueous medium than  $\text{Br}^\ominus$ . Thus, HI has a stronger tendency to release protons to water molecules and hence, is a stronger acid.
- False** : The reason of accepting electrons by fluorine more readily than chlorine is as follows:  
(i)  $\text{F}_2$  has lower enthalpy of dissociation (+159 kJ mol<sup>-1</sup>) due to weak F-F bond than  $\text{Cl}_2$  (+243 kJ mol<sup>-1</sup>).  
(ii)  $\text{F}_2$  has higher enthalpy of hydration due to the smaller size of  $\text{F}^-$  ion (-513 kJ/mol) than that of  $\text{Cl}^-$  ion (-370 kJ mol<sup>-1</sup>). Thus,  $\text{F}_2$  is stronger reductant than  $\text{Cl}_2$  both in solution and when dry.  
Similarly,  $\text{Cl}_2$  will displace  $\text{Br}^-$  and  $\text{I}^-$  ions from their solutions and  $\text{Br}_2$  will displace  $\text{I}^-$  ions.  
In general, a halogen of low atomic number will oxidise the halide ion of higher atomic number.
- True** :  $\text{Fe} + 2\text{HCl} \xrightarrow{+2} \text{FeCl}_2 + \text{H}_2; E_{\text{Fe}^{2+}/\text{Fe}}^\ominus = -044\text{V}$
- (a, b, d)

(a) Acidic order :  $\text{HClO} < \text{HClO}_3 < \text{HClO}_4$   
Conjugate base order :  $\text{ClO}^- > \text{ClO}_3^- > \text{ClO}_4^-$




(b) Hypochlorite ion ( $\text{ClO}^-$ ) :  $:\ddot{\text{Cl}}-\ddot{\text{O}}:$  Linear

Chlorate ion : ( $\text{ClO}_3^-$ ) Trigonal pyramidal

Perchlorate ion : ( $\text{ClO}_4^-$ ) Tetrahedral

In chlorate ion bond angle changes due to presence of lone pair on chlorine atom. While there is no effect of lone pair on hypochlorite ion and perchlorate ion.

(c) Disproportionation reaction of




23. (c, d) Energy,  $E = \frac{hc}{\lambda}$


On moving down the group, the colour of the  $X_2$  molecule of group 17 elements changes gradually from yellow to violet. This happens because the amount of energy required for the excitation of the halogen atom decreases down the group. HOMO ( $\pi^*$ )-LUMO ( $\sigma^*$ ) gap decreases down the group that makes  $\pi^*$  to  $\sigma^*$  excitation easier. Lesser the energy gap, more is the wavelength of light absorbed and hence, lesser is the wavelength of light emitted.

24. (a, b, d)

(a) In both the acids central atom is  $sp^3$  hybridized.




(b)  $\text{HClO}_4$  is more acidic than  $\text{HClO}$  because  $\text{ClO}_4^-$  is more stable than  $\text{ClO}^-$  due to resonance.




(d)  $\text{HClO}_4$  is a stronger acid than  $\text{H}_3\text{O}^+$ , hence conjugate base of  $\text{HClO}_4$ , i.e.  $\text{ClO}_4^-$  is weaker base than  $\text{H}_2\text{O}$ .

25. (a, d)  $\text{NH}_3$  and  $\text{CF}_2\text{Cl}_2$  (freon-12) are used as refrigerants.

26. (a)  $\text{Ca}(\text{OCl})\text{Cl} \rightarrow \text{Ca}^{2+} + \text{OCl}^- + \text{Cl}^-$   
 $\text{OCl}^-$  (Hypochlorite ion) is anion of the acid  $\text{HOCl}$  which on dehydration gives  $\text{Cl}_2\text{O}$ .



27. (c) Bleach +  $2\text{KI} \rightarrow \text{I}_2$  + Products



Number of millimole of hypo =  $0.25 \times 48$   
 $= 2 \times \text{millimole of } \text{I}_2$

$$\therefore \text{Number of millimole of } \text{I}_2 = \frac{0.25 \times 48}{2} = 6$$

millimole of  $\text{I}_2$  = millimole of bleach


Molarity of bleaching solution

$$\frac{\text{Millimoles of bleach}}{\text{Vol. (in mL) of bleach}} = \frac{6}{25} = 0.24$$

28. (c) F has slightly less electron affinity than chlorine because F has very small atomic size (only two shells).

Hence, there is a tendency of electron-electron repulsion, which results in less evolution of energy in the formation of  $\text{F}^-$  ion. Assertion is correct but reason is incorrect.

29. More electronegative halogen displaces lesser electronegative halogen from its halide. Thus,




30. (a)  $2\text{KI} + \text{Cl}_2 \rightarrow 2\text{KCl} + \text{I}_2$

Since,  $\text{Cl}_2$  is more powerful oxidising agent than  $\text{I}_2$ ,  $\text{Cl}_2$  is able to displace  $\text{I}^-$  to form  $\text{I}_2$ .





On subtracting eq. (i) from eq. (ii), we get




(b)  $2\text{KClO}_3 + \text{I}_2 \rightarrow 2\text{KIO}_3 + \text{Cl}_2$





Here,  $\text{ClO}_3^-$  is more powerful oxidising agent than  $\text{IO}_3^-$ , so Cl is displaced by I.





On subtracting eq. (i) from eq. (ii), we get



31. At first,  $\text{Bi}(\text{NO}_3)_3$  hydrolyses to give nitric acid which being an oxidising agent, oxidises potassium iodide liberating free iodine responsible for dark brown precipitate. Iodine dissolves in excess of potassium iodide forming soluble  $\text{KI}_3$  imparting yellow colour to solution.



32.  $2\text{NH}_3 + \text{NaOCl} \rightarrow \text{H}_2\text{N.NH}_2 + \text{NaCl} + \text{H}_2\text{O}$   
Hydrazine



34. (i)  $\text{HOCl} < \text{HOClO} < \text{HOClO}_2 < \text{HOClO}_3$





As the number of oxygen atoms increase, the -ve charge dispersal becomes more and more from Cl atom due to more electronegativity of oxygen atom. Due to more double bond character between Cl and O atoms, the bond length decreases and thus, bond strength increases. Hence, thermal stability increases.






(ii)  $\text{HI} < \text{HBr} < \text{HCl} < \text{HF}$

The strength of H-X bond decreases from HF to HI. The larger is H-X bond length, lower is the bond energy, lesser is the bond strength.




36. (i) The repulsive forces between fluorine atoms are high due to its small size and high electronegativity. It makes dissociation of F-F bond easy. So, bond dissociation energy of  $\text{F}_2$  is less than  $\text{Cl}_2$ .

(ii) The standard reduction potential of fluorine is highest and thus, it cannot be oxidized by any reagent.



(iii) Anhydrous HCl, being a covalent compound, is a bad conductor however, an aqueous solution of HCl is ionised to give  $\text{H}^+$  and  $\text{Cl}^-$  ions and is a good conductor.






(iii) HI cannot be prepared by heating hydrogen iodide with conc.  $\text{H}_2\text{SO}_4$  because it is a strong oxidising agent and oxidises HI to  $\text{I}_2$ .



Hence, HI is prepared by heating iodides with conc. phosphoric acid.



$\text{H}_3\text{PO}_4$  is not a strong oxidising agent.




38. (i) HBr is a reducing agent and it reduces  $\text{H}_2\text{SO}_4$  to  $\text{SO}_2$ .

(ii) Acids turn blue litmus red, so HClO also turns blue litmus red. The colour of litmus is decolourised because HClO is also a strong oxidising agent.




### Topic-4: Group-18 Elements (Noble Gases)

1. (c)





2. (d)  $\text{XeO}_2\text{F}_2$  has trigonal bipyramidal geometry, but due to presence of lone pair of electrons on equatorial position, its shape is *see-saw*.



3. (b) In  $\text{XeOF}_4$ , Xenon is  $sp^3d^2$  hybridised and has one lone pair.





4. (19)






Shape of  $\text{XeF}_6$  is distorted octahedral contains one lone pair  $e^-$ s on central atom 3 lone pair  $e^-$ s on each F atom surrounded by Xe.

Total no. of lone pairs:  $1 + 18 = 19$

5. (d) (P)




6. (A)-(p), (s)  $(\text{CH}_3)_2\text{SiCl}_2 + 2\text{H}_2\text{O}$



glass




Soluble hexafluorosilicic(IV) acid





7 (a) Argon, being a noble gas, will not react with the metals, thus, can be used in arc welding.





8. (c) In  $\text{XeO}_3$  there are total of 4 electron pairs around central atom. Out of which 3 are bonding electron pair and one is non-bonding electron pair. This combination provides  $sp^3$ -hybridization and pyramidal shape.



9. (a) All xenon fluorides are strongly oxidizing,  $\text{XeF}_4$  can act as reducing agent (with  $\text{F}_2$ ) as well as oxidizing agent but  $\text{XeF}_6$  can only function as an oxidizing agent.



(ii)  $\text{CaNCN} + 3\text{H}_2\text{O} \longrightarrow \text{CaCO}_3 \downarrow + 2\text{NH}_3$   
 Ammonia formed dissolves in water to form  $\text{NH}_4\text{OH}$



11. Use the formula

$H$  (hybridisation),  $H = \frac{1}{2}(V + M - C + A)$  where

$V$  = number of electron in valence shell of central atom  
 $M$  = number of monovalent atoms surrounding the central atom  
 $C$  = Charge on cation  
 $A$  = Charge on anion

$\text{XeF}_2$ :  $H = \frac{1}{2}(8 + 2 - 0 + 0) = 5$  Hence, hybridisation is  $sp^3d$ , and thus its shape is linear.

$\text{XeF}_4$ :  $H = \frac{1}{2}(8 + 4 - 0 + 0) = 6$ , Hence, hybridisation is  $sp^3d^2$ , and thus its shape is square planar.

$\text{XeO}_2\text{F}_2$ :  $H = \frac{1}{2}(8 + 2 - 0 + 0) = 5$ , Hence, hybridisation is  $sp^3d$ , and shape is see saw.

