

01

RATE OF A CHEMICAL REACTION

decrease in conc of reactant OR Increase in conc of reactant
Time taken Time taken

AVERAGE RATE

Consider a reaction: $A + B \rightarrow C + D$

$$-\frac{\Delta[A]}{\Delta t}, -\frac{\Delta[B]}{\Delta t}, \frac{\Delta[C]}{\Delta t}, \frac{\Delta[D]}{\Delta t}$$

INSTANTANEOUS RATE

Consider a reaction: $aA + bB \rightarrow cC + dD$

$$-\frac{1}{a} \frac{d[A]}{dt}, -\frac{1}{b} \frac{d[B]}{dt}, \frac{1}{c} \frac{d[C]}{dt}, \frac{1}{d} \frac{d[D]}{dt}$$

(Unit of Rate = $\text{mol litre}^{-1} \text{s}^{-1}$)

02

FACTORS INFLUENCING RATE OF REACTION

Factors	Effect on reaction rate
Increase in concentration	Increases
Increase in temperature	Increases
Presence of catalyst	Increases

03

RATE LAW

Consider a general reaction,
 $aA + bB \rightarrow \text{product}$
Rate = $k[A]^x[B]^y$ (law of mass action)
Rate = $k[A]^x[B]^y$ (rate law expression)
 x & y are determined experimentally and may or may not be equal to a & b
 x & y represents the order of reaction with respect to A & B

RATE CONSTANT

- Larger the value of k , faster is the reaction.
- The value of k changes only with temperature for given reaction.

(unit of rate constant = $(\text{mol})^{-x-y} \text{L}^{x+y} \text{s}^{-1}$)

04

ORDER AND MOLECULARITY

Consider a general reaction,
 $aA + bB \rightarrow \text{product}$

Rate = $k[A]^x[B]^y$

molecularity = $a + b$


order = $x + y$

Molecularity	Order
Theoretical concept.	An experimentally determined quantity.
It cannot be zero, fractional, infinite and imaginary.	It can be equal to zero, positive, negative and fractional.

05

PSEUDO ORDER REACTIONS

Consider the reaction



In these reactions, concentration of water (one of the reactants) is in excess and its concentration remains constant throughout the reaction.

Thus, rate $\propto [C_2H_5O_2]$

Therefore, order = 1

Q During the decomposition of H_2O_2 , 48 g O_2 is formed per minute at a certain point of time.

The rate of formation of water at this point is

- (a) $0.75 \text{ mol min}^{-1}$ (b) 1.5 mol min^{-1}
(c) $2.25 \text{ mol min}^{-1}$ (d) 3.0 mol min^{-1}

AFMC 2012

Q Which of the following will lead to an increase in the rate of the reaction?

- (a) Decrease in temperature
(b) Decreasing concentration of reactants
(c) Addition of catalyst
(d) Addition of inhibitor

AMU 2010

Q The rate constant of a zero-order reactions has the unit

- (a) s^{-1} (b) $\text{mol L}^{-1} \text{s}^{-1}$
(c) $\text{L}^2 \text{mol}^{-2} \text{s}^{-1}$ (d) $\text{L mol}^{-1} \text{s}^{-1}$

AIMS 2009

Q When the rate of the reaction is equal to the rate constant, the order of the reaction is

- (a) zero order
(b) first order
(c) second order
(d) third order

Q For a pseudo first-order reaction, what is the unit of the rate of the reaction?

- (a) s^{-1}
(b) $\text{mol L}^{-1} \text{s}^{-1}$
(c) $\text{mol}^{-1} \text{L s}^{-1}$
(d) $\text{mol}^{-2} \text{L}^2 \text{s}^{-1}$

AMU 2006

CHEMICAL KINETICS

06

ELEMENTARY & COMPLEX REACTIONS


• Reactions occurring only in one step are called elementary reactions while that involving a sequence of elementary reactions, are called complex reactions.

• In case of complex reactions, the slowest step is called rate determining step.


(Note) Consider the reaction



Step - 1



Step - 2



From slow step

$r = k [O] [O_2]$

Here

$[O] \propto [O_2]$; From fast step

$[O] = \frac{[O_2]}{K}$

$r = k [O_2]^2 [O_2]^{-1}$

07

INTEGRATED RATE EQUATIONS

Zero order

$$k = \frac{[A]_0 - [A]_t}{t}$$

First order

$$k = \frac{2.303 \log \frac{[A]_0}{[A]_t}}{t}$$

Second order

$$k = \frac{1}{t} \left[\frac{1}{[A]_t} - \frac{1}{[A]_0} \right]$$

08

HALF LIFE PERIOD

Zero order

$$t_{1/2} = \frac{[A]_0}{2k}$$

First order

$$t_{1/2} = \frac{0.693}{k}$$

Second order

$$t_{1/2} = \frac{1}{k[A]_0}$$

FIRST ORDER TRICKS

$$t_{75\%} = 2t_{1/2}$$

$$t_{90\%} = 2.303 \frac{k}{k}$$

$$t_{99.9\%} = 10t_{1/2}$$

09

GRAPHICAL REPRESENTATION

1. Concentration - time graph

2. Rate - concentration graph

3. Half life - concentration graph

10

ARRHENIUS EQUATION


$$k = A e^{-E_a/RT}$$

$$\log k = \log A - \frac{E_a}{2.303R} \left(\frac{1}{T} \right)$$




(Note)

• For every 10° rise in temperature, rate becomes double and hence, rate constant becomes double.

• A reaction with higher value of E_a will have smaller value of rate constant.



Q Suppose the reaction: $A + 2B \rightarrow AB_2$ occurs by the following mechanism:



- (a) $k[A]$ (b) $k[B]$ (c) $k[A][B]$ (d) $k[B]^2$

Q A first order reaction has a specific reaction rate of 10^{-2} sec^{-1} . How much time will it take for 20 g of the reactant to reduce to 5 g?

- (a) 138.6 sec (b) 346.5 sec
(c) 693.0 sec (d) 238.6 sec

NEET 2017

Q The half-life period of zero order reaction is directly proportional to the _____

- (a) Rate constant
(b) Initial concentration of reactants
(c) Final concentration of reactants
(d) Concentration of products

AMU 2012

Q The graph of $t_{1/2}$ versus initial concentration 'a' is for

- (a) First order
(b) Second order
(c) Zero order
(d) Can't predict

Q The slope of Arrhenius plot ($\ln k$ vs $\frac{1}{T}$) of first order reaction is $-5 \times 10^3 \text{ K}$. The value of E_a of the reaction is [Given: $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$]

- (a) -83 kJ mol^{-1} (b) 41.5 kJ mol^{-1}
(c) 83 kJ mol^{-1} (d) 166 kJ mol^{-1}

NEET 2021