

# SIMPLE HARMONIC MOTION

## Time Period

- (i) Simple pendulum:  $T = 2\pi\sqrt{\frac{l}{g}}$
- (ii) Physical pendulum:  $T = 2\pi\sqrt{\frac{I}{mgl}}$
- (iii) Torsional pendulum:  $T = 2\pi\sqrt{\frac{I}{C}}$

## Combination of Springs

- (i) Series :  $\frac{1}{k_{\text{eff}}} = \frac{1}{k_1} + \frac{1}{k_2}$
- (ii) Parallel :  $k_{\text{eff}} = k_1 + k_2$
- (iii) Spring cut into two parts in ratio m:n  
 $k_1 = \frac{(m+n)k}{m}$ ,  $k_2 = \frac{(m+n)k}{n}$

## Linear SHM

- (i) Displacement :  $x = A \sin(\omega t + \phi)$
- (ii) Velocity :  $\frac{dx}{dt} = A\omega \cos(\omega t + \phi)$   
 $= \omega\sqrt{A^2 - x^2}$
- (iii) Acceleration :  $\frac{d^2x}{dt^2} = -A\omega^2 \sin(\omega t + \phi)$   
 $= -\omega^2 x$
- (iv) Phase :  $\omega t + \phi$
- (v) Phase Constant :  $\phi$

## Equation of SHM

- (i) Linear :  $a = -\omega^2 x$
- (ii) Angular :  $\alpha = -\omega^2 \theta$

## Mass Spring system

- (i)  $T = 2\pi\sqrt{\frac{m}{k}}$
- (ii) Two bodies system  $T = 2\pi\sqrt{\frac{\mu}{K}}$   
 Where  $(\mu) = \frac{m_1 m_2}{m_1 + m_2}$

## Composition of 2 SHM

- $x_1 = A_1 \sin \omega t$
- $x_2 = A_2 \sin(\omega t + \phi)$
- $x = A \sin(\omega t + \delta)$  Where
- $A = \sqrt{A_1^2 + A_2^2 + 2A_1 A_2 \cos \phi}$
- and  $\tan \delta = \frac{A_2 \sin \phi}{A_1 + A_2 \cos \phi}$

## Angular SHM

- (i) Displacement :  $\theta = \theta_0 \sin(\omega t + \phi)$
- (ii) Angular Velocity :  $\frac{d\theta}{dt} = \theta_0 \omega \cos(\omega t + \phi)$
- (iii) Acceleration :  $\frac{d^2\theta}{dt^2} = -\theta_0 \omega^2 \sin(\omega t + \phi)$   
 $= -\omega^2 \theta$
- (iv) Phase :  $\omega t + \phi$
- (v) Phase Constant :  $\phi$

## Energy in SHM

- (i) K.E. =  $\frac{1}{2} m\omega^2(A^2 - x^2)$
- (ii) U =  $\frac{1}{2} m\omega^2 x^2$
- (iii) E = K+U =  $\frac{1}{2} m\omega^2 A^2$   
 = Constant.

# SHM