



# 3D COORDINATE GEOMETRY




## Distance between two points



$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

## Section Formula



$$C \left( \frac{kx_2 + x_1}{k+1}, \frac{ky_2 + y_1}{k+1}, \frac{kz_2 + z_1}{k+1} \right)$$

## Direction Cosines & Ratios

### Direction Cosines

$$\cos \alpha = \frac{a_1}{|\vec{a}|}, \quad \cos \beta = \frac{a_2}{|\vec{a}|}, \quad \cos \gamma = \frac{a_3}{|\vec{a}|}$$

Note :  $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$

$\cos \alpha \equiv l$  ;  $\cos \beta \equiv m$  ;  $\cos \gamma \equiv n$

So  $l^2 + m^2 + n^2 = 1$


### Direction Ratios

If a, b and c are direction ratios then  $a \propto l$  ;  $b \propto m$  ;  $c \propto n$


$$\frac{l}{a} = \frac{m}{b} = \frac{n}{c} = \lambda \text{ (say)}$$

$$\therefore (a^2 + b^2 + c^2) \lambda^2 = l^2 + m^2 + n^2 = 1$$

Therefore,  $l = \pm \frac{a}{\sqrt{a^2 + b^2 + c^2}}$  ;  $m = \pm \frac{b}{\sqrt{a^2 + b^2 + c^2}}$  ;  $n = \pm \frac{c}{\sqrt{a^2 + b^2 + c^2}}$



# THREE DIMENSIONAL LINES



Line passing through point (a, b, c) parallel to line having direction cosines  $\ell, m, n$  is

$$L_1 : \frac{x-a}{\ell} = \frac{y-b}{m} = \frac{z-c}{n}$$

Equation of a line passing through two points A & B

$$L_2 : \frac{x-x_1}{x_2-x_1} = \frac{y-y_1}{y_2-y_1} = \frac{z-z_1}{z_2-z_1}$$

## Angle between two lines

$$\cos \theta = \frac{a_1a_2 + b_1b_2 + c_1c_2}{\sqrt{a_1^2+b_1^2+c_1^2} \sqrt{a_2^2+b_2^2+c_2^2}}$$



$$\cos \theta = \ell_1\ell_2 + m_1m_2 + n_1n_2$$

If  $L_1$  and  $L_2$  are Perpendicular ( $\theta = 90^\circ$ )

$$\ell_1\ell_2 + m_1m_2 + n_1n_2 = 0 \quad \text{Or} \quad a_1a_2 + b_1b_2 + c_1c_2 = 0$$

If  $L_1$  and  $L_2$  are Parallel

$$\frac{\ell_1}{\ell_2} = \frac{m_1}{m_2} = \frac{n_1}{n_2} \quad \text{Or} \quad \frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$$



## Distance between two skew lines

$$d = \frac{|(\vec{b}_1 \times \vec{b}_2) \cdot (\vec{a}_2 - \vec{a}_1)|}{|\vec{b}_1 \times \vec{b}_2|}$$

Cartesian form


$$\text{Line } L_1 : \frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1}$$

$$\text{Line } L_2 : \frac{x-x_2}{a_2} = \frac{y-y_2}{b_2} = \frac{z-z_2}{c_2}$$

$$d = \frac{\begin{vmatrix} x_2-x_1 & y_2-y_1 & z_2-z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix}}{\sqrt{(b_1c_2 - b_2c_1)^2 + (c_1a_2 - c_2a_1)^2 + (a_1b_2 - a_2b_1)^2}}$$



# THREE DIMENSIONAL PLANES



## Equation of a plane in Normal form

$$\text{Equation : } \vec{r} \cdot \hat{n} = d$$

unit normal vector along  $\vec{ON}$       Perpendicular distance of plane from 'O'

### Cartesian form

$$\text{Equation : } \ell x + my + nz = d$$

Here  $\ell, m, n$  are the direction cosines of  $\hat{n}$

## Intercept form of the equation of a plane ABC

$$\text{Equation : } \frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$


$a, b$  and  $c$  are the direction ratios.

## Equation of a plane perpendicular to a given vector and passing through a given point

Equation : Cartesian Form


$$(\vec{r} - \vec{a}) \cdot \vec{n} = 0 \quad \text{if } \vec{n} = a\hat{i} + b\hat{j} + c\hat{k}$$

Plane :  $a(x - x_1) + b(y - y_1) + c(z - z_1) = 0$



## The Distance of a Point P From Plane $P_1: \vec{r} \cdot \hat{n} = d$

$$\text{Perpendicular distance} = \frac{|\vec{p} \cdot \hat{n} - d|}{|\hat{n}|}$$



## Equation of a plane passing through three non - collinear points

$$\text{Equation : } (\vec{r} - \vec{a}) \cdot [(\vec{b} - \vec{a}) \times (\vec{c} - \vec{a})] = 0$$

### Cartesian Form

If  $R = (x_1, y_1, z_1)$ ,  $S = (x_2, y_2, z_2)$  &  $T = (x_3, y_3, z_3)$

$$\text{Equation : } \begin{vmatrix} x - x_1 & y - y_1 & z - z_1 \\ x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ x_3 - x_1 & y_3 - y_1 & z_3 - z_1 \end{vmatrix} = 0$$



Plane P passing through the Intersection of two given planes  $P_1$  &  $P_2$

$P_1 : \vec{r} \cdot \hat{n}_1 = d_1$  Or  $P_2 : \vec{r} \cdot \hat{n}_2 = d_2$

Equation of Plane P


$\vec{r} \cdot (\vec{n}_1 + \lambda \vec{n}_2) = d_1 + \lambda d_2 ; \lambda : \text{Constant}$

Cartesian form

$P_1 : a_1x + b_1y + c_1z = d_1$  Or  $P_2 : a_2x + b_2y + c_2z = d_2$

Equation of plane P :

$(a_1x + b_1y + c_1z - d_1) + \lambda(a_2x + b_2y + c_2z - d_2) = 0$



If Angle between two planes  $P_1$  &  $P_2$  is  $\theta$

$\cos \theta = \frac{|\vec{n}_1 \cdot \vec{n}_2|}{|\vec{n}_1| \cdot |\vec{n}_2|}$

Cartesian form

If  $P_1 : a_1x + b_1y + c_1z + d_1 = 0, P_2 : a_2x + b_2y + c_2z + d_2 = 0$

$\cos \theta = \frac{|a_1a_2 + b_1b_2 + c_1c_2|}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}}$

• If  $P_1 \perp P_2 \Rightarrow \theta = 90^\circ$



$a_1a_2 + b_1b_2 + c_1c_2 = 0$

• If  $P_1 \parallel P_2 \Rightarrow \theta = 0^\circ$

$\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$

Angle Between a Line and a Plane

Coplanarity of Two Lines



Line :  $\vec{r} = \vec{a} + \lambda \vec{b}$       Plane :  $\vec{r} \cdot \vec{n} = d$

Therefore  $\cos \theta = \frac{|\vec{b} \cdot \vec{n}|}{|\vec{b}| \cdot |\vec{n}|}$

If  $L_1$  &  $L_2$  are coplaner, then

$(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) = 0$