



# DERIVATIVE

Part I

## Derivative by First Principle





$$\frac{dy}{dx} = \lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x} = f'(x) \rightarrow \text{Instantaneous rate of change of } y \text{ w.r.t. } x.$$

## Fundamental Rules for Differentiation

|   |                      |  |
|---|----------------------|--|
| 1 | <b>PRODUCT RULE</b>  |  $\frac{d}{dx} [f(x).g(x)] = f(x) \frac{d}{dx} \{g(x)\} + g(x) \frac{d}{dx} \{f(x)\}$   |
| 2 | <b>QUOTIENT RULE</b> | $\frac{d}{dx} \left\{ \frac{f(x)}{g(x)} \right\} = \frac{g(x) \cdot \frac{d}{dx} \{f(x)\} - f(x) \cdot \frac{d}{dx} \{g(x)\}}{\{g(x)\}^2}$  |
| 3 | <b>CHAIN RULE</b>    | if $y = f(u)$ & $u = g(x)$ then $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$  |

## Derivative of Standard Functions

For function  $y = f(x)$ , the derivative of the function is  $\frac{dy}{dx}$

|   |   |
|---|---|
| $\frac{d}{dx} (\sin x) = \cos x$                     | $\frac{d}{dx} (\sec^{-1} x) = \frac{1}{ x \sqrt{x^2-1}},  x  > 1$   |
| $\frac{d}{dx} (\cos x) = -\sin x$   | $\frac{d}{dx} (\operatorname{cosec}^{-1} x) = \frac{-1}{ x \sqrt{x^2-1}},  x  > 1$  |
| $\frac{d}{dx} (\tan x) = \sec^2 x$  |  $\frac{d}{dx} (\cot^{-1} x) = \frac{-1}{1+x^2}, x \in \mathbb{R}$ |
| $\frac{d}{dx} (\sec x) = \sec x \cdot \tan x$   | $\frac{d}{dx} (x^n) = n \cdot x^{n-1}; x \in \mathbb{R}, n \in \mathbb{R}, x > 0$   |
| $\frac{d}{dx} (\operatorname{cosec} x) = -\operatorname{cosec} x \cdot \cot x$  | $\frac{d}{dx} (a^x) = a^x \cdot \ln a; a > 0, a \neq 1$   |
| $\frac{d}{dx} (\cot x) = -\operatorname{cosec}^2 x$  | $\frac{d}{dx} (e^{ax}) = ae^{ax}$   |
| $\frac{d}{dx} (\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}, -1 < x < 1$   | $\frac{d}{dx} (\log_a  x ) = \frac{1}{x} \log_a e$  |
| $\frac{d}{dx} (\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}, -1 < x < 1$  | $\frac{d}{dx} (\ln  x ) = \frac{1}{x}$                           |
| $\frac{d}{dx} (\tan^{-1} x) = \frac{1}{1+x^2}, x \in \mathbb{R}$  | $\frac{d}{dx} (\text{constant}) = 0$  |



## L' Hopital Rule

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{0}{0} \text{ or } \frac{\infty}{\infty}$$

$f(x)$  &  $g(x)$  are differentiable at  $x = a$



$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} = \lim_{x \rightarrow a} \frac{f''(x)}{g''(x)} = \dots \text{till the indeterminate form vanishes.}$$



Guillaume de L' Hopital

## Logarithmic Differentiation

$$\text{If } y = [f(x)]^{g(x)} \Rightarrow \ln y = g(x) \ln [f(x)]$$

$$\Rightarrow \frac{1}{y} \cdot \frac{dy}{dx} = \frac{d}{dx} \{g(x) \cdot \ln[f(x)]\} \Rightarrow \frac{dy}{dx} = [f(x)]^{g(x)} \cdot \left\{ \frac{d}{dx} [g(x) \ln f(x)] \right\}$$

## Parametric Differentiation



$$\text{If } x = f(t) \text{ \& } y = g(t) \text{ then } \frac{dy}{dx} = \frac{dy/dt}{dx/dt}$$

## Differentiation of Inverse Function

$$y = f(x) \text{ and } x = g(y) \text{ are inverse function of each other } \frac{dx}{dy} = \frac{1}{dy/dx} \text{ or } g'(y) = \frac{1}{f'(x)} \quad \text{Hourglass icon}$$

## Derivative of a Determinant

$$\text{If } F(x) = \begin{vmatrix} f(x) & g(x) & h(x) \\ l(x) & m(x) & n(x) \\ u(x) & v(x) & w(x) \end{vmatrix}$$

where  $f, g, h, l, m, n, u, v, w$  are differentiable function of  $x$  then

$$F'(x) = \begin{vmatrix} f'(x) & g'(x) & h'(x) \\ l(x) & m(x) & n(x) \\ u(x) & v(x) & w(x) \end{vmatrix} + \begin{vmatrix} f(x) & g(x) & h(x) \\ l'(x) & m'(x) & n'(x) \\ u(x) & v(x) & w(x) \end{vmatrix} + \begin{vmatrix} f(x) & g(x) & h(x) \\ l(x) & m(x) & n(x) \\ u'(x) & v'(x) & w'(x) \end{vmatrix}$$