

Topics : Projectile Motion, Rectilinear Motion

Type of Questions

Single choice Objective ('-1' negative marking) Q.1 to Q.6

(3 marks, 3 min.)

M.M., Min.

[18, 18]

Subjective Questions ('-1' negative marking) Q.7 to Q.8

(4 marks, 5 min.)


[8, 10]

Match the Following (no negative marking) (2 × 4) Q.10

(8 marks, 10 min.)

[8, 10]

- For ground to ground projectile motion equation of path is $y = 12x - \frac{3}{4}x^2$. Given that $g = 10 \text{ ms}^{-2}$. What is the range of the projectile?
(A) 36m (B) 30.6 m (C) 16 m (D) 12.4 m
- The vertical height of the projectile at time t is given by $y = 4t - t^2$ and the horizontal distance covered is given by $x = 3t$. What is the angle of projection with the horizontal?
(A) $\tan^{-1} 3/5$ (B) $\tan^{-1} 4/5$ (C) $\tan^{-1} 4/3$ (D) $\tan^{-1} 3/4$
- A particle A is projected with speed V_A from a point making an angle 60° with the horizontal. At the same instant, second particle B (lie in the same horizontal plane) is thrown vertically upwards from a point directly below the maximum height point of parabolic path of A, with velocity V_B . If the two particles collide then the ratio of V_A/V_B should be ;
(A) 1 (B) $2/\sqrt{3}$ (C) $\sqrt{3}/2$ (D) $\sqrt{3}$
- A car accelerates from rest at a constant rate α for some time after which it decelerates at a constant rate β to come to rest. If total time taken by car is t , then maximum velocity V will be :
(A) $V = t \frac{\alpha\beta}{\alpha - \beta}$ (B) $V = t \left(\frac{\beta^2}{\alpha - \beta} \right)$ (C) $V = t \left(\frac{\alpha^2}{\alpha + \beta} \right)$ (D) $V = t \left(\frac{\alpha\beta}{\alpha + \beta} \right)$
- A lift is moving in upward direction with speed 20 m/s and having acceleration 5 m/s^2 in downward direction. A bolt drops from the ceiling of lift at that moment. Just after the drop, the :
(A) velocity of bolt with respect to ground is zero
(B) velocity of bolt with respect to ground is 20 m/s in upward direction
(C) acceleration of bolt with respect to ground is 5 m/s^2
(D) none of these
- A balloon is moving with constant upward acceleration of 1 m/s^2 . A stone is thrown from the balloon downwards with speed 10 m/s with respect to the balloon. At the time of projection balloon is at height 120 m from the ground and is moving with speed 20 m/s upward. The time required to fall on the ground by the stone after the projection will be-



- (A) 4 sec.
(C) 6 sec.


- (B) 5 sec.
(D) None of these




7. A particle is projected under gravity at an angle of projection 45° with horizontal. Its horizontal range is 36 m. Find maximum Height attained by particle.
8. A bullet is fired with speed 50 m/s at 45° angle with horizontal. Find the height of the bullet when its direction of motion makes angle 30° with the horizontal.
9. In the column-I, the path of a projectile (initial velocity 10 m/s and angle of projection with horizontal 60° in all cases) is shown in different cases. Range 'R' is to be matched in each case from column-II. Take $g = 10 \text{ m/s}^2$. Arrow on the trajectory indicates the direction of motion of projectile.

Column-I


Column-II




(p) $R = \frac{15\sqrt{3}}{2} \text{ m}$



(q) $R = \frac{40}{3} \text{ m}$



(r) $R = 5\sqrt{3} \text{ m}$



(s) $R = \frac{20}{3} \text{ m}$



Answers Key

DPP NO. - 16


1. (C) 2. (C) 3. (B) 4. (D) 5. (B)
6. (C) 7. 9
8. $h = \frac{125}{3}$ m above point of projection
9. (A) r (B) p (C) s (D) q

Hint & Solutions

DPP NO. - 16

1. $y = x \tan \theta \left(1 - \frac{x}{R}\right) \Rightarrow y = (12x) \left(1 - \frac{x}{16}\right)$

\Rightarrow Range = 16 m Ans.



$$y = 4t - t^2, \quad x = 3t$$

$$V_y = \frac{dy}{dt} = 4 - 2t, \quad V_x = \frac{dx}{dt} = 3$$

$$\Rightarrow u_y = v_y \Big|_{t=0} = 4, \quad u_x = v_x \Big|_{t=0} = 3$$

The angle of projection :

$$\tan \theta = \frac{V_y}{V_x} = \frac{4}{3} \Rightarrow \theta = \tan^{-1} \left(\frac{4}{3} \right) \text{ Ans.}$$

3. $V_A \sin 60^\circ = V_B$


$$\Rightarrow \frac{V_A}{V_B} = \frac{2}{\sqrt{3}}$$



4. $t = t_1 + t_2$

slope of OA curve = $\tan\theta = \alpha = \frac{V_{\max}}{t_1}$

slope of AB curve = $\beta = \frac{V_{\max}}{t_2}$



$$\Rightarrow t = \frac{V_{\max}}{\alpha} + \frac{V_{\max}}{\beta} \Rightarrow V_{\max} = \left(\frac{\alpha \beta}{\alpha + \beta} \right) t$$

5. The velocity of an object released in a moving frame is equal to that of the frame as observed from the frame.

6. velocity of ball w.r.t. ground = 20 – 10 = 10 m/sec upwards.

$$x = ut + \frac{1}{2} at^2$$

$$120 = -10t + \frac{1}{2} \times 10t^2$$

$$24 = -2t + t^2$$


$$t^2 - 2t - 24 = 0$$

$$t = 6 \text{ sec.}$$

7. $\frac{H}{R} = \frac{\tan\theta}{4}$

$$\theta = 45^\circ \text{ \& } R = 36 \text{ m}$$

$$H = 9 \text{ m}$$



h = height of the point where velocity makes 30° with horizontal.

As the horizontal component of velocity remain same
 $50 \cos 45^\circ = v \cos 30^\circ$

$$v = 50 \sqrt{\frac{2}{3}}$$

Now by equation

$$v^2 = u^2 + 2a_y y$$

$$\left(50 \times \sqrt{\frac{2}{3}}\right)^2 = 50^2 - 2gh$$

$$\Rightarrow 2gh = 50^2 - 50^2 \times \frac{2}{3}$$

$$\Rightarrow 2gh = \frac{1}{3} \times 50^2$$

$$\Rightarrow h = \frac{2500}{60} = \frac{125}{3}$$

$$h = \frac{125}{3} \text{ m above point of projection}$$

9. (A) $R = \frac{u^2 \sin 2\theta}{g} = \frac{100\sqrt{3}}{2(10)} = 5\sqrt{3} \text{ m}$

(B) $11.25 = -10 \sin 60^\circ t + \frac{1}{2} (10) t^2$


$$\Rightarrow 5t^2 - 5\sqrt{3} t - 11.25 = 0$$

$$t = \frac{5\sqrt{3} \pm \sqrt{25(3) + 4(5)(11.25)}}{10}$$

$$= \frac{5\sqrt{3} \pm \sqrt{3}(10)}{10}$$

$$= \frac{15}{10}\sqrt{3} = \frac{3}{2}\sqrt{3}$$

$$R = 10 \cos 60 \left(\frac{3}{2}\sqrt{3} \right) = 7.5\sqrt{3} \text{ m}$$




$$(C) t = \frac{2u \sin 30^\circ}{g \cos 30^\circ} = \frac{2(10)\left(\frac{1}{2}\right)}{10\left(\frac{\sqrt{3}}{2}\right)} = \frac{2}{\sqrt{3}} \text{ sec.}$$

$$R = 10 \cos 30^\circ t - \frac{1}{2} g \sin 30^\circ t^2$$

$$= \frac{10\sqrt{3}}{2} \left(\frac{2}{\sqrt{3}} \right) - \frac{1}{2} (10) \left(\frac{1}{2} \right) \frac{4}{3}$$

$$= 10 - \frac{10}{3} = \frac{20}{3} \text{ m}$$

$$(D) T = \frac{2(10)}{g \cos 30} = \frac{2(10)}{10\left(\frac{\sqrt{3}}{2}\right)} = \frac{4}{\sqrt{3}} \text{ sec.}$$



$$R = \frac{1}{2} g \sin 30^\circ t^2$$

$$= \frac{1}{2} (10) \left(\frac{1}{2} \right) \frac{16}{3} = \frac{40}{3} \text{ m}$$