

Topics : Newton's Law of Motion, Relative Motion

Type of Questions

Single choice Objective ('-1' negative marking) Q.1 to Q.5

(3 marks, 3 min.)

M.M., Min.

[15, 15]

Multiple choice objective ('-1' negative marking) Q.6

(4 marks, 4 min.)


[4, 4]

Subjective Questions ('-1' negative marking) Q.7 to Q.8

(4 marks, 5 min.)

[8, 10]


1. A man of mass m stands on a platform of equal mass m and pulls himself by two ropes passing over pulleys as shown. If he pulls each rope with a force equal to half his weight, his upward acceleration would be :



- (A) $g/2$
(C) g

- (B) $g/4$
(D) zero

2. Two blocks of masses m_1 and m_2 are connected as shown in the figure. The acceleration of the block m_2 is (pulleys and strings are ideal) :




(A) $\frac{m_2 g}{m_1 + m_2}$

(B) $\frac{m_1 g}{m_1 + m_2}$

(C) $\frac{4m_2 g - m_1 g}{m_1 + m_2}$

(D) $\frac{m_2 g}{m_1 + 4m_2}$

3. In the system shown in figure assume that cylinder remains in contact with the two wedges. The velocity of cylinder is -



(A) $\sqrt{19 - 4\sqrt{3}} \frac{u}{2}$ m/s


(B) $\frac{\sqrt{13} u}{2}$ m/s

(C) $\sqrt{3} u$ m/s

(D) $\sqrt{7} u$ m/s




4. A system is as shown in the figure. All speeds shown are with respect to ground. Then the speed of Block B with respect to ground is :



- (A) 5 m/s (B) 10 m/s (C) 15 m/s (D) 7.5 m/s

5. A wedge of height 'h' is released from rest with a light particle P placed on it as shown. The wedge slides down an incline which makes an angle θ with the horizontal. All the surfaces are smooth, P will reach the surface of the incline in time:




- (A) $\sqrt{\frac{2h}{g \sin^2 \theta}}$ (B) $\sqrt{\frac{2h}{g \sin \theta \cos \theta}}$ (C) $\sqrt{\frac{2h}{g \tan \theta}}$ (D) $\sqrt{\frac{2h}{g \cos^2 \theta}}$





6. A block of weight 9.8N is placed on a table. The table surface exerts an upward force of 10 N on the block. Assume $g = 9.8 \text{ m/s}^2$.

- (A) The block exerts a force of 10N on the table
 (B) The block exerts a force of 19.8N on the table
 (C) The block exerts a force of 9.8N on the table
 (D) The block has an upward acceleration.

7. A block of mass $M_1 = 3 \text{ kg}$ on a smooth fixed inclined plane of angle 30° is connected by a cord over a small frictionless pulley to a second block of mass 2 kg hanging vertically. The tension in the cord and the acceleration of each block are _____ and _____ respectively.



8. In which of the following cases the magnitude of acceleration of the block A will be maximum (Neglect friction, mass of pulley and string)



Answers Key


DPP NO. - 23

1. (D) 2. (A) 3. (D) 4. (B) 5. (A)
6. (A,D)

Hint & Solutions

DPP NO. - 23

1. for (man + platform) system :
 $2mg - 4T = 2m(a)$



$$\Rightarrow 2mg - 4 \left(\frac{mg}{2} \right) = 2m(a) \quad [\because T = \frac{mg}{2}]$$

$$\Rightarrow a = 0$$

2. Let a = acceleration of m_1

$$\text{then acceleration of pulley} = \frac{a+0}{2} = \frac{a}{2}$$

If acceleration of $m_2 = b$

$$\text{Then } 0 + \frac{b}{2} = \frac{a}{2}$$

Hence $a = b$

$$T = m_1 a, \quad m_2 g - T = m_2 a$$


$$\therefore a = \frac{m_2 g}{m_1 + m_2}$$



3. Method - I

As cylinder will remain in contact with wedge A

$$V_x = 2u$$



As it also remain in contact with wedge B

$$u \sin 30^\circ = V_y \cos 30^\circ - V_x \sin 30^\circ$$

$$V_y = V_x \frac{\sin 30^\circ}{\cos 30^\circ} + \frac{U \sin 30^\circ}{\cos 30^\circ}$$


$$V_y = V_x \tan 30^\circ + u \tan 30^\circ$$

$$V_y = 3u \tan 30^\circ = \sqrt{3} u$$

$$V = \sqrt{V_x^2 + V_y^2} = \sqrt{7} u \text{ Ans.}$$

Method - II

In the frame of A



$$3u \sin 30^\circ = V_y \cos 30^\circ$$


$$\Rightarrow V_y = 3u \tan 30^\circ = \sqrt{3} u$$

$$\text{and } V_x = 2u$$

$$\Rightarrow V = \sqrt{V_x^2 + V_y^2} = \sqrt{7} u \text{ Ans.}$$

4. $l_1 + 2l_2 = \text{constant}$

$$\therefore \frac{dl_1}{dt} + \frac{2dl_2}{dt} = 0$$



$$(5 + 5) + 2(5 + v_B) = 0 \text{ or } v_B = 10 \text{ m/s}$$

5. Assume that acceleration of particle is a_p and acceleration of wedge is a_w

Then, $a_w = g \sin \theta$


From wedge constant

$$a_p = a_w \sin \theta = g \sin^2 \theta$$

$$h = \frac{1}{2} g \sin^2 \theta t^2$$

$$t = \sqrt{\frac{2h}{g \sin^2 \theta}}$$

6. From Newton's third law, the force exerted by table on block is equal to that exerted by block on the table. Therefore block exerts a 10 N force on table. Since the upward force on the block is larger than downward force, it moves upwards.




FBD of block $M_2 = 2 \text{ kg}$



$$20 - T = 2a \quad \dots\dots\dots(i)$$

FBD of block $M_1 = 3 \text{ kg}$



$$= 3 \times \frac{10\sqrt{3}}{2}$$

$$= 15\sqrt{3} \text{ N.}$$

$$= 15 \text{ N}$$

$$T - 15 = 3a \quad \dots\dots\dots(ii)$$

$$(i) + (ii)$$

$$5 = 5a$$

$$\Rightarrow a = 1 \text{ m/s}^2 \quad ; \quad T = 18 \text{ N.}$$

8. (i) $a = \frac{2mg - mg}{3m} = \frac{g}{3}$

(ii) $a = \frac{2mg - mg}{m} = g$

(iii) $a = \frac{2mg}{m} = 2g$

(iv) $a = \frac{2g}{3}$