






4. A particle of mass 1 kg moves from rest along a straight line due to action of a force  $F$  which varies with the displacement  $x$  as shown in graph - (Use  $\frac{1}{\sqrt{2}} = 0.7$  if needed)



- (A) maximum K.E. of particle is 25 J  
 (B) Total work done by force on particle up to  $x = 6\text{ m}$  is  $-5\text{ J}$   
 (C) There will be no power delivered by the particle at  $x = 3, 5.3$  and  $6\text{ m}$   
 (D) None of these
5. A particle is projected from ground with an initial velocity  $20\text{ m/sec}$  making an angle  $60^\circ$  with horizontal. If  $R_1$  and  $R_2$  are radius of curvatures of the particle at point of projection and highest point respectively, then find the value of  $\frac{R_1}{R_2}$
6. A block of mass  $m_1 = 1\text{ kg}$  is attached to a spring of force constant  $k = 24\text{ N/cm}$  at one end and attached to a string tensioned by mass  $m_2 = 5\text{ kg}$ . Deduce the frequency of oscillations of the system. If  $m_2$  is initially supported in hand and then suddenly released, find



- (a) instantaneous tension just after  $m_2$  is released.  
 (b) the maximum displacement of  $m_1$ .  
 (c) the maximum and minimum tensions in the string during oscillations.
7. A mass  $M$  is in static equilibrium on a massless vertical spring as shown in the figure. A ball of mass  $m$  dropped from certain height sticks to the mass  $m$  after colliding with it. The oscillations they perform reach to height ' $a$ ' above the original level of spring & depth ' $b$ ' below it.



- (a) Find the force constant of the spring.  
 (b) Find the oscillation frequency.  
 (c) What is the height above the initial level from which the mass  $m$  was dropped ?





$$mgh = \frac{1}{2}mv^2 + \frac{1}{5}mv^2$$

$$\begin{aligned} \text{Translational kinetic energy} &= \frac{mgh \times \frac{1}{2}}{\frac{1}{2} + \frac{1}{5}} \\ &= \frac{mgh \times \frac{1}{2}}{\frac{7}{10}} = \frac{5mgh}{7} \end{aligned}$$

$$\text{Rotational kinetic energy} = \frac{2mgh}{7}$$

4. (A) Maximum kinetic energy at  $x = 3\text{m}$ .  
 (B)  $\text{KE} = \text{work done} = \text{area under the curve}$


$$= 10 \times 2 + \frac{1}{2} \times 1 \times 10 = 25 \text{ J}$$

- (C)  $w_{\text{ret}} = \text{area under the curve}$

$$= 25 - \frac{1}{2} \times 3 \times 20 = -5 \text{ J}$$

- (D) Power  $P = F \cdot v$

5.  $R_1 = \frac{v_0^2}{g \cos \theta}$



$R_2 = \frac{(v_0 \cos \theta)^2}{g}$

$$\therefore \frac{R_1}{R_2} = \frac{1}{(\cos \theta)^3} = 8$$

**Ans. 8**

6.  $T = 2\pi \sqrt{\frac{m_1 + m_2}{K}}$

$$= 2\pi \sqrt{\frac{6}{2400}} = \frac{\pi}{10}$$

$$\Rightarrow f = \frac{10}{\pi}$$

Instantaneous tension just after  $m_2$  is released will be zero as the spring is unstressed.


Amplitude of  $m_1 = m_2g / K = 25 / 12 \text{ cm}$ , hence maximum displacement of  $m_1$  will be  $25/6 \text{ cm}$ .

7. [Ans: (a)  $K = \frac{2mg}{b-a}$

(b)  $\frac{1}{2\pi} \sqrt{\frac{k}{M+m}}$



$$(c) \left( \frac{M+m}{b-a} \right) \frac{ab}{m}$$



$$\text{Amplitude} = b - \frac{mg}{K} = a + \frac{mg}{K} \quad (\text{by diagram})$$

$$\Rightarrow K = \frac{2mg}{b-a}$$

$$(b) f = \frac{1}{T} = \frac{1}{2\pi} \sqrt{\frac{K}{M+m}}$$

(c) If the ball was dropped from 'h'.

$$\Rightarrow V_{\text{before collision}} = \sqrt{2gh}$$

conserving momentum :

$$m\sqrt{2gh} = (m+M)V' \quad \text{Where } V'$$

$$= \omega \sqrt{A^2 - x^2} = \sqrt{\frac{K}{M+m}} \cdot \sqrt{\left(b - \frac{mg}{K}\right)^2 - \left(\frac{mg}{K}\right)^2}$$

Squaring both sides and putting

$$K = \frac{2mg}{b-a}, \text{ get } h = \left( \frac{M+m}{b-a} \right) \frac{ab}{m}$$