

Topics : Calorimetry & Thermal Expansion, Electrostatics, Work, Power and Energy, Rigid Body Dynamics, String Wave, Geometrical Optics

Type of Questions

Type of Questions	M.M., Min.
Single choice Objective ('-1' negative marking) Q.1 to Q.5	(3 marks, 3 min.) [15, 15]
Subjective Questions ('-1' negative marking) Q.6	(4 marks, 5 min.) [4, 5]
Comprehension ('-1' negative marking) Q.7 to Q.9	(3 marks, 3 min.) [9, 9]
Match the Following (no negative marking) (2 × 4)	(8 marks, 10 min.) [8, 10]


1. Water of mass $m_2 = 1$ kg is contained in a copper calorimeter of mass $m_1 = 1$ kg. Their common temperature $t = 10^\circ\text{C}$. Now a piece of ice of mass $m_3 = 2$ kg and temperature is -11°C dropped into the calorimeter. Neglecting any heat loss, the final temperature of system is. [specific heat of copper = 0.1 Kcal/kg $^\circ\text{C}$, specific heat of water = 1 Kcal/kg $^\circ\text{C}$, specific heat of ice = 0.5 Kcal/kg $^\circ\text{C}$, latent heat of fusion of ice = 78.7 Kcal/kg]

- (A) 0°C (B) 4°C (C) -4°C (D) -2°C

2. Two identical spheres of same mass and specific gravity (which is the ratio of density of a substance and density of water) 2.4 have different charges of Q and $-3Q$. They are suspended from two strings of same length ℓ fixed to points at the same horizontal level, but distant ℓ from each other. When the entire set up is transferred inside a liquid of specific gravity 0.8 , it is observed that the inclination of each string in equilibrium remains unchanged. Then the dielectric constant of the liquid is

- (A) 2 (B) 3 (C) 1.5 (D) None of these

3. A block of mass 10 kg is released on a fixed wedge inside a cart which is moved with constant velocity 10 m/s towards right. Take initial velocity of block with respect to cart zero. Then work done by normal reaction (with respect to ground) on block in two seconds will be: ($g = 10$ m/s 2).




- (A) zero (B) 960 J
(C) 1200 J (D) none of these

4. If the frequency of a wave is increased by 25% , then the change in its wavelength will be: (medium not changed)


- (A) 20% increase (B) 20% decrease (C) 25% increase (D) 25% decrease

5. The moment of inertia of a thin sheet of mass M of the given shape about the specified axis is



- (A) $(3/2)MR^2$ (B) $(3/4)MR^2$
(C) $MR^2(1 + 1/\sqrt{2})$ (D) $MR^2/(2\sqrt{2})$

6. AB wire is vibrating in its fundamental mode. Wire AB is in resonance with resonance tube in which air column is also vibrating with its fundamental mode. Sound speed is 400 m/sec and linear mass density of AB wire is 10^{-4} kg/m and $g = 10$ m/sec 2 , value of mass $m = [\beta(10^{-1})]$ kg, then find value of β . Neglect the masses of wires in comparison to block's mass 'm'.







COMPREHENSION

A glass prism with a refracting angle of 60° has a refractive index 1.52 for red and 1.6 for violet light. A parallel beam of white light is incident on one face at an angle of incidence, which gives minimum deviation for red light. Find :

[Use: $\sin(50^\circ) = 0.760$; $\sin(31.6^\circ) = 0.520$; $\sin(28.4^\circ) = 0.475$; $\sin(56^\circ) = 0.832$; $\pi = 22/7$]

7. The angle of incidence at the prism is :
 (A) 30° (B) 40° (C) 50° (D) 60°
8. The angular width of the spectrum is :
 (A) 6° (B) 4.8° (C) 9.6° (D) 12°
9. The length of the spectrum if it is focussed on a screen by a lens of focal length 100 cm is :
 (A) $\frac{10\pi}{3}$ cm (B) $\frac{10\pi}{3}$ m (C) $\frac{5\pi}{3}$ cm (D) $\frac{5\pi}{3}$ m
10. The column I gives the two point charge system separated by $2a$ and the column II gives the variation of magnitude of electric field intensity at point on the x-axis. Match the situation in Column I with the results in Column II and indicate your answer by darkening appropriate bubbles in the 4×4 matrix given in the OMR.

Column – I



Column – II

(p) Increases as x increases
in the interval $0 \leq x < a$

(q) Decreases as x increases
in the interval $0 \leq x < a$

(r) Zero at $x = 0$



(s) Decreases as x increases
in the interval $a < x < \infty$

Answers Key

1. (A) 2. (C) 3. (B) 4. (B)
 5. (B) 6. 6 7. (C) 8. (A)
 9. (A) 10. (A) → (p, r, s), (B) → (p, s),
 (C) → (r, s), (D) → (q, s)

Hints & Solutions

1. Loss in heat from calorimeter + water as temperature changes from 10°C to 0°C
 $= m_1 C_1 10 + m_2 C_2 10$
 $= 1 \times 1 \times 10 + 1 \times 0.1 \times 10 = 11 \text{ kcal}$
 Gain in heat of ice as its temperature changes from -11°C to 0°C
 $= m_3 C_3 \times 11 = 2 \times 0.5 \times 11 = 11 \text{ kcal}$
 Hence ice and water will coexist at 0°C without any phase change.



$$\tan\theta = \frac{F_e}{V\rho g} = \frac{F_e/k}{V(\rho - \sigma)g}$$


$$\Rightarrow k = \frac{\rho}{\rho - \sigma} = \frac{2.4}{2.4 - 0.8} = 1.5$$

$$k = \frac{\rho}{\rho - \sigma} = \frac{2.4}{2.4 - 0.8} = 1.5$$

3. Because the acceleration of wedge is zero, the normal reaction exerted by wedge on block is

$$N = mg \cos 37^\circ .$$

The acceleration of the block is $g \sin 37^\circ$ along the incline and initial velocity of the block is $v = 10$ m/s horizontally towards right as shown in figure.



The component of velocity of the block normal to the incline is $v \sin 37^\circ$. Hence the displacement of the block normal to the incline in $t = 2$ second is

$$S = v \sin 37^\circ \times 2 = 10 \times \frac{3}{5} \times 2 = 12 \text{ m.}$$

\therefore The work done by normal reaction

$$W = mg \cos 37^\circ \quad S = 100 \times \frac{4}{5} \times 12 = 960 \text{ J}$$

4. Since, the medium has not changed, speed of wave remains same.

$$\Rightarrow v = f\lambda = \text{constant}$$

$$f_1 \lambda_1 = f_2 \lambda_2$$

$$\Rightarrow f_1 \lambda_1 = (1.25f_1) \lambda_2$$

(\because frequency increased by 25%)


$$\Rightarrow \lambda_2 = \frac{\lambda_1}{1.25}$$

$\Rightarrow \lambda_2$ decreases.

\Rightarrow % change in wavelength

$$= \frac{\lambda_1 - \lambda_2}{\lambda_1} \times 100 = \frac{\lambda_1 - \frac{\lambda_1}{1.25}}{\lambda_1} \times 100$$

$$= \frac{0.25}{1.25} \times 100 = \frac{100}{5} = 20\%$$



$$T_1 = 2T_0 = 2 \left[\frac{2m(2m)}{a} \right]_a$$

$$T_1 = \frac{8m}{3}g = \frac{80m}{3} \dots\dots\dots(i)$$

In resonance,

$$f_{\text{wire}} = f_{\text{tube}}$$

$$\frac{(1)V_1}{2l_1} = \frac{(1)V_2}{4l_2}$$


$$\frac{\left(\sqrt{\frac{T_1}{\mu}}\right)}{2(x)} = \frac{(400)}{4\left(\frac{x}{2}\right)}$$

$$\Rightarrow T_1 = \mu(16 \times 10^4)$$

From (i),

$$\frac{80}{3}m = 10^{-4}(16 \times 10^4)$$

$$m = 0.6 \text{ kg.}$$



$$\mu_R = 1.52$$

$$\mu_V = 1.6$$

Minimum deviation condition for red is $r = 30^\circ$

$$\Rightarrow (1) \sin i = (1.52) \sin 30^\circ$$

$$i = 50^\circ,$$

$$\delta_R = (50^\circ) 2 - 60^\circ$$

$$= 40^\circ$$

8. For violet light

$$(1) \sin 50^\circ = (1.6) \sin r$$

$$\therefore r = 28.4^\circ$$

$$r' = 31.6^\circ \quad (\because r + r' = A)$$

$$(1) \sin e = (1.6) \sin 31.6^\circ$$

$$\therefore e = 56^\circ,$$

$$\Rightarrow \delta_V = i + e - A = 50^\circ + 56^\circ - 60^\circ$$

$$= 46^\circ$$

$$\therefore \text{angular width} = \delta_V - \delta_R = 6^\circ$$

9. The length of the spectrum if it is focussed on a screen by a lens of focal length 100 cm is :

$$(A^*) \frac{10\pi}{3} \text{ cm}$$

$$(B) \frac{10\pi}{3} \text{ m}$$

$$(C) \frac{5\pi}{3} \text{ cm}$$

$$(D) \frac{5\pi}{3} \text{ m}$$

Sol. if $\theta = 100 \times 6 \times \frac{\pi}{180} \text{ cm} = \frac{10\pi}{3} \text{ cm}$

10. (A) \rightarrow (p, r, s), (B) \rightarrow (p, s), (C) \rightarrow (r, s), (D) \rightarrow (q, s)