

Topics : Friction, Electrostatics, Geometrical Optics, Relative Motion, Rigid Body Dynamics, Newton's Law of Motion

Type of Questions

Single choice Objective ('-1' negative marking) Q.1 to Q.3

(3 marks, 3 min.)

M.M., Min.

[9, 9]

Subjective Questions ('-1' negative marking) Q.4

(4 marks, 5 min.)

[4, 5]

Comprehension ('-1' negative marking) Q.5 to Q.7

(3 marks, 3 min.)


[9, 9]

Match the Following (no negative marking) (2 × 4) Q.8

(8 marks, 10 min.)

[8, 10]


1. A plank is held at an angle α to the horizontal (Fig.) on two fixed supports A and B. The plank can slide against the supports (without friction) because of its weight Mg . With what acceleration and in what direction, a man of mass m should move so that the plank does not move.



- (A) $g \sin \alpha \left(1 + \frac{m}{M}\right)$ down the incline
- (B) $g \sin \alpha \left(1 + \frac{M}{m}\right)$ down the incline
- (C) $g \sin \alpha \left(1 + \frac{m}{M}\right)$ up the incline
- (D) $g \sin \alpha \left(1 + \frac{M}{m}\right)$ up the incline
2. Two small electric dipoles each of dipole moment $p \hat{j}$ are situated at $(0, 0, 0)$ and $(r, 0, 0)$. The electric potential at a point $\left(\frac{r}{2}, \frac{\sqrt{3}r}{2}, 0\right)$ is :
- (A) $\frac{p}{4\pi \epsilon_0 r^2}$
- (B) 0
- (C) $\frac{p}{2\pi \epsilon_0 r^2}$
- (D) $\frac{p}{8\pi \epsilon_0 r^2}$
3. A mango tree is at the bank of a river and one of the branch of tree extends over the river. A tortoise lives in river. A mango falls just above the tortoise. The acceleration of the mango falling from tree appearing to the tortoise is (Refractive index of water is $\frac{4}{3}$ and the tortoise is stationary)
- (A) g
- (B) $\frac{3g}{4}$
- (C) $\frac{4g}{3}$
- (D) None of these
4. A balloon is ascending vertically with an acceleration of 0.4 m/s^{-2} . Two stones are dropped from it at an interval of 2 sec. Find the distance between them 1.5 sec. after the second stone is released. ($g = 10 \text{ m/sec}^2$)

COMPREHENSION

In the given figure $F=10\text{N}$, $R=1\text{m}$ mass of the body is 2kg and moment of inertia of the body about an axis passing through O and perpendicular to plane of body is 4kgm^2 . O is the centre of mass of the body.



5. Find the frictional force acting on the body if it performs pure rolling.
- (A) $\frac{20}{3}$ (B) $\frac{10}{3}$ (C) $\frac{5}{3}$ (D) None of these
6. The kinetic energy of the body in the above question after 3 seconds will be.
- (A) 75J (B) 80J (C) 82J (D) 85J
7. If ground is smooth, then the total kinetic energy after 3 seconds will be :
- (A) 105.5J (B) 112.5J (C) 115.5J (D) None of these
8. In the column-I, a system is described in each option and corresponding time period is given in the column-II. Suitably match them.

Column-I

- (A) A simple pendulum of length ' ℓ ' oscillating with small amplitude in a lift moving down with retardation $g/2$.
- (B) A block attached to an end of a vertical spring, whose other end is fixed to the ceiling of a lift, stretches the spring by length ' ℓ ' in equilibrium. It's time period when lift moves up with an acceleration $g/2$ is
- (C) The time period of small oscillation of a uniform rod of length ' ℓ ' smoothly hinged at one end. The rod oscillates in vertical plane.
- (D) A cubical block of edge ' ℓ ' and specific gravity $1/2$ is in equilibrium with some volume inside water filled in a large fixed container. Neglect viscous forces and surface tension. The time period of small oscillations of the block in vertical direction is

Column-II

$$(p) T = 2\pi \sqrt{\frac{2\ell}{3g}}$$

$$(q) T = 2\pi \sqrt{\frac{\ell}{g}}$$

$$(r) T = 2\pi \sqrt{\frac{2\ell}{g}}$$

$$(s) T = 2\pi \sqrt{\frac{\ell}{2g}}$$




Answers Key

1. (B) 2. (B) 3. (C) 4. 52 m
 5. (B) 6. (A) 7. (B)
 8. (A) p (B) q (C) p (D) s

Hints & Solutions

1. F.B.D. of man and plank are -



For plank to be at rest, applying Newton's second law to plank along the incline

$$Mg \sin \alpha = f \quad \dots\dots\dots(1)$$

and applying Newton's second law to man along the incline.

$$mg \sin \alpha + f = ma \quad \dots\dots\dots(2)$$

$$a = g \sin \alpha \left(1 + \frac{M}{m} \right) \text{ down the incline}$$

Alternate Solution :

If the friction force is taken up the incline on man, then application of Newton's second law to man and plank along incline yields.

$$f + Mg \sin \alpha = 0 \quad \dots\dots\dots(1)$$

$$mg \sin \alpha - f = ma \quad \dots\dots\dots(2)$$

Solving (1) and (2)

$$a = g \sin \alpha \left(1 + \frac{M}{m} \right) \text{ down the incline}$$


Alternate Solution :

Application of Newton's second law to system of man + plank along the incline yields

$$mg \sin \alpha + Mg \sin \alpha = ma$$

$$a = g \sin \alpha \left(1 + \frac{M}{m} \right) \text{ down the incline}$$

2. As $ON = MN = OM = r$
 So it is equilateral triangle :
 \therefore Potential at N due to two dipoles ;
 $V = V_1 + V_2$



$$= \frac{Kp \cos 60^\circ}{r^2} + \frac{Kp \cos(\pi - 60^\circ)}{r^2} = 0$$

3. $\frac{x}{1} = \frac{x_{rel}}{\mu} \quad x_{rel} = \mu x$

$$\frac{d^2 x_{rel}}{dt^2} = \mu \frac{d^2 x}{dt^2}$$

$$a_{rel} = \mu g$$

4. At position A balloon drops first particle
 So, $u_A = 0, a_A = -g, t = 3.5 \text{ sec.}$

$$S_A = \left(\frac{1}{2} g t^2 \right) \quad \dots\dots\dots(i)$$

Balloon is going upward from A to B in 2 sec. so distance travelled by balloon in 2 second.

$$\left(S_B = \frac{1}{2} a_B t^2 \right) \quad \dots\dots\dots(ii)$$

$$a_B = 0.4 \text{ m/s}^2, \quad t = 2 \text{ sec.}$$

$$S_1 = BC = (SB + SA) \quad \dots\dots\dots(iii)$$

Distance travelled by second stone which is dropped from balloon at B


$$u_2 = u_B = a_B t = 0.4 \times 2 = 0.8 \text{ m/s}$$


$$t = 1.5 \text{ sec.}$$

$$\left(S_2 = u_2 t - \frac{1}{2} g t^2 \right) \quad \dots\dots\dots(iv)$$

Distance between two stone

$$\Delta S = S_1 - S_2.$$





$$f = ma \quad \dots(i)$$

$$F2R - FR - fR = I\alpha \quad \dots(ii)$$

$$a = R\alpha \quad \dots(iii)$$

$$FR - fR = I \cdot \frac{a}{R}$$

$$F - ma = \frac{Ia}{R^2}$$

$$F = \left(m + \frac{I}{R^2} \right) a.$$

$$a = \frac{F}{m + I/R^2}$$

$$f = ma = \frac{mF}{m + \frac{I}{R^2}}$$

$$f = \frac{mF}{m + \frac{I}{R^2}}$$

$$f = \frac{10}{3} \text{ N}$$

6. $a = \frac{F}{m + \frac{I}{R^2}} = \frac{5}{3}$

$$\alpha = \frac{a}{2} = \frac{5}{6}$$

$$v = 0 + \frac{5}{6} \times 3 = \frac{5}{2}$$

$$\omega = \omega_0 + \alpha t = 0 + \frac{5}{6} \times 3 = \frac{5}{2}$$

$$KE = \frac{1}{2}mv^2 + \frac{1}{2}I\omega^2$$

$$= \frac{1}{2} \times 2 \times 5 \times 5 + \frac{1}{2} \times 4 \times \frac{5}{2} \times \frac{5}{2}$$

$$= 25 + \frac{25}{2} = \frac{75}{2} \text{ J}$$

$$7. F2R - FR = I\alpha$$

$$\alpha = \frac{FR}{I}$$

$$\omega = 0 + \left(\frac{FR}{I}\right)t$$

$$KE = \frac{1}{2}I\omega^2$$

$$= \frac{1}{2} \times I \left(\frac{F^2 R^2}{I^2}\right) t^2$$

$$= \frac{F^2 R^2}{2I} = \frac{100 \times 1 \times 3 \times 3}{2 \times 4} = \frac{25 \times 9}{2} = 112.5 \text{ J.}$$

8. (A) p (B) q (C) p (D) s

(A) In frame of lift effective acceleration due to

gravity is $g + \frac{g}{2} = \frac{3g}{2}$ downwards

$$\therefore T = 2\pi \sqrt{\frac{2\ell}{3g}}$$

$$(B) K\ell = mg \quad \therefore \frac{k}{m} = \frac{g}{L}$$

constant acceleration of lift has no effect in time period of oscillation.

$$\therefore T = 2\pi \sqrt{\frac{m}{k}} = 2\pi \sqrt{\frac{\ell}{g}}$$

$$(C) T = 2\pi \sqrt{\frac{I}{mgd}} = 2\pi \sqrt{\frac{\frac{m\ell^2}{3}}{mg\frac{\ell}{2}}} = 2\pi \sqrt{\frac{2\ell}{3g}}$$

$$(D) T = 2\pi \sqrt{\frac{m}{\rho Ag}} = 2\pi \sqrt{\frac{\rho/2A\ell}{\rho Ag}} = 2\pi \sqrt{\frac{\ell}{2g}}$$

