


**Topics : Heat, Emf, Rotation, Center of Mass, Viscosity, Geometrical Optics, Current Electricity**


**Type of Questions**

Type of Questions	M.M., Min.
Single choice Objective ('-1' negative marking) Q.1 to Q.6	(3 marks, 3 min.) [18, 18]
Subjective Questions ('-1' negative marking) Q.7	(4 marks, 5 min.) [4, 5]
Match the Following (no negative marking) (2 × 4) Q.8	(8 marks, 10 min.) [8,10]
Assertion and Reason (no negative marking) Q. 9	(3 marks, 3 min.) [3, 3]

- A simple microscope has a focal length of 5 cm. The magnification at the least distance of distinct vision is-  
(A) 1 (B) 5 (C) 4 (D) 6
- Two identical solid spheres have the same temperature. One of the sphere is cut into two identical pieces. The intact sphere radiates an energy Q during a given small time interval. During the same interval, the two hemispheres radiate a total energy Q'. The ratio Q'/Q is equal to :  
(A) 2.0 (B) 4.0 (C)  $\frac{2}{3}$  (D) 1.5
- A ring of radius 5 m is lying in the x-y plane and is carrying current of 1 A in anti-clockwise sense. If a uniform magnetic field  $\vec{B} = 3\hat{i} + 4\hat{j}$  is switched on, then the co-ordinates of point about which the loop will lift up is:




- (A) (3, 4) (B) (4, 3)  
(C) (3, 0) (D) (0, 3)
- A ring of radius R rolls without slipping on a rough horizontal surface with a constant velocity. The radius of curvature of the path followed by any particle of the ring at the highest point of its path will be :



- (A) (B) 2 R  
(C) 4 R (D) none of these



5. A small bob of mass 'm' is suspended by a massless string from a cart of the same mass 'm' as shown in the figure. The friction between the cart and horizontal ground is negligible. The bob is given a velocity  $V_0$  in horizontal direction as shown. The maximum height attained by the bob is,



- (A)  $\frac{2V_0^2}{g}$       (B)  $\frac{V_0^2}{g}$       (C)  $\frac{V_0^2}{4g}$       (D)  $\frac{V_0^2}{2g}$
6. Two identical spherical drops of water are falling (vertically downwards) through air with a steady velocity of 5 cm/sec. If both the drops coalesce (combine) to form a new spherical drop, the terminal velocity of the new drop will be- (neglect buoyant force on the drops.)
- (A)  $5 \times 2$  cm/sec      (B)  $5 \times \sqrt{2}$  cm/sec      (C)  $5 \times (4)^{1/3}$  cm/sec      (D)  $\frac{5}{\sqrt{2}}$  cm/sec.
7. A steel wire of length  $l$  has a magnetic moment  $M$ . It is then bent into a semicircular arc. What is the new magnetic moment ?
8. In each situation of column-I a statement regarding a point object and its image is given. In column-II four optical instruments are given which form the image of that object. Match the statement in column-I with the optical instruments in column-II.

**Column-I**

- (A) Real image of a real point object may be formed by  
 (B) Virtual image of a real point object may be formed by  
 (C) Real image of a virtual point object may be formed by  
 (D) Virtual image of a virtual point object may be formed by

**Column-II**

- (p) concave mirror  
 (q) convex mirror  
 (r) convex lens (surrounded by air)  
 (s) concave lens (surrounded by air)

9. **STATEMENT-1** : Two cells of unequal emf  $E_1$  and  $E_2$  having internal resistances  $r_1$  and  $r_2$  are connected as shown in figure. Then the potential difference across any cell cannot be zero.



**STATEMENT-2** : If two cells having nonzero internal resistance and unequal emf are connected across each other as shown, then the current in the circuit cannot be zero.



- (A) Statement-1 is True, Statement-2 is True; Statement-2 is a correct explanation for Statement-1  
 (B) Statement-1 is True, Statement-2 is True; Statement-2 is **NOT** a correct explanation for Statement-1  
 (C) Statement-1 is True, Statement-2 is False  
 (D) Statement-1 is False, Statement-2 is True.

# Answers Key

1. (D)    2. (D)    3. (A)    4. (C)
5. (C)    6. (C)    7.  $M' = m \times 2r = \frac{M}{l} \times \frac{2l}{\pi} = \frac{2M}{\pi}$
8. (A) p,r (B) p,q,r,s (C) p,q,r,s (D) q,s
9. (B)


# Hints & Solutions

1.  $MP = \left(1 + \frac{D}{f}\right) = \left(1 + \frac{25}{5}\right) = 6$

2. Heat radiated (at temp same temp)  $\propto A$   
 $\Rightarrow Q \propto 4\pi R^2$  and  $Q' \propto (4\pi R^2 + 2 \times \pi R^2)$

$$\Rightarrow \frac{Q'}{Q} = \frac{6\pi R^2}{4\pi R^2} = 1.5$$

Here  $\pi R^2$  is extra surface area of plane surface of one of the hemisphere.




Magnetic moment  $\vec{M} = \pi r^2 \hat{i}$  &  $\vec{B} = 3\hat{i} + 4\hat{j}$

$$\therefore \vec{\tau} = \vec{M} \times \vec{B} = \pi r^2 (3\hat{i} - 4\hat{j})$$

$\vec{\tau}$  will be along the direction shown .

Hence , the point about which the loop will be lift up will be : (3, 4)



$$\text{Radius of Curvature} = \frac{(\text{velocity})^2}{\text{Normal Acceleration}}$$

$$= \frac{(2v)^2}{v^2/R} = 4R$$



5. By linear momentum conservation in horizontal direction = for (bob + string + cart)

$$mV_0 = (m + m)v$$

$$v = \frac{V_0}{2}$$

By mechanical energy conservation for (bob + string + cart + earth)

$$\frac{1}{2} mV_0^2 + 0 + 0 = \frac{1}{2} (2m)v^2 + mgh + 0$$

$$\frac{1}{2} mV_0^2 - \frac{1}{2} (2m) \frac{V_0^2}{4} = mgh$$

Solving it,

$$h = \frac{V_0^2}{4g}$$

6. When two drops of radius  $r$  each combine to form a big drop, the radius of big drop will be given by

$$\frac{4}{3} \pi R^3 = \frac{4\pi}{3} r^2 + \frac{4\pi}{3} r^3$$

$$\text{or } R^3 = 2r^3 \quad \text{or } R = 2^{1/3}r \quad \text{Now}$$

$$\frac{V_R}{V_r} = \left(\frac{R}{r}\right)^2 = 2^{2/3} = 4^{1/3}$$

$$\therefore V_R = 5 \times 4^{1/3} \text{ cm/s}$$

7. If  $m$  is pole strength, then

$$m = m = \frac{M}{l}$$

When the wire is bent into a semicircular arc, the separation between the two poles changes from  $l$  to  $2l$ , where new magnetic moment of the steel wire,

$$M' = m \times 2r = \frac{M}{l} \times \frac{2l}{\pi} = \frac{2M}{\pi}$$

8. (A) Real image of a real object is formed by concave mirror and convex lens.  
 (B) Virtual image of a real object is formed by all four.  
 (C) Real image of a virtual object may be formed by all four.  
 (D) Virtual image of a virtual object may be formed by convex mirror and concave lens.  
 (A) p,r (B) p,q,r,s (C) p,q,r,s (D) q,s

9. Let  $E_1 < E_2$  and a current  $i$  flows through the circuit. Then the potential difference across cell of emf  $E_1$  is  $E_1 + ir_1$  which is positive, hence potential difference across this cell cannot be zero. Hence statement 1 is correct.

For current in the circuit to be zero, emf of both the cells should be equal. But  $E_1 \neq E_2$ . Hence statement 2 is correct but it is not a correct explanation of statement 1.

