

**Topics : Electromagnet Induction, Rotation, Center of Mass, Magnetic Effect of Current and Magnetic Force on Charge/current**

**Type of Questions**

**Single choice Objective ('-1' negative marking) Q.1 to Q.5**

**(3 marks, 3 min.)**

**M.M., Min.**

**[15, 15]**

**Comprehension ('-1' negative marking) Q.6 to Q.8**

**(3 marks, 3 min.)**

**[9, 9]**


1. In the circuit shown switch S is connected to position 2 for a long time and then joined to position 1. The total heat produced in resistance  $R_1$  is :

(A)  $\frac{LE^2}{2R_2^2}$

(B)  $\frac{LE^2}{2R_1^2}$


(C)  $\frac{LE^2}{2R_1R_2}$

(D)  $\frac{LE^2(R_1+R_2)^2}{2R_1^2R_2^2}$



2. A tungsten bulb radiates 2 W of energy. The filament of the bulb has surface area  $2 \text{ mm}^2$  and an emissivity of 0.9. The temperature of the bulb is: (Stefan's Constant  $\sigma = 5.6 \times 10^{-8} \text{ S.I. units}$ )  
 (A) 3500K (B) 4210K (C) 2110K (D) 211K
3. A person with a defective sight is using a lens having a power of +2D. The lens he is using is  
 (A) concave lens with  $f = 0.5 \text{ m}$  (B) convex lens with  $f = 2.0 \text{ m}$   
 (C) concave lens with  $f = 0.2 \text{ m}$  (D) convex lens with  $f = 0.5 \text{ m}$

4. A uniform solid cylinder is given an angular speed  $\omega$  and placed on a rough plate of negligible thickness. The horizontal surface below the plate is smooth. Then the angular speed of the cylinder when it starts pure rolling on the plate will be: [ Assume sufficient length of plate ]



(A)  $\frac{\omega}{2}$

(B)  $\frac{\omega}{3}$

(C)  $\frac{2\omega}{3}$

(D) none of these

5. Four blocks of masses  $M_1, M_2, M_3$  and  $M_4$  are placed on a smooth horizontal surface along a straight line as shown. It is given that  $M_1 \gg M_2 \gg M_3 \gg M_4$ . All the blocks are initially at rest.  $M_1$  is given initial velocity  $v_0$  towards right such that it will collide with  $M_2$ . Consider all collisions to be perfectly elastic. The speed of  $M_4$  after all collision are over is



(A)  $v_0$

(B)  $4 v_0$

(C)  $8 v_0$

(D)  $16 v_0$

**COMPREHENSION**

A uniform and constant magnetic field  $\vec{B} = (20\hat{i} - 30\hat{j} + 50\hat{k})$  Tesla exists in space. A charged particle with charge to mass ratio  $\left(\frac{q}{m}\right) = \frac{10^3}{19} \text{ C/kg}$  enters this region at time  $t = 0$  with a velocity  $\vec{V} = (20\hat{i} + 50\hat{j} + 30\hat{k}) \text{ m/s}$ . Assume that the charged particle always remains in space having the given magnetic field. (Use  $\sqrt{2} = 1.4$ )

6. During the further motion of the particle in the magnetic field, the angle between the magnetic field  $\vec{B}$  and velocity of the particle  
 (A) remains constant (B) increases  
 (C) decreases (D) may increase or decrease.

7. The frequency of the revolution of the particle in cycles per second will be

(A)  $\frac{10^3}{\pi\sqrt{19}}$

(B)  $\frac{10^4}{\pi\sqrt{38}}$

(C)  $\frac{10^4}{\pi\sqrt{19}}$

(D)  $\frac{10^4}{2\pi\sqrt{19}}$

8. The pitch of the helical path of the motion of the particle will be

(A)  $\frac{\pi}{125} \text{ m}$

(B)  $\frac{\pi}{125} \text{ m}$

(C)  $\frac{\pi}{215} \text{ m}$

(D)  $\frac{\pi}{250} \text{ m}$




# Answers Key

1. (A)      2. (C)      3. (D)      4. (D)  
 5. (C)      6. (A)      7. (B)      8. (D)


## Hints & Solutions

1. When the key is at position (B) for a long time ;  
 the energy stored in the inductor is :

$$U_B = \frac{1}{2} Li_0^2 = \frac{1}{2} \cdot L \cdot \left( \frac{E}{R_2} \right)^2 = \frac{L \cdot E^2}{2R_2^2}$$



This whole energy will be dissipated in the form of heat when the inductor is connected to  $R_1$  and no source is connected.



Hence (A).


2.  $P = Ae\sigma T^4$   
 $2 = 2 \times 10^{-6} \times 0.9 \times 5.6 \times 10^{-8} \times T^4$


$$T^4 = \frac{10^{14}}{0.9 \times 5.6}$$

$$T = 2110 \text{ k}$$

3.  $f = \frac{1}{p} = \frac{1}{2} \text{ metre}$

$f = 0.5 \text{ m}$  this is positive so lense is convex lense.





Condition for pure rolling

$$V - \omega R = V' \quad \dots\dots\dots(i)$$

Momentum conservation

$$m'V' = mV$$

$$\Rightarrow V' = \frac{mV}{m'} \quad \dots\dots\dots(ii)$$

From (i) and (ii)  $V = \frac{m'\omega_1 V}{m' - m}$

Torque of friction about point P is zero Angular momentum will remain conserved about this point

$$\frac{mR^2}{2} \omega = \frac{mR^2}{2} \omega_1 + mVR$$

Solving this we get  $\omega' = \frac{m' - m}{3m' - m} \cdot \omega$

5.  $M_1$  is very large as compared to  $M_2$ . Hence for collision between  $M_1$  and  $M_2$ ,  $M_1$  can be considered equivalent to a wall and  $M_2$  as a small block. Thus the velocity of  $M_2$  will be  $2v_o$  after collision with  $M_1$ . Similarly after collision between  $M_2$  and  $M_3$ , the velocity of  $M_3$  will be  $2(2v_o)$ . In sequence, the velocity of  $M_4$  shall be  $2(2(2v_o)) = 8v_o$  after collision with  $M_3$ .

6. The component of velocity of charged particle along the magnetic field does not change. The component of velocity of charged particle normal to magnetic field only changes in direction but always remains normal to magnetic field. Hence angle between velocity and magnetic field remains same.

$$7. f = \frac{qB}{2\pi m} = \frac{10^3}{19} \times \frac{\sqrt{3800}}{2\pi}$$

$$= \frac{10^4}{\pi\sqrt{38}}$$

8. Pitch =  $T \cdot V_{\parallel}$

$$= \frac{1}{f} \cdot \frac{\vec{V} \cdot \vec{B}}{|\vec{B}|} = \frac{\pi \sqrt{38} \cdot 400}{10^4 \cdot \sqrt{3800}} = \frac{4\pi}{10^3} \text{ m}$$

$$= \frac{\pi}{250} \text{ m}$$

