

Ordinary Thinking

Objective Questions

Oxidation, Reduction

- H_2O_2 reduces MnO_4^- ion to [KCET (Med.) 2000]
 - Mn^+
 - Mn^{2+}
 - Mn^{3+}
 - Mn^-
- When a sulphur atom becomes a sulphide ion [AMU 1999]
 - There is no change in the composition of atom
 - It gains two electrons
 - The mass number changes
 - None of these
- The ultimate products of oxidation of most of hydrogen and carbon in food stuffs are [DCE 2001]
 - H_2O alone
 - CO_2 alone
 - H_2O and CO_2
 - None of these
- When P reacts with caustic soda, the products are PH_3 and NaH_2PO_2 . This reaction is an example of [IIT 1980; Kurukshetra CEE 1993; CPMT 1997]
 - Oxidation
 - Reduction
 - Oxidation and reduction (Redox)
 - Neutralization
- Which one of the following does not get oxidised by bromine water [MP PET/PMT 1988]
 - Fe^{+2} to Fe^{+3}
 - Cu^+ to Cu^{+2}
 - Mn^{+2} to MnO_4^-
 - Sn^{+2} to Sn^{+4}
- In the reaction $H_2S + NO_2 \rightarrow H_2O + NO + S$. H_2S is
 - Oxidised
 - Reduced
 - Precipitated
 - None of these
- The conversion of PbO_2 to $Pb(NO_3)_2$ is
 - Oxidation
 - Reduction
 - Neither oxidation nor reduction
 - Both oxidation and reduction
- In the course of a chemical reaction an oxidant [MP PMT 1986]
 - Loses electrons
 - Gains electrons
 - Both loses and gains electron
 - Electron change takes place
- $2CuI \rightarrow Cu + CuI_2$, the reaction is [RPMT 1997]
 - Redox
 - Neutralisation
 - Oxidation
 - Reduction
- H_2S reacts with halogens, the halogens [JIPMER 2000]
 - Form sulphur halides
 - Are oxidised
 - Are reduced
 - None of these
- H_2O_2 reduces $K_4Fe(CN)_6$ [MP PMT 1985]
 - In neutral solution
 - In acidic solution
 - In non-polar solvent
 - In alkaline solution
- Max. number of moles of electrons taken up by one mole of NO_3^- when it is reduced to [DPMT 2002]
 - NH_3
 - NH_2OH
 - NO
 - NO_2
- In the reaction $3Mg + N_2 \rightarrow Mg_3N_2$ [MP PMT 1999]
 - Magnesium is reduced
 - Magnesium is oxidized
 - Nitrogen is oxidized
 - None of these
- When sodium metal is dissolved in liquid ammonia, blue colour solution is formed. The blue colour is due to [NCERT 1981]
 - Solvated Na^+ ions
 - Solvated electrons
 - Solvated NH_2^- ions
 - Solvated protons
- Following reaction describes the rusting of iron $4Fe + 3O_2 \rightarrow 4Fe^{3+} + 6O^{2-}$

Which one of the following statement is incorrect [NCERT 1981; MNR 1991; AIIMS 1998]

 - This is an example of a redox reaction
 - Metallic iron is reduced to Fe^{3+}
 - Fe^{3+} is an oxidising agent
 - Metallic iron is a reducing agent
- $SnCl_2$ gives a precipitate with a solution of $HgCl_2$. In this process $HgCl_2$ is [CPMT 1983]
 - Reduced
 - Oxidised
 - Converted into a complex compound containing both Sn and Hg
 - Converted into a chloro complex of Hg
- Oxidation involves [NCERT 1971, 81; CPMT 1980, 82, 83; MP PMT 1983]
 - Loss of electrons
 - Gain of electrons
 - Increase in the valency of negative part
 - Decrease in the valency of positive part
- Incorrect statement regarding rusting is [MP PET 2000]
 - Metallic iron is oxidised to Fe^{3+} ions
 - Metallic iron is reduced to Fe^{2-} ions
 - Oxygen gas is reduced to oxide ion
 - Yellowish - brown product is formed
- When copper turnings are added to silver nitrate solution, a blue coloured solution is formed after some time. It is because, copper [CPMT 1974, 79; DPMT 2000]
 - Displaces silver from the solution



552 Redox Reactions

- (b) Forms a blue coloured complex with $AgNO_3$
 (c) Is oxidised to Cu^{2+}
 (d) Is reduced to Cu^{2+}
20. Solution of sodium metal in liquid ammonia is strongly reducing due to the presence of the following in the solution
 [NCERT 1977; KCET (Med.) 2000]
 (a) Sodium atoms (b) Solvated electrons
 (c) Sodium hydride (d) Sodium amide
21. When Sn^{2+} changes to Sn^{4+} in a reaction [CPMT 1981]
 (a) It loses two electrons (b) It gains two electrons
 (c) It loses two protons (d) It gains two protons
22. Oxidation of thiosulphate ($S_2O_3^{2-}$) ion by iodine gives
 [NCERT 1976]
 (a) SO_3^{2-} (b) SO_4^{2-}
 (c) $S_4O_6^{2-}$ (d) $S_2O_6^{2-}$
23. $Zn^{2+}(aq) + 2e \rightarrow Zn(s)$. This is [CPMT 1985]
 (a) Oxidation (b) Reduction
 (c) Redox reaction (d) None of these
24. One gas bleaches the colour of flowers by reduction while the other by oxidation [EAMCET 1980]
 (a) CO and Cl_2 (b) SO_2 and Cl_2
 (c) H_2S and Br_2 (d) NH_3 and SO_2
25. Reduction involves [NCERT 1972]
 (a) Loss of electrons
 (b) Gain of electrons
 (c) Increase in the valency of positive part
 (d) Decrease in the valency of negative part
26. In a reaction between zinc and iodine, in which zinc iodide is formed, what is being oxidised [NCERT 1975]
 (a) Zinc ions (b) Iodide ions
 (c) Zinc atom (d) Iodine
27. Which one of the following reactions does not involve either oxidation or reduction [EAMCET 1982]
 (a) $VO_2^+ \rightarrow V_2O_5$ (b) $Na \rightarrow Na^+$
 (c) $CrO_4^{2-} \rightarrow Cr_2O_7^{2-}$ (d) $Zn^{2+} \rightarrow Zn$
28. In the following reaction,
 $3Br_2 + 6CO_3^{2-} + 3H_2O = 5Br^- + BrO_3^- + 6HCO_3^-$
 [MP PMT 1994, 95]
 (a) Bromine is oxidised and carbonate is reduced
 (b) Bromine is reduced and water is oxidised
 (c) Bromine is neither reduced nor oxidised
 (d) Bromine is both reduced and oxidised
29. In the following reaction,
 $4P + 3KOH + 3H_2O \rightarrow 3KH_2PO_2 + PH_3$ [Pb. PMT 2002]
 (a) P is oxidized as well as reduced
 (b) P is reduced only
 (c) P is oxidised only
 (d) None of these
30. In the following reaction
 $Cr_2O_7^{2-} + 14H^+ + 6I^- \rightarrow 2Cr^{3+} + 3H_2O + 3I_2$
 Which element is reduced [CPMT 1976]
 (a) Cr (b) H
 (c) O (d) I
31. The conversion of sugar $C_{12}H_{22}O_{11} \rightarrow CO_2$ is
 (a) Oxidation
 (b) Reduction
 (c) Neither oxidation nor reduction
 (d) Both oxidation and reduction
32. Which halide is not oxidised by MnO_2
 [MNR 1985; JIPMER 2000]
 (a) F (b) Cl
 (c) Br (d) I
33. When Fe^{2+} changes to Fe^{3+} in a reaction
 (a) It loses an electron (b) It gains an electron
 (c) It loses a proton (d) It gains a proton
34. In acid solution, the reaction $MnO_4^- \rightarrow Mn^{2+}$ involves
 [MP PMT 1989]
 (a) Oxidation by 3 electrons
 (b) Reduction by 3 electrons
 (c) Oxidation by 5 electrons
 (d) Reduction by 5 electrons
35. When iron or zinc is added to $CuSO_4$ solution, copper is precipitated. It is due to [CPMT 1974, 79]
 (a) Oxidation of Cu^{+2} (b) Reduction of Cu^{+2}
 (c) Hydrolysis of $CuSO_4$ (d) Ionization of $CuSO_4$
36. In the reaction, $4Fe + 3O_2 \rightarrow 4Fe^{3+} + 6O^{2-}$ which of the following statement is incorrect [UPSEAT 2001, 02]
 (a) A Redox reaction
 (b) Metallic iron is a reducing agent
 (c) Fe^{3+} is an oxidising agent
 (d) Metallic iron is reduced to Fe^{3+}
37. Which of the following is redox reaction [CBSE PMT 1997]
 (a) H_2SO_4 with $NaOH$
 (b) In atmosphere, O_3 from O_2 by lightning
 (c) Evaporation of H_2O
 (d) Nitrogen oxides form nitrogen and oxygen by lightning

Oxidizing and Reducing agent

1. Equation $H_2S + H_2O_2 \rightarrow S + 2H_2O$ represents
 [UPSEAT 2001]
 (a) Acidic nature of H_2O_2

- (b) Basic nature of H_2O_2 [CPMT 1996]
 (c) Oxidising nature of H_2O_2
 (d) Reducing nature of H_2O_2
2. In the reaction
 $C_2O_4^{2-} + MnO_4^- + H^+ \rightarrow Mn^{2+} + CO_2 + H_2O$
 the reductant is [EAMCET 1991]
 (a) $C_2O_4^{2-}$ (b) MnO_4^-
 (c) Mn^{2+} (d) H^+
3. A reducing agent is a substance which can [CPMT 1971, 74, 76, 78, 80; NCERT 1976]
 (a) Accept electron (b) Donate electrons
 (c) Accept protons (d) Donate protons
4. Which of the following is the most powerful oxidizing agent [MNR 1990; CPMT 2003]
 (a) F_2 (b) Cl_2
 (c) Br_2 (d) I_2
5. Of the four oxyacids of chlorine the strongest oxidising agent in dilute aqueous solution is [MP PET 2000]
 (a) $HClO_4$ (b) $HClO_3$
 (c) $HClO_2$ (d) $HOCl$
6. Identify the correct statement about H_2O_2 [AIIMS 1996]
 (a) It acts as reducing agent only [NCERT 1979; CPMT 1988]
 (b) It acts as both oxidising and reducing agent
 (c) It is neither an oxidiser nor reducer
 (d) It acts as oxidising agent only
7. Several blocks of magnesium are fixed to the bottom of a ship to [AIEEE 2003]
 (a) Keep away the sharks
 (b) Make the ship lighter
 (c) Prevent action of water and salt
 (d) Prevent puncturing by under-sea rocks
8. Which of the following behaves as both oxidising and reducing agents [AFMC 1995]
 (a) H_2SO_4 (b) SO_2
 (c) H_2S (d) HNO_3
9. The reaction $H_2S + H_2O_2 \rightarrow 2H_2O + S$ shows [JIPMER 2001]
 (a) Oxidizing action of H_2O_2
 (b) Reducing action of H_2O_2
 (c) Alkaline nature of H_2O_2
 (d) Acidic nature of H_2O_2
10. Which of the following is not a reducing agent [EAMCET 1987]
 (a) $NaNO_2$ (b) $NaNO_3$
 (c) HI (d) $SnCl_2$
11. Which of the following cannot work as oxidising agent
 (a) O_2 (b) $KMnO_4$
 (c) I_2 (d) None of these
12. H_2O_2 is used as [CPMT 1994]
 (a) An oxidant only
 (b) A reductant only
 (c) An acid only
 (d) An oxidant, a reductant and an acid
13. In $C + H_2O \rightarrow CO + H_2$, H_2O acts as [AFMC 1988]
 (a) Oxidising agent (b) Reducing agent
 (c) (a) and (b) both (d) None of these
14. Strongest reducing agent is [CPMT 1977; BHU 1984, 96; MP PET 1990; AMU 1999]
 (a) F^- (b) Cl^-
 (c) Br^- (d) I^-
15. A solution of sulphur dioxide in water reacts with H_2S precipitating sulphur. Here sulphur dioxide acts as [NCERT 1980]
 (a) As oxidising agent (b) A reducing agent
 (c) An acid (d) A catalyst
16. Which of these substances is a good reducing agent [NCERT 1979; CPMT 1988]
 (a) $NaOCl$ (b) HI
 (c) $FeCl_3$ (d) KBr
17. The strongest reducing agent is [MNR 1982]
 (a) HNO_2 (b) H_2S
 (c) H_2SO_3 (d) $SnCl_2$
18. Which one is an oxidising agent [DPMT 1996]
 (a) $FeSO_4$
 (b) HNO_3
 (c) $FeSO_4 \cdot (NH_4)_2SO_4 \cdot 6H_2O$
 (d) H_2SO_4
19. In which of the following reactions H_2O_2 is a reducing agent [CPMT 1981; NCERT 1981; BHU 1999]
 (a) $2FeCl_2 + 2HCl + H_2O_2 \rightarrow 2FeCl_3 + 2H_2O$
 (b) $Cl_2 + H_2O_2 \rightarrow 2HCl + O_2$
 (c) $2HI + H_2O_2 \rightarrow 2H_2O + I_2$
 (d) $H_2SO_3 + H_2O_2 \rightarrow H_2SO_4 + H_2O$
20. When $NaCl$ is dissolved in water the sodium ion becomes [NCERT 1976]
 (a) Oxidised (b) Reduced
 (c) Hydrolysed (d) Hydrated
21. Strongest reducing agent is [MNR 1984, 89]
 (a) K (b) Mg



554 Redox Reactions

- (c) *Al* (d) *Br*
(e) *Na*
22. Which substance is serving as a reducing agent in the following reaction
 $14H^+ + Cr_2O_7^{2-} + 3Ni \rightarrow 2Cr^{3+} + 7H_2O + 3Ni^{2+}$
 [CBSE PMT 1994; AFMC 2000; DPMT 2001]
 (a) H_2O (b) *Ni*
 (c) H^+ (d) $Cr_2O_7^{2-}$
23. Which of the following acid possesses oxidising, reducing and complex forming properties [MNR 1985]
 (a) HNO_3 (b) H_2SO_4
 (c) HCl (d) HNO_2
24. Which one is oxidising substance [CPMT 1997]
 (a) $C_2H_2O_2$ (b) CO
 (c) H_2S (d) CO_2
25. The compound that can work both as oxidising and reducing agent is [CPMT 1986; MP PET 2000]
 (a) $KMnO_4$ (b) H_2O_2
 (c) BaO_2 (d) $K_2Cr_2O_7$
26. Which one is oxidising agent in the reaction below
 $2CrO_4^{2-} + 2H^+ \rightarrow Cr_2O_7^{2-} + H_2O$ [CPMT 1997]
 (a) H^+ (b) $Cr_2O_4^-$
 (c) Cr^{++} (d) None of these
27. Which is the best description of the behaviour of bromine in the reaction given below
 $H_2O + Br_2 \rightarrow HOBr + HBr$ [CBSE PMT 2004]
 (a) Oxidised only
 (b) Reduced only
 (c) Proton acceptor only
 (d) Both oxidised and reduced
28. What is the oxidising agent in chlorine water [JEE Orissa 2004]
 (a) HCl (b) $HClO_2$
 (c) $HOCl$ (d) None of these
29. In the reaction
 $Ag_2O + H_2O_2 \rightarrow 2Ag + H_2O + O_2$, the H_2O_2 acts as [BHU 2004]
 (a) Reducing agent (b) Oxidising agent
 (c) Bleaching agent (d) None of the above
30. In the reaction
 $HAsO_2 + Sn^{2+} \rightarrow As + Sn^{4+} + H_2O$ oxidising agent is [BVP 2004]
 (a) Sn^{2+} (b) Sn^{4+}
 (c) *As* (d) $HAsO_2$
31. Which of the following substances acts as an oxidising as well as a reducing agent [UPSEAT 2004; DCE 2004]
 (a) Na_2O (b) $SnCl_2$
 (c) Na_2O_2 (d) $NaNO_2$
32. In the reaction
 $P + NaOH \rightarrow PH_3 + NaH_2PO_2$ [MP PET 2004]
 (a) *P* is oxidised only
 (b) *P* is reduced only
 (c) *P* is oxidized as well as reduced
 (d) *Na* is reduced

Oxidation number and Oxidation state

1. The oxidation number of *C* in CO_2 is [MP PET 2001]
 (a) - 2 (b) + 2
 (c) - 4 (d) + 4
2. The oxidation number of *As* is [RPMT 1997]
 (a) + 2 and + 3 (b) + 3 and + 5
 (c) + 3 and + 4 (d) None of these
3. The oxidation number of *Ba* in barium peroxide is [Pb. PMT 2002]
 (a) + 6 (b) + 2
 (c) 1 (d) + 4
4. HNO_2 acts both as reductant and oxidant, while HNO_3 acts only as oxidant. It is due to their [AIIMS 2000]
 (a) Solubility ability
 (b) Maximum oxidation number
 (c) Minimum oxidation number
 (d) Minimum number of valence electrons
5. Chlorine is in +1 oxidation state in [MP PMT 1981; NCERT 1974; CPMT 1971, 78]
 (a) HCl (b) $HClO_4$
 (c) ICl (d) Cl_2O
6. The valency of *Cr* in the complex $[Cr(H_2O)_4Cl_2]^+$ [MP PMT 2000]
 (a) 1 (b) 3
 (c) 5 (d) 6
7. In the conversion $Br_2 \rightarrow BrO_3^-$, the oxidation state of bromine changes from [EAMCET 1990; AMU 1999; RPMT 2002]
 (a) - 1 to - 1 (b) 0 to - 1
 (c) 0 to + 5 (d) 0 to - 5
8. In the chemical reaction $Cl_2 + H_2S \rightarrow 2HCl + S$, the oxidation number of sulphur changes from [MP PMT 1999]
 (a) 0 to 2 (b) 2 to 0
 (c) - 2 to 0 (d) - 2 to - 1
9. Oxidation number of cobalt in $K[Co(CO)_4]$ is [KCET 1996]
 (a) + 1 (b) + 3
 (c) - 1 (d) - 3

10. When $K_2Cr_2O_7$ is converted to K_2CrO_4 , the change in the oxidation state of chromium is [NCERT 1981]
 (a) 0 (b) 6
 (c) 4 (d) 3
11. The oxidation number of chlorine in $HOCl$
 (a) -1 (b) 0
 (c) +1 (d) +2
12. Oxidation number of S in S^{2-} is [CPMT 1979]
 (a) -2 (b) 0
 (c) -6 (d) +2
13. Oxidation number of N in $(NH_4)_2SO_4$ is [CPMT 1996]
 (a) -1/3 (b) -1
 (c) +1 (d) -3
14. In which compound, oxidation state of nitrogen is 1
 [MP PMT 1989]
 (a) NO (b) N_2O
 (c) NH_2OH (d) N_2H_4
15. Oxidation number of nickel in $Ni(CO)_4$
 [AIIMS 1984; MNR 1985; CPMT 1997;
 MP PET/PMT 1998; AMU 2000; 01]
 (a) 0 (b) +4
 (c) -4 (d) +2
16. The oxidation number of sulphur in H_2SO_4 is
 [CPMT 1979Pb. CET 2002]
 (a) -2 (b) +2
 (c) +4 (d) +6
17. Oxidation state of chlorine in perchloric acid is
 [EAMCET 1989]
 (a) -1 (b) 0
 (c) -7 (d) +7
18. Oxidation number of N in HNO_3 is
 [BHU 1997]
 (a) -3.5 (b) +3.5
 (c) -3, +5 (d) +5
19. The oxidation number of Mn in MnO_4^{-1} is
 (a) +7 (b) -5
 (c) +6 (d) +5
20. Sn^{++} loses two electrons in a reaction. What will be the oxidation number of tin after the reaction
 (a) +2 (b) Zero
 (c) +4 (d) -2
21. The oxidation state of Mn in K_2MnO_4
 [CPMT 1982, 83, 84; DPMT 1982;
 NCERT 1973; AMU 2000]
 (a) +2 (b) +7
 (c) -2 (d) +6
22. Oxidation number of oxygen in O_2 molecule is
 [CPMT 1984]
 (a) +1 (b) 0
 (c) +2 (d) -2
23. Maximum oxidation state of Cr is [RPMT 2002]
 (a) 3 (b) 4
 (c) 6 (d) 7
24. In which of the following compound transition metal has zero oxidation state [CBSE PMT 1999; BHU 2000]
 (a) CrO_5 (b) $NH_2.NH_2$
 (c) $NOClO_4$ (d) $[Fe(CO)_5]$
25. Carbon is in the lowest oxidation state in
 [NCERT 1979; MH CET 1999]
 (a) CH_4 (b) CCl_4
 (c) CF_4 (d) CO_2
26. Oxidation number of carbon in $H_2C_2O_4$ is
 [CPMT 1982]
 (a) +4 (b) +3
 (c) +2 (d) -2
27. The oxidation number of Pt in $[Pt(C_2H_4)Cl_3]^-$ is
 [MNR 1993]
 (a) +1 (b) +2
 (c) +3 (d) +4
28. The oxidation number of carbon in CH_2Cl_2 is
 [CPMT 1976; Pb. PET 1999; AFMC 2004]
 (a) 0 (b) +2
 (c) -2 (d) +4
29. The oxidation states of phosphorus vary from
 [CPMT 1976]
 (a) -3 to +5 (b) -1 to +1
 (c) -3 to +3 (d) -5 to +1
30. The process in which oxidation number increases is known as
 [CPMT 1976]
 (a) Oxidation (b) Reduction
 (c) Auto-oxidation (d) None of the above
31. The oxidation number of S in $H_2S_2O_8$ is [MP PET 2002]
 (a) +2 (b) +4
 (c) +6 (d) +7
32. The oxidation state of nitrogen in N_3H is
 [NCERT 1977, 81]
 (a) $+\frac{1}{3}$ (b) +3
 (c) -1 (d) $-\frac{1}{3}$
33. Which of the following statements is correct [AFMC 1997]
 (a) Hydrogen has oxidation number -1 and +1
 (b) Hydrogen has same electronegativity as halogens
 (c) Hydrogen will not be liberated at anode
 (d) Hydrogen has same ionization potential as alkali metals
34. The oxidation state of Cr in $[Cr(NH_3)_4Cl_2]^+$ is
 [AIEEE 2005]



556 Redox Reactions

35. Sulphur has highest oxidation state in
 (a) +3 (b) +2 (c) +1 (d) 0
 [EAMCET 1991]
36. The oxidation number of *Fe* and *S* in iron pyrites are
 (a) SO_2 (b) H_2SO_4
 (c) $Na_2S_2O_3$ (d) $Na_2S_4O_6$
 [RPMT 1997]
37. The oxidation number of nitrogen in NO_3^- is
 (a) 4, - 2 (b) 2, - 1
 (c) 3, - 1.5 (d) 3, - 1
 [CPMT 1982]
38. Oxidation state of elemental carbon is [MNR 1983]
 (a) - 1 (b) + 2
 (c) + 3 (d) + 5
39. The sum of the oxidation numbers of all the carbons in C_6H_5CHO is [EAMCET 1986]
 (a) 0 (b) 1
 (c) 2 (d) 3
40. Which one of the following has the highest oxidation number of iodine [CPMT 1982]
 (a) + 2 (b) 0
 (c) + 4 (d) - 4
41. The oxidation number of *N* in $N_2H_5^+$ [Pb. PMT 2001]
 (a) KI_3 (b) KI
 (c) IF_5 (d) KIO_4
42. In which of the following compounds the oxidation number of carbon is maximum
 (a) - 3 (b) (- 2)
 (c) - 1 (d) + 2
43. The oxidation state of chlorine in $KClO_4$ is [CPMT 1985]
 (a) $HCHO$ (b) $CHCl_3$
 (c) CH_3OH (d) $C_{12}H_{22}O_{11}$
44. The oxidation state of *I* in $H_4IO_6^-$ is [CBSE PMT 1994]
 (a) - 1 (b) + 1
 (c) + 7 (d) - 7
45. An element which never has a positive oxidation number in any of its compounds [AIIMS 1981]
 (a) + 7 (b) + 5
 (c) + 1 (d) - 1
46. In an oxidation process, oxidation number [CPMT 1976]
 (a) Boron (b) Oxygen
 (c) Chlorine (d) Fluorine
47. If HNO_3 changes into N_2O , the oxidation number is changed by [BHU 1997; AFMC 2001]
 (a) Decreases (b) Increases
 (c) Does not change (d) First increases then decreases
48. The characteristic oxidation number of atoms in free metals is [NCERT 1975]
 (a) 0 (b) + 2
 (c) + 1 (d) 0
49. In which one of the following changes there are transfer of five electrons [NCERT 1982]
 (a) $MnO_4^- \rightarrow Mn^{2+}$ (b) $CrO_4^{2-} \rightarrow Cr^{3+}$
 (c) $MnO_4^{2-} \rightarrow MnO_2$ (d) $Cr_2O_7^{2-} \rightarrow 2Cr^{3+}$
50. Oxidation number of *C* in $C_6H_{12}O_6$ is [KCET 1992]
 (a) + 6 (b) - 6
 (c) 0 (d) + 4
51. In which of the following compounds iron has lowest oxidation state [MNR 1984]
 (a) $FeSO_4 \cdot (NH_4)_2SO_4 \cdot 6H_2O$
 (b) $K_4Fe(CN)_6$
 (c) $Fe(CO)_5$
 (d) Fe_2O
 (e) K_2FeO_4
52. The oxidation number of hydrogen in MH_2 is [CPMT 1976]
 (a) + 1 (b) - 1
 (c) + 2 (d) - 2
53. Oxidation number of iodine varies from [CPMT 1982]
 (a) - 1 to +1 (b) - 1 to +7
 (c) +3 to +5 (d) - 1 to +5
54. When SO_2 is passed through acidic solution of potassium dichromate, then chromium sulphate is formed. Change in valency of chromium is [CPMT 1979]
 (a) +4 to +2 (b) +5 to +3
 (c) +6 to +3 (d) +7 to +2
55. The oxidation states of the most electronegative element in the products of the reaction of BaO_2 with dilute H_2SO_4 are [IIT 1991; CBSE PMT 1992; BHU 2000]
 (a) 0 and - 1 (b) - 1 and - 2
 (c) - 2 and 0 (d) - 2 and + 1
56. The highest oxidation state of *Mn* is shown by [MNR 1983; RPMT 1999]
 (a) K_2MnO_4 (b) $KMnO_4$
 (c) MnO_2 (d) Mn_2O_2
 (e) MnO
57. The oxidation number of carbon in CH_2O is [IIT 1982; EAMCET 1985; MNR 1990; UPSEAT 2001 CPMT 1997, 2004]
 (a) - 2 (b) + 2
 (c) 0 (d) + 4
58. Oxidation state of oxygen in hydrogen peroxide is [DPMT 1984; 91; CPMT 1988; MNR 1994; UPSEAT 2001; RPMT 2002; JEE Orissa 2004]
 (a) - 1 (b) + 1
 (c) 0 (d) - 2

59. The oxidation number of Cr in $K_2Cr_2O_7$ is
[CPMT 1981, 85, 90, 93, 99; KCET 1992; BHU 1988, 98; AFMC 1991, 99; EAMCET 1986; MP PMT 1996, 99, 2002; MP PET/PMT 1998; Bihar CEE 1995; RPET 2000]
(a) +6 (b) -7
(c) +2 (d) -2
60. In which of the following compounds transition metal is in oxidation state zero [NCERT 1982]
(a) $[Co(NH_3)_6]Cl_2$ (b) $[Fe(H_2O)_6]SO_4$
(c) $[Ni(CO)_4]$ (d) $[Fe(H_2O)_3](OH)_2$
61. Oxidation number of osmium (Os) in OsO_4 is [AIIMS 1999]
(a) +4 (b) +6
(c) +7 (d) +8
62. The atomic number of an element which shows the oxidation state of +3 is [CPMT 1989, 94]
(a) 13 (b) 32
(c) 33 (d) 17
63. The oxidation number of iron in the compound $K_4[Fe(CN)_6]$ is [NCERT 1976; MNR 1986; AIIMS 2000]
(a) +6 (b) +4
(c) +3 (d) +2
64. The brown ring complex compound is formulated as $[Fe(H_2O)_5NO]SO_4$. The oxidation state of iron is [EAMCET 1987; IIT 1987; MP PMT 1994; AIIMS 1997; DCE 2000]
(a) 1 (b) 2
(c) 3 (d) 0
65. Oxidation state of oxygen in F_2O is [BHU 1982; UPSEAT 2001; MH CET 2002]
(a) +1 (b) +2
(c) -1 (d) -2
66. Phosphorus has the oxidation state of +3 in [NCERT 1982; RPMT 1999]
(a) Orthophosphoric acid (b) Phosphorus acid
(c) Metaphosphoric acid (d) Pyrophosphoric acid
67. Oxidation number of P in $Mg_2P_2O_7$ is [CPMT 1989; MP PMT 1995]
(a) +3 (b) +2
(c) +5 (d) -3
68. The oxidation state of nitrogen is highest in [MP PMT 2001; BHU 2002]
(a) N_3H (b) NH_2OH
(c) N_2H_4 (d) NH_3
69. Oxidation number of P in KH_2PO_2 is [CPMT 1987; MH CET 1999]
(a) +1 (b) +3
(c) +5 (d) -4
70. The most common oxidation state of an element is -2. The number of electrons present in its outermost shell is
(a) 4 (b) 2
(c) 6 (d) 8
71. Sulphur has lowest oxidation number in [EAMCET 1993]
(a) H_2SO_3 (b) SO_2
(c) H_2SO_4 (d) H_2S
72. The oxidation number and covalency of sulphur in the sulphur molecule (S_8) are respectively [NCERT 1977]
(a) 0 and 2 (b) 6 and 8
(c) 0 and 8 (d) 6 and 2
73. In ferrous ammonium sulphate oxidation number of Fe is [CPMT 1988]
(a) +3 (b) +2
(c) +1 (d) -2
74. The oxidation number of nitrogen in NH_2OH is [NCERT 1981]
(a) +1 (b) -1
(c) -3 (d) -2
75. The oxidation number of phosphorus in $Ba(H_2PO_2)_2$ is [Kurukshetra CEE 1998; DCE 2004]
(a) -1 (b) +1
(c) +2 (d) +3
76. A compound is in its low oxidation state. Then its will be [DCE 2001]
(a) Highly acidic
(b) Highly basic
(c) Highest oxidising property
(d) Half acidic, half basic
77. The oxidation number and the electronic configuration of sulphur in H_2SO_4 is [KCET 2002]
(a) +4; $1s^2 2s^2 2p^6 3s^2$
(b) +2; $1s^2 2s^2 2p^6 3s^2 3p^2$
(c) +3; $1s^2 2s^2 2p^6 3s^2 3p^1$
(d) +6; $1s^2 2s^2 2p^6$
78. The oxidation number of Mn in $KMnO_4$ is [CPMT 1982, 83; EAMCET 1992, 93; RPET 1999]
(a) +7 (b) -7
(c) +1 (d) -1
79. Oxidation number of As atoms in H_3AsO_4 is [DPMT 2001]
(a) -3 (b) +4
(c) +6 (d) +5
80. In XeO_3 and XeF_6 the oxidation state of Xe is [MP PET 2003]
(a) +4 (b) +6
(c) +1 (d) +3
81. Oxidation number of carbon in CH_3-Cl is






- (a) + 2 (b) + 3
(c) + 1 (d) + 4
105. Oxidation number of N in NH_3 is [CPMT 1979; Pb CET 2004]
(a) - 3 (b) + 3
(c) 0 (d) + 5
106. What is the net charge on ferrous ion [AFMC 2004]
(a) + 2 (b) + 3
(c) + 4 (d) + 5
107. Which of the following elements never show positive oxidation number [CPMT 2004]
(a) O (b) Fe
(c) Ga (d) F
108. The oxidation state shown by silicon when it combines with strongly electropositive metals is [MH CET 2004]
(a) - 2 (b) - 4
(c) + 4 (d) - 2
109. The oxidation number of sulphur in H_2S is [Pb. CET 2002]
(a) - 2 (b) + 3
(c) + 2 (d) - 3
110. Oxidation number of nitrogen in $NaNO_2$ is [Pb. CET 2000]
(a) + 2 (b) + 3
(c) + 4 (d) - 3
111. Oxidation number of S in SO_4^{2-} [BCECE 2005]
(a) + 6 (b) + 3
(c) + 2 (d) - 2
112. The oxidation state of chromium in the final product formed by the reaction between KI and acidified potassium dichromate solution is [AIEEE 2005]
(a) +4 (b) +6
(c) +2 (d) +3
113. The oxidation state of I in IPO_4 is [Orissa JEE 2005]
(a) +1 (b) +3
(c) +5 (d) +7
4. When $KMnO_4$ acts as an oxidising agent and ultimately forms $[MnO_4]^{-2}$, MnO_2 , Mn_2O_3 , Mn^{+2} then the number of electrons transferred in each case respectively is [AIEEE 2002]
(a) 4, 3, 1, 5 (b) 1, 5, 3, 7
(c) 1, 3, 4, 5 (d) 3, 5, 7, 1
5. Starch paper is used to test for the presence of [NCERT 1979]
(a) Iodine (b) Oxidising agent
(c) Iodide ion (d) Reducing agent
6. How many moles of $K_2Cr_2O_7$ can be reduced by 1 mole of Sn^{2+} [MP PMT 2003]
(a) 1/3 (b) 1/6
(c) 2/3 (d) 1
7. $2MnO_4^- + 5H_2O_2 + 6H^+ \rightarrow 2Z + 5O_2 + 8H_2O$. In this reaction Z is [RPMT 2002]
(a) Mn^{+2} (b) Mn^{+4}
(c) MnO_2 (d) Mn
8. What is 'A' in the following reaction [MP PET 2003]
 $2Fe^{3+}_{(aq)} + Sn^{2+}_{(aq)} \rightarrow 2Fe^{2+}_{(aq)} + A$
(a) $Sn^{3+}_{(aq)}$ (b) $Sn^{4+}_{(aq)}$
(c) $Sn^{2+}_{(aq)}$ (d) Sn
9. For the redox reaction
 $MnO_4^- + C_2O_4^{2-} + H^+ \rightarrow Mn^{2+} + CO_2 + H_2O$
the correct coefficients of the reactants for the balanced reaction are [IIT 1988, 92; BHU 1995; CPMT 1997; RPMT 1999; DCE 2000; MP PET 2003]
- | | MnO_4^- | $C_2O_4^{2-}$ | H^+ |
|-----|-----------|---------------|-------|
| (a) | 2 | 5 | 16 |
| (b) | 16 | 5 | 2 |
| (c) | 5 | 16 | 2 |
| (d) | 2 | 16 | 5 |
10. Which of the following is a redox reaction [AIEEE 2002]
(a) $NaCl + KNO_3 \rightarrow NaNO_3 + KCl$
(b) $CaC_2O_4 + 2HCl \rightarrow CaCl_2 + H_2C_2O_4$
(c) $Mg(OH)_2 + 2NH_4Cl \rightarrow MgCl_2 + 2NH_4OH$
(d) $Zn + 2AgCN \rightarrow 2Ag + Zn(CN)_2$
11. Which of the following reaction is a redox reaction [MP PMT 2003]
(a) $P_2O_5 + 2H_2O \rightarrow H_4P_2O_7$
(b) $2AgNO_3 + BaCl_2 \rightarrow 2AgCl + Ba(NO_3)_2$
(c) $BaCl_2 + H_2SO_4 \rightarrow BaSO_4 + 2HCl$
(d) $Cu + 2AgNO_3 \rightarrow 2Ag + Cu(NO_3)_2$
12. Which of the following reactions involves oxidation-reduction

Redox reaction and Method for balancing Redox reaction

1. The value of x in the partial redox equation
 $MnO_4^- + 8H^+ + xe \rightleftharpoons Mn^{2+} + 4H_2O$ is
(a) 5 (b) 3
(c) 1 (d) 0
2. $C_2H_6(g) + nO_2 \rightarrow CO_2(g) + H_2O(l)$
In this equation, the ratio of the coefficients of CO_2 and H_2O is [KCET 1992]
(a) 1 : 1 (b) 2 : 3
(c) 3 : 2 (d) 1 : 3
3. The number of electrons involved in the reduction of $Cr_2O_7^{2-}$ in acidic solution to Cr^{3+} is [EAMCET 1983]
(a) 0 (b) 2
(c) 3 (d) 5





560 Redox Reactions

[NCERT 1972; AFMC 2000; Pb. CET 2004; CPMT 2004]




13. Which of the following is the strongest oxidising agent

[Pb. CET 2000]



14. In the balanced chemical reaction,



a, b, c and d respectively correspond to [AIIMS 2005]



15. The number of moles of KMnO_4 reduced by one mole of


KI in alkaline medium is: [CBSE PMT 2005]




Auto oxidation and Disproportionation

1. In the equation $\text{H}_2\text{S} + 2\text{HNO}_3 \rightarrow 2\text{H}_2\text{O} + 2\text{NO}_2 + \text{S}$

The equivalent weight of hydrogen sulphide is [BVP 2003]



2. If 1.2 g of metal displace 1.12 litre hydrogen at normal temperature and pressure, equivalent weight of metal would be [DPMT 2001]



3. Which one of the following nitrates will leave behind a metal on strong heating [AIEEE 2003]




4. To prevent rancidification of food material, which of the following is added [CPMT 1996]



5. Prevention of corrosion of iron by zinc coating is called

[MP PMT 1993; CPMT 2002]






6. The metal used in galvanizing of iron is

[MP PET 1985, 96]



7. In which of the following reactions there is no change in valency [NCERT 1974; CPMT 1978]




8. The equivalent weight of phosphoric acid (H_3PO_4) in the reaction $\text{NaOH} + \text{H}_3\text{PO}_4 \rightarrow \text{NaH}_2\text{PO}_4 + \text{H}_2\text{O}$ is


[AIIMS 1999]





9. What is the equivalent mass of IO_4^- when it is converted into I_2 in acid medium [Kerala PMT 2004]



10. For decolourization of 1 mole of KMnO_4 , the moles of H_2O_2 required is [AIIMS 2004]




11. In the reaction $\text{I}_2 + 2\text{S}_2\text{O}_3^{2-} \rightarrow 2\text{I}^- + \text{S}_4\text{O}_6^{2-}$ equivalent weight of iodine will be equal to [MP PET 2004]



12. The equivalent weight of KIO_3 in the reaction $2\text{Cr}(\text{OH})_3 + 4\text{OH}^- + \text{KIO}_3 \rightarrow 2\text{CrO}_4^{2-} + 5\text{H}_2\text{O} + \text{KI}$ is

[MP PMT 2004]



13. The product of oxidation of I^- with MnO_4^- in alkaline medium is [IIT-JEE Screening 2004]



14. In alkaline medium ClO_2 oxidize H_2O_2 in O_2 and reduced itself in Cl^- then how many mole of H_2O_2 will oxidize by one mole of ClO_2 [Kerala CET 2005]

- (a) 1.0 (b) 1.5
(c) 2.5 (d) 3.5
(e) 5.0

Critical Thinking

Objective Questions

1. In which of the following acid, which acid has oxidation reduction and complex formation properties
[UPSEAT 2001]
(a) HNO_3 (b) H_2SO_4
(c) HCl (d) HNO_2
2. The compound which could not act both as oxidising as well as reducing agent is [IIT Screening 1991]
(a) SO_2 (b) MnO_2
(c) Al_2O_3 (d) CrO
3. H_2S acts only as a reducing agent while SO_2 can act both as a reducing and oxidizing agent because [AMU 1999]
(a) S in H_2S has - 2 oxidation state
(b) S in SO_2 has oxidation state + 4
(c) Hydrogen in H_2S more +ve than oxygen
(d) Oxygen is more - ve in SO_2
4. Of all the three common mineral acids, only sulphuric acid is found to be suitable for making the solution acidic because [Kurukshetra CEE 2002]
(a) It does not react with $KMnO_4$ or the reducing agent
(b) Hydrochloric acid reacts with $KMnO_4$
(c) Nitric acid is an oxidising agent which reacts with reducing agent
(d) All of the above are correct
5. For H_3PO_3 and H_3PO_4 the correct choice is [IIT Screening 2003]
(a) H_3PO_3 is dibasic and reducing
(b) H_3PO_3 is dibasic and non-reducing
(c) H_3PO_4 is tribasic and reducing
(d) H_3PO_3 is tribasic and non-reducing
6. Match List I with List II and select the correct answer using the codes given below the lists
List I (Compound) List II (Oxidation state of N)
(A) NO_2 (1) + 5
(B) HNO (2) - 3
(C) NH_3 (3) + 4
(D) N_2O_5 (4) + 1
- Codes :
(a) A B C D
2 3 4 1
(b) A B C D
3 1 2 4
(c) A B C D
3 4 2 1
(d) A B C D
2 3 1 4
7. M^{+3} ion loses $3e^-$. Its oxidation number will be [CPMT 2002]
(a) 0 (b) + 3
(c) + 6 (d) - 3
8. In the reaction $Zn + 2H^+ + 2Cl^- \rightarrow Zn^{2+} + 2Cl^- + H_2$, the spectator ion is [AIIMS 2001]
(a) Cl^- (b) Zn^{2+}
(c) H^+ (d) All of these
9. The oxidation number of sulphur in $H_2S_2O_7$ and iron in $K_4Fe(CN)_6$ is respectively [AIIMS 2000]
(a) + 6 and + 2 (b) + 2 and + 2
(c) + 8 and + 2 (d) + 6 and + 4
10. Oxidation number of oxygen in potassium superoxide (KO_2) is [UPSEAT 1999, 2002]
(a) - 2 (b) - 1
(c) - 1/2 (d) - 1/4
11. One mole of N_2H_4 loses 10 mol of electrons to form a new compound Y. Assuming that all nitrogen appear in the new compound, what is the oxidation state of N_2 in Y? (There is no change in the oxidation state of hydrogen) [IIT 1981; Pb. PMT 1998]
(a) + 3 (b) - 3
(c) - 1 (d) + 5
12. Amongst the following identify the species with an atom in + 6 oxidation state [IIT Screening 2000]
(a) MnO_4^- (b) $Cr(CN)_6^{3-}$
(c) NiF_6^{2-} (d) CrO_2Cl_2
13. In which of the following compounds, is the oxidation number of iodine is fractional [BVP 2003]
(a) IF_3 (b) IF_2
(c) I_3^- (d) IF_7
14. The compound $YBa_2Cu_3O_7$ which shows superconductivity has copper in oxidation state Assume that the rare earth element Yttrium is in its usual +3 oxidation state [IIT 1994]
(a) 3/7 (b) 7/3
(c) 3 (d) 7



562 Redox Reactions

15. The oxidation number of sulphur in S_8, S_2F_2, H_2S respectively, are [IIT 1999]
 (a) 0, +1 and -2 (b) +2, +1 and -2
 (c) 0, +1 and +2 (d) -2, +1 and -2
16. Which one of the following reactions is not an example of redox reaction [Kurukshetra CEE 1998]
 (a) $Cl_2 + 2H_2O + SO_2 \rightarrow 4H^+ + SO_4^{2-} + 2Cl^-$
 (b) $Cu^{++} + Zn \rightarrow Zn^{++} + Cu$
 (c) $2H_2 + O_2 \rightarrow 2H_2O$
 (d) $HCl + H_2O \rightarrow H_3O^+ + Cl^-$
17. For the reactions, $C + O_2 \rightarrow CO_2; \Delta H = -393 J$
 $2Zn + O_2 \rightarrow 2ZnO; \Delta H = -412 J$ [AIEEE 2002]
 (a) Carbon can oxidise Zn
 (b) Oxidation of carbon is not feasible
 (c) Oxidation of Zn is not feasible
 (d) Zn can oxidise carbon
18. In the reaction $B_2H_6 + 2KOH + 2X \rightarrow 2Y + 6H_2$, X and Y are respectively [EAMCET 2003]
 (a) H_2, H_3BO_3 (b) HCl, KBO_3
 (c) H_2O, KBO_3 (d) H_2O, KBO_2
19. In a balanced equation $H_2SO_4 + xHI \rightarrow H_2S + yI_2 + zH_2O$, the values of x, y, z are [EAMCET 2003]
 (a) $x = 3, y = 5, z = 2$
 (b) $x = 4, y = 8, z = 5$
 (c) $x = 8, y = 4, z = 4$
 (d) $x = 5, y = 3, z = 4$
20. Which of the following can act as an acid and as a base [AMU 1999]
 (a) $HClO_3^-$ (b) $H_2PO_4^-$
 (c) HS^- (d) All of these
21. MnO_4^{2-} (1 mole) in neutral aqueous medium is disproportionate to [AIIMS 2003]
 (a) $2/3$ mole of MnO_4^- and $1/3$ mole of MnO_2
 (b) $1/3$ mole of MnO_4^- and $2/3$ mole of MnO_2
 (c) $1/3$ mole of Mn_2O_7 and $1/3$ mole of MnO_2
 (d) $2/3$ mole of Mn_2O_7 and $1/3$ mole of MnO_2
22. The conductivity of a saturated solution of $BaSO_4$ is $3.06 \times 10^{-6} \text{ ohm}^{-1} \text{ cm}^{-1}$ and its equivalent conductance is $1.53 \text{ ohm}^{-1} \text{ cm}^{-1} \text{ equivalent}^{-1}$. The K_{sp} of the $BaSO_4$ will be [KCET 1996]
 (a) 4×10^{-12} (b) 2.5×10^{-9}
 (c) 2.5×10^{-13} (d) 4×10^{-6}
23. When MnO_2 is fused with KOH , a coloured compound is formed, the product and its colour is [IIT Screening 2003]
 (a) K_2MnO_4 , purple green
 (b) $KMnO_4$, purple
 (c) Mn_2O_3 , brown
 (d) Mn_3O_4 black

Assertion & Reason

For AIIMS Aspirants

Read the assertion and reason carefully to mark the correct option out of the options given below :

- (a) If both assertion and reason are true and the reason is the correct explanation of the assertion.
 (b) If both assertion and reason are true but reason is not the correct explanation of the assertion.
 (c) If assertion is true but reason is false.
 (d) If the assertion and reason both are false.
 (e) If assertion is false but reason is true.

1. Assertion : SO_2 and Cl_2 both are bleaching agents.
 Reason : Both are reducing agents. [AIIMS 1995]
2. Assertion : Fluorine exists only in -1 oxidation state.
 Reason : Fluorine has $2s^2 2p^5$ configuration. [AIIMS 2001]
3. Assertion : Stannous chloride is a powerful oxidising agent which oxidises mercuric chloride to mercury.
 Reason : Stannous chloride gives grey precipitate with mercuric chloride, but stannic chloride does not do so. [AIIMS 2001]
4. Assertion : $HClO_4$ is a stronger acid than $HClO_3$.
 Reason : Oxidation state of Cl in $HClO_4$ is +VII and in $HClO_3$ +V. [AIIMS 2004]
5. Assertion : In a reaction $Zn(s) + CuSO_4(aq) \rightarrow ZnSO_4(aq) + Cu(s)$, Zn is a reductant but itself get oxidized.
 Reason : In a redox reaction, oxidant is reduced by accepting electrons and

Oxidation number and Oxidation state

1	d	2	b	3	b	4	b	5	d
6	b	7	c	8	c	9	c	10	a
11	c	12	a	13	d	14	b	15	a
16	d	17	d	18	d	19	a	20	c
21	d	22	b	23	c	24	d	25	a
26	b	27	b	28	a	29	a	30	a
31	c	32	d	33	a	34	a	35	b
36	a	37	d	38	a	39	d	40	d
41	b	42	b	43	c	44	a	45	d
46	b	47	d	48	d	49	a	50	c
51	c	52	b	53	b	54	c	55	b
56	b	57	c	58	a	59	a	60	c
61	d	62	a	63	d	64	b	65	b
66	b	67	c	68	a	69	a	70	c
71	d	72	a	73	b	74	b	75	b
76	c	77	d	78	a	79	d	80	b
81	b	82	c	83	c	84	b	85	c
86	c	87	d	88	d	89	b	90	c
91	c	92	a	93	d	94	c	95	d
96	b	97	d	98	b	99	b	100	c
101	a	102	d	103	a	104	b	105	a
106	a	107	d	108	b	109	a	110	b
111	a	112	d	113	b				

Redox reaction and Method for balancing Redox reaction

1	a	2	b	3	c	4	c	5	a
6	a	7	a	8	b	9	a	10	d
11	d	12	c	13	c	14	a	15	d

Auto oxidation and Disproportionation

1	d	2	a	3	d	4	b	5	a
6	b	7	c	8	d	9	b	10	c
11	a	12	d	13	a	14	c		

Critical Thinking Questions

1	d	2	c	3	a,b	4	d	5	a
6	c	7	c	8	a	9	a	10	c
11	a	12	d	13	c	14	b	15	a
16	d	17	d	18	d	19	c	20	d
21	a	22	d	23	a				

Assertion & Reason

- reductant is oxidized by losing electrons.
6. Assertion : Oxidation number of carbon in CH_2O is zero.
Reason : CH_2O formaldehyde, is a covalent compound.
7. Assertion : The oxidation numbers are artificial, they are useful as a 'book-keeping' device of electrons in reactions.
Reason : The oxidation numbers do not usually represent real charges on atoms, they are simply conventions that indicate what the maximum charge could possibly be on an atom in a molecule.
8. Assertion : H_2SO_4 cannot act as reducing agent.
Reason : Sulphur cannot increase its oxidation number beyond + 6.
9. Assertion : Equivalent weight of NH_3 in the reaction $N_2 \rightarrow NH_3$ is $17/3$ while that of N_2 is $28/6$.
Reason : Equivalent weight = $\frac{\text{Molecular weight}}{\text{number of } e^- \text{ lost or gained}}$.

Answers

Oxidation, Reduction

1	b	2	b	3	c	4	c	5	c
6	a	7	b	8	b	9	a	10	c
11	b	12	a	13	b	14	b	15	b
16	a	17	a	18	b	19	c	20	b
21	a	22	c	23	b	24	b	25	b
26	c	27	c	28	d	29	a	30	a
31	a	32	a	33	a	34	d	35	b
36	d	37	d						

Oxidizing and Reducing agent




1	c	2	a	3	b	4	a	5	d
6	b	7	c	8	b	9	a	10	b
11	c	12	d	13	a	14	d	15	a
16	b	17	b	18	bd	19	b	20	d
21	a	22	b	23	d	24	d	25	b
26	d	27	d	28	c	29	a	30	d
31	d	32	c						

564 Redox Reactions




1	c	2	b	3	e	4	b	5	a
6	b	7	a	8	a	9	a		

AS Answers and Solutions


Oxidation, Reduction




It shows oxidation and reduction (Redox) properties.




9. (a) $2CuI \rightarrow \overset{+1}{Cu} + \overset{+2}{Cu}I_2$. Oxidation and Reduction both occur so the reaction is redox.
10. (c) $H_2S + X_2(Cl, Br, I = X) \rightarrow 2HX + S$. Here the halogen are reduced.
11. (b) When H_2O_2 reduces with $K_4[Fe(CN)_6]$. It is present in acidic solution.



13. (b) In the given reaction oxidation state of Mg is changing from 0 to +2 while in nitrogen it is changing from 0 to -3. So oxidation of Mg and reduction of nitrogen takes place.
14. (b) When sodium metal is dissolved in liquid ammonia to form coloured solution. Dilute solutions are bright blue in colour due to the presence of solvated electrons.





15. (b) The metallic iron is oxidised to Fe^{+3} .




In this reaction $HgCl_2$ is reduced in Hg .

17. (a) It is the process in which electrons are lost (de-electronation).
18. (b) $4Fe + 3O_2 \rightarrow 4Fe^{3+} + 6O^{2-}$
19. (c) Cu is above of Ag in electrochemical series and thus $Cu + 2Ag^+ \rightarrow Cu^{2+} + 2Ag$ reaction occurs.
21. (a) $Sn^{2+} \rightarrow Sn^{4+} + 2e^-$. In this reaction Sn^{2+} change in Sn^{4+} it is called an oxidation reaction.




24. (b) SO_2 bleaches by reduction while chlorine bleaches colour of flowers by oxidation.

25. (b) It is the process in which electrons are gained (electronation).




In this reaction Zn atom oxidised to Zn^{2+} ion and iodine reduced to I^- .



In this reaction oxidation and reduction are not involved because there is no change in oxidation number.

28. (d) $3Br_2 + 6CO_3^{2-} + 3H_2O \rightarrow 5Br^- + BrO_3^- + 6HCO_3^-$. In this reaction bromine is oxidised as well as reduced.

29. (a) P is oxidized as well as reduced (as in option a).





30. (a) In this reaction oxidation occur.

31. (a) Fluorine has highest E° - value and more reactive than MnO_2 .


32. (a) $Fe^{2+} \rightarrow Fe^{3+} + e^-$ oxidation.

33. (a) $MnO_4^- \rightarrow Mn^{2+}$. In this reaction $5e^-$ are needed for the reduction of Mn^{2+} as:




In this reaction Cu^{2+} change in Cu^0 , hence it is called as reduction reaction.

34. (d) $4Fe + 3O_2 \rightarrow 4Fe^{3+} + 6O^{2-}$, in this reaction metallic iron is oxidised to Fe^{3+} .




Here O.N. of N increases from 0 in N_2 to +2 in NO , 2- and that of decreased from 0 in O_2 to -2 in O , therefore, it is a redox reaction.

Oxidizing and Reducing agent



564 Redox Reactions

The oxidation of S shows oxidising nature of H_2O_2 .



In this reaction $C_2O_4^{2-}$ act as a reducing agent.

3. (b) A substance which is capable of reducing other substances and is capable of donating electrons during reduction is called a reducing agent or reductant.

4. (a) Fluorine is a most powerful oxidizing agent because it consist of $E^\circ = +2.5 \text{ volt}$.

5. (d) $HClO$ is the strongest oxidising agent. The correct order of oxidising power is ${}^{+1}HClO > {}^{+3}HClO_2 > {}^{+5}HClO_3 > {}^{+7}HClO_4$.

6. (b) It acts both oxidizing and reducing agent.

7. (c) Prevent action of water and salt.

9. (a) In this reaction H_2O_2 acts as a oxidizing agent.

10. (b) $NaNO_2$, $SnCl_2$ and HI have reducing and oxidizing properties but $NaNO_3$ have only oxidizing property.

11. (c) Because I_2 is a reducing agent.

13. (a) In this reaction H_2O acts as oxidising agent.


14. (d) I^- act as a more reducing agent than other ions.

15. (a) When sulphur dioxide is react with H_2S here SO_2 act as an oxidising agent and H_2S act as reducing agent.

16. (b) HI (Hydrogen Iodide) is a good reducing agent than other compound.

17. (b) Hydrogen sulphide (H_2S) acts as strong reducing agent as it decomposes by evolving hydrogen.

19. (b) $Cl_2 + H_2O_2 \rightarrow 2HCl + O_2$. In this reaction chlorine reduced from zero to -1 oxidation state.



Sodium ion hydrated in water.


21. (a) Potassium has higher negative value of reduction potential hence it shows more reducing properties.

22. (b) The oxidation number of Ni changes from 0 to +1



23. (d) HNO_2 (Nitrous acid) acid acts as a oxidising, reducing agent and has complex formation properties.

24. (d) CO_2 is an oxidizing agent.


25. (b) Hydrogen peroxide (H_2O_2) act as a both oxidising and reducing agent.




In the above reaction the oxidation number of Br_2 increases from zero (in Br_2) to +1 (in $HOBr$) and decrease from zero (Br_2) to -1 (in HBr). Thus Br_2 is oxidised as well as reduced & hence it is a redox reaction.



$HOCl$ can furnish, nascent oxygen.




30. (d) Oxidizing agent itself, undergoes reduction during a redox reaction





Hence, here $HAsO_2$ is acting as oxidizing agent.

31. (d) $NaNO_2$ (Sodium nitrite) act both as oxidising as well as reducing agent because in it N atom is in +3 oxidation state (intermediate oxidation state)


Oxidising property



Reducing property



Oxidation number and Oxidation state




$x + 2(-2) = 0$; $x - 4 = 0$; $x = +4$.

3. (b) +2 it is a second group element.

4. (b) In *HNO_2 oxidation number of $N = +3$

In *HNO_3 oxidation number of $N = +5$.

5. (d) In case of Cl_2O chlorine shows +1 oxidation state.



$$x + 0 + 2(-1) = +1; \quad x - 2 = +1$$

$$x = +3 \text{ for Cr in complex.}$$

7. (c) $Br_2 \rightarrow BrO_3^-$, in this reaction oxidation state change from 0 to +5.

8. (c) Oxidation state of sulphur in H_2S is -2, while it is zero in 'S' i.e. in this reaction oxidation of sulphur and reduction of chlorine is takes place.

9. (c) $K[Co(CO)_4]$

$$1 + x + 0 = 0; \quad x = -1.$$

10. (a) $K_2Cr_2O_7 \rightarrow K_2CrO_4$. In this reaction no change in oxidation state of chromium.

11. (c) In hypochlorous acid chlorine atom has + 1 oxidation number.

12. (a) $S \rightarrow S^{2-}$ O.N. of S = -2.

13. (d) $(NH_4)_2SO_4 \rightleftharpoons 2NH_4^+ + SO_4^{2-}$



$$x + 4 = +1; \quad x = 1 - 4 = -3.$$

14. (b) In N_2O nitrogen have +1 oxidation state.

15. (a) If any central metal atom combined with carbonyl group than central metal atom shows always zero oxidation state.

16. (d) H_2SO_4

$$2 + x - 2 \times 4 = 0, \quad x = 8 - 2 = +6.$$

17. (d) $HClO_4$

$$1 + x - 2 \times 4 = 0; \quad 1 + x - 8 = 0$$

$$x = 8 - 1 = +7 \text{ oxidation state.}$$

18. (d) HNO_3 ; $1 + x - 6 = 0$; $x = +5$.

19. (a) Mn shows + 7 oxidation state in MnO_4^-



$$x + (-2 \times 4) = -1$$

$$x - 8 = -1$$

$$x = -1 + 8 = +7.$$

20. (c) $Sn^{2+} \rightarrow Sn^{4+} + 2e^-$

21. (d) K_2MnO_4

$$2 + x - 2 \times 4 = 0$$

$$x = 8 - 2 = +6.$$

22. (b) Each molecule always show zero oxidation state.

23. (c) Maximum oxi. state for Cr is + 6.

24. (d) In $[Fe(CO)_5]$, transition metal Fe has zero oxidation state.

25. (a) In (b, c, d) carbon show + 4 oxidation state while in (a) carbon show - 4 oxidation state.

26. (b) $H_2C_2O_4$

$$2 + 2x - 2 \times 4 = 0; \quad 2x = 8 - 2 = 6$$

$$x = \frac{6}{2} = +3.$$


27. (b) In complex $[Pt(C_2H_4)Cl_3]^-$ Pt have + 2 oxidation state.

28. (a) CH_2Cl_2

$$x + 2 - 2 = 0; \quad x = 0.$$

29. (a) Phosphorus shows - 3 to + 5 oxidation state.

31. (c) The chemical structure of $H_2S_2O_8$ is as follows:-



So the oxidation number of S should be :

$$2 \times (+1) + 2 \times X + 6 \times (-2) + 2 \times (-1) = 0 \text{ or } X = +6.$$

(for H) (for S) (for O) (for O-O)

32. (d) In hydrazoic acid (N_3H) nitrogen shows $-\frac{1}{3}$ oxidation state.



$$3x + 1 = 0, \quad 3x = -1, \quad x = -\frac{1}{3}.$$

33. (a) Hydrogen have oxidation no. + 1 and - 1.

34. (a) $[Cr(NH_3)_4Cl_2]^+$

$$x + 4 \times (0) - 2 = 1 \Rightarrow x + 0 - 2 = 1$$

$$\Rightarrow x = 1 + 2 = +3.$$

35. (b) $SO_2 = +4$

$$H_2SO_4 = +6$$

$$Na_2S_2O_3 = +2$$

$$Na_2S_4O_6 = +\frac{5}{2}.$$

36. (a) FeS_2 FeS_2

$$x - 4 = 0 \quad 4 + 2x = 0$$

$$x = +4 \quad 2x = -4$$

$$x = \frac{-4}{2} = -2.$$

37. (d) NO_3^-

$$x - 2 \times 3 = -1; \quad x = 6 - 1 = +5.$$

38. (a) Every element always shows zero oxidation state.

39. (d) In benzaldehyde all carbon atoms show - 4 oxidation state.



71. (d) $H_2\overset{*}{S}O_3 = +4$; $\overset{*}{S}O_2 = +4$
 $H_2\overset{*}{S}O_4 = +6$; $H_2\overset{*}{S} = -2$.
72. (a) The oxidation number of sulphur in the sulphur molecule (S_8) is 0 and 2.
73. (b) In ferrous ammonium sulphate Fe shows +2 oxidation state.
74. (b) $\overset{*}{N}H_2OH$
 $x + 2(+1) - 2 + 1 = 0$
 $x + 2 - 2 + 1 = 0$; $x = -1$.
75. (b) $Ba(H_2\overset{*}{P}O_2)_2$; $BaH_4\overset{*}{P}_2O_4$
 $2 + 4 + 2x - 8 = 0$; $2x = 2$
 $x = \frac{2}{2} = +1$.
77. (d) $H_2\overset{*}{S}O_4$
 $2 \times (+1) + x + 4 \times (-2) = 0$
 $+2 + x - 8 = 0$; $x = 8 - 2 = +6$
 Electronic configuration of sulphur in $H_2\overset{*}{S}O_4$ is
 $1s^2, 2s^2, 2p^6$.
78. (a) $\overset{*}{K}MnO_4$
 $1 + x - 2 \times 4 = 0$; $x = 8 - 1 = +7$.
79. (d) $H_3\overset{*}{As}O_4$
 $+3 + x - 2 \times 4 = 0$; $x = 8 - 3 = +5$.
80. (b) The oxidation state of Xe in both XeO_3 and XeF_6 is +6
 $\overset{*}{Xe}O_3$ $\overset{*}{Xe}F_6$
 $x - 2 \times 3 = 0$ $x - 6 = 0$
 $x = +6$ $x = +6$.
81. (b) $\overset{*}{C}H_3-Cl$
 $x + 3(+1) + (-1) \times 1 = 0$
 $x + 3 - 1 = 0$; $x + 2 = 0$
 $x = -2$.
82. (c) $Cr_2\overset{*}{O}_7^{2-}$
 $2x - 2 \times 7 = -2$; $2x = 14 - 2 = 12$
 $x = \frac{12}{2} = +6$.
83. (c) $H_2\overset{*}{S}O_3$
 $+2 + x - 2 \times 3 = 0$; $x = 6 - 2 = +4$.
84. (b) Two Cl atom shows +1 and -1 oxidation state.
85. (c) $\overset{*}{H}ClO_3$
 $1 + x - 2 \times 3 = 0$; $x = 6 - 1 = +5$.
86. (c) $5 \begin{array}{c} \overset{COOH}{|} \\ \overset{COOH}{|} \end{array} + 2\overset{+7}{KMnO}_4 + 3H_2\overset{*}{S}O_4 \rightarrow$
 $\overset{+2}{K}_2\overset{*}{S}O_4 + 2\overset{+2}{MnSO}_4 + 10CO_2 + 8H_2O$
 In this reaction oxidation state of Mn change from +7 to +2.
87. (d) Oxygen have +2 oxidation state in OF_2 .
89. (b) $\overset{*}{S}_2O_3^{2-}$
 $2x + 3(-2) = -2$; $x = +2$.
90. (c) $x + 2 \times (+1) + 2(-1) = 0$
 $x + 2 - 2 = 0$; $x = 0$ in CH_2Cl_2 .
91. (c) In potassium superoxide (KO_2) oxygen shows, $-\frac{1}{2}$ oxidation state.
92. (a) $\overset{*}{S}_2Cl_2$
 $2x + 2(-1) = 0$; $2x - 2 = 0$
 $x = +1$.
93. (d) $Na_2\overset{*}{S}_4O_6$
 $2 + 4x - 12 = 0$
 $4x = 10$ $x = \frac{10}{4}$ $x = \frac{5}{2}$.
94. (c) $\overset{+2}{CuSO}_4 + 2KI \rightleftharpoons \overset{+2}{K}_2\overset{*}{S}O_4 + CuI_2$
 $2CuI_2 \longrightarrow \overset{+1}{Cu}_2I_2 + I_2$
95. (d) $NH_4Cl \rightleftharpoons \overset{*}{NH}_4^+ + Cl^-$
 $x + 4 = +1$; $x = 1 - 4 = -3$.
96. (b) $2\overset{+4}{NO}_2 + H_2O \rightarrow \overset{+3}{HNO}_2 + \overset{+5}{HNO}_3$. In this reaction oxidation state changes.
97. (d) $\overset{*}{Fe}_3O_4$
 $3x + (-8) = 0$; $3x - 8 = 0$
 $3x = 8$; $x = \frac{8}{3}$.
99. (b) $\overset{*}{K}_2\overset{*}{Mn}O_4$ $\overset{*}{MnSO}_4$
 $2 + x - 8 = 0$ $x + 6 - 8 = 0$
 $x = +6$ $x = +2$.
100. (c) Chlorine have oxidation state -1 to +7.
101. (a) $\overset{*}{[Co(NH}_3)_4ClNO_2]$
 $x + 4(0) + 1(-1) + 1(-1) = 0$
 $x + 0 - 1 - 1 = 0$
 $x - 2 = 0$; $x = +2$.
102. (d) $\overset{*}{K}_4[Ni(CN)_4]$
 $4 \times (+1) + x + 4 \times (-1) = 0$



568 Redox Reactions

$$+4 + x - 4 = 0 \Rightarrow x = 0.$$

103. (a) Fluorine always shows -1 oxidation state in oxides.

104. (b) $K_3[Fe(CN)_6]$



$$1 \times 3 + x + (-1 \times 6) = 0$$

$$3 + x - 6 = 0; \quad x = +3.$$

105. (a) NH_3

$$x + 3(+1) = 0, \quad x = -3.$$

106. (a) ${}_{26}Fe \longrightarrow [Ar]3d^6 4s^2$



In +2 state *Fe* is called Ferrous & in +3 state as ferric.

107. (d) Fluorine is the most electronegative element in the periodic table so it never shows positive oxidation state.

108. (b) Silicon forms silicides with strongly electropositive metals (like *Na, Mg, K* etc.) In these compounds. It has oxidation number = -4.

109. (a) H_2S [O.N. of *H* = +1]

$$(+1) \times 2 + x = 0$$

$$2 + x = 0; \quad x = -2$$



110. (b) Let the oxidation number of *N* in $NaNO_2$ be *x*

$$+1 + x + (-2) \times 2 = 0$$

$$1 + x - 4 = 0; \quad x = +3$$

111. (a) $x = 8 - 2 = +6$

112. (d) $K_2Cr_2O_7 + 6KI + 7H_2SO_4 \rightarrow 4K_2SO_4 + Cr_2(SO_4)_3$



113. (b) Let the oxidation number of *I* in $IPO_4 = x$

$$\text{Oxidation number of } PO_4 = -3$$

$$x + (-3) = 0 \Rightarrow x = +3$$

Redox reaction and Method for balancing Redox reaction

1. (a) $MnO_4^- + 8H^+ + 5e^- \rightleftharpoons Mn^{++} + 4H_2O$.

2. (b) The balanced equation is $2C_2H_6 + 7O_2 \rightarrow 4CO_2 + 6H_2O$. Ratio of the coefficients of CO_2 and H_2O is 4 : 6 or 2 : 3.


3. (c) $Cr_2O_7^{2-} + 3e^- \rightarrow Cr^{3+}$

Reduction



In this reaction three electrons are required for the reduction of $Cr_2O_7^{2-}$ into Cr^{3+} .

4. (c) Number of e^- transferred in each case is 1, 3, 4, 5.

5. (a) Starch paper are used for iodine test



6. (a) $Cr_2O_7^{2-} + 14H^+ + 6e^- \rightarrow 2Cr^{3+} + 7H_2O$



It is clear from this equation that 3 moles of Sn^{2+} reduce one mole of $Cr_2O_7^{2-}$, hence 1 mol. of Sn^{2+} will reduce $\frac{1}{3}$ moles of $Cr_2O_7^{2-}$.



7. (a) $2MnO_4^- + 5H_2O_2 + 6H^+ \rightarrow 2Mn^{2+} + 5O_2 + 8H_2O$.

Reduction

8. (b) $2Fe^{3+} + Sn^{2+} \rightarrow 2Fe^{2+} + Sn^{4+}$

Oxidation

9. (a) $MnO_4^- + 8H^+ + 5e^- \rightarrow Mn^{2+} + 4H_2O \times 2$



Thus the coefficient of MnO_4^- , $C_2O_4^{2-}$ and H^+ in the above balanced equation respectively are 2, 5, 16.

10. (d). $Zn + 2AgCN \rightarrow 2Ag + Zn(CN)_2$.

Reductio

Oxidatio

Oxidation Reduction

11. (d) $Cu + 2AgNO_3 \rightarrow Cu(NO_3)_2 + 2Ag$. This is a redox reaction.

Oxidatio

12. (c) $H_2 + Br_2 \rightarrow 2H^+ + 2Br^-$

Reductio



13. (c) Higher is the reduction potential stronger is the oxidising agent. Hence in the given options. MnO_4^- is strongest oxidising agent.

14. (a) $IO_3^- + aI^- + bH^+ \rightarrow cH_2O + dI_2$

Step 1 : $I^- \rightarrow I_2$ (oxidation)

$IO_3^- \rightarrow I_2$ (reduction)

Step 2 : $2IO_3^- + 12H^+ \rightarrow I_2 + 6H_2O$

Step 3 : $2IO_3^- + 12H^+ + 10e^- \rightarrow I_2 + 6H_2O$

$2I^- \rightarrow I_2 + 2e^-$

Step 4 : $2IO_3^- + 12H^+ + 10e^- \rightarrow I_2 + 6H_2O$

$[2I^- \rightarrow I_2 + 2e^-] \times 5$

Step 5 : $2IO_3^- + 10I^- + 12H^+ \rightarrow 6I_2 + 6H_2O$

$IO_3^- + 5I^- + 6H^+ \rightarrow 3I_2 + 3H_2O$

On comparing, $a = 5$, $b = 6$, $c = 3$, $d = 3$

15. (d) In alkaline medium

$2KMnO_4 + KI + H_2O \rightarrow 2MnO_2 + 2KOH + KIO_3$.

Auto oxidation and Disproportionation

1. (d) $H_2S \rightarrow S + 2e^-$

Equivalent wt. = $\frac{\text{Mol. wt.}}{2} = \frac{34}{2} = 17$.

2. (a) $1.12 \text{ ltr } H_2 = 1.2 \text{ g}$; $\therefore 22.4 \text{ ltr } H_2 = 24 \text{ g}$.

3. (d) $2AgNO_3 \xrightarrow{\Delta} 2Ag + 2NO_2 + O_2$.

4. (b) To prevent rancidification of food material we add anti-oxidant which are called oxidation inhibitor.

6. (b) Zn^{2+} / Zn . $E^\circ = -0.76 \text{ V}$

Al^{3+} / Al $E^\circ = -1.662$

Sn^{2+} / Sn $E^\circ = -0.136$

Pb^{2+} / Pb $E^\circ = -0.126$

In galvanizing action Zn is coated over iron.

8. (d) Molecular weight of H_3PO_4 is 98 and change in

its valency = 1 equivalent wt. of H_3PO_4

= $\frac{\text{Molecular weight}}{\text{Change in valency}} = \frac{98}{1} = 98$.

9. (b) Equivalent mass

$$= \frac{\text{Molecular weight}}{\text{Change in oxidation number per mole}}$$


Suppose molecular weight is M

Oxidation number of I_2 in IO_4^- in

Acidic medium i.e., $I \times (-8) + 1e^- = +7$

So eq. wt. = $M/7$.

10. (c) $2KMnO_4 + 3H_2SO_4 \longrightarrow K_2SO_4 + 2MnSO_4 + 3H_2O + 5O$



11. (a) $\frac{\text{Molecular weight}}{2} = \text{Equivalent weight of Iodine.}$

12. (d) $\frac{\text{Molecular weight}}{3}$ Because in KIO_3 effective oxidation number is 3.

13. (a) $6MnO_4^- + I^- + 6OH^- \longrightarrow 6MnO_4^{2-} + IO_3^- + 3H_2O$

14. (c) $ClO_2 \rightarrow Cl^-$

$ClO_2 + 2H_2O + 5e^- \rightarrow Cl^- + 4OH^-$

$H_2O_2 \rightarrow O_2$

$H_2O_2 + 2OH^- \rightarrow O_2 + 2H_2O + 2e^-$

$ClO_2 + 2H_2O + 5e^- \rightarrow Cl^- + 4OH^- \times 2$

$H_2O_2 + 2OH^- \rightarrow O_2 + 2H_2O + 2e^- \times 2$

$2ClO_2 + 5H_2O_2 + 2OH^- \rightarrow 2Cl^- + 5O_2 + 5H_2O$

$2ClO_2 \equiv 5H_2O_2$

$\therefore ClO_2 = 2.5H_2O_2$

Critical Thinking Questions

- (d) HNO_2 shows both oxidation and reduction properties.
- (c) Al_2O_3 could not act as a oxidising and reducing agent.
- (a, b) In H_2S sulphur shows -2 oxidation state and in SO_2 shows +4 oxidation state. Hence SO_2 shows both oxidising and reducing properties.
- (d) All the given statements are true.

570 Redox Reactions

5. (a)
$$\begin{array}{c} H \\ | \\ H-O-P-OH \\ | \\ O \end{array}$$
, hence it is dibasic. It acts as reducing agent also.
6. (c) (a) $\overset{*}{N}O_2$; $x-4=0$; $x=+4$
 (b) $H\overset{*}{N}O$; $1+x-2=0$; $x=+1$
 (c) NH_3 ; $x+3=0$; $x=-3$
 (d) $N_2\overset{*}{O}_5$; $2x-10=0$; $2x=10$; $x=\frac{10}{2}$; $x=5$.
7. (c) $2 \times \text{No. of } e^- \text{ losses} = \text{Oxi. no.}$
 $2 \times 3e^- = +6$.
8. (a) The ion which is not affected during the course of reaction is known as spectator ion.
9. (a) $H_2S_2O_7$
 $2 \times (+1) + 2 \times x + 7 \times (-2) = 0$
 $+2 + 2x - 14 = 0$
 $2x = 14 - 2 = 12$
 $x = \frac{12}{2} = +6$ for S
 $K_4\overset{*}{Fe}(CN)_6$
 $4 \times (+1)x + 6 \times (-1) = 0$
 $4 + x - 6 = 0$
 $x = 6 - 4 = +2$ for Fe.
10. (c) $\overset{*}{K}O_2$, $+1 + 2x = 0$, $x = -\frac{1}{2}$.
11. (a) $N_2^{2-} \rightarrow 2N^{a+} + 10e^-$
 $\therefore 2a - [2 \times (-2)] = 10$
 $\therefore a = +3$.
12. (d) CrO_2Cl_2 , $x-4-2=0$, $x=+6$.
13. (c) $3x = -1$, $x = -1/3$.
14. (b) $Ba_2\overset{*}{Cu}_3O_7$
 $3 + 2 \times 2 + 3x - (2 \times 7) = 0$
 $3 + 4 + 3x - 14 = 0$
 $3x = 7$
 $x = \frac{7}{3}$.
15. (a) $\overset{*}{S}_8 = 0$
 $\overset{*}{S}_2F_2 = +1$
 $H_2S^* = -2$.
16. (d) In reaction $HCl + H_2O \rightarrow H_3O^+ + Cl^-$, only reduction has taken place not oxidation.
17. (d) Zn can oxidise carbon because heat of combustion of $Zn < C$.
18. (d) $B_2H_6 + 2KOH + 2H_2O \rightarrow 2KBO_2 + 6H_2$.
19. (c) The values of x, y, z are 8, 4, 4 respectively hence the reaction is
 $H_2SO_4 + 8HI \rightarrow H_2S + 4I_2 + 4H_2O$
20. (d)

Acid	Base
$HClO_3^-$	ClO_3^{2-}
HS^-	S^{2-}
$H_2PO_4^-$	HPO_4^{2-}
21. (a) MnO_4^{2-} in neutral aqueous medium is disproportionate to $\frac{2}{3}$ mole of MnO_4^- and $\frac{1}{3}$ mole of MnO_2 .
22. (d) $\lambda m = \frac{1000 K}{S} = \frac{1000 \times 3.06 \times 10^{-6}}{S} = 1.53$
 $S = 2 \times 10^{-3} \frac{\text{mol}}{\text{litre}}$
 $K_{sp}(BaSO_4) = S^2 = (2 \times 10^{-3})^2 = 4 \times 10^{-6}$.
23. (a) $2MnO_2 + 4KOH + O_2 \xrightarrow{\Delta} 2K_2MnO_4 + 2H_2O$
purple green

Assertion & Reason


1. (c) It is true that SO_2 and Cl_2 both are bleaching agents. But Cl_2 is an oxidising agent while SO_2 is a reducing agent. Therefore, in this questions assertion is true while reason is false.
2. (b) It is correct that fluorine exists only in -1 oxidation state because it has $1s^2 2p^5$ electronic configuration and thus shows only -1 oxidation state in order to complete its octet. Hence, both assertion and reason are true and reason is not a correct explanation of assertion.
3. (e) Here, assertion is false, because stannous chloride is a strong reducing agent not strong oxidising agent. Stannous chlorides gives Grey precipitate with mercuric chloride. Hence, reason is true.



$$NH_3 = \frac{14 + 3}{3} = \frac{17}{3} \text{ (M. wt. of } NH_3 \text{)}$$

$$\text{while for } N_2 = \frac{14 \times 2}{6} = \frac{28}{6}$$

4. (b) Both assertion and reason are true but reason is not the correct explanation of assertion. Greater the number of negative atoms present in the oxy-acid make the acid stronger. In general, the strengths of acids that have general formula $(HO)_mZO_n$ can be related to the value of n . As the value of n increases, acidic character also increases. The negative atoms draw electrons away from the Z-atom and make it more positive. The Z-atom, therefore, becomes more effective in with drawing electron density away from the oxygen atom that bonded to hydrogen. In turn, the electrons of $H-O$ bond are drawn more strongly away from the H-atom. The net effect makes it easier from the proton release and increases the acid strength.
5. (a) Both assertion and reason are true and reason is the correct explanation of assertion.



6. (b) Both assertion and reason are true but reason is not the correct explanation of assertion. Oxidation number can be calculated using some rules. H is assigned +1 oxidation state and O has oxidation number -2
- \therefore O. No. of C in CH_2O :
- O. no. of C + 2(+1) + (-2) = 0
- \therefore O. No. of C = 0
7. (a) Both assertion and reason are true and reason is the correct explanation of assertion.
8. (a) Both assertion and reason are true and reason is the correct explanation of assertion. Maximum oxidation state of S is +6, it cannot exceed it. Therefore it can't be further oxidised as S^{-2} can't be reduced further.
9. (a) Both assertion and reason are true and reason is the correct explanation of assertion.
- $$N_2^0 + 6e^- \longrightarrow 2N^{3-}$$
- \therefore equivalent weight of

