

# Ordinary Thinking

## Objective Questions

### Alkali metals

- As compared to potassium, sodium has [MP PMT 1985]
  - Lower electronegativity
  - Higher ionization potential
  - Greater atomic radius
  - Lower melting point
- Potassium is kept in [CPMT 1976]
  - Alcohol
  - Water
  - Kerosene
  - Liquid ammonia
- The product obtained on fusion of  $BaSO_4$  and  $Na_2CO_3$  is [AFMC 2005]
  - $BaCO_3$
  - $BaO$
  - $Ba(OH)_2$
  - $BaHSO_4$
- Which of the following statement is correct regarding alkali metals [NCERT 1981]
  - Cation is less stable than the atom
  - Cation is smaller than the atom
  - Size of cation and atom is the same
  - Cation is greater in size than the atom
- Valency electrons in alkali metals are [CPMT 1972]
  - 1
  - 7
  - 4
  - 2
- Magnitude of which of the following property of alkali metals increases with the increase of atomic number [MP PMT 1987]
  - Electronegativity
  - Ionic radius
  - First ionization energy
  - Melting point
- As compared to lithium, sodium reacts quickly with water because [NCERT 1978, 80]
  - Its molecular weight is less
  - It is stronger electronegative
  - It is stronger electropositive
  - It is a metal
- Which is an ore of potassium [DPMT 1984; CPMT 1986; Kurukshetra CEE 1998]
  - Carnellite
  - Cryolite
  - Bauxite
  - Dolomite
- $Na_2CO_3$  can be manufactured by Solvey's process but  $K_2CO_3$  cannot be prepared because [MP PMT 1993]
  - $K_2CO_3$  is more soluble
  - $K_2CO_3$  is less soluble
  - $KHCO_3$  is more soluble than  $NaHCO_3$
  - $KHCO_3$  is less soluble than  $NaHCO_3$
- Which of the following alkali metals is smallest in size [CPMT 1990]
  - Rb
  - K
  - Na
  - Li
- When potassium dichromate crystal are heated with conc.  $HCl$ 
  - $O_2$  is evolved
  - Chromyl chloride vapours are evolved
  - $Cl_2$  is evolved
  - No reaction takes place
- Which of the following does not illustrate the anomalous properties of lithium [MP PET 1993]
  - The melting point and boiling point of  $Li$  are comparatively high
  - $Li$  is much softer than the other group I metals
  - $Li$  forms a nitride  $Li_3N$  unlike group I metals
  - The ion of  $Li$  and its compounds are more heavily hydrated than those of the rest of the group
- Correct order of increasing activity is
  - $Cu, Mg, Na$
  - $Na, Mg, Cu$
  - $Mg, Na, Cu$
  - $Cu, Na, Mg$
- On heating anhydrous  $Na_2CO_3$ , ..... is evolved [CPMT 1971, 79]
  - $CO_2$
  - Water vapour
  - $CO$
  - No gas
- Chile saltpetre is [DPMT 1984; CPMT 1986, 89; CET Pune 1998; MP PMT 2003]
  - $NaNO_3$
  - $Na_2SO_4$
  - $KNO_3$
  - $Na_2SO_3$
- A mixture of  $KCl$  and  $KF$  is added to sodium chloride
  - To increase the conductivity of  $NaCl$
  - To decrease the melting point of  $NaCl$
  - To suppress the degree of dissociation of  $NaCl$
  - To decrease the volatility of  $NaCl$
- A well known reagent which contains copper sulphate, sodium potassium tartrate and sodium hydroxide is
  - Fenton's reagent
  - Schiff's reagent
  - Fehling's solution
  - Nessler's reagent
- Sodium metal can be stored under [CPMT 1972, 85; BHU 1983]
  - Benzene
  - Kerosene
  - Alcohol
  - Toluene



## 756 s and p-Block Elements

19. The most dangerous method of preparing hydrogen would be by the action of  $HCl$  and [JIPMER 2000]  
 (a)  $Al$  (b)  $K$   
 (c)  $Fe$  (d)  $Zn$
20. Based on lattice energy and other considerations which one of the following alkali metal chlorides is expected to have the highest melting point [AIIEE 2005]  
 (a)  $LiCl$  (b)  $NaCl$   
 (c)  $KCl$  (d)  $RbCl$
21. The correct formula of hypo is  
 (a)  $Na_2S_2O_3 \cdot 5H_2O$  (b)  $Na_2SO_4$   
 (c)  $Na_2S_2O_3 \cdot 4H_2O$  (d)  $Na_2S_2O_3 \cdot 3H_2O$
22. The reagent commonly used to determine hardness of water titrimetrically is [AIIMS 2003]  
 (a) Oxalic acid  
 (b) Disodium salt of  $EDTA$   
 (c) Sodium citrate  
 (d) Sodium thiosulphate
23.  $K_2CS_3$  can be called potassium [CPMT 1972, 74]  
 (a) Thiocyanate (b) Thiocarbonate  
 (c) Thiocarbide (d) Sulphocyanide
24. Which is most basic in character [BHU 1982]  
 (a)  $RbOH$  (b)  $KOH$   
 (c)  $NaOH$  (d)  $LiOH$
25. When washing soda is heated [AFMC 2005]  
 (a)  $CO$  is released  
 (b)  $CO + CO_2$  is released  
 (c)  $CO_2$  is released  
 (d) Water vapour is released
26. Which of the following is correct [CPMT 1971]  
 (a) All carbonates are soluble in water  
 (b) Carbonates of  $Na$ ,  $K$  and  $NH_4$  are soluble in water  
 (c) Carbonates of  $Ca$ ,  $Sr$ ,  $Ba$  are soluble in water  
 (d) All carbonates are insoluble
27. Nitre is [CPMT 1986]  
 (a)  $AgNO_3$  (b)  $KNO_3$   
 (c)  $NH_4NO_3$  (d)  $NaNO_3$
28. Nelson cell is used for the preparation of [CPMT 1985]  
 (a) Slaked lime (b) Baryta  
 (c) Sodium (d) Caustic soda
29. Potash alum is a [CPMT 1986; MNR 1981]  
 (a) Complex salt (b) Acid salt  
 (c) Double salt (d) Normal salt
30. The process of industrial manufacturing of sodium carbonate is known as [CPMT 1978, 86; MP PMT 1995]  
 (a) Castner process (b) Haber's process  
 (c) Le-blanc process (d) Chamber process
31. The colour of hydrogen is [IIT 1980]  
 (a) Black (b) Yellow  
 (c) Orange (d) None of these
32. Which one of the following salts gives aqueous solution which is weakly basic [Bihar CEE 1995]  
 (a)  $NaHCO_3$  (b)  $NaHSO_4$   
 (c)  $NaCl$  (d)  $NH_4HCO_3$
33. An example for a double salt is [KCET 2002]  
 (a) Silver nitrate (b) Mohr's salt  
 (c) Potassium ferricyanide (d) Cupromonium sulphate
34. The elements of group IA provide a colour to the flame of Bunsen burner due to [AIIMS 1987]  
 (a) Low ionization potential  
 (b) Low melting point  
 (c) Softness  
 (d) Presence of one electron in the outermost orbit
35. Which of the following is the smallest cation [MP PMT 1993]  
 (a)  $Na^+$  (b)  $Mg^{+2}$   
 (c)  $Ca^{+2}$  (d)  $Al^{+3}$
36.  $K$ ,  $Ca$  and  $Li$  metals may be arranged in the decreasing order of their standard electrode potentials as [CPMT 1990]  
 (a)  $K$ ,  $Ca$ ,  $Li$  (b)  $Li$ ,  $K$ ,  $Ca$   
 (c)  $Li$ ,  $Ca$ ,  $K$  (d)  $Ca$ ,  $Li$ ,  $K$
37. Alkali metals lose electrons in [CBSE PMT 1990]  
 (a) s-orbitals (b) p-orbitals  
 (c) d-orbitals (d) f-orbitals
38. The alkali metal that reacts with nitrogen directly to form nitride is [Roorkee 1992; MP PMT 2000; BHU 2000]  
 (a)  $Li$  (b)  $Na$   
 (c)  $K$  (d)  $Rb$
39. Which of the following has density greater than water [MP PET 1994]  
 (a)  $Li$  (b)  $Na$   
 (c)  $K$  (d)  $Rb$
40. The reactivity of the alkali metal sodium with water, is made use of [MP PMT 1994]  
 (a) In drying of alcohols  
 (b) In drying of benzene  
 (c) In drying of ammonia solution  
 (d) As a general drying agent
41. Which of the following has smaller size [RPET 2003]  
 (a)  $H$  (b)  $He^+$   
 (c)  ${}_1H^2$  (d)  $Li^{2+}$

42.  $KF$  combines with  $HF$  to form  $KHF_2$ . The compound contains the species [IIT 1996]  
 (a)  $K^+$ ,  $F^-$  and  $H^+$  (b)  $K^+$ ,  $F^-$  and  $HF$   
 (c)  $K^+$  and  $[HF_2]^-$  (d)  $[KHF]^+$  and  $F^-$
43. Which alkali metal is most metallic in character [MH CET 2001]  
 (a)  $K$  (b)  $Cs$   
 (c)  $Na$  (d)  $Li$
44. The property of hydrogen which distinguishes it from other alkali metals is [MP PET 1996]  
 (a) Its electropositive character  
 (b) Its affinity for non-metals  
 (c) Its reducing character  
 (d) Its non-metallic character
45. Which of the following reacts with water with high rate [AFMC 1995]  
 (a)  $Li$  (b)  $K$   
 (c)  $Na$  (d)  $Rb$
46. The valence shell electronic configuration of alkali metals is [MP PET 1996; UPSEAT 2001]  
 (a)  $ns^2np^1$  (b)  $ns^1$   
 (c)  $(n-1)p^6ns^2$  (d)  $(n-1)d^2ns^2$
47. Alkali metals are [MP PMT 1996]  
 (a)  $Li, Na, Be, Mg, Cs$  (b)  $Li, Na, K, Rb, Cs$   
 (c)  $Na, K, Mg, Ca, Rb$  (d)  $K, Rb, Cs, Ba, Sr$
48. The atomic number of an element is 11. Its oxide will be [MP PMT 1996]  
 (a) Acidic (b) Basic  
 (c) Acid and basic both (d) Neutral
49. The commercial production of sodium carbonate is done by [CPMT 1982; MP PMT 1996]  
 (a) Lead-chamber process  
 (b) Haber's process  
 (c) Solvay's process  
 (d) Castner's process
50. Alkali metals are strong reducing because  
 (a) These are monovalent  
 (b) Their ionisation potential are very high  
 (c) Their standard electrode potential are very much negative  
 (d) These are metals
51. Which of the following statement about  $LiCl$  and  $NaCl$  is correct [Kurukshetra CEE 2002]  
 (a)  $LiCl$  has higher melting point than  $NaCl$   
 (b)  $LiCl$  dissolves in water whereas  $NaCl$  does not  
 (c)  $LiCl$  would ionize in water more than  $NaCl$   
 (d) Fused  $LiCl$  would be less conducting than fused  $NaCl$
52. In the Castner's process for the extraction of sodium, the anode is made of.....metal. [EAMCET 2003]  
 (a) Copper (b) Iron  
 (c) Sodium (d) Nickel
53. Which of the following s-block elements forms nitride [RPET 2003]  
 (a)  $Ba$  (b)  $Be$   
 (c)  $Ca$  (d)  $Li$
54. Tincal is [Pb. PMT 2001]  
 (a)  $Na_2CO_3 \cdot 10H_2O$  (b)  $NaNO_3$   
 (c)  $NaCl$  (d)  $Na_2B_4O_7 \cdot 10H_2O$
55. Which has minimum solubility [BHU 2003]  
 (a)  $Br_2S_3$  (b)  $Ag_2S$   
 (c)  $CoS$  (d)  $PbS$
56. Cryolite helps in [BHU 2003]  
 (a) Lowering the melting point  
 (b) Increasing the melting point  
 (c) Increasing the electrical conductivity  
 (d) Decreasing the electrical conductivity
57. In certain matters lithium differs from other alkali metals, the main reason for this is [MP PET/PMT 1996]  
 (a) Small size of  $Li$  atom and  $Li^+$  ion  
 (b) Extremely high electropositivity of  $Li$   
 (c) Greater hardness of  $Li$   
 (d) Hydration of  $Li^+$  ion
58. Acidified potassium permanganate solution is decolourised by  
 (a) Bleaching powder (b) Microcosmic salt  
 (c) Mohr salt (d) White vitriol
59. Which one of the following is used as a disinfectant in water treatment [NDA 1999]  
 (a) Alum (b) Charcoal  
 (c) Kieselguhr (d) Potassium permanganate
60. Sodium thiosulphate is used in photography [UPSEAT 1999]  
 (a) To convert metallic silver into silver salt  
 (b)  $AgBr$  grain is reduced to non-metallic silver  
 (c) To remove reduced silver  
 (d) To remove undecomposed  $AgBr$  in the form of  $Na_3[Ag[S_2O_3)_2]$  (a complex salt)
61. Composition of borax is [UPSEAT 2001;04]  
 (a)  $Na_2B_4O_7 \cdot 4H_2O$  (b)  $Na_2B_4O_7 \cdot 10H_2O$



## 758 s and p-Block Elements

- (c)  $NaBO_2$  (d)  $Na_2BO_3$
62. When sodium dicarbonate is heated strongly for calcined in a kiln, it forms [CPMT 2000; KCET (Med.) 2000]  
 (a)  $Na$  (b)  $Na_2CO_3$   
 (c)  $NaCO_3$  (d)  $NaHCO_3$
63. The strongest reducing agent is [MP PET 2001]  
 (a)  $K$  (b)  $Al$   
 (c)  $Mg$  (d)  $Br$
64. The word 'alkali' is used for alkali metals indicates [RPMT 1999]  
 (a) Ash of the plants (b) Metallic nature  
 (c) Silvery lusture (d) Active metal
65. Potassium nitrate is called [RPMT 1999]  
 (a) Mohr's salt (b) Gypsum  
 (c) Indian salt petre (d) Chile salt petre
66. Which of the following chemicals, in addition to water, are used for the manufacture of  $Na_2CO_3$  by Solvay process [Roorkee 1999]  
 (a)  $NaCl, CO$  and  $NH_3$   
 (b)  $NaCl, CO_2$  and  $NH_3$   
 (c)  $NaCl, NH_4Cl$  and  $CO_2$   
 (d)  $NaHCO_3, CO$  and  $NH_3$
67. Which metal forms amide with  $NH_3$  at  $300^\circ C$  [CPMT 1994]  
 (a)  $Mg$  (b)  $Pb$   
 (c)  $Al$  (d)  $Na$
68. When sodium is heated with moist air, then the product obtained is [AIIMS 1999]  
 (a)  $Na_2O$  (b)  $NaOH$   
 (c)  $Na_2CO_3$  (d)  $Na_2O_2$
69. An inorganic compound first melts then resolidifies and then liberates a gas. It may be [DPMT 2002]  
 (a)  $MnO_2$  (b)  $Al_2O_3$   
 (c)  $KMnO_4$  (d)  $KClO_3$
70. On dissolving moderate amount of sodium metal in liquid  $NH_3$  at low temperature, which one of the following does not occur [AIIMS 2003]  
 (a) Blue coloured solution is obtained  
 (b)  $Na^+$  ions are formed in the solution  
 (c) Liquid  $NH_3$  becomes good conductor of electricity  
 (d) Liquid ammonia remains diamagnetic
71. The solubility of the alkali metal carbonates [Pune CET 1998]  
 (a) Increases at first and then decreases  
 (b) Does not show regular variation  
 (c) Increases as we go down the group  
 (d) Decreases as we go down the group
72. Which of the following properties is not true for an alkali metal [Pune CET 1998]  
 (a) Low atomic volume  
 (b) Low ionization energy  
 (c) Low density  
 (d) Low electronegativity
73. Which of the following alkali metals has the biggest tendency for the half reaction,  $M(g) \rightarrow M^+(aq) + e^-$  [DPMT 2001]  
 (a) Lithium (b) Sodium  
 (c) Cesium (d) Potassium
74. Which one of the following metallic hydroxides does not dissolve in sodium hydroxide solution [KCET (Me)]  
 (a)  $Zn(OH)_2$  (b)  $Al(OH)_3$   
 (c)  $Fe(OH)_3$  (d)  $Pb(OH)_2$
75. Which one of the following on heating will not give  $CO_2$  [NDA 1999; BHU 2000]  
 (a)  $CaCO_3$  (b)  $Na_2CO_3$   
 (c)  $PbCO_3$  (d)  $Li_2CO_3$
76.  $NaOH$  is prepared by the method [AFMC 2005]  
 (a) Down's cell (b) Castner cell  
 (c) Solvay process (d) Castner Kellner cell
77. Sodium gives blue colour with  $NH_3$  solution, this blue colour is due to [UPSEAT 2000,02; AMU 2002; RPMT 2002]  
 (a) Ammoniated  $Na^\oplus$  (b) Ammoniated  $Na^\ominus$   
 (c) Ammoniated  $e^-$  (d)  $Na^+ / Na^-$  pair
78. The strongest reducing agent of the alkali metal is [CPMT 1999; Pb.CET 2001]  
 (a)  $Li$  (b)  $Na$   
 (c)  $K$  (d)  $Cs$
79. With the increase in atomic weights, melting points of the alkali metals [MP PMT 1995]  
 (a) Increase  
 (b) Decrease  
 (c) Remain constant  
 (d) Do not show definite trend
80. The reaction of water with sodium and potassium is [BHU 1999]  
 (a) Exothermic  
 (b) Endothermic  
 (c) Reversible  
 (d) Irreversible and endothermic

81. When potassium ferrocyanide crystals are heated with concentrated sulphuric acid, the gas evolved is  
[CBSE PMT PMT 1999; KCET 2000]  
(a) Ammonia (b) Sulphur dioxide  
(c) Carbon dioxide (d) Carbon monoxide
82. Characteristic feature of alkali metals is  
[RPMT 2000; MP PMT 2004]  
(a) Good conductor of heat and electricity  
(b) High melting points  
(c) Low oxidation potentials  
(d) High ionization potentials
83. A substance  $X$  is a compound of an element of group IA the substance  $X$  gives a violet colour in flame test,  $X$  is  
[MP PMT 1980, 85, 86; CPMT 1985; DCE 2000]  
(a)  $LiCl$  (b)  $NaCl$   
(c)  $KCl$  (d) None
84. Which of the following alkali metal ions has lowest ionic mobility in aqueous solution [KCET 2000]  
(a)  $Rb^+$  (b)  $Cs^+$   
(c)  $Li^+$  (d)  $Na^+$
85. Lithium shows similarities to magnesium in its chemical behaviour because [Pb. PMT 2000]  
(a) Similar size, greater electronegativity and similar polarizing power.  
(b) Similar size same electronegativity and lower polarizing power  
(c) Similar size, same electronegativity and similar high polarizing power  
(d) None of these
86. Which one of the following is the most electropositive element [Pb. PMT 2000]  
(a) Calcium (b) Chlorine  
(c) Potassium (d) Carbon
87. Electrolysis of molten sodium chloride leads to the formation of [KCET 1990]  
(a)  $Na$  and  $H_2$  (b)  $Na$  and  $O_2$   
(c)  $H_2$  and  $O_2$  (d)  $Na$  and  $Cl_2$
88. When sodium bicarbonate is heated the product obtained is  
[Pb. CET 2000; DCE 2004]  
(a)  $Na$  (b)  $Na_2CO_3$   
(c)  $NaCO_3$  (d)  $Na_2(HCO_3)$
89. Which of the following is a use of alum [CPMT 2004]  
(a) Making explosives (b) Bleaching clothes  
(c) Water softening (d) All of the above
90. Which of the following salt does not get hydrolysed in water  
[CPMT 2004]  
(a)  $KClO_4$  (b)  $NH_4Cl$   
(c)  $CH_3COONa$  (d) None of these
91. A fire of lithium, sodium and potassium can be extinguished by [DCE 2003]  
(a)  $H_2O$  (b) Nitrogen  
(c)  $CO_2$  (d) Asbestos blanket
92. Which of the following metal has stable carbonates  
[AFMC 2004]  
(a)  $Na$  (b)  $Mg$   
(c)  $Al$  (d)  $Si$
93. Aluminium reacts with caustic soda to form [DCE 2004]  
(a) Aluminium hydroxide  
(b) Aluminium oxide  
(c) Sodium meta-aluminate  
(d) Sodium tetra aluminate
94. Alkaline earth metals are denser than alkali metals, because metallic bonding in alkaline earth's metal, is [BHU 2004]  
(a) Stronger (b) Weaker  
(c) Volatile (d) Not present
95. Which of the following is a false statement [CPMT 2004]  
(a) Fluorine is more electronegative than chlorine  
(b) Nitrogen has greater  $IE_1$  than oxygen  
(c) Lithium is amphoteric  
(d) Chlorine is an oxidising agent
96. Which is most basic in character [UPSEAT 2004]  
(a)  $CsOH$  (b)  $KOH$   
(c)  $NaOH$  (d)  $LiOH$
97. Photoelectric effect is maximum in [AFMC 2004]  
(a)  $Cs$  (b)  $Na$   
(c)  $K$  (d)  $Li$
98. A metal  $M$  reacts with  $N_2$  to give a compound ' $A$ ' ( $M_3N$ ). ' $A$ ' on heating at high temperature gives back ' $M$ ' and ' $A$ ' on reacting with  $H_2O$  gives a gas ' $B$ '. ' $B$ ' turns  $CuSO_4$  solution blue on passing through it.  $A$  and  $B$  can be [DCE 2003]  
(a)  $Al$  and  $NH_3$  (b)  $Li$  and  $NH_3$   
(c)  $Na$  and  $NH_3$  (d)  $Mg$  and  $NH_3$
99. A solid compound ' $X$ ' on heating gives  $CO_2$  gas and a residue. The residue mixed with water forms ' $Y$ '. On passing an excess of  $CO_2$  through ' $Y$ ' in water, a clear solution, ' $Z$ ' is obtained. On boiling ' $Z$ ', compound ' $X$ ' is reformed. The compound ' $X$ ' is [CBSE PMT 2004]  
(a)  $Na_2CO_3$  (b)  $K_2CO_3$   
(c)  $Ca(HCO_3)_2$  (d)  $CaCO_3$
100. Amongst  $LiCl$ ,  $RbCl$ ,  $BeCl_2$  and  $MgCl_2$  the compounds with the greatest and least ionic character respectively are



## 760 s and p-Block Elements

- [Pb. CET 2004]
- (a)  $LiCl$  and  $RbCl$  (b)  $MgCl_2$  and  $BeCl_2$   
 (c)  $RbCl$  and  $BeCl_2$  (d)  $RbCl$  and  $MgCl_2$
101. Salt cake is  
 (a) Sodium sulphate  
 (b) Sodium chloride  
 (c) Sodium bisulphite  
 (d) Sodium sulphate and Sodium chloride
102. Gobar salt is [BHU 1983; CPMT 1988, 91; IIT 1985; MP PET 2000]  
 (a)  $MgSO_4 \cdot 7H_2O$  (b)  $CuSO_4 \cdot 5H_2O$   
 (c)  $FeSO_4 \cdot 7H_2O$  (d)  $Na_2SO_4 \cdot 10H_2O$
103. The colour given to the flame by sodium salts is [CPMT 1980; MP PET 1986]  
 (a) Light red (b) Golden yellow  
 (c) Green (d) Pink
104. Solvay's process is used for the preparation of [CPMT 1982; AIIMS 1987]  
 (a) Ammonia (b) Sodium bicarbonate  
 (c) Sodium carbonate (d) Calcium carbonate
105. Sodium when heated in a current of dry ammonia gives [NCERT 1981; KCET 2000]  
 (a) Sodium nitrite (b) Sodium hydride  
 (c) Sodium amide (d) Sodium azide
106. Washing soda is [CPMT 1982; DPMT 1982; CBSE PMT 1990; MP PMT 1987, 96]  
 (a)  $Na_2CO_3 \cdot 10H_2O$  (b)  $Na_2CO_3 \cdot H_2O$   
 (c)  $Na_2CO_3 \cdot 5H_2O$  (d)  $Na_2CO_3$
107. The substance used to decolourise and purify oils is [MP PMT 1987]  
 (a) Sodium carbonate (b) Sodium chloride  
 (c) Sodium hydroxide (d) Sodium sulphate
108. The main salt soluble in sea water is [MP PMT 1998]  
 (a)  $MgCl_2$  (b)  $NaCl$   
 (c)  $MgSO_4$  (d)  $CaSO_4$
109. The metallic lustre exhibited by sodium is explained by [IIT 1987]  
 (a) Diffusion of sodium ions  
 (b) Oscillation of loose electrons  
 (c) Excitation of free protons  
 (d) Existence of body centred cubic lattice
110. The metal which reacts with water at room temperature is [CPMT 1985; MP PMT 1996; MP PET 1998]  
 (a) Copper (b) Iron  
 (c) Magnesium (d) Sodium
111. When  $NaCl$  is dissolved in water, the sodium ion is [CPMT 1989]  
 (a) Oxidised (b) Reduced  
 (c) Hydrolysed (d) Hydrated
112. Sodium metal cannot be stored under [CPMT 1985, 88, 94]  
 (a) Benzene (b) Kerosene  
 (c) Alcohol (d) Toluene
113. Causticization process is used for the preparation of [CPMT 1985; BHU 1986]  
 (a) Caustic soda (b) Caustic potash  
 (c) Baryta (d) Slaked lime
114. When  $CO$  is passed over solid  $NaOH$  heated to  $200^\circ C$ , it forms [MP PMT 1985]  
 (a)  $Na_2CO_3$  (b)  $NaHCO_3$   
 (c)  $HCOONa$  (d) None
115.  $NaOH$  is manufactured by electrolysis of brine solution. The products of the reaction are [KCET 1990]  
 (a)  $Cl_2$  and  $H_2$  (b)  $Cl_2$  and  $Na - Hg$   
 (c)  $Cl_2$  and  $Na$  (d)  $Cl_2$  and  $O_2$
116. Sodium carbonate is manufactured by Solvay process, the products that are recycled are [KCET 1993; DCIT 1993]  
 (a)  $CO_2$  and  $NH_3$  (b)  $CO_2$  and  $NH_4Cl$   
 (c)  $NaCl, CaO$  (d)  $CaCl_2, CaO$
117. The useful by-products, obtained in the Solvay process of manufacturing sodium carbonate, are [KCET 1988]  
 (a) Quick lime and  $CO_2$   
 (b)  $NaHCO_3$  and  $NH_4Cl$   
 (c)  $NH_4Cl$  solution and quick lime  
 (d)  $NaHCO_3$  and  $CO_2$
118. In the preparation of sodium carbonate, which of the following is used [AFMC 1992]  
 (a) Slaked lime (b) Quick lime  
 (c) Lime stone (d)  $NaOH$
119. When  $NaOH$  crystals are left in open air, they acquire a fluid layer around each crystal as [CPMT 1974]  
 (a) They start melting  
 (b) They absorb moisture from air  
 (c) They react with air to form a liquid compound  
 (d) They absorb  $CO_2$  from air
120. Sodium carbonate reacts with  $SO_2$  in aqueous medium to give [MP PMT 1982, 85]  
 (a)  $NaHSO_3$  (b)  $Na_2SO_3$   
 (c)  $NaHSO_4$  (d)  $Na_2SO_4$

- 121.** Baking soda is [CPMT 1974, 78, 79, 91; BHU 1979; Manipal MEE 1995; AIIMS 1996; CPMT 1973; RPET 1999; AFMC 2001, 05; Pb. CET 2002]  
 (a)  $Na_2CO_3$  (b)  $NaHCO_3$   
 (c)  $Na_2SO_4$  (d)  $K_2CO_3$
- 122.** Soda ash is [KCET 1993]  
 (a)  $Na_2CO_3 \cdot H_2O$  (b)  $NaOH$   
 (c)  $Na_2CO_3$  (d)  $NaHCO_3$
- 123.** Soda lime is [KCET 1993]  
 (a)  $NaOH$  (b)  $CaO$   
 (c)  $NaOH$  and  $CaO$  (d)  $Na_2CO_3$
- 124.** Molten sodium is used in nuclear reactors to [KCET 1989]  
 (a) Absorb neutrons in order to control the chain reaction  
 (b) Slow down the fast neutrons  
 (c) Absorb the heat generated by nuclear fission  
 (d) Extract radio-isotopes produced in the reactor
- 125.** Squashes are stored by adding [AFMC 1989]  
 (a) Citric acid (b)  $KCl$   
 (c)  $Na_2SO_3$  (d) Sodium metabisulphite
- 126.** Sodium thiosulphate ( $Na_2S_2O_3 \cdot 5H_2O$ ) is used in photography to [CPMT 1972, 74, 79; DPMT 1983; Bihar CEE 1995; MNR 1995]  
 (a) Reduce silver bromide to metallic silver  
 (b) Convert metallic silver to silver salt  
 (c) Remove undecomposed  $AgBr$  as a soluble silver thiosulphate complex  
 (d) Remove unreduced silver
- 127.** Which of the following pair can't exist in solution [IIT 1986; DCE 1999]  
 (a)  $NaHCO_3$  and  $NaOH$  (b)  $Na_2CO_3$  and  $NaOH$   
 (c)  $Na_2CO_3$  and  $NaCl$  (d)  $NaHCO_3$  and  $NaCl$
- 128.** Sodium thiosulphate is prepared by [IIT 1996]  
 (a) Reducing  $Na_2SO_4$  solution with  $H_2S$   
 (b) Boiling  $Na_2SO_3$  solution with  $S$  in alkaline medium  
 (c) Neutralising  $H_2S_2O_3$  solution with  $NaOH$   
 (d) Boiling  $Na_2SO_3$  solution with  $S$  in acidic medium
- 129.** When  $NaOH$  is prepared, the gas released is [CPMT 1996]  
 (a)  $Cl_2$  (b)  $H_2$   
 (c)  $O_2$  (d)  $H_2O$
- 130.** What is lye [BHU 1997]  
 (a) 10% solution of  $NaCl$  (b) 10% solution of  $KOH$   
 (c) 10% solution of  $Ca(OH)_2$   
 (d) 10% solution of  $Na_2CO_3$
- 131.**  $Na$  imparts yellow colour to Bunsen flame because of [RPMT 1997]  
 (a) Low ionisation potential  
 (b) Sensitivity  
 (c) Sublimation  
 (d) Absorbed high radiation
- 132.**  $Sn$  is dissolved in excess of  $NaOH$  solution, the compound obtained is [RPMT 1997]  
 (a)  $Sn(OH)_2$  (b)  $Na_2SnO_3$   
 (c)  $Na_2SnO_2$  (d)  $SnO_2$
- 133.** Identify the correct statement [CPMT 1997]  
 (a) Elemental sodium can be prepared and isolated by electrolysis of an aqueous solution of sodium chloride  
 (b) Elemental sodium is a strong oxidizing agent  
 (c) Elemental sodium is insoluble in ammonia  
 (d) Elemental sodium is easily oxidized
- 134.** Calcium is obtained by [CBSE PMT 1997]  
 (a) Roasting of limestone  
 (b) Electrolysis of solution of calcium chloride in  $H_2O$   
 (c) Reduction of calcium chloride with carbon  
 (d) Electrolysis of molten anhydrous calcium chloride
- 135.** When sodium chloride solution is electrolysed, the gas that is liberated at the cathode is [Kurukshetra CEE]  
 (a) Oxygen (b) Hydrogen  
 (c) Chlorine (d) Air
- 136.** During the electrolysis of fused sodium chloride, the anodic reaction is [KCET 1998]  
 (a) Reduction of sodium ions  
 (b) Oxidation of sodium ions  
 (c) Reduction of chloride ions  
 (d) Oxidation of chloride ions
- 137.** Which of the following does not participate in the Solvay's process for the manufacture of  $Na_2CO_3$  [EAMCET]  
 (a)  $NH_3$  (b)  $NaCl$  solution  
 (c)  $CO_2$  (d)  $H_2SO_4$
- 138.** The colour of the precipitate produced by adding  $NaOH$  solution to  $HgCl_2$  is [KCET 1998]  
 (a) Yellow (b) Black  
 (c) Brown (d) White
- 139.** The cell used for the electrolysis of fused  $NaCl$  is [AFMC 1999; Kerala (Mea.) 2002]  
 (a) Down's cell (b) Castner cell

## 762 s and p-Block Elements

- (c) Solvay cell (d) Nelson cell
140. Slaked lime [ $Ca(OH)_2$ ] is used in the manufacture  
[UPSEAT 2000]  
(a) Cement (b) Fire bricks  
(c) Pigment (d) Medicine
141. The alum used for purifying water is [KCET (Med.) 2001]  
(a) Ferric alum (b) Chrome alum  
(c) Potash alum (d) Ammonium alum
142. Which one of the following metallic hydroxides does not dissolve in sodium hydroxide solution [KCET (Med.) 2001]  
(a)  $Zn(OH)_2$  (b)  $Al(OH)_3$   
(c)  $Fe(OH)_3$  (d)  $Pb(OH)_2$
143. In which of the following processes, fused sodium hydroxide is electrolysed at a  $330^\circ C$  temperature for extraction of sodium [CBSE PMT 2000; AFMC 2001]  
(a) Castner's process (b) Down's process  
(c) Cyanide process (d) Both (b) and (c)
144. Excess of  $Na^+$  ions in our system causes  
[KCET (Med.) 2001]  
(a) High B.P. (b) Low B.P.  
(c) Diabetes (d) Anaemia
145. Ferric alum has the composition  
 $(NH_4)_2SO_4 \cdot Fe_2(SO_4)_3 \cdot xH_2O$  [Orissa JEE 2002]  
(a) 7 (b) 24  
(c) 6 (d) 15
146. If  $Na$  is heated in presence of air, it forms [AFMC 2002]  
(a)  $Na_2CO_3$  (b)  $Na_2O_2$   
(c)  $Na_2O$  (d) Both (b) and (c)
147. Which of the following is most reducing agent [RPMT 2002]  
(a)  $HNO_3$  (b)  $Na$   
(c)  $Cl_2$  (d)  $Cr$
148. Pyrolusite is [DPMT 2002]  
(a) Carbonate ore (b) Sulphur ore  
(c) Silicon ore (d) None of these
149. In the manufacture of metallic sodium by the fused salt electrolysis (Down's process) a small amount of calcium chloride is added to [MP PET 1993; MP PMT 1994]  
(a) Improve the electrical conduction  
(b) Increase the temperature of electrolysis  
(c) Bring down the melt temperature  
(d) Stabilize the metallic sodium
150. Sodium metal is extracted by [MP PMT 1996]  
(a) Electrolysis of aqueous solution of sodium chloride  
(b) Electrolysis of fused sodium chloride  
(c) Heating sodium oxide with carbon  
(d) Heating sodium oxide with hydrogen
1. To remove last traces of water from alcohol, the metal used is  
(a) Sodium (b) Potassium  
(c) Calcium (d) Aluminium
2. Plaster of Paris is  
[CPMT 1972, 76, 78, 83, 87, 88, 90, 91, 93, 94; JIPMER 2002; MP PET 1986, 2001; BHU 1992, 95, 2000; MNR 1982; DCE 2000; Manipal MEE 1995; NCERT 1976; Bihar MEE 1997; EAMCET 1978; AMU 1982, 84; DPMT 1982, 83]  
(a)  $CaSO_4 \cdot 2H_2O$  (b)  $CaSO_4 \cdot 3H_2O$   
(c)  $CaSO_4 \cdot H_2O$  (d)  $CaSO_4 \cdot \frac{1}{2}H_2O$
3. Which of the following substance is used as dehydrating agent in laboratory [MP PMT 1987]  
(a) Calcium chloride (b) Sodium chloride  
(c) Sodium carbonate (d) Potassium nitrate
4. The metal that is extracted from sea water is  
[EAMCET 1978; CPMT 1988; CET Pune 1998; MP PET 2000]  
(a)  $Ba$  (b)  $Mg$   
(c)  $Ca$  (d)  $Sr$
5. Which of the following ore contains both magnesium and calcium [MDAT Bihar 1984; MP PET 2003]  
(a) Magnesite (b) Dolomite  
(c) Carnellite (d) Phosphorite
6. Epsom salt is  
[EAMCET 1978, 80; BHU 1979; MP PET 1999; CPMT 1988, 89, 90; Bihar MEE 1996]  
(a)  $CaSO_4 \cdot 2H_2O$  (b)  $BaSO_4 \cdot 2H_2O$   
(c)  $MgSO_4 \cdot 2H_2O$  (d)  $MgSO_4 \cdot 7H_2O$
- Setting of plaster of paris is [MP PMT 1985; CPMT 1989]  
(a) Oxidation with atmospheric oxygen  
(b) Combination with atmospheric  $CO_2$   
(c) Dehydration  
(d) Hydration to yield another hydrate
8. To prevent magnesium from oxidation in electrolytic extraction process  
(a) Some calcium fluoride is added  
(b) Some chlorides are added  
(c) Metal is taken out by spoons  
(d) The whole process is done in an atmosphere of coal gas
9. Which of the following metal is found in green colouring pigment chlorophyll of plants  
[KCET 1993; RPMT 1999; MP PET 2002]  
(a)  $Fe$  (b)  $Mg$   
(c)  $Na$  (d)  $Al$
10. Which of the following metal carbonate is decomposed on heating [MNR 1985; MP PET 1994; Pb. CET 2002]  
(a)  $MgCO_3$  (b)  $Na_2CO_3$   
(c)  $K_2CO_3$  (d)  $Rb_2CO_3$

## Alkaline earth metals



11. The outer electronic configuration of alkaline earth metal is [NCERT 1982]  
[BHU 1980; CPMT 1985, 93; MP PAT 1993]  
(a)  $ns^2$  (b)  $ns^1$   
(c)  $np^6$  (d)  $nd^{10}$
12. Metallic magnesium is prepared by [BHU 1973, 77]  
(a) Reduction of  $MgO$  by coke  
(b) Electrolysis of aqueous solution of  $Mg(NO_3)_2$   
(c) Displacement of  $Mg$  by iron from  $MgSO_4$  solution  
(d) Electrolysis of molten  $MgCl_2$
13. Of the metals  $Be$ ,  $Mg$ ,  $Ca$  and  $Sr$  of group II A. In the periodic table the least ionic chloride would be formed by [NCERT 1980; CPMT 1980]  
(a)  $Be$  (b)  $Mg$   
(c)  $Ca$  (d)  $Sr$
14. Which one of the following is fluorspar  
(a)  $CaF_2$  (b)  $CaO$   
(c)  $H_2F_2$  (d)  $CaCO_3$
15. Which one is known as barytes [CPMT 1987]  
(a)  $BaSO_4$  (b)  $BaCl_2 \cdot 2H_2O$   
(c)  $BaO$  (d)  $BaCO_3$
16. Which of the following sulphates have the highest solubility in water [EAMCET 1980, 84, 85; MP PMT 1994; Kurukshetra CEE 1998; AFMC 1990; MP PET 1994]  
(a)  $MgSO_4$  (b)  $BaSO_4$   
(c)  $CaSO_4$  (d)  $BeSO_4$
17. The composition formulae of gypsum is [CPMT 1975, 78, 82; DPMT 1982; IIT 1978; MNR 1981; MP PMT 1996; RPMT 1997]  
(a)  $(CaSO_4)_2 \cdot H_2O$  (b)  $2CaSO_4$   
(c)  $CaSO_4 \cdot 2H_2O$  (d)  $2CaSO_4 \cdot H_2O$
18. Mortar is a mixture of [EAMCET 1998; AIIMS 2000]  
(a)  $CaCO_3$ , sand and water  
(b) Slaked lime and water  
(c) Slaked lime, sand and water  
(d)  $CaCO_3$  and  $CaO$
19. Gypsum  $CaSO_4 \cdot 2H_2O$  on heating to about  $120^\circ C$  forms a compound which has the chemical composition represented by [CPMT 1978, 82, 88, 90; EAMCET 1978; DPMT 1982, 83; NCERT 1979]  
(a)  $CaSO_4$  (b)  $2CaSO_4 \cdot H_2O$   
(c)  $CaSO_4 \cdot H_2O$  (d)  $2CaSO_4 \cdot 3H_2O$
20. The highly efficient method of obtaining beryllium is  
(a) Dissociation of beryllium carbide  
(b) Electrolysis of fused beryllium chloride  
(c) Reduction of beryllium oxide with carbon  
(d) Reduction of beryllium halide with magnesium
21. Mark the incorrect statement  
(a) Lithopone is cheap and possess good covering power  
(b) Lithopone is yellow pigment  
(c) Lithopone is prepared by mixing barium sulphide and zinc sulphate  
(d) Lithopone is a mixture of barium sulphate and zinc sulphate
22. Pure anhydrous  $MgCl_2$  can be prepared from the hydrated salt by [CPMT 1986; MP PMT 1989]  
(a) Heating the hydrate with coke  
(b) Heating the hydrate with  $Mg$  ribbon  
(c) Melting the hydrate  
(d) Heating the hydrate to red heat in an atmosphere of  $HCl$  gas
23. Bleaching powder is obtained by the interaction of chlorine and [CPMT 1972, 78, 89; 2002; DPMT 1983]  
(a) Conc. solution of  $Ca(OH)_2$   
(b) Dilute solution of  $Ca(OH)_2$   
(c) Dry calcium oxide  
(d) Dry slaked lime
24. Deep pink colour is given to flame by the salts of  
(a) Strontium (b) Potassium  
(c) Zinc (d) Barium
25. Calcium salts give which colour when put in a flame  
(a) Brick red (b) Green  
(c) White (d) Pink
26. Phosphine is obtained from the following ore [Roorkee 1995]  
(a) Calcium superphosphate (b) Calcium phosphide  
(c) Potassium phosphide (d) Calcium hypophosphite
27. Calcium is obtained by [DPMT 1980; IIT 1980; CPMT 1996; AIIMS 2001]  
(a) Roasting of lime stone  
(b) Reduction of  $CaCl_2$  with carbon  
(c) Electrolysis of a solution of  $CaCl_2$  in water  
(d) Electrolysis of molten  $CaCl_2$
28. Which element possesses biggest atomic radii  
(a)  $P$  (b)  $Si$   
(c)  $Al$  (d)  $Mg$
29. Magnesia is



## 764 s and p-Block Elements

- (a)  $MgCO_3$  (b)  $MgO$   
 (c)  $MgSO_4$  (d)  $MgCl_2$
30. Mixture of  $MgCl_2$  and  $MgO$  is called [DPMT 1984]  
 (a) Double salt (b) Sorrel cement  
 (c) Portland cement (d) None of these
31. Lithopone is [AFMC 1992; BHU 1983, 86, 95; JIPMER 1999; RPET/PMT 1999]  
 (a)  $BaO + ZnSO_4$  (b)  $ZnO + BaSO_4$   
 (c)  $BaS + ZnSO_4$  (d)  $ZnS + BaSO_4$
32. For bleaching powder, which is incorrect [EAMCET 1984; CPMT 1985]  
 (a) Reacts with dilute acid to release chlorine  
 (b) Oxidising agent  
 (c) Light yellow coloured powder  
 (d) Highly soluble in water
33. Bleaching powder is a compound having the molecular formula [CPMT 1986, 89, 90, 93; MP PMT 1996; BHU 2005]  
 (a)  $CaOCl_3$  (b)  $CaOCl_2$   
 (c)  $CaClO$  (d)  $CaClO_3$
34. Calcium cyanamide is [CPMT 1986, 93]  
 (a)  $CaCHNH_2$  (b)  $CaCN_2$   
 (c)  $CaC_2N_2$  (d)  $Ca(CN)_2$
35. Which one of the following is a true peroxide [RPET 1999; CPMT 1981; Roorkee 1995]  
 (a)  $SO_2$  (b)  $BaO_2$   
 (c)  $MnO_2$  (d)  $NO_2$
36. Which of the following is not a water absorber and dehydrating substance [CBSE 1989; JIPMER 2002]  
 (a) Silica gel (b)  $P_2O_5$   
 (c) Conc.  $H_2SO_4$  (d) Aqueous  $CaCl_2$
37. The dark red colour of bombs in fireworks is due to the presence of [Roorkee 1989; DPMT 2001; MP PMT 1985; AFMC 1989; Roorkee 1989]  
 (a)  $Na$  (b)  $Ba$   
 (c)  $Sr$  (d)  $K$
38. The most electropositive amongst the alkaline earth metals is [MP PMT 1993]  
 (a) Beryllium (b) Magnesium  
 (c) Calcium (d) Barium
39. Which of the following salts is insoluble in water at room temperature but soluble in boiling water [MP PMT 1993]  
 (a)  $CaCl_2$  (b)  $BaCl_2$   
 (c)  $SrCl_2$  (d)  $PbCl_2$
40. Electronegativity of beryllium is approximately equal to that of [MP PMT 1993]  
 (a) Aluminium (b) Boron  
 (c) Magnesium (d) Sodium
41. The right order of the solubility of sulphates of alkaline earth metals in water is [MP PET 1993; Pb. CET 2000; DPMT 2004]  
 (a)  $Be > Ca > Mg > Ba > Sr$   
 (b)  $Mg > Be > Ba > Ca > Sr$   
 (c)  $Be > Mg > Ca > Sr > Ba$   
 (d)  $Mg > Ca > Ba > Be > Sr$
42. Which of the following has highest electrode potential [CPMT 1990]  
 (a)  $Be$  (b)  $Mg$   
 (c)  $Ca$  (d)  $Ba$
43. The alkaline earth metals  $Ba$ ,  $Sr$ ,  $Ca$  and  $Mg$  may be arranged in the order of their decreasing first ionisation potential as  
 (a)  $Mg, Ca, Sr, Ba$ , (b)  $Ca, Sr, Ba, Mg$   
 (c)  $Sr, Ba, Mg, Ca$  (d)  $Ba, Mg, Ca, Sr$ ,
44. Which of the following alkaline earth metals shows some properties similar to aluminium [BHU 1983]  
 (a)  $Be$  (b)  $Ca$   
 (c)  $Sr$  (d)  $Ba$
45. Which of the following ions forms highly soluble hydroxide in water [CPMT 1974, 76, 79, 82]  
 (a)  $K^+$  (b)  $Zn^{++}$   
 (c)  $Al^{+++}$  (d)  $Ca^{++}$
46. Sodium sulphate is soluble in water whereas barium sulphate is sparingly soluble because [IITJEE 1989]  
 (a) The hydration energy of  $Na_2SO_4$  is less than its lattice energy  
 (b) The hydration energy of  $Na_2SO_4$  is more than its lattice energy  
 (c) The lattice energy of  $BaSO_4$  is more than its hydration energy  
 (d) The lattice energy has no role to play in solubility
47. Which one of the following is most basic [CPMT 1977, 83]  
 (a)  $Al_2O_3$  (b)  $MgO$   
 (c)  $SiO_2$  (d)  $P_2O_5$
48. Alloys of ..... metal are light and strong and so are used in the manufacture of aeroplane parts [EAMCET 1993]  
 (a)  $Cr$  (b)  $Sn$   
 (c)  $Fe$  (d)  $Mg$
49. In India at the occasion of marriages, the fire works used give green flame. Which one of the following radicals may be present [CPMT 1980; AFMC 1989; MP PET 2002]  
 (a)  $Na$  (b)  $K$

- (c) *Ba* (d) *Ca*
50.  $\text{CaCO}_3 \xrightarrow{\square} \text{CaO} + \text{CO}_2$  reaction in a line goes to completion because [AFMC 2005]  
 (a) *CaO* does not react to  $\text{CO}_2$  to give  $\text{CaCO}_3$   
 (b) Backward reaction is very slow  
 (c)  $\text{CO}_2$  formed escapes out  
 (d) None of these
51. The wire of flash bulb is made of [CPMT 1988]  
 (a) *Mg* (b) *Cu*  
 (c) *Ba* (d) *Ag*
52. Bone ash contains [KCET 1992]  
 (a) *CaO* (b)  $\text{CaSO}_4$   
 (c)  $\text{Ca}_3(\text{PO}_4)_2$  (d)  $\text{Ca}(\text{H}_2\text{PO}_4)_2$
53. A substance absorbs  $\text{CO}_2$  and violently reacts with water. That substance is  
 (a)  $\text{CaCO}_3$  (b) *CaO*  
 (c)  $\text{H}_2\text{SO}_4$  (d) *ZnO*
54. Setting of cement is an [DPMT 1984]  
 (a) Exothermic reaction  
 (b) Endothermic reaction  
 (c) Neither exothermic nor endothermic  
 (d) None of these
55. Which is quick lime [EAMCET 1993]  
 (a)  $\text{Ca}(\text{OH})_2$  (b) *CaO*  
 (c)  $\text{CaCO}_3$  (d)  $\text{Ca}(\text{OH})_2 + \text{H}_2\text{O}$
56. A major constituent of portland cement except lime is [CPMT 1982]  
 (a) Silica (b) Alumina  
 (c) Iron oxide (d) Magnesia
57. Portland cement is manufactured by using [CPMT 1986]  
 (a) Lime stone, clay and sand  
 (b) Lime stone, gypsum and sand  
 (c) Lime stone, gypsum and alumina  
 (d) Lime stone, clay and gypsum
58. Identify the correct statement [CBSE PMT 1995]  
 (a) Gypsum contains a lower percentage of plaster of calcium than plaster of paris  
 (b) Gypsum is obtained by heating plaster of paris  
 (c) Plaster of paris can be obtained by hydration of gypsum  
 (d) Plaster of paris is obtained by partial oxidation of gypsum
59. Which of the following decreases on going gradually from *Be* to *Ba* (in periodic table)  
 (a) Basic character of hydroxides  
 (b) Solubility of sulphates in water  
 (c) Solubility of hydroxides in water  
 (d) Strength of elements as reducing agent
60. Alkaline earth metals are [MP PMT 1996]  
 (a) *Li, Be, K, Mg, Ca* (b) *Be, Mg, Ca, Sr, Ba*  
 (c) *Be, K, Mg, Ca, Sr* (d) *Be, Mg, Ca, K, Rb*
61. Which of the following substances is used in the laboratory for fast drying of neutral gases [AIIMS 1998; AFMC 1998]  
 (a) Sodium phosphate  
 (b) Phosphorus pentoxide  
 (c) Sodium sulphate  
 (d) Anhydrous calcium chloride
62. Which of the following can be represented by the configuration  $[\text{Kr}]5s^2$ ? [MP PMT 1997]  
 (a) *Ca* (b) *Sr*  
 (c) *Ba* (d) *Ra*
63. Point out the incorrect statement regarding *Be* (Group-IIA) [MP PMT 1997]  
 (a) It forms an ionic carbide [AFMC 1988]  
 (b) Its carbonate decomposes on heating  
 (c) Its halides are covalent  
 (d) It is easily attacked by water
64. Beryllium differs from rest of the members of its family (Group-IIA) in many ways. The reason for this is its [MP PMT 1997]  
 (a) Small size and higher electronegativity  
 (b) Small size and lower electronegativity  
 (c) Large size and lower ionisation energy  
 (d) Large size and largest ionic radius
65. The oxide, which is best soluble in  $\text{H}_2\text{O}$  is [BHU 2001]  
 (a)  $\text{Ba}(\text{OH})_2$  (b)  $\text{Mg}(\text{OH})_2$   
 (c)  $\text{Sr}(\text{OH})_2$  (d)  $\text{Ca}(\text{OH})_2$
66. The property of the alkaline earth metals that increases with their atomic number is [BHU 2001]  
 (a) Ionisation energy  
 (b) Electronegativity  
 (c) Solubility of their sulphates  
 (d) Solubility of their hydroxides
67. In the Alkaline earth metals, the element forming predominantly covalent compound is [BHU 2001]  
 (a) *Be* (b) *Mg*  
 (c) *Sr* (d) *Ca*
68. A mixture of lime paste is sand, water and [RPMT 1997]  
 (a) Gypsum (b) Slacked lime  
 (c) Quick lime (d) Lime stone
69. The formula for calcium chlorite is [CBSE PMT 1994, 96]  
 (a)  $\text{Ca}(\text{ClO}_4)_2$  (b)  $\text{Ca}(\text{ClO}_3)_2$   
 (c)  $\text{CaClO}_2$  (d)  $\text{Ca}(\text{ClO}_2)_2$
70. Which pair of substances gives same gaseous product, when these react with water [CBSE PMT 1994]  
 (a) *Ca* and  $\text{CaH}_2$  (b) *Na* and  $\text{Na}_2\text{O}_2$   
 (c) *K* and  $\text{KO}_2$  (d) *Ba* and  $\text{BaO}_2$



## 766 s and p-Block Elements

71. Magnesium does not decompose the [AFMC 1999]  
 (a) Steam (b) Hot water  
 (c) Cold Water (d) Semi hot water
72. Alkaline earth metals are denser than alkali metals because metallic bonding is [AIIMS 1999]  
 (a) Stronger (b) Weaker  
 (c) Not present (d) Volatile
73. Property of the alkaline earth metals that increases with their atomic number is [IIT 1997]  
 (a) Ionisation energy  
 (b) Solubility of their hydroxides  
 (c) Solubility of their sulphates  
 (d) Electronegativity
74. A metal is burnt in air and the ash on moistening smells of  $NH_3$ . The metal is [AIIMS 1999]  
 (a) Na (b) Fe  
 (c) Mg (d) Al
75. Alkaline earth metals come under [Bihar MEE 1996]  
 (a) Halogens (b) Representative elements  
 (c) Transition elements (d) Inner transition elements  
 (e) None of these
76. Which of the following alkaline-earth metal hydroxides is the strongest base [CPMT 1996]  
 (a)  $Be(OH)_2$  (b)  $Mg(OH)_2$   
 (c)  $Ca(OH)_2$  (d)  $Ba(OH)_2$
77. Which one of the following is the strongest base [Pb. PMT 1998]  
 (a)  $Be(OH)_2$  (b)  $Mg(OH)_2$   
 (c)  $Al(OH)_3$  (d)  $Si(OH)_4$
78. Lime stone is [RPMT 1997]  
 (a) CaO (b)  $Ca(OH)_2$   
 (c) Both (a) and (b) (d) None of these
79. Which of the alkaline earth metals is strongest reducing agent [MP PMT 1995]  
 (a) Ca (b) Sr  
 (c) Ba (d) Mg
80. Plaster of paris hardens by [CPMT 1994]  
 (a) Giving off  $CO_2$  (b) Changing into  $CaCO_3$   
 (c) Uniting with water (d) Giving out water
81. Which is not soluble in water [CPMT 1994]  
 (a)  $CaCO_3$  (b)  $BaCO_3$   
 (c)  $SrCO_3$  (d) All of these
82. The correct order of the increasing ionic character is [MNR 1991; AFMC 1998]  
 (a)  $BeCl_2 < MgCl_2 < CaCl_2 < BaCl_2$   
 (b)  $BeCl_2 < MgCl_2 < BaCl_2 < CaCl_2$   
 (c)  $BeCl_2 < BaCl_2 < MgCl_2 < CaCl_2$   
 (d)  $BaCl_2 < CaCl_2 < MgCl_2 < BeCl_2$
83.  $MgCl_2 \cdot 6H_2O$  when heated gives [CPMT 1997]  
 (a) Magnesium oxychloride [AIIMS 1999]  
 (b) Magnesium dichloride  
 (c) Magnesium oxide  
 (d) Magnesium chloride
84. Which of the following hydroxide is insoluble in water [AIIMS 2001]  
 (a)  $Be(OH)_2$  (b)  $Mg(OH)_2$   
 (c)  $Ca(OH)_2$  (d)  $Ba(OH)_2$
85. Which of the following statements is false [BHU 2005]  
 (a)  $CaOCl_2$  gives  $OH^-$ ,  $Cl^-$  and  $OCl^-$  in aqueous solution [KCET 1996]  
 (b) Diamond and graphite are allotrops of carbon  
 (c) Bleaching action of  $Cl_2$  in moist condition is not permanent  
 (d) Calomel is  $Hg_2Cl_2$
86. A metal  $M$  readily forms its sulphate  $MSO_4$  which is water-soluble. It forms its oxide  $MO$  which becomes inert on heating. It forms its insoluble hydroxide  $M(OH)_2$  which is soluble in  $NaOH$  solution. Then  $M$  is [AIEEE 2002]  
 (a) Mg (b) Ba  
 (c) Ca (d) Be
87. In the lime (kiln), the reaction  $CaCO_3(s) \rightarrow CO_2(g)$  goes to completion because [Kerala (Engg.) 2002]  
 (a) Of high temperature  
 (b) CaO is more stable than  $CaCO_3$   
 (c)  $CO_2$  escapes simultaneously  
 (d) CaO is not dissociated
88. The ionic compound  $BaSO_4$  is insoluble in water due to [CPMT 1999]  
 (a) High lattice energy (b) Low lattice energy  
 (c) Low hydration energy (d) Both (a) and (c)
89. which is used to reduced the acidity of soil [DPMT 2001]  
 (a) Calcium hydroxide (b) Ammonium sulphate  
 (c) Ammonium nitrate (d) Ammonium chloride
90. Alkaline earth metals belong to the [KCET (Med.) 2001]  
 (a)  $s$  - block in periodic table  
 (b)  $p$  - block in periodic table  
 (c)  $d$  - block in periodic table  
 (d)  $f$  - block in periodic table
91. The element having atomic number 56 belongs to [AFMC 2002]

- (a) Actinides (b) Alkaline earth metals  
(c) Transition series (d) Lanthanides
92. The thermal stability of alkaline earth metal carbonates  $MgCO_3, CaCO_3, BaCO_3$  and  $SrCO_3$  decreases as  
[MP PMT 2002]  
(a)  $CaCO_3 > SrCO_3 > MgCO_3 > BaCO_3$   
(b)  $BaCO_3 > SrCO_3 > MgCO_3 > CaCO_3$   
(c)  $BaCO_3 > SrCO_3 > CaCO_3 > MgCO_3$   
(d)  $MgCO_3 > CaCO_3 > SrCO_3 > BaCO_3$
93. A certain metal M is used to prepare an antacid, which is used as a medicine in acidity. This metal accidentally catches fire which can not be put out by using  $CO_2$  based extinguishers. The metal M is  
(a) Ca (b) C  
(c) Mg (d) All of these
94.  $Be(OH)_2$  is insoluble in water while  $Ba(OH)_2$  is highly soluble due to [AMU 2002]  
(a) Bond order (b) Lattice energy difference  
(c) Common ion effect (d) Hard acid
95. Which of the following gives a green colour to flame [AFMC 2001]  
(a) Barium (b) Calcium  
(c) Strontium (d) None of these
96. Sparingly soluble salt is [RPMT 1999]  
(a)  $KCl$  (b)  $NaCl$   
(c)  $NH_4Cl$  (d)  $BaSO_4$
97. Among the alkaline earth metals the element forming predominantly covalent compound is [MP PET 1999]  
(a) Barium (b) Strontium  
(c) Calcium (d) Beryllium
98. Peroxide bond is present in [RPET 2003]  
(a)  $MgO$  (b)  $CaO$   
(c)  $Li_2O$  (d)  $BaO_2$
99. Least ionic character is found in [CPMT 1993]  
(a)  $Mg$  (b)  $Sr$   
(c)  $Ca$  (d)  $Ra$
100. The number of water molecules in gypsum and plaster of paris respectively are  
(a) 1/2 and 2 (b) 2 and 1/2  
(c) 2 and 1 (d) 5 and 2
101. Which of the following is formed when calcium combines with oxygen [MH CET 2000]  
(a) Ca (b)  $CaO$   
(c)  $CaO_2$  (d)  $Ca_2O_2$
102. Slow acting nitrogenous fertilizer among the following is [DCE 2003]  
(a)  $NH_2CONH_2$  (b)  $NH_4NO_3$   
(c)  $CaNCN$  (d)  $KNO_3$
103. Plaster of paris is used [Pb. CET 2000; CPMT 2000]  
(a) In surgery and dentistry  
(b) As a white wash  
(c) As a constituent of tooth paste  
(d) For the preparation of RCC
104. Iron pipes lying under acidic soil are often attached to blocks of magnesium for protection from rusting. Magnesium offers protection to iron against corrosion because it [DPMT 2004; BHU 2004]  
(a) Prevents air from reaching the surface of iron  
(b) is more readily converted into positive ions [BHU 2002]  
(c) Is higher than iron  
(d) Forms a corrosion-resistance alloy with iron
105. Among  $K, Ca, Fe,$  and  $Zn$ , the element which can form more than one binary compound with chlorine is [CBSE PMT 2004]  
(a)  $K$  (b)  $Ca$   
(c)  $Fe$  (d)  $Zn$
106.  $Li$  shows the diagonal relationship with [Pb. CET 2001]  
(a)  $Mg$  (b)  $B$   
(c)  $Al$  (d)  $C$
107. A sodium salt on treatment with  $MgCl_2$  gives white precipitate only on heating. The anion of the sodium salt is [IIT JEE Screening 2004]  
(a)  $HCO_3^-$  (b)  $CO_3^{2-}$   
(c)  $NO_3^-$  (d)  $SO_4^{2-}$
108.  $MgCl_2 \cdot 6H_2O$ . When heated gives [MHCET 2003]  
(a) Magnesium oxide  
(b) Magnesium oxychloride  
(c) Magnesium dichloride  
(d) Magnesium chloride
109.  $Mg$  burns in  $CO$  to produce [Pb. PMT 2001]  
(a)  $MgO_2$  (b)  $MgCO_3$   
(c)  $MgO + CO$  (d)  $MgO + C$
110. Sorel's cement is [Pb. PMT 1999] [Pb. CET 2003]  
(a) Portland cement +  $MgO$   
(b)  $MgCl_2 \cdot CaSiO_3 \cdot 2H_2O$   
(c)  $CaSiO_3 \cdot MgCO_3$   
(d)  $MgCl_2 \cdot 5MgO \cdot xH_2O$
111. Colemanite is [AFMC 2004]  
(a)  $Ca[B_3O_4(OH)_2] \cdot 2H_2O$   
(b)  $Ca_2B_6O_{11} \cdot 5H_2O$   
(c)  $Ca(OH)_2$



## 768 s and p-Block Elements

(d)  $Na_2B_4O_7 \cdot 2H_2O$

### Boron family

- Which of the following statements about  $H_3BO_3$  is not correct [CBSE PMT 1994]
  - It is a strong tribasic acid
  - It is prepared by acidifying an aqueous solution of borax
  - It has a layer structure in which planar  $BO_3$  units are joined by hydrogen bonds
  - It does not act as proton donor but acts as a Lewis acid by accepting hydroxyl ion
- The type of hybridisation of boron in diborane is [CPMT 1999]
  - $sp$ -hybridisation
  - $sp^2$  - hybridisation
  - $sp^3$  - hybridisation
  - $sp^3d^2$  - hybridisation
- In the reaction  $B_2O_3 + C + Cl_2 \rightarrow A + CO$ . The A is [Pb. PMT 2000]
  - $BCl_3$
  - $BCl_2$
  - $B_2Cl_2$
  - $CCl_2$
- The molecular formula of felspar is [MP PMT 2003]
  - $K_2O \cdot Al_2O_3 \cdot 6SiO_2$
  - $K_2O \cdot 3Al_2O_3 \cdot 6SiO_2$
  - $Na_3AlF_6$
  - $CaSO_4 \cdot 2H_2O$
- The most acidic of the following compounds is [Bihar CEE 1995]
  - $P_2O_3$
  - $Sb_2O_3$
  - $B_2O_3$
  - $As_2O_3$
- Identify the statement that is not correct as far as structure of diborane is concerned [Pb. PMT 1998]
  - There are two bridging hydrogen atoms in diborane
  - Each boron atom forms four bonds in diborane
  - The hydrogen atoms are not in the same plane in diborane
  - All B - H bonds in diborane are similar
- Soft heavy metal melts at  $30^\circ C$  and is used in making heat sensitive thermometers the metal is [RPET 2000]
  - Galium
  - Sodium
  - Potassium
  - Caesium
- Which of the following is formed when aluminium oxide and carbon is strongly heated in dry chlorine gas [AFMC 2000]
  - Aluminium chloride
  - Hydrate aluminium chloride
  - Anhydrous aluminium chloride
  - None of these
- Which metal burn in air at high temperature with the evolution of much heat
  - Cu
  - Hg
  - Pb
  - Al
- Aluminium hydroxide is soluble in excess of sodium hydroxide forming the ion [AMU 2001]
  - $AlO_2^{+3}$
  - $AlO_2^{-3}$
  - $AlO_2^-$
  - $AlO_3^-$
- Boron form covalent compound due to [Pb. PMT 2000]
  - Higher ionization energy
  - Lower ionization energy
  - Small size
  - Both (a) and (c)
- In diborane, the two  $H - B - H$  angles are nearly [AIIMS 2005]
  - $60^\circ, 120^\circ$
  - $95^\circ, 120^\circ$
  - $95^\circ, 150^\circ$
  - $120^\circ, 180^\circ$
- Which of the following is a non-metal [MP PMT 1999]
  - Gallium
  - Indium
  - Boron
  - Aluminium
- Which of the following is most acidic [BHU 1998]
  - $Na_2O$
  - $MgO$
  - $Al_2O_3$
  - $CaO$
- When orthoboric acid ( $H_3BO_3$ ) is heated, the residue left is [Pb. PMT 2002]
  - Metaboric acid
  - Boron
  - Boric anhydride
  - Borax
- Which of the following form dimeric halides [Roorkee Qualifying 1998]
  - Al
  - Mg
  - In
  - Ga
- The liquid field metal expanding on solidification is [AIIMS 2004]
  - Ga
  - Al
  - Zn
  - Cu
- Aluminium chloride exists as dimer,  $Al_2Cl_6$  in solid state as well as in solution of non-polar solvents such as benzene. When dissolved in water, it gives [AIEEE 2004]
  - $[Al(OH)_6]^{3-} + 3HCl$
  - $[Al(H_2O)_6]^{3+} + 3Cl^-$
  - $Al^{3+} + 3Cl^-$
  - $Al_2O_3 + 6HCl$
- The hardest substance amongst the following is [Kerala PMT 2004]
  - $Be_2C$
  - Graphite
  - Titanium
  - SiC
  - $B_4C$

20. Which of the following is known as inorganic benzene  
[Pb. CET 2001]  
(a) Borazine (b) Boron nitride  
(c) *p*-dichlorobenzene (d) Phosphonitrilic acid
21. Which of the following is only acidic in nature  
[AIIMS 2004]  
(a)  $Be(OH)_2$  (b)  $Mg(OH)_2$   
(c)  $B(OH)_3$  (d)  $Al(OH)_3$
22. Moissan boron is [DCE 2003]  
(a) Amorphous boron of ultra purity  
(b) Crystalline boron of ultra purity  
(c) Amorphous boron of low purity  
(d) Crystalline boron of low purity
23. Which of the following does not exist in free form  
[Kerala PMT 2004]  
(a)  $BF_3$  (b)  $BCl_3$   
(c)  $BBr_3$  (d)  $BH_3$   
(e) None of these
24. Alumina is [DCE 2002]  
(a) Acidic (b) Basic  
(c) Amphoteric (d) None of these
25. The most abundant metal in the earth crust is  
[Pb. CET 2004]  
(a) *Al* (b) *Ca*  
(c) *Fe* (d) *Na*
26. Crystalline metal can be transformed into metallic glass by  
[NCERT 1984]  
(a) Alloying  
(b) Pressing into thin plates  
(c) Slow cooling of molten metal  
(d) Very rapid cooling of a spray of the molten metal
27. Which metal is protected by a layer of its own oxide  
[NCERT 1981; DPMT 1983; BHU 1998]  
(a) *Al* (b) *Ag*  
(c) *Au* (d) *Fe*
28. Aluminium is a self-preserving metal, because  
(a) It is not tarnished by air  
(b) A thin film of basic carbonate on its surface  
(c) A non-porous layer of oxide is formed on its surface  
(d) It is not affected by salt water
29. Anhydrous  $AlCl_3$  cannot be obtained from which of the following reactions [CPMT 1987]  
(a) Heating  $AlCl_3 \cdot 6H_2O$   
(b) By passing dry *HCl* over hot aluminium powder  
(c) By passing dry  $Cl_2$  over hot aluminium powder  
(d) By passing dry  $Cl_2$  over a hot mixture of alumina and coke
30. An element *A* dissolves both in acid and alkali. It is an example of [NCERT 1972]  
(a) Allotropic nature of *A* (b) Dimorphic nature of *A*  
(c) Amorphous nature of *A* (d) Amphoteric nature of *A*
31. Hydrogen gas will not reduce [IIT 1984]  
(a) Heated cupric oxide  
(b) Heated ferric oxide  
(c) Heated stannic oxide  
(d) Heated aluminium oxide
32. Conc.  $HNO_3$   
(a) Reacts with aluminium vigorously  
(b) Reacts with aluminium to form aluminium nitrate  
(c) Does not react with aluminium  
(d) Reacts with platinum
33. Anhydrous  $AlCl_3$  is obtained from [BHU 1980; CPMT 1982]  
(a) *HCl* and aluminium metal  
(b) Aluminium and chlorine gas  
(c) Hydrogen chloride gas and aluminium metal  
(d) None of the above
34. Which is true for an element *R* present in III group of the periodic table [EAMCET 1991]  
(a) It is gas at room temperature  
(b) It has oxidation state of +4  
(c) It forms  $R_2O_3$   
(d) It forms  $RX_2$
35. When *Al* is added to *KOH* solution [NCERT 1974, 76; CPMT 1977]  
(a) No action takes place  
(b) Oxygen is evolved  
(c) Water is produced  
(d) Hydrogen is evolved
36. Aluminium is more reactive than iron. But aluminium is less easily corroded than iron because [KCET 1993]  
(a) Aluminium is a noble metal  
(b) Oxygen forms a protective oxide layer  
(c) Iron undergoes reaction easily with water  
(d) Iron forms mono and divalent ions
37. Aluminium vessels should not be washed with materials containing washing soda since [KCET 1993]  
(a) Washing soda is expensive



## 770 s and p-Block Elements

- (b) Washing soda is easily decomposed  
 (c) Washing soda reacts with aluminium to form soluble aluminate  
 (d) Washing soda reacts with aluminium to form insoluble aluminium oxide
38. Which of the statements about anhydrous aluminium chloride is correct [IIT 1981]  
 (a) It exists as  $AlCl_3$  molecule  
 (b) It is not easily hydrolysed  
 (c) It sublimes at  $100^\circ C$  under vacuum  
 (d) It is a strong Lewis base
39. Common alum is [DPMT 1982; CPMT 1978; AMU 1982, 83]  
 (a)  $K_2SO_4 \cdot Al_2(SO_4)_3 \cdot 24H_2O$   
 (b)  $K_2SO_4 \cdot Cr_2(SO_4)_3 \cdot 24H_2O$   
 (c)  $K_2SO_4 \cdot Fe_2(SO_4)_3 \cdot 24H_2O$   
 (d)  $(NH_4)_2SO_4 \cdot FeSO_4 \cdot 6H_2O$
40. Which of the following is not true about potash alum [MNR 1993; UPSEAT 2002]  
 (a) Its empirical formula is  $KAl(SO_4)_2 \cdot 12H_2O$   
 (b) Its aqueous solution is basic  
 (c) It is used in dyeing industries  
 (d) On heating it melts in its water of crystallization
41. Which one of the following is correct statement  
 (a) The hydroxide of aluminium is more acidic than that of boron  
 (b) The hydroxide of boron is basic, while that of aluminium is amphoteric  
 (c) The hydroxide of boron is acidic, while that of aluminium is amphoteric  
 (d) The hydroxide of boron and aluminium are amphoteric
42.  $AlCl_3$  is [AFMC 1995]  
 (a) Anhydrous and covalent (b) Anhydrous and ionic  
 (c) Covalent and basic (d) Coordinate and acidic
43. Aluminium (III) chloride forms a dimer because [CBSE PMT 1995]  
 (a) Higher coordination number can be achieved by aluminium  
 (b) Aluminium has high ionization energy  
 (c) Aluminium belongs to III group  
 (d) It cannot form a trimer
44. Aluminium has a great affinity for oxygen and its oxidation is an exothermic process. This fact is made use of in [MP PMT 1997]  
 (a) Preparing thin foils of aluminium  
 (b) Making utensils  
 (c) Preparing duralumin alloy  
 (d) Thermite welding
45. Number of water molecules in Mohr's salt is [CPMT 1997; AIIMS 2001; JIPMER 2001]  
 (a) 7 (b) 6  
 (c) 5 (d) 8
46. Which of the following is an amphoteric oxide [BHU 2001]  
 (a)  $MgO$  (b)  $Al_2O_3$   
 (c)  $Cl_2O_7$  (d)  $Ti_2O_2$
47. Aluminium oxide is not reduced by chemical reactions since [KCET 2002]  
 (a) Aluminium oxide is reactive  
 (b) Reducing agents contaminate  
 (c) Aluminium oxide is highly stable  
 (d) The process pollutes the environment
48. Aluminium is not used [DPMT 2002]  
 (a) In silvery paints  
 (b) For making utensils  
 (c) As a reducing agent  
 (d) As oxidizer in metallurgy
49. In the thermite process the reducing agent is [Pb. PMT 2002]  
 (a)  $Al$  (b)  $C$   
 (c)  $Mg$  (d)  $Na$
50. In Goldschmidt aluminothermic process, thermite contains [KCET 2003]  
 (a) 3 parts of  $Al_2O_3$  and 4 parts of  $Al$   
 (b) 3 parts of  $Fe_2O_3$  and 2 parts of  $Al$   
 (c) 3 parts of  $Fe_2O_3$  and 1 part of  $Al$   
 (d) 1 part of  $Fe_2O_3$  and 1 part of  $Al$
51. Bauxite containing impurities of iron oxide is purified by [CPMT 1987; AIIMS 1998]  
 (a) Hoop's process (b) Serpeck's process  
 (c) Baeyer's process (d) Electrolytic process
52. In the purification of bauxite by Hall's process  
 (a) Bauxite ore is heated with  $NaOH$  solution at  $50^\circ C$   
 (b) Bauxite ore is fused with  $Na_2CO_3$   
 (c) Bauxite ore is fused with coke and heated at  $1800^\circ C$  in a current of nitrogen  
 (d) Bauxite ore is heated with  $NaHCO_3$
53. Which one is used as a bye-product in Serpeck's process  
 (a)  $NH_3$  (b)  $CO_2$   
 (c)  $N_2$  (d)  $PH_3$
54. In the metallurgy of aluminium, cryolite is mixed in the molten state because it [Roorkee 1995]  
 (a) Increases the melting point of alumina  
 (b) Oxidises alumina



- (c) Reduces alumina  
(d) Decreases the melting point of alumina
55. In the electrolytic extraction of aluminium, cryolite is used  
[NCERT 1981; CPMT 1989; RPMT 2000; MP PMT 2000, 02]
- (a) To obtain more aluminium  
(b) To decrease temperature to dissolve bauxite  
(c) To protect the anode  
(d) As reducing agent
56. In the extraction of aluminium, bauxite is dissolved in cryolite because  
(a) It acts as a solvent  
(b) It reduces melting point of aluminium oxide  
(c) It increases the resistance of aluminium oxide  
(d) Bauxite becomes active
57. In the extraction of aluminium the electrolyte is  
[CBSE PMT 1989; AIEEE 2002]
- (a) Fused cryolite with felspar  
(b) Fused cryolite with fluorspar  
(c) Pure alumina in molten cryolite  
(d) Pure alumina with bauxite and molten cryolite
58. Aluminium is obtained by [KCET 1992; RPMT 2002]
- (a) Reducing  $Al_2O_3$  with coke  
(b) Electrolysing  $Al_2O_3$  dissolved in  $Na_3AlF_6$   
(c) Reducing  $Al_2O_3$  with chromium  
(d) Heating alumina and cryolite
59. In the electrolysis of alumina, cryolite is added to  
[IIT 1986; BHU 1987]
- (a) Increase the melting point of alumina  
(b) Increase the electrical conductivity  
(c) Minimise the anodic effect  
(d) Remove impurities from alumina
60. The function of fluorspar in the electrolytic reduction of alumina dissolved in fused cryolite ( $Na_3AlF_6$ ) is  
[KCET 1993; IIT 1993]
- (a) As a catalyst  
(b) To lower the temperature of the melt and to make the fused mixture very conducting  
(c) To decrease the rate of oxidation of carbon at the anode  
(d) None of the above
61. For purification of alumina, the modern processes most useful when (i) the impurity present is a lot of iron oxides and (ii) the impurity present is a lot of silica, are  
(a) For (i) Hall's process; for (ii) Baeyer's process  
(b) For (i) Hall's process; for (ii) Serpeck's process  
(c) For (i) Serpeck's process; for (ii) Baeyer's process  
(d) For (i) Baeyer's process; for (ii) Serpeck's process
62. For the electrolytic production of aluminium, (i) the cathode and (ii) the anode are made of  
(a) (i) Platinum and (ii) Iron  
(b) (i) Copper and (ii) Iron  
(c) (i) Copper and (ii) Carbon  
(d) (i) Carbon and (ii) Carbon
63. In the commercial electrochemical process for aluminium extraction, the electrolyte used is [IIT-JEE 1999]
- (a)  $Al(OH)_3$  in  $NaOH$  solution  
(b) An aqueous solution of  $Al_2(SO_4)_3$   
(c) A molten mixture of  $Al_2O_3$  and  $Na_3AlF_6$   
(d) A molten mixture of  $AlO(OH)$  and  $Al(OH)_3$
64. In electrolysis of aluminium oxide which of the following is added to accelerate the process [AFMC 1999; C]
- (a) Silica (b) Cryolite  
(c) Nickel (d) Silicate
65. The purification of alumina is called  
[CPMT 1997; AFMC 1998; AIIMS 1999]
- (a) Bosch process (b) Caster process  
(c) Baeyer's process (d) Hoop's process
66. Electrolytic reduction of alumina to aluminium by Hall-Heroult process is carried out in the presence of  
[IIT-JEE (Screening) 2000]
- (a)  $NaCl$   
(b) Fluorite  
(c) Cryolite which forms a melt with lower melting temperature  
(d) Cryolite which forms a melt with higher melting temperature
67. In the electrolytic method of obtaining aluminium from purified bauxite, cryolite is added to the charge in order to  
[KCET 2004]
- (a) Minimize the heat loss due to radiation  
(b) Protect aluminium produced from oxygen  
(c) Dissolve bauxite and render it conductor of electricity  
(d) Lower the melting point of bauxite
68. Hoop's process is used for the purification of the metal  
[MP PET 1995; MP PMT 2001]
- (a)  $Al$  (b)  $Zn$   
(c)  $Ag$  (d)  $Cu$
69. Purification of aluminium done by electrolytic refining is known as  
[CPMT 1989; CBSE PMT 1999; RPET 2003; BCECE 2005]
- (a) Serpeck's process (b) Hall's process  
(c) Baeyer's process (d) Hoop's process
70. In the Hoop's process for refining of aluminium, the fused materials form three different layers and they remain separated during electrolysis also. This is because  
[MP PET 1996]



## 772 s and p-Block Elements

- (a) The upper layer is kept attracted by the cathode and the lower layer is kept attracted by the anode  
(b) There is special arrangement in the cell to keep the layers separate  
(c) The 3 layers have different densities  
(d) The 3 layers are maintained at different temperatures

71. During metallurgy of aluminium bauxite is dissolved in cryolite because

- (a) Bauxite is non-electrolyte  
(b) Cryolite is a flux  
(c) Cryolite acts as an electrolyte  
(d) All are correct

72. For the electrolytic refining of aluminium, the three fused layers consist of

	Bottom Layer	Middle Layer	Upper Layer
(a)	Cathode of pure Al	Cryolite and fluorspar	Anode of Al and Cu alloy
(b)	Cathode of Al and Cu alloy	Bauxite and cryolite	Anode of pure Al
(c)	Anode of Al and Cu alloy	Cryolite and barium fluoride	Cathode of pure Al
(d)	Anode of impure Al	Bauxite, cryolite and fluorspar	Cathode of pure Al

73. Heating an aqueous solution of aluminium chloride to dryness will give [AIEEE 2005]

- (a)  $AlCl_3$  (b)  $Al_2Cl_6$   
(c)  $Al_2O_3$  (d)  $Al(OH)Cl_2$

74. The structure of diborane ( $B_2H_6$ ) contains [AIEEE 2005]

- (a) Four 2c-2e bonds and two 3c-2e bonds  
(b) Two 2c-2e bonds and four 3c-2e bonds  
(c) Two 2c-2e bonds and two 3c-3e bonds  
(d) Four 2c-2e bonds and four 3c-2e bonds

75. Which of the following is the electron deficient molecule

[CBSE PMT 2005]

- (a)  $B_2H_6$  (b)  $C_2H_6$   
(c)  $PH_3$  (d)  $SiH_4$

76. In Hall's process, the main reagent is mixed with

[AFMC 2005]

- (a) NaF (b)  $Na_3AlF_6$   
(c)  $AlF_3$  (d) None of these

77. Acidity strength of Boron trihalide are in order of

[Kerala CET 2005]

- (a)  $BF_3 < BCl_3 < BBr_3 < BI_3$   
(b)  $BI_3 < BBr_3 < BCl_3 < BF_3$   
(c)  $BBr_3 < BCl_3 < BF_3 < BI_3$   
(d)  $BF_3 < BI_3 < BCl_3 < BBr_3$



## Carbon family

1. Carbon and silicon belong to (IV) group. The maximum coordination number of carbon in commonly occurring compounds is 4, whereas that of silicon is 6. This is due to  
[CBSE PMT 1994]
- (a) Large size of silicon  
(b) More electropositive nature of silicon  
(c) Availability of low lying *d*-orbitals in silicon  
(d) Both (a) and (b)
2. The ionic carbide is [JIPMER 2000]
- (a)  $ZnC$  (b)  $TiC$   
(c)  $SiC$  (d)  $CaC_2$
3.  $PbO_2$  is [JIPMER 2000]
- (a) Basic (b) Acidic  
(c) Neutral (d) Amphoteric
4. Lead pipes are not suitable for drinking water because  
[JIPMER 2000]
- (a) A layer of lead dioxide is deposited over pipes  
(b) Lead reacts with air to form litharge  
(c) Lead reacts with water containing air to form  $Pb(OH)_2$   
(d) Lead forms basic lead carbonate
5. Silicon dioxide is formed by the reaction of [KCET (Med.) 2001]
- (a)  $SiCl_4 + 2H_2O$  (b)  $SiO_2 + 4HF$   
(c)  $SiO_2 + NaOH$  (d)  $SiCl_4 + NaOH$
6. Which alkali metal carbonate decomposes on heating to liberate  $CO_2$  gas
- (a)  $Li_2CO_3$  (b)  $CaCO_3$   
(c)  $Na_2CO_3$  (d)  $Al_2CO_3$
7. Which of the following gives propyne on hydrolysis  
[AIIMS 2005]
- (a)  $Al_4C_3$  (b)  $Mg_2C_3$   
(c)  $B_4C$  (d)  $La_4C_3$
8. Which one of the following statements is not correct  
[CBSE PMT 1994]
- (a) Zinc dissolves in sodium hydroxide solution  
(b) Carbon monoxide reduces iron (III) oxide to iron  
(c) Mercury (II) iodide dissolves in excess of potassium iodide solution  
(d) Tin (IV) chloride is made by dissolving tin solution in concentrated hydrochloric acid
9. In laboratory silicon can be prepared by the reaction  
[Pb. PMT 1999]
- (a) By heating carbon in electric furnace  
(b) By heating potassium with potassium dichromate  
(c) Silica with magnesium  
(d) None of these
10. Which of the following is the correct statement for red lead  
[AIIMS 2000]
- (a) It is an active form of lead  
(b) Its molecular formula is  $Pb_2O_3$   
(c) It decomposes into  $Pb$  and  $CO_2$   
(d) It decomposes into  $PbO$  and  $O_2$
11. Suppose you have to determine the percentage of carbon dioxide in a sample of a gas available in a container. Which is the best absorbing material for the carbon dioxide  
[Pb. PMT 2001]
- (a) Heated copper oxide  
(b) Cold, solid calcium chloride  
(c) Cold, solid calcium hydroxide  
(d) Heated charcoal
12. The number and type of bonds between 2 carbon atoms in  $CaC_2$  [UPSEAT 2001]
- (a) One sigma ( $\sigma$ ) and one pi ( $\pi$ ) bond  
(b) One sigma ( $\sigma$ ) and two pi ( $\pi$ ) bond  
(c) One sigma ( $\sigma$ ) and half pi ( $\pi$ ) bond  
(d) One sigma ( $\sigma$ ) bond
13. Metalloid among the following is [DPMT 2001]
- (a)  $Si$  (b)  $C$   
(c)  $Pb$  (d)  $Ge$
- [Pb. PMT 2000]
14. 'Lead pencil' contains [DPMT 2001; IIT 1990]
- (a)  $PbS$  (b) Graphite  
(c)  $FeS$  (d)  $Pb$
15. Nitrogen gas is absorbed by [DPMT 2001]
- (a) Calcium hydroxide (b) Ferrous sulphate  
(c) Calcium carbide (d) Aluminium carbide
16. In laboratory silicon can be prepared by the reaction  
[Pb. PMT 1999; AFMC 2002]
- (a) Silica with magnesium  
(b) By heating carbon in electric furnace  
(c) By heating potassium fluosilicate with potassium  
(d) None of these
17. Formation of in-numberable compounds of carbon is due to its
- (a) High reactivity  
(b) Catenation tendency



## 772 s and p-Block Elements

- (c) Covalent and ionic tendency  
(d) Different valency
18. Colour is imparted to glass by mixing [Pb. PMT 2002]  
(a) Synthetic dyes (b) Metal oxide  
(c) Oxides of non-metal (d) Coloured salt
19. Which of the following is insoluble in water [MP PET 2002]  
(a)  $Na_2CO_3$  (b)  $CaCO_3$   
(c)  $ZnCO_3$  (d)  $Al_2(CO_3)_3$
20. In which of the following the inert pair effect is most prominent [MP PMT 2000]  
(a) C (b) Si  
(c) Ge (d) Pb
21. Plumbosolvency implies dissolution of lead in [DCE 1999]  
(a) Bases (b) Acids  
(c) Ordinary water (d)  $CuSO_4$  sol
22. Which of the following glass is used in making wind screen of automobiles [AIIMS 1999; Pb. CET 2000]  
(a) Crook's (b) Jena  
(c) Safety (d) Pyrex
23. Glass reacts with HF to produce [KCET 2000; CBSE PMT PMT 2000]  
(a)  $SiF_4$  (b)  $H_2SiF_6$   
(c)  $H_2SiO_3$  (d)  $Na_3AlF_6$
24. The type of glass used in making lenses and prisms is [JIPMER 1999]  
(a) A flint glass (b) Jena glass  
(c) Pyrex glass (d) Quartz glass
25. When carbon monoxide is passed over solid caustic soda heated to  $200^\circ C$ , it forms [KCET (Med.) 1999]  
(a)  $Na_2CO_3$  (b)  $NaHCO_3$   
(c)  $H-COONa$  (d)  $CH_3COONa$
26. Which is used to produce smoke screens [AFMC 2005]  
(a) Calcium phosphide (b) Zinc sulphide  
(c) Sodium carbonate (d) Zinc phosphide
27. Sodium oxalate on heating with conc.  $H_2SO_4$  gives [Roorkee 2000]  
(a) CO only (b)  $CO_2$  only  
(c) CO and  $CO_2$  (d)  $SO_2$  and  $SO_3$
28. Extraction of lead by reduction methods is done by [AMU 2000]  
(a) Adding more galena into reverberatory furnace  
(b) Adding more lead sulphate into reverberatory furnace  
(c) Adding more galena and coke into the reverberatory furnace  
(d) Self reduction of oxide from sulphide present in the furnace
29. Which gas is used in excess water [BVP 2003]  
(a)  $CO_2$  (b)  $SO_2$   
(c) CO (d) Water vapours
30. The compound which does not possess a peroxide linkage is  
(a)  $Na_2O_2$  (b)  $CrO_5$   
(c)  $H_2SO_5$  (d)  $PbO_2$
31. Silicon is an important constituent of [MH CET 2001]  
(a) Rocks (b) Amalgams  
(c) Chlorophyll (d) Haemoglobin
32. Carborundum is [AFMC 2002; MH CET 2003; BHU 2003, 05]  
(a) SiC (b)  $AlCl_3$   
(c)  $Al_2(SO_4)_3$  (d)  $Al_2O_3 \cdot 2H_2O$
33.  $SiF_4$  gets hydrolysed giving ..... [Orissa JEE 2002]  
(a)  $SiO_2$  (b)  $Si(OH)_2F_2$   
(c)  $H_2SiF_6$  (d)  $Si(OH)_4$
34. Glass is a [AIEEE 2003; RPET 2003]  
(a) Micro-crystalline solid  
(b) Super cooled liquid  
(c) Gel  
(d) Polymeric mixture
35.  $H_2O_2$  on reaction with PbS gives [RPET 2003]  
(a) PbO (b)  $PbSO_4$   
(c)  $PbO_2$  (d)  $PbHSO_4$
36. Soldiers of Napoleon army while at Alps during freezing winter suffered a serious problem as regards to the tin buttons of their uniforms. White metallic tin buttons got converted to grey powder. This transformation is related to [AIEEE 2004]  
(a) A change in the partial pressure of oxygen in the air  
(b) A change in the crystalline structure of tin  
(c) An interaction with nitrogen of the air at very low to temperatures  
(d) An interaction with water vapour contained in the humid air
37. Solid  $CO_2$  is known as dry ice, because [Pb. CET 2000]  
(a) It melts at  $0^\circ C$   
(b) It evaporates at  $40^\circ C$   
(c) It evaporates at  $-78^\circ C$  without melting  
(d) Its boiling point is more than  $199^\circ C$
38. Which one of the following statements about the zeolites is false [CBSE PMT 2004]  
(a) Zeolites are aluminosilicates having three dimensional network  
(b) Some of the  $SiO_4^{-4}$  units are replaced by  $AlO_4^{-5}$  and  $AlO_6^{9-}$  ions in zeolites

- (c) They are used as cation exchangers  
(d) They have open structure which enables them to take up small molecules
39. Which of the following cuts ultraviolet rays [AFMC 2004]  
(a) Soda glass (b) Crooke's glass  
(c) Pyrax (d) None of these
40. In IIIA group, *Tl* (thallium) shows +1 oxidation state while other members show +3 oxidation state. Why [JEE Orissa 2004]  
(a) Presence of lone pair of electron in *Tl*  
(b) Inert pair effect  
(c) Large ionic radius of *Tl* ion  
(d) None of these
41. Carbon suboxide  $C_3O_2$  has [DCE 2003]  
(a) Linear structure  
(b) Bent structure  
(c) Trigonal planar structure  
(d) Distorted tetrahedral structure
42. Which of the following is a mixed oxide [Pb. CET 2003]  
(a)  $Fe_2O_3$  (b)  $PbO_2$   
(c)  $Pb_3O_4$  (d)  $BaO_2$
43. Noble gases are absorbed on [BVP 2004]  
(a) Anhydrous  $CaCl_2$  (b) Charcoal  
(c) Conc.  $H_2SO_4$  (d) Coconut
44. Lapis lazuli is [AFMC 2004]  
(a) Ferrous sulphate (b) Copper sulphate  
(c) Sodium aluminosilicate (d) Zinc sulphate
45. Which of the following statement is correct with respect to the property of elements in the carbon family with an increase in atomic number, their [Pb. CET 2002]  
(a) Atomic size decreases  
(b) Ionization energy increases  
(c) Metallic character decreases  
(d) Stability of +2 oxidation state increases
46. When tin is treated with concentrated nitric acid [DCE 2004]  
(a) It is converted into stannous nitrate  
(b) It is converted into stannic nitrate  
(c) It is converted into metastannic acid  
(d) It becomes passive
47. Solder is an alloy of [Pb. CET 2003]  
(a)  $Pb + Zn + Sn$  (b)  $Pb + Zn$   
(c)  $Pb + Sn$  (d)  $Sn + Zn$
48. A metal used in storage batteries is  
(a) Copper (b) Lead  
(c) Tin (d) Nickel
49. Name of the structure of silicates in which three oxygen atoms of  $[SiO_4]^{4-}$  are shared is [IIT 2005]  
(a) Pyrosilicate  
(b) Sheet silicate  
(c) Linear chain silicate  
(d) Three dimensional silicate
50. Red lead is [CPMT 1972, 74, 94; MNR 1985; DPMT 1982, 2002; Bihar CEE 1995; MP PET 1995]  
(a)  $Pb_3O_4$  (b)  $PbO$   
(c)  $PbO_2$  (d)  $Pb_4O_3$
51. White lead is [CPMT 1983, 93, 2002; MNR 1984; MP PMT 1995; UPSEAT 1999; DCE 2000]  
(a)  $PbCO_3$  (b)  $PbCO_3.PbO$   
(c)  $2PbCO_3.Pb(OH)_2$  (d)  $2PbSO_4.PbO$
52. Lead pipes are corroded quickly by [AFMC 1981]  
(a) Dil.  $H_2SO_4$  (b) Conc.  $H_2SO_4$   
(c) Acetic acid (d) Water
53. In silicon dioxide [AIIEEE 2005]  
(a) Each silicon atom is surrounded by four oxygen atoms and each oxygen atom is bonded to two silicon atoms  
(b) Each silicon atom is surrounded by two oxygen atoms and each oxygen atom is bonded to two silicon atoms  
(c) Silicon atom is bonded to two oxygen atoms  
(d) There are double bonds between silicon and oxygen atoms
54. Litharge is chemically [DPMT 1984; JIPMER 2001]  
(a)  $PbO$  (b)  $PbO_2$   
(c)  $Pb_3O_4$  (d)  $Pb(CH_3COO)_2$
55. The element of  $s^2p^2$  configuration is of ..... group  
(a) IV (b) III  
(c) V (d) II
56. Which of the following compounds of elements in group IV would you expect to be most ionic in character [NCERT 1978]  
(a)  $CCl_4$  (b)  $SiCl_4$   
(c)  $PbCl_2$  (d)  $PbCl_4$
57. Which of the following compounds of lead is used in match industry  
(a)  $PbO$  (b)  $PbO_2$   
(c)  $PbCl_2$  (d) None of these
58. Type metal is an alloy of *Pb*, *Sb* and *Sn*. It consists of  
(a) Equal amounts of the three metals  
(b) More amount of lead  
(c) More amount of antimony  
(d) More amount of tin
59. Which is correct oxidation state of lead [AFMC 1987]  
(a) + 2, + 4 (b) + 1, + 2  
(c) + 3, + 4 (d) + 4
60. Sugar of lead is  
(a)  $2PbSO_4.PbO$  (b)  $(CH_3COO)_2Pb$   
(c)  $PbCO_3$  (d)  $PbCO_3.Pb(OH)_2$



## 774 s and p-Block Elements

61. Which of the following compounds has peroxide linkage  
[CPMT 1988]  
(a)  $Pb_2O_3$  (b)  $SiO_2$   
(c)  $CO_2$  (d)  $PbO_2$
62. Percentage of lead in lead pencil is  
[CBSE PMT 1999]  
(a) Zero (b) 20  
(c) 80 (d) 70
63. Which of the following has most density  
[CPMT 1996]  
(a)  $Fe$  (b)  $Cu$   
(c)  $B$  (d)  $Pb$
64. Red lead is an example of a/an .... oxide  
[JIPMER 2001]  
(a) Basic (b) Super  
(c) Mixed (d) Amphoteric
65. Which of the following lead oxides is 'Sindhur'  
[MP PET 2002]  
(a)  $PbO$  (b)  $PbO_2$   
(c)  $Pb_2O_3$  (d)  $Pb_3O_4$
66. Element showing the phenomenon of allotropy is  
[MP PMT 1999]  
(a) Aluminium (b) Tin  
(c) Lead (d) Copper
67. Which of the following element is a metalloid  
[CPMT 2004]  
(a)  $Bi$  (b)  $Sn$   
(c)  $Ge$  (d)  $C$
68. Which gas is liberated when  $Al_4C_3$  is hydrolysed  
[AFMC 2005]  
(a)  $CH_4$  (b)  $C_2H_2$   
(c)  $C_2H_6$  (d)  $CO_2$
69. Which of the following attacks glass  
[NCERT 1976; AFMC 2005]  
(a)  $HCl$  (b)  $HF$   
(c)  $HI$  (d)  $HBr$
- Pb. PMT 2000; AFMC 1988, 92; MP PET 1997, 2000, 01  
(a)  $N_2O$  (b)  $NO$   
(c)  $N_2O_3$  (d)  $N_2O_5$
4. Metaphosphoric acid has the formula [CPMT 1973, 89, 93]  
(a)  $H_3PO_4$  (b)  $HPO_3$   
(c)  $H_2PO_3$  (d)  $H_3PO_2$
5. Which of the following is the most suitable drying agent for ammonia gas  
[MP PMT 1989; CBSE PMT 1989; DPMT 1982; CPMT 1974, 78, 91; BHU 1986, 96; 2001; IIT (Screening) 2000]  
(a) Calcium oxide  
(b) Anhydrous calcium chloride  
(c) Phosphorus pentoxide  
(d) Conc. sulphuric acid
6. Each of the following is true for white and red phosphorus except that they  
(a) Are both soluble in  $CS_2$   
(b) Can be oxidised by heating in air  
(c) Consists of same kind of atoms  
(d) Can be converted into one another
7. Which of the following is a tetrabasic acid [CPMT 1988]  
(a) Orthophosphorus acid (b) Orthophosphoric acid  
(c) Metaphosphoric acid (d) Pyrophosphoric acid
8. Phosphine is prepared by the reaction of  
[MP PET/PMT 1988]  
(a)  $P$  and  $H_2SO_4$  (b)  $P$  and  $NaOH$   
(c)  $P$  and  $H_2S$  (d)  $P$  and  $HNO_3$
9. Which of the following is not known [MP PET/PMT 1988; NCERT 1982; CBSE PMT 1989; MP PET 1993]  
(a)  $NCl_5$  (b)  $NI_3$   
(c)  $SbCl_3$  (d)  $NCl_3$
10. Chemical formula for the phosphorus molecule is  
[CPMT 1976, 80, 84, 90; BHU 1984, 86; NCERT 1977]  
(a)  $P$  (b)  $P_4$   
(c)  $P_2$  (d)  $P_5$
11. White phosphorus ( $P_4$ ) has [IIT 1998]  
[CBSE PMT 1989, 94]  
(a) Six  $P-P$  single bonds  
(b) Four  $P-P$  single bonds  
(c) Four lone pairs of electrons  
(d)  $PPP$  angle of  $60^\circ$
12. Ammonium nitrate decomposes on heating into  
[NCERT 1974, 75; CPMT 1973, 78, 88, 94; AMU 1984]  
(a) Ammonia and nitric acid  
(b) Nitrous oxide and water  
(c) Nitrogen, hydrogen and ozone

## Nitrogen family

1. Which of the following elements does not form stable diatomic molecules  
(a) Iodine (b) Phosphorus  
(c) Nitrogen (d) Oxygen
2. Producer gas is a mixture of [DPMT 1982; CPMT 1978]  
(a)  $CO$  and  $N_2$  (b)  $CO_2$  and  $H_2$   
(c)  $CO$  and  $H_2$  (d)  $CO_2$  and  $N_2$
3. Which one of the following combines with  $Fe(II)$  ions to form a brown complex  
[AIIMS 1982, 83, 87; BHU 1998; CBSE PMT 2000;

- (d) Nitric oxide, nitrogen dioxide and hydrogen
13. In Birkeland-Eyde process, the raw material used is  
[CPMT 1982, 86]  
(a) Air (b)  $NH_3$   
(c)  $NO_2$  (d)  $HNO_3$
14. Among the following nitrates, Lead nitrate, Silver nitrate and Ammonium nitrate; the one that decomposes without leaving any solid residue is [NCERT 1983]  
(a) Lead nitrate (b) Ammonium nitrate  
(c) Silver nitrate (d) Sodium nitrate
15. Of the different allotropes of phosphorus, the one which is most reactive is [CPMT 1983; NCERT 1978; CBSE PMT 1999; Kurukshetra CEE 1998]  
(a) Violet phosphorus (b) Scarlet phosphorus  
(c) Red phosphorus (d) White phosphorus
16. Phosphine is generally prepared in the laboratory [CPMT 1983, 2003]  
(a) By heating phosphorus in a current of hydrogen  
(b) By heating white phosphorus with aqueous solution of caustic potash  
(c) By decomposition of  $P_2H_4$  at  $110^\circ C$   
(d) By heating red phosphorus with an aqueous solution of caustic soda
17. Which of the following elements is most metallic [CPMT 1983; MP PMT 1993]  
(a) Phosphorus (b) Arsenic  
(c) Antimony (d) Bismuth
18. The basicity of orthophosphoric acid is [CPMT 1984, 91]  
(a) 2 (b) 3  
(c) 4 (d) 5
19.  $HNO_2$  acts as [AFMC 1992]  
(a) Oxidising agent (b) Reducing agent  
(c) Both (a) and (b) (d) Its solution is stable
20. Nitrogen dioxide cannot be obtained by heating [CPMT 1989; IIT 1985; CPMT 1993]  
(a)  $KNO_3$  (b)  $Pb(NO_3)_2$   
(c)  $Cu(NO_3)_2$  (d)  $AgNO_3$
21. When heated  $NH_3$  is passed over  $CuO$  gas evolved is [BCECE 2005]  
(a)  $N_2$  (b)  $N_2O$   
(c)  $HNO_3$  (d)  $NO_2$
22. Non-combustible hydride is [CPMT 1979]  
(a)  $NH_3$  (b)  $PH_3$   
(c)  $AsH_3$  (d)  $SbH_3$
23. On heating a mixture of  $NH_4Cl$  and  $KNO_2$  we get [CPMT 1972, 79; NCERT 1977]  
(a)  $NH_4NO_3$  (b)  $N_2$   
(c)  $N_2O$  (d)  $NO$
24. Which of the following oxide of nitrogen is the anhydride of  $HNO_3$  [CPMT 1979, 80, 89, 97; MP PET/PMT 1988; KCET 1991; CBSE PMT 1989, 91, 99; EAMCET 1991; NCERT 1975; MP PET 1989; MP PMT 1994]  
(a)  $NO$  (b)  $N_2O_3$   
(c)  $N_3O_4$  (d)  $N_2O_5$
25. Phosphorus is manufactured by heating in a electric furnace a mixture of [NCERT 1977; CPMT 1974, 78, 81]  
(a) Bone ash and coke  
(b) Bone ash and silica  
(c) Bone ash, silica and coke  
(d) None of these
26. A certain element forms a solid oxide which when dissolved in water forms an acidic solution, the element is [CPMT 1972, 78]  
(a) Argon (b) Potassium  
(c) Phosphorus (d) Sulphur
27. Dissociation of  $H_3PO_4$  occurs in following stages [CPMT 1976]  
(a) 1 (b) 2  
(c) 3 (d) 4
28. Nitrogen forms how many oxides  
(a) 3 (b) 4  
(c) 5 (d) 6
29. The  $P-P-P$  bond angle in white phosphorus is [MP PET 1991]  
(a)  $120^\circ$  (b)  $109^\circ 28'$   
(c)  $90^\circ$  (d)  $60^\circ$
30. Ammonium dichromate on heating gives [BHU 1973, 78; CBSE PMT 1993; MP PMT 1993]  
(a) Chromium oxide and ammonia  
(b) Chromic acid and nitrogen  
(c) Chromium oxide and nitrogen  
(d) Chromic acid and ammonia
31. When concentrated nitric acid is heated, it decomposes to give  
(a)  $O_2$  and  $N_2$  (b)  $NO$   
(c)  $N_2O_5$  (d)  $NO_2$  and  $O_2$
32. The element which catches fire in air at  $30^\circ C$  and is stored under water is [BHU 1973; MP PET 1989, 99]  
(a) Calcium (b) Sodium  
(c) Phosphorus (d) Zinc
33. A solution of ammonia in water contains



## 776 s and p-Block Elements

- (a)  $H^+$   
 (b)  $OH^-$   
 (c) Only  $NH_4^+$   
 (d)  $OH^-$ ,  $NH_4^+$  and  $NH_4OH$  molecules
34. Among the members of V A group (*N, P, As, Sb* and *Bi*), which of the following properties shows an increase as we go down from nitrogen to bismuth  
 [CBSE PMT 1991]  
 (a) Stability of +3 oxidation state  
 (b) Reducing character of hydrides  
 (c) Electronegativity  
 (d) Acidic nature of the pentoxide
35. The important method for the fixation of nitrogen is  
 [CPMT 1985, 94]  
 (a) Haber (b) Solvay  
 (c) Deacon (d) Fischer method
36. Which of the following is oxidised in air  
 [AFMC 1987; KCET 1991]  
 (a) White phosphorus (b)  $CH_4$   
 (c)  $H_2O$  (d)  $NaCl$
37. A pure sample of nitrogen is prepared by heating  
 (a) Calcium cyanamide (b) Barium azide  
 (c) Ammonium hydroxide (d) Ammonium nitrite
38. Nitrous oxide  
 (a) Is a mixed oxide  
 (b) Is an acidic oxide  
 (c) Is highly soluble in hot water  
 (d) Supports the combustion of sulphur
39. Which of the following represents laughing gas  
 [CPMT 1986, 89; Manipal MEE 1995; MP PMT 1990; MP PET 1995; RPMT 1999; AFMC 2002]  
 (a)  $NO$  (b)  $N_2O$   
 (c)  $NO_2$  (d)  $N_2O_3$
40.  $NO_2$  is a mixed oxide is proved by the first that with  $NaOH$ , it forms  
 (a) Nitrites salt  
 (b) Nitrates salt  
 (c) Mixture of nitrate and nitrite  
 (d) Ammonia
41. Which of the following metal produces nitrous oxide with dil.  $HNO_3$   
 (a) *Fe* (b) *Zn*  
 (c) *Cu* (d) *Ag*
42. Which of the following acid exist in polymeric form  
 (a)  $HPO_3$  (b)  $H_4P_2O_7$   
 (c)  $H_3PO_4$  (d) None of these
43. Superphosphate of lime is [AMU 1985]  
 (a) A mixture of normal calcium phosphate and gypsum  
 (b) A mixture of primary calcium phosphate and gypsum  
 (c) Normal calcium phosphate  
 (d) Soluble calcium phosphate
44. If phosphoric acid is allowed to react with sufficient quantity of  $NaOH$ , the product obtained is  
 [DPMT 1983; MP PMT 1983]  
 (a)  $NaHPO_3$  (b)  $Na_2HPO_4$   
 (c)  $NaH_2PO_4$  (d)  $Na_3PO_4$
45. White phosphorus contains [CPMT 1978; KCET (Med.) 2000; MP PET 1990]  
 (a)  $P_5$  molecules (b)  $P_4$  molecules  
 (c)  $P_6$  molecules (d)  $P_2$  molecules
46. In the catalytic oxidation of ammonia an oxide is formed which is used in the preparation of  $HNO_3$ . This oxide is  
 [CPMT 1984; KCET 1990; AIIMS 1996]  
 (a)  $N_2O_5$  (b)  $N_2O_4$   
 (c)  $NO_2$  (d)  $NO$
47. Nitric acid oxidises phosphorus to [CPMT 1984; JIPMER 2002]  
 (a)  $H_2P_2O_7$  (b)  $H_3PO_3$   
 (c)  $P_2O_5$  (d)  $H_3PO_4$
48. Which one of the following statements is true for  $HNO_2$   
 [CPMT 1980, 84]  
 (a) It is very stable in aqueous solution  
 (b) It cannot act both as an oxidant and as a reductant  
 (c) It cannot act as an oxidising agent  
 (d) It cannot act as reducing agent
49. Which oxide is alkaline [MP PET 1990]  
 (a)  $P_2O_3$  (b)  $Bi_2O_3$   
 (c)  $As_2O_3$  (d)  $B_2O_3$
50. Which acid is formed by  $P_2O_3$  [MP PET 1991]  
 (a)  $H_3PO_4$  (b)  $H_3PO_3$   
 (c)  $HPO_3$  (d)  $H_4P_2O_7$
51. Which nitrogen trihalides is least basic [IIT 1987; Kurukshetra CEE 1998; CPMT 1999]  
 (a)  $NF_3$  (b)  $NCl_3$   
 (c)  $NBr_3$  (d)  $NI_3$
52. Dehydrated phosphorus trichloride in water gives [MP PET 1990]  
 (a)  $HPO_3$  (b)  $H_3PO_4$   
 (c)  $H_3PO_2$  (d)  $H_3PO_3$
53. Which is used in the manufacture of safe matchsticks [DPMT 1982, CPMT 1974, 75]  
 (a) White phosphorus (b) Sulphur  
 (c) Red phosphorus (d) Selenium



54. Which oxide of nitrogen is coloured gas  
[IIT 1987; Kurukshetra CEE 1998]  
(a)  $N_2O$  (b)  $NO$   
(c)  $N_2O_5$  (d)  $NO_2$
55. Which oxide do not act as a reducing agent  
[MP PET 1990]  
(a)  $NO$  (b)  $NO_2$   
(c)  $N_2O$  (d)  $N_2O_5$
56. In  $NH_4NO_2$ , the oxidation number of nitrogen will be  
[MP PET 1990]  
(a) +3 (b) +5  
(c) -3 and +3 (d) +3 and +5
57. In which compound, the oxidation state of phosphorus is +4  
[MP PET 1991]  
(a)  $P_4O_{11}$  (b)  $P_4O_8$   
(c)  $P_4O_6$  (d)  $H_3PO_4$
58. In which compound, the oxidation state of nitrogen is -1  
[MP PMT 1989]  
(a)  $NO$  (b)  $N_2O$   
(c)  $NH_2OH$  (d)  $N_2H_4$
59. Which of the following oxide is least acidic  
[MP PMT 1990; CBSE PMT 1996]  
(a)  $P_4O_6$  (b)  $P_4O_{10}$   
(c)  $As_4O_6$  (d)  $As_4O_{10}$
60. The basic character of hydrides of the V-group elements decreases in the order [CBSE PMT 1996]  
(a)  $SbH_3 > PH_3 > AsH_3 > NH_3$   
(b)  $NH_3 > SbH_3 > PH_3 > AsH_3$   
(c)  $NH_3 > PH_3 > AsH_3 > SbH_3$   
(d)  $SbH_3 > AsH_3 > PH_3 > NH_3$
61. Which is least stable [MP PET 1989]  
(a)  $BiH_3$  (b)  $SbH_3$   
(c)  $AsH_3$  (d)  $PH_3$
62. Which of the following is not hydrolysed [DPMT 2005]  
(a)  $AsCl_3$  (b)  $PF_3$   
(c)  $SbCl_3$  (d)  $NF_3$
63. Electrolysis temperature is maximum for [MP PET 1990]  
(a)  $AsH_3$  (b)  $NH_3$   
(c)  $PH_3$  (d)  $SbH_3$
64. Which of the following is kept in water? [BCECE 2005]  
(a) White phosphorous (b) Sodium  
(c) Potassium (d) Calcium
65. Which of the following substances is used as a fertilizer  
(a)  $K_2SO_4 \cdot Al_2(SO_4)_3 \cdot 24H_2O$   
(b)  $Ca(H_3PO_4)_2 \cdot H_2O + CaSO_4$   
(c)  $NaAlO_2$  (d)  $CaC_2$
66. On adding water to  $BiCl_3$  solution in  $HCl$ , the compound produced is  
(a)  $Bi_2O_3$  (b)  $Bi(OH)_3$   
(c)  $BiOCl$  (d)  $BiOCl_2$
67. V-A group precipitate was dissolved in  $HNO_3$  and treated with excess of  $NH_4OH$ . It gives a white ppt. because of  
(a)  $Cu(OH)_2$  (b)  $Cd(OH)_2$   
(c)  $Bi(OH)_3$  (d)  $Hg(OH)_2$
68. N, P, As, Sb, Bi elements belong to [DPMT 1982]  
(a) VA group (b) IVA group  
(c) VIIA group (d) VB group
69. Which one of the following elements occur free in nature [CPMT 1988]  
(a) Nitrogen (b) Phosphorus  
(c) Arsenic (d) Antimony
70. Which of the following elements of group VA does not show allotropy [CPMT 1980]  
(a) N (b) Bi  
(c) P (d) As
71. Which does not form complex [CPMT 1986]  
(a) N (b) P  
(c) As (d) Bi
72. The strongest base is [IIT 1989; CPMT 1997; MP PET 2001, 03]  
(a)  $NH_3$  (b)  $PH_3$   
(c)  $AsH_3$  (d)  $SbH_3$
73. The most stable hydride is [EAMCET 1988]  
(a)  $NH_3$  (b)  $PH_3$   
(c)  $AsH_3$  (d)  $SbH_3$
74. Which has the lowest boiling point [CBSE PMT 1989]  
(a)  $NH_3$  (b)  $PH_3$   
(c)  $AsH_3$  (d)  $SbH_3$
75. Which is the most explosive [BHU 1984; Roorkee 1989; AIIMS 1996; MP PMT 1985, 2001]  
(a)  $NCl_3$  (b)  $PCl_3$   
(c)  $AsCl_3$  (d) All of these
76. Of the following, the most acidic is [EAMCET 1980]  
(a)  $As_2O_3$  (b)  $P_2O_3$   
(c)  $Sb_2O_3$  (d)  $Bi_2O_3$
77. Of the following, non-existent compound is [NCERT 1975, 79]  
(a)  $PH_4I$  (b)  $As_2O_3$   
(c)  $SbCl_2$  (d)  $As_2H_3$
78. Pure  $N_2$  gas is obtained from [CBSE PMT 1991]  
(a)  $NH_3 + NaNO_2$  (b)  $NH_4Cl + NaNO_2$   
(c)  $N_2O + Cu$  (d)  $(NH_4)_2Cr_2O_7$
79. Pure nitrogen can be prepared from



## 778 s and p-Block Elements

- [KCET 1991; AFMC 1993; AMU 1985]
- (a)  $NH_4OH$  (b)  $Ca_3N_2$   
(c)  $NH_4NO_2$  (d)  $Ba(NO_3)_2$
80. Nitrogen combines with metals to form [CPMT 1981, 93]  
(a) Nitrites (b) Nitrates  
(c) Nitrosyl chloride (d) Nitrides
81. Nitrogen is relatively inactive element because [CBSE PMT 1992]  
(a) Its atom has a stable electronic configuration  
(b) It has low atomic radius  
(c) Its electronegativity is fairly high  
(d) Dissociation energy of its molecule is fairly high
82. The cyanide ion,  $CN^-$  and  $N_2$  are isoelectronic. But in contrast to  $CN^-$ ,  $N_2$  is chemically inert because of [IIT 1992]  
(a) Low bond energy  
(b) Absence of bond polarity  
(c) Unsymmetrical electron distribution  
(d) Presence of more number of electrons in bonding orbitals
83. Which statement is not correct for nitrogen [AIIMS 1991]  
(a) It has a small size  
(b) It does not readily react with  $O_2$   
(c) It is a typical non-metal  
(d)  $d$ -orbitals are available for bonding
84. The element which is essential in nitrogen fixation is [NCERT 1981]  
(a) Zinc (b) Copper  
(c) Molybdenum (d) Boron
85. Laughing gas is prepared by heating [EAMCET 1980]  
(a)  $NH_4Cl$  (b)  $(NH_4)_2SO_4$   
(c)  $NH_4Cl + NaNO_3$  (d)  $NH_4NO_3$
86. Nitrogen (I) oxide is produced by [IIT 1989]  
(a) Thermal decomposition of ammonium nitrate  
(b) Disproportionation of  $N_2O_4$   
(c) Thermal decomposition of ammonium nitrite  
(d) Interaction of hydroxyl amine and nitrous acid
87. Which of the following is not correct for  $N_2O$  [CPMT 1984]  
(a) It is called laughing gas  
(b) It is nitrous oxide  
(c) It is not a linear molecule  
(d) It is least reactive in all oxides of nitrogen
88. Which of the following oxides of nitrogen is the anhydride of nitrous acid  
(a)  $NO$  (b)  $N_2O_3$   
(c)  $N_2O_4$  (d)  $N_2O_5$
89. Which of the following is a true acidic anhydride [NCERT 1977]  
(a)  $CO$  (b)  $NO$   
(c)  $ClO_2$  (d)  $N_2O_5$
90. On strongly heating  $Pb(NO_3)_2$  crystals, the gas formed is [NCERT 1980; CPMT 1997]  
(a)  $NO_2$  (b)  $O_2$   
(c)  $NO_2 + O_2$  (d)  $NO$
91. Nitrogen dioxide is released by heating [AFMC 1992]  
(a)  $Pb(NO_3)_2$  (b)  $KNO_3$   
(c)  $NaNO_2$  (d)  $NaNO_3$
92. Nitric oxide is prepared by the action of  $HNO_3$  on [AFMC 1990]  
(a)  $Fe$  (b)  $Cu$   
(c)  $Zn$  (d)  $Sn$
93. When lightning flash is produced, which gas may form [EAMCET 1992; AFMC 1989]  
(a) Nitrous oxide (b) Nitrogen dioxide  
(c) Dinitrogen pentoxide (d) Nitric oxide
94. Of the following, which has three electron bond in its structure [CPMT 1986]  
(a) Nitrous oxide (b) Nitric oxide  
(c) Dinitrogen trioxide (d) Nitrogen pentoxide
95. Which of the following oxides of nitrogen is neutral [CPMT 1988]  
(a)  $N_2O_5$  (b)  $N_2O_3$   
(c)  $N_2O_4$  (d)  $N_2O$
96. Oxidation of  $NO$  in air produces [KCET 1992]  
(a)  $N_2O$  (b)  $N_2O_3$   
(c)  $NO_2$  (d)  $N_2O_5$
97. The reddish brown coloured gas formed when nitric oxide is oxidised by air is [IIT 1979]  
(a)  $N_2O_5$  (b)  $N_2O_4$   
(c)  $NO_2$  (d)  $N_2O_3$
98. When  $AgNO_3$  is heated strongly, the products formed are [Roorkee 1990]  
(a)  $NO$  and  $NO_2$  (b)  $NO_2$  and  $O_2$   
(c)  $NO_2$  and  $N_2O$  (d)  $NO$  and  $O_2$
99. Which of the nitrates on strong heating leaves the metal as the residue [KCET 1990]  
(a)  $AgNO_3$  (b)  $Pb(NO_3)_2$   
(c)  $Cu(NO_3)_2$  (d)  $Al(NO_3)_3$
- [NCERT 1975; AIIMS 1991]
100. Nitrogen dioxide [KCET 1989]  
(a) Dissolves in water forming nitric acid  
(b) Does not dissolve in water

- (c) Dissolves in water to form nitrous acid and gives off oxygen  
(d) Dissolves in water to form a mixture of nitrous and nitric acids
- 101.** Concentrated nitric acid oxidises cane sugar to  
[CBSE PMT 1991]  
(a)  $CO_2$  and  $H_2O$  (b)  $CO$  and  $H_2O$   
(c)  $CO$ ,  $CO_2$  and  $H_2O$  (d) Oxalic acid and water
- 102.** A mixture of ammonia and air at about  $800^\circ C$  in the presence of *Pt* gauze forms [Pb. CET 1989]  
(a)  $N_2O$  (b)  $NO$   
(c)  $NH_2OH$  (d)  $N_2O_3$
- 103.** Which of the following acid possesses oxidising, reducing and complex forming properties [MNR 1985]  
(a)  $HNO_3$  (b)  $H_2SO_4$   
(c)  $HCl$  (d)  $HNO_2$
- 104.** Nitrogen is essential constituent of all [MP PMT 1990]  
(a) Proteins (b) Fats  
(c) Proteins and fats (d) None of these
- 105.** Ammonia gas can be collected by the displacement of  
[NCERT 1989, 90]  
(a) Conc.  $H_2SO_4$  (b) Brine  
(c) Water (d) Mercury
- 106.** The chemical used for cooling in refrigeration is  
[CPMT 1981, 88]  
(a)  $CO_2$  (b)  $NH_4OH$   
(c)  $NH_4Cl$  (d) Liquid  $NH_3$
- 107.** A hydride of nitrogen which is acidic is  
[NCERT 1978, 80; CPMT 1980; BHU 1986]  
(a)  $NH_3$  (b)  $N_2H_4$   
(c)  $N_2H_2$  (d)  $N_3H$
- 108.**  $PCl_5$  exists but  $NCl_5$  does not because  
[EAMCET 1977, 82]  
(a) Nitrogen has no vacant orbitals  
(b)  $NCl_5$  is unstable  
(c) Nitrogen atom is much smaller  
(d) Nitrogen is highly inert
- 109.** Phosphide ion has the electronic structure similar to that of  
[CPMT 1988]  
(a) Nitride ion (b) Fluoride ion  
(c) Sodium ion (d) Chloride ion
- 110.** Which of the following phosphorus is most stable  
[AFMC 1992]  
(a) Red (b) White  
(c) Black (d) All stable
- 111.** Red phosphorus can be obtained from white phosphorus by  
(a) Heating it with a catalyst in an inert atmosphere  
(b) Distilling it in an inert atmosphere  
(c) Dissolving it in carbon disulphide and crystallizing  
(d) Melting it and pouring the liquid into water
- 112.** Bones glow in the dark because [EAMCET 1980]  
(a) They contain shining material  
(b) They contain red phosphorus  
(c) White phosphorus undergoes slow combustion in contact with air  
(d) White phosphorus changes into red form
- 113.** Which of the following properties of white phosphorus are shared by red phosphorus [NCERT 1973, 74]  
(a) It shows phosphorescence in air  
(b) It reacts with hot aqueous  $NaOH$  to give phosphine  
(c) It dissolves in carbon disulphide  
(d) It burns when heated in air
- 114.** Mixture used for the tips of match stick is [DPMT 1984]  
(a)  $S + K$   
(b)  $Sb_2S_3$   
(c)  $K_2Cr_2O_7 + S + \text{red } P$   
(d)  $K_2Cr_2O_7 + K + S$
- 115.** In modern process phosphorus is manufactured by  
[CPMT 1974, 78, 81]  
(a) Heating a mixture of phosphorite mineral with sand and coke in electric furnace  
(b) Heating calcium phosphate with coke  
(c) Heating bone ash with coke  
(d) Heating the phosphate mineral with sand
- 116.** White phosphorus when boiled with strong solution of caustic soda produces [CPMT 1989, 03]  
(a) Phosphine (b) Phosphoric acid  
(c) Phosphorus acid (d) No reaction
- 117.** White phosphorus reacts with caustic soda. The products are  $PH_3$  and  $NaH_2PO_2$ . This reaction is an example of  
[IIT 1980; KCET 1993]  
(a) Oxidation (b) Reduction  
(c) Oxidation and reduction (d) Neutralisation
- 118.** Phosphine is not obtained by the reaction [MP PMT 1989]  
(a) White *P* is heated with  $NaOH$   
(b) Red *P* is heated with  $NaOH$   
(c)  $Ca_3P_2$  reacts with water  
(d) Phosphorus trioxide is boiled with water
- 119.**  $PH_4I + NaOH$  forms [CBSE PMT 1991]  
(a)  $PH_3$  (b)  $NH_3$   
(c)  $P_4O_6$  (d)  $P_4O_{10}$
- 120.** Phosphine is produced by adding water to [KCET 1991]







## 780 s and p-Block Elements

- (a)  $CaC_2$  (b)  $HPO_3$  (a)  $N_2$  (b)  $HNO_3$   
 (c)  $Ca_3P_2$  (d)  $P_4O_{10}$  (c)  $NH_3$  (d)  $PH_3$
- 121.** When aluminium phosphide is treated with dil. sulphuric acid  
 [KCET 1989]  
 (a)  $SO_2$  is liberated (b)  $PH_3$  is evolved  
 (c)  $H_2S$  is evolved (d)  $H_2$  is evolved
- 122.** With reference to protonic acids, which of the following statements is correct  
 (a)  $PH_3$  is more basic than  $NH_3$   
 (b)  $PH_3$  is less basic than  $NH_3$   
 (c)  $PH_3$  is equally basic as  $NH_3$   
 (d)  $PH_3$  is amphoteric while  $NH_3$  is basic
- 123.** One of the acid listed below is formed from  $P_2O_3$  and the rest are formed from  $P_2O_5$ . The acid formed from phosphorus (III) oxide is  
 (a)  $HPO_3$  (b)  $H_4P_2O_7$   
 (c)  $H_3PO_4$  (d)  $H_3PO_3$
- 124.**  $P_2O_5$  is heated with water to give  
 [CBSE PMT 1991; DPMT 2000]  
 (a) Hypophosphorus acid (b) Orthophosphorus acid  
 (c) Hypophosphoric acid (d) Orthophosphoric acid
- 125.** Hypophosphorus acid is [NCERT 1977; MP PMT 1992]  
 (a) A tribasic acid (b) A dibasic acid  
 (c) A monobasic acid (d) Not acidic at all
- 126.**  $PCl_3$  reacts with water to form  
 [KCET 1990; CBSE PMT 1991; CPMT 2003]  
 (a)  $PH_3$  (b)  $H_3PO_3$ ,  $HCl$   
 (c)  $POCl_3$  (d)  $H_3PO_4$
- 127.**  $H_3PO_3$  is [CPMT 1977, 79, 94; NCERT 1981; MP PMT 1980]  
 (a) A tribasic acid (b) A dibasic acid  
 (c) Neutral (d) A monobasic acid
- 128.** Oxidation state of + 1 for phosphorus is found in [MP PMT 1991; MP PET 2001]  
 (a)  $H_3PO_3$  (b)  $H_3PO_4$   
 (c)  $H_3PO_2$  (d)  $H_4P_2O_7$
- 129.** Which is not an acid salt [MNR 1989]  
 (a)  $NaH_2PO_3$  (b)  $NaH_2PO_2$   
 (c)  $Na_3HP_2O_6$  (d)  $Na_4P_2O_7$
- 130.** By the action of hot conc.  $H_2SO_4$ , phosphorus changes to [Roorkee 1992]  
 (a) Phosphorus acid  
 (b) Orthophosphoric acid  
 (c) Metaphosphoric acid  
 (d) Pyrophosphoric acid
- 131.** Cyanamide process is used in the formation of [BHU 1995]  
 (a)  $N_2$  (b)  $HNO_3$   
 (c)  $NH_3$  (d)  $PH_3$
- 132.** The number of hydroxyl group in pyrophosphoric acid is [KCET 1993]  
 (a) 3 (b) 4  
 (c) 5 (d) 7
- 133.** There is very little difference in acid strength in the series  $H_4PO_4$ ,  $H_3PO_3$ , and  $H_3PO_2$  because [KCET 1990; CPMT 1990]  
 (a) Phosphorus in these acids exists in different oxidation states  
 (b) The hydrogen in these acids are not all bounded to the phosphorus  
 (c) Phosphorus is not a highly electronegative element  
 (d) Phosphorus oxides are less basic
- 134.**  $BiCl_3$  on hydrolysis forms a white precipitate of [NCERT 1975] [CPMT 1988]  
 (a) Bismuthio acid  
 (b) Bismuth oxychloride  
 (c) Bismuth pentachloride  
 (d) Bismuth hydroxide
- 135.** At high temperature nitrogen combines with calcium carbide ( $CaC_2$ ) to give [DPMT 1981, 85; AFMC 1998; MP PET 2000]  
 (a) Calcium cyanide (b) Calcium cyanamide  
 (c) Calcium carbonate (d) Calcium nitride
- 136.** Calcium cyanamide on treatment with steam under pressure gives ammonia and [EAMCET 1984, 88; CPMT 1990]  
 (a) Calcium carbonate (b) Calcium hydroxide  
 (c) Calcium oxide (d) Calcium bicarbonate
- 137.** Which one has the highest percentage of nitrogen [KCET 1991; CBSE PMT 1993; AIIMS 1996, MP PET 2001; RPET 2003]  
 (a) Urea (b) Ammonium sulphate  
 (c) Ammonium nitrate (d) Calcium nitrate
- 138.** Superphosphate of lime contains [CPMT 1984]  
 (a)  $Ca_3(PO_4)_2$  (b)  $CaHPO_4$   
 (c)  $Ca_3(PO_4)_2 + H_3PO_4$  (d)  $Ca(H_2PO_4)_2$
- 139.** Thomas slag is [CPMT 1988]  
 (a)  $Ca_3(PO_4)_2$  (b)  $MnSiO_3$   
 (c)  $CaSiO_3$  (d)  $FeSiO_3$
- 140.** When equal weights of the two fertilizers, urea and ammonium sulphate are taken, urea contains [KCET 1990]  
 (a) Less nitrogen than ammonium sulphate  
 (b) As much nitrogen as ammonium sulphate  
 (c) Twice the amount of nitrogen present in ammonium sulphate  
 (d) More than twice the amount of nitrogen present in ammonium sulphate

141. Which statement is wrong for  $NO$  [DPMT 2005]  
 (a) It is anhydride of nitrous acid  
 (b) Its dipole moment is  $0.22 D$   
 (c) It forms dimer  
 (d) it is paramagnetic
142. The resonance hybrid of nitrate ion is [AFMC 2002]  
 (a)  $O \overset{1/2}{\cdots} \overset{\cdot\cdot}{N} \overset{\cdot\cdot}{\cdots} \overset{-1/2}{O}$  (b)  $O \overset{-2/3}{\cdots} \overset{\cdot\cdot}{N} \overset{\cdot\cdot}{\cdots} \overset{2/3}{O}$   
 (c)  $O \overset{-1/3}{\cdots} \overset{\cdot\cdot}{N} \overset{\cdot\cdot}{\cdots} \overset{-1/3}{O}$  (d)  $O \overset{-2/3}{\cdots} \overset{\cdot\cdot}{N} \overset{\cdot\cdot}{\cdots} \overset{-2/3}{O}$
143. Blasting of TNT is done by mixing [AFMC 1993]  
 (a)  $NH_4Cl$  (b)  $NH_4NO_3$   
 (c)  $NH_4NO_2$  (d)  $(NH_4)_2SO_4$
144. Sodium hydroxide solution reacts with phosphorus to give phosphine. To bring about this reaction, we need [KCET 1989]  
 (a) White phosphorus and dil.  $NaOH$   
 (b) White phosphorus and conc.  $NaOH$   
 (c) Red phosphorus and dil.  $NaOH$   
 (d) Red phosphorus and conc.  $NaOH$
145. Which of the following exhibits highest solubility in water [MP PET 1994]  
 (a)  $NH_3$  (b)  $PH_3$   
 (c)  $AsH_3$  (d)  $SbH_3$
146. Which of the following has highest boiling point [MP PET 1994]  
 (a)  $NH_3$  (b)  $PH_3$   
 (c)  $AsH_3$  (d)  $SbH_3$
147. In the following reaction  
 $P_4 + 3NaOH + 3H_2O \rightarrow PH_3 + 3NaH_2PO_2$  [MP PMT 1994]  
 (a) Phosphorus is oxidised  
 (b) Phosphorus is oxidised and reduced  
 (c) Phosphorus is reduced  
 (d) Sodium is oxidised
148.  $HNO_3$  in aqueous solution yields [AMU 2000]  
 (a)  $NO_3^-$  and  $H^+$  (b)  $NO_3^-$  and  $H_3O^+$   
 (c)  $NO_2^-$  and  $OH^-$  (d)  $N_2O_5$  and  $H_2O$
149. The oxyacid of phosphorus, in which phosphorus has the lowest oxidation state, is [KCET (Med.) 2001]  
 (a) Hypophosphorous acid (b) Orthophosphoric acid  
 (c) Pyrophosphoric acid (d) Metaphosphoric acid
150. Superphosphate is a mixture of [KCET (Med.) 2001]  
 (a)  $Ca(H_2PO_4)_2 \cdot H_2O + CaCl_2 \cdot 2H_2O$   
 (b)  $Ca_3(PO_4)_2 \cdot H_2O + CaCl_2 \cdot 2H_2O$   
 (c)  $Ca_3(PO_4)_2 \cdot H_2O + 2CaSO_4 \cdot 2H_2O$   
 (d)  $Ca(H_2PO_4)_2 \cdot H_2O + 2CaSO_4 \cdot 2H_2O$
151. Solid  $PCl_5$  exists as [JIPMER 2002]  
 (a)  $PCl_5$  (b)  $PCl_4^+$   
 (c)  $PCl_6^-$  (d)  $PCl_4^+$  and  $PCl_6^-$
152. In the reaction,  $P_2O_5 + 3CaO \rightarrow Ca_3(PO_4)_2; P_2O_5$  acts as..... [Orissa JEE 2002]  
 (a) Acidic flux (b) Basic flux  
 (c) Basic impurity (d) Acidic impurity
153. Atoms in a  $P_4$  molecule of white phosphorus are arranged regularly in space in which of the following way [Kerala (Engg.) 2002]  
 (a) At the corners of tetrahedron  
 (b) At the corners of a cube  
 (c) At the corners of a four membered ring  
 (d) At the centre and corners of an equilateral triangle
154. The most common minerals of phosphorus are [Kerala (Med.) 2002]  
 (a) Hydroxy apatite and kernite  
 (b) Colemanite and fluorapatite  
 (c) Borax and fluorapatite  
 (d) Hydroxy apatite and colemanite  
 (e) Hydroxy apatite and fluorapatite
155. The three important oxidation states of phosphorus are [Kerala (Med.) 2002]  
 (a)  $-3, +3$  and  $+5$  (b)  $-3, +3$  and  $-5$   
 (c)  $-3, +4$  and  $-4$  (d)  $-3, +3$  and  $+4$
156. In case of nitrogen,  $NCl_3$  is possible but not  $NCl_5$  while in case of phosphorous,  $PCl_3$  as well as  $PCl_5$  are possible. It is due to  
 (a) Availability of vacant  $d$ -orbital in  $P$  but not in  $N$   
 (b) Lower electronegativity of  $P$  than  $N$   
 (c) Lower tendency of  $H$  bond formation in  $P$  than  $N$   
 (d) Occurrence of  $P$  in solid while  $N$  in gaseous state at room temperature
157. When ammonia is passed over heated copper oxide, the metallic copper is obtained. the reaction shows that ammonia is  
 (a) A dehydrating agent (b) An oxidising agent  
 (c) A reducing agent (d) A nitrating agent
158. Urea is preferred to ammonium sulphate as a nitrogenous fertilizer because [KCET 2003]  
 (a) It is more soluble in water  
 (b) It is cheaper than ammonium sulphate

## 782 s and p-Block Elements

- (c) It is quite stable  
(d) It does not cause acidity in the soil
159. Liquid ammonia is used for refrigeration because [MP PET 2002]  
(a) It has a high dipole moment  
(b) It has a high heat of vapourisation  
(c) It is basic  
(d) It is a stable compound
160. Action of concentrated nitric acid ( $HNO_3$ ) on metallic tin produces [BHU 2002]  
(a) Stannic nitrate (b) Stannous nitrate  
(c) Stannous nitrite (d) Meta stannic acid
161. How can you synthesize nitric oxide in the laboratory [Orissa JEE 2003]  
(a) Zinc with cold and dilute  $HNO_3$   
(b) Zinc with concentrated  $HNO_3$   
(c) Copper with cold and dilute  $HNO_3$   
(d) Heating  $NH_4NO_3$
162. What would happen when a solution of potassium chromate is treated with an excess of dilute nitric acid [AIEEE 2003]  
(a)  $Cr^{3+}$  and  $Cr_2O_7^{2-}$  are formed  
(b)  $Cr_2O_7^{2-}$  and  $H_2O$  are formed  
(c)  $CrO_4^{2-}$  is reduced to +3 state of Cr  
(d)  $CrO_4^{2-}$  is oxidized to +7 state of Cr
163. The pentavalence in phosphorus is more stable as compared to that of nitrogen even though they belong to the same group. It is due to  
(a) Inert nature of nitrogen  
(b) Reactivity of phosphorus  
(c) Larger size of phosphorus atom  
(d) Dissimilar electronic configuration
164. A neutral fertilizer among the following compounds is [KCET 2002]  
(a) Urea  
(b) Ammonium nitrate  
(c) Ammonium sulphate  
(d) Calcium ammonium nitrate
165. Which is true with regard to the properties of  $PH_3$  [BHU 2000]  
(a)  $PH_3$  is not much stable  
(b)  $PH_3$  is neutral towards litmus  
(c)  $PH_3$  has fishy smell  
(d)  $PH_3$  is insoluble in water
166. Nitrogen is obtained when  $NaNO_2$  reacts with [UPSEAT 2003]  
(a)  $NH_4Cl$  (b)  $NH_4NO_3$   
(c)  $(NH_4)_2CO_3$  (d)  $NH_4OH$
167.  $N_2$  combines with metal to form [JIPMER 2000]  
(a) Nitride (b) Nitrate  
(c) Nitrite (d) Nitrosyl chloride
168. The number of  $P-O-P$  bridges in the structure of phosphorous pentoxide and phosphorous trioxide are respectively [AIIMS 2005]  
(a) 6, 6 (b) 5, 5  
(c) 5, 6 (d) 6, 5
169. Sodium pyrophosphate is [CPMT 2003]  
(a)  $Na_2P_2O_7$  (b)  $Na_4P_2O_7$   
(c)  $NaPO_4$  (d)  $Na_2PO_2$
170. Which of the following is solid in nature [UPSEAT 2003; AFMC 2004]  
(a)  $NO$  (b)  $N_2O$   
(c)  $N_2O_3$  (d)  $N_2O_5$
171. Which of the following is a cyclic phosphate [KCET 1996]  
(a)  $H_5P_3O_{10}$  (b)  $H_6P_4O_{13}$   
(c)  $H_5P_5O_{15}$  (d)  $H_7P_5O_{16}$
172. The reaction, which forms nitric oxide, is [KCET (Med.) 2001]  
(a) C and  $N_2O$  (b) Cu and  $N_2O$   
(c) Na and  $NH_3$  (d) Cu and  $HNO_3$
173. Which one of the following can be used as an anaesthetic [EAMCET 1998]  
(a)  $N_2O$  (b)  $NO$   
(c)  $NCl_3$  [KCET 2002] (d)  $NO_2$
174. Solution of sodium metal in liquid ammonia is strongly reducing due to the presence in the solution of the following [KCET 2000; MP PMT 2001]  
(a) Sodium hydride (b) Sodium amide  
(c) Sodium atoms (d) Solvated electrons
175. What may be expected to happen when phosphine gas is mixed with chlorine gas [AIEEE 2003]  
(a) The mixture only cools down  
(b)  $PCl_3$  and  $HCl$  are formed and the mixture warms up  
(c)  $PCl_5$  and  $HCl$  are formed and the mixture cools down  
(d)  $PH_3.Cl_2$  is formed with warming up
176.  $P_4O_{10}$  is not used with to dry  $NH_3$  gas because [KCET 2001]  
(a)  $P_4O_{10}$  is basic and  $NH_3$  is acidic  
(b)  $P_4O_{10}$  is acidic and  $NH_3$  is basic  
(c)  $P_4O_{10}$  is not a drying agent  
(d)  $P_4O_{10}$  reacts with moisture in  $NH_3$

177. When ammonia reacts with sodium hypochlorite, product containing nitrogen is [AFMC 2000]  
 (a)  $N_2$  (b)  $N_2O$   
 (c)  $NH_2OH$  (d)  $H_2N.NH_2$
178.  $P_2O_5$  is used extensively as a [BVP 2003]  
 (a) Reducing agent (b) Oxidising agent  
 (c) Dehydrating agent (d) Preservative
179. Inertness of  $N_2$  gas is due to [DCE 2000; MP PET 2001]  
 (a) No vacant  $d$ -orbital  
 (b) High dissociation energy  
 (c) High electronegativity  
 (d) None
180. Which show maximum valency [CPMT 2003]  
 (a) Phosphorus (b) Tin  
 (c) Antimony (d) Bismuth
181. Which is used in the Haber process for the manufacture of  $NH_3$  [MH CET 2001]  
 (a)  $Pt$  (b)  $Fe + Mo$   
 (c)  $CuO$  (d)  $Al_2O_3$
182. On adding excess of ammonium hydroxide to a copper chloride solution.  
 (a) A deep blue solution is obtained  
 (b) No change is observed  
 (c) Blue precipitate of copper hydroxide is obtained  
 (d) Black precipitate of copper oxide is obtained
183. The product obtained by heating  $(NH_4)_2SO_4$  and  $KCNO$  is [DPMT 2000]  
 (a) Hydrocyanic acid (b) Ammonia  
 (c) Ammonium cyanide (d) Urea
184. The number of  $P-O-P$  bonds in cyclic metaphosphoric acid is [IIT-JEE (Screening) 2000]  
 (a) Zero (b) Two  
 (c) Three (d) Four
185. When  $HNO_3$  is dropped into the palm and washed with water, it turns into yellow. It shows the presence of [CPMT 1997]  
 (a)  $NO_2$  (b)  $N_2O$   
 (c)  $NO$  (d)  $N_2O_5$
186. Which of the following is nitrogenous fertilizers [CPMT 1999]  
 (a) Bone meal (b) Thomas meal  
 (c) Nitro phosphate (d) Ammonium sulphate
187. Which compound is related to Haber's process [RPET 1999]  
 (a)  $CO_2$  (b)  $H_2$   
 (c)  $NO_2$  (d)  $NH_3$
188. Ammonia is dried over [CPMT 2002; JIPMER 2002]  
 (a) Quick lime (b) Slaked lime  
 (c) Anhy.  $CaCl_2$  (d) None of these
189. Which of the following compounds is sparingly soluble in ammonia [JIPMER 1999]  
 (a)  $AgI$  (b)  $AgBr$   
 (c)  $AgCl$  (d)  $CuCl_2$
190. The carbonate which does not leave a residue on heating is [JIPMER 1999; DCE 1999]  
 (a)  $Na_2CO_3$  (b)  $Ag_2CO_3$   
 (c)  $CuCO_3$  (d)  $(NH_4)_2CO_3$
191. Which of the following has the highest dipole moment [CBSE PMT 1997]  
 (a)  $NH_3$  (b)  $PH_3$   
 (c)  $SbH_3$  (d)  $AsH_3$
192. The structural formula of hypophosphorus acid is [CBSE PMT 1997; AIIMS 2001; BCECE 2005; Pb. CET 2002]  
 (a)  (b)   
 (c)  [MH CET 1999] (d) 
193. Repeated use of which of the following fertilizers would increase the acidity of the soil [CBSE PMT 1998]  
 (a) Urea (b) Potassium nitrate  
 (c) Ammonium sulphate (d) Superphosphate of lime
194. Aqua-regia is [Orissa JEE 2005]  
 (a) 1 : 3 conc.  $HNO_3$  and conc.  $HCl$   
 (b) 1 : 2 conc.  $HNO_3$  and conc.  $HCl$   
 (c) 3 : 1 conc.  $HNO_3$  and conc.  $HCl$   
 (d) 2 : 1 conc.  $HNO_3$  and conc.  $HCl$
195. Which oxide of nitrogen is obtained on heating ammonium nitrate at  $250^\circ C$   
 (a) Nitric oxide (b) Nitrous oxide  
 (c) Nitrogen dioxide (d) Dinitrogen oxide
196. The oxidation number of phosphorus vary from [Kurukshetra CEE 1998; DCE 2001]  
 (a) -1 to +3 (b) -3 to +3  
 (c) -3 to +5 (d) -5 to +1
197. Inorganic graphite is [KCET 2003]  
 (a)  $B_3N_3H_6$  (b)  $B_3N_3$   
 (c)  $SiC$  (d)  $Fe(CO)_5$
198. Conc.  $HNO_3$  can be stored in container of [MH CET 2001]  
 (a)  $Al$  (b)  $Sn$   
 (c)  $Cu$  (d)  $Zn$
199. Which of the following compounds does not exist [JIPMER 1997]  
 (a)  $SbCl_3$  (b)  $BiCl_5$   
 (c)  $SbCl_5$  (d)  $AsCl_5$

## 784 s and p-Block Elements

200. The formula of an oxyacid of phosphorus is  $H_3PO_4$ . It is a  
[MP PMT 1996; CPMT 1999; J & K CET 2005]  
(a) Dibasic acid (b) Monobasic acid  
(c) Tribasic acid (d) Tetrabasic acid
201. Which salt can be classified as an acid salt [CPMT 1993]  
(a)  $Na_2SO_4$  (b)  $BiOCl$   
(c)  $Pb(OH)Cl$  (d)  $Na_2HPO_4$
202. In  $NH_3$  and  $PH_3$ , the common is [AFMC 1995]  
(a) Odour (b) Combustibility  
(c) Basic nature (d) None of these
203. Which one of the following hydrides is least stable  
(a)  $AsH_3$  (b)  $SbH_3$   
(c)  $NH_3$  (d)  $PH_3$
204. Which element from V group, gives most basic compound with hydrogen  
(a) Nitrogen (b) Bismuth  
(c) Arsenic (d) Phosphorus
205. The acid which forms two series of salts is [KCET 1996]  
(a)  $H_3PO_4$  (b)  $H_3PO_3$   
(c)  $H_3BO_3$  (d)  $H_3PO_2$
206. Which gas is obtained when urea is heated with  $HNO_2$   
[CPMT 1996]  
(a)  $N_2$  (b)  $H_2$   
(c)  $O_2$  (d)  $NH_3$
207. Atomic number of  $N$  is 7. The atomic number of IIIrd member of nitrogen family is [CPMT 1996]  
(a) 23 (b) 15  
(c) 33 (d) 43
208. Which of the following have least covalent  $P-H$  bond  
[CPMT 1996]  
(a)  $PH_3$  (b)  $P_2H_6$   
(c)  $P_2H_5$  (d)  $PH_6^+$
209. Sides of match box have coating of [BHU 1995]  
(a) Potassium chlorate, red lead  
(b) Potassium chlorate, antimony sulphide  
(c) Antimony sulphide, red phosphorus  
(d) Antimony sulphide, red lead
210. Which of the following compound is tribasic acid  
[AIIMS 2002]  
(a)  $H_3PO_2$  (b)  $H_3PO_3$   
(c)  $H_3PO_4$  (d)  $H_4P_2O_7$
211. Which of the following is manufactured from the molecular nitrogen by bacteria [MP PET 1999]  
(a)  $NO_3$  (b)  $NO_2$   
(c) Amino acids (d) Ammonia
212. One mole of calcium phosphide on reaction with excess water gives [IIT-JEE 1999]  
(a) One mole of phosphine  
(b) Two moles of phosphoric acid  
(c) Two moles of phosphine  
(d) One mole of phosphorous pentoxide
213. On heating ammonium dichromate, the gas evolved is [IIT-JEE 1999]  
(a) Oxygen (b) Ammonia  
(c) Nitrous oxide (d) Nitrogen
214. In compounds of type  $ECl_3$ , where  $E = B, P, As$  or  $Bi$ , the angles  $Cl-E-Cl$  for different  $E$  are in the order [IIT-JEE 1999]  
(a)  $B > P = As = Bi$  (b)  $B > P > As > Bi$   
(c)  $B < P = As = Bi$  (d)  $B < P < As < Bi$
215. Ammonia on reaction with hypochlorite anion, can form [IIT-JEE 1999]  
(a)  $NO$  (b)  $NH_4Cl$   
(c)  $N_2H_4$  (d)  $HNO_2$
216. Orthophosphoric acid represents the molaysis condition due to  
(a) Hydrogen bonding  
(b) Phosphorous group  
(c) Maximum oxygen group  
(d) Tribasicity
217. Which of the following elements forms a strongly acidic oxide  
(a)  $P$  (b)  $As$   
(c)  $Sb$  (d)  $Bi$
218. In group V-A of the periodic table nitrogen forms only a trihalide but other elements form pentahalides also. The reason is  
(a) Nitrogen has less affinity towards halogens  
(b) Nitrogen halides are covalent  
(c) Nitrogen undergoes  $d^2sp^3$  hybridization  
(d) Nitrogen does not have  $d$ -orbitals
219. In the nitrogen family the  $H-M-H$  bond angle in the hydrides  $MH_3$  gradually becomes closer to  $90^\circ$  on going from  $N$  to  $Sb$ . This shows that gradually [MP PET/PMT 1998; MP PMT 2000]  
(a) The basic strength of hydrides increases  
(b) Almost pure  $p$ -orbitals are used for  $M-H$  bonding  
(c) The bond energies of  $M-H$  bond increase  
(d) The bond pairs of electrons become nearer to the central atom
220. An element ( $X$ ) forms compounds of the formula  $XCl_3$ ,  $X_2O_5$  and  $Ca_3X_2$ , but does not form  $XCl_5$ , which of the following is the element  $X$  [MP PET 1997]  
(a)  $B$  (b)  $Al$



- (c) N (d) P
221. Which of the following tendencies remains unchanged on going down in the nitrogen family (Group-VA) ? [MP PMT 1997]
- (a) Highest oxidation state  
(b) Non-metallic character  
(c) Stability of hydrides  
(d) Physical state
222. Which of the following oxy acids of phosphorus is a reducing agent and monobasic
- (a)  $H_3PO_2$  (b)  $H_3PO_3$   
(c)  $H_3PO_4$  (d)  $H_4P_2O_6$
223. Bone black is a polymorphic form of [DCE 2003]
- (a) Phosphorus (b) Sulphur  
(c) Carbon (d) Nitrogen
224. Nitrous oxide is known as [AFMC 2004]
- (a) Breathing gas (b) Laughing gas  
(c) exercising gas (d) Laboratory gas
225. When lead nitrate is heated, it gives [MH CET 2003]
- (a)  $NO_2$  (b)  $NO$   
(c)  $N_2O_5$  (d)  $N_2O$
226. Which element exist as a solid at  $25^\circ C$  and 1 atmospheric pressure among the following [DCE 2003]
- (a) Br (b) Cl  
(c) Hg (d) P
227. In the reaction  $HNO_3 + P_4O_{10} \rightarrow 4HPO_3 + x$ , the product x is [MH CET 2003; DPMT 2004]
- (a)  $N_2O_3$  (b)  $N_2O_5$   
(c)  $NO_2$  (d)  $H_2O$
228. The number of hydrogen atom (s) attached to phosphorus atom in hypophosphorous acid is [AIEEE 2005]
- (a) Zero (b) Two  
(c) One (d) Three
229. Which blue liquid is obtained on reacting equimolar amounts of two gases at  $-30^\circ C$  [IIT-JEE (Screening) 2005]
- (a)  $N_2O$  (b)  $N_2O_3$   
(c)  $N_2O_4$  (d)  $N_2O_5$
230. Which is the most thermodynamically stable allotropic form of phosphorus [IIT-JEE (Screening) 2005]
- (a) Red (b) White  
(c) Black (d) Yellow
231.  $(NH_4)_2Cr_2O_7$  on heating liberates a gas. The same gas will be obtained by [IIT JEE (Screening) 2004; BVP 2004]
- (a) Heating  $NH_4NO_2$   
(b) Heating  $NH_4NO_3$   
(c) Treating  $H_2O_2$  with  $NaNO_2$   
(d) Treating  $Mg_3N_2$  with  $H_2O$
232. The element which forms oxides in all oxidation states +I to +V is [AIIMS 2004]
- (a) N (b) P  
(c) As (d) Sb
233. The boiling points of the following hydrides follow the order of [DPMT 2004]
- (a)  $NH_3 > AsH_3 > PH_3 > SbH_3$   
(b)  $SbH_3 > AsH_3 > PH_3 > NH_3$   
(c)  $SbH_3 > NH_3 > AsH_3 > PH_3$   
(d)  $NH_3 > PH_3 > AsH_3 > SbH_3$
234. Which of the following halides is most acidic [MP PMT 2004]
- (a)  $PCl_3$  (b)  $BiCl_3$   
(c)  $SbCl_3$  (d)  $CCl_4$
235. In the electrothermal process, the compound displaced by silica from calcium phosphate is [KCET 2004]
- (a) Calcium (b) Phosphine  
(c) Phosphorus (d) Phosphorus pentoxide
236. Which of the following compound show sublimation [AFMC 1995; Pb. CET 2000]
- (a)  $NH_4Cl$  (b)  $CaCO_3$   
(c)  $BaSO_4$  (d)  $CaHPO_3$
237. Number of P-O bonds in  $P_4O_{10}$  is [DCE 2002]
- (a) 17 (b) 16  
(c) 15 (d) 6
238. Most acidic oxide is [Pb. CET 2004]
- (a)  $Na_2O$  (b)  $ZnO$   
(c)  $MgO$  (d)  $P_2O_5$
239. Which of the following is a mixed anhydride [Pb. CET 2003]
- (a)  $NO$  (b)  $NO_2$   
(c)  $N_2O_5$  (d)  $N_2O$
240. Oxidation number of As in  $H_2AsO_4^-$  is [CPMT 2001]
- (a) 6 (b) 7  
(c) 5 (d) 9
241. When plants and animals decay, the organic nitrogen is converted into inorganic nitrogen. The inorganic nitrogen is in the form of [KCET 2005]
- (a) Ammonia (b) Elements of nitrogen  
(c) Nitrates (d) Nitrides

### Oxygen family

1. Which element is found in free state [CPMT 1972, 81, 91; DPMT 1986]
- (a) Iodine (b) Sulphur  
(c) Phosphorus (d) Magnesium

## 786 s and p-Block Elements

2. Which of the elements listed below occurs in allotropic forms  
[CPMT 1972]  
(a) Iodine (b) Copper  
(c) Sulphur (d) Silver
3. Which forms new compound in air [AFMC 1987]  
(a)  $H_2O$  in air (b)  $O_2$  in air  
(c)  $N_2$  in air (d) Phosphorus in air
4. Which of the following after burning at room temperature gives gaseous oxide  
[NCERT 1973; CPMT 1981; DPMT 1982; JIPMER 2001]  
(a)  $H$  (b)  $Na$   
(c)  $S$  (d)  $He$
5. Sulphur molecule is converted into sulphur ion, when it  
[DPMT 1980]  
(a) Gains two electrons (b) Loses two electrons  
(c) Gains two protons (d) Shares two electrons
6. The element which liberates oxygen gas from water is  
[MP PMT 1993]  
(a)  $P$  (b)  $Na$   
(c)  $F$  (d)  $I$
7. The highest catenation ability is shown by  
[AIIMS 1983; MP PET 1993; CPMT 1997]  
(a) Oxygen (b) Sulphur  
(c) Selenium (d) Tellurium
8. Ozone belongs to which group of the periodic table  
(a) V group (b) VI group  
(c) VII group (d) None of these
9. The number of unpaired electrons in the  $p$ -subshell of oxygen atom is  
(a) 1 (b) 2  
(c) 3 (d) 4
10. Most abundant element on earth is [MP PET/PMT 1988]  
(a)  $O$  (b)  $S$   
(c)  $Se$  (d)  $Te$
11. Which of the following is most electronegative [BHU 1978]  
(a)  $O$  (b)  $S$   
(c)  $Te$  (d)  $Se$
12. The ability of a substance to assume two or more crystalline structures is called  
(a) Isomerism (b) Polymorphism  
(c) Isomorphism (d) Amorphism
13.  $SO_2$  is obtained when [Roorkee 1995]  
(a) Oxygen reacts with dilute sulphuric acid  
(b) Hydrolysis of dilute  $H_2SO_4$   
(c) Concentrated  $H_2SO_4$  reacts with  $Na_2SO_3$   
(d) All of these
14. Which shows polymorphism [BHU 1982; MP PMT 1985]  
(a)  $O$  (b)  $S$   
(c)  $Se$  (d) All the above
15. Bond angle is minimum for [DPMT 1990]  
(a)  $H_2O$  (b)  $H_2S$   
(c)  $H_2Se$  (d)  $H_2Te$
16. Oxygen was discovered by [BHU 1987]  
(a) Priestley (b) Boyle  
(c) Scheele (d) Cavandish
17. The compound which gives off oxygen on moderate heating is  
[IIT-JEE 1986; MP PMT 1991; MADT Bihar 1995]  
(a) Cupric oxide  
(b) Mercuric oxide  
(c) Zinc oxide  
(d) Aluminium oxide
18. It is possible to obtain oxygen from air by fractional distillation because  
(a) Oxygen is in a different group of the periodic table from nitrogen  
(b) Oxygen is more reactive than nitrogen  
(c) Oxygen has higher b.p. than nitrogen  
(d) Oxygen has a lower density than nitrogen
19. Oxygen is denser than air so it is collected over  
[CPMT 1980; MP PET 1999]  
(a)  $H_2O$  (b) Ethanol  
(c) Mercury (d) Kerosene oil
20. Oxygen molecule exhibits  
[CPMT 1991, 99, 2002; AIIMS 1983; BHU 1986; NCERT 1980, 81; MP PMT 1985, 92; AFMC 2004]  
(a) Paramagnetism (b) Diamagnetism  
(c) Ferromagnetism (d) Ferrimagnetism
21. When oxygen is passed through a solution of  $Na_2SO_3$  we get  
(a)  $Na_2SO_4$  (b)  $Na_2S$   
(c)  $NaHSO_4$  (d)  $NaH$
22. Oxygen does not react with [CBSE PMT 1989]  
[Kurukshetra CEE 1991]  
(a)  $P$  (b)  $Na$   
(c)  $S$  (d)  $Cl$
23. The formula of ozone is  $O_3$ , it is  
[CPMT 1989, 91; Manipal MEE 1995; RPET 1999, 2000]  
(a) An allotrope of oxygen (b) Compound of oxygen  
(c) Isotope of oxygen (d) None of these
24. Ozone is obtained from oxygen [CPMT 1982]  
(a) By oxidation at high temperature  
(b) By oxidation using a catalyst  
(c) By silent electric discharge  
(d) By conversion at high pressure
25. Which of the following statement is true about ozone layer  
[NCERT 1980]  
(a) It is harmful because ozone is dangerous to living organism  
(b) It is beneficial because oxidation reaction can proceed faster in the presence of ozone  
(c) It is beneficial because ozone cuts out the ultraviolet radiation of the sun

- (d) It is harmful because ozone cuts out the important radiation of the sun which are vital for photosynthesis
26. Identify the incorrect statement with respect to ozone  
[AIIMS 1992]
- (a) Ozone is formed in the upper atmosphere by a photochemical reaction involving dioxygen  
(b) Ozone is more reactive than oxygen  
(c) Ozone is diamagnetic whereas dioxygen is paramagnetic  
(d) Ozone protects the earth's inhabitants by absorbing  $\gamma$  radiations
27. Which one of the following property is not correct for ozone  
[CPMT 1984]
- (a) It oxidises lead sulphide  
(b) It oxidises potassium iodide  
(c) It oxidises mercury  
(d) It cannot act as bleaching agent
28. Ozone with  $KI$  solution produces [CPMT 1987]
- (a)  $Cl_2$  (b)  $I_2$   
(c)  $HI$  (d)  $IO_3$
29. The gases respectively absorbed by alkaline pyrogallol and oil of cinnamon is [CBSE PMT 1989]
- (a)  $O_3$ ,  $CH_4$  (b)  $O_2$ ,  $O_3$   
(c)  $SO_2$ ,  $CH_4$  (d)  $N_2O$ ,  $O_3$
30. Ozone turns trimethyl paper [CPMT 1989]
- (a) Green (b) Violet  
(c) Red (d) Black
31. No. of atoms in one molecule of sulphur is [AFMC 1987, 91; AMU 1985]
- (a) 8 (b) 4  
(c) 3 (d) None of these
32. When  $H_2S$  is passed through acidified  $KMnO_4$ , we get [CPMT 1979, 91; MP PMT 1987]
- (a)  $K_2SO_3$  (b)  $MnO_2$   
(c)  $KHSO_3$  (d) Sulphur
33. Copper turnings when heated with concentrated sulphuric acid will give [AFMC 1987; BHU 1999; CBSE PMT 2000]
- (a)  $SO_2$  (b)  $SO_3$   
(c)  $H_2S$  (d)  $O_2$
34. Which of the following is used to absorb sulphur dioxide [EAMCET 1980]
- (a) Conc.  $H_2SO_4$  (b)  $KOH$  solution  
(c) Water (d) Anhydrous  $CaCl_2$
35. Which compound acts as an oxidising as well as reducing agent [IIT 1991]
- (a)  $SO_2$  (b)  $MnO_2$   
(c)  $Al_2O_3$  (d)  $CrO_3$
36. A solution of sulphur dioxide in water reacts with  $H_2S$  precipitating sulphur. Here sulphur dioxide acts as [NCERT 1980; MP PMT 1994]
- (a) An oxidising agent (b) A reducing agent  
(c) An acid (d) A catalyst
37. When  $SO_2$  is passed through acidified  $K_2Cr_2O_7$  solution [CPMT 1989, 94]
- (a) The solution turns blue  
(b) The solution is decolourised  
(c)  $SO_2$  is reduced  
(d) Green  $Cr_2(SO_4)_3$  is formed
38. When  $SO_2$  is passed through cupric chloride solution [CPMT 1979, 81, 89, 94]
- (a) A white precipitate is obtained  
(b) The solution becomes colourless  
(c) The solution becomes colourless and a white precipitate of  $Cu_2Cl_2$  is obtained  
(d) No visible change takes place
39. Which of the following is oxidised by  $SO_2$  [BHU 1986]
- (a)  $Mg$  (b)  $K_2Cr_2O_7$   
(c)  $KMnO_4$  (d) All of these
40. Bleaching action of  $SO_2$  is due to [CPMT 1971, 79, 86]
- (a) Reduction (b) Oxidation  
(c) Hydrolysis (d) Its acidic nature
41. A salt of sulphurous acid is called [NCERT 1978]
- (a) Sulphate (b) Sulphurate  
(c) Sulphite (d) Sulphide
42. Which of the following is acidic [AFMC 1990; JIPMER 1997]
- (a)  $SO_3$  (b)  $N_2O$   
(c)  $BeO$  (d)  $HgO$
43. The final acid obtained during the manufacture of  $H_2SO_4$  by contact process is
- (a)  $H_2SO_4$  (conc.) (b)  $H_2SO_4$  (dil.)  
(c)  $H_2SO_4$  (d)  $H_2S_2O_7$
44. About  $H_2SO_4$  which is incorrect [CPMT 1985]
- (a) Reducing agent (b) Dehydrating agent  
(c) Sulphonating agent (d) Highly viscous
45. In the reaction  
 $2Ag + 2H_2SO_4 \rightarrow Ag_2SO_4 + 2H_2O + SO_2$   
 $H_2SO_4$  acts as [CPMT 1981]
- (a) Reducing agent (b) Oxidising agent  
(c) Catalytic agent (d) Dehydrating agent
46. In the reaction  
 $HCOOH \xrightarrow{H_2SO_4} CO + H_2O$ ;  $H_2SO_4$  acts as
- (a) Dehydrating agent (b) Oxidising agent  
(c) Reducing agent (d) All of these



## 788 s and p-Block Elements

47. When conc.  $H_2SO_4$  comes in contact with sugar, it becomes black due to  
[CPMT 1989; BHU 1986; MDAT Bihar 1980]  
(a) Hydrolysis (b) Hydration  
(c) Decolourisation (d) Dehydration
48. Oxalic acid when heated with conc.  $H_2SO_4$ , gives out  
[DPMT 1981; AFMC 1998]  
(a)  $H_2O$  and  $CO_2$  (b)  $CO$  and  $CO_2$   
(c) Oxalic sulphate (d)  $CO_2$  and  $H_2S$
49. Which one is known as **oil of vitriol**  
[CPMT 1988; MP PMT 2004]  
(a)  $H_2SO_3$  (b)  $H_2SO_4$   
(c)  $H_2S_2O_7$  (d)  $H_2S_2O_8$
50. The acid used in lead storage cells is  
[NCERT 1971; Roorkee 1989]  
(a) Phosphoric acid (b) Nitric acid  
(c) Sulphuric acid (d) Hydrochloric acid
51. Which one of the gas dissolves in  $H_2SO_4$  to give oleum  
[CPMT 1988]  
(a)  $SO_2$  (b)  $H_2S$   
(c)  $S_2O$  (d)  $SO_3$
52. Oleum is [CBSE PMT 1991; MP PMT 2002; CPMT 2004]  
(a) Castor oil (b) Oil of vitriol  
(c) Fuming  $H_2SO_4$  (d) None of them
53. There is no S-S bond in  
[IIT 1991; CPMT 1999; DCE 2000]  
(a)  $S_2O_4^{2-}$  (b)  $S_2O_5^{2-}$   
(c)  $S_2O_3^{2-}$  (d)  $S_2O_7^{2-}$
54. Which of the following sulphate is insoluble in water  
[MP PMT 2000]  
(a)  $CuSO_4$  (b)  $CdSO_4$   
(c)  $PbSO_4$  (d)  $Bi_2(SO_4)_3$
55. When sulphur is boiled with  $Na_2SO_3$  solution, the compound formed is [CPMT 1979; Roorkee 1992]  
(a) Sodium sulphide (b) Sodium sulphate  
(c) Sodium persulphate (d) Sodium thiosulphate
56. The products of the chemical reaction between  $Na_2S_2O_3$ ,  $Cl_2$  and  $H_2O$  are [EAMCET 1989]  
(a)  $S + HCl + Na_2S$  (b)  $S + HCl + Na_2SO_4$   
(c)  $S + HCl + Na_2SO_3$  (d)  $S + NaClO_3 + H_2O$
57. Hypo is used in photography for [Roorkee 1989]  
(a) Developing a picture  
(b) Picture printing  
(c) The colour of picture  
(d) The fixation of the picture
58. Hypo is used in photography to  
[CBSE PMT 1988; Pb. CET 1989]  
(a) Reduce  $AgBr$  grains to metallic silver  
(b) Convert the metallic silver to silver salt  
(c) Remove undecomposed silver bromide as a soluble complex  
(d) Remove reduced silver
59. Hypo is used in photography because of its  
[IIT 1981; EAMCET 1988; MADT Bihar 1995]  
(a) Reducing behaviour  
(b) Oxidising behaviour  
(c) Complex forming behaviour  
(d) Reaction with light
60. Aqueous solutions of hydrogen sulphide and sulphur dioxide when mixed together, yield [KCET 2002]  
(a) Sulphur and water  
(b) Sulphur trioxide and water  
(c) Hydrogen peroxide and sulphur  
(d) Hydrogen and sulphurous acid
61. An example of a neutral oxide is [KCET 2003]  
(a)  $NO$  (b)  $CO_2$   
(c)  $CaO$  (d)  $ZnO$
62. Which of the following is the best scientific method to test presence of water in a liquid [AIIMS 1999]  
(a) Taste  
(b) Smell  
(c) Use of litmus paper  
(d) Use of anhydrous copper sulphate
63.  $H_2S$  react with  $O_2$  to form [AFMC 1995]  
(a)  $H_2O + S$  (b)  $H_2O + SO_2$   
(c)  $H_2O + SO_3$  (d)  $H_2SO_4 + S$
64. Which of the following mixture is chromic acid [Pb. PMT 2000]  
(a)  $K_2Cr_2O_7$  and conc.  $H_2SO_4$   
(b)  $K_2Cr_2O_7$  and  $HCl$   
(c)  $K_2SO_4$  and conc.  $H_2SO_4$   
(d)  $H_2SO_4$  and  $HCl$
65. At room temperature  $H_2O$  is a liquid while  $H_2S$  is a gas. The reason is [RPET 1999]  
(a) Electronegativity of O is greater than S  
(b) Difference in the bond angles of both the molecules  
(c) Association takes place in  $H_2O$  due to H-bonding while no H-bonding in  $H_2S$   
(d) O and S belong to different periods
66. Which of the following products is formed on boiling tin with an alkali solution [Roorkee 2000]  
(a)  $Sn(OH)_2$  (b)  $Sn(OH)_4$   
(c)  $SnO_3^{2-}$  (d)  $SnO_2$
67. Amongst  $H_2O$ ,  $H_2S$ ,  $H_2Se$  and  $H_2Te$  the one with the highest boiling point is [IIT-JEE (Screening) 2000]  
(a)  $H_2O$  because of hydrogen bonding  
(b)  $H_2Te$  because of higher molecular weight  
(c)  $H_2S$  because of hydrogen bonding

- (d)  $H_2Se$  because of lower molecular weight
68. Among the hydrides formed by the group VI-A elements, only  $H_2O$  has an abnormally low volatility (high boiling point). This is so because
- $H_2O$  molecules are associated due to intermolecular hydrogen bonds
  - $H_2O$  is covalent in nature
  - The  $O-H$  bond in  $H_2O$  is very strong
  - The electronegativity difference of  $H$  and  $O$  is very large
69. Sulphuric acid reacts with  $PCl_5$  to give  
[KCET 1996; JIPMER 2000]
- Thionyl chloride
  - Sulphur monochloride
  - Sulphuryl chloride
  - Sulphur tetrachloride
70. Carbogen is [EAMCET 1998]
- Pure form of carbon
  - $COCl_2$
  - Mixture of  $CO$  and  $CO_2$
  - Mixture of  $O_2$  and  $CO_2$
71. Which of the following dissociates to give  $H^+$  most easily [MP PET 1994]
- $H_2O$
  - $H_2S$
  - $H_2Te$
  - $H_2Se$
72. Superphosphate is the mixture of [CPMT 1993]
- Calcium phosphate and dil.  $H_2SO_4$
  - Sodium phosphate and dil.  $H_2SO_4$
  - Potassium phosphate and dil.  $H_2SO_4$
  - None of these
73. Among  $KO_2$ ,  $NO_2^-$ ,  $BaO_2$  and  $NO_2^+$  unpaired electron is present in [IIT 1997]
- $NO_2^+$  and  $BaO_2$
  - $KO_2$  and  $BaO_2$
  - $KO_2$  only
  - $BaO_2$  only
74. Which is not easily soluble in water [CPMT 1994]
- $H_2$
  - $O_2$
  - $SO_2$
  - $CO_2$
75. Point out in which of the following properties oxygen differs from the rest of the members of its family (Group-VIA) [MP PMT 1997]
- High value of ionisation energies
  - Oxidation states (2, 4, 6)
  - Polymorphism
  - Formation of hydrides



## 788 s and p-Block Elements

76. In the preparation of sulphuric acid,  $V_2O_5$  is used in the reaction, which is [CBSE PMT 2001; AFMC 2001]  
 (a)  $S + O_2 \rightarrow SO_2$   
 (b)  $2SO_2 + O_2 \rightarrow 2SO_3$   
 (c)  $SO_2 + H_2O \rightarrow H_2SO_4$   
 (d)  $N_2 + 3H_2 \rightarrow 2NH_3$
77. Which of the following hydrides has the lowest boiling point [MP PET 1997]  
 (a)  $H_2O$  (b)  $H_2S$   
 (c)  $H_2Se$  (d)  $H_2Te$
78. The catalyst used in the manufacture of  $H_2SO_4$  by contact process is [UPSEAT 1999]  
 (a)  $Al_2O_3$  (b)  $Cr_2O_3$   
 (c)  $V_2O_5$  (d)  $MnO_2$
79. The molecular formula of sulphur is [MP PMT 1996; MP PET/PMT 1998]  
 (a)  $S$  (b)  $S_2$   
 (c)  $S_4$  (d)  $S_8$
80. Which of the following is not suitable for use in a desiccator to dry substances  
 (a) Conc.  $H_2SO_4$  (b)  $Na_2SO_4$   
 (c)  $CaCl_2$  (d)  $P_4O_{10}$
81. Which shows polymorphism [DCE 2000]  
 (a)  $O$  (b)  $S$   
 (c)  $Se$  (d) All
82. All the elements of oxygen family are [MP PET/PMT 1998]  
 (a) Non-metals (b) Metalloids  
 (c) Radioactive (d) Polymorphic
83. The triatomic species of elemental oxygen is known as [Kerala (Med.) 2002]  
 (a) Azone (b) Polyzone  
 (c) Triozone (d) Ozone
84. When  $H_2S$  gas is passed through nitric acid, the product is [Kerala (Engg.) 2002]  
 (a) Rhombic  $S$  (b) Prismatic  $S$   
 (c) Amorphous  $S$  (d) Monoclinic  $S$   
 (e) None of these
85. Shape of  $O_2F_2$  is similar to that of [AIIMS 2004]  
 (a)  $C_2F_2$  (b)  $H_2O_2$   
 (c)  $H_2F_2$  (d)  $C_2H_2$
86. Which of the following bonds has the highest energy [CBSE PMT 1996]  
 (a)  $Se - Se$  (b)  $Te - Te$   
 (c)  $S - S$  (d)  $O - O$
87. Which of the following is not a chalcogen [CPMT 1999]  
 (a)  $O$  (b)  $S$
- (c)  $Se$  (d)  $Na$
88. Which of the following is a suboxide [DPMT 2001]  
 (a)  $Ba_2O$  (b)  $Pb_2O$   
 (c)  $C_3O_2$  (d)  $ZnO$
89. In the manufacture of sulphuric acid by contact process, Tyndall box is used to [KCET 2003]  
 (a) Filter dust particles  
 (b) Remove impurities  
 (c) Convert  $SO_2$  to  $SO_3$   
 (d) Test the presence of dust particles
90. Permono sulphuric acid is known as [Bihar CEE 1995]  
 (a) Marshall's acid (b) Caro's acid  
 (c) Sulphuric acid (d) None of these
91.  $KO_2 + CO_2 \rightarrow ?$  (gas) [CPMT 1997]  
 (a)  $H_2$  (b)  $N_2$   
 (c)  $O_2$  (d)  $CO$
92.  $H_2SO_4$  acts as dehydrating agent in its reaction with [JIPMER 2001]  
 (a)  $H_2C_2O_4$  (b)  $Ba(OH)_2$   
 (c)  $KOH$  (d)  $Zn$
93. Which of the following group shows the highest boiling point [MP PET 2002]  
 (a)  $H_2O$  (b)  $H_2S$   
 (c)  $H_2Se$  (d)  $H_2Te$
94. In presence of moisture,  $SO_2$  can [BVP 2003]  
 (a) Act as oxidant (b) Lose electron  
 (c) Gain electron (d) Not act as reductant
95. A gas that cannot be collected over water is [Kurukshetra CEE 1998]  
 (a)  $N_2$  (b)  $O_2$   
 (c)  $SO_2$  (d)  $PH_3$
96. Which of the following is formed by the action of water on sodium peroxide  
 (a)  $H_2$  (b)  $N_2$   
 (c)  $O_2$  (d)  $CO_2$
97. Sulphur on boiling with  $NaOH$  solution gives [Roorkee 1999]  
 (a)  $Na_2S_2O_3 + NaHSO_3$  (b)  $Na_2S_2O_3 + Na_2S$   
 (c)  $Na_2SO_3 + H_2S$  (d)  $Na_2SO_3 + SO_2$
98. Quartz is a crystalline variety of [Pb. CET 2002; Pb. PMT 2000, 04]  
 (a) Silicon carbide (b) Sodium silicate  
 (c) Silica (d) Silicon
99. The most efficient agent for the absorption of  $SO_3$  is [BHU 2004; DPMT 2004]  
 (a) 80%  $H_2SO_4$  (b) 98%  $H_2SO_4$

- (c) 50%  $H_2SO_4$  (d) 20%  $H_2S_2O_7$
100. Conc.  $H_2SO_4$  is diluted [Pb. CET 2001]  
 (a) By adding water in  $H_2SO_4$   
 (b) By adding  $H_2SO_4$  in water  
 (c) By adding glacial acetic acid in  $H_2SO_4$   
 (d) None of the above
101. The smog is essentially caused by the presence of [AIEEE 2004]  
 (a) Oxides of sulphur and nitrogen  
 (b)  $O_2$  and  $N_2$   
 (c)  $O_2$  and  $O_3$   
 (d)  $O_3$  and  $N_2$
102. Bleaching action of  $SO_2$  is due to its [CPMT 2004]  
 (a) Oxidising property (b) Acidic property  
 (c) Basic property (d) Reducing property
103. Oxygen is not evolved on reaction of ozone with [Pb. PMT 2004]  
 (a)  $H_2O_2$  (b)  $SO_2$   
 (c)  $Hg$  (d)  $KI$
104. When  $PbO_2$  reacts with conc.  $HNO_3$  the gas evolved is [IIT-JEE (Screening) 2005]  
 (a)  $NO_2$  (b)  $O_2$   
 (c)  $N_2$  (d)  $N_2O$
105. Sulphur in +3 oxidation state is present in [DCE 2003]  
 (a) Sulphurous acid (b) Pyrosulphuric acid  
 (c) Dithionous acid (d) Thiosulphuric acid
106.  $SO_2 + H_2S \rightarrow$  product. the final product is [Orissa JEE 2005]  
 (a)  $H_2O + S$  (b)  $H_2SO_4$   
 (c)  $H_2SO_3$  (d)  $H_2S_2O_3$
107. Oncontrolled hydrolysis and condensation,  $R_3SiCl$  yields [Orissa JEE 2005]  
 (a)  $R_3Si - O - SiR_3$  (b)  $\left( R_3Si - O - SiR_3 \right)_n$   
 (c)  $R_3SiOH$  (d)  $\begin{array}{c} R \quad R \\ | \quad | \\ -Si - O - Si - \\ | \quad | \\ O \quad O \\ -Si - O - Si - \\ | \quad | \end{array}$
108. Ozone deplect due to the formation of following compound in Antarctica [Kerala CET 2005]  
 (a) Acrolein (b) Peroxy acetyl nitrate  
 (c)  $SO_2$  and  $SO_3$  (d) Chlorine nitrate  
 (e) Formaldehyde
1. The correct order of the thermal stability of hydrogen halides ( $H-X$ ) is [AIEEE 2005]  
 (a)  $HI > HBr > HCl > HF$   
 (b)  $HF > HCl > HBr > HI$   
 (c)  $HCl < HF < HBr < HI$   
 (d)  $HI > HCl < HF < HBr$
2. Phosgene is the common name of [CPMT 1974, 86; DPMT 1989; MP PMT 1994]  
 (a) Carbonyl chloride (b) Phosphine  
 (c) Phosphorus oxychloride (d) Phosphorus trichloride
3. The solubility of iodine in water increases in the presence of [CPMT 1973, 74, 78, 86, 89, 91; NCERT 1973; AFMC 1995]  
 (a) Alcohol (b) Chloroform  
 (c) Sodium hydroxide (d) Potassium iodide
4. When thiosulphate ion is oxidised by iodine, which one of the following ion is produced [CPMT 1989; AFMC 1990; CBSE PMT 1996]  
 (a)  $SO_3^{2-}$  (b)  $SO_4^{2-}$   
 (c)  $S_4O_6^{2-}$  (Tetrathionate) (d)  $S_2O_6^{2-}$
5. Bromine is liberated when an aqueous solution of potassium bromide is treated with [CBSE PMT 1989]  
 (a)  $Cl_2$  (b)  $I_2$   
 (c) Dilute  $H_2SO_4$  (d)  $SO_2$
6. Which of the following has greatest reducing power [CPMT 1984, 88, 89, 94]  
 (a)  $HI$  (b)  $HBr$   
 (c)  $HCl$  (d)  $HF$
7. Chlorine was discovered by [BHU 1988]  
 (a) Davy (b) Priestley  
 (c) Rutherford (d) Sheele
8. Bad conductor of electricity is [MP PET/PMT 1988]  
 (a)  $H_2F_2$  (b)  $HCl$   
 (c)  $HBr$  (d)  $HI$
9. Which of the following will not occur [MP PET/PMT 1988]  
 (a)  $Fe + H_2SO_4 \rightarrow FeSO_4 + H_2$   
 (b)  $Cu + 2AgNO_3 \rightarrow Cu(NO_3)_2 + 2Ag$   
 (c)  $2KBr + I_2 \rightarrow 2KI + Br_2$   
 (d)  $CuO + H_2 \rightarrow Cu + H_2O$
10. Bromine is obtained on commercial scale from [CPMT 1985]  
 (a) Caliche (b) Carnellite  
 (c) Common salt (d) Cryolite
11. Which one of the halogen acids is a liquid [MP PMT 1985]  
 (a)  $HF$  (b)  $HCl$   
 (c)  $HBr$  (d)  $HI$
12. Which one of the following acids is the weakest [MP PMT 1985]  
 (a)  $HClO$  (b)  $HBr$   
 (c)  $HClO_3$  (d)  $HCl$

Halogen family

## 790 s and p-Block Elements

13. Chlorine reacts with sodium hydroxide under various condition to give  
 (a) Sodium chloride (b) Sodium hypochlorite  
 (c) Sodium chlorate (d) All of these
14.  $Br_2$  gas turns starch iodide paper  
 [CPMT 1987; AFMC 1987; AMU 1999]  
 (a) Blue (b) Red  
 (c) Colourless (d) Yellow
15. When  $Cl_2$  gas is passed through hot and conc. solution of  $KOH$ , following compound is formed  
 [CPMT 1971, 79; BVP 2003]  
 (a)  $KCl$  (b)  $KClO_3$   
 (c)  $KClO_2$  (d)  $KClO_4$
16. Deacon's process is used in the manufacture of  
 [BHU 1979]  
 (a) Bleaching powder (b) Sulphuric acid  
 (c) Nitric acid (d) Chlorine
17. Which of the following is the weakest acid  
 [BHU 1984, 86; CPMT 1988, 2000; MP PMT 1995;  
 MP PET 1989, 90; Kurukshetra CEE 1998;  
 Roorkee 2000; RPMT 2000]  
 (a)  $HF$  (b)  $HCl$   
 (c)  $HBr$  (d)  $HI$
18. Which is the most volatile compound  
 [CPMT 1979; AIIMS 1980; DPMT 1982;  
 Kurukshetra CEE 1998; J & K CET 2005; DPMT 2002]  
 (a)  $HF$  (b)  $HCl$   
 (c)  $HBr$  (d)  $HI$
19. On boiling an aqueous solution of  $KClO_3$  with iodine, the following product is obtained [NCERT 1980]  
 (a)  $KIO_3$  (b)  $KClO_4$   
 (c)  $KIO_4$  (d)  $KCl$
20. Colour of iodine solution is disappeared by shaking it with aqueous solution of [CPMT 1979, 81; MP PET/PMT 1988;  
 MP PMT 1986; RPMT 1997, 2002]  
 (a)  $H_2SO_4$  (b)  $Na_2S$   
 (c)  $Na_2S_2O_3$  (d)  $Na_2SO_4$
21. A quick supply of  $Cl_2$  gas may be made by reacting crystals of  $KMnO_4$  with a concentrated solution of  
 [CPMT 1973]  
 (a) Potassium chloride (b) Sodium chloride  
 (c) Bleaching powder (d) Hydrochloric acid
22. The strongest acid amongst the following is [IIT 1989]  
 (a)  $HClO_4$  (b)  $HClO_3$   
 (c)  $HClO_2$  (d)  $HClO$
23. Iodine deficiency in diet causes  
 (a) Nightblindness (b) Rickets  
 (c) Goitre (d) Beri-beri
24. Which of the following is correct [CPMT 1985]  
 (a) Iodine is a solid  
 (b) Chlorine is insoluble in water  
 (c) Iodine is more reactive than bromine  
 (d) Bromine is more reactive than chlorine
25. [NCERT 1973]  $HBr$  is treated with concentrated  $H_2SO_4$  redish brown gas evolved, gas is [Pb. CET 2003]  
 (a) Mixture of bromine and  $HBr$   
 (b)  $HBr$   
 (c) Bromine  
 (d) None of these
26. Sea weed is employed as a source of manufacture of  
 [CPMT 1982, 86, 2002; MP PET 2002]  
 (a)  $F$  (b)  $I$   
 (c)  $Br$  (d)  $Cl$
27. Which of the following is most active halogen  
 [MP PET 1990]  
 (a)  $Cl_2$  (b)  $Br_2$   
 (c)  $I_2$  (d)  $F_2$
28. Which of the following represents clear electropositive properties  
 [MP PET/PMT 1988; MP PMT 1991]  
 (a)  $F$  (b)  $Cl$   
 (c)  $Br$  (d)  $I$
29. Which statement is false [MP PET 1991]  
 (a) Electronegativity of fluorine is maximum  
 (b) Electron affinity of fluorine is maximum  
 (c) Melting point of fluorine is minimum  
 (d) Boiling point of fluorine is maximum
30. Strongest reducing agent is [MP PMT 1990]  
 (a)  $F^-$  (b)  $Cl^-$   
 (c)  $Br^-$  (d)  $I^-$
31. Which of the following represents outermost shell electronic configuration of halogens  
 [MP PET 1991; Manipal MEE 1995; MP PMT 1996]  
 (a)  $s^2 p^3$  (b)  $s^2 p^6$   
 (c)  $s^2 p^5$  (d)  $s^2 p^5$
32. Chlorine can remove [MP PET 1990]  
 (a)  $Br$  from  $NaBr$  solution  
 (b)  $F$  from  $NaF$  solution  
 (c)  $Cl$  from  $NaCl$  solution  
 (d)  $F$  from  $CaF_2$  solution
33. Hydrolysis of which of the following does not occur  
 [AIIMS 1982]  
 (a)  $VCl_4$  (b)  $TiCl_4$   
 (c)  $SiCl_4$  (d)  $CCl_4$
34. Nitric acid converts iodine into [MP PMT 1990]  
 (a) Iodic acid (b) Hydroiodic acid  
 (c) Iodine nitrate (d) Iodine pentaoxide
35. In  $KI$  solution,  $I_2$  readily dissolves and forms  
 [MP PMT 1989; EAMCET 1992]  
 (a)  $I^-$  (b)  $KI_2$   
 (c)  $KI_2^-$  (d)  $KI_3$
36. Which reaction cannot be used for the production of halogen acid [MP PMT 1989]



- (a)  $2KBr + H_2SO_4 \rightarrow K_2SO_4 + 2HBr$   
 (b)  $NaHSO_4 + NaCl \rightarrow Na_2SO_4 + HCl$   
 (c)  $NaCl + H_2SO_4 \rightarrow NaHSO_4 + HCl$   
 (d)  $CaF_2 + H_2SO_4 \rightarrow CaSO_4 + 2HF$
37. In dark, which of the following reacts with hydrogen  
 [MP PMT/PET 1988; Kurukshetra CEE 1998]  
 (a)  $Br_2$  (b)  $F_2$   
 (c)  $I_2$  (d)  $Cl_2$
38. The more activeness of fluorine is due to [MP PMT 1990]  
 (a)  $F-F$  bond has less energy  
 (b)  $F_2$  is gas at normal temperature  
 (c) Its electronic bond is maximum  
 (d)  $F-F$  bond has more energy
39. Which of the following after reacting with  $KI$  do not remove iodine [MP PET 1989]  
 (a)  $CuSO_4$  (b)  $K_2Cr_2O_7$   
 (c)  $HNO_3$  (d)  $HCl$
40. Aqueous solution of which of the following acids cannot be kept in a bottle of glass [MP PET 1989]  
 (a)  $HF$  (b)  $HCl$   
 (c)  $HBr$  (d)  $HI$
41. Which of the following pairs is not correctly matched  
 [MP PET 1993]  
 (a) A halogen which is liquid at room temperature—*Bromine*  
 (b) The most electronegative element—*Fluorine*  
 (c) The most reactive halogen—*Fluorine*  
 (d) The strongest oxidizing halogen—*Iodine*
42. Iodine is formed when potassium iodide reacts with a solution of [MNR 1984; MP PET/PMT 1998]  
 (a)  $ZnSO_4$  (b)  $CuSO_4$   
 (c)  $(NH_4)_2SO_4$  (d)  $Na_2SO_4$
43. As the atomic number of halogens increases, the halogens  
 [MP PMT 1991]  
 (a) Lose the outermost electrons less readily  
 (b) Become lighter in colour  
 (c) Become less denser  
 (d) Gain electrons less readily
44. Which statement is correct about halogens  
 [EAMCET 1991]  
 (a) They are all diatomic and form univalent ions  
 (b) They are all capable of exhibiting several oxidation states  
 (c) They are all diatomic and form divalent ions  
 (d) They can mutually displace each other from the solution of their compounds with metals
45. Mark the smallest atom [CPMT 1984, 89]  
 (a)  $F$  (b)  $Cl$  (c)  $Br$  (d)  $I$
46. Mark the element which shows only one oxidation state  
 [BHU 1988; MP PET 2002]  
 (a)  $F$  (b)  $Cl$   
 (c)  $Br$  (d)  $I$
47. Which of the following arrangement for the three halogens  $Cl$ ,  $Br$  and  $I$  when placed in the order of their increasing electron affinity is correct [CPMT 1990]  
 (a)  $Cl$ ,  $Br$ ,  $I$  (b)  $I$ ,  $Br$ ,  $Cl$   
 (c)  $Br$ ,  $Cl$ ,  $I$  (d)  $I$ ,  $Cl$ ,  $Br$
48. Which of the following is strongest oxidising agent  
 [CPMT 1978, 91, 94; MNR 1990; AMU 1983, 84; MP PMT 1991, 92, 96; IIT 1992; UPSEAT 2000]  
 (a)  $F_2$  (b)  $Cl_2$   
 (c)  $Br_2$  (d)  $I_2$
49. Fluorine is a better oxidising agent than  $Br_2$ . It is due to  
 [EAMCET 1992]  
 (a) Small size of fluorine  
 (b) More electron repulsion in fluorine  
 (c) More electronegativity of fluorine  
 (d) Non-metallic nature of fluorine
50. Fluorine is a stronger oxidising agent than chlorine in aqueous solution. This is attributed to many factors except  
 [Pb. CET 1989]  
 (a) Heat of dissociation (b) Electron affinity  
 (c) Heat of hydration (d) Ionisation potential
51. Mark the element which displaces three halogens from their compounds  
 [MP PMT 1980, 82; BHU 1984; NCERT 1987]  
 (a)  $F$  (b)  $Cl$   
 (c)  $Br$  (d)  $I$
52. Which one of the following is the most basic  
 [CPMT 1975, 77; MP PMT 2001]  
 (a)  $I$  (b)  $Br$   
 (c)  $Cl$  (d)  $F$
53. Which of the following will displace the halogen from the solution of the halide [EAMCET 1979]  
 (a)  $Br_2$  added to  $NaCl$  solution  
 (b)  $Cl_2$  added to  $KCl$  solution  
 (c)  $KCl$  added to  $NaF$  solution  
 (d)  $Br_2$  added to  $KI$  solution
54. Fluorine does not form positive oxidation states because  
 [AIIMS 1987]  
 (a) It is most electronegative element  
 (b) It forms only anions in ionic compounds  
 (c) It cannot form multiple bonding



## 792 s and p-Block Elements

- (d) It shows non-bonded electron pair repulsion due to small size
55. In the isolation of fluorine, a number of difficulties were encountered. Which statement is correct  
[NCERT 1983, 86]
- (a) The potential required for the discharge of the fluoride ions is the lowest  
(b) Fluorine reacts with most glass vessels  
(c) Fluorine has great affinity for hydrogen  
(d) Electrolysis of aqueous  $HF$  gives ozonised oxygen
56. Fluorine reacts with water to give [BHU 1988, 89]
- (a)  $HF$  and  $O_2$  (b)  $HF$  and  $OF_2$   
(c)  $HF$  and  $O_3$  (d)  $HF, O_2$  and  $O_3$
57. In which of the following, oxygen has + 2 oxidation number [EAMCET 1986]
- (a)  $F_2O$  (b)  $Cl_2O$   
(c)  $Na_2O_2$  (d)  $Na_2O$
58. The electrolysis of a certain liquid resulted in the formation of hydrogen at the cathode and chlorine at the anode. The liquid is
- (a) Pure water  
(b)  $H_2SO_4$  solution  
(c)  $NaCl$  solution in water  
(d)  $CuCl_2$  solution in water
59. In the preparation of chlorine from  $HCl$ ,  $MnO_2$  acts as [CPMT 1981]
- (a) Oxidising agent (b) Reducing agent  
(c) Catalytic agent (d) Dehydrating agent
60. Chlorine gas is dried over [CPMT 1980]
- (a)  $CaO$  (b)  $NaOH$   
(c)  $KOH$  (d) Conc.  $H_2SO_4$
61. Chlorine can be manufactured from [CPMT 1989]
- (a) Electrolysis of  $NaCl$   
(b) Electrolysis of brine  
(c) Electrolysis of bleaching powder  
(d) All of these
62. When chlorine water is exposed to sunlight,  $O_2$  is liberated. Hence [AFMC 1989]
- (a) Hydrogen has little affinity to  $O_2$   
(b) Hydrogen has more affinity to  $O_2$   
(c) Hydrogen has more affinity to  $Cl_2$   
(d) It is a reducing agent
63. When cold  $NaOH$  reacts with  $Cl_2$  which of the following is formed [AFMC 1992]
- (a)  $NaClO$  (b)  $NaClO_2$   
(c)  $NaClO_3$  (d) None of these
64. Chlorine is used in water for [CBSE PMT 1988]
- (a) Killing germs (b) Prevention of pollution  
(c) Cleansing (d) Removing dirt
65. Chlorine cannot be used [MP PET/PMT 1988]
- (a) As bleaching agent  
(b) In sterilisation  
(c) In preparation of antiseptic  
(d) For extraction of silver and copper
66. Chlorine acts as a bleaching agent only in presence of [IIT 1983; DCE 2002]
- (a) Dry air (b) Moisture  
(c) Sunlight (d) Pure oxygen
67. Euchlorine is a mixture of [CPMT 1988]
- (a)  $Cl_2$  and  $SO_2$  (b)  $Cl_2$  and  $ClO_2$   
(c)  $Cl_2$  and  $CO$  (d) None of these
68. A gas reacts with  $CaO$ , but not with  $NaHCO_3$ . The gas is [AFMC 1987]
- (a)  $CO_2$  (b)  $Cl_2$   
(c)  $N_2$  (d)  $O_2$
69. When chlorine is passed over dry slaked lime at room temperature, the main reaction product is [CBSE PMT 1992]
- (a)  $Ca(ClO_2)_2$  (b)  $CaCl_2$   
(c)  $CaOCl_2$  (d)  $Ca(OCl_2)_2$
70. Bromine is obtained commercially from sea water by adding [CPMT 1988]
- (a)  $AgNO_3$  solution (b) Crystals of  $NaBr$   
(c)  $Cl_2$  (d)  $C_2H_4$
71. In the manufacture of bromine from sea water, the mother liquor containing bromides is treated with [CBSE PMT 1992; MP PMT 2001; BHU 2002; JIPMER 2002]
- (a)  $CO_2$  (b)  $Cl_2$   
(c)  $I_2$  (d)  $SO_2$
72.  $Br^-$  is converted into  $Br_2$  by using [CPMT 1987]
- (a)  $Cl_2$  (b) Conc.  $HCl$   
(c)  $HBr$  (d)  $H_2S$
73. A salt, which on heating with conc.  $H_2SO_4$  gives violet vapours, is [CPMT 1971]
- (a) Iodide (b) Nitrate  
(c) Sulphate (d) Bromide
74. When  $I_2$  is dissolved in  $CCl_4$ , the colour that results is [AFMC 1993]
- (a) Brown (b) Violet  
(c) Colourless (d) Bluish green
75. Which of the following halogen oxides is ionic [CPMT 1989]

- (a)  $ClO_2$  (b)  $BrO_2$  NCERT 1974; CPMT 1976, 90  
 (c)  $I_2O_5$  (d)  $I_4O_9$
76.  $KI$  when heated with conc.  $H_2SO_4$  gives [MP PET/PMT 1988]  
 (a)  $HI$  (b)  $I_2$   
 (c)  $HIO_3$  (d)  $KIO_3$
77. The type of bonding in  $HCl$  molecule is [AIIMS 1992]  
 (a) Pure covalent (b) Polar covalent  
 (c) Highly covalent (d)  $H$ -bonding
78.  $HCl$  is a gas, but  $HF$  is a low boiling liquid. This is because [EAMCET 1981, 89]  
 (a)  $H-F$  bond is strong  
 (b)  $H-F$  bond is weak  
 (c) The molecules aggregate because of hydrogen bonding in  $HF$   
 (d)  $HF$  is a weak acid
79.  $HI$  cannot be prepared by the action of conc.  $H_2SO_4$  on  $KI$  because [MNR 1984]  
 (a)  $HI$  is stronger than  $H_2SO_4$   
 (b)  $HI$  is more volatile than  $H_2SO_4$   
 (c)  $H_2SO_4$  is an oxidising agent  
 (d)  $H_2SO_4$  forms complex
80. A solution of  $HCl$  in water is good conductor while gaseous hydrogen chloride is not. This is due to the reason that [NCERT 1980]  
 (a) Water is a good conductor of electricity  
 (b)  $HCl$  in water ionises  
 (c) Gas can not conduct electricity but water can  
 (d) None of these
81. Sodium chloride when heated with conc.  $H_2SO_4$  and solid potassium dichromate gives [CPMT 1981, 84]  
 (a) Chromic chloride (b) Chromyl chloride  
 (c) Chromous chloride (d) None of these
82. Hydrogen bonding does not play any role in boiling of [AFMC 1992]  
 (a)  $NH_3$  (b)  $H_2O$   
 (c)  $HI$  (d)  $C_2H_5OH$
83.  $HBr$  and  $HI$  reduce sulphuric acid,  $HCl$  can reduce  $KMnO_4$  and  $HF$  can reduce [IIT 1981; MP PET 1993]  
 (a)  $H_2SO_4$  (b)  $KMnO_4$   
 (c)  $K_2Cr_2O_7$  (d) None of these
84. Which has the highest molar heat of vaporisation [CPMT 1991]  
 (a)  $HF$  (b)  $HCl$   
 (c)  $HBr$  (d)  $HI$
85. Mark the strongest acid [Bihar MEE 1996; MP PET/PMT 1998;  
 (a)  $HF$  (b)  $HCl$   
 (c)  $HBr$  (d)  $HI$
- (a)  $HF$  (b)  $HCl$   
 (c)  $HBr$  (d)  $HI$
86. Which of the following hydrogen halides has the highest boiling point [AIIMS 1980; DPMT 1983; MP PMT 1983]  
 (a)  $HF$  (b)  $HCl$   
 (c)  $HBr$  (d)  $HI$
87. Hydrogen bonding is present in [MP PMT 1989; DPMT 1990; Roorkee 1995]  
 (a)  $HF$  (b)  $HCl$   
 (c)  $HBr$  (d)  $HI$
88. The type of hybrid orbitals used by chlorine atom in  $ClO_2^-$  is  
 (a)  $sp^3$  (b)  $sp^2$   
 (c)  $sp$  (d) None of these
89. Which one is the anhydride of  $HClO_4$  [AIIMS 1983; BHU 1983; AMU 1984]  
 (a)  $Cl_2O$  (b)  $ClO_2$   
 (c)  $Cl_2O_6$  (d)  $Cl_2O_7$
90. Which of the following halogens is solid at room temperature [MP PET 1999; AFMC 1999]  
 (a) Chlorine (b) Iodine  
 (c) Bromine (d) Fluorine
91. Which of the following chemical contains chlorine [MP PET 1999]  
 (a) Fischer salt (b) Epsom salt  
 (c) Fremy's salt (d) Spirit of salt
92. The element which never acts as reducing agent in a chemical reaction is [Bihar CEE 1995]  
 (a)  $O$  (b)  $Li$   
 (c)  $F$  (d)  $C$
93. Concentrated  $HNO_3$  reacts with  $I_2$  to give [IIT 1989; Roorkee 1995; JIPMER 2001]  
 (a)  $HI$  (b)  $HOI$   
 (c)  $HOIO_3$  (d)  $HOIO_2$
94. The formula of some fluorides are given below. Which of them will combine further with fluorine [NCERT]  
 (a)  $IF_5$  (b)  $NaF$   
 (c)  $CaF_2$  (d)  $SF_6$
95. Which one below is a pseudohalide [AIIMS 1982]  
 (a)  $CN^-$  (b)  $ICl$   
 (c)  $IF_5$  (d)  $I_3^-$
96. Which one is highest melting halide [AIIMS 1982; BHU 1999]  
 (a)  $NaCl$  (b)  $NaBr$   
 (c)  $NaF$  (d)  $NaI$
97. The above answer is correct because the chosen halide has [AIIMS 1982]  
 (a) Minimum ionic character  
 (b) Maximum ionic character

## 794 s and p-Block Elements

- (c) Highest oxidising power  
(d) Lowest polarity
- 98.** Which of the following oxidizes  $H_2O$  to oxygen  
[MP PET 1994]  
(a) Chlorine (b) Fluorine  
(c) Bromine (d) Iodine
- 99.** The bleaching action of the bleaching powder is due to the liberation of  
[MP PMT 1994]  
(a) Chlorine (b) Molecular oxygen  
(c) Nascent oxygen (d) Calcium carbonate
- 100.** Which of the following element is extracted commercially by the electrolysis of an aqueous solution of its compound  
[KCET 2002]  
(a) Chlorine (b) Bromine  
(c) Aluminium (d) Calcium
- 101.** The effective component of bleaching powder is ..... of calcium  
[Kerala (Engg.) 2002]  
(a) Chlorine (b) Bromine  
(c) Aluminium (d) Calcium
- 102.**  $Na_2S_2O_3 + I_2 \rightarrow$  Product is  
[BHU 2003]  
(a)  $Na_2S$  (b)  $NaI$   
(c)  $Na_2S_4O_6$  (d)  $S_2$
- 103.** Which of the following is prepared by electrolytic method  
[CBSE PMT 2001]  
(a)  $Ca$  (b)  $Sn$   
(c)  $S$  (d)  $F_2$
- 104.** Beilstein test is used for  
[AFMC 1995]  
(a)  $N_2$  (b)  $Cl$   
(c)  $Na$  (d)  $CO_2$
- 105.** Which one will liberate  $Br_2$  from  $KBr$   
[BVP 2003]  
(a)  $I_2$  (b)  $SO_2$   
(c)  $HI$  (d)  $Cl_2$
- 106.** Chlorine dioxide is best prepared by passing dry  
[Kerala PMT 2003]  
(a) Chlorine gas over hot  $HgO$   
(b) Chlorine and oxygen gas over hot Pt catalyst  
(c) Hydrogen chloride and oxygen over silver oxide  
(d) Hydrogen chloride over phosphorus pentoxide  
(e) Chlorine over hot silver chlorate
- 107.** The mixture of concentrated  $HCl$  and  $HNO_3$  made in 3 : 1 ratio contains  
[AIIMS 2003]  
(a)  $ClO_2$  (b)  $NOCl$   
(c)  $NCl_3$  (d)  $N_2O_4$
- 108.** On exciting  $Cl_2$  molecule by UV light, we get  
[UPSEAT 2003]  
(a)  $Cl$  (b)  $Cl^+$   
(c)  $Cl^-$  (d) All
- 109.** Which of the following statements is not true  
[CBSE PMT 2003]  
(a)  $HF$  is a stronger acid than  $HCl$   
(b) Among halide ions, iodide is the most powerful reducing agent  
(c) Fluorine is the only halogen that does not show a variable oxidation state  
(d)  $HOCl$  is a stronger acid than  $HOBr$
- 110.** The correct order of acidic strength [Pb. CET 2004]  
(a)  $Cl_2O_7 > SO_2 > P_4O_{10}$  (b)  $K_2 > CaO > MgO$   
(c)  $CO_2 > N_2O_5 > SO_3$  (d)  $Na_2O > MgO > Al_2O_3$
- 111.** Halogen acid used in the preparation of aqua regia is  
[DPMT 2002]  
(a)  $HBr$  (b)  $HI$   
(c)  $HCl$  (d)  $HF$
- 112.**  $NaOCl$  is used as a bleaching agent and sterilising agent. It can be synthesized by the action of [RPET 2003]  
(a)  $NaCl$  with  $H_2O$   
(b)  $NH_4Cl$  with  $NaOH$   
(c)  $Cl_2$  with cold and dilute  $NaOH$   
(d)  $Cl_2$  with hot and concentrated  $NaOH$
- 113.** Metal halide which is insoluble in water is [AIIMS 1996]  
(a)  $AgI$  (b)  $KBr$   
(c)  $CaCl_2$  (d)  $AgF$
- 114.** Which one among the following non-metals liquid at  $25^\circ C$   
[MP PMT 1999]  
(a) Bromine (b) Carbon  
(c) Phosphorus (d) Sulphur
- 115.** Bleaching action of chlorine is due to [Bihar CEE 1995]  
(a) Oxidation (b) Reduction  
(c) Hydrolysis (d) Its acidic nature
- 116.** Hydrogen iodide cannot be prepared by the action of conc.  $H_2SO_4$  on potassium iodide because [Bihar CEE 1995]  
(a)  $HI$  is stronger than  $H_2SO_4$   
(b)  $HI$  is more volatile than  $H_2SO_4$   
(c)  $H_2SO_4$  is an oxidising agent  
(d)  $H_2SO_4$  forms complex
- 117.** White enamel of our teeth is [Bihar CEE 1995]  
(a)  $Ca_3(PO_4)_2$  (b)  $CaF_2$   
(c)  $CaCl_2$  (d)  $CaBr_2$
- 118.** The least active halogen with hydrogen is [DPMT 1996]  
(a)  $Cl$  (b)  $I$   
(c)  $Br$  (d)  $F$
- 119.** Iodine dissolves readily in [BHU 1996]  
(a) Water (b) Potassium iodide  
(c) Carbon tetrachloride (d) Alcohol

120. Which one of the following compounds in aqueous solution gives a white precipitate with perchloric acid  
[EAMCET 1997]  
(a)  $NaCl$  (b)  $KCl$   
(c)  $MgCl_2$  (d)  $FeCl_3$
121. Which of the following sequence is correct with reference to the oxidation number of iodine[EAMCET 1997]  
(a)  $I_2 > ICl < HI < HIO_4$  (b)  $HIO_4 < ICl < I_2 < HI$   
(c)  $I_2 < HI < ICl < HIO_4$  (d)  $HI < I_2 < ICl < HIO_4$
122. The correct order of acidic strength is  
[Pb. PMT 1998; AFMC 1998; KCET 2000; Orissa JEE 2005]  
(a)  $HF < HCl < HBr < HI$   
(b)  $HCl < HBr < HF < HI$   
(c)  $HBr < HCl < HI < HF$   
(d)  $HI < HBr < HCl < HF$
123. The chief source of iodine in which it is present as sodium iodate is [JIPMER 1997]  
(a) Sea weeds  
(b) Caliche  
(c) Carnallite  
(d) Iodine never exists as sodium iodate
124. The lattice energy of the lithium halides is in the following order [Roorkee Qualifying 1998]  
(a)  $LiF > LiCl > LiBr > LiI$   
(b)  $LiCl > LiF > LiBr > LiI$   
(c)  $LiBr > LiCl > LiF > LiI$   
(d)  $LiI > LiBr > LiCl > LiF$
125. Which has the strong bond [DCE 2001]  
(a)  $F - F$  (b)  $F - Cl$   
(c)  $F - Br$  (d)  $Cl - B$
126. Iodine and hypo react to produce [DPMT 2001]  
(a)  $Na_2S$  (b)  $Na_2SO_4$   
(c)  $Na_2S_4O_6$  (d)  $Na_2SO_3$
127. Bleaching powder is correctly represented as [RPMT 1997]  
(a)  $CaOCl_2$  (b)  $CaO$   
(c)  $CaO(Cl)$  (d)  $CaCl(OCl)$
128. When chlorine reacts with cold and dilute solution of sodium hydroxide, the products obtained are [CBSE PMT 1998]  
(a)  $Cl^- + ClO^-$  (b)  $Cl^- + ClO_2^-$   
(c)  $Cl^- + ClO_3^-$  (d)  $Cl^- + ClO_4^-$
129. A one litre flask is full of brown bromine vapour. The intensity of brown colour of vapour will not decrease appreciably on adding to the flask some [CBSE PMT 1998]  
(a) Pieces of marble (b) Carbon disulphide  
(c) Carbon tetrachloride (d) Animal charcoal powder
130. Which of the following statements is correct [BHU 1997]  
(a) Only chlorine and bromine form oxy acids  
(b) All halogens form oxy acids  
(c) All halogens except fluorine form oxy acids  
(d) Only iodine form oxy acids
131. When iodine reacts with  $NaF$ ,  $NaBr$  and  $NaCl$  [CPMT 1997]  
(a) It gives mixture of  $F_2$ ,  $Cl_2$  and  $Br_2$   
(b) It gives chlorine  
(c) It gives bromine  
(d) None of these
132. Which is the strongest of the following acids [JIPMER 1999]  
(a)  $HClO_4$  (b)  $H_2SO_4$   
(c)  $HCl$  (d)  $HNO_3$
133. Hydrogen has a tendency to gain one electron to acquire helium configuration. In this respect it resembles [JIPMER 1999]  
(a) Halogens (b) Actinides  
(c) Transition elements (d) Alkali metals
134. What is the product obtained in the reaction of  $HgCl_2$  and  $Hg(CN)_2$  [MP PET 2002]  
(a)  $(CN)_2$   
(b) Addition compound  $HgCl_2 \cdot Hg(CN)_2$   
(c)  $Hg(CN)Cl$   
(d)  $Hg[Hg(CN)_2Cl_2]$
135. The weakest acid  $HX$  ( $X = F, Cl, Br, I$ ) is [BHU 2000]  
(a)  $HF$  (b)  $HCl$   
(c)  $HBr$  (d)  $HI$
136. Bleaching powder is obtained by passing chlorine on [KCET 2002]  
(a) Lime stone (b) Quick lime  
(c) Slaked lime (d) Pure lime
137. Chlorine is liberated, when we heat [AFMC 1998]  
(a)  $KMnO_4 + NaCl$  (b)  $K_2Cr_2O_7 + MnO_2$   
(c)  $Pb_2(NO_3)_4 + MnO_2$  (d)  $K_2Cr_2O_7 + HCl$
138. Which of the following silver compounds finds maximum use in photography  
(a)  $AgCl$  (b)  $AgBr$   
(c)  $AgI$  (d)  $AgNO_3$
139. Which of the following halogen does not exhibit positive oxidation state in its compounds [EAMCET 1997; AIIMS 2000]  
(a)  $Cl$  (b)  $Br$   
(c)  $I$  (d)  $F$
140. The strength of oxy acids of chlorine follows the order [AIIMS 2000; CBSE PMT 2005]  
(a)  $HClO < HClO_2 < HClO_3 < HClO_4$   
(b)  $HClO_4 < HClO_3 < HClO_2 < HClO$



## 796 s and p-Block Elements

- (c)  $HClO_4 < HClO_3 < HClO < HClO_2$   
 (d) None of these
- 141.** Bleaching powder is obtained by treating chlorine with  
 [Pb. PMT 1999]  
 (a)  $CaO$  (b)  $CaCO_3$   
 (c)  $CaSO_4$  (d)  $Ca(OH)_2$
- 142.** Which statement is not true [MP PET 2000]  
 (a)  $Ni(CO)_4$  is diamagnetic  
 (b)  $BI_3$  is a stronger Lewis acid than  $BF_3$   
 (c) Graphite conducts electricity whereas diamond does not  
 (d)  $CCl_4$  is hydrolysed whereas  $BCl_3$  is inert
- 143.** Bleaching powder loses its power on keeping for a long time because [KCET 2000]  
 (a) It changes into calcium hypochlorate  
 (b) It changes into calcium chloride and calcium hydroxide  
 (c) It absorbs moisture  
 (d) It changes into calcium chloride and calcium chlorate
- 144.** The compound which forms a dative bond with ammonia  
 [JIPMER 2001]  
 (a)  $CCl_4$  (b)  $BCl_3$   
 (c)  $MgCl_2$  (d)  $NaCl$
- 145.** The bleaching action of bleaching powder is due to the formation of [Roorkee 1999]  
 (a)  $CaCl_2$  (b)  $CaSO_4$   
 (c)  $HClO$  (d)  $Ca(ClO_3)_2$
- 146.** Fluorine with dilute  $NaOH$  gives [MH CET 2000]  
 (a)  $OF_2$  (b)  $O_3$   
 (c)  $O_2$  (d)  $HF$  and  $O_2$
- 147.** Which is not oxidised by  $MnO_2$  [DCE 2003]  
 (a)  $F$  (b)  $Cl$   
 (c)  $I_2$  (d)  $I$
- 148.** Bromine water reacts with  $SO_2$  to form [AFMC 1995]  
 (a)  $H_2O$  and  $HBr$  (b)  $H_2SO_4$  and  $HBr$   
 (c)  $HBr$  and  $S$  (d)  $S$  and  $H_2O$
- 149.** Which of the following reaction is not feasible  
 [CBSE PMT PMT 2002]  
 (a)  $2KI + Br_2 \rightarrow 2KBr + I_2$   
 (b)  $2H_2O + 2F_2 \rightarrow 2HF + O_2$   
 (c)  $2KBr + I_2 \rightarrow 2KI + Br_2$   
 (d)  $2KBr + Cl_2 \rightarrow 2KCl + Br_2$
- 150.** Which of the following has the lowest solubility  
 [Roorkee 2000]  
 (a)  $CaF_2$  (b)  $CaCl_2$
- (c)  $CaBr_2$  (d)  $CaI_2$
- 151.** Which one of the following pairs of substances when mixed, produces chlorine gas at room temperature [IIT 1995]  
 (a)  $NaCl$  and  $MnO_2$   
 (b)  $NaCl$  and  $HNO_3$  (conc.)  
 (c)  $NaCl$  and  $H_2SO_4$  (conc.)  
 (d)  $HCl$  (conc.) and  $KMnO_4$
- 152.** Concentrated  $H_2SO_4$  cannot be used to prepare  $HBr$  from  $NaBr$ , because it  
 (a) Reduces  $HBr$  (b) Oxidises  $HBr$   
 (c) Disproportionates  $HBr$  (d) Reacts slowly with  $NaBr$
- 153.** Which of the following halides is least stable and has doubtful existence [IIT 1996]  
 (a)  $Cl_4$  (b)  $GeI_4$   
 (c)  $SnI_4$  (d)  $PbI_4$
- 154.** Chlorine cannot displace [MP PET 1996]  
 (a) Fluorine from  $NaF$  (b) Iodine from  $NaI$   
 (c) Bromine from  $NaBr$  (d) None of these
- 155.** When fluoride is heated with conc.  $H_2SO_4$  and  $MnO_2$  the gas evolved is [DPMT 2000]  
 (a)  $F_2$  (b)  $SF$   
 (c)  $HF$  (d) None
- 156.**  $Cl_2$  reacts with  $CS_2$  in presence of  $I_2$  catalyst to form [AFMC 1995]  
 (a)  $CHCl_3$  (b)  $CCl_4$   
 (c)  $C_2H_5Cl$  (d)  $C_2H_6$
- 157.** Amongst  $LiCl, RbCl, BeCl_2$  and  $MgCl_2$ . Maximum and minimum ionic character will be shown by the compounds [RPMT 1999]  
 (a)  $LiCl, MgCl_2$  (b)  $RbCl, BeCl_2$   
 (c)  $RbCl, MgCl_2$  (d)  $MgCl_2, BeCl_2$
- 158.** Which is formed when fluorine react with hot and concentrated sodium hydroxide  
 (a)  $O_2$  (b)  $O_3$   
 (c)  $NaO$  (d)  $HF$
- 159.** Which of the following condition is used to find atomic  $Cl_2$  from molecular  $Cl_2$  [CPMT 1996]  
 (a) High temperature, high pressure  
 (b) Low temperature, high pressure  
 (c) High temperature, low pressure  
 (d) Low temperature, low pressure
- 160.** Which one is least basic [JIPMER 2000]  
 (a)  $BI_3$  (b)  $BBr_3$   
 (c)  $BCl_3$  (d)  $BF_3$
- 161.** On heating  $NaCl + K_2Cr_2O_7 + \text{conc. } H_2SO_4$ , the gas comes out is [JIPMER 2000]

- (a)  $O_2$  (b)  $Cl_2$   
(c)  $CrOCl_2$  (d)  $CrO_2Cl_2$
- 162.** Aqua regia is a mixture of [KCET (Med.) 2001]  
(a)  $3HCl + 1HNO_3$  (b)  $H_3PO_4 + H_2SO_4$   
(c)  $3HNO_3 + 1HCl$  (d)  $HCl + CH_3COOH$
- 163.** Unlike other halogens fluorine does not show higher oxidation states because [MP PET 1997]  
(a) It is highly electronegative  
(b) It has no *d*-orbitals  
(c) Its atomic radius is very small  
(d) The  $F^-$  ion is stable and isoelectronic with neon
- 164.** Which halogen does not show variable oxidation state [UPSEAT 2003]  
(a)  $F_2$  (b)  $Cl_2$   
(c)  $Br_2$  (d)  $I_2$
- 165.** To purify fluorine gas, fumes of  $HF$  are removed by [MH CET 2002]  
(a) Solid  $NaF$  (b)  $H_2$  gas  
(c) Solid  $KHF_2$  (d) None of these
- 166.** Fluorine is prepared by  
(a) Oxidation of  $HF$   
(b) Electrolysis of  $KF$   
(c) Electrolysis of fused  $KHF_2$   
(d) Decomposition of  $HgF_2$
- 167.** Amongst halogens fluorine is most oxidising because  
(a) Fluorine has highest electron affinity  
(b) Fluorine is most electronegative  
(c) Dissociation energy for fluorine molecule is lowest  
(d) All are correct
- 168.** The alkali metal halides are soluble in water but  $LiF$  is insoluble because  
(a) It is amphoteric  
(b) The  $Li-F$  bond is highly ionic  
(c) Its lattice energy is high  
(d)  $Li^+$  ion is least hydrated
- 169.** In which of the following pairs does the first gas bleaches flowers by reduction while the second gas does so by oxidation [Manipal MEE 1995]  
(a)  $CO$  and  $Cl_2$  (b)  $SO_2$  and  $Cl_2$   
(c)  $H_2$  and  $Br_2$  (d)  $NH_3$  and  $SO_2$
- 170.** Which of the following halogens does not form oxyacid [MP PET 1997]  
(a) Fluorine (b) Chlorine  
(c) Bromine (d) Iodine
- 171.** Which of the following molecule is theoretically not possible [BHU 2002]  
(a)  $OF_4$  (b)  $OF_2$   
(c)  $SF_4$  (d)  $O_2F_2$
- 172.** Iodine is released when potassium iodide reacts with [UPSEAT 1999]  
(a)  $ZnSO_4$  (b)  $CuSO_4$   
(c)  $FeSO_4$  (d)  $(NH_4)_2SO_4$
- 173.** Which of the following is used in the preparation of chlorine [CBSE PMT 1999]  
(a) Only  $MnO_2$   
(b) Only  $KMnO_4$   
(c) Both  $MnO_2$  and  $KMnO_4$   
(d) Either  $MnO_2$  or  $KMnO_4$
- 174.** Among  $Cl^-$ ,  $Br^-$ ,  $I^-$ , the correct order for being oxidised to dihalogen is [CPMT 1999]  
(a)  $I^- > Cl^- > Br^-$  (b)  $Cl^- > Br^- > I^-$   
(c)  $I^- > Br^- > Cl^-$  (d)  $Br^- > I^- > Cl^-$
- 175.** On heating  $KClO_3$ , we get [CPMT 1999]  
(a)  $Cl_2O$  (b)  $ClO_2$   
(c)  $ClO_3$  (d)  $Cl_2O_7$
- 176.** For which one of the following properties of halogens the sequence  $F > Cl > Br > I$  holds good [MP PET/1999]  
(a) Electron affinity (b) Electronegativity  
(c) Atomic radius (d) Boiling point
- 177.** Which of the following properties increases on going down from  $F$  to  $I$  in Group VII-A of the periodic table? [MP PMT 1997]  
(a) Electronegativity (b) Volatile nature  
(c) Ionic radius (d) Oxidising power
- 178.** Among the halogens, the one which is oxidised by nitric acid is [KCET 2004]  
(a) Fluorine (b) Iodine  
(c) Chlorine (d) Bromine
- 179.** The reaction of the type  $2X_2 + S \rightarrow SX_4$  is shown by sulphur when  $X$  is [DCE 2003]  
(a) Fluorine or chlorine  
(b) Chlorine only  
(c) Chlorine and bromine only  
(d)  $F, Cl, Br$  all
- 180.** When  $I_2$  is passed through  $KCl$ ,  $KF$  and  $KBr$  solutions [CPMT 2004]  
(a)  $Cl_2$  and  $Br_2$  are evolved  
(b)  $Cl_2$  is evolved



## 798 s and p-Block Elements

- (c)  $Cl_2$ ,  $Br_2$  and  $F_2$  are evolved  
(d) None of these
181. The solubility of  $I_2$  increases in water in the presence of  
[Pb. CET 2002]  
(a)  $KI$  (b)  $H_2SO_4$   
(c)  $KMnO_4$  (d)  $NH_3$
182. Which of the hydrogen halides forms salts like  $KHX_2$  (where  $X$  is a halogen atom) [Kerala PMT 2004]  
(a)  $HF$  (b)  $HCl$   
(c)  $HI$  (d)  $HBr$   
(e) All of these
183. With cold and dilute sodium hydroxide fluorine reacts to give [MH CET 2004]  
(a)  $NaF$  and  $OF_2$  (b)  $NaF + O_3$   
(c)  $O_2$  and  $O_3$  (d)  $NaF + O_2$
184. Which one of the following oxides is expected exhibit paramagnetic behaviour  
(a)  $CO_2$  (b)  $SO_2$   
(c)  $ClO_2$  (d)  $SiO_2$
185. Of the following acids, the one that is strongest is [DPMT 2004]  
(a)  $HBrO_4$  (b)  $HOCl$   
(c)  $HNO_2$  (d)  $H_3PO_3$
186. Which of the following is anhydride of perchloric acid [CPMT 2004]  
(a)  $Cl_2O_7$  (b)  $Cl_2O_5$   
(c)  $Cl_2O_3$  (d)  $HClO$
187.  $I_2$  dissolves in  $KI$  solution due to the formation of [CPMT 2004]  
(a)  $KI_2$  and  $I^-$  (b)  $K^+$ ,  $I^-$  and  $I_2$   
(c)  $KI_3^-$  (d) None of these
- (a) 5 (b) 1  
(c) 0 (d) None of these
5. Helium was discovered by  
(a) Crooks (b) Rutherford  
(c) Frankland and Lockyer (d) Dorn
6. The inert gases are [CPMT 1984]  
(a) Polyatomic (b) Triatomic  
(c) Diatomic (d) Monoatomic
7. The charcoal maintained at  $100^\circ C$  absorbs  
(a)  $Ne$  and  $Kr$  (b)  $He$  and  $Ar$   
(c)  $Ar$ ,  $Kr$ ,  $Xe$  (d)  $He$  and  $Ne$
8. Every inert gas atom  
(a) Has a saturated outermost shell  
(b) Has one electron in outermost shell  
(c) Has eight electrons in outermost shell  
(d) Has two electrons in outermost shell
9. Argon was discovered by [CPMT 1991]  
(a) Rayleigh (b) Frankland and Lockyer [CBSE PMT 2005]  
(c) Jansen (d) Ramsay
10. Deep sea divers used to respire is a mixture of  
(a) Oxygen and argon (b) Oxygen and helium  
(c) Oxygen and nitrogen (d) Oxygen and hydrogen
11. The noble gas which forms maximum number of compounds is [NCERT 1976; BHU 1980; CPMT 1982, 91; AMU 1985; DPMT 1996; MP PMT 2001; Pb.CET 2003]  
(a)  $Ar$  (b)  $He$   
(c)  $Xe$  (d)  $Ne$
12. Which of the following gases exist more abundantly in nature than the others [BHU 1982; DPMT 1982,02; CPMT 1983, 89; EAMCET 1993; Manipal MEE 1995; MHCET 2003]  
(a) Helium (b) Neon  
(c) Argon (d) Krypton
13. Which of the following is monoatomic [NCERT 1976, 77; CPMT 1983, 86, 90]  
(a) Nitrogen (b) Fluorine  
(c) Neon (d) Oxygen
14. Nuclear fusion produces  
(a) Argon (b) Deuterium  
(c) Helium (d) Krypton
15. Among the fluorides below, the one which does not exist is [NCERT 1977; CPMT 1988]  
(a)  $XeF_4$  (b)  $HeF_4$   
(c)  $SF_4$  (d)  $CF_4$

## Noble gases

1. Which of the following outer electronic configuration represents argon [DPMT 1982; CPMT 1976; NCERT 1987; Kurukshetra CEE 1998]  
(a)  $ns^2$  (b)  $ns^2np^6$   
(c)  $ns^2np^5$  (d)  $ns^2np^4$
2. Which mineral was used in isolation of radium [CPMT 1978, 81, 91]  
(a) Lime stone (b) Pitch blende  
(c) Rutile (d) Haematite
3. Which is the lightest gas  
(a) Hydrogen (b) Oxygen  
(c) Helium (d) Nitrogen
4. The valency of inert gases is
- (a) 6 (b) 2  
(c) 18 (d) 8
16. The last orbit of argon would have electrons [CPMT 1971, 78]
17. The electronic configuration of neon is [CPMT 1974, 80, 81; DPMT 1982; MNR 1995]



- (a)  $1s^2, 2s^2 2p^2$  (b)  $1s^2, 2s^2 2p^6$   
 (c)  $1s^2, 2s^2$  (c)  $1s^2$
18. The colour discharge tubes for advertisement mainly contain  
 [CPMT 1980, 89; MP PET 2002]  
 (a) Argon (b) Neon  
 (c) Helium (d) Xenon
19. Least chemical activity is shown by [CPMT 1973, 79]  
 (a) Nitrogen (b) Argon  
 (c) Methane (d) Ammonia
20. Noble gases do not react with other elements because  
 [CPMT 1981]  
 (a) They have completely paired up and stable electron shells  
 (b) The sizes of their atoms are very small  
 (c) Are not found in abundance  
 (d) Are monoatomic
21. Monazite is source of  
 (a) *He* (b) *Kr*  
 (c) *Ar* (d) *Ne*
22. Which of the following fluorides of Xenon is impossible  
 [CPMT 1982; Kurukshetra CEE 1998; RPET 1999]  
 (a)  $XeF_6$  (b)  $XeF_4$   
 (c)  $XeF_3$  (d)  $XeF_2$
23.  $XeF_2$  molecule is  
 (a) Square planer  
 (b) Trigonal bipyramidal  
 (c) Trigonal planer  
 (d) Linear
24.  $XeF_4$  on partial hydrolysis produces [AFMC 1995]  
 (a)  $XeF_2$  (b)  $XeOF_2$   
 (c)  $XeOF_4$  (d)  $XeO_3$
25. In  $XeF_2$  hybridisation of *Xe* is  
 (a)  $sp^2$  (b)  $sp^3 d$   
 (c)  $sp^3$  (d)  $sp^3 d^2$
26. Which one of the following noble gases is the least polarizable [AIIMS 1983; MP PET 1999; Pb. PMT 2001; JIPMER (Med.) 2002]  
 (a) *Xe* (b) *Ar*  
 (c) *Ne* (d) *He*
27. Which one of the following noble gases is not found in the atmosphere [MP PMT 1993]  
 (a) *Rn* (b) *Kr*  
 (c) *Ne* (d) *Ar*
28. Helium is added to the oxygen supply used by deep sea divers because [MP PMT 1993; MP PET 1997]  
 (a) It is less soluble in blood than nitrogen at high pressure  
 (b) It is lighter than nitrogen  
 (c) It is readily miscible with oxygen  
 (d) It is less poisonous than nitrogen
29. Which of the following statements is not correct for a noble gas  
 (a) *Ar* is used in electric bulbs  
 (b) *Kr* is obtained during radioactive disintegration  
 (c) Half life of *Rn* is only 3.8 days  
 (d) *He* is used in producing very low temperature
30. Which one of the following configuration represents a noble gas  
 [CPMT 1976, 83, 89; BHU 1982; Pb. CET 2000 NCERT 1979; IIT Screening 1993; EAMCET 1993]  
 (a)  $1s^2, 2s^2 2p^6, 3s^2$   
 (b)  $1s^2, 2s^2 2p^6, 3s^1$   
 (c)  $1s^2, 2s^2 2p^6$   
 (d)  $1s^2, 2s^2 2p^6, 3s^2 3p^6, 4s^2$
31. Which of the following has zero valency [DPMT 1985]  
 (a) Sodium (b) Beryllium  
 (c) Aluminium (d) Krypton
32. The forces acting between noble gas atoms are [NCERT 1989]  
 (a) Vander Waals forces  
 (b) Ion-dipole forces  
 (c) London dispersion forces  
 (d) Magnetic forces
33. Which of the following is the correct sequence of the noble gases in their group in the periodic table  
 [Manipal MEE 1995]  
 (a) *Ar, He, Kr, Ne, Rn, Xe* (b) *He, Ar, Ne, Kr, Xe, Rn*  
 (c) *He, Ne, Ar, Kr, Xe, Rn* (d) *He, Ne, Kr, Ar, Xe, Rn*
34. Which of the following represent noble gas configuration  
 [BHU 1995]  
 (a)  $1s^2, 2s^2 2p^6, 3s^2 3p^6 3d^{10}, 4s^2 4p^6 4d^{10}, 5s^2 5p^6$   
 (b)  $1s^2, 2s^2 2p^6, 3s^2 3p^6 3d^{10}, 4s^2 4p^6 4d^{10} 4f^{14}, 5s^2 5p^6 5d^1, 6s^2$   
 (c)  $1s^2, 2s^2 2p^6, 3s^2 3p^6 3d^{10}, 4s^2 4p^6 4d^{10}, 5s^2 5p^6 5d^1, 6s^2$   
 (d)  $1s^2, 2s^2 2p^6, 3s^2 3p^6 3d^{10}, 4s^2 4p^6 4d^{10}$
35.  $XeF_6$  on hydrolysis gives [MP PET 2000; Pb. PMT 2000; DCE 2002]  
 (a)  $XeO_3$  (b)  $XeO$   
 (c)  $XeO_2$  (d)  $Xe$
36. The correct order of solubility in water for *He, Ne, Ar, Kr, Xe*, is [AIIMS 2002]  
 (a)  $He > Ne > Ar > Kr > Xe$



## 800 s and p-Block Elements

- (b)  $Ne > Ar > Kr > He > Xe$   
 (c)  $Xe > Kr > Ar > Ne > He$   
 (d)  $Ar > Ne > He > Kr > Xe$
37. In  $XeF_2, XeF_4, XeF_6$  the number of lone pairs on Xe is respectively [AIEEE 2002]  
 (a) 2, 3, 1 (b) 1, 2, 3  
 (c) 4, 1, 2 (d) 3, 2, 1
38. Noble gases are group of elements which exhibit very [Kerala (Med.) 2002]  
 (a) High chemical activity  
 (b) Low chemical activity  
 (c) Minimum electronegativity  
 (d) Much paramagnetic properties
39. Which noble gas is most soluble in water [CPMT 2002]  
 (a) He (b) Ar  
 (c) Ne (d) Xe
40. Gradual addition of electronic shells in the noble gases causes a decrease in their [MP PET 1997]  
 (a) Ionization energy (b) Atomic radius  
 (c) Boiling point (d) Density
41. Which of the following noble gas does not have an octet of electrons in its outermost shell [MP PET 1996]  
 (a) Neon (b) Radon  
 (c) Argon (d) Helium
42. The low chemical reactivity of the rare gases can be attributed to their [Pune CET 1998]  
 (a) Being non-metals  
 (b) Having high ionization energies  
 (c) Being gases  
 (d) Found in nature in small quantities
43. Percentage of Ar in air is about [CPMT 1989]  
 (a) 1% (b) 2%  
 (c) 3% (d) 4%
44. Which of the following is not obtained by direct reaction of constituent elements [MP PET 1994]  
 (a)  $XeF_2$  (b)  $XeF_4$   
 (c)  $XeO_3$  (d)  $XeF_6$
45. Fluorine forms chemical compounds with [MP PMT 1994]  
 (a) He (b) Ne  
 (c) Ar (d) Xe
46. Which of the following has  $sp^3$  hybridisation [DCE 2001]  
 (a)  $XeO_3$  (b)  $BCl_3$   
 (c)  $XeF_4$  (d)  $BBr_3$
47. Which element out of He, Ar, Kr, and Xe forms least number of compounds [MP PMT 1995]  
 (a) He (b) Ar  
 (c) Kr (d) Xe
48. Which of the following exhibits the weakest intermolecular forces [KCET (Med.) 2001]  
 (a) He (b) HCl  
 (c)  $NH_3$  (d)  $H_2O$
49. Which of the following are formed by Xenon [Roorkee 2000]  
 (a)  $XeF_3$  (b)  $XeF_4$   
 (c)  $XeF_5$  (d)  $XeF_6$
50. Among the following molecule (i)  $XeO_3$  (ii)  $XeOF_4$  (iii)  $XeF_6$   
 Those having same number of lone pairs on Xe are [AIIMS 2005]  
 (a) (i) and (ii) only (b) (i) and (iii) only  
 (c) (ii) and (iii) only (d) (i),(ii) and (iii)
51. Who among the following first prepared a stable compound of noble gas [MP PET 1999]  
 (a) Rutherford (b) Rayleigh  
 (c) Ramsay (d) Neil Bartlett
52. The last member of inert gas elements is [MP PMT 1999]  
 (a) Helium (b) Neon  
 (c) Argon (d) Radon
53. Which of the following gas is/are called rare gas [CPMT 2000; Pb. CET 2002]  
 (a) Ne (b) He  
 (c) Kr (d) All of these
54. Which one of the following statements regarding helium is incorrect [AIEEE 2004]  
 (a) It is used to produce and sustain powerful superconducting magnets  
 (b) It is used as a cryogenic agent for carrying out experiments at low temperatures  
 (c) It is used to fill gas balloons instead of hydrogen because it is lighter and non-inflammable  
 (d) It is used in gas-cooled nuclear reactors
55. Which of the following inert gas liquifies easily [Pb. CET 2002]  
 (a) Kr (b) He  
 (c) Ne (d) Ar
56. The oxidation number of xenon in  $XeOF_2$  is [J & K 2005]  
 (a) Zero (b) 2  
 (c) 4 (d) 3
57. Which inert gas having highest boiling point [BCECE 2005]  
 (a) Xe (b) Ar  
 (c) Kr (d) He
58. Which of the following is an inert gas [AFMC 2005]  
 (a)  $H_2$  (b)  $O_2$   
 (c)  $N_2$  (d) Argon
59. Which of the following is most polarised [DPMT 2005]

- (a) *Kr* (b) *He*  
 (c) *Ar* (d) *Xe*
60. Which of the following is planar [J & K 2005]  
 (a)  $XeF_2$  (b)  $XeO_3F$   
 (c)  $XeO_2F_2$  (d)  $XeF_4$

- (d) Decreasing degree of hydration
7. Fusion mixture is [CPMT 2002]  
 (a)  $Na_2CO_3 + K_2CO_3$  (b)  $Na_2CO_3 + NaHCO_3$   
 (c)  $Na_2CO_3 + NaOH$  (d)  $Na_2CO_3 + K_2SO_4$

8. Concentrated hydrochloric acid when kept in open air sometimes produces a cloud of white fumes. The explanation for it is that  
 (a) Concentrated hydrochloric acid emits strongly smelling *HCl* gas all the time  
 (b) Oxygen in air reacts with the emitted *HCl* gas to form a cloud of chlorine gas  
 (c) Strong affinity of *HCl* gas for moisture in air results in forming of droplets of liquid solution which appears like a cloudy smoke  
 (d) Due to strong affinity for water, concentrated hydrochloric acid pulls moisture of air towards itself. This moisture forms droplets of water and hence the cloud

9. Match List I with List II and select the correct answer using the codes given below the lists

**List I**

- (a) Peroxide  
 (b) Superoxide  
 (c) Dioxide  
 (d) Trioxide

**List II**

- (1)  $C_3O_2$   
 (2)  $PbO_2$   
 (3)  $KO_2$   
 (4)  $H_2O_2$

Codes :

- (a) A B C D  
 4 3 2 1  
 (b) A B C D  
 3 2 1 4  
 (c) A B C D  
 4 2 3 1  
 (d) A B C D  
 4 1 2 3

[NDA 1999]

10. The most efficient agent for the absorption of  $SO_3$  is

- (a) 98%  $H_2SO_4$  (b) 80%  $H_2SO_4$   
 (c) 20% oleum (d) 90%  $H_2SO_4$

[KCET 1998]

11. Mark the oxide which is amphoteric in character

- (a)  $CO_2$  (b)  $SiO_2$   
 (c)  $SnO_2$  (d)  $CaO$

[MP PMT 2000]

12. Concentrated aqueous sodium hydroxide can separate mixture of

- (a)  $Al^{3+}$  and  $Sn^{2+}$  (b)  $Al^{3+}$  and  $Fe^{3+}$   
 (c)  $Al^{3+}$  and  $Zn^{2+}$  (d)  $Zn^{2+}$  and  $Pb^{2+}$

[MNR 1995]

13. The composition of the common glass is

- (a)  $Na_2O.CaO.6SiO_3$  (b)  $Na_2O.Al_2O_3.SiO_2$   
 (c)  $CaO.Al_2O_3.SiO_2$  (d)  $Na_2O.CaO.6SiO_2$

[DCE 2004]

## Critical Thinking

### Objective Questions

1. The correct sequence in decreasing order of the percentage of nitrogen in the given compounds is [NDA 1999]  
 (a) Urea > Ammonium chloride > Ammonium nitrate > Ammonium nitrite  
 (b) Urea > Ammonium nitrate > Ammonium nitrite > Ammonium chloride  
 (c) Urea > Ammonium nitrite > Ammonium nitrate > Ammonium chloride  
 (d) Urea > Ammonium nitrite > Ammonium chloride > Ammonium nitrate
2. As the alkaline earth metals (except Be) tend to lose their valence electrons readily they act as [Kerala (Med.) 2003]  
 (a) Weak oxidising agent (b) Weak reducing agent  
 (c) Strong oxidising agent (d) Strong reducing agent
3. The first ionisation energies of alkaline earth metals are higher than those of the alkali metals. This is because [UPSEAT 2001]  
 (a) There is increases in the nuclear charge of the alkaline earth metals  
 (b) There is decreases in the nuclear charge of the alkaline earth metals  
 (c) There is no change in the nuclear charge  
 (d) None of these
4. Lead is maximum in [BVP 2004]  
 (a) Soda glass (b) Jena glass  
 (c) Pyrex glass (d) Flint glass
5.  $BaSO_4$  and carbon on heating reacts to produce [Pb. PMT 2004]  
 (a)  $Ba + SO_2 + CO_2$  (b)  $BaS + CO$   
 (c)  $BaS + O_2 + SO_2$  (d)  $BaCO_3 + S + O_2$
6. The atomic radii of alkali metals (*M*) lie in the order  $Li < Na < K < Rb$  but the radii of  $M^+$  ions in aqueous solution lie in the reverse order  $Li^+ > Na^+ > K^+ > Rb^+$ . What is the reason for this reverse order (on going from *Li* to *Rb*)? [MP PMT 1997]  
 (a) Gradual increase in ionisation energy  
 (b) Increasing weakness of the metallic bond  
 (c) Increasing electropositive character


## 802 s and p-Block Elements

14. The metal which does not form ammonium nitrate by reaction with dilute nitric acid is  
 (a) *Al* (b) *Fe*  
 (c) *Pb* (d) *Mg*
15. Total number of lone pair of electrons in  $XeOF_4$  is  
 [IIT-JEE (Screening) 2004]  
 (a) 0 (b) 1  
 (c) 2 (d) 3
16. What is the correct relationship between the *pH*s of isomolar solutions of sodium oxide ( $pH_1$ ), sodium sulphide ( $pH_2$ ), sodium selenide ( $pH_3$ ) and sodium telluride ( $pH_4$ )  
 [CBSE PMT 2005]  
 (a)  $pH_1 > pH_2 = pH_3 > pH_4$   
 (b)  $pH_1 < pH_2 < pH_3 < pH_4$   
 (c)  $pH_1 < pH_2 < pH_3 = pH_4$   
 (d)  $pH_1 > pH_2 > pH_3 > pH_4$
17. Which one of the following is not an amphoteric substance  
 [KCET 2004]  
 (a)  $HNO_3$  (b)  $HCO_3^-$   
 (c)  $H_2O$  (d)  $NH_3$
18. Which group is called buffer group of the periodic table  
 [Pb. CET 2004]  
 (a) I (b) VII  
 (c) VIII (d) Zero
19. Which of the following salt is insoluble in water  
 [MP PET 2004]  
 (a)  $CuSO_4$  (b)  $CdSO_4$   
 (c)  $PbSO_4$  (d)  $Bi_2(SO_4)_3$
20. Which of the following oxides is the most acidic  
 [CBSE PMT 1999; MP PMT 2002]  
 (a)  $N_2O_5$  (b)  $P_2O_5$   
 (c)  $As_2O_5$  (d)  $Sb_2O_5$
21. Whose bond energy is maximum  
 [CPMT 1988; MP PMT 1990]  
 (a)  $F_2$  (b)  $Cl_2$   
 (c)  $Br_2$  (d)  $I_2$
22. Calcium cyanide on treatment with steam under pressure gives  $NH_3$  and  
 [DPMT 2002]  
 (a)  $CaHCO_3$  (b)  $CaO$   
 (c)  $Ca(OH)_2$  (d)  $CaCO_3$
23. Six volumes of oxygen, on complete ozonisation, form ..... volumes of ozone  
 (a) 4 (b) 3  
 (c) 2 (d) 6
24. The substance not likely to contain  $CaCO_3$  is  
 [AIEEE 2003]  
 (a) A marble statue (b) Calcined gypsum  
 (c) Sea shells (d) Dolomite
25. [KCET 2004] Which of the following statements is false for alkali metals  
 [MNR 1994; MP PET 2001]  
 (a) Lithium is the strongest reducing agent  
 (b) *Na* is amphoteric in nature  
 (c)  $Li^+$  is exceptionally small  
 (d) All alkali metals give blue solution in liquid ammonia
26. Solubility of iodine in water is greatly increased by the addition of iodide ions because of the formation of .....  
 [IIT 1994]  
 (a)  $I_2$  (b)  $I_3$   
 (c)  $I_3^-$  (d)  $I^-$
27. The solubility in water of sulphates down the *Be* group is  $Be > Mg \gg Ca > Sr > Ba$ . This is due to  
 [CBSE PMT 1995]  
 (a) High heat of solvation for smaller ions like  $Be^{2+}$   
 (b) Increasing molecular weight  
 (c) Decreasing lattice energy  
 (d) Increase in melting points
28. Magnesium burns in air to give  
 [CPMT 1988, 89; AFMC 1987]  
 (a)  $MgO$  (b)  $Mg_3N_2$   
 (c)  $MgCO_3$  (d)  $MgO$  and  $Mg_3N_2$
- both
29. Philosopher's wool when heated with  $BaO$  at  $1100^\circ C$  gives a compound. Identify the compound [CPMT 1987]  
 (a)  $BaZnO_2$  (b)  $Ba + ZnO_2$   
 (c)  $BaCdO_2$  (d)  $BaO_2 + Zn$
30. Which of the following oxide is diamagnetic  
 [MP PET 1990]  
 (a)  $NO$  (b)  $N_2O_4$   
 (c)  $NO_2$  (d)  $N_2O_5$
31. Which of the following salt becomes plaster of paris on being appropriately hydrated [CPMT 1985]  
 (a)  $ZnCO_3$  (b)  $CaSO_4$   
 (c)  $MgSO_4$  (d)  $CaCO_3$
32. The number of electron and proton in the third alkaline earth metal ion will be  
 (a)  $\frac{e}{20}, \frac{p}{20}$  (b)  $\frac{e}{18}, \frac{p}{20}$   
 [DPMT 2000]  
 (c)  $\frac{e}{18}, \frac{p}{18}$  (d)  $\frac{e}{19}, \frac{p}{20}$
33. The compounds of alkaline earth metals have the following magnetic nature  
 [MP PET/PMT 1998; RPMT 2000; JIPMER 2002]  
 (a) Diamagnetic (b) Paramagnetic

- (c) Ferromagnetic (d) Diaferromagnetic
34. Which of the following is the life saving mixture for an asthma patient [MP PMT 2001]  
 (a) Mixture of helium and oxygen  
 (b) Mixture of neon and oxygen  
 (c) Mixture of xenon and nitrogen  
 (d) Mixture of argon and oxygen
35. Which would quickly absorb oxygen [CBSE PMT 1992; MP PET 1995]  
 (a) Alkaline solution of pyrogallol  
 (b) Conc.  $H_2SO_4$   
 (c) Lime water  
 (d) Alkaline solution of  $CuSO_4$
36. Nitrogen is liberated by the thermal decomposition of only [IIT 1991]  
 (a)  $NH_4NO_2$  (b)  $NaN_3$   
 (c)  $(NH_4)_2Cr_2O_7$  (d) All the three
37. Red phosphorus is less reactive than yellow phosphorus because [DPMT 1982; JIPMER 1999; CBSE PMT 1999; RPET 2003]  
 (a) Its colour is red  
 (b) It is highly polymerised  
 (c) It is hard  
 (d) It is insoluble in  $C_2H_5OH$
38. Carbon differs from other elements of the group. Which is the false statement  
 (a) Due to its marked tendency to form long chains (catenation)  
 (b) Due to its unique ability to form multiple bonds  
 (c) Due to  $d$ -orbital in penultimate shell  
 (d) Due to its limitation of co-ordination number 4
39. Which of the following oxide does not form acidic aqueous solution [CPMT 2004]  
 (a)  $N_2O_3$  (b)  $NO_2$   
 (c)  $N_2O_5$  (d)  $NO$
40. Which of the following is in the increasing order of the ionic character [JIPMER 2002]  
 (a)  $PbCl_4 < PbCl_2 < CaCl_2 < NaCl$   
 (b)  $PbCl_2 < PbCl_4 < CaCl_2 < NaCl$   
 (c)  $PbCl_2 < PbCl_4 < NaCl < CaCl_2$   
 (d)  $PbCl_4 < PbCl_2 < NaCl < CaCl_2$
41. Silicon chloroform is prepared by [MH CET 1999]  
 (a)  $Si + HCl$  (b)  $SiCl_4 + H_2O$   
 (c)  $SiF_4 + NaF$  (d)  $H_2SiF_6 + Cl_2$
42.  $KO_2$  (potassium superoxide) is used in oxygen cylinders in space and submarines because it [AIEEE 2002]  
 (a) Absorbs  $CO_2$  and increases  $O_2$  content  
 (b) Eliminates moisture  
 (c) Absorbs  $CO_2$   
 (d) Produces ozone
43. Fire extinguishers contain  $H_2SO_4$  and [AFMC 1980]  
 (a)  $CaCO_3$  (b)  $Na_2CO_3$   
 (c)  $NaHCO_3$  (d)  $NaHCO_3$  and  $Na_2CO_3$
44. Which is insoluble in water [CPMT 2003]  
 (a)  $H_2S$  (b)  $HgCl_2$   
 (c)  $Ca(NO_3)_2$  (d)  $CaF_2$
45. Which of the following halides is most acidic [KCET 1996]  
 (a)  $PCl_3$  (b)  $SbCl_3$   
 (c)  $BiCl_3$  (d)  $CCl_4$
46. The stability of the following alkali metal chlorides follows the order  
 (a)  $LiCl > KCl > NaCl > CsCl$   
 (b)  $CsCl > KCl > NaCl > LiCl$   
 (c)  $NaCl > KCl > LiCl > CsCl$   
 (d)  $KCl > CsCl > NaCl > LiCl$
47. The reaction of  $Na_2S_2O_3$  with iodine gives [CPMT 1971, 80, 81; DPMT 1983, 90; MP PMT 1985; EAMCET 1990; BHU 1980]  
 (a) Sodium sulphide (b) Sodium sulphite  
 (c) Sodium sulphate (d) Sodium tetrathionate
48. Which one of the following is the true covalent oxide of iodine [MP PET/PMT 1988]  
 (a)  $I_2O_4$  (b)  $I_2O_5$   
 (c)  $I_2O_7$  (d)  $I_2O_9$
49. Lithium aluminium hydride acts as [CPMT 1994]  
 (a) Oxidising agent (b) Reducing agent  
 (c) Both the above (d) None of these
50. The mixture of conc.  $HCl$  and potassium chlorate on heating gives [Roorkee 2000]  
 (a)  $Cl_2$  only (b)  $ClO_2$  only  
 (c)  $Cl_2 + ClO_2$  (d)  $Cl_2 + ClO_2 + ClO_3$
51. When  $SO_2$  is passed through acidified solution of  $H_2S$  [CPMT 1973, 81, 93]  
 (a)  $H_2SO_4$  is formed (b)  $H_2SO_3$  is formed  
 (c) Sulphur is precipitated (d) None of these
52. Four reactions are given below  
 (i)  $2Li + 2H_2O \rightarrow 2LiOH + H_2$   
 (ii)  $2Na + 2H_2O \rightarrow 2NaOH + H_2$   
 (iii)  $2LiNO_3 \xrightarrow{\text{Heat}} 2LiNO_2 + O_2$



## 804 s and p-Block Elements



Which of the above, if any, is wrong

- (a) (iv) (b) (iii)  
(c) (i) (d) None of these
53. Increasing order of solubility is [AFMC 1987]  
(a)  $CaCO_3, KHCO_3, NaHCO_3$   
(b)  $NaHCO_3, KHCO_3, CaCO_3$   
(c)  $KHCO_3, NaHCO_3, CaCO_3$   
(d)  $CaCO_3, NaHCO_3, KHCO_3$
54. Nitrolim is [CPMT 1976, 78, 2000; BHU 1987; DCE 1999; RPMT 2000]  
(a)  $Ca(NO_3)_2$  (b)  $Ca(CN)_2$   
(c)  $CaCN_2 + C$  (d)  $CaCN_2$
55. The following acids have been arranged in the order of decreasing acid strength. Identify the correct order.  
(I)  $ClOH$  (II)  $BrOH$  (III)  $IOH$   
(a)  $I > II > III$  (b)  $II > I > III$   
(c)  $III > II > I$  (d)  $I > III > II$
56. Which of the following element does not belong to the family of elements indicated [Orissa JEE 1997]  
(a) Rubidium ( $Rb, Z = 37$ ) : Alkali metals  
(b) Barium ( $Ba, Z = 56$ ) : Alkaline earth metals  
(c) Iridium ( $I, Z = 77$ ) : Nobel gases  
(d) Argon ( $Ar, Z = 18$ ) : Nobel gases
57.  $H_3PO_2$  is the formula for one of the phosphorus acid. Its name and basicity are respectively [CBSE PMT 1992; BHU 1999; KCET 1999]  
(a) Phosphorus acid and two  
(b) Hypophosphorus acid and two  
(c) Hypophosphorus acid and one  
(d) Hypophosphoric acid and two
58. Which of the following oxides of nitrogen is paramagnetic [CPMT 1984; CBSE PMT 1994; AIIMS 2000]  
(a)  $N_2O_3$  (b)  $N_2O$   
(c)  $NO_2$  (d)  $N_2O_5$
59. Nessler's reagent is [CPMT 2002]  
(a) Potassium in mercuric iodide  
(b)  $TiCl_4$   
(c) Anhydrous  $AlCl_3$   
(d)  $Al_2O_3 / Cr_2O_3$
60. The noble gas was first time discovered by  
(a) Cavandish (b) William Ramsay  
(c) Rayleigh (d) Frankland
61. The ratio of  $\frac{C_p}{C_v}$  for inert gases is [DCE 1999]  
(a) 1.99 (b) 2.13  
(c) 1.66 (d) 1.33
62. White  $P$  reacts with caustic soda, the products are  $PH_3$  and  $NaH_2PO_2$ . This reaction is an example of [DCE 2000]  
(a) Oxidation (b) Reduction  
(c) Disproportionation (d) Neutralisation
63. The oxide, which cannot act as a reducing agent, is [Pb. CET 2002]  
(a)  $NO_2$  (b)  $SO_2$   
(c)  $CO_2$  (d)  $ClO_2$
64. Which of the following product is formed when  $SiF_4$  reacts with water [Pb. CET 2003]  
(a)  $SiF_3$  (b)  $H_4SiO_4$   
(c)  $H_2SO_4$  (d)  $H_2SiF_4$
65. Ozone with dry iodine give [Pb. CET 2003]  
(a)  $I_4O_4$  (b)  $I_2O_3$   
(c)  $I_2O_5$  (d)  $I_2O_4$
66. The microcosmic salt is [Pb. CET 2004; Pb. PMT 2004]  
(a)  $Na(NH_4)H_2O$  (b)  $K(NH_4)HPO_3 \cdot 2H_2O$   
(c)  $Na(NH_4)HPO_4 \cdot 4H_2O$  (d)  $Na(NH_3)HPO_4 \cdot 4H_2O$
67. Thermitite is a mixture of [Pb. CET 2004]  
(a)  $Cr_2O_3 + Al_2O_3$  (b)  $Fe_2O_3 + Al$   
(c)  $Fe_2O_3 + Al_2O_3$  (d)  $Al_2O_3 + 2Cr$
68. The colour of liquid  $O_2$  is [BVP 2004]  
(a) Red (b) Dark blue  
(c) Pale yellow (d) Pale blue
69. Which of the following gas mixture is used by the divers inside the sea [AFMC 2004]  
(a)  $O_2 + He$  (b)  $O_2 + Xe$   
(c)  $O_2 + Ar$  (d)  $O_2 + N_2$
70. One mole of magnesium nitride on the reaction with an excess of water gives  
(a) Two moles of ammonia (b) One mole of nitric acid  
(c) One mole of ammonia (d) Two moles of nitric acid
71. Calcium cyanamide on treatment with steam produce [Pb. PMT 2004]  
(a)  $CaCO_3 + NH_3$  (b)  $CaHCO_3 + NH_3$   
(c)  $CaO + NH_3$  (d)  $Ca(OH)_2 + NH_3$

## Assertion & Reason

For AIIMS Aspirants



Read the assertion and reason carefully to mark the correct option out of the options given below :

- (a) If both assertion and reason are true and the reason is the correct explanation of the assertion.  
 (b) If both assertion and reason are true but reason is not the correct explanation of the assertion.  
 (c) If assertion is true but reason is false.  
 (d) If the assertion and reason both are false.  
 (e) If assertion is false but reason is true.

1. Assertion : Sulphate is estimated as  $BaSO_4$  and not as  $MgSO_4$ .  
Reason : Ionic radius of  $Mg^{2+}$  is smaller than that of  $Ba^{2+}$ . [IIT 1998]
2. Assertion : Amongst the halogens fluorine can oxidise the elements to highest oxidation state.  
Reason : Due to small size of fluoride ion, it is difficult to oxidise fluoride ion to fluorine. Hence reverse reaction takes place more easily. [IIT 1996]
3. Assertion :  $HNO_3$  is a stronger acid than  $HNO_2$ .  
Reason : In  $HNO_3$  there are two nitrogen-to-oxygen bonds whereas in  $HNO_2$  there is only one.
4. Assertion : The Value of Vander Waal's constant 'a' is larger for ammonia than for nitrogen.  
Reason : Hydrogen bonding is present in ammonia. [IIT 1998]
5. Assertion : Xenon forms fluorides.  
Reason : Due to the strong electronegativity of fluorine. [AIIMS 2001]
6. Assertion : Chlorine and sulphur dioxide both are bleaching agents.  
Reason : The bleaching action of chlorine and sulphur dioxide is performed through the process of oxidation.[AIIMS 2000]
7. Assertion : Nitrogen is unreactive at room temperatures but becomes reactive at elevated temperatures (On heating) or in presence of catalyts.  
Reason : In nitrogen molecule, there is extensive delocalization of electrons.
8. Assertion : Covalency of oxygen is three.  
Reason : Dinegative anion of oxygen ( $O^{2-}$ ) is quite common but dinegative anion of sulphur ( $S^{2-}$ ) is less common.[AIIMS 2001]
9. Assertion : At room temperature, oxygen exists as a diatomic gas, whereas sulphur exists as solid.

- Reason : The catenated  $-O-O-O-$  changes are less stable as compared to  $O=O$  molecule. [AIIMS 2001]
10. Assertion : Potassium and caesium are used in photo-electric cells.  
Reason : Potassium and caesium emit electrons on exposure to light.[AIIMS 2002]
  11. Assertion : The fluorine has lower reactivity.  
Reason :  $F-F$  bond has low bond dissociation energy. [AIIMS 2002]
  12. Assertion : Halogens do not occur in free state.  
Reason : Halogens are highly reactive. [AIIMS 1994]
  13. Assertion : Lithium forms Lithium oxide ( $LiO_2$ ).  
Reason :  $N_2$  molecule have unpaired electrons. [AIIMS 1995]
  14. Assertion : Liquid  $NH_3$  is used for refrigeration.  
Reason : Liquid  $NH_3$  quickly vaporises. [AIIMS 1995]
  15. Assertion :  $Al(OH)_3$  is insoluble in  $NH_4OH$  but soluble in  $NaOH$ .  
Reason :  $NaOH$  is strong alkali. [AIIMS 1997]
  16. Assertion : Boron is metalloid.  
Reason : Boron shows metallic nature. [AIIMS 1997]
  17. Assertion : Inert gases are monoatomic.  
Reason : Inert gases have stable configuration. [AIIMS 1999]
  18. Assertion : Magnesium continue to burn in nitric oxide.  
Reason : During burning heat evolved do not decompose  $NO$ . [AIIMS 2001]
  19. Assertion : Anhydrous  $BaO_2$  is used for preparing  $H_2O_2$ .  
Reason : Hydrated  $BaO_2$  is not available. [AIIMS 2001]
  20. Assertion : Benzene is reactive while inorganic benzene is unreactive compound.  
Reason : Inorganic benzene is, borazine,  $B_3N_3H_6$ . [AIIMS 2002]
  21. Assertion : Halogens absorb visible light.  
Reason : All halogens are coloured.[AIIMS 2002]
  22. Assertion : Barium is not required for normal biological function in human.  
Reason : Barium does not show variable oxidation state. [AIIMS 2003]
  23. Assertion : The  $O-O$  bond length in  $H_2O_2$  is shorter than that of  $O_2F_2$ .  
Reason :  $H_2O_2$  is an ionic compound.[AIIMS 2003]

## 806 s and p-Block Elements

24. Assertion :  $PbI_4$  is a stable compound.  
Reason : Iodide stabilizes higher oxidation state.  
[AIIMS 2003]
25. Assertion :  $Mg$  is not present in enamel of human teeth.  
Reason :  $Mg$  is an essential element for biological functions of human. [AIIMS 2004]
26. Assertion : Radium is most abundant s-block element.  
Reason : s-block elements are non-radioactive in nature.
27. Assertion :  $LiCl$  is predominantly a covalent compound.  
Reason : Electronegativity difference between  $Li$  and  $Cl$  is too small.
28. Assertion : The first ionization energy of  $Be$  is greater than that of  $B$ .  
Reason :  $2p$ -orbital is lower in energy than  $2s$ -orbital.
29. Assertion : The alkali metals can form ionic hydrides which contains the hydride ion.  
Reason : The alkali metals have low electronegativity, their hydrides conduct electricity when fused and liberate hydrogen at the anode.
30. Assertion :  $Be$  does not impart any characteristic colour to the bunsen flame.  
Reason : Due to its very high ionization energy, beryllium requires a large amount of energy for excitation of the electrons.
31. Assertion : Potassium is not obtained by the electrolysis of fused  $KCl$ .  
Reason : Potassium vapourises at the melting point of  $KCl$ .
32. Assertion : Helium and beryllium have similar outer electronic configuration of the type  $ns^2$ .  
Reason : Both are chemically inert.
33. Assertion :  $Na_2SO_4$  is soluble while  $BaSO_4$  is insoluble.  
Reason : Lattice energy of  $BaSO_4$  exceeds its hydration energy.
34. Assertion : Alkali metals impart colour to the flame.  
Reason : Their ionisation energies are low.
35. Assertion : Superoxides of alkali metals are paramagnetic.  
Reason : Superoxides contain the ion  $O^{2-}$  which has one unpaired electron.
36. Assertion : Although  $PF_5$ ,  $PCl_5$  and  $PBr_3$  are known, the pentahalides of nitrogen have not been observed.  
Reason : Phosphorus has lower electronegativity than nitrogen.
37. Assertion : The electronic structure of  $O_3$  is  
  
  
Reason : Structure is not allowed because octet around  $O$  cannot be expanded.
38. Assertion : Sulphuric acid is more viscous than water.  
Reason : Concentrated sulphuric acid has a great affinity for water.
39. Assertion :  $PCl_5$  is covalent in gaseous and liquid states but ionic in solid state.  
Reason :  $PCl_5$  in solid state consists of tetrahedral  $PCl_4^+$  cation and octahedral  $PCl_6^-$  anion.
40. Assertion : Among nitrogen halides  $NX_3$ , the dipole moment is highest for  $NI_3$  and lowest for  $NF_3$ .  
Reason : Nitrogen halides  $NX_3$ , have trigonal pyramidal structure.
41. Assertion : White phosphorus is stored under water.  
Reason : White phosphorous is highly reactive and catches fire spontaneously in air.
42. Assertion :  $Al$  forms  $[AlF_6]^{3-}$  but  $B$  does not form  $[BF_6]^{3-}$ .  
Reason :  $B$  does not react with  $F_2$ .
43. Assertion :  $NO_3^-$  is planar while  $NH_3$  is pyramidal.  
Reason :  $N$  in  $NO_3^-$  is  $sp^2$  hybridized but in  $NH_3$  it is  $sp^3$ - hybridized.
44. Assertion :  $Si-Si$  bonds are much stronger than  $Si-O$  bonds.  
Reason : Silicon forms double bonds with itself.
45. Assertion : The  $S-S-S$  bond angle in  $S_8$  molecule is  $105^\circ$ .  
Reason :  $S_8$  has a V-shape.





46. Assertion : Caro's acid has S atom in +6 oxidation state.  
Reason : Caro's acid contains one peroxo  $O_2^{2-}$  group.
47. Assertion : The m.p./b.p. of noble gases are quite high.  
Reason : The interparticle forces among noble gases in their liquid state are covalent forces.
48. Assertion : In  $SO_2$ , the bond angle is  $119^\circ$  whereas in  $SO_3$ , the bond angle is  $120^\circ$ .  
Reason : S atom in both  $SO_2$  and  $SO_3$  is  $sp^2$ -hybridized.
49. Assertion : Calcium carbide on hydrolysis gives methane.  
Reason : Calcium carbide contains  $C^{4-}$  anions.
50. Assertion : Xenon forms fluorides.  
Reason : Because 5 d-orbitals are available for valence shell expansion.
51. Assertion : Hydrogen cannot be prepared in laboratory.  
Reason : Hydrogen of high purity is obtained by electrolysis of warm aqueous barium hydroxide between nickel electrodes.
52. Assertion : Diprothium shows relatively inert behaviour at room temperature.  
Reason : The values of melting point and boiling point for dideuterium are higher as compared to diprothium.
53. Assertion : Water can be transformed from liquid to solid state only.  
Reason : The distribution of water over the earth surface is uniform.
54. Assertion : Ice is less dense than water.  
Reason : Ice is a solid whereas water is liquid.
55. Assertion :  $HF, NH_3$  and  $H_2O$  form intermolecular hydrogen bonding.  
Reason :  $HF, NH_3$  and  $H_2O$  molecules are bonded in same manner.
56. Assertion : Hard water does not lather with soap.  
Reason : In hard water, the sodium stearate of soap changes to the corresponding calcium magnesium salt which precipitates out.
57. Assertion :  $H_2O_2$  is stored in wax-lined glass.  
Reason : Presence of metal surfaces, traces of alkali (present in glass) etc. increases its decomposition.
58. Assertion : A nearly tetrahedral arrangement of the orbitals about the oxygen atom allows each water molecule to form hydrogen bonds with as many as four neighbouring water molecules.  
Reason : In ice each water molecule forms four hydrogen bonds as each molecule is fixed in the space.
59. Assertion : Calgon is used for removing  $Ca^{2+}$  and  $Mg^{2+}$  ions from hard water.  
Reason : Calgon forms precipitate with  $Ca^{2+}$  and  $Mg^{2+}$  ions.
60. Assertion : Reaction of  $SO_2$  and  $H_2S$  in the presence of  $Fe_2O_3$  catalyst gives elemental sulphur.  
Reason :  $SO_2$  is a reducing agent. [AIIMS 2005]
61. Assertion :  $SiF_6^{2-}$  is known but  $SiCl_6^{2-}$  is not.  
Reason : Size of fluorine is small and its lone pair of electrons interacts with d-orbitals of Si strongly. [AIIMS 2005]
62. Assertion : Borax bead test is not suitable for Al(III).  
Reason :  $Al_2O_3$  is insoluble in water. [AIIMS 2005]
63. Assertion :  $SeCl_4$  does not have a tetrahedral structure.  
Reason : Se in  $SeCl_4$  has two lone pairs. [AIIMS 2005]
64. Assertion : Ozone is a powerful oxidizing agent in comparison to  $O_2$ .  
Reason : Ozone is diamagnetic but  $O_2$  is paramagnetic. [AIIMS 2005]

# Answers

## Alkali metals

1	b	2	c	3	a	4	b	5	a
6	b	7	c	8	a	9	c	10	d
11	c	12	b	13	a	14	d	15	a
16	b	17	c	18	b	19	b	20	b
21	a	22	b	23	b	24	a	25	d
26	b	27	b	28	d	29	c	30	c
31	d	32	a	33	b	34	a	35	d
36	b	37	a	38	a	39	d	40	b
41	d	42	c	43	b	44	d	45	d

## 808 s and p-Block Elements

46	b	47	b	48	b	49	c	50	c
51	d	52	c	53	d	54	d	55	a
56	c	57	a	58	c	59	d	60	a
61	b	62	b	63	a	64	a	65	c
66	b	67	d	68	a	69	d	70	d
71	c	72	a	73	a	74	c	75	b
76	d	77	c	78	a	79	b	80	a
81	d	82	a	83	c	84	b	85	c
86	c	87	d	88	b	89	c	90	a
91	c	92	a	93	c	94	a	95	c
96	a	97	a	98	b	99	d	100	c
101	a	102	d	103	b	104	c	105	c
106	a	107	c	108	b	109	b	110	d
111	d	112	c	113	a	114	c	115	a
116	a	117	c	118	c	119	b	120	a
121	b	122	c	123	c	124	c	125	d
126	c	127	a	128	b	129	a,b	130	b
131	d	132	b	133	d	134	d	135	b
136	d	137	d	138	a	139	a	140	b
141	c	142	c	143	a	144	a	145	b
146	d	147	b	148	d	149	c	150	b

### Alkaline earth metals

1	c	2	d	3	a	4	b	5	b
6	d	7	d	8	d	9	b	10	a
11	a	12	d	13	a	14	a	15	a
16	d	17	c	18	c	19	b	20	b
21	b	22	d	23	d	24	a	25	a
26	bc	27	d	28	d	29	b	30	b
31	d	32	d	33	b	34	b	35	b
36	d	37	c	38	d	39	d	40	a
41	c	42	a	43	a	44	a	45	a
46	c	47	b	48	d	49	c	50	c
51	a	52	c	53	b	54	a	55	b
56	a	57	d	58	a	59	b	60	b
61	d	62	b	63	d	64	a	65	a
66	d	67	a	68	c	69	d	70	a
71	c	72	a	73	b	74	c	75	b
76	d	77	b	78	d	79	c	80	c
81	d	82	a	83	c	84	a	85	c
86	d	87	c	88	d	89	a	90	a
91	b	92	c	93	c	94	b	95	a
96	d	97	d	98	d	99	a	100	b

101	b	102	c	103	a	104	b	105	c
106	a	107	a	108	c	109	d	110	d
111	b								

### Boron family

1	a	2	c	3	a	4	a	5	c
6	d	7	a	8	c	9	d	10	c
11	d	12	b	13	c	14	c	15	c
16	acd	17	a	18	d	19	e	20	a
21	c	22	c	23	d	24	c	25	a
26	d	27	a	28	c	29	a	30	d
31	d	32	c	33	c	34	c	35	d
36	b	37	c	38	c	39	a	40	a
41	c	42	a	43	a	44	d	45	b
46	b	47	c	48	d	49	a	50	c
51	c	52	b	53	a	54	d	55	b
56	b	57	c	58	b	59	b	60	b
61	d	62	d	63	c	64	b	65	c
66	c	67	c	68	a	69	d	70	c
71	c	72	c	73	c	74	a	75	a
76	b	77	a						

### Carbon family

1	c	2	d	3	d	4	c	5	a
6	a	7	b	8	d	9	c	10	d
11	c	12	b	13	d	14	b	15	c
16	a	17	b	18	b	19	d	20	d
21	c	22	c	23	a	24	a	25	c
26	a	27	c	28	a	29	a	30	d
31	a	32	a	33	d	34	b	35	b
36	b	37	c	38	b	39	b	40	b
41	a	42	c	43	b	44	c	45	d
46	c	47	c	48	b	49	d	50	a
51	c	52	c	53	a	54	a	55	a
56	c	57	b	58	b	59	a	60	b
61	d	62	a	63	d	64	c	65	d
66	b	67	c	68	a	69	b		

### Nitrogen family

1	b	2	a	3	b	4	b	5	a
6	a	7	d	8	b	9	a	10	b
11	ad	12	b	13	a	14	b	15	d

## s and p-Block Elements 809

16	b	17	d	18	b	19	c	20	a
21	a	22	a	23	b	24	d	25	c
26	c	27	c	28	c	29	d	30	c
31	d	32	c	33	d	34	ab	35	a
36	a	37	b	38	d	39	b	40	c
41	b	42	a	43	b	44	d	45	b
46	d	47	d	48	a	49	b	50	b
51	a	52	d	53	c	54	d	55	d
56	c	57	b	58	c	59	c	60	c
61	a	62	d	63	b	64	a	65	b
66	c	67	c	68	a	69	a	70	b
71	a	72	a	73	a	74	b	75	a
76	b	77	c	78	b	79	c	80	d
81	d	82	d	83	d	84	c	85	d
86	d	87	c	88	b	89	d	90	c
91	a	92	b	93	d	94	b	95	d
96	c	97	c	98	b	99	a	100	d
101	d	102	b	103	d	104	a	105	d
106	d	107	d	108	a	109	d	110	a
111	a	112	c	113	d	114	c	115	a
116	a	117	c	118	b	119	a	120	c
121	b	122	b	123	d	124	d	125	c
126	b	127	b	128	c	129	d	130	b
131	c	132	b	133	b	134	b	135	b
136	a	137	a	138	d	139	a	140	d
141	a	142	c	143	b	144	a	145	a
146	d	147	b	148	b	149	a	150	d
151	d	152	d	153	a	154	e	155	a
156	a	157	c	158	d	159	b	160	d
161	c	162	b	163	c	164	b	165	d
166	a	167	a	168	a	169	b	170	d
171	c	172	d	173	a	174	d	175	c
176	b	177	d	178	c	179	b	180	a
181	b	182	a	183	d	184	c	185	a
186	d	187	d	188	a	189	a	190	d
191	a	192	a	193	c	194	a	195	b
196	c	197	b	198	a	199	b	200	c
201	d	202	c	203	b	204	a	205	b
206	a	207	c	208	d	209	c	210	c
211	a	212	c	213	d	214	b	215	b,c
216	b	217	a	218	d	219	d	220	c
221	a	222	a	223	a	224	b	225	a
226	d	227	b	228	b	229	b	230	c

231	a	232	a	233	c	234	a	235	d
236	a	237	b	238	d	239	b	240	c
241	a								

### Oxygen family

1	b	2	c	3	b	4	c	5	a
6	c	7	b	8	d	9	b	10	a
11	a	12	b	13	bc	14	d	15	d
16	a	17	b	18	c	19	a	20	a
21	a	22	d	23	a	24	c	25	c
26	d	27	d	28	b	29	b	30	b
31	a	32	d	33	a	34	b	35	a
36	a	37	d	38	c	39	a	40	a
41	c	42	a	43	d	44	a	45	b
46	a	47	d	48	b	49	b	50	c
51	d	52	c	53	d	54	c	55	d
56	b	57	a	58	c	59	c	60	a
61	a	62	d	63	a	64	a	65	c
66	c	67	a	68	a	69	c	70	d
71	c	72	a	73	c	74	c	75	b
76	b	77	b	78	c	79	d	80	a
81	d	82	d	83	d	84	e	85	b
86	c	87	d	88	c	89	d	90	b
91	c	92	a	93	a	94	b	95	c
96	c	97	b	98	c	99	b	100	b
101	a	102	d	103	b	104	b	105	c
106	a	107	a	108	d				

### Halogen family

1	b	2	a	3	d	4	c	5	a
6	a	7	a	8	a	9	c	10	b
11	a	12	a	13	d	14	a	15	b
16	d	17	a	18	b	19	a	20	c
21	d	22	a	23	c	24	a	25	c
26	b	27	d	28	d	29	bd	30	d
31	d	32	a	33	d	34	a	35	d
36	a	37	b	38	a	39	d	40	a
41	d	42	b	43	d	44	a	45	a
46	a	47	b	48	a	49	d	50	b
51	a	52	a	53	d	54	a	55	a
56	d	57	a	58	c	59	a	60	d
61	b	62	c	63	a	64	a	65	d
66	b	67	b	68	b	69	c	70	c

## 810 s and p-Block Elements

71	b	72	a	73	a	74	b	75	d
76	b	77	b	78	c	79	c	80	b
81	b	82	c	83	d	84	d	85	d
86	a	87	a	88	a	89	d	90	b
91	d	92	c	93	c	94	a	95	a
96	c	97	b	98	b	99	c	100	a
101	a	102	c	103	d	104	b	105	d
106	e	107	b	108	a	109	a	110	a
111	c	112	c	113	a	114	a	115	a
116	c	117	b	118	b	119	b	120	b
121	d	122	a	123	b	124	a	125	a
126	c	127	d	128	a	129	a	130	c
131	d	132	a	133	a	134	a	135	a
136	c	137	d	138	b	139	d	140	a
141	d	142	d	143	d	144	b	145	a
146	a	147	a	148	b	149	c	150	a
151	d	152	b	153	d	154	a	155	c
156	b	157	b	158	a	159	c	160	d
161	d	162	a	163	b	164	a	165	a
166	c	167	c	168	c	169	b	170	a
171	a	172	b	173	c	174	c	175	b
176	b	177	c	178	b	179	a	180	d
181	a	182	a	183	a	184	c	185	a
186	a	187	c						

16	d	17	a	18	d	19	c	20	a
21	b	22	d	23	a	24	b	25	b
26	c	27	a	28	d	29	a	30	b
31	b	32	b	33	a	34	a	35	a
36	d	37	b	38	c	39	d	40	a
41	a	42	a	43	d	44	d	45	a
46	b	47	d	48	b	49	b	50	c
51	c	52	b	53	d	54	c	55	a
56	c	57	c	58	c	59	a	60	b
61	c	62	c	63	c	64	b	65	a
66	c	67	b	68	d	69	a	70	a
71	a								

## Noble gases

1	b	2	b	3	a	4	c	5	c
6	d	7	c	8	a	9	d	10	b
11	c	12	c	13	c	14	c	15	b
16	d	17	b	18	b	19	b	20	a
21	a	22	c	23	d	24	b	25	b
26	d	27	a	28	a	29	b	30	c
31	d	32	a	33	c	34	a	35	a
36	c	37	d	38	b	39	d	40	a
41	d	42	b	43	a	44	c	45	d
46	a	47	a	48	a	49	b	50	d
51	d	52	d	53	d	54	c	55	a
56	c	57	a	58	d	59	d	60	d

## Critical Thinking Questions

1	c	2	d	3	a	4	d	5	b
6	d	7	a	8	b	9	a	10	a
11	c	12	b	13	d	14	c	15	b

Assertion and Reason

1	b	2	b	3	a	4	a	5	a
6	c	7	b	8	e	9	a	10	a
11	e	12	a	13	a	14	a	15	a
16	c	17	b	18	c	19	d	20	d
21	a	22	b	23	d	24	d	25	b
26	d	27	c	28	c	29	a	30	a
31	a	32	c	33	b	34	a	35	a
36	b	37	a	38	b	39	b	40	b
41	a	42	c	43	a	44	d	45	c
46	a	47	d	48	b	49	d	50	a
51	e	52	b	53	d	54	b	55	c
56	a	57	a	58	a	59	d	60	b
61	a	62	b	63	c	64	b		

# AS Answers and Solutions

Alkali metals

- (b) Element 

Na	K	
$IE_1$	496	419
$IE_2$	4562	3051

  
Sodium has higher I.E. because of smaller atomic size.
- (c) Alkali metals are highly reactive metals. They react with  
Alcohol -  $2C_2H_5OH + 2K \rightarrow 2C_2H_5OK + H_2$   
Water -  $2K + 2H_2O \rightarrow 2KOH + H_2$   
Ammonia -  $K + (x + y)NH_3 \rightarrow [K(NH_3)_x]^+ + [e(NH_3)_y]^-$   
Ammoniated cation  
Ammoniated electron  
But they do not react with kerosene.
- (b) After removal of an electron the effective nuclear charge per electron increases hence the size decreases.
- (a) Alkali metals valence shell configuration =  $ns^1$
- (b) Element - 

Li	Na	K	Rb	Cs	
Ionic radius	76	102	138	152	167
- (pm)					

  
as the atomic no. increases the no. of shells increases hence, atomic radius increases.
- (c) On moving down the group electropositive character increases.

- (a) Carnellite -  $KCl \cdot MgCl_2 \cdot 6H_2O$   
Cryolite -  $Na_3AlF_6$   
Bauxite -  $(Al_2O_3 \cdot 2H_2O)$   
Dolomite -  $MgCO_3 \cdot CaCO_3$
- (d) Element - 




Li	Na	K	Rb	
Atomic radius (pm)	152	186	227	248
- (b) Li is much softer than the other group I metals. Actually Li is harder than other alkali metals
- (a)  $Cu^{+2} + 2e^- \rightarrow Cu, E^\circ = +0.34 V$   
 $Mg^{+2} + 2e^- \rightarrow Mg, E^\circ = -2.37 V$   
 $Na^+ + e^- \rightarrow Na, E^\circ = -2.71 V$
- (d) Anhydrous form of  $Na_2CO_3$  does not decompose on heating even to redness. It is an amorphous powder called soda ash.
- (c) Fehling's solution is a mixture of Alk.  $CuSO_4 + Na - K$  tartarate (Rochelle salt)
- (b)  $2K + 2HCl \rightarrow 2KCl + H_2$  (violent reaction).
- (b) Although lattice energy of LiCl higher than NaCl but LiCl is covalent in nature and NaCl ionic there after, the melting point decreases as we move NaCl because the lattice energy decreases as a size of alkali metal atom increases (lattice energy  $\propto$  melting point of alkali metal halide)
- (b) It form calcium and magnesium complex with EDTA salt.
- (a)  $LiOH < NaOH < KOH < RbOH$   
Down the group basic character increases
- (d)  $Na_2CO_3 \cdot 10H_2O \xrightarrow{\Delta} Na_2CO_3 \cdot H_2O \xrightarrow{\Delta}$   
washing powder  
 $Na_2CO_3 + H_2O \uparrow$
- (b)  $Na_2CO_3, K_2CO_3$  and  $(NH_4)_2CO_3$  are soluble in water because hydration energy is more than lattice energy
- (c)  $K_2SO_4 \cdot Al_2(SO_4)_3 \cdot 24H_2O$  potash alum it is a double salt.
- (d) It is a colourless gas.
- (a)  $NaHCO_3 \rightarrow Na^+ + HCO_3^-$   
(Salt of strong base & weak acid)  
 $\downarrow$   
 $OH^- + CO_2$
- (b)  $FeSO_4 \cdot (NH_4)_2SO_4 \cdot 6H_2O$  Mohr's salt.
- (d)  $Ca^{+2} > Na^+ > Mg^{+2} > Al^{+3}$
- (b)  $Li^+ + e^- \rightarrow Li, E^\circ = -3.05 V$   
 $K^+ + e^- \rightarrow K, E^\circ = -2.93 V$   
 $Ca^{+2} + 2e^- \rightarrow Ca, E^\circ = -2.87 V$
- (a) Because their valence electrons are present in s- orbitals.
- (a)  $6Li + N_2 \rightarrow 2Li_3N$  Lithium nitride.

## 810 s and p-Block Elements




39. (d) *Li, Na, K* are lighter than water but *Rb* is heavier than water.
42. (c)  $KF + HF \rightarrow KHF_2 \rightleftharpoons K^+ + HF_2^-$
43. (b)  $Cs > Rb > K > Na > Li$   
Metallic character decreasing order.
45. (d)  $2Rb + 2H_2O \rightarrow 2RbOH + H_2$   
 $\frac{Li < Na < K < Rb < Cs}{\rightarrow}$   
As we go down the group reactivity with  $H_2O$  increases.
48. (b) Atomic number 11  $\rightarrow Na \rightarrow Na_2O$   
 $Na_2O + H_2O \rightarrow 2NaOH$   
(base)
51. (d) Generally ionic character decreasing from *LiCl* to *NaCl*.
52. (c) In castner process *Na* metal is made of anode.
55. (a) Fajan's rule is applied.
57. (a) Small atomic and ionic size leads to high electronegativity and hydration energy. Small atomic and ionic size leads to high electronegativity and hydration energy.
58. (c) Mohr salt is  $(FeSO_4)(NH_4)_2SO_4 \cdot 6H_2O$ .
60. (a) Sodium thiosulphate is a reducing agent which convert metallic silver into silver salt.
64. (a) In alkali metal group elements alkali means plant ash.
67. (d)  $2Na + 2NH_3 \xrightarrow{\text{heat}} 2NaNH_2 + H_2$
68. (a,b)  $2Na + \frac{1}{2}O_2 \xrightarrow{\text{moist air}} Na_2O$   
 $Na_2O + 2H_2O \longrightarrow 2NaOH + H_2$ .
69. (d)  $2KClO_3 \rightarrow 2KCl + 3O_2$ .
70. (d) Due to free electron liquid ammonia becomes paramagnetic.
72. (a) They possess highest atomic volume in their respective periods.
74. (c)  $Fe(OH)_3$  is soluble in sodium hydroxide solution.
76. (d) The cell involves the following reaction,  
 $NaCl \rightleftharpoons Na^+ + Cl^-$   
At anode :  $2Cl^- \rightarrow 2Cl + 2e \rightarrow Cl_2$   
At cathode :  $Na^+ + e \rightarrow Na$   
 $Na + Hg \rightarrow \text{amalgam}$   
At anode :  $Na - \text{amalgam} \rightarrow Na^+ + Hg + e$   
At cathode :  $2H_2O + 2e \rightarrow H_2 \uparrow + 2OH^-$
78. (a) *Li* is a more reducing agent compare to other element.
79. (b) Element - 

<i>Li</i>	<i>Na</i>	<i>K</i>	<i>Rb</i>	<i>Cs</i>
M.pt in K - 4535	370.8	336.2	312	301.5
80. (a)  $2Na + 2HOH \rightarrow 2NaOH + H_2 \uparrow$   
 $2K + 2HOH \rightarrow 2KOH + H_2 \uparrow$
82. (a) Alkali metal are good conductor of heat and electricity.
83. (c) Potassium react with halogens (chlorine) to gives violet colour flame.
84. (b) Mobility decreases from top to bottom because of the atomic size is increases.
85. (c) Lithium shows digonal relationships with *Mg*.
86. (c)  $K > Ca > C > Cl$   
Electropositive character in decreasing order.
87. (d)  $2NaCl \xrightarrow[\text{Molten}]{\text{Electrolysis}} 2Na + Cl_2$   
Cathode Anode
88. (b) When sodium bicarbonate ( $NaHCO_3$ ) is heated, sodium carbonate,  $CO_2$  and water are formed.  
 $2NaHCO_3 \xrightarrow{\Delta} Na_2CO_3 + CO_2 \uparrow + H_2O$   
Sodium carbonate
89. (c) Alum is used for softning of water.
90. (a) Only salts of (weak acid + strong base) and (strong acid + weak base) get hydrolysed (*i.e.*, show alkalinity or acidity in water).  $KClO_4$  is a salt of strong acid and strong base therefore it does not get hydrolysed in water.  
 $KClO_4 \rightleftharpoons K^+ + ClO_4^-$ ;  $H_2O \rightleftharpoons \underset{\substack{\downarrow \\ \text{Strong}}}{OH^-} + \underset{\substack{\downarrow \\ \text{Strong}}}{H^+}$   
 $\text{Strong} \quad \text{Strong}$
91. (c) Carbon dioxide does not help in burning, also it forms carbonate with alkali metals.
92. (a) When carbonate are heated they decompose to form the oxide. Sodium carbonate and potassium carbonate do not decompose. The carbonate become more difficult to decompose as we go down the group.
93. (c) Aluminium reacts with caustic soda to form sodium meta aluminate.  
 $2Al + 2NaOH + 2H_2O \rightarrow 2NaAlO_2 + 3H_2 \uparrow$   
Sodium meta aluminate
94. (a) Alkaline earth metals ( $ns^2$ ) are denser than alkali metal ( $ns^1$ ) because metallic bonding in alkaline earth metal is stronger.
95. (c) Lithium is basic in nature and hence it is not amphoteric.
96. (a)  $CsOH$  of the following is most basic in character due to increase electropositive character in a group of alkali.
97. (a) Group I element are so highly electropositive that they emit electrons even when exposed to light (Photoelectric effect) and this character increase on moving down the group from lithium towards cesium.
98. (b) Lithium form nitride on heating with nitrogen. Lithium nitride gives ammonia when


heated with  $H_2O$ . Ammonia gas form tetrammine copper complex with  $CuSO_4$  solution.



99. (d) The given compound x must be  $CaCO_3$ . It can be explained by following reactions,






100. (c) According to Fajan's rule  $RbCl$  has greatest ionic character due to large ionic size of  $Rb^+$  ion.  $BeCl_2$  has least ionic (Maximum covalent) due to small size of  $Be^{+2}$  ion which has highly polarising.












112. (c) It reacts with alcohol to form sodium alkoxide  
 $2C_2H_5OH + 2Na \rightarrow 2C_2H_5ONa + H_2$

113. (a) Causticizing process (Gossage process) it is an old process and involves heating of 10% solution of  $Na_2CO_3$  with a little excess of milk of lime  $Ca(OH)_2$



119. (b)  $NaOH$  is a deliquescent white crystalline solid. It absorbs moisture from the atmosphere.



139. (a) Down's cell is used for the electrolysis of fused  $NaCl$

142. (c)  $Fe(OH)_3$  does not dissolve in  $NaOH$

143. (a) Castner's process used to obtain  $Na$ , by electrolysis of sodium hydroxide.

144. (a) Excess of  $Na^+$  ion causes high B.P.

145. (b) Ferric alum is  $(NH_4)_2SO_4 \cdot Fe_2(SO_4)_3 \cdot 24H_2O$

146. (d) When  $Na$  is heated in presence of air or oxygen,  $Na$  burns to form sodium oxide and sodium peroxide.  
 $CaO + H_2O \rightarrow Ca(OH)_2$

148. (d) Pyrolusite or Manganese dioxide ( $MnO_2$ ) is a mineral of manganese.


149. (c)  $CaCl_2$  bring down the melt temperature from 1075 K to 850 K

### Alkaline earth metals

2. (d)  $CaSO_4 \cdot \frac{1}{2}H_2O$  or  $(CaSO_4)_2 \cdot H_2O$

3. (a)  $CaCl_2$  because it is hygroscopic

7. (d) Setting of plaster of paris is exothermic process




The setting is due to formation of another hydrate

10. (a)  $MgCO_3 \xrightarrow{\text{Heat}} MgO + CO_2$

The metal whose oxide is stable, its carbonate is unstable

12. (d)  $MgCl_2 \xrightarrow{\text{Electrolysis}} Mg^{+2} + 2Cl^-$   
(Molten) Cation Anion



13. (a) Because of small atomic size and high I.E. Be forms covalent chloride.

16. (d)  $BeSO_4$  is most soluble because hydration energy is more than lattice energy.



Hydration energy decreases hence, solubility decreases.

19. (b)  $2(CaSO_4 \cdot 2H_2O) \xrightarrow[\text{Plaster of paris}]{120^\circ C} 2CaSO_4 \cdot H_2O + 3H_2O$   
Gypsum Dehydration

21. (b) Lithopone ( $ZnS + BaSO_4$ ) is used as a white pigment.

23. (d)  $Ca(OH)_2 + Cl_2 \rightarrow CaOCl_2 + H_2O$   
Slaked lime Bleaching powder

24. (a) Strontium  $\rightarrow$  Crimson or pink colour

## 812 s and p-Block Elements

26. (b,c)  
 $Ca_3P_2 + 6H_2O \rightarrow 3Ca(OH)_2 + 2PH_3$   
 $K_3P + 3H_2O \rightarrow 3KOH + PH_3$
27. (d)  $CaCl_2 \rightarrow Ca^{+2} + 2Cl^-$   
Cathode Anode  
 Cathode :  $Ca^{+2} + 2e^- \rightarrow Ca$   
 Anode :  $2Cl^- \rightarrow 2e^- + Cl_2$
28. (d) Element - 

Mg	Al	Si	P	
Atomic radii (Å)	1.60	1.43	1.32	1.28

  
 as we move across the period nuclear charge increases, hence, size decreases.
30. (b)  $MgCl_2 \cdot 6H_2O + 5MgO + xH_2O \rightarrow$   
 $MgCl_2 \cdot 5MgO \cdot xH_2O$   
 Magnesia cement or sorrel cement
31. (d)  $ZnS + BaSO_4$  is lithopone used as white pigment.
36. (d) Aqueous  $CaCl_2$  or hydrated  $CaCl_2$  can not act as dehydrating agent.
38. (d) As we go down the group electropositive character increases because I.E. decreases.  
*Ba* is most electropositive element in the group.
39. (d) Due to the inert pair effect.
40. (a) Element - 

Be	Al	
Electronegativity -	1.5	1.5
41. (c)  $Be > Mg > Ca > Sr > Ba$   
 On moving down the group lattice energy remains almost constant as the sulphate is so big that small increase in the size of the cations from *Be* to *Ba* does not make any difference. However the hydration energy decreases from  $Be^{+2}$  to  $Ba^{+2}$ . This causes decrease in the solubility of the sulphates as the ionic size increases.
42. (a) Element - 

Be	Mg	Ca	Sr	
Electrode potential -	1.70	-2.37	-2.87	-2.89

  
 -2.90
43. (a) Element - 

Mg	Ca	Sr	Ba	
I.E -	737	590	549	503
44. (a) *Be* due to diagonal relationship
45. (a)  $K^+$  is highly soluble because of high hydration energy.
47. (b) 


$\underbrace{MgO}_{\text{Basic}}$	$\underbrace{Al_2O_3}_{\text{Amphoteric}}$	$\underbrace{SiO_2}_{\text{Acidic}}$	$\underbrace{P_2O_5}_{\text{Acidic}}$
-----------------------------------	--	--------------------------------------	---------------------------------------

  
 $MgO + H_2O \rightarrow Mg(OH)_2$  Base or alkali
48. (d)  
 Duralium (*Al* = 95%, *Cu* = 4%, *Mn* = 0.5%, *Mg* = 0.5%)  
 being light, tough and durable is used for the manufacture of aeroplanes and automobile parts.
49. (c) 

Na	K	Ba	Ca	Sr
Yellow	Pale violet	Apple green	Brick red	Crimson
51. (a) Magnesium burns with an intense light. Therefore *Mg* is used in flash bulbs for photography, fireworks and signal fires.
53. (b)  $CaO + CO_2 \rightarrow CaCO_3$   
 $CaO + H_2O \rightarrow Ca(OH)_2$
54. (a) When water is added to cement an exothermic reaction occurs. During this process, the cement reacts with water to form a gelatinous mass which slowly sets into a hard mass having three dimensional network structure involving  $-Si-O-Si-$  and  $-Si-O-Al-$  chains.
55. (b) *CaO* - (quick lime)  
 $Ca(OH)_2$  - (slaked lime)  
 $Ca(OH)_2 + H_2O$  - an aqueous suspension of  $Ca(OH)_2$  in water is called lime water.  
 $CaCO_3$  (lime stone).
57. (d) Lime stone -  $CaCO_3$   
 Clay - silica and alumina  
 Gypsum -  $CaSO_4 \cdot 2H_2O$
59. (b) Because hydration energy decreases down the group.
63. (d) *Be* does not react with water.
64. (a) (i) Small atomic size.  
 (ii) High electronegativity  
 (iii) Absence of *d* orbitals
65. (a)  $Ba(OH)_2 > Sr(OH)_2 > Ca(OH)_2 > Mg(OH)_2$   
 Solubility decreasing order.
66. (d) Solubility increasing top to bottom.
67. (a) *Be* to *Ba* ionic character increasing.
70. (a)  $Ca + 2H_2O \rightarrow Ca(OH)_2 + H_2$   
 $CaH_2 + 2H_2O \rightarrow Ca(OH)_2 + 2H_2$
72. (a) They are denser than alkali metals because they can be packed more tightly to their greater charge and smaller radii.
76. (d)  $Be(OH)_2 < Mg(OH)_2 < Ca(OH)_2 < Sr(OH)_2 < Ba(OH)_2$   
 On moving down the group basic character increases.
77. (b)  $Mg(OH)_2$  *Mg* is most electropositive element amongst the given elements.
78. (d) Lime stone =  $CaCO_3$   
 Quick lime = *CaO*  
 Slaked lime =  $Ca(OH)_2$




79. (c) As we go down the group I.E. decreases. Hence, Ba can easily give electrons. Therefore strongest reducing agent.






82. (a)  $BeCl_2 < MgCl_2 < CaCl_2 < BaCl_2$

As we go down the group I.E. decreases. Hence ionic character increases.



84. (a) The solubility of hydroxides of alkaline earth metals in water increases on moving down the group.

85. (c) Bleaching action of  $Cl_2$  in moist condition is permanent.



Coloured matter + nascent oxygen → colourless matter

86. (d) The solubility of hydroxides of alkaline earth metals in water increases on moving down the group.

87. (c)  $CO_2$  escapes simultaneously.

88. (d) It consist of high lattice energy and lowest hydration energy.

90. (a) It is a s-block elements.


91. (b) Barium  $Ba_{56}$  is a alkali earth metal.

92. (c) Thermal stability increasing from top to bottom.

94. (b) On moving down the group; Lattice energy decreases with increase in size of cation.



96. (d)  $BaSO_4$  is sparingly soluble in water because the solubility of second group sulphates decreases with increasing atomic size. Because of hydration energy decreases.

97. (d) Beryllium because of small atomic size and high ionization energy.






as we go down the group ionic nature increases because I.E. decreases.


100. (b)  $CaSO_4 \cdot 2H_2O$  - Gypsum



102. (c) Calcium cyanamide is the slow acting nitrogenous fertilizer as it decompose very slowly.





103. (a) Plaster of paris  $[(CaSO_4)_2 \cdot H_2O]$  is used in surgery for setting of bones, dentistry and manufacturing of statues. It is prepared as follows,




104. (b) Due to electropositive and reactive in nature, magnesium is readily converted into positive ions on contact with iron pipes and hence, iron pipes remains as it is.

105. (c) A binary compound is one made of two different elements. These can be one of each element such as  $CuCl$  or  $FeO$ . These can also be several of each element such as  $Fe_2O_3$  or  $SnBr_4$ . Metal which have variable oxidation number can form more than one type of binary compound like  $Fe$  shows the oxidation state +2 and +3. Hence it forms two type of binary compound e.g.,  $FeCl_2, FeCl_3$ .

106. (a) Diagonal relationship: elements of 2<sup>nd</sup> period often show resemblance to the element of the III<sup>rd</sup> period diagonally placed to it. This type of behaviour is called as diagonal relationship  $Li$  shows the diagonal relationship with  $Mg$ .




108. (c) We know that



Thus in this reaction magnesium dichloride is produced.


109. (d) Magnesium burns in  $CO$  to produce



110. (d) Sorel's cement is,  $MgCl_2 \cdot 5MgO \cdot xH_2O$


111. (b) Colemnite is a mineral of boron having composition as  $Ca_2B_6O_{11} \cdot 5H_2O$ .

### Boron family



$BCl_3$  is obtained by passing chlorine over the heated mixture of  $B_2O_3$  and powdered charcoal.

6. (d)  $B_2H_6$  has two types of B-H bonds



119 pm  
+++++


134 pm  
+++++

## 814 s and p-Block Elements


B H (Terminal bond)

B H (Bridge bond)

12. (b) Dilthey in 1921 proposed a bridge structure for diborane. Four hydrogen atoms, two on the left and two on the right, known as terminal hydrogens and two boron atoms lie in the same plane. Two hydrogen atoms forming bridges, one above and other below, lie in a plane perpendicular to the rest of molecule.





15. (c)  $2H_3BO_3 \rightarrow B_2O_3 + 3H_2O$ .
16. (a,c,d)  $Al_2Cl_6$ ,  $In_2Cl_6$ ,  $Ga_2Cl_6$
17. (a) Liquified Ga expand on solidification Ga is less electropositive in nature, It has the weak metallic bond so it expand on solidification.
18. (d)  $Al_2Cl_6 + 12H_2O \rightleftharpoons 2[Al(H_2O)_6]^{3+} + 6Cl^-$
19. (e)  $B_4C$  is the hardest substance along with diamond.
20. (a) Borazine  $B_3N_3H_6$ , is isoelectronic to benzene and hence, is called inorganic benzene some physical properties of benzene and borazine are also similar.



21. (c) Except  $B(OH)_3$  all other hydroxide are of metallic hydroxide having the basic nature  $B(OH)_3$  are the hydroxide of nonmetal showing the acidic nature.
22. (c) Moissan boron is amorphous boron, obtained by reduction of  $B_2O_3$  with Na or Mg. It has 95-98% boron and is black in colour.
23. (d) Boron form different hydride of general formula  $B_nH_{n+4}$  and  $B_nH_{n+6}$  but  $BH_3$  is unknown.

24. (c) Alumina is amphoteric oxide, which reacts acid as well as base.
25. (a) Al is the most abundant metal in the earth crust.
29. (a)  $AlCl_3 \cdot 6H_2O \xrightarrow{\Delta} Al(OH)_3 + 3HCl + 3H_2O$   
Thus  $AlCl_3$  can not be obtained by this method
30. (d) Amphoteric substance can react with both acid and base.
33. (c)  $2Al + 6HCl \rightarrow 2AlCl_3 + 3H_2$
34. (c) Al  $\rightarrow$  III group  $\rightarrow$  Forms  $Al_2O_3$
35. (d)  $2KOH + 2Al + 2H_2O \rightarrow 2KAlO_2 + 3H_2$
37. (c)  $Na_2CO_3 + H_2O \rightarrow 2NaOH + CO_2$   
 $2NaOH + 2Al + 6H_2O \rightarrow 2Na[Al(OH)_4] + 3H_2$
41. (c)  $B(OH)_3 \Rightarrow H_3BO_3$  Boric acid  
 $Al(OH)_3 \Rightarrow$  Amphoteric
45. (b)  $FeSO_4 \cdot (NH_4)_2SO_4 \cdot 6H_2O$
46. (b)  $Al_2O_3$  is an amphoteric oxide.
47. (c) Aluminium oxide is highly stable therefore, it is not Reduced by chemical reactions.
48. (d) Aluminium is used as reducing agent in metallurgy.
49. (a) Al is used as reducing agent in thermite process.
50. (c) In Goldschmidt aluminothermic process, thermite contains 3 parts of  $Fe_2O_3$  and 1 part of Al.
51. (c) For the purification of red bauxite which contains iron oxide as impurity  $\rightarrow$  Baeyer's process. For the purification of white bauxite which contains silica as the main impurity Serpeck's process.
52. (b) In Hall's process  
 $Al_2O_3 \cdot 2H_2O + Na_2CO_3 \rightarrow 2NaAlO_2 + CO_2 + 2H_2O$   
 $2NaAlO_2 + 3H_2O + CO_2 \xrightarrow{333 K}$   
 $2Al(OH)_3 \downarrow + Na_2CO_3$   
 $2Al(OH)_3 \xrightarrow{1473 K} Al_2O_3 + 3H_2O$
54. (d) Cryolite  $Na_3AlF_6$   
(1) Decreases the melting point of alumina  
(2) Increases conductivity of the solution
55. (b) Cryolite  $Na_3AlF_6$  is added  
(1) To decrease the melting temp from 2323 K to 1140 K  
(2) To increase the electrical conductivity of solution
61. (d) Iron oxide impurity - Baeyer's process  
Silica impurity - Serpeck's process
64. (b) Cryolite is added to lower the melting point of alumina and to increase the electrical conductivity.
65. (c) The purification of alumina can be done by Baeyer's process.

67. (c) In electrolytic method of obtaining aluminium from purified bauxite, cryolite is added to charge because it reduces the melting point of Bauxite (from  $1200^{\circ}\text{C}$  to  $800^{\circ} - 900^{\circ}\text{C}$ ) and also it increases electrical conductivity of mixture.
68. (a) Hoop's process  $\Rightarrow$  Purification of Al  
Hall and Heroult process  $\Rightarrow$  Reduction of  $\text{Al}_2\text{O}_3$   
Baeyer's and Serpeck's process  $\Rightarrow$  Concentration of Bauxite ore
74. (a)   
 $3c - 2e : \text{B} - \text{H} - \text{B}; \quad 2c - 2e : \text{H} - \text{B} - \text{H}$
75. (a)  $\text{B}_2\text{H}_6$   
  
Empty  $sp^3$  orbital of B
76. (b) Pure alumina is a bad conductor of electricity and the fusion temperature of pure alumina is about  $2000^{\circ}\text{C}$  and at this temperature when the electrolysis is carried of fused mass the metal formed vapoureses as the boiling point of Al is  $1800^{\circ}\text{C}$ .  
To overcome this difficulty,  $\text{Na}_3\text{AlF}_6$  and  $\text{CaF}_2$  are mixed with alumina.
77. (a) Concentration of Lewis acid of boron trihalides is increased in following order.  
 $\text{BF}_3 < \text{BCl}_3 < \text{BBr}_3 < \text{BI}_3$ .

### Carbon family

3. (d) It react with alkali as well as acid.
6. (a) Among alkali metal carbonates only  $\text{Li}_2\text{CO}_3$  decomposes.  $\text{Li}_2\text{CO}_3 \xrightarrow{\Delta} \text{Li}_2\text{O} + \text{CO}_2 \uparrow$
7. (b) Propyne can be prepared by the hydrolysis of magnesium carbide.  
 $\text{Mg}_2\text{C}_3 + 4\text{H}_2\text{O} \rightarrow \text{CH}_3\text{C} \equiv \text{CH} + 2\text{Mg}(\text{OH})_2$
10. (d) Generally red lead decompose into  $\text{PbO}$  and  $\text{O}_2$ .
11. (c)  $\text{CO}_2$  is acidic oxide and thus more effectively absorbed by an alkali.
12. (b)  $\text{CaC}_2$  have one sigma and two  $\pi$  bond.
13. (d) C and Si are non-metal and Pb is a metal.
16. (a)  $\text{SiO}_2 + 2\text{Mg} \rightarrow \text{Si} + 2\text{MgO}$ .
17. (b) Generally IV group element shows catenation tendency and carbon has more catenation power.
18. (b) Metal oxides or some salts are fused with glass to imported colour of glass.
19. (d)  $\text{Al}_2(\text{CO}_3)_3$  is less soluble in water than  $\text{Na}_2\text{CO}_3$ ,  $\text{ZnCO}_3$ .
20. (d) The inert pair effect is most prominent in Pb because from top to bottom due to increase in number of shells.
25. (c)  $\text{Co} + \text{NaOH} \xrightarrow{200^{\circ}\text{C}} \text{HCOONa}$   
Sod. formate
27. (c) Sodium oxalate react with conc.  $\text{H}_2\text{SO}_4$  to form CO and  $\text{CO}_2$  gas.
33. (d) It is hydrolysed with water to form a  $\text{Si}(\text{OH})_4$ .
35. (b) When hydrogen peroxide react with PbS then they form  $\text{PbSO}_4$ .
36. (b) Grey tin is very brittle and easily crumbles down to a powder in very cold climates.  
Grey tin  $\rightleftharpoons$  White tin  
(Cubic) (Tetragonal)
- The change of white tin to grey tin is accompanied by increase in volume. This is called tin disease or tin plague.
37. (c) Solid  $\text{CO}_2$  is knows as dry ice because it evaporates at  $-78^{\circ}\text{C}$  without changing in the liquid state.
38. (b) Zeolite have  $\text{SiO}_4$  and  $\text{AlO}_4$  tetrahedrons linked together in a three dimensional open structure in which four or six membered ring predominate. Due to open chain structure they have cavities and can take up water and other small molecules.
39. (b) Crook's glass is a special type of glass containing cerium oxide. It does not allow the passage of ultra violet ray and is used for making lenses.
40. (b) Inert pair effect become significant for the 6<sup>th</sup> and 7<sup>th</sup> period of p-block element.
41. (a) Carbon suboxide has linear structure with C-C bond length equal to  $130 \text{ \AA}$  and C-O bond length equal to  $120 \text{ \AA}$ .  
 $\text{O} = \text{C} = \text{C} = \text{C} = \text{O} \leftrightarrow \text{O}^- - \text{C} \equiv \text{C} - \text{C} \equiv \text{O}^+$
42. (c)  $\text{Pb}_3\text{O}_4$  is a mixed oxide. It can be represented as  $2\text{PbO} - \text{PbO}_2$ .
43. (b) Noble gases are found in very minute amount in atmosphere. These are separated from each other by using coconut charcoal. Which adsorb different gas at different temperature.
44. (c) Lapis Lazuli is a rock composed mainly of the following mineral, lazurite, hauynite sodalite, nosean, calcite, pyrite, lapis lazuli is actually sulphur containing, sodium aluminium silicate having chemical composition  $3\text{Na}_2\text{O} \cdot 0.3\text{Al}_2 \cdot 6\text{SiO}_2 \cdot 2\text{Na}_2\text{S}$ .



## 816 s and p-Block Elements

45. (d) In carbon family stability +2 oxidation state increases on moving down the group in the periodic table with an increase in atomic number due to screening effect.
46. (c) Tin is oxidised to meta stannic acid when it is treated with nitric acid.  

$$\text{Sn} + 4\text{HNO}_3 \rightarrow \text{H}_2\text{SnO}_3 + 4\text{NO}_2 + \text{H}_2\text{O}$$
47. (c)  $\text{Pb} + \text{Sn}$
49. (d) Three dimensional sheet structures are formed when three oxygen atoms of each  $[\text{SiO}_4]^{4-}$  tetrahedral are shared.
50. (a)  $\text{Pb}_3\text{O}_4 \Rightarrow$  Red lead (Sindhur)
51. (c) White lead  $\Rightarrow 2\text{PbCO}_3 \cdot \text{Pb}(\text{OH})_2$
52. (c) Organic acids dissolve lead in presence of oxygen  

$$\text{Pb} + 2\text{CH}_3\text{COOH} + \frac{1}{2}\text{O}_2 \rightarrow \text{Pb}(\text{CH}_3\text{COO})_2 + \text{H}_2\text{O}$$
53. (a) 
$$\begin{array}{c} | \quad | \\ \text{O} \quad \text{O} \\ | \quad | \\ -\text{O}-\text{Si}-\text{O}-\text{Si}-\text{O}- \\ | \quad | \\ \text{O} \quad \text{O} \\ | \quad | \end{array}$$
55. (a)  $s^2p^2$  Total 4 valence electrons  $\Rightarrow$  IV group
56. (c)  $\text{PbCl}_2$  is most ionic because on going down the group the metallic character increases and also the inert pair effect predominates.
58. (b) Type metal  $\text{Pb} = 82\%$ ,  $\text{Sb} = 15\%$ ,  $\text{Sn} = 3\%$
60. (b) Sugar of lead  $(\text{CH}_3\text{COO})_2\text{Pb} \Rightarrow$  lead acetate
63. (d)  $\text{Pb} \Rightarrow 11.34 \text{ g/ml}$  Heaviest
64. (c)  $\text{Pb}_3\text{O}_4$  is a mixed oxide of  $2\text{PbO} + \text{PbO}_2$
67. (c) Boron (B), Si, Ge, As, Sb, and At are the metalloid elements. Bismuth (Bi) and tin (Sn) are metals while carbon (C) is non-metal.
68. (a)  $\text{Al}_4\text{C}_3 + 12\text{H}_2\text{O} \rightarrow 3\text{CH}_4 + 4\text{Al}(\text{OH})_3$
69. (b) Glass being a mixture of sodium and calcium silicates reacts with hydrofluoric acid forming sodium and calcium fluorosilicates respectively.  

$$\text{Na}_2\text{SiO}_3 + 3\text{H}_2\text{F}_2 \rightarrow \text{Na}_2\text{SiF}_4 + 3\text{H}_2\text{O}$$


$$\text{CaSiO}_3 + 3\text{H}_2\text{F}_2 \rightarrow \text{CaSiF}_4 + 3\text{H}_2\text{O}$$
 The etching of glass is based on these reactions.

### Nitrogen family

3. (b)  $\text{FeSO}_4 + \text{NO} \rightarrow \text{FeSO}_4 \cdot \text{NO}$   
 (Brown)
4. (b)  $\text{HPO}_3$ , metaphosphoric acid  


$$\begin{array}{c} \text{O} \\ || \\ \text{P} \\ / \quad \backslash \\ \text{O} \quad \text{OH} \end{array}$$
6. (a) White phosphorus is soluble in  $\text{CS}_2$  whereas red phosphorus is insoluble in it.

7. (d)  $\text{H}_4\text{P}_2\text{O}_7$  pyrophosphoric acid



Tetrabasic (4-OH groups)

8. (b)  $\text{P}_4 + 3\text{NaOH} + 3\text{H}_2\text{O} \rightarrow \text{PH}_3 + 3\text{NaH}_2\text{PO}_2$   
 (White) Phosphine Sod. hypophosphite
9. (a)  $\text{NCl}_5$  is not known because of absence of d-orbitals in nitrogen.
11. (a,d)  $\text{P}_4$  molecule



Bond angle =  $60^\circ$

Six P-P = Single bonds

Lone pairs = 4

12. (b)  $\text{NH}_4\text{NO}_3 \xrightarrow{\Delta} 2\text{H}_2\text{O} \uparrow + \text{N}_2\text{O} \uparrow$   
 (s) Nitrous oxide  
 (Laughing gas)

13. (a) Birkeland - Eyde process  
 Dinitrogen is prepared commercially from air by liquification and fractional distillation. When liquid air is allowed to distil, dinitrogen having lower b.pt (77K) distils over first leaving behind liquid oxygen (bpt 90K). World wide production of dinitrogen from liquid air is more than 50 million tons per year.

14. (b)  $\text{NH}_4\text{NO}_3 \xrightarrow{\Delta} 2\text{H}_2\text{O} \uparrow + \text{N}_2\text{O} \uparrow$   
 (s)
- $$\text{NaNO}_3 \xrightarrow{\Delta} \text{NaNO}_2 + \text{O}_2 \uparrow$$
- (s) (s)
- $$2\text{AgNO}_3(\text{s}) \rightarrow 2\text{Ag}(\text{s}) + 2\text{NO}_2(\text{g}) + \text{O}_2(\text{g})$$
- Lunar caustic
- $$2\text{Pb}(\text{NO}_3)_2 \rightarrow 2\text{PbO} + 4\text{NO}_2 \uparrow + \text{O}_2 \uparrow$$
- (s)

16. (b)  $\text{P}_4 + 3\text{NaOH} + 3\text{H}_2\text{O} \rightarrow \text{PH}_3 + 3\text{NaH}_2\text{PO}_2$   
 White Phosphine


17. (d)  $\begin{array}{ccc} \text{N} & \text{P} & \text{As} & \text{Sb} & \text{Bi} \\ \underbrace{\hspace{1.5cm}} & \underbrace{\hspace{1.5cm}} & \underbrace{\hspace{1.5cm}} & \underbrace{\hspace{1.5cm}} & \underbrace{\hspace{1.5cm}} \\ \text{Non-metals} & \text{Metalloids} & \text{Meta} & & \end{array}$

18. (b) 
$$\begin{array}{c} \text{O} \\ || \\ \text{P} \\ / \quad \backslash \\ \text{HO} \quad \text{OH} \quad \text{OH} \end{array}$$













3 - OH groups are present hence it is tribasic.

19. (c) Nitrous acid behaves as reducing as well as an oxidising agent. It reduces potassium permanganate, potassium dichromate,  $\text{H}_2\text{O}_2$  and other strong oxidising agents. It oxidises strong reducing agents such as hydroiodic acid, sulphurous acid etc.


It oxidises  $\text{Fe}^{+2}$  into  $\text{Fe}^{+3}$  in acidic medium;




It reduces acidified  $\text{KMnO}_4$ .




32. (c) Because of its very low ignition temperature (303K) it is always kept under water.




34. (a,b) Stability of + 3 oxidation states increases on account of inert pair effect.  
Reducing character of hydrides increases down the group because bond dissociation energy decreases down the group.






36. (a)  $P_4 + 5O_2 \rightarrow P_4O_{10}$ ; white phosphorus gets easily oxidized because it is highly reactive.




39. (b) When  $N_2O$  is inhaled in moderate quantities, it produces hysterical laughter, hence the name laughing gas.







When dissolved in water, gives a mixture of nitrous acid and nitric acid.









42. (a)  $(HPO_3)_n$  Polymetaphosphoric acid



43. (b) Superphosphate of lime - It is a mixture of calcium dihydrogen phosphate and gypsum and is obtained by treating phosphatic rock will conc.  $H_2SO_4$



$NO$  is used in the preparation of  $HNO_3$




It is least basic because of the high electronegativity of 3F atoms. The lone pair present on nitrogen atom is not easily available for donation.





53. (c) Due to less reactivity of red phosphorus

54. (d)  $NO_2$  brown coloured gas.



Therefore, it can act only as oxidising agent.




## 818 s and p-Block Elements


58. (c)  $NH_2OH$   
 $x + 2 + (-2) + 1 = 0$   
 $x + 2 - 2 + 1 = 0$   
 $x = -1$
60. (c)  $NH_3 > PH_3 > AsH_3 > SbH_3$   
 On moving down the group atomic size increases and availability of lone pair decreases. Hence, basic character decreases.
61. (a)  $PH_3 > AsH_3 > SbH_3 > BiH_3$   
 On moving down the group bond energy decreases. Hence, stability decreases.
62. (d) Due to absence of  $d$ -orbitals in  $N$  atom, it cannot accept electrons from  $H_2O$  for hydrolysis of  $NF_3$ .
63. (b)  $NH_3$  is most thermally stable hydride. Hence, electrolysis temperature is maximum.
64. (a) Phosphorus is kept in water due to it burt at  $30^\circ C$ .
66. (c)  $BiCl_3 + H_2O \rightarrow BiOCl + 2HCl$
67. (c) When the black ppt. of  $Bi_2S_3$  is dissolved in 50%  $HNO_3$  and a solution of  $NH_4OH$  is added. A white ppt. of  $Bi(OH)_3$  is obtained.
69. (a) Atmospheric nitrogen is inert and unreactive because of very high bond energy ( $945 \text{ kJ/mole}$ ).
70. (b) Bismuth does not show allotropy other elements show allotropy.  
 Nitrogen  $\rightarrow \alpha$ -nitrogen and  $\beta$ -nitrogen (solid crystalline forms)  
 Phosphorus  $\rightarrow$  White, Red and Black forms  
 Arsenic  $\rightarrow$  Yellow and Grey forms  
 Antimony  $\rightarrow$  Yellow and Grey forms
71. (a) Nitrogen does not form complexes because of the absence of  $d$ -orbitals.
72. (a)  $NH_3$  is a strongest base because Lone pair is easily available for donation.
74. (b) Hydride  $NH_3$   $PH_3$   $AsH_3$   $SbH_3$   $BiH_3$   
 Boling point 238.5 185.5 210.6 254.6  
 290
75. (a)  $NCl_3$  is highly reactive and unstable. Hence it is explosive.
76. (b)  $N_2O_3$   $P_2O_3$   $As_2O_3$   $Sb_2O_3$   $Bi_2O_3$   
 Acidic Oxides Amphoteri Basic  
 Acidic character decreases down the group  $\rightarrow$
77. (c)  $SbCl_2$  is not exists because  
 $V^{th}$  group elements normally show +3 and +5 oxidation state.
78. (b)  $NH_4Cl + NaNO_2 \rightarrow NH_4NO_2 + NaCl$   
 $(aq) \quad (aq)$   
 $NH_4NO_2 \xrightarrow{\text{heat}} N_2 + 2H_2O$   
 $(g) \quad (l)$
79. (c)  $NH_4NO_2 \rightarrow N_2 + 2H_2O$
80. (d)  $6Li + N_2 \rightarrow 2Li_3N$  Lithium nitride  
 $3Mg + N_2 \rightarrow Mg_3N_2$  Magnesium nitride
81. (d)  $N \equiv N$  bond energy is very high  $945 \text{ kJ mol}^{-1}$ .
83. (d)  $N_7 \rightarrow 1s^2, 2s^2, 2p^3$   
 $d$ -orbitals are absent in nitrogen.
85. (d)  $NH_4NO_3 \xrightarrow{\text{heat}} N_2O + 2H_2O$   
 (Laughing gas)
86. (d)  $NH_2OH + HNO_2 \rightarrow H_2N_2O_2 + H_2O$
87. (c)  $N_2O$  is a linear molecule
88. (b)  $2HNO_2 \rightarrow H_2O + N_2O_3$
89. (d)  $2HNO_3 \rightarrow H_2O + N_2O_5$
90. (c)  $2Pb(NO_3)_2 \rightarrow 2PbO + 4NO_2 + O_2$
91. (a)  $2Pb(NO_3)_2 \rightarrow 2PbO + 4NO_2 + O_2$
92. (b)  $3Cu + 8HNO_3 \rightarrow 3Cu(NO_3)_2 + 4H_2O + 2NO$
93. (d) In upper atmosphere  $NO$  is formed by lightning flash.  
 $N_2 + O_2 = 2NO$
96. (c)  $2NO + O_2 \rightarrow 2NO_2$
98. (b)  $2AgNO_3 \rightarrow 2AgNO_2 + O_2$   
 $\downarrow$   
 $2Ag + 2NO_2$
100. (d)  $2NO_2 + H_2O \rightarrow HNO_3 + HNO_2$
101. (d)  $C_{12}H_{22}O_{11} \xrightarrow{\text{conc. } HNO_3} \begin{matrix} COOH \\ | \\ COOH \\ \text{Oxalic acid} \end{matrix} + H_2O$
102. (b)  $4NH_3 + 5O_2 \xrightarrow[800^\circ C]{Pt} 4NO + 6H_2O$
103. (d)  $HNO_2$  can be either reduced to nitric oxide ( $NO$ ) or oxidised to nitric acid and hence it acts both as an oxidising as well as a reducing agent.  
 $2HNO_2 \rightarrow 2NO + H_2O + [O]$   
 $HNO_2 + [O] \rightarrow HNO_3$
106. (d)  $NH_3$  is highly volatile compound. When vapourized, liquid ammonia causes intense cooling. Hence used as a coolant in ice factories and cold storages.
107. (d)  $N_3H \rightleftharpoons N_3^- + H^+$   
 Hy drazoic acid
108. (a)  $d$ -orbitals are absent in nitrogen.
109. (d) Phosphide ion Chloride ion  
 $(P^{3-}) \quad (Cl^-)$   
 Total electrons 18  
 $P^{3-}$  and  $Cl^-$  are isoelectronic.
110. (a) Due to the less reactivity.
116. (a)  $P_4 + 3H_2O + 3NaOH \rightarrow PH_3 + 3NaH_2PO_2$   
 Phosphine Sod. hypophosphite
117. (c) Both oxidation and reduction  
 (Disproportionation)

Oxidation

B.pt in (K) 238.5 185.5 210.6 254.6




118. (b)  $P_4 + NaOH \rightarrow$  No reaction  
 120. (c)  $Ca_3P_2 + 6H_2O \rightarrow 3Ca(OH)_2 + 2PH_3$   
 122. (b)  $PH_3$  is less basic because lone pair is not easily available for donation.  
 123. (d)  $P_2O_3 + 3H_2O \rightarrow 2H_3PO_3$   
 124. (d)  $P_2O_5 + 3H_2O \rightarrow 2H_3PO_4$  orthophosphoric acid.  
 125. (c)  $H_3PO_2$  Monobasic acid



126. (b)  $PCl_3 + 3H_2O \rightarrow H_3PO_3 + 3HCl$   
 127. (b)  $H_3PO_3$   


$$\begin{array}{c} O \\ || \\ H - P - OH \\ | \\ OH \end{array}$$
  
 128. (c)  $H_3PO_2$   
 $3 + x + (-2 \times 2) = 0$   
 $x = +1$   
 129. (d)  $Na_4P_2O_7$  Salt of strong acid and strong base.  
 130. (b)  $P_4 + 6H_2SO_4 \rightarrow 4H_3PO_4 + 6SO_2$   
 131. (c)  $CaCN_2 + 3H_2O \rightarrow CaCO_3 + 2NH_3$   
 132. (b)  $H_4P_2O_7$



4  $-OH$  group are present.

134. (b)  $BiCl_3 + H_2O \rightarrow BiOCl + 2HCl$   
 135. (b)  $CaC_2 + N_2 \xrightarrow[6-8 \text{ atm}]{500-600^\circ C} CaCN_2 + C$   
 136. (a)  $CaCN_2 + 3H_2O \rightarrow CaCO_3 + 2NH_3$   
 137. (a)  $NH_2CONH_2$   


$$\% \text{ of } N = \frac{\text{Mass of } N}{\text{Mass of compound}} \times 100 = \frac{28}{60} \times 100 = 46\%$$
  
 141. (a) Anhydride of nitrous acid is  $N_2O_3$ .  
 144. (b)  $P_4 + 3NaOH + 3H_2O \rightarrow PH_3 + NaH_2PO_4$   
 145. (a)  $NH_3$  is highly soluble due to H-bonding.




146. (d)  $NH_3$   $PH_3$   $AsH_3$   $SbH_3$

149. (a)  $H_3PO_2$   
 $3 + x - 4 = 0$   
 $x = +1$


151. (d) Solid  $PCl_5$  exists as  $PCl_4^+$  and  $PCl_6^-$ .



153. (a)  
 154. (e) Phosphorus minerals is called as hydroxyapatite and fluorapatite.  
 156. (a) Nitrogen does not have  $d$ -orbitals.  
 157. (c)  $3CuO + 2NH_3 \rightarrow 3Cu + N_2 + 3H_2O$ .  
 159. (b) Liquid ammonia is used in refrigeration because it has high heat of vaporisation.  
 160. (a)  $Sn + \text{conc. } 4NHO_3 \rightarrow H_2SnO_3 + 4NO_2 + H_2O$   
Meta stannic acid  
 161. (c)  $3Cu + 8NHO_3 \rightarrow 3Cu(NO_3)_2 + 4H_2O + 2NO$   
Nitric oxide  
 163. (c) Pentavalency in phosphorus is more stable than that of nitrogen due to large size of phosphorus atom.  
 164. (b) Ammonium nitrate is neutral fertilizer.  
 165. (d)  $PH_3$  insoluble in water because does not consist of hydrogen bond.




167. (a) Nitrogen react with metal to form a nitride.  
 $6Li + N_2 \xrightarrow{450^\circ C} 2Li_3N$  (Lithium nitride)



169. (b) It is a salt of pyrophosphoric acid  $H_4P_2O_7$ .  
 172. (d) Copper react with conc. nitric acid to form a nitric oxide.  
 173. (a)  $N_2O$  on account of stimulating effect on nervous system.

## 820 s and p-Block Elements

174. (d) Sodium metal in liq.  $NH_3$  solution shows strong reducing power due to solvated electron.



175. (c)  $PH_3 + 4Cl_2 \rightarrow PCl_5 + 3HCl$


178. (c) Generally  $P_2O_5$  are used as a dehydrating agent.

180. (a) Phosphorus show + 5 valency.

181. (b) In the Haber process for the manufacture of  $NH_3$ ,  $Fe$  is used catalyst and  $Mo$  as a promotre.

182. (a) On adding excess of ammonium hydroxide to a copper chloride solution a deep blue solution of  $[Cu(NH_3)_4]^{2+}$  ion is formed.

183. (d)  $(NH_4)_2SO_4 + KCNO \rightarrow$




185. (a) Nitric acid turns the skin yellow because it reacts with protein giving a yellow compound called xanthoprotein.

186. (d) Ammonium sulphate is a nitrogenous fertilizers.

187. (d) Ammonia generally prepared by the Haber's process.

192. (a)  $H_3PO_2$  is hypophosphorus acid

193. (c)  $(NH_4)_2SO_4$  is a salt of weak base & strong acid





194. (a) One part of concentrated  $HNO_3$  and 3 parts concentrated  $HCl$  form aquaregia.

196. (c) -3 to +5  $PH_3(-3)$  and  $H_3PO_4(+5)$

199. (b)  $BiCl_5$  does not exist because +3 oxidation state of  $Bi$  is more stable than +5 due to inert pair effect.


200. (c)  $H_3PO_3 \rightarrow$  Tribasic acid  $\rightarrow$  3 - OH groups are present



201. (d)  $Na_2HPO_4 \rightarrow Na_2PO_4^- + H^+$

It can give  $H^+$  ion in solution.

202. (c)  $\ddot{N}H_3$  and  $\ddot{P}H_3$  both are basic because of the presence of lone pair of electrons.



203. (b)  $\xrightarrow{\hspace{10em}}$

Stability decreases down the group because bond energy decreases down the group.

204. (a) Nitrogen forms  $NH_3$  which is most basic.

205. (b)  $H_3PO_3$  is a dibasic acid. It forms two types of salts  $NaH_2PO_3$  and  $Na_2HPO_3$ .

206. (a)  $NH_2 - CO - NH_2 + 2HNO_2 \rightarrow CO_2 + 3H_2O + 2N_2$

	I	II	III	IV
Element -	P	As	Sb	Bi
N				
Atomic no.	15	33	51	83
7				

210. (c)  $HO - \overset{\overset{O}{||}}{P} - OH$  it is ionizes in three steps because three -OH group are present.

212. (c)  $Ca_3P_2 + 3H_2O \rightarrow 3Ca(OH)_2 + 2PH_3$

213. (d)  $(NH_4)_2Cr_2O_7 \rightarrow N_2 + Cr_2O_3 + 4H_2O$

214. (b)  $B > P > As > Bi$

As we go down the group bond angle decreases because repulsion between bonded pairs of electron decreases.

215. (b,c)  $3NH_3 + OCl^- \rightarrow NH_2 - NH_2 + NH_4Cl + OH^-$

217. (a) Acidic character of oxides decreases down the group.

218. (d)  $N_7 - 1s^2, 2s^2, 2p^3$

d-orbitals absent in second sub-shell.

220. (c)  $N_2$  can form  $NCl_3$ ,  $N_2O_5$  and  $Ca_3N_2$  but does not form  $NCl_5$ .


221. (a) Highest oxidation state is +5 which remains unchanged.

222. (a) Hypophosphorus acid ( $H_3PO_2$ ) is a monobasic acid which act as reducing agent. In this molecule two P-H bonds are responsible for its reducing character and one O-H bond is responsible for its monobasic acid character.

223. (a) Bone black is the polymorphic form of phosphorus.

224. (b) Nitrous oxide is known as Laughing gas.


225. (a) We know that,



So nitric oxide ( $NO_2$ ) is produced.

226. (d) Phosphorus exist as solid at  $27^\circ C$  and 1 atmospheric pressure (m.p. of white phosphorus =  $44^\circ C$ )

227. (b) We know that,  $4HNO_3 + P_4O_{10} \rightarrow 4HPO_3 + 2N_2O_5$   
The product is dinitrogen pentaoxide ( $N_2O_5$ ).





228. (b) Hypophosphorous acid is  $H_3PO_2$ .
229. (b)  $NO(g) + NO_2(g) \xrightarrow{-30^\circ C} N_2O_3(l)$   
(Blue)
230. (c) The ignition temperature of black phosphorus is highest among all allotropes.
231. (a)  $(NH_4)_2Cr_2O_7 \xrightarrow{\Delta} N_2 \uparrow + Cr_2O_3 + 4H_2O$   
 $NH_4NO_2 \xrightarrow{\Delta} N_2 \uparrow + 2H_2O$
232. (a) Nitrogen shows +I to +V, all oxidation states.
233. (c) Boiling points of  $SbH_3$  (254 K),  $NH_3$  (238 K),  $AsH_3$  (211 K) and  $PH_3$  (185 K) therefore boiling points are of the order  $SbH_3 > NH_3 > AsH_3 > PH_3$ .
234. (a) Because phosphorous is most electronegative element out of P, Bi, Sb and C.
235. (d)  $Ca_3(PO_4)_2 + 3SiO_2 \rightarrow 3CaSiO_3 + P_2O_5$
236. (a) When a solid compound on heating change into gaseous state without changing into liquid state, the phenomenon is known as sublimation. e.g.,  $I_2$ ,  $NH_4Cl$  and camphor.
237. (b) 16 bond by its structure.
238. (d) Phosphorus is a non-metallic element. It form's acidic oxide.
239. (b)  $2NO_2 + H_2O \rightarrow HNO_2 + HNO_3$   
Mixed acid anhydride
240. (c) Oxidation number of As in  $H_2AsO_4^-$   
 $2 + x - 8 = -1$   
 $x - 6 = -1$   
 $x = 5$
241. (a) The inorganic nitrogen exists in the form of ammonia, which may be lost as gas to the atmosphere, may be acted upon by nitrifying bacteria, or may be taken up directly by plants.


### Oxygen family

2. (c) Sulphur -  
(1) Monoclinic (2) Rhombic (3) Plastic
4. (c)  $S + O_2 \rightarrow SO_2$
5. (a)  $\frac{1}{8}S_8 + 2e^- \rightarrow S^{2-}$
6. (c)  $2H_2O + 2F_2 \rightarrow 4HF + O_2$
9. (b)  $O_8 -$   $1s^2$   $2s^2$   $2p^4$   
 $\uparrow\downarrow$   $\uparrow\downarrow$   $\uparrow\downarrow$   $\uparrow$   $\uparrow$   $\uparrow$   
2 unpaired electrons
11. (a) Element - O S Sc Te  
Po

Electronegativity - 3.5 2.5 2.4 2.1  
2.0

13. (b,c)  
 $Na_2SO_3 + H_2SO_4 \rightarrow Na_2SO_4 + H_2O + SO_2$   
(s) (aq) (aq) (l) (g)
15. (d)  $H_2O$   $H_2S$   $H_2Se$   $H_2Te$   
104.5° 92.1° 91° 90°  
As we go down the group electronegativity decreases due to which repulsion between bonded pairs of electron also decreases. Hence, bond angle decreases.
20. (a) Paramagnetism because of two unpaired electrons in the antibonding molecular orbitals.
21. (a)  $2Na_2SO_3 + O_2 \rightarrow 2Na_2SO_4$
24. (c)  $3O_2 \xrightleftharpoons[\text{electric discharge}]{\text{silent}} 2O_3$
28. (b)  $O_3 \rightarrow O_2 + [O]$   
 $2KI + H_2O + [O] \rightarrow 2KOH + I_2$   
 $2KI + H_2O + O_3 \rightarrow 2KOH + I_2 + O_2$
32. (d)  $2KMnO_4 + 3H_2SO_4 + 5H_2S \rightarrow$   
 $K_2SO_4 + 2MnSO_4 + 8H_2O + 5S$
33. (a)  $Cu + 2H_2SO_4 \rightarrow CuSO_4 + 2H_2O + SO_2$
35. (a) The minimum and maximum oxidation number of S are -2 and +6 respectively. Since the oxidation number of S in  $SO_2$  is +4, therefore it can be either increased or decreased. Therefore  $SO_2$  behaves both as an oxidising as well as reducing agent.
36. (a)  $2H_2S + SO_2 \rightarrow 2H_2O + 3S$   
reducing agent oxidising agent
37. (d)  $K_2Cr_2O_7 + H_2SO_4 + 3SO_2 \rightarrow$   
 $K_2SO_4 + Cr_2(SO_4)_3 + 3H_2O$   
Green
39. (a)  $SO_2 + 2Mg \rightarrow 2MgO + S$
40. (a)  $2H_2O + SO_2 \rightarrow H_2SO_4 + 2[H]$  (nascent hydrogen)  
Coloured flower +  $2[H] \rightarrow$  Colourless flower
41. (c)  $H_2SO_3 + 2NaOH \rightarrow Na_2SO_3 + 2H_2O$   
Sodium sulphite
42. (a)  $H_2O + SO_3 \rightarrow H_2SO_4$
43. (d)  $H_2SO_4 + SO_3 \rightarrow H_2S_2O_7$   
(oleum)
45. (b)  $2Ag + 2H_2SO_4 \rightarrow Ag_2SO_4 + 2H_2O + SO_2$   
Reducing agent Oxidising agent
46. (a) Only dehydrating agent  
 $HCOOH \xrightarrow{H_2SO_4} CO + H_2O$

## 822 s and p-Block Elements

48. (b)  $\begin{array}{c} \text{COOH} \\ | \\ \text{COOH} \end{array} \xrightarrow{\text{conc. } H_2SO_4} H_2O + CO + CO_2$
51. (d)  $H_2SO_4 + SO_3 \rightarrow H_2S_2O_7$   
(Oleum or Fuming  $H_2SO_4$ )
53. (d)  $S_2O_7^{2-}$
- $$\begin{array}{c} O \quad O \\ || \quad || \\ ^-O-S-O-S-O^- \\ || \quad || \\ O \quad O \end{array}$$
55. (d)  $Na_2SO_3 + S \rightarrow Na_2S_2O_3$
56. (b)  $Na_2S_2O_3 + Cl_2 + H_2O \rightarrow Na_2SO_4 + 2HCl + S$
58. (c)  $AgBr + 2Na_2S_2O_3 \rightarrow Na_3[Ag(S_2O_3)_2] + NaBr$
60. (a)  $SO_2 + 2H_2S \rightarrow 3S + 2H_2O$
63. (a)  $2H_2S + O_2 \rightarrow 2H_2O + 2S$
64. (a) Mixture of  $K_2Cr_2O_7$  and conc.  $H_2SO_4$  is known as chromic acid.
65. (c)  $H_2O$  contain hydrogen bond while no hydrogen bonding is present in  $H_2S$ .
67. (a)  $H_2O$  containing hydrogen bond.
69. (c)  $HO-SO_2-OH + PCl_5 \rightarrow Cl-SO_2-OH + POCl_3$   
+HCl  
 $HO-SO_2-OH + 2PCl_5 \rightarrow Cl-SO_2-Cl + 2POCl_3 + 2HCl$   
Sulphuryl chloride
70. (d) Mixture of  $O_2$  and  $CO_2$ .
71. (c) The bond between ( $H_2Te$ ) is weakest hence it gives  $H^+$  ion easily.
73. (c)  $KO_2$  because in  $O_2^-$  (superoxide ion) One unpaired electron is present in the antibonding orbital.
75. (b) Oxidation states are 2, 4, 6
77. (b)  $H_2O$   $H_2S$   $H_2Se$   $H_2Te$   
373K 213K 269K 232K  
 $H_2S$  has lowest boiling point and  $H_2O$  has highest boiling point because if any compound has hydrogen bond. Its boiling point is high.
78. (c)  $V_2O_5$  catalyst are used for the manufacture of  $H_2SO_4$  by Contact process.
81. (d) O, S, Se shows polymorphism.
84. (e)  $H_2S + 2HNO_3 \rightarrow 2NO_2 + S + 2H_2O$ .
85. (b)  $O_2F_2$  is similar to that of  $H_2O_2$
- 
87. (d) 16<sup>th</sup> group called chalcogens (oxygen family) while Na is a 1<sup>st</sup> group element which is called alkali metal.
90. (b) Caro's acid ( $H_2SO_5$ )
91. (c)  $2KO_2 + CO_2 \rightarrow K_2CO_3 + \frac{3}{2}O_2$
93. (a)  $H_2O$  consist of highest boiling point than other hydride (Due to presence of the hydrogen bonding).
94. (b) Lose of electron.
95. (c)  $SO_2$  is soluble in water  
 $H_2O + SO_2 \rightarrow H_2SO_3$   
Sulphurous acid
96. (c)  $2Na_2O_2 + 2H_2O \rightarrow 4NaOH + O_2$ .  
In this reaction oxygen ( $O_2$ ) is formed.
97. (b)  $3S + 4NaOH \xrightarrow{\text{boiling}} Na_2S_2O_3 + Na_2S$
98. (c) Quartz is a crystalline variety of silica.
99. (b) 98%  $H_2SO_4$  is used for absorbing dense fog of acid which is formed by dissolving  $SO_3$  in water. Hence 98%  $H_2SO_4$  is the most efficient agent for the absorption of  $SO_3$ .
100. (b) Concentrated  $H_2SO_4$  is diluted by adding the conc.  $H_2SO_4$  in the water drop by drop with constant stirring because it is an exothermic reaction and by doing so heat is generated slowly and dissipated in the atmosphere.
101. (a) Photochemical smog is caused by oxides of sulphur and nitrogen.
102. (d)  $SO_2$  act as bleaching agent due to its reducing property.  
 $SO_2 + 2H_2O \rightarrow H_2SO_4 + 2H$   
Coloured matter + H → Colourless matter
103. (b) Ozone on treatment with  $SO_2$  produce  $SO_3$ .  
 $3SO_2 + O_3 \rightarrow 3SO_3$
104. (b)  $PbO_2$  is a powerful oxidizing agent and liberate  $O_2$  when treated with acids  $SO, O_2$  gas will be evolved.
105. (c) Dithionous acid ( $H_2S_2O_4$ ) has sulphur in +3 oxidation state.  
 $HO-\overset{\overset{O}{||}}{S}-\overset{\overset{O}{||}}{S}-OH$  ;  $2(+1) + 2x + 4(-2) = 0$   
 $2x = 8 - 2 = 6$  ;  $x = +3$
106. (a)  $SO_2$  acts as an oxidising agent particularly when treated with stronger reducing agents.  $SO_2$  oxidises  $H_2S$  into S.
107. (a)  $R_3SiCl$  on hydrolysis forms only a dimer.  
 $R_3SiOH + HOSiR_3 \rightarrow R_3Si-O-SiR_3$ .

108. (d) Formation of chlorine nitrate is the main cause of ozone depletion.

### Halogen family

- (b)  $HF > HCl > HBr > HI$  (Thermal stability).
- (a)  $CHCl_3 + \frac{1}{2}O_2 \rightarrow COCl_2 + HCl$   
Phosgene or carbonyl chloride
- (d) Iodine has the least affinity for water and is only slightly soluble in it. However, it dissolves in 10% aq. solution of KI due to the formation of a complex ion i.e.  $I_3^-$ .  
 $I_2 + KI = KI_3$  or  $I_2 + I^- = I_3^-$  (complex ion)
- (c)  $2Na_2S_2O_3 + I_2 \rightarrow 2NaI + Na_2S_4O_6$
- (a)  $Cl_2 + 2KBr \rightarrow 2KCl + Br_2$   
A more electronegative halogen can displace less electronegative halogen.
- (a) HI is the strongest reducing agent among halogen acids because of lowest bond dissociation energy.
- (a) Due to H-Bonding free ions are not present in aq. solution. Hence, bad conductor.
- (c) Electronegativity of  $I_2$  is less than  $Br_2$ . Therefore unable to displace bromine.
- (b) Carnallite is  $KCl \cdot MgCl_2 \cdot 6H_2O$ . The mother liquor left after crystallisation of KCl from carnallite contains about 0.25% of bromine as  $MgBr_2$  and  $KBr$ .
- (a) HF is liquid because of intermolecular H-Bonding.
- (a)  $HClO \rightleftharpoons H^+ + ClO^-$   
Weak acid                      Strong conjugate base
- (d)  $2NaOH + Cl_2 \xrightarrow{\text{Cold}} NaCl + NaClO + H_2O$   
(dil.)                                      Sod. hypochlorite  
 $6NaOH + 3Cl_2 \xrightarrow{\text{heat}} 5NaCl + NaClO_3 + 3H_2O$   
(conc.)                                      Sodium chlorate
- (b)  $6KOH + 3Cl_2 \rightarrow 5KCl + KClO_3 + 3H_2O$ .
- (a) HF is the weakest acid. Since it is unable to give  $H^+$  ions which are trapped in H-Bonding.
- (b) Hydride -  $HF$      $HCl$      $HBr$      $HI$   
B.pt (in K) - 293    189    206    238  
Because of having low b.p. HCl is more volatile.
- (a)  $2KClO_3 + I_2 \rightarrow 2KIO_3 + Cl_2$
- (c)  $2Na_2S_2O_3 + I_2 \rightarrow 2NaI + Na_2S_4O_6$
- (d)  $2KMnO_4 + 16HCl \rightarrow 2KCl + 2MnCl_2 + 5Cl_2 + 8H_2O$
- (a)        +7        +5        +3        +1  
 $HClO_4 > HClO_3 > HClO_2 > HClO$

As the oxidation no. of halogen increases acidic character increases.

- (c)  $2KBr + 3H_2SO_4 + MnO_2 \xrightarrow{\Delta} 2KHSO_4 + MnSO_4 + 2H_2O + Br_2$
- (b,d) Electron affinity of  $Cl_2$  is maximum  

Element -	F	Cl	Br	I
E.A. kJ/mole	332.6	348.5	324.7	295.5
Boiling pt (°C)	-	-34.6	59.5	185.2
	188.1			
- (a)  $Cl_2 + 2NaBr \rightarrow 2NaCl + Br_2$
- (d)  $CCl_4 + H_2O \rightarrow$  No reaction  
 $d$ -orbitals are absent in carbon atom.
- (a)  $I_2 + 10HNO_3 \rightarrow 2HIO_3 + 10NO_2 + 4H_2O$
- (d)  $KI + I_2 \rightarrow KI_3$
- (a)  $2KBr + H_2SO_4 \rightarrow K_2SO_4 + 2HBr$
- (b)  $H_2 + F_2 \rightarrow 2HF$
- (b)  $CuSO_4 + 2KI \rightarrow CuI_2 + K_2SO_4$   
 $2CuI_2 \rightarrow 2CuI + I_2$   
Cuprous iodide
- (d) As the atomic number increases electronegativity decreases. Hence, tendency to gain electron decreases.
- (a)  $F_2 + 2Cl^- \rightarrow Cl_2 + 2F^-$   
 $F_2 + 2Br^- \rightarrow Br_2 + 2F^-$   
 $F_2 + 2I^- \rightarrow I_2 + 2F^-$
- (d)  $Br_2 + 2KI \rightarrow I_2 + 2KBr$
- (d)  $2F_2 + 2H_2O \rightarrow 4HF + O_2$   
 $3F_2 + 3H_2O \rightarrow 6HF + O_3$
- (c)  $2NaCl + 2H_2O \rightarrow 2NaOH + Cl_2 + H_2$   
(anode) (cathode)
- (a)  $MnO_2 + 4HCl \rightarrow MnCl_2 + 2H_2O + Cl_2$   
Reduction  
+4                      +2
- (b)  $2NaCl + 2H_2O \xrightarrow{\text{Electrolysis}} 2NaOH + Cl_2 + H_2$   
(aq)                      (g)                      (g)
- (c)  $H_2O + Cl_2 \rightarrow HCl + HClO$   
Exposed to air  
 $HClO \rightarrow HCl + [O]$  or  $2HClO \rightarrow 2HCl + O_2$
- (a)  $2NaOH + Cl_2 \rightarrow NaClO + NaCl + H_2O$
- (b)  $Cl_2 + H_2O \rightarrow 2HCl + [O]$  Nascent oxygen  
Coloured flower    + [O]    Bleaching agent    Colourless flower (Oxidized)
- (b)  $CaO + Cl_2 \rightarrow CaOCl_2$   
 $NaHCO_3 + Cl_2 \rightarrow$  No reaction
- (c)  $Ca(OH)_2 + Cl_2 \rightarrow CaOCl_2 + H_2O$   
Slaked lime
- (b)  $MgBr_2 + Cl_2 \rightarrow MgCl_2 + Br_2$


## 824 s and p-Block Elements

72. (a)  $Cl_2 + 2Br^- \rightarrow 2Cl^- + Br_2$
73. (a)  $KI + H_2SO_4 \xrightarrow[\text{Conc.}]{\Delta} KHSO_4 + HI$   
 $\text{Conc. } H_2SO_4 + 2HI \rightarrow 2H_2O + I_2 + SO_2 \uparrow$   
Violet vapour
74. (b)  $2NaI + Cl_2 \rightarrow 2KCl + I_2$   
 $CCl_4 + I_2 \rightarrow \text{Violet colour}$
76. (b)  $KI + H_2SO_4 \xrightarrow{\Delta} KHSO_4 + HI$   
 $H_2SO_4 + 2HI \rightarrow 2H_2O + I_2 + SO_2 \uparrow$   
Violet vapour
79. (c)  $KI + H_2SO_4 \rightarrow KHSO_4 + HI$   
 $HI$  is formed but it is further oxidised by conc.  $H_2SO_4$  into  $I_2$   
 $2HI + H_2SO_4 \rightarrow 2H_2O + I_2 + SO_2 \uparrow$   
Violet vapour
80. (b)  $HCl \xrightarrow{H_2O} H^+ + Cl^-$   
(aq)            (aq)
81. (b)  $4NaCl + K_2Cr_2O_7 + 3H_2SO_4 \rightarrow$   
 $K_2SO_4 + 2Na_2SO_4 + 2CrO_2Cl_2 + 3H_2O$   
Chromylchloride
82. (c) Hydrogen bonding is absent in  $HI$  while it is present in  $NH_3$ ,  $H_2O$  and  $C_2H_5OH$ .
84. (d) In case of  $HI$  due to large size of iodine strong Vander Waal forces are present. Hence, it has highest molar heat of vaporisation.
85. (d)  $HI$  is strongest acid because  $H-I$  bond is weakest among halogen acids.
86. (a) In  $HF$  the molecules aggregate because of intermolecular hydrogen bonding. Hence, it has highest boiling point.
88. (a)  $ClO_2^-$  has  $sp^3$ -hybridization and two lone pairs on halogen which produces V-shape Bent structure
- 
89. (d)  $2HClO_4 \rightarrow H_2O + Cl_2O_7$ .
90. (b)  $F_2$  } gases  
 $Cl_2$  }  
 $Br_2$  } liquid  
 $I_2$  } solid
- As we go down the group Vander Waal forces increases. Hence, physical state changes.
92. (c)  $F$  can not act as reducing agent because it has highest reduction potential  
 $F_2 + 2e^- \rightarrow 2F^-; E^\circ = +2.87 V$
93. (c)  $I_2 + 10HNO_3 \rightarrow 2HIO_3 + 10NO_2 + 4H_2O$
94. (a)  $IF_5 + F_2 \rightarrow IF_7$ .
95. (a) Pseudohalide ions and Pseudohalogens  
 There are certain monovalent negative ions made up of two or more electronegative atoms which exhibit properties similar to these of halide ions. Such ions are known as pseudo halide ions just as halide ions, pseudo halide ions have also corresponding dimeric molecules. These are called pseudo halogens and show properties similar to those of halogens.
- |                     |                        |
|---------------------|------------------------|
| Pseudohalide        | Pseudohalogens         |
| $CN^-$ cyanide      | $(CN)_2$ Cyanogen      |
| $SCN^-$ Thiocyanate | $(SCN)_2$ Thiocyanogen |
96. (c)  $NaF$  is highest melting halide because it is most ionic in nature.
98. (b)  $2F_2 + 2H_2O \rightarrow 4HF + O_2$
99. (c)  $CaOCl_2 \rightarrow CaCl_2 + [O]$  Nascent oxygen
100. (a) Generally alkali metals and alkali earth metals elements extracted by the fused electrolysis method.
102. (c)  $2Na_2S_2O_3 + I_2 \rightarrow 2NaI + Na_2S_4O_6$ .
104. (b) Beilstein test - In this test organic compound is heated on a copper wire in a flame. The appearance of a green or bluish green flame due to the formation of volatile cupric halides indicate the presence of halogens in the organic compound. (It does not tell which halogen is actually present).
- Reduction
105. (d)  $Cl_2 + 2KBr \rightarrow Br_2 + 2KCl$   
Oxidation
107. (b)  $3HCl + HNO_3 \rightarrow NOCl + 2H_2O + Cl_2$ .
108. (a)  $Cl-Cl \xrightarrow{U.V} Cl + Cl$   
Free radical
109. (a)  $HF$  is a weak acid due to intermolecular hydrogen bonding.
110. (a) Acidic nature of oxide  $\propto$  Non metallic nature of element  
 Non metallic nature decrease in the order  $Cl > S > P$ .
111. (c) Aqua regia is 1 part of  $HNO_3$  and 3 part of  $HCl$ .
113. (a)  $AgI$  is a covalent compound.
114. (a) Bromine is a liquid at room temperature.
115. (a)  $Cl_2 + H_2O \rightarrow 2HCl + [O]$   
Nascent oxygen
117. (b) The enamel of our teeth is the hardest substance in the body made up of  $CaF_2$  and dentine below it made of  $Ca_3(PO_4)_2$ .



118. (b) As the electronegativity decreases reactivity also decreases.
119. (b)  $KI + I_2 \rightarrow KI_3$   
(soluble complex)
121. (d)  $HI < I_2 < ICl < HIO_4$   
-1      0      +1      +7
122. (a)  $HF < HCl < HBr < HI$   
As we go down the group bond energy decreases hence, acidic nature increases.
123. (b) Caliche is crude chile salt petre ( $NaNO_3$ ) which contains about 0.02% iodine as sodium iodate ( $NaIO_3$ ).
124. (a)  $LiF > LiCl > LiBr > LiI$   
Lattice energy depends on the size and charge of the ion.
125. (a) F-F more strong bond compare to F-Cl, F-Br and Cl-Br bond.
126. (c)  $2Na_2S_2O_3 + I_2 \rightarrow Na_2S_4O_6 + 2NaI$ .
127. (d)  $CaCl(OCl) \Rightarrow Ca \begin{matrix} \swarrow Cl \\ \searrow O-Cl \end{matrix}$
128. (a)  $2NaOH + Cl_2 \rightarrow NaCl + NaOCl + H_2O$   
 $Na^+ + Cl^- \quad Na^+ + OCl^-$
131. (d)  $I_2 + NaF \longrightarrow$   
 $I_2 + NaBr \longrightarrow$   
 $I_2 + NaCl \longrightarrow$  } No reaction  
Because  $I_2$  is least electronegative among halogens.
132. (a)  $HClO_4 > HCl > H_2SO_4 > HNO_3$   
Decreasing order of acidic character.
134. (a)  $HgCl_2 + Hg(CN)_2 \rightarrow HgCl_2 \cdot Hg(CN)_2$   
Mercuric chloride    Mercuric cyanide
135. (a)  $HI > HBr > HCl > HF$   
Acidic character decreasing order.
136. (c)  $Ca(OH)_2 + Cl_2 \rightarrow CaOCl_2 + HOCl$   
Slaked lime                      Bleaching poeder
137. (d)  $K_2Cr_2O_7 + 14HCl \rightarrow 2KCl + 2CrCl_3 + 7H_2O + 3Cl_2$
139. (d) Fluorine does not gives positive oxidation state it is always show -1 oxidation state.
140. (a)  $HClO_4 > HClO_3 > HClO_2 > HCl$   
+7      +5      +3      +1
141. (d)  $Ca(OH)_2 + Cl_2 \rightarrow CaOCl_2 + H_2O$   
bleaching powder
146. (a)  $F_2 + (dil.)2NaOH \rightarrow 2NaF + OF_2 + H_2O$
147. (a) Fluorine can not be oxidised by even strongest oxidising agent.
148. (b)  $Br_2 + 2H_2O + SO_2 \rightarrow H_2SO_4 + 2HBr$
149. (c)  $2KBr + I_2 \rightarrow 2KI + Br_2$   
Iodine is a less electronegative compare to Bromine hence iodine does not change  $Br^-$  to  $Br_2$ .
150. (a)  $CaI_2$  are show covalent properties than other  $CaF_2, CaCl_2, CaBr_2$  compound.
151. (d)  $2KMnO_4 + 3H_2SO_4 + 10HCl \rightarrow K_2SO_4 + 2MnSO_4 + 8H_2O + 5Cl_2$
153. (d)  $PbI_4$  is least stable because of two reasons  
(1) Size of iodine is biggest.  
(2) +2 oxidation state of Pb is more stable than +4 state because of inert pair effect.
154. (a)  $Cl_2 + NaF \rightarrow$  No reaction  
Since  $Cl_2$  is less electronegative then  $F_2$ . Therefore unable to displace fluorine from its salt.
156. (b)  $CS_2 + 3Cl_2 \xrightarrow{I_2} CCl_4 + S_2Cl_2$
157. (b) According to the Fajan's rule largest cation and smallest anion.
158. (a)  $2F_2 + 4NaOH \rightarrow 4HF + 2H_2O + O_2$
159. (c)  $Cl_2 \rightarrow 2Cl \Delta H = +ve$   
1 mole      2 moles  
High temperature and low pressure is favourable.
160. (d)  $BF_3$  accept lone pair of electrons.
161. (d)  $CrO_2Cl_2$  is a orange red gas.
164. (a) Florine always show -1 oxidation state.
165. (a) Solid  $NaF$  is used to purify fluorine i.e. by removing of  $HF$  fumes.
166. (c)  $KHF_2 \rightarrow KF + HF$   
 $KF \rightarrow K^+ + F^-$   
At cathode :  $K^+ + e^- \rightarrow K$   
 $2K + 2HF \rightarrow 2KF + H_2$   
At anode :  $F^- \rightarrow F + e^-$   
 $F + F \rightarrow F_2$
168. (c) Small atomic size of  $Li$  and  $F$  lattice energy is highest.
169. (b)  $SO_2$  bleaches flower by reduction  
 $2H_2O + SO_2 \rightarrow H_2SO_4 + 2[H]$   
 $2[H] + \text{Coloured flower} \xrightarrow{\text{Reduction}} \text{Colourless reduced flower}$   
This bleaching is temporary because reduced flower again oxidised by air to form coloured flower  
 $Cl_2 + H_2O \rightarrow 2HCl + [O]$   
 $[O] + \text{Coloured flower} \xrightarrow{\text{Oxidation}} \text{Colourless Oxidised flower}$   
This bleaching is permanent because oxidised flower remains colourless.
170. (a) Fluorine does not form oxyacids because it is more electronegative than oxygen.
173. (c)  $MnO_2 + 4HCl \rightarrow MnCl_2 + 2H_2O + Cl_2$   
 $2KMnO_4 + 3H_2SO_4 + 10HCl \rightarrow K_2SO_4 + 2MnSO_4 + 8H_2O + 5Cl_2$

## 826 s and p-Block Elements

175. (b)  $3KClO_3 + 3H_2SO_4 \xrightarrow{\text{Heat}} 3KHSO_4 + HClO_4 + 2ClO_2 + H_2O$
176. (b)  $F > Cl > Br > I$ . As the size increases electronegativity decreases.
177. (c) Ionic radius increases on going down the group because no. of shells increases.
178. (b) Reducing properties increase from  $F$  to  $I$  so it oxidise by nitric acid.  
 $I_2 + 10HNO_3 \rightarrow 2HIO_3 + 10NO_2 + 4H_2O$
179. (a) Fluorine and chlorine are more electronegative than sulphur.
180. (d) Upper halogen can replace lower halogen from their compounds solution.
181. (a) Iodine ( $I_2$ ) is slightly soluble in water but it dissolves in 10% aqueous solution of  $KI$  due to the formation of potassium triiodide ( $KI_3$ ).
182. (a) Due to highest electronegativity of fluorine the anion  $[F \cdots H - F]^-$  exists as a result of strong hydrogen bond by which  $K^+$  associate to form  $KHF_2$ .
183. (a) Fluorine is the most electronegative element. It does not form oxyfluorides like other halogens. If reacts with  $NaOH$  to form sodium fluoride and oxygen fluoride.  
 $2NaOH + 2F_2 \rightarrow 2NaF + OF_2 + H_2O$
184. (c) Due to unpaired  $e^-$   $ClO_2$  is paramagnetic.
185. (a) Oxidation number of  $HBrO_4$  is more than that of  $HOCl$ ,  $HNO_2$  and  $H_3PO_3$  so it is the strongest acid among these.
186. (a) Chlorine heptachloride ( $Cl_2O_7$ ) is the anhydride of perchloric acid.  
 $2HClO_4 \xrightarrow{\Delta} Cl_2O_7 + H_2O$
187. (c)  $I_2$  forms complex ion  $I_3^-$  in  $KI$  solution due to which it dissolves in it.
10. (b) An oxygen-helium mixture is used artificial respiration in deep sea diving instead of air because nitrogen present in air dissolves in blood under high pressure when sea diver goes into deep sea. When he comes to the surface, nitrogen bubbles out of the blood due to decrease in pressure, causing pains. This disease is called "bends".
11. (c)  $XeF_2, XeOF_2, XeF_4, XeOF_4, XeF_6, XeO_3$ .
12. (c) **Gas** (Abundance in air by volume (ppm))
- |         |      |
|---------|------|
| Helium  | 5.2  |
| Neon    | 18.2 |
| Argon   | 93.4 |
| Krypton | 1.1  |
| Xenon   | 0.09 |
13. (c) Neon  $\rightarrow Ne$  is monoatomic and others are diatomic  $N_2, F_2$  and  $O_2$ .
14. (c)  ${}_1H^2 + {}_1H^2 \rightarrow {}_2He^4$
15. (b)  $HeF_4$  does not exist.
16. (d)  $Ar_{18} \rightarrow 2, 8, 8$
17. (b)  $Ne_{10} \rightarrow 1s^2 2s^2 2p^6$
23. (d)  $XeF_2$  has  $sp^3d$ -hybridization with linear shape



24. (b) Partial hydrolysis;  $XeF_4 + H_2O \rightarrow XeOF_2 + 2HF$   
 Complete hydrolysis;  
 $2XeF_4 + 3H_2O \rightarrow Xe + XeO_3 + F_2 + 6HF$
26. (d)  $He$  is least polarizable because of small atomic size.
27. (a)  $Rn$  because it is radioactive element obtained by the disintegration of radium  
 ${}_{88}Ra^{206} \rightarrow {}_{86}Rn^{202} + {}_2He^4$
30. (c)  $1s^2 2s^2 2p^6 \rightarrow Neon$
35. (a)  $XeF_6 + 3H_2O \rightarrow XeO_3 + 6HF$
36. (c)  $Xe > Kr > Ar > Ne > He$   
 Solubility in decreasing order.
37. (d)




## Noble gases

3. (a) Gas -  $H_2 \quad O_2 \quad He \quad N_2$   
 Mol. mass - 2 32 4 28
5. (c) Helium (In Greek Helios = Sun)
6. (d) All the noble gases are monoatomic, colourless and odourless gases. Their monoatomic nature is due to the stable outer configuration  $ns^2 np^6$  of their atoms. As a result, they do not enter into chemical combination even amongst themselves.
7. (c) Except  $He$ , all other noble gases are adsorbed by coconut charcoal at low temperatures. The extent of adsorption increases as the atomic size of the noble gas increases.

$sp^3d^2$  hybridization       $sp^3d^3$  hybridization



38. (b) Zero group element are show less chemically activity because this group element have 8 electron.

39. (d) Xe is formed following compounds.




40. (a) As the number of shells increases, size increases and the effective nuclear charge on the outermost electron decreases. Thus, I.E. decreases.

44. (c)  $XeF_2, XeF_4$  &  $XeF_6$  can be directly prepared



$XeO_3$  is obtained by the hydrolysis of  $XeF_6$



46. (a)  $XeO_3$  shows  $sp^3$  hybridization.


47. (a) It is because

- (1) Small atomic size
- (2) High Ionization energy
- (3) Absence of  $d$ -orbitals


48. (a) Zero group element are attached with weak intermolecular force.

49. (b)  $XeF_2, XeF_4, XeF_6$ .


50. (d)  $XeO_3$  :



$XeOF_4$  :



$XeF_6$  :



51. (d) Neil Bartlett prepared first noble gas compound. Xenon hexafluoroplatinate (IV).

53. (d) He, Ne, and Kr all are found in very little amount in atmosphere, so all are called rare gas.

54. (c) Helium is twice as heavy as hydrogen, its lifting power is 92% of that of hydrogen. Helium has the lowest melting and boiling

point of any element which makes liquid helium an ideal coolant for many extremely low temperature application such as super conducting magnet and cryogenic research where temperature close to absolute zero are needed.

55. (a) The maximum temperature at which gas can be liquified is called its critical temperature. The gas which have high boiling point will change into liquid first and so critical temperature of the gas will be more.

$$T_c \propto \text{B.P.} \text{ and } \text{B.P.} \propto \text{Molecular weight}$$

So Kr liquifies first.

56. (c) Suppose the oxidation state of Xe is  $x$ .  
 $XeOF_2$

$$x + (-2) + 2(-1) = 0 \Rightarrow x - 2 - 2 = 0 \Rightarrow x = 4.$$

57. (a) 

He	Ne	Ar	Kr	Xe
Rn				


Boiling point of - 269 -246 - 186 -153.6 - 108.1 -62

Inert gases

59. (d) Xe is highly polar since the ionisation potential of xenon is quite close to the ionisation potential of oxygen.

60. (d) In the formation of  $XeF_4, sp^3d^2$  hybridisation


occurs which gives the molecule an octahedral structure. The xenon and four fluorine atoms are coplanar while the two equatorial positions are occupied by the two lone pairs of electrons.



### Critical Thinking Questions

1. (c) Correct sequence is  
 $NH_2CONH_2 > NH_4N_3 > NH_3NO_3 > NH_4Cl$ .
2. (d) Second group elements are show strong reducing properties but less 1<sup>st</sup> group element show less.
3. (a) The size of alkaline earth metals is smaller as compared to its corresponding alkali metals and its effective nuclear charge is also more than that of its corresponding alkali metals.
4. (d) Lead is maximum in flint glass.
5. (b)  $BaSO_4 + 4C \xrightarrow{\text{Heat}} BaS + 4CO$   
Thus, on heating they produce  $BaS + 4CO$ .
6. (d) Smaller the atomic size tendency of hydration is more as the size increases tendency for hydration also decreases.
7. (a) Fusion mixture is  $Na_2CO_3 + K_2CO_3$ .
8. (b)  $HCl$  is a gas.
9. (a) (A) Peroxide is  $H_2O_2(4)$ ; (B) Super oxide is  $KO_2(3)$


## 828 s and p-Block Elements

- (C) Dioxide is  $PbO_2(2)$ ; (D) Suboxide is  $C_3O_2(1)$
10. (a)  $H_2SO_4 + SO_3 \rightarrow H_2S_2O_7$   
98% Oleum (Fuming  $H_2SO_4$ )
11. (c)  $SnO_2 + 2NaOH \rightarrow Na_2SnO_3 + H_2O$   
 $SnO_2 + 4HCl \rightarrow SnCl_4 + 2H_2O$
12. (b)  $NaOH + Al^{+3} \rightarrow NaAlO_2$   
Sod. metaaluminate (Soluble in  $NaOH$ )  
 $NaOH + Fe^{+3} \rightarrow$  No reaction (Insoluble in  $NaOH$ )
13. (d) The composition of the common glass is  $Na_2O.CaO.6SiO_2$ .
14. (c) Lead form nitric oxide with dil.  $HNO_3$   
 $3Pb + 8HNO_3 \rightarrow 3Pb(NO_3)_2 + 2NO + 4H_2O$
15. (b)  Therefore, total number of lone pair of electron on central atom Xenon = 1
16. (d) Order of acidic strength is  $H_2Te > H_2Se > H_2S > H_2O$   
 $Na_2O$  is a salt of  $NaOH + H_2O$  and  $H_2O$  is least acidic among given acids hence  $pH$  in this case will be maximum.
17. (a)  $HNO_3$  is the strong oxidising acid so it react with alkali while rest can be react with both and alkali.
18. (d) Zero group is called a buffer group because it lies between highly electronegative halogens and highly electropositive alkali metal element.
19. (c)  $PbSO_4$  is insoluble in water.
20. (a)  $N_2O_5$  is most acidic.
21. (b) Element - Fluorine Chlorine  
 Bromine Iodine  
 B.E. in  $kJ\ mole^{-1}$  - 158.8 242.6 192.8 151.1
22. (d)  $CaCN_2 + 3H_2O \xrightarrow[\text{pressure}]{\text{under}}$   $CaCO_3 + 2NH_3$ .
23. (a)  $6O_2 \xrightarrow{\text{Ozonolysis}}$   $4O_3$   
 Complete ozonolysis gives 4 volume of ozone.
24. (b) Marble statue  $\rightarrow CaCO_3$ ; Calcinated Gypsum  $\rightarrow CaSO_4.2H_2O$ ; Sea shell  $\rightarrow CaCO_3$ ; Dalomite  $\rightarrow CaCO_3.MgCO_3$ .
25. (b) Sodium is basic in nature.
26. (c)  $I_2 + I^- \rightarrow I_3^-$
27. (a) The hydration energy decreases from  $Be^{+2}$  to  $Ba^{+2}$ . This causes the decrease in the solubilities of the sulphates in this order. In other words, the solubilities decreases with increase in the ionic size.
28. (d)  $2Mg + O_2 \rightarrow 2MgO$   
 $3Mg + N_2 \rightarrow Mg_3N_2$
29. (a)  $ZnO + BaO \xrightarrow{1100^\circ C} BaZnO_2$
30. (b) There are no free electron in  $N_2O_4$ , so it is dimagnetic in nature.
31. (b)  $CaSO_4.2H_2O \xrightarrow{120^\circ C} CaSO_4 + \frac{1}{2}H_2O + 1\frac{1}{2}H_2O$   
Gypsum Plaster of paris
32. (d) Third alkaline earth metal is calcium  ${}_{20}Ca^{40}$ .  
 No. of electron are 20 & No. of proton are 20;  $e/20, p/20$ .
33. (a) In the compounds of alkaline earth metals all the electrons are paired. Hence, they are diamagnetic in nature.
34. (a) Mixture of helium and oxygen is used in artificial respiration.
35. (a) Alkaline solution of pyrogallol quickly absorbs oxygen.
36. (d)  $NH_4NO_2 \rightarrow N_2 + 2H_2O$   
 $2NaN_3 \rightarrow 2Na + 3N_2$   
 $(NH_4)_2Cr_2O_7 \rightarrow N_2 + Cr_2O_3 + 4H_2O$
37. (b) Because yellow phosphorus is most reactive form of phosphorus and is highly polymerised.
38. (c) Carbon has 2 electrons in their penultimate shell configuration so due to  $d$ -orbital in penultimate shell is false statement.
39. (d) All other oxides of nitrogen except  $N_2O$  and  $NO$  are acidic in nature.
40. (a)  $PbCl_4 < PbCl_2 < CaCl_2 < NaCl$  is the increasing order of ionic character.
41. (a)  $Si + 3HCl \rightarrow SiHCl_3 + H_2$   
silicon chloroform
42. (a)  $KO_2$  is used in oxygen cylinder because it absorbs  $CO_2$  and increases  $O_2$  content. Super oxides reacts with water to give  $H_2O_2$  &  $O_2$ .
43. (d)  $NaHCO_3$  and  $Na_2CO_3$  decomposes to give  $CO_2$  which extinguish fire.
44. (d)  $CaF_2$  is insoluble in water.
45. (a)  $PCl_3$  is most acidic.
46. (b)  $CsCl > KCl > NaCl > LiCl$
47. (d)  $2Na_2S_2O_3 + I_2 \rightarrow Na_2S_4O_8 + 2NaI$ .
48. (b) In general, higher the oxidation state, more is the covalent character of the oxide.  $I_2O_7$  &  $I_2O_5$  do not exist.  $I_2O_4$  is ionic in






## 830 s and p-Block Elements



16. (c) Boron is metalloid. Thus assertion is correct. Metalloids possess, metallic as well as non-metallic nature. Hence, reason is false.
17. (b) It is correct that inert gases are monoatomic because for inert gases  $C_p / C_v = 1.66$ .
18. (c) When *Mg* is burnt in nitric oxide it continues to burn because during burning the heat evolved decomposes *NO* to *N<sub>2</sub>* and *O<sub>2</sub>*. Oxygen thus, produced helps *Mg* to burn.  
Here assertion is true but reason is false.
19. (d) Anhydrous *BaO<sub>2</sub>* is not used for preparing *H<sub>2</sub>O<sub>2</sub>* because it reacts with *H<sub>2</sub>SO<sub>4</sub>* and the reaction ceases after some time due to formation of *BaSO<sub>4</sub>* on *BaO<sub>2</sub>*. Therefore, assertion and reason both are false.
20. (d) Inorganic benzene, borazine is highly reactive while benzene is much less reactive.  
Here, assertion is false, but reason is true.
21. (a) The halogens absorb visible light due to which all halogens are coloured. Hence, both assertion and reason are true and reason is correct explanation.
22. (b) It is true that barium is not required for normal biological function in human beings and it is also true that it does not show variable oxidation state. It only shows +2 oxidation state.
23. (d) The *O–O* bond length is shorter in *O<sub>2</sub>F<sub>2</sub>* than in *H<sub>2</sub>O<sub>2</sub>* due to higher electronegativity. *H<sub>2</sub>O<sub>2</sub>* is a non-ionic compound. Here both assertion and reason are false.
24. (d) Here both assertion and reason are false because *PbI<sub>4</sub>* is not a stable compound and iodine cannot stabilize higher oxidation states. *Pb* shows (II) oxidation state more frequently than *Pb* (IV) due to inert pair effect.
25. (b) Both assertion and reason are true but reason is not correct explanation of the assertion. Enamel, the hardest substance of the body is composed of fluorine, not magnesium. Magnesium is an essential element as it acts as a factor of many enzymes of glycolysis and a number of other metabolic reactions dependent upon ATP.
26. (d) Both assertion and reason are false.  
Radium is the rarest of all s-block elements comprising only  $10^{-10}$  percent of igneous rocks. Francium (s-block member) is radioactive; its long-lived isotope <sup>223</sup>*Fr* has a half-life of only 21 minutes.
27. (c) Assertion is true but reason is false.  
Due to high polarizing power of *Li<sup>+</sup>*, *LiCl* is a covalent compound.
28. (c) Assertion is true but reason is false.  
*Be* has fully filled  $2s^2$  – orbital which gives a relatively more stable electronic configuration.
29. (a) Both assertion and reason are true and reason is the correct explanation of assertion.
30. (a) Both assertion and reason are true and reason is the correct explanation of assertion.
31. (a) Both assertion and reason are true and reason is the correct explanation of assertion.
32. (c) Assertion is true but reason is false.  
Helium is a noble gas (chemically inactive) but beryllium is a member of alkaline earth metals (chemically active).
33. (b) Both assertion and reason are true but reason is not the correct explanation of assertion.  
Lattice energy of *Na<sub>2</sub>SO<sub>4</sub>* is less than its hydration energy but the lattice energy of *BaSO<sub>4</sub>* exceeds its hydration energy.
34. (a) Both assertion and reason are true and reason is the correct explanation of assertion.
35. (a) Both assertion and reason are true and reason is the correct explanation of assertion.  
Presence of unpaired electrons in superoxides of alkali metals makes them paramagnetic.
36. (b) Both assertion and reason are true but reason is not the correct explanation of assertion.  
Nitrogen cannot expand its octet due to the non-availability of *d*-orbital.
37. (a) Both assertion and reason are true and reason is the correct explanation of assertion.  
Ozone is considered to be a resonance hybrid of the following two forms.



38. (b) Both assertion and reason are true but reason is not the correct explanation of assertion.  
Molecules of sulphuric acid are associated due to a large number of intermolecular hydrogen bonds.
39. (b) Both assertion and reason are true but reason is not the correct explanation of assertion.  
*PCl<sub>5</sub>* is trigonal bipyramidal containing  $sp^3d$  hybridized *P* atom in liquid and gaseous state. Whereas in solid state it consists of tetrahedral *PCl<sub>4</sub><sup>+</sup>* cation and octahedral *PCl<sub>6</sub><sup>-</sup>* anion.

40. (b) Both assertion and reason are true but reason is not the correct explanation of assertion.

In case of  $NI_3$ , the lone pair moment adds on the resultant of the  $N-I$  moments but in case of  $NF_3$ , the lone pair moment on partly cancels the resultant  $N-F$  moments.

41. (a) Both assertion and reason are true and reason is the correct explanation of assertion.

The ignition temperature of white phosphorus is low (About  $30^\circ\text{C}$ ). In air it readily catches fire giving dense fumes of phosphorus pentoxide. It is therefore, kept in water.

42. (c) Assertion is true but reason is false.

$B$  does not have vacant  $d$ -orbitals as second shell is the outermost shell.

43. (a) Both assertion and reason are true and reason is the correct explanation of assertion.

44. (d) Both assertion and reason are false.


$Si-Si$  bonds are weaker than  $Si-O$  bonds  $Si$  has no tendency to form double bonds with itself.

45. (c) Assertion is true but reason is false.

$S$  atoms in  $S_8$  molecule undergo  $sp^3$  hybridization and contain two lone pairs of electrons on each and exists as staggered 8 atom rings.

46. (a) Both assertion and reason are true and reason is the correct explanation of assertion.

This can be explained through structure of Caro's acid (Peroxomonosulphuric acid).



Oxidation no. of  $S = x$ , oxidation no. of  $H = +1$ ,

Oxidation no. of oxygen in peroxo linkage =  $-1$ ,

Oxidation no. of other oxygen atoms =  $-2$  (each)  $2 + x - 6 - 2 = 0$  or  $x = +6$ .

47. (d) Both assertion and reason are false.

The melting point / boiling point of noble gases are quite low. The inter particle forces among noble gases are weak Vander Waal's forces.

48. (b) If both assertion and reason are true and reason is the correct explanation of assertion.

$S$  atom in both  $SO_2$  and  $SO_3$  is  $sp^2$  hybridized but it contains a lone pair of electrons in  $SO_2$ .

49. (d) Both assertion and reason are false.

Calcium carbide on hydrolysis gives acetylene. Calcium carbide contains  $C_2^{2-}$  anion.

50. (a) Both assertion and reason are true and reason is the correct explanation of assertion.

51. (e) Assertion is false but reason is true.

In lab, hydrogen is generally prepared by the reaction of zinc with dilute hydrochloric acid.

52. (b) Both assertion and reason are true but reason is not the correct explanation of assertion.

The relatively inert behaviour of dihydrogen at room temperature is due to the high enthalpy of  $H-H$  bond, being the highest for a single bond between any two elements.

53. (d) Both assertion and reason are false.

Water can be easily transformed from liquid solid and to gaseous states. The distribution of water over the earth's surface is not uniform. The desert region have no permanent surface water while the oceans cover vast areas.

54. (b) Both assertion and reason are true but reason is not the correct explanation of assertion.

The structure of ice is open due to hydrogen bonding which makes ice less dense than liquid water at the same temperature.

55. (c) Assertion is true but reason is false.

The water molecules are joined together in an extensive three dimensional network in which each oxygen atom is bonded to four hydrogen atoms two by hydrogen bonds and two by normal covalent bonds in a near tetrahedral configuration. This situation does not exist for molecules like  $NH_3$  and  $HF$ .

56. (a) Both assertion and reason are true and reason is the correct explanation of assertion.

Hard water contain soluble calcium and magnesium salt like bicarbonates, chlorides and sulphates.

57. (a) Both assertion and reason are true and reason is the correct explanation of assertion.

To stop decomposition  $H_2O_2$  is stored in wax-lined glass or plastic vessels in the presence of stabilizers like urea.

58. (a) Both assertion and reason are true and reason is the correct explanation of assertion.

At any given instant in liquid water at room temperature, each water molecule forms hydrogen bonds with an average 3.4 other water molecules. The  $H_2O$  molecules are in continuous motion so hydrogen bonds are constantly and rapidly broken and formed. In ice  $H_2O$  molecules are, however fixed in the space lattice.

59. (d) Both assertion and reason are false.




## 832 s and p-Block Elements

Calgon is used for making  $Ca^{2+}$  and  $Mg^{2+}$  ions present in hard water ineffective. It forms soluble complex with  $Ca^{2+}$  and  $Mg^{2+}$  ions.

60. (b)  $SO_2$  shows both oxidising as well as reducing nature.

The reaction given in assertion is due to oxidising nature of  $SO_2$ .



61. (a) (i) Due to smaller size of  $F$ ; steric repulsions will be less in  $SiF_6^{2-}$ .


(ii) Interaction of  $F$  lone pair electrons with  $Si$  is stronger than that of chlorine lone pairs.

62. (b) Borax bead test is not suitable for  $Al(III)$  because its oxidising as well as reducing flame is colourless in both hot as well as cold.

Alumina is insoluble in water as they exist in hydrated form like  $Al_2O_3 \cdot 2H_2O$ ,  $Al_2O_3 \cdot H_2O$  etc.

63. (c)  $SeCl_4$  possess see saw geometry, which can be regarded as a distorted trigonal bipyramidal structure having one lone pair (lp) of electrons in the basal position of the trigonal bipyramidal. See-saw geometry of  $SeCl_4$  molecules arises due to the  $sp^3d$  hybridisation of the central atom. The distortion in shape is due to the presence of one lone pair of electrons.

64. (b) Due to the ease with which it can liberate nascent oxygen,  $O_3$  acts as a powerful oxidising agent.



$:O \cdot \cdot \cdot \cdot$   
paramagnetic  
(due to presence of  
two unpaired

