





Ordinary Thinking

Objective Questions

General Introduction of Carboxylic Acids and Their Derivatives





- Identify the wrong statement from the following
[Tamil Nadu CET 2002]
 - Salicylic acid's a monobasic acid
 - Methyl salicylate is an ester
 - Salicylic acid gives violet colour with neutral ferric chloride as well as brisk effervescence with sodium bicarbonate
 - Methyl salicylate does not occur in natural oils
- Which of the following is optically active[BHU 1997]
 - Ethylene glycol
 - Oxalic acid
 - Glycerol
 - Tartaric acid
- Palmitic acid is
[BHU 1997]
 - $C_{16}H_{31}COOH$
 - $C_{17}H_{35}COOH$
 - $C_{15}H_{31}COOH$
 - $C_{17}H_{31}COOH$
- Which one among the following represents an amide
[MP PMT 1993]
 - 
 - 
 - 
 - 
- The name of the compound having the structure $ClCH_2CH_2COOH$ is
[MP PET 1993]
 - 3-chloropropanoic acid
 - 2-chloropropanoic acid
 - 2-chloroethanoic acid
 - Chlorosuccinic acid
- Fats and oils are mixture of
[CPMT 1993]
 - Glycerides and saturated fatty acids
 - Glycerides and unsaturated fatty acids
 - Glycerides of saturated and unsaturated fatty acids
 - Only saturated and unsaturated fatty acids
- Which one is not a glyceride
[CPMT 1994]
 - Fat
 - Oil
 - Phospholipid
 - Soaps
- $(RCO)_2NH$ is
 - Primary amine
 - Secondary amine
 - Secondary amide
 - Tertiary amide
- Which of the following is the formula of tartaremetic
[Manipal MEE 1995]
 - $CH(OH)COOH$
 - $CH(OH)COONa$
 - $CH(OH)COOK$
 - $CH(OH)COO(SbO)$
- Which compound is known as oil of winter green
[MP PET/PMT 1998; CPMT 2002]
 - Phenyl benzoate
 - Phenyl salicylate
 - Phenyl acetate
 - Methyl salicylate
- Which of the following structure of carboxylic acid accounts for the acidic nature
 - $R-C \begin{matrix} \nearrow O \\ \searrow OH \end{matrix}$
 - $R-C^+ \begin{matrix} \nearrow OH \\ \searrow OH \end{matrix}$
 - $R-C \begin{matrix} \nearrow O \\ \searrow H \end{matrix}$
 - None of these
- Acetoacetic ester behaves as
[CPMT 1988]
 - An unsaturated hydroxy compound
 - A keto compound
 - Both of these ways
 - None of these
- The general formula $(RCO)_2O$ represents
[CPMT 1974; DPMT 1982; MP PMT 1996]
 - An ester
 - A ketone
 - An ether
 - An acid anhydride
- A tribasic acid is
 - Oxalic acid
 - Tartaric acid
 - Lactic acid
 - Citric acid
- Amphiphilic molecules are normally associated with
[Orissa JEE 1997]
 - Isoprene based polymers
 - Soaps and detergents
 - Nitrogen based fertilizers e.g. urea
 - Pain relieving medicines such as aspirin
- Wax are long chain compounds belonging to the class
[CPMT 1982, 93]
 - Acids
 - Alcohols
 - Esters
 - Ethers
- Glycine may be classed as all of the following except
[JIPMER 1997]
 - A base
 - An acid
 - A zwitter ion
 - Optically active acid
- Which of the following is not a fatty acid[CPMT 1988]
 - Stearic acid
 - Palmitic acid
 - Oleic acid
 - Phenyl acetic acid
- Vinegar obtained from canesugar contains
[CPMT 1980; DPMT 1982; KCET 1992; MP PMT 1994; AIIMS 1999]
 - Citric acid
 - Lactic acid
 - Acetic acid
 - Palmitic acid
- The general formula for monocarboxylic acids is
[CPMT 2003]
 - C_nH_nCOOH
 - $C_nH_{2n+1}COOH$

1338 Carboxylic acids and Their derivatives


- (c) $C_nH_{2n-1}COOH$ (d) $C_nH_{2n}O_2$
21. Number of oxygen atoms in a acetamide molecule is
(a) 1 (b) 2
(c) 3 (d) 4
22. Urea is [CPMT 1984]
(a) Monoacidic base (b) Diacidic base
(c) Neutral (d) Amphoteric
23. Fats and oils are [CPMT 1990]
(a) Acids (b) Alcohols
(c) Esters (d) Hydrocarbons
24. The general formulas $C_nH_{2n}O_2$ could be for open chain [AIIEE 2003]
(a) Diketones (b) Carboxylic acids
(c) Diols (d) Dialdehydes
25. $H - \overset{O}{\parallel} C - Cl$ is called
(a) Acetyl chloride (b) Formyl chloride
(c) Chloretone (d) Oxochloromethane
26. Urea
(a) Is an amide of carbonic acid
(b) It is diamide of carbonic acid
(c) Gives carbonic acid on hydrolysis
(d) Resembles carbonic acid
27. Which of the following acids is isomeric with phthalic acid
(a) Succinic acid
(b) Salicylic acid
(c) 1, 4-benzene dicarboxylic acid
(d) Methyl benzoic
28. The ester among the following is [Kerala PMT 2003]
(a) Calcium lactate (b) Ammonium acetate
(c) Sodium acetate (d) None of these
29. Sodium or potassium salts of higher fatty acids are called [MP PET 2003]
(a) Soaps (b) Terpenes
(c) Sugars (d) Alkaloids
30. Formamide is
(a) $HCONH_2$ (b) CH_3CONH_2
(c) $HCOONH_4$ (d) $(HCHO + NH_3)$
31. Oleic, stearic and palmitic acids are [CPMT 1997]
(a) Nucleic acids (b) Amino acids
(c) Fatty acids (d) None of these
32. Which one is called ethanoic acid [CPMT 1997]
(a) $HCOOH$ (b) CH_3COOH
(c) CH_3CH_2COOH (d) $CH_3CH_2CH_2COOH$
33. Vinegar is [CPMT 1997]
(a) $HCHO$ (b) $HCOOH$
(c) CH_3CHO (d) CH_3COOH
34. Which of these do not contain $-COOH$ group [CPMT 1997]
(a) Aspirin (b) Benzoic acid
(c) Picric acid (d) Salicylic acid
35. Vinegar obtained from sugarcane has [AFMC 2005]
(a) CH_3COOH (b) $HCOOH$
(c) C_6H_5COOH (d) CH_3CH_2COOH
36. Carbolic acid is [AFMC 2005]
(a) C_6H_5CHO (b) C_6H_6
(c) C_6H_5COOH (d) C_6H_5OH
37. The most acidic of the following is [J & K 2005]
(a) $ClCH_2COOH$ (b) C_6H_5COOH
(c) CD_3COOH (d) CH_3CH_2COOH
38. Which is most reactive of the following [J & K 2005]
(a) Ethyl acetate (b) Acetic anhydride
(c) Acetamide (d) Acetyl chloride

Preparation of Carboxylic Acids and Their Derivatives

1. Hydrolysis of $CH_3CH_2NO_2$ with 85% H_2SO_4 gives [KCET 1996]
(a) CH_3CH_2OH (b) C_2H_6
(c) $CH_3CH = NOH$ (d) CH_3COOH
2. When formic acid reacts with PCl_5 it forms [MNR 1982]
(a) Formyl chloride (b) Acetyl chloride
(c) Methyl chloride (d) Propionyl chloride
3. Laboratory method for the preparation of acetyl chloride is [RPMT 2003]
(a) $CH_3COOH + SOCl_2 \rightarrow CH_3COCl$
(b) $CH_3COOH + PCl_3 \rightarrow CH_3COCl$
(c) $CH_3COONa + PCl_3 \rightarrow CH_3COCl$
(d) All of these
4. Tischenko reaction yields ester in the presence of catalyst which is
(a) $LiAlH_4$ (b) N -bromosuccinamide
(c) $Al(OC_2H_5)_3$ (d) $Zn - Hg / HCl$
5. Acetic acid is obtained when [NCERT 1975; CPMT 1977]
(a) Methyl alcohol is oxidised with potassium permanganate
(b) Calcium acetate is distilled in the presence of calcium formate
(c) Acetaldehyde is oxidised with potassium dichromate and sulphuric acid
(d) Glycerol is heated with sulphuric acid
6. Acetic acid is manufactured by the fermentation of [CPMT 1985]
(a) Ethanol (b) Methanol
(c) Ethanal (d) Methanal
7. $CO + NaOH \rightarrow$ [CPMT 1997; KCET 1999]
(a) $HCOONa$ (b) $C_2H_2O_4$
(c) $HCOOH$ (d) CH_3COOH

8. Carboxylic acids react with diazomethane to form
 [MP PMT/PET 1988; MP PMT 1990]
 (a) Amine (b) Alcohol
 (c) Ester (d) Amide
9. $C_2H_2 \xrightarrow[H_2SO_4]{HgOH\ 1\%} A \xrightarrow{[O]} B$, B is
 [CBSE PMT 1991; BHU 1995]
 (a) An acid (b) An aldehyde
 (c) A ketone (d) Ethanol
10. Reimer-Tiemann reaction involves a
 [MP PET 1997]
 (a) Carbonium ion intermediate
 (b) Carbene intermediate
 (c) Carbanion intermediate
 (d) Free radical intermediate
11. The product D of the reaction
 $CH_3Cl \xrightarrow{KCN} (A) \xrightarrow{H_2O} (B) \xrightarrow{NH_3} (C) \xrightarrow{\Delta} (D)$ is
 [MP PET 1997]
 (a) $CH_3CH_2NH_2$ (b) CH_3CN
 (c) $HCONH_2$ (d) CH_3CONH_2
12. Which of the following on hydrolysis forms acetic acid
 [BHU 1997]
 (a) CH_3CN (b) CH_3OH
 (c) C_2H_5OH (d) $C_2H_5NH_2$
13. When benzyl alcohol is oxidised with $KMnO_4$, the product obtained is
 [SCRA 1991]
 (a) Benzaldehyde (b) Benzoic acid
 (c) CO_2 and H_2O (d) None of these
14. Which of the following gives benzoic acid on oxidation
 [CBSE PMT 1996]
 (a) Chlorophenol (b) Chlorotoluene
 (c) Chlorobenzene (d) Benzyl chloride
15. $(CH_3)_2CO \xrightarrow[HCl]{NaCN} A \xrightarrow[\Delta]{H_3O^+} B$ In the above sequence of reactions A and B are [CPMT 2000]
 (a) $(CH_3)_2C(OH)CN, (CH_3)_2C(OH)COOH$
 (b) $(CH_3)_2C(OH)CN, (CH_3)_2C(OH)_2$
 (c) $(CH_3)_2C(OH)CN, (CH_3)_2CHCOOH$
 (d) $(CH_3)_2C(OH)CN, (CH_3)_2C=O$
16. Two moles of acetic acid are heated with P_2O_5 . The product formed is [MP PET/PMT 1988]
 (a) 2 moles of ethyl alcohol
 (b) Formic anhydride
 (c) Acetic anhydride
 (d) 2 moles of methyl cyanide
17. Formic acid is obtained when [NCERT 1974]
 (a) Calcium acetate is heated with conc. H_2SO_4
 (b) Calcium formate is heated with calcium acetate
 (c) Glycerol is heated with oxalic acid at $110^\circ C$
 (d) Acetaldehyde is oxidised with $K_2Cr_2O_7$ and H_2SO_4
18. Acetyl chloride cannot be obtained by treating acetic acid with
 (a) $CHCl_3$ (b) $SOCl_2$
 (c) PCl_3 (d) PCl_5
19. o-xylene when oxidised in presence of V_2O_5 the product is
 (a) Benzoic acid (b) Phenyl acetic acid
 (c) Phthalic acid (d) Acetic acid
20. The reaction
 $CH_3CH=CH_2 \xrightarrow[H^+]{CO+H_2O} CH_3-\underset{\substack{| \\ COOH}}{CH}-CH_3$
 is known as [MP PMT 2002]
 (a) Wurtz reactions
 (b) Koch reaction
 (c) Clemenson's reduction
 (d) Kolbe's reaction
21. By aerial oxidation, which one of the following gives phthalic acid [Tamil Nadu CET 2002]
 (a) Naphthalene (b) Benzene
 (c) Mesitylene (d) Toluene
22.  In the reaction, product P is [CBSE PMT 2002]
 (a)  (b) 
 (c)  (d) $C_6H_5-\overset{O}{\parallel}C-C_6H_5$
23. Glacial acetic is obtained by [KCET 2002]
 (a) Distilling vinegar
 (b) Crystallizing separating and melting acetic acid
 (c) Treating vinegar with dehydrating agent
 (d) Chemically separating acetic acid
24. In esterification, OH^- ion for making H_2O comes from [CPMT 1996]
 (a) Acid (b) Alcohol
 (c) Ketone (d) Carbohydrate
25. Heating a mixture of ethyl alcohol and acetic acid in presence of conc. H_2SO_4 produces a fruity smelling compound. This reaction is called
 (a) Neutralisation (b) Ester hydrolysis

1340 Carboxylic acids and Their derivatives

- (c) Esterification synthesis (d) Williamson's synthesis
26. Product formed by heating a mixture of ammonium chloride and potassium cyanate is
 (a) N_2O (b) NH_3
 (c) CH_3NH_2 (d) H_2NCONH_2
27. Rearrangement of an oxime to an amide in the presence of strong acid is called [Kerala CET 2000]
 (a) Curtius rearrangement (b) Fries rearrangement
 (c) Backman rearrangement (d) Sandmeyer reaction
28. Which reagent will bring about the conversion of carboxylic acids into esters
 (a) C_2H_5OH (b) Dry $HCl + C_2H_5OH$
 (c) $LiAlH_4$ (d) $Al(OC_2H_5)_3$
29. The acid formed when propyl magnesium bromide is treated with carbon dioxide is [CPMT 1982, 84, 86; Pb. PMT 1998]
 (a) C_3H_7COOH (b) C_2H_5COOH
 (c) Both (a) and (b) (d) None of the above
30. CO_2 on reaction with ethyl magnesium bromide gives [BHU 1983]
 (a) Ethane (b) Propanoic acid
 (c) Acetic acid (d) None of these
31. Acetic anhydride is obtained from acetyl chloride by the reaction of [CPMT 1985, 93]
 (a) P_2O_5 (b) H_2SO_4
 (c) CH_3COONa (d) CH_3COOH
32. Hydrolysis of acetamide produces [DPMT 1984; MP PMT 1994; MP PET 2001]
 (a) Acetic acid (b) Acetaldehyde
 (c) Methylamine (d) Formic acid
33. Ethyl acetate is obtained when methyl magnesium iodide reacts with [Tamil Nadu CET 2002]
 (a) Ethyl formate (b) Ethyl chloroformate
 (c) Acetyl chloride (d) Carbon dioxide
34. Sodium acetate reacts with acetyl chloride to form [BIT 1992]
 (a) Acetic acid (b) Acetone
 (c) Acetic anhydride (d) Sodium formate
35. Ammonium acetate reacts with acetic acid at $110^\circ C$ to form [MP PET 1991]
 (a) Acetamide (b) Formamide
 (c) Ammonium cyanate (d) Urea
36. Tischenko reaction is used for preparation of
 (a) Ether (b) Ester
 (c) Amide (d) Acid anhydride
37. The silver salt of a fatty acid on refluxing with an alkyl halide gives an [KCET 2004]
 (a) Acid (b) Ester
 (c) Ether (d) Amine
38. Which reaction is used for the preparation of α -Bromoacetic acid? [MP PET 2004; MP PET/PMT 1998]
 (a) Kolbe's Reaction
 (b) Reimer-Tiemann Reaction
 (c) Hell volhard Zelinsky Reaction
 (d) Perkin's Reaction
39. Tertiary alcohols (3°) having atleast four carbon atoms upon drastic oxidation yield carboxylic acid with [MH CET 2004]
 (a) One carbon atom less
 (b) Two carbon atoms less
 (c) Three carbon atom less
 (d) All the above three options are correct
40. When succinic acid is heated, product formed is [CBSE PMT 2000] [Pb. CET 2000]
 (a) Succinic anhydride (b) Acetic acid
 (c) CO_2 and methane (d) Propionic acid
41. In the reaction, $C_6H_5OH \xrightarrow{NaOH} (A) \xrightarrow[140^\circ C, (4-7 \text{ atm})]{CO_2} (B) \xrightarrow{HCl} (C)$, the compound (C) is
 (a) Benzoic acid (b) Salicylaldehyde
 (c) Chlorobenzene (d) Salicylic acid
42. When an acyl chloride is heated with Na salt of a carboxylic acid, the product is
 (a) An ester (b) An anhydride
 (c) An alkene (d) An aldehyde
43. The compound X, in the reaction, is
 $X \xrightarrow{CH_3MgI} Y \xrightarrow{\text{hydrolysis}} Mg(OH)I + CH_3COOH$ [Pb. CET 2003]
 (a) CH_3CHO (b) CO_2
 (c) $(CH_3)_2CO$ (d) $HCHO$
44. $CH_3CONH_2 \xrightarrow{NaNO_2 / HCl} X$ [Pb. CET 2003]
 (a) CH_3COOH (b) $CH_3CO \overset{+}{N} H_3 Cl^-$
 (c) CH_3NH_2 (d) CH_3CHO
45. Primary aldehyde on oxidation gives [DPMT 20004]
 (a) Esters (b) Carboxylic acid
 (c) Ketones (d) Alcohols
46. Toluene is oxidised to benzoic acid by [BHU 2004; CPMT 1985]
 (a) $KMnO_4$ (b) $K_2Cr_2O_7$
 (c) H_2SO_4 (d) Both (a) and (b)
47.  [IIT 2005]
 The compound (X) is
 (a) CH_3COOH (b) $BrCH_2 - COOH$
 (c) $(CH_3CO)_2O$ (d) $CHO - COOH$
48. Salicylic acid is prepared from phenol by [AFMC 2005]
 (a) Reimer Tiemann reaction
 (b) Kolbe's reaction
 (c) Kolbe-electrolysis reaction
 (d) None of these
49. Acetic acid will be obtained on oxidation of [J & K 2005]

- (a) Ethanol (b) Propanal
(c) Methanal (d) Glyoxal

Properties of Carboxylic Acids and Their Derivatives

1. Which of the following acids has the smallest dissociation constant [IIT-JEE (Screening) 2002]

- (a) $CH_3CHFCOOH$ (b) FCH_2CH_2COOH
(c) $BrCH_2CH_2COOH$ (d) $CH_3CHBrCOOH$





2. What is obtained, when propene is treated with *N*-bromo succinimide [MP PMT 2003]

- (a) $CH_3 - \underset{\text{Br}}{\text{C}} = CH_2$ (b) $BrCH_2 - CH = CH_2$

- (c) $BrCH_2 - CH = CHBr$ (d) $BrCH_2 - \underset{\text{Br}}{\text{CH}} - CH_2Br$

3. What will be the product, when carboxy phenol, obtained by Reimer Tiemann's process, is deoxidised with *Zn* powder

[MP PMT 2003]



4. The vapour of a carboxylic acid *HA* when passed over MnO_2 at 573 K yields propanone. The acid *HA* is

- (a) Methanoic acid (b) Ethanoic acid
(c) Propanoic acid (d) Butanoic acid

5. Which acid is strongest or Which is most acidic [CPMT 1982, 89; BIT 1992; MP PET 1996; MP PMT/PET 1988; MP PMT 1995, 97; RPMT 1997]

- (a) $Cl_2CH.COOH$ (b) $ClCH_2COOH$
(c) CH_3COOH (d) $Cl_3C.COOH$

6. Ethyl acetate at room temperature is a

- (a) Solid (b) Liquid
(c) Gas (d) Solution

7. Urea is a better fertilizer than ammonium sulphate because

- (a) It has greater percentage of nitrogen
(b) It is more soluble
(c) It is weakly basic
(d) It does not produce acidity in soil


8. The reaction of acetamide with water is an example of

[Kurukshetra CEE 1998; RPMT 2000]

- (a) Alcoholysis (b) Hydrolysis
(c) Ammonolysis (d) Saponification

9. The acid which reduces Fehling solution is [KCET 1998]

- (a) Methanoic acid (b) Ethanoic acid
(c) Butanoic acid (d) Propanoic acid



The above shown polymer is obtained when a carbon compound is allowed to stand. It is a white solid. The polymer is [CBSE PMT 1989]

- (a) Trioxane (b) Formose
(c) Paraformaldehyde (d) Metaldehyde

11. What will happen if $LiAlH_4$ is added to an ester

[CBSE PMT 2000]

- (a) Two units of alcohol are obtained
(b) One unit of alcohol and one unit of acid is obtained
(c) Two units of acids are obtained
(d) None of these

12. When anisole is heated with *HI*, the product is

[CET Pune 1998]

- (a) Phenyl iodide and methyl iodide
(b) Phenol and methanol
(c) Phenyl iodide and methanol
(d) Methyl iodide and phenol

13. When CH_3COOH reacts with $CH_3 - Mg - X$ [BVP 2003]

- (a) CH_3COX is formed (b) Hydrocarbon is formed
(c) Acetone is formed (d) Alcohol is formed

14. Which class of compounds shows *H*-bonding even more than in alcohols

- (a) Phenols (b) Carboxylic acids
(c) Ethers (d) Aldehydes

15. When propanamide reacts with Br_2 and $NaOH$

then which of the following compound is formed [Manipal]

- (a) Ethyl alcohol (b) Propyl alcohol
(c) Propyl amine (d) Ethylamine

16. Hydrolysis of an ester gives a carboxylic acid which on Kolbe's electrolysis yields ethane. The ester is

[EAMCET 1997; Manipal PMT 2001]

- (a) Ethyl methanoate (b) Methyl ethanoate
(c) Propylamine (d) Ethylamine

17. On prolonged heating of ammonium cyanate or urea, we get [DPMT 1982; CPMT 1979; MP PMT 1996]


- (a) N_2 (b) CO_2
(c) Biurette (d) Ammonium carbonate

18. In the Gabriel's phthalimide synthesis, phthalimide is treated first with

- (a) C_2H_5I / KOH (b) Ethanolic *Na*
(c) Ethanol and H_2SO_4 (d) Ether and $LiAlH_4$

19. Which of the following is the strongest acid [NCERT 1984]

1342 Carboxylic acids and Their derivatives

- (a) CH_3COOH (b) $BrCH_2COOH$
 (c) $ClCH_2COOH$ (d) FCH_2COOH
20. Which of the following reduces Tollen's reagent
 [MP PMT 1991]
 (a) Acetic acid (b) Citric acid
 (c) Oxalic acid (d) Formic acid
21. Oxalic acid may be distinguished from tartaric acid by
 (a) Sodium bicarbonate solution
 (b) Ammonical silver nitrate solution
 (c) Litmus paper
 (d) Phenolphthalein
22. The reaction of $HCOOH$ with conc. H_2SO_4 gives
 [DPMT 1982, CPMT 1989; MP PET 1995; AIIMS 2000; Manipal 2001; Pb. CET 2002]
 (a) CO_2 (b) CO
 (c) Oxalic acid (d) Acetic acid
23. Sulphonation of benzoic acid produces mainly [CPMT 1982]
 (a) *o*-sulphobenzoic acid
 (b) *m*-sulphobenzoic acid
 (c) *p*-sulphobenzoic acid
 (d) *o*- and *p*-sulphobenzoic acid
24. Which one is strongest acid [MP PMT 1992]
 (a) CH_2FCOOH (b) $CH_2ClCOOH$
 (c) $CHCl_2COOH$ (d) CHF_2COOH
25. Which does not give silver mirror with ammoniacal $AgNO_3$
 [MP PET 1992]
 (a) $HCHO$ (b) CH_3CHO
 (c) CH_3COOH (d) $HCOOH$
26. $2CH_3COOH \xrightarrow[300^\circ C]{MnO} A$, product 'A' in the reaction is
 [RPMT 2003]
 (a) CH_3CH_2CHO (b) CH_3-CH_2-OH
 (c) CH_3COCH_3 (d) $CH_3-\overset{\parallel}{O}-C-\overset{\parallel}{O}-C-CH_3$
27. Acetic acid is weak acid than sulphuric acid because
 [CPMT 2003]
 (a) It decompose on increasing temperature
 (b) It has less degree of ionisation
 (c) It has $-COOH$ group
 (d) None of these
28. In CH_3COOH and $HCOOH$, $HCOOH$ will be
 [CPMT 1975; DPMT 1982]
 (a) Less acidic (b) Equally acidic
 (c) More acidic (d) None
29. Acetic anhydride reacts with excess of ammonia to form
 [MP PET 1992]
 (a) $2CH_3COONH_4$
 (b) $2CH_3CONH_2$
- (c) $CH_3CONH_2 + CH_3COONH_4$
 (d) $2CH_3COOH$
30. In the following sequence of reactions, what is D

 [UPSEAT 2002]
 (a) Primary amine
 (b) An amide
 (c) Phenyl isocyanate
 (d) A chain lengthened hydrocarbon
31. Hydrolytic reaction of fats with caustic soda is known as
 [MP PMT/PET 1988; AMU 1988; KCET 2000; MP PET 2001]
 (a) Esterification (b) Saponification
 (c) Acetylation (d) Carboxylation
32. In the reaction
 $CH_3COOH \xrightarrow{LiAlH_4} (A) \xrightarrow{I_2+NaOH} (B) \xrightarrow{Ag(Dust)} (C)$
 the final product (C) is
 (a) C_2H_5I (b) C_2H_5OH
 (c) C_2H_2 (d) CH_3COCH_3
33. Reaction of ethyl formate with excess of CH_3MgI followed by hydrolysis gives [IIT (Screening) 1992]
 (a) *n*-propyl alcohol (b) Ethanal
 (c) Propanal (d) Isopropyl alcohol
34. Of the following four reactions, formic and acetic acids differ in which respect
 (a) Replacement of hydrogen by sodium
 (b) Formation of ester with alcohol
 (c) Reduction of Fehling solution
 (d) Blue litmus reaction
35. Formaldehyde and formic acid can be distinguished using
 [AFMC 1993]
 (a) Tollen's reagent (b) Fehling solution
 (c) Ferric chloride (d) Sodium bicarbonate
36. Ester and acetamide are distinguished by [BHU 1996]
 (a) Hydrolysis with strong acids or alkali
 (b) Derivatives of fatty acids
 (c) Both (a) and (b)
 (d) None of these
37. Acetic acid exists as a dimer in benzene solution. This is due to
 [MP PMT 1989; CPMT 1982]
 (a) Condensation
 (b) Presence of $-COOH$ group
 (c) Presence of α -hydrogen
 (d) Hydrogen bonding
38. Which of the following compounds will react with $NaHCO_3$ solution to give sodium salt and carbon dioxide
 [CBSE PMT 1999; BHU 1983, 2002]
 (a) Phenol (b) *n*-hexanol

- (c) Acetic acid (d) Both (a) and (b)
39. Acetic acid dissolved in benzene shows a molecular mass of [MP PET 1993]
 (a) 30 (b) 60
 (c) 120 (d) 240
40. The reaction

$$2\text{CH}_3 - \overset{\text{O}}{\parallel}{\text{C}} - \text{OC}_2\text{H}_5 \xrightarrow{\text{C}_2\text{H}_5\text{ONa}}$$





$$\text{CH}_3 - \overset{\text{O}}{\parallel}{\text{C}} - \text{CH}_2 - \overset{\text{O}}{\parallel}{\text{C}} - \text{OC}_2\text{H}_5 + \text{C}_2\text{H}_5\text{OH}$$
 is called [MP PMT 2003; KCET 1996]
 (a) Etard reaction (b) Perkin's reaction
 (c) Claisen condensation (d) Claisen Schmidt reaction
41. Which is the strongest acid? (pK_a value is given in the bracket) [MP PMT 1997; BHU 2003]
 (a) HCOOH (3.77) (b) $\text{C}_6\text{H}_5\text{COOH}$ (4.22)
 (c) CH_3COOH (4.71) (d) $\text{CH}_3\text{CH}_2\text{COOH}$ (4.88)
42. In the presence of iodine catalyst, chlorine reacts with acetic acid to form [MP PMT 1997]
 (a) $\text{CH}_3 - \overset{\text{O}}{\parallel}{\text{C}} - \text{Cl}$ (b) $\text{CH}_2\text{Cl} - \overset{\text{O}}{\parallel}{\text{C}} - \text{OH}$
 (c) $\text{CH}_3 - \overset{\text{Cl}}{\mid}{\text{C}} - \text{OH}$ (d) $\text{CH}_3 - \overset{\text{O}}{\parallel}{\text{C}} - \text{O} - \text{Cl}$
43. The acid showing salt-like character in aqueous solution is [MP PET/PMT 1998]
 (a) Acetic acid (b) Benzoic acid
 (c) Formic acid (d) α -amino acetic acid
44. $\text{CH}_3\text{COOH} \xrightarrow[\text{P}_2\text{O}_5]{\Delta} \text{X}$. Identify X [JIPMER 2000; CPMT 2003]
 (a) CH_3COCH_3 (b) CH_3CHO
 (c) $(\text{CH}_3\text{CO})_2\text{O}$ (d) CH_4
45. Formic acid [MP PET/PMT 1988]
 (a) Is immiscible with water
 (b) Reduces the ammoniacal silver nitrate
 (c) Is a weak acid nearly three and a half times weaker than acetic acid
 (d) Is prepared by heating potassium hydroxide
46. Given below are some statements concerning formic acid, which of them is true [CPMT 1983]
 (a) It is a weaker acid than acetic acid
 (b) It is a reducing agent
 (c) When its calcium salt is heated, it forms a ketone
 (d) It is an oxidising agent
47. Which decolourises the colour of acidic KMnO_4 [CPMT 1991]
 (a) CH_3COOH (b) $\text{CH}_3\text{CH}_2\text{COOH}$
- (c) $\text{COOH} \cdot \text{COOH}$ (d) $\text{CH}_3\text{COOC}_2\text{H}_5$
48. A colourless water soluble organic liquid decomposes sodium carbonate and liberates carbon dioxide. It produces black precipitate with Tollen's reagent. The liquid is [KCET 1989]
 (a) Acetaldehyde (b) Acetic acid
 (c) Formaldehyde (d) Formic acid
49. The end product B in the sequence of reactions

$$\text{R} - \text{X} \xrightarrow{\text{CN}^-} \text{A} \xrightarrow{\text{NaOH}} \text{B}$$
 is [CPMT 1985]
 (a) An alkane
 (b) A carboxylic acid
 (c) Sodium salt of carboxylic acid
 (d) A ketone
50. $\text{CH}_3\text{CH}_2\text{COOH} \xrightarrow{\text{Cl}_2/\text{Fe}} \text{X} \xrightarrow[\text{KO}_4]{\text{Alcoholic}} \text{Y}$
 Compound Y is [DPMT 1981; JIPMER 2000; AIEEE 2002]
 (a) $\text{CH}_3\text{CH}_2\text{OH}$ (b) $\text{CH}_3\text{CH}_2\text{CN}$
 (c) $\text{CH}_2 = \text{CHCOOH}$ (d) $\text{CH}_3\text{CHClCOOH}$
51. In the precipitation of soap, which can be used instead of NaCl [CPMT 1979]
 (a) Na (b) CH_3COONa
 (c) Na_2SO_4 (d) Sodium silicate
52. Which of the following can possibly be used as analgesic without causing addiction and moon modification [CBSE PMT 1997]
 (a) Morphine
 (b) *N*-acetyl-para-aminophenol
 (c) Drazepam
 (d) Tetrahydrocannabinol
53. Which of the following esters cannot undergo Claisen self condensation
 (a) $\text{CH}_3 - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \text{COOC}_2\text{H}_5$
 (b) $\text{C}_6\text{H}_5\text{COOC}_2\text{H}_5$
 (c) $\text{C}_6\text{H}_5\text{CH}_2\text{COOC}_2\text{H}_5$
 (d) $\text{C}_6\text{H}_{11}\text{CH}_2\text{COOC}_2\text{H}_5$
54. When acetic acid is dissolved in benzene its molecular mass [AFMC 1991]
 (a) Decreases
 (b) Increases
 (c) Either increases or decreases
 (d) Suffers no change
55. Benzoic acid has higher molecular weight in benzene and less in water because
 (a) Water has lower freezing point and higher boiling point than benzene
 (b) It dissociates to a greater extent in benzene than in water
 (c) It associates in water and dissociates in benzene







1344 Carboxylic acids and Their derivatives









- (d) It dissociates in water and associates in benzene
56. What is the main reason for the fact that carboxylic acids can undergo ionization [MNR 1993; Pb. PMT 2004]
- (a) Absence of α - hydrogen
 (b) Resonance stabilisation of the carboxylate ion
 (c) High reactivity of α - hydrogen
 (d) Hydrogen bonding
57. Which of the following compounds will evolve hydrogen on treatment with metal [CPMT 1974]
- (a) C_2H_5OH (b) CH_3COOH
 (c) (a) and (b) both (d) None of these
58. When urea is heated, it forms biurette, alkaline solution of which forms with $CuSO_4$ solution [AFMC 1980]
- (a) Violet colour (b) Red colour
 (c) Green colour (d) Black colour
59. Which of the following would be expected to be most highly ionised in water
- (a) $CH_2ClCH_2CH_2COOH$ (b) $CH_3CHCl.CH_2.COOH$
 (c) $CH_3.CH_2.CCl_2.COOH$ (d) $CH_3.CH_2.CHCl.COOH$
60. Alkaline hydrolysis of esters is known as [CPMT 1986, 88, 93; MNR 1986; MP PET 1993]
- (a) Saponification (b) Hydration
 (c) Esterification (d) Alkalisiation
61. Which of the following undergoes hydrolysis when dissolved in water [CPMT 1989]
- (a) CH_3COONa (b) CH_3CONH_2
 (c) Both (a) and (b) (d) $C_6H_5CH_3$
62. Name the end product in the following series of reactions
- $$CH_3COOH \xrightarrow{NH_3} A \xrightarrow[\text{P}_2\text{O}_5]{\Delta} B$$
- [DPMT 1984]
- (a) CH_4 (b) CH_3OH
 (c) Acetonitrile (d) Ammonium acetate
63. Reduction of carboxylic acids gives
- (a) Alcohol with hydrogen in presence of palladium
 (b) Alcohol with $LiAlH_4$
 (c) Aldehyde with $LiAlH_4$
 (d) Alcohol with $2HI(P)$
64. Which of the following substances when boiled with caustic soda solution will evolve ammonia [BHU 1983]
- (a) Ethylamine (b) Aniline
 (c) Acetamide (d) Acetoxime
65. $CH_2 = CH - (CH_2)_5 COOH \xrightarrow[HBr]{\text{Peroxide}} Z$
- where Z is [CPMT 1996]
- (a) $CH_3 - \underset{\text{Br}}{\text{CH}} - (CH_2)_5 COOH$
 (b) $BrCH_2 - (CH_2)_6 COOH$
- (c) $CH_2 = CH - (CH_2)_5 - CH_2OH$
 (d) C_6H_5COOH
66. $HCOOH$ shows all tests of aldehyde because [CPMT 1996]
- (a) It has one aldehyde group
 (b) It is member of aldehyde
 (c) All acids show tests of aldehyde
 (d) Does not show any test
67. Which one of the following orders of acid strength is correct [CBSE PMT 2003]
- (a) $RCOOH > HC \equiv CH > HOH > ROH$
 (b) $RCOOH > ROH > HOH > HC \equiv CH$
 (c) $RCOOH > HOH > ROH > HC \equiv CH$
 (d) $RCOOH > HOH > HC \equiv CH > ROH$
68. The order of decreasing rate of reaction with ammonia is [AIIMS 1982] [Pb. PMT 1998]
- (a) Anhydrides, esters, ethers
 (b) Anhydrides, ethers, esters
 (c) Ethers, anhydrides, esters
 (d) Esters, ethers, anhydrides
69. Oxidation of toluene with CrO_3 in the presence of $(CH_3CO)O_2$ gives a product 'A' which on treatment with aqueous $NaOH$ produces
- (a) C_6H_5CHO (b) $(C_6H_5CO)_2O$
 (c) C_6H_5COONa (d) 2, 4-diacetyl toluene
70. $CH_3COOCH_3 + \text{excess } PhMgBr \rightarrow \text{product} \xrightarrow{H^+} X$
- The product X is [Orissa JEE 2005]
- (a) 1, 1-diphenylethanol
 (b) 1, 1-diphenylmethanol
 (c) Methyl phenylethanol
 (d) Methyl phenylketone
71. Which of the following is most acidic [MP PMT 1995]
- (a) Formic acid (b) Chloroacetic acid
 (c) Propionic acid (d) Acetic acid
72. Urea on slow heating gives
- (a) $NH_2CON.HNO_2$ (b) $NH_2CONHCONH_2$
 (c) $HCNO$ (d) $NH_2CONH_2.HNO_3$
73. The principal organic product formed in the following reaction is
- $$CH_2 = CH(CH_2)_8 COOH + HBr \xrightarrow{\text{peroxide}}$$
- [Pb. PMT 1998]
- (a) $CH_3CHBr(CH_2)_8 COOH$
 (b) $CH_2 = CH(CH_2)_8 COBr$
 (c) $CH_2BrCH_2(CH_2)_8 COOH$

- (d) $CH_2 = CH(CH_2)_7CHBrCOOH$
74. Which one of the following compound gives aspirin on reacting with acetic anhydride in presence of H_2SO_4 [EAMCET 2003]
- (a)  (b) 
- (c)  (d) 
75. An acyl halide is formed when PCl_5 reacts with an [CBSE PMT 1994; AIIMS 1998; CBSE PMT 2002]
- (a) Acid (b) Alcohol
(c) Amide (d) Ester
76. Which one of the following orders is wrong with respect to the property indicated [CBSE PMT 1994]
- (a) Formic acid > acetic acid > propanoic acid (acid strength)
(b) Fluoroacetic acid > chloroacetic acid > bromoacetic acid (acid strength)
(c) Benzoic acid > phenol > cyclohexanol (acid strength)
(d) Aniline > cyclohexylamine > benzamide (basic strength)
77. A fruity smell is produced by the reaction of C_2H_5OH with [AFMC 2000]
- (a) PCl_5 (b) CH_3COCH_3
(c) CH_3COOH (d) None of these
78. Which of the following orders of relative strengths of acids is correct
- (a) $FCH_2COOH > ClCH_2COOH > BrCH_2COOH$
(b) $ClCH_2COOH > BrCH_2COOH > FCH_2COOH$
(c) $BrCH_2COOH > ClCH_2COOH > FCH_2COOH$
(d) $ClCH_2COOH > FCH_2COOH > BrCH_2COOH$
79. When acetamide is treated with $NaOBr$, the product formed is [Haryana CEET 2000]
- (a) CH_3CN (b) $CH_3CH_2NH_2$
(c) CH_3NH_2 (d) None of the above
80. The fatty acid which shows reducing property is [Kerala CET 2000]
- (a) Acetic acid (b) Ethanoic Acid
(c) Oxalic acid (d) Formic acid
81. The reagent that can be used to distinguish between methanoic acid and ethanoic acid is [Kerala CET 2001, 02]
- (a) Ammoniacal silver nitrate solution
(b) Neutral ferric Chloride solution
(c) Sodium carbonate solution
(d) Phenolphthalein
82. Hydrolysis of an ester gives acid A and alcohol B. A reduces Fehling solution and oxidation of B gives A. The ester is [MP PMT 1999]
- (a) Methyl formate (b) Ethyl formate
(c) Methyl acetate (d) Ethyl acetate
83. Order of reactivity is [RPMT 2003]
- (a) $R-\overset{O}{\parallel}C-X > RCONH_2 > RCOOCOR > RCOOR$
(b) $RCOX > RCOOCOR > RCOOR > RCONH_2$
(c) $RCOOR > RCONH_2 > RCOX > RCOOCOR$
(d) $RCOOCOR > RCOOR > RCOX > RCONH_2$
84. Right order of acidic strength is [RPMT 2003]
- (a) $CH_2ClCOOH > HCOOH > C_2H_5COOH > CH_3COOH$
(b) $CH_2ClCOOH > HCOOH > CH_3COOH > C_2H_5COOH$
(c) $C_2H_5COOH > CH_3COOH > HCOOH > CH_2ClCOOH$
(d) $HCOOH > CH_2ClCOOH > CH_3COOH > C_2H_5COOH$
85. Saponification of ethyl benzoate with caustic soda as alkali gives [Kerala (Med.) 2001]
- (a) Benzyl alcohol and ethanoic acid
(b) Sodium benzoate and ethanol
(c) Benzoic acid and sodium ethoxide
(d) Phenol and ethanoic acid
(e) Sodium benzoate and ethanoic acid
86. Lactic acid on oxidation by alkaline potassium permanganate gives [Tamil Nadu CET 2002]
- (a) Tartaric acid (b) Pyruvic acid
(c) Cinnamic acid (d) Propionic acid
87. $RCOOH \longrightarrow RCH_2OH$ [CPMT 2000]
- This mode of reduction of an acid to alcohol can be affected only by [CBSE PMT 1989]
- (a) Zn/HCl
(b) Na -alcohol
(c) Aluminium isopropoxide and isopropyl alcohol
(d) $LiAlH_4$
88. Which one of the following compounds forms a red coloured solution on treatment with neutral $FeCl_3$ solution [EAMCET 2003]
- (a) CH_3COCH_3 (b) CH_3OCH_3
(c) CH_3CH_2OH (d) CH_3COOH
89. Urea can be tested by [UPSEAT 1999; BVP 2003]
- (a) Benedict test (b) Mullicken test
(c) Ninhydrin (d) Biuret test
90. What are the organic products formed in the following reaction [IIT 1995]
- $$C_6H_5-COO-CH_3 \xrightarrow[2.H_2O]{1.LiAlH_4}$$
- (a) C_6H_5-COOH and CH_4



1346 Carboxylic acids and Their derivatives

- (b) $C_6H_5-CH_2-OH$ and CH_4
 (c) $C_6H_5-CH_3$ and CH_3-OH
 (d) $C_6H_5-CH_2-OH$ and CH_3-OH
91. Reaction between an acid and alcohol will give
 [Roorkee 1995]
 (a) Higher C containing acid (b) Secondary alcohol
 (c) Alkane (d) Ester
92. Benzoic acid gives benzene on being heated with X and phenol gives benzene on being heated with Y. Therefore X and Y are respectively [CBSE PMT 1992]
 (a) Sodalime and copper (b) Zn dust and NaOH
 (c) Zn dust and sodalime (d) Sodalime and zinc dust
93. The product obtained when acetic acid is treated with phosphorus trichloride is [CPMT 1989, 93, 94; RPMT 1997; AIIMS 1998; EAMCET 1998]
 (a) $CH_3COOPCl_3$ (b) CH_3COOCl
 (c) CH_3COCl (d) $ClCH_2COOH$
94. Acetyl chloride is reduced with $LiAlH_4$ the product formed is
 (a) Methyl alcohol (b) Ethyl alcohol
 (c) Acetaldehyde (d) Acetone
95. In the preparation of an ester, the commonly used dehydrating agent is [KCET 1992]
 (a) Phosphorus pentoxide
 (b) Anhydrous calcium carbide
 (c) Anhydrous aluminium chloride
 (d) Concentrated sulphuric acid
96. In the esterification reaction of alcohols [KCET 1984]
 (a) OH^- is replaced by C_6H_5OH
 (b) H^+ is replaced by sodium metal
 (c) OH^- is replaced by chlorine
 (d) OH^- is replaced by CH_3COO^- group
97. Lower carboxylic acids are soluble in water due to [MP PET 1999]
 (a) Low molecular weight (b) Hydrogen bonding
 (c) Dissociation into ions (d) Easy hydrolysis
98. Acetamide reacts with P_2O_5 (phosphorus pentoxide) to give
 (a) Methyl cyanide (b) Methyl cyanate
 (c) Ethyl cyanide (d) Ethyl isocyanate
99. The reaction
 $CH_3COOH + Cl_2 \xrightarrow{P} ClCH_2COOH + HCl$ is called [NSE 2001; MP PET 2003]
 (a) Hell-Volhard-Zelinsky reaction
 (b) Birch reaction
 (c) Rosenmund reaction
 (d) Hunsdiecker reaction
100. An aqueous solution of urea [CPMT 1983]
 (a) Is neutral
 (b) Is acidic
 (c) Is basic
 (d) Can act as an acid and a base
101. Nitration of benzoic acid gives [MP PMT 1997]
 (a) 3-nitrobenzoic acid (b) 2-nitrobenzoic acid
 (c) 2, 3-dinitrobenzoic acid (d) 2, 4-dinitrobenzoic acid
102. The reagent used for converting ethanoic acid to ethanol is [KCET 1996; EAMCET 1998]
 (a) $LiAlH_4$ (b) $KMnO_4$
 (c) PCl_3 (d) $K_2Cr_2O_7 / H^+$
103. Which one of the following has the maximum acid strength [NCERT 1983]
 (a) o-nitrobenzoic acid (b) m-nitrobenzoic acid
 (c) p-nitrobenzoic acid (d) p-nitrophenol
104. When benzoic acid is treated with PCl_5 at $100^\circ C$, it gives [Orissa JEE 2003]
 (a) Benzoyl chloride (b) o-chlorobenzoic acid
 (c) p-chlorobenzoic acid (d) Benzyl chloride
105. Oxalic acid on being heated upto $90^\circ C$ with conc. H_2SO_4 forms [SCRA 1990] [AFMC 1989; MP PET 1994; MP PMT 1989]
 (a) $HCOOH + CO_2$ (b) $CO_2 + H_2O$
 (c) $CO_2 + CO + H_2O$ (d) $HCOOH + CO$
106. Benzoic acid is less acidic than salicylic acid because of [Bihar MEE 1997]
 (a) Hydrogen bond (b) Inductive effect
 (c) Resonance (d) All of these
 (e) None of these
107. Lactic acid on heating with conc. H_2SO_4 gives [MP PET 1996]
 (a) Acetic acid (b) Propionic acid
 (c) Acrylic acid (d) Formic acid
108. Acetamide is [MP PET 1990; RPMT 1999]
 (a) Acidic (b) Basic
 (c) Neutral (d) Amphoteric
109. Silver benzoate reacts with bromine to form [KCET 1996]
 (a)  [AFMC 1996]
 (b) 
 (c) 
 (d) 
110. Acetic anhydride reacts with diethyl ether in presence of anhydrous $AlCl_3$ to form [MP PMT 1992]
 (a) Ethyl acetate (b) Methyl propionate
 (c) Methyl acetate (d) Propionic acid
111. Treatment of benzoic acid with $Cl_2 / FeCl_3$ will give [KCET 1998; CET Pune 1998]
 (a) p-chlorobenzoic acid (b) o-chlorobenzoic acid
 (c) 2, 4-dichlorobenzoic acid (d) m-chlorobenzoic acid

112. Hinsberg's reagent is [MP PMT 2003]
- (a)  (b) 
 (c)  (d) 
113. Which of the following is the correct order of increasing strengths of carboxylic acids
- (a) $CH_2FCOOH < CH_3COOH < CH_2ClCOOH < CCl_3COOH$
 (b) $CH_3COOH < CH_2ClCOOH < CH_2FCOOH < CCl_3COOH$
 (c) $CH_2ClCOOH < CH_2FCOOH < CCl_3COOH < CH_3COOH$
 (d) $CCl_3COOH < CH_2ClCOOH < CH_2FCOOH < CH_3COOH$
114. The weakest acid among the following is [CPMT 1976, 82, 89; BHU 1982; CBSE PMT 1991; MP PMT 1989; Roorkee 1992; RPET 1999]
- (a) CH_3COOH (b) $Cl_2CHCOOH$
 (c) $ClCH_2COOH$ (d) Cl_3CCOOH
115. Consider the acidity of the carboxylic acids
- (a) $PhCOOH$ (b) $o-NO_2C_6H_4COOH$
 (c) $p-NO_2C_6H_4COOH$ (d) $m-NO_2C_6H_4COOH$
- Which of the following order is correct? [AIIEEE 2004]
- (a) $b > d > a > c$ (b) $b > d > c > a$
 (c) $a > b > c > d$ (d) $b > c > d > a$
116. On mixing ethyl acetate with aqueous sodium chloride, the composition of the resultant solution is [AIIEEE 2004]
- (a) $CH_3COCl + C_2H_5OH + NaOH$
 (b) $CH_3COONa + C_2H_5OH$
 (c) $CH_3COOC_2H_5 + NaCl$
 (d) $CH_3Cl + C_2H_5COONa$
117. *o*-Toluic acid on reaction with $Br_2 + Fe$, gives [AIIMS 2004]
- (a)  (b) 
 (c)  (d) 
118. The reaction of an ester $RCOOR'$ with an alcohol $R''OH$ in the presence of an acid gives [Kerala PMT 2004]
- (a) $RCOOH$ (b) $R'COOH$
 (c) $R''COOR$ (d) $RCOOR''$
 (e) $R'COOR''$
119. $RCOOH$ on treatment with PCl_5 and KCN , is subjected to hydrolysis followed by Clemmensen's reduction, the product obtained is [Kerala PMT 2004]
- (a) RCH_2COCl (b) RCH_2COOH
 (c) $RCOCl$ (d) RCN
 (e) $R-OH$
120. The reagent which does not give acid chloride on treating with a carboxylic acid is [KCET 2004]
- (a) PCl_5 (b) Cl_2
 (c) $SOCl_2$ (d) PCl_3
121. An organic compound is boiled with alcoholic potash. The product is cooled and acidified with HCl . A white solid separates out. The starting compound may be [KCET 2004]
- (a) Ethyl benzoate (b) Ethyl formate
 (c) Ethyl acetate (d) Methyl acetate
122. The OH group of an alcohol or the $-COOH$ group of a carboxylic acid can be replaced by $-Cl$ using [CBSE PMT 2004]
- (a) Chlorine
 (b) Hydrochloric acid
 (c) Phosphorus pentachloride
 (d) Hypochlorous acid
123. Which of the following is most acidic [MP PET 2004]
- (a) Picric acid (b) *p*-nitrophenol
 (c) *m*-nitrophenol (d) *o-p* dinitrophenol
124. Benedict's solution is not reduced by [CPMT 2004]
- (a) Formaldehyde (b) Acetaldehyde
 (c) Glucose (d) Acetic anhydride
125. CH_3COOH is reacted with $CH \equiv CH$ in presence of Hg^{++} , the product is [DPMT 2004; BHU 1998]
- (a) $CH_3(OOCCH_3)$ (b) CH_3
 $|$ $|$
 $CH_2(OOCH_3)$ $CH_2-(OOC-CH_3)$
 (c) CH_3 (d) None of these
 $|$
 $CH(OOC-CH_3)_2$
126. Acetic acid reacts with PCl_5 to form [Pb. CET 2001]
- (a) CH_3COCl (b) $CHCl_2COOH$
 (c) $CH_2ClCOOH$ (d) CH_3COOCl
127. $CH_3COOC_2H_5$ with excess of C_2H_5MgBr and hydrolysis gives [MH CET 2004]
- (a) $CH_3-C=O$ (b) CH_3-C-OH
 $|$ $|$
 C_2H_5 C_2H_5
 (c) $CH_3-C=O$ (d) $CH_3-C=O$
 $|$ $|$
 CH_3 CH_3
128. Urea upon hydrolysis yields [Pb. CET 2001]
- (a) Acetamide (b) Carbonic acid
 (c) Ammonium hydroxide (d) NO_2
129. $CH_3CHO \xrightarrow{HCN} A \xrightarrow{HOH} B$. The product B is [Pb. CET 2003]
- (a) Malonic acid (b) Glycolic acid
 (c) Lactic acid (d) Malic acid

1348 Carboxylic acids and Their derivatives

- 130.** What is the % of acetic acid present in vinegar?
[AFMC – 2004; MH CET 2003; CPMT 1974, 75]
(a) 6–10% (b) 70–80%
(c) 7–8% (d) 90–100%
- 131.** Fruity smell is given by [MH CET 2004]
(a) Esters (b) Alcohols
(c) Chloroform (d) Acid anhydrides
- 132.** Lactic acid molecule has [MH CET 2004]
(a) One chiral carbon atom
(b) Two chiral carbon atoms
(c) No chiral carbon atom
(d) As asymmetric molecule
- 133.** 4-methyl benzene sulphonic acid reacts with sodium acetate to give [IIT-JEE (Screening) 2005]
- (a) ; CH_3COOH

(b) ; SO_3
- (c) ; SO_3


(d) ; $NaOH$
- 134.** In a set reactions acid yielded a product *D*
 $CH_3COOH \xrightarrow{SOCl_2} A \xrightarrow[\text{Anhr. AlCl}_3]{\text{Benzene}} B \xrightarrow{HCN} C \xrightarrow{HOH} D$.
[CBSE PMT 2005]
- (a)

(b)
- (c)

(d)
- 135.** $C_6H_5CHO + HCN \rightarrow C_6H_5 - \overset{\overset{H}{|}}{\underset{\underset{OH}{|}}{C}} - CN$
The product would be [Pb. PMT 1998]
(a) A racemate
(b) Optically active
(c) A meso compound
(d) A mixture of diastereomers
- 136.** What happens when 2-hydroxy benzoic acid is distilled with zinc dust, it gives [MP PET/PMT 1998]
(a) Phenol (b) Benzoic acid
(c) Benzaldehyde (d) A polymeric compound
- 137.** $CH_3CO_2C_2H_5$ on reaction with sodium ethoxide in ethanol gives *A*, which on heating in the presence of acid gives *B* compound *B* is [AIIMS 2005]
(a) CH_3COCH_2COOH (b) CH_3COCH_3
(c) (d) $CH_2 = C \begin{matrix} \swarrow OC_2H_5 \\ \searrow OC_2H_5 \end{matrix}$
- 138.** $C_6H_5CONHCH_3$ can be converted into $C_6H_5CH_2NHCH_3$ by [AIIMS 2005]
(a) $NaBH_4$ (b) $H_2 - Pd / C$
(c) $LiAlH_4$ (d) $Zn - Hg / HCl$
- 139.** Among the following acids which has the lowest *pKa* value [AIEEE 2005]
(a) CH_3COOH (b) $HCOOH$
(c) $(CH_3)_2CH - COOH$ (d) CH_3CH_2COOH
- 140.** *X* is heated with soda lime and gives ethane. *X* is [AFMC 2005]
(a) Ethanoic acid (b) Methanoic acid
(c) Propanoic acid (d) Either (a) or (c)
- 141.** Which of the following is an amphoteric acid [KCET 2005]
(a) Glycine (b) Salicylic acid
(c) Benzoic acid (d) Citric acid
- 142.** Colouration of Br_2 / CCl_4 will be discharged by [Orissa JEE 2005]
(a) Cinnamic acid (b) Benzoic acid
(c) *o*-phthalic acid (d) Acetophenone
- 143.** Order of hydrolysis for the following
(I) $RCOCl$ (II) $RCOOR$
(III) $RCOONa$ (IV) $(RCO)_2O$
(a) I>IV>II>III (b) I>II>III>IV
(c) I>III>II>IV (d) IV>III>II>I
- 144.** If the enolate ion combines with carbonyl group of ester, we get [DPMT 2005]
(a) Aldol (b) α, β -unsaturated ester
(c) β -keto aldehyde (d) Acid
- 145.** Which of the following compounds will react with $NaHCO_3$ solution to give sodium salt and carbon dioxide [DPMT 2005]
(a) Acetic acid (b) *n*-hexanol
(c) Phenol (d) Both (a) and (c)
- 146.** A carboxylic acid is converted into its anhydride using [J & K 2005]
(a) Thionyl chloride (b) Sulphur chloride
(c) Sulphuric acid (d) Phosphorus pentoxide

Uses of Carboxylic Acids and Their Derivatives

1. The following compound is used as



[KCET 1996]





- (a) An anti-inflammatory agent
 (b) Analgesic
 (c) Hypnotic
 (d) Antiseptic
2. To which of the following groups does soap belongs
 [NCERT 1979; RPET 2000]
 (a) Esters
 (b) Amines
 (c) Salts of organic higher fatty acids
 (d) Aldehydes
3. Aspirin is an acetylation product of [CBSE PMT 1998]
 (a) *o*-hydroxybenzoic acid (b) *o*-dihydroxybenzene
 (c) *m*-hydroxybenzoic acid (d) *p*-dihydroxybenzene
4. Which one is used as a food preservative
 [MP PET 1989; KCET 1999]
 (a) Sodium acetate (b) Sodium propionate
 (c) Sodium benzoate (d) Sodium oxalate
5. What makes a lemon sour
 [CPMT 1972; CBSE PMT 1991; RPET 1999]
 (a) Tartaric acid (b) Oxalic acid
 (c) Citric acid (d) Hydrochloric acid
6. The reagent used for protection of amino group during the nitration of aniline is [JIPMER 1997]
 (a) $SOCl_2$ / Pyridine (b) PCl_5
 (c) Acetic acid (d) Acetic anhydride
- (c) $C_6H_5CHOHCH_2OH$ (d) $C_6H_{11}CH_2COOH$
3. Which of the following has the most acidic proton
 [Roorkee Qualifying 1998]
 (a) CH_3COCH_3 (b) $(CH_3)_2C=CH_2$
 (c) $CH_3COCH_2COCH_3$ (d) $(CH_3CO)_3CH$
4. In the anion $HCOO^-$ the two carbon-oxygen bonds are found to be of equal length. What is the reason for it
 (a) Electronic orbitals of carbon atom are hybridised
 (b) The $C=O$ bond is weaker than the $C-O$ bond
 (c) The anion $HCOO^-$ has two resonating structures
 (d) The anion is obtained by removal of a proton from the acid molecule
5. An organic compound of molecular formula $C_4H_{10}O$ does not react with sodium. With excess of HI , it gives only one type of alkyl halide. The compound is [SCRA 2001]
 (a) Ethoxyethane (b) 2-Methoxypropane
 (c) 1-Methoxypropane (d) 1-Butanol
6. When $CH_2=CH-COOH$ is reduced with $LiAlH_4$, the compound obtained will be [AIEEE 2003]
 (a) CH_3-CH_2-COOH (b) $CH_2=CH-CH_2OH$
 (c) $CH_3-CH_2-CH_2OH$ (d) CH_3-CH_2-CHO
7. In a set of the given reactions, acetic acid yielded a product C

$$CH_3COOH + PCl_5 \rightarrow A \xrightarrow[\text{anh. } AlCl_3]{C_6H_6} B \xrightarrow[\text{ether}]{C_5H_5MgBr} C$$

 Product C would be [CBSE PMT 2003]
 (a) $CH_3-\overset{C_2H_5}{C}(OH)C_6H_5$ (b) $CH_3CH(OH)C_2H_5$
 (c) $CH_3COC_6H_5$ (d) $CH_3CH(OH)C_6H_5$

Critical Thinking

Objective Questions

1. Ethyl ester $\xrightarrow[\text{excess}]{CH_3MgBr} P$. The product P will be
 [IIT-JEE 2003]
- (a) 
- (b) 
- (c) 
- (d) 
2. Hydrogenation of $C_6H_5CHOH-COOH$ over $Rh-Al_2O_3$ catalyst in methanol gives
 [Roorkee Qualifying 1998]
 (a) $C_6H_5CH_2COOH$ (b) $C_6H_{11}CHOHCOOH$
8. Carboxylic acids are more acidic than phenol and alcohol because of [Tamil Nadu CET 2001]
 (a) Intermolecular hydrogen bonding
 (b) Formation of dimers
 (c) Highly acidic hydrogen
 (d) Resonance stabilization of their conjugate base
9. $R-CH_2-CH_2OH$ can be converted into RCH_2CH_2COOH . The correct sequence of the reagents is [AIIMS 1997]
 (a) PBr_3, KCN, H_3O^+ (b) PBr_3, KCN, H_2
 (c) HCN, PBr_3, H^+ (d) KCN, H^+
10. When propionic acid is treated with aqueous sodium bicarbonate CO_2 is liberated. The 'C' of CO_2 comes from [IIT-JEE (Screening) 1999]
 (a) Methyl group (b) Carboxylic acid group
 (c) Methylene group (d) Bicarbonate

1350 Carboxylic acids and Their derivatives

11. Benzoyl chloride is prepared from benzoic acid by
[IIT-JEE (Screening) 2000]
(a) Cl_2, hv (b) SO_2Cl_2
(c) $SOCl_2$ (d) Cl_2, H_2O
12. Identify the correct order of boiling points of the following compounds
 $CH_3CH_2CH_2CH_2OH$, $CH_3CH_2CH_2CHO$,
(1) (2)
 $CH_3CH_2CH_2COOH$ [IIT-JEE (Screening) 2002]
(3)
(a) $1 > 2 > 3$ (b) $3 > 1 > 2$
(c) $1 > 3 > 2$ (d) $3 > 2 > 1$
13. The compound not soluble in acetic acid is
[UPSEAT 2003; IIT-JEE 1986]
(a) $CaCO_3$ (b) CaO
(c) CaC_2O_4 (d) $Ca(OH)_2$
14. The *ortho/para* directing group among the following is
[AIIMS 2003]
(a) $COOH$ (b) CN
(c) $COCH_3$ (d) $NHCOCH_3$
15. Iodoform test is not given by [BHU 1995]
(a) Acetone (b) Ethyl alcohol
(c) Acetic acid (d) None of these
16. How will you convert butan-2-one to propanoic acid
[IIT-JEE (Screening) 2005]
(a) Tollen's reagent (b) Fehling's solution
(c) $NaOH / I_2 / H^+$ (d) $NaOH / NaI / H^+$
17. Which of the acids cannot be prepared by Grignard reagent
[MH CET 2004]
(a) Acetic acid (b) Succinic acid
(c) Formic acid (d) All of these
- Reason : Electron withdrawing substituents decrease the activity.
3. Assertion : First four aliphatic monocarboxylic acids are colourless.
Reason : Carboxylic acids with more than five carbon atoms are insoluble in water.
4. Assertion : Carboxylic acids do not give characteristic reactions of carbonyl group.
Reason : Carboxylic acids exist as cyclic dimers in solid, liquid and even in vapour state.
5. Assertion : Pure acetic acid is converted into ice like solid called glacial acetic acid.
Reason : Acetic acid is stronger than $HCOOH$.
6. Assertion : The second dissociation constant of maleic acid is greater than fumaric acid.
Reason : Higher the dissociation constant of acid more is acidic character.
7. Assertion : Lower acids on reacting with strong electropositive metals give effervescences of H_2 .
Reason : $MeCOOC_4H_9$ hydrolyses rapidly than $MeCOOCH_3$.
8. Assertion : Melting point of carboxylic acids shows a regular pattern.
Reason : Carboxylic acids are reduced to alkanes on reduction with HI in presence of red phosphorus.
9. Assertion : Electron withdrawing groups decrease the acidity of carboxylic acids.
Reason : Substituents affect the stability of the conjugate base and acidity of carboxylic acids.
10. Assertion : Fluoroacetic acid is stronger acid than bromoacetic acid.
Reason : Acidity depends upon the electron withdrawing effects of the fluorine and chlorine.
11. Assertion : Aminoacetic acid is less acidic than acetic acid.
Reason : Amino group is electron donating in nature.
12. Assertion : Carboxylic acids have higher boiling points than alkanes.
Reason : Carboxylic acids are resonance hybrids.
13. Assertion : Both formic acid and oxalic acid decolourize $KMnO_4$ solution.
1. Assertion : Carboxylic acid exist as dimer.
Reason : Carboxylic acid shows hydrogen bonding.
2. Assertion : Trichloroacetic acid is stronger than acetic acid.

Assertion & Reason

For AIIMS Aspirants

Read the assertion and reason carefully to mark the correct option out of the options given below :

- (a) If both assertion and reason are true and the reason is the correct explanation of the assertion.
(b) If both assertion and reason are true but reason is not the correct explanation of the assertion.
(c) If assertion is true but reason is false.
(d) If the assertion and reason both are false.
(e) If assertion is false but reason is true.

1. Assertion : Carboxylic acid exist as dimer.
Reason : Carboxylic acid shows hydrogen bonding.
2. Assertion : Trichloroacetic acid is stronger than acetic acid.

Reason : Both are easily oxidised to CO_2 and H_2O .

14. Assertion : Esters which contain α -hydrogens undergo Claisen condensation.

Reason : $LiAlH_4$ reduction of esters gives acids.

Answers

General Introduction of Carboxylic Acids and Their Derivatives

1	d	2	d	3	c	4	d	5	a
6	c	7	d	8	c	9	c	10	d
11	a	12	c	13	d	14	d	15	b
16	c	17	d	18	d	19	c	20	b,d
21	a	22	a	23	c	24	b	25	b
26	b	27	c	28	d	29	a	30	a
31	c	32	b	33	d	34	c	35	a
36	d	37	a	38	d				

Preparation of Carboxylic Acids and Their Derivatives

1	d	2	a	3	a	4	c	5	c
6	a	7	a	8	c	9	a	10	b
11	d	12	a	13	b	14	d	15	a
16	c	17	c	18	a	19	c	20	b
21	a	22	b	23	b	24	a	25	c
26	d	27	c	28	b	29	a	30	b
31	c	32	a	33	b	34	c	35	a
36	b	37	b	38	c	39	b	40	a
41	d	42	b	43	b	44	a	45	b
46	d	47	c	48	a	49	a		

Properties of Carboxylic Acids and Their Derivatives

1	c	2	b	3	d	4	b	5	d
6	b	7	ad	8	b	9	a	10	a
11	a	12	d	13	b	14	b	15	d
16	b	17	c	18	a	19	d	20	d

21	b	22	b	23	b	24	d	25	c
26	c	27	b	28	c	29	c	30	c
31	b	32	c	33	d	34	c	35	d
36	c	37	d	38	c	39	c	40	c
41	a	42	b	43	d	44	c	45	b
46	b	47	c	48	d	49	c	50	c
51	c	52	b	53	b	54	b	55	d
56	b	57	c	58	a	59	c	60	a
61	c	62	c	63	b	64	c	65	b
66	a	67	c	68	b	69	c	70	a
71	b	72	b	73	c	74	b	75	a
76	d	77	c	78	a	79	c	80	d
81	a	82	a	83	b	84	b	85	b
86	b	87	d	88	d	89	d	90	d
91	d	92	d	93	c	94	b	95	d
96	d	97	b	98	a	99	a	100	a
101	a	102	a	103	a	104	a	105	c
106	a	107	c	108	d	109	d	110	a
111	d	112	b	113	b	114	a	115	d
116	c	117	c	118	d	119	b	120	b
121	a	122	c	123	a	124	d	125	c
126	a	127	b	128	b	129	c	130	a
131	a	132	a	133	a	134	a	135	b
136	b	137	c	138	d	139	b	140	c
141	a	142	a	143	a	144	c	145	a
146	d								

Uses of Carboxylic Acids and Their Derivatives

1	b	2	c	3	a	4	c	5	c
6	d								

Critical Thinking Questions

1	a	2	b	3	d	4	c	5	a
6	b	7	a	8	d	9	a	10	d
11	c	12	b	13	c	14	d	15	c
16	c	17	c						


Assertion & Reason

1	a	2	c	3	c	4	b	5	c
6	e	7	c	8	e	9	e	10	a
11	c	12	b	13	a	14	c		

AS Answers and Solutions

General Introduction of Carboxylic Acids and Their Derivatives







1. (d) Methyl salicylate occurs in natural essential oils like winter green.
2. (d) Tartaric acid have the chiral carbon (*) atom. So it is optically active.
















3. (c) Formula of palmitic acid is $C_{15}H_{31}COOH$.




Preparation of Carboxylic Acids and Their Derivatives












10. (b) Reimer-Tiemann reaction involves a carbene intermediate.




23. (b) Acetic acid freezes at 16.6°C while water freezes at 0°C. So glacial acetic acid is obtained by crystallizing, separating and melting acetic acid.



It is called esterification reaction.




37. (b) $RCOOAg + R'I \rightarrow RCOOR' + AgI$
ester
38. (c) When Cl_2 or Br_2 is react with carboxylic acid in the presence of red phosphorus then α -hydrogen of carboxylic acid is replaced by Cl_2 or Br_2




This reaction is known as Hell Volhard Zelinsky reaction.


39. (b) Tertiary alcohol are not oxidised easily but on drastic conditions, these oxidise to give first ketone and then acid by losing one carbon at each step




40. (a) When succinic acid is heated it forms. Succinic anhydride




41. (d) Treatment of sodium salt of phenol with CO_2 under pressure bring about substitution of the carbonyl group $-COOH$, for the hydrogen of the ring. This is called as Kolbe's reaction




Phenol Sodium salt of phenol Sodium salicylate




42. (b) When an acyl halide is heated with acid salt, anhydrides are formed



43. (b) CO_2 adds to Grignard's reagent to yield acids.




44. (a) Amide, on treating with HNO_2 , give acids.



45. (b) Aldehydes are easily oxidised to carboxylic acids on treatment with common oxidising


agents like nitric acid, potassium permanganate and dichromate etc.



Toluene Benzoic acid

47. (c) This is an example of Perkin's reaction.

Therefore, (X) is Acetic anhydride.



The above given reaction is known as Reimer-Tiemann reaction.

49. (a) $C_2H_5OH \xrightarrow{[O]} CH_3COOH$


Properties of carboxylic acids and Their derivatives

1. (c) $BrCH_2CH_2COOH$ is least acidic or has less K_a i.e., dissociation constant. It is (a) due to lesser $-I$ effect of Br than F and (b) Br atom further away from $-COOH$ group.


2. (b) $CH_3 - CH = CH_2 + \text{N-Br} \rightarrow$

Propene

N-Bromosuccinimide



allyl bromide


3. (d) 

Salicylic acid

Benzoic acid


4. (b) $2CH_3COOH \xrightarrow[573 K]{MnO} CH_3COCH_3 + CO_2 + H_2O$

5. (d) Presence of $-I$ effect chlorine atom increases the acidic nature by withdrawing electrons








8. (b) The reaction of acetamide with water is an example of hydrolysis.


9. (a) Methanoic acid resemble with aldehyde due to its structure. So it reduce fehling's reagent.



1354 Carboxylic acids and Their derivatives

11. (a) $R-\overset{\overset{O}{\parallel}}{C}-R' \xrightarrow{LiAlH_4} R-CH_2OH + R'OH$
Two units of alcohols
12. (d)  Anisole + HI → CH₃I + Phenol
Methyl Iodide
13. (b) $CH_3COOH + CH_3-Mg-X \rightarrow CH_3-CH_3$
14. (b) Forms H-bonding by means two highly electronegative atoms present in it.
15. (d) $CH_3CH_2CONH_2 \xrightarrow{Br_2/KOH} CH_3CH_2NH_2$
Propionamide Hofmann bromamide reaction Ethyl amine
16. (b) $CH_3COOCH_3 \xrightarrow[-CH_3OH]{NaOH} CH_3COONa$
Methylethanoate Sodium acetate
 $CH_3-CH_3 \xleftarrow[-CO_2, -NaOH, -H_2]{Kolbe's\ electrolysis}$
17. (c) $NH_4CNO \xrightarrow{\Delta} NH_2-CO-NH_2$
Ammonium cyanate Urea
 $NH_2-CO-NH_2 + H-NH-CO-NH_2 \xrightarrow[-NH_3]{\Delta}$
Urea
 $NH_2-CO-NH-CO-NH_2$
Biuret
19. (d) $F-CH_2-COOH > Cl-CH_2-COOH >$
 $Br-CH_2-COOH > CH_3COOH$
20. (d) Formic acid resemble with aldehyde due to its structure so it reduce Tollen's reagent.

Aldehydic group
22. (b) $HCOOH \xrightarrow{conc. H_2SO_4} CO + H_2O$
23. (b)  + conc. H₂SO₄ →  + H₂O
m-sulpho benzoic acid
24. (d) CHF₂-COOH. Difluoroacetic acid is strongest because presence of two F atoms increases its acidic nature.
25. (c) CH₃COOH does not give silver mirror test.
26. (c) $2CH_3COOH \xrightarrow[300^\circ C]{MnO} CH_3COCH_3 + CO_2 + H_2O$
Acetone
27. (b) CH₃COOH is slightly ionised than H₂SO₄.
28. (c) Presence of methyl group decreases the acidic character of acetic acid due to positive inductive effect (+I).
29. (c) $CH_3CO > O + 2NH_3 \rightarrow CH_3CONH_2 + CH_3COONH_4$
Acetamide Ammonium acetate

(A)

30. (c)



32. (c) $CH_3COOH \xrightarrow{LiAlH_4} CH_3CH_2OH$
 $CH_3CH_2OH \xrightarrow{I_2/NaOH} CHI_3 \xrightarrow{Ag} C_2H_2$

34. (c) $HCOOH + 2Cu^{+2} \xrightarrow[Solution]{Fehling} Cu_2O + H_2O + CO_2$
Red ppt

Whereas $CH_3COOH \xrightarrow[Solution]{Fehling}$ No reaction

35. (d) $HCOOH + NaHCO_3 \rightarrow HCOONa + H_2O + CO_2$
 $HCHO + NaHCO_3 \rightarrow$ No reaction

37. (d) $CH_3-C \begin{matrix} \diagup O \\ \diagdown OH \end{matrix} \begin{matrix} \cdots HO \\ \cdots O \end{matrix} =C-CH_3$

Due to H-bonding

38. (c) $CH_3COOH + NaHCO_3 \rightarrow CH_3COONa + H_2O + CO_2$

39. (c) Acetic acid forms dimer in benzene due to which molecular mass becomes doubles.


42. (b) $CH_3COOH \xrightarrow{I_2/Red\ p} CH_2Cl-COOH$

44. (c) $CH_3COOH \xrightarrow{\Delta/P_2O_5} (CH_3CO)_2O$

47. (c) $\begin{matrix} COOH \\ | \\ COOH \end{matrix} + KMnO_4 + H_2SO_4 \rightarrow 2CO_2 + H_2O$
 $+K_2SO_4 + MnSO_4$



49. (c) $R-X \xrightarrow{KCN} R-CN \xrightarrow[H_2O]{NaOH} R-COONa + NH_3$

50. (c) $CH_3CH_2COOH \xrightarrow{Cl_2/Fe} CH_3-\underset{\underset{Cl}{|}}{CH}-COOH$
 $CH_2=CH-COOH \xleftarrow[KOH]{Alcohol}$




52. (b)  N-acetyl para amino phenol
NH-COCH₃

53. (b) Because it does not have α-hydrogen atom.











54. (b) Molecular Mass increases due to dimer




In benzene solution



59. (c) $CH_3-CH_2-CCl_2-COOH$; α, α -dichloro butanoic acid is most acidic. Hence it will easily loose H^+ ions in solution.




74. (b)











77. (c) Alcohols react with acids to form esters which have fruity smell.

78. (a) Acidity decreases with the decrease in electronegativity of halogen i.e.,
 $FCH_2COOH > ClCH_2COOH > BrCH_2COOH$





80. (d) Formic acid, $HCOOH$ shows reducing property.

81. (a) $HCOOH$ reduces ammoniacal silver nitrate solution, i.e., Tollen's reagent but acetic acid does not.




99. (a) In this reaction α -H is replaced by chlorine.


100. (a) Urea are neutral in nature in aqueous solution.








1356 Carboxylic acids and Their derivatives



106. (a) Due to intramolecular hydrogen bonding.




108. (d) Acetamide can behave as weak acid as well as base.




112. (b) Benzene sulphonyl chloride is called Hinsberg's reagent.

114. (a) $\text{CH}_3\text{COOH} < \text{ClCH}_2\text{COOH} < \text{Cl}_2\text{CH}_2\text{COOH} < \text{Cl}_3\text{CCOOH}$
Increasing order of acidic nature.



Electron withdrawing group, increases the acidity of benzoic acid, *O*-isomer will have higher acidity than corresponding *m* and *p*-isomer due to ortho effect.


116. (c) $\text{CH}_3\text{COOC}_2\text{H}_5 + \text{NaCl}_{(aq)} \rightarrow$ no reaction
($\text{CH}_3\text{COOC}_2\text{H}_5 + \text{NaCl}_{(aq)}$)






118. (d) $\text{R}-\text{COOR}' + \text{R}''\text{OH} \xrightleftharpoons{\text{H}^+} \text{RCOOR}'' + \text{R}'\text{OH}$

The exchange of alcohol residue known as alcoholysis or transesterification


119. (b) $\text{RCOOH} + \text{PCl}_5 \rightarrow \text{RCOCl} + \text{POCl}_3 + \text{HCl} \xrightarrow{\text{KCN}} \text{RCON} + \text{KCl} \xrightarrow{2\text{H}_2\text{O}} \text{RCO}_2\text{COH} + 2\text{NH}_3$



120. (b) $\text{CH}_3\text{COOH} + \text{Cl}_2 \rightarrow \text{CH}_2\text{ClCOOH} + \text{HCl}$



121. (a) $\text{C}_6\text{H}_5\text{COOC}_2\text{H}_5 + \text{KOH} (\text{alc}) \rightarrow \text{C}_6\text{H}_5\text{COOK} +$




122. (c) $\text{ROH} + \text{PCl}_5 \rightarrow \text{R}-\text{Cl} + \text{POCl}_3 + \text{HCl}$







123. (a) It is picric acid because it has three $-\text{NO}_2$ group are arranged which are ortho and para position

124. (d) Benedict solution is readily reduced by aldehyde. It does not oxidise anhydrides



This reaction is an example of addition reaction.

126. (a) When acetic acid is reacted with PCl_5 the product formed are acetyl chloride, phosphoryl chloride and hydrochloric acid




130. (a) Vinegar is 6 - 10% aqueous solution of acetic acid



Carboxylic acids and Their derivatives 1357

131. (a) All esters are pleasant liquid having pleasant fruity smell. Many of them are used in perfumery, e.g. Benzyl acetate etc.

132. (a) Lactic acid has one asymmetric (chiral) carbon atom, hence it has $(2^n = 2)$ optical isomers.






133. (a) 4-methyl benzene sulphonic acid is stronger than acetic acid thus it will release acetic acid from sodium acetate.



135. (b) $C_6H_5CHO + HCN \rightarrow C_6H_5-CH(OH)CN$ is optically active.

137. (c) Claisen condensation




138. (d) $C_6H_5CONHCH_3 \xrightarrow[HCl]{Zn-Hg} C_6H_5CH_2NHCH_3$

This reaction is known as Clemmenson reduction.


139. (b) $K_a \propto \frac{1}{pK_a}$; The value of K_a is highest for $HCOOH$.

140. (c) $X \xrightarrow{NaOH / CaO} C_2H_6$




141. (a) Glycine is a amphoteric acid as it contains both acidic as well as basic groups.

142. (a) Cinnamic acid reacts with bromine in carbon tetrachloride to give dibromocinnamic acid.





143. (a) OF all the acid derivatives, acid chlorides, i.e. CH_3COCl is most reactive.

The order of reactivity of acid derivatives decreases in the following order,



144. (c) Reaction can be explained as follows




145. (a) $CH_3COOH + NaHCO_3 \rightarrow CH_3COONa + CO_2 + H_2O$




146. (d) $RCOOH \xrightarrow{P_2O_5} RCO > O + H_2O$

P_2O_5 acts as a dehydrating agent.


Uses of Carboxylic Acids and Their Derivatives







Critical Thinking Questions







4. (c) Anion $HCOO^-$ has two resonating structure (identical).



5. (a) Since a single alkyl halide is formed on treatment with HI , it must be a symmetrical ether i.e., ethoxyethane.



8. (d) Resonance stabilization of their conjugate base i.e., carboxylate ion.



11. (c)

12. (b) $-COOH$ and $-OH$ group form the hydrogen bond by which they have high boiling point. $-COOH$ group show strong hydrogen bonding so it form dimer and have more boiling point than $-OH$ group. While $-CHO$ group do not form hydrogen bond. Thus the reactivity order are as $3 > 1 > 2$.



13. (c) CaC_2O_4 is a salt of oxalic acid which is more acidic than acetic acid, so it is insoluble in acetic acid.

14. (d) $-COOH$ and $COCH_3$ are meta directing group



due to the presence of $-\overset{\text{O}}{\parallel}{C}-$, similarly CN is also meta directing due to the presence of multiple bond while $NHCOCH_3$ is ortho/para directing group because of less electron

density over $-\overset{\text{O}}{\parallel}{C}-$ group.

15. (c) Iodoform test is given by all the compounds having CH_3-C- or CH_3-C- group.




17. (c) Formic acid can not be prepared by grignard's reagent. Higher acids are prepared by the reaction of CO_2 on grignard's reagent




Assertion & Reason

1. (a) Carboxylic acid exist as dimer due to the formation of hydrogen bonding.




2. (c) Trichloro acetic acid is stronger than acetic acid because the electron withdrawing group withdraw electrons from the carboxylate base and thus increasing the acid strength.




- I effect
(Stronger acid)

+ I effect
(Weaker acid)

3. (c) Carboxylic acids ($RCOOH$) dissolves in water due to hydrogen bonding between H -atom of $-COOH$ group and O -atom of water. As alkyl portion R is non polar and lyophobic, this effect predominates as $-R$ gets larger (over five carbon atoms).
4. (b) As carboxylic acids are resonance stabilized they do not contain true carbonyl group as is present in carbonyl compounds.





5. (c) Formic acid is stronger than acetic acid



Presence of CH_3 group in acetate ion shows $+I.E.$, and there by intensifying charge on O^{-} of acetate ion than formate ion or acetate ion is destabilized. Thus formate ion is more stable than acetate ion or $HCOOH$ loses proton more easily than CH_3COOH .

6. (e) Both fumaric and maleic acids have two ionisable H^{+} i.e. protons. The maleate monoanion shows intramolecular H -bonding and thus requires more energy to give maleate dianion. It is therefore second dissociation of fumaric acid is more than maleic acid since former does not show intramolecular H -bonding.
7. (c) Larger is the size of alkyl group of ester, greater is the steric effect and thus lesser will be rate of hydrolysis.



8. (e) The melting point of an aliphatic carboxylic acid containing an even number of carbon atoms is higher than the next lower and next higher homologue containing odd number of carbon atoms.
9. (e) Electron withdrawing groups increase the acidity of carboxylic acids by stabilising the conjugate base through delocalisation of the negative charge by inductive and resonance effects.

10. (a) The larger the electron withdrawing inductive effect the greater is the acidity.
11. (c) In aminoacetic acid, NH_2 group is electron repelling in nature.
12. (b) Boiling points of carboxylic acids are higher due to their tendency to associate and form dimers to a greater extent by hydrogen bonding.
13. (a) Both formic acid and oxalic acid behave as reducing agent and decolourise acidified $KMnO_4$ solution.
- $$2KMnO_4 + 3H_2SO_4 \rightarrow K_2SO_4 + 2MnSO_4 + 3H_2O + 5[O]$$
14. (c) Esters containing α -hydrogens on treatment with a base form a carbanion which brings about nucleophilic acyl substitution at the carbonyl group of the other molecule of the ester to form β -keto esters.