

# Tips & Tricks

- ✍ Alkyl nitrites are the esters of nitrous acid.
- ✍ Nitroparaffins are used as solvents for oils, fats, resins, esters, rubbers and cellulose etc. nitromethane is used as high power fuel in racing automobiles.
- ✍ Nitrobenzene is good solvent in Friedel Crafts reaction because it dissolves  $AlCl_3$
- ✍ All amines have basic properties. The basic property, that is, the tendency of primary, secondary and tertiary amines to bind a proton, is due to the unshared pair of electrons on the nitrogen. When a proton is bound, positive ion is formed and originally electrically neutral amine takes on the charge of the proton. When ions are formed in this way, they are called onium ions. The ion formed in case of amines are substituted ammonium ions. The hydronium ion,  $H_3O^+$  is also the onium ion, which belongs to the class of oxonium ions.
- ✍ Some derivatives of ammonia arranged in order of decreasing basicity are  $(CH_3)_4N^+OH^-$ ,  $(CH_3)_2NH$ ,  $CH_3NH_2$ ,  $(CH_3)_3N$ ,  $NH_3$ ,  $C_6H_5NH_2$ ,  $C_6H_5NHCH_3$ ,  $C_6H_5NH_2$ ,  $(C_6H_5)_2NH$ ,  $CH_3CONH_2$ .
- ✍ In water the basicity follows the order : Primary < Tertiary < Secondary amine, with reference to hydronium ion,  $H_3O^+$ . In this case solvation factor and steric effect alter, to some extent, the order of basicity because of inductive effect alone.
- ✍ In a non-polar solvent such as benzene, using trichloroacetic acid as the reference acid, the basicity follows the order Tertiary < Secondary < Primary amines. The solvation factor is absent but steric effect upsets the inductive effect of alkyl groups.
- ✍ Carbylamine test is specific for primary amines.

## Ordinary Thinking

### Objective Questions

#### Introduction of Nitrogen Containing Compounds

1. Cyanide ion is
 

(a) Nucleophilic	(b) Electrophilic
(c) Strongly acidic	(d) Non-reactive

 and neutral
2. Compounds containing both amino and COOH groups are known as
 

(a) Diamines	(b) Unknown
(c) Amino acids	(d) Enzymes
3. Which of the following is  $1^\circ$  amine
 

(a) Ethylene diamine	(b) Dimethyl amine
(c) Trimethyl amine	(d) <i>N</i> -methyl aniline
4.  $C_3H_9N$  represents [AMU 1988]

(a) Primary amine	(b) Secondary amine
(c) Tertiary amine	(d) All of these
5.  $(CH_3)_2C \cdot CH_2 \cdot CO \cdot CH_3$  is [MP PET/PMT 1988]

$$\begin{array}{c} | \\ NH_2 \end{array}$$

(a) Diacetone	(b) Acetoneamine
(c) Diacetoneamine	(d) Aminoacetone
6. A secondary amine is [KCET 1992]

(a) An organic compound with two $-NH_2$ groups	(b) A compound with two carbon atoms and an $-NH_2$ group
(c) A compound with an $-NH_2$ group on the carbon atom in number 2 position	(d) A compound in which two of the hydrogens of $NH_3$ have been replaced by organic groups
7. The structural formula of methyl aminomethane is [MP PMT 1991]

(a) $(CH_3)_2CHNH_2$	(b) $(CH_3)_3N$
(c) $(CH_3)_2NH$	(d) $CH_3NH_2$
8. Allyl isocyanide has [IIT 1995]

(a) 9 sigma bonds and 4 pi bonds	(b) 8 sigma bonds and 5 pi bonds
(c) 8 sigma bonds, 3 pi bonds and 4 non-bonding electrons	(d) 9 sigma bonds, 3 pi bonds and 2 non-bonding electrons
9. Triaminobenzene is a [BHU 1996]

(a) $2^\circ$ amine	(b) $3^\circ$ amine
(c) $1^\circ$ amine	(d) Quarternary salt
10.  $CH_2 = CH - CH_2 - NH - CH_3$  is a [RPET 2000]

(a) Secondary amine	(b) Primary amine
(c) Tertiary amine	(d) None of these
11. Leakage of which gas was responsible for the Bhopal tragedy in 1984 [MP PET 2001]

(a) $CH_3 - N = C = O$	(b) $CH_3 - C - N = S$
(c) $CHCl_3$	(d) $C_6H_5COCl$
12. Which of the following is not a nitro-derivative [DCE 2004]

(a) $C_6H_5NO_2$	(b) $CH_3CH_2ONO$
(c) $CH_3CH - N \begin{array}{l} \nearrow O \\ \searrow O \end{array}$	(d) $C_6H_4(OH)NO_2$
13. Acetonitrile is: [MP PMT 2004]

(a) $C_2H_5CN$	(b) $CH_3CN$
----------------	--------------





- (c)  $CH_3COCN$  (d)  $C_6H_5CH_2CN$
14. In alkyl cyanide alkyl group attached with [BCECE 2005]  
 (a) C of CN group  
 (b) N of CN group  
 (c) Either C or N of CN group  
 (d) Both C and N of CN group
15. Number of isomeric primary amines obtained from  $C_4H_{11}N$  are  
 (a) 3 (b) 4  
 (c) 5 (d) 6

### Preparation of Nitrogen Containing Compounds


1. Amides may be converted into amines by reaction named after [CPMT 1974; MP PET 1992; CBSE PMT 1999]  
 (a) Perkin (b) Claisen  
 (c) Hoffmann (d) Kolbe
2. Reaction  $CH_3CONH_2 \xrightarrow{NaOBr}$  gives [CPMT 1983, 93, 97]  
 (a)  $CH_3Br$  (b)  $CH_4$   
 (c)  $CH_3COBr$  (d)  $CH_3NH_2$
3. Acetamide is treated separately with the following reagents. Which would give methyl amine [IIT 1983; CPMT 1988, 94; MP PET 1993; MP PMT 1996; AIIMS 1998]  
 (a)  $PCl_5$  (b)  $NaOH + Br_2$   
 (c) Sodalime (d) Hot conc.  $H_2SO_4$
4. The amine formed from an amide by means of bromine and alkali has  
 (a) Same number of C atoms as that of amide  
 (b) One less C atom than that of amide  
 (c) One more C atom than that of amide  
 (d) Two more C atoms than that of amide
5.  $CH_3CN \xrightarrow{Na+C_2H_5OH} X$   
 The compound X is [MP PMT 1983; BHU 1984]  
 (a)  $CH_3CONH_2$  (b)  $CH_3CH_2NH_2$   
 (c)  $C_2H_6$  (d)  $CH_3NHCH_3$
6. Ethylamine can be prepared by the action of bromine and caustic potash on [CPMT 1994]  
 (a) Acetamide (b) Propionamide  
 (c) Formamide (d) Methyl cyanide
7. Ethylamine can be obtained by the [CPMT 1985]  
 (a) Action of  $NH_3$  on ethyl iodide  
 (b) Action of  $NH_3$  on ethyl alcohol  
 (c) Both (a) and (b)  
 (d) None of the above
8. Aniline is usually purified by [CPMT 1983, 93; JIPMER 1997]  
 (a) Steam distillation (b) Simple distillation  
 (c) Vacuum distillation (d) Extraction with a solvent
9. Reduction of nitroalkanes yields  
 (a) Acid (b) Alcohol  
 (c) Amine (d) Diazo compounds
10. Acetamide changes into methylamine by  
 (a) Hofmann bromamide reaction  
 (b) Hofmann reaction  
 (c) Friedel-Craft's reaction  
 (d) Hofmann isomerization
11. When methyl iodide is heated with ammonia, the product obtained is  
 (a) Methylamine  
 (b) Dimethylamine  
 (c) Trimethylamine  
 (d) A mixture of the above three amines
12. Acetanilide can be prepared from aniline and which of the following  
 (a) Ethanol (b) Acetaldehyde  
 (c) Acetone (d) Acetic anhydride
13. Reduction of nitroalkanes in neutral medium (e.g.  $Zn / NH_4Cl$ ) forms mainly  
 (a)  $R-NH_2$  (b)  $R-NHOH$   
 (c)  $R-N=N-Cl$  (d) All of these
14. Nitrosobenzene can be prepared by oxidizing aniline from  
 (a)  $H_2SO_4$  (b)  $H_2SO_5$   
 (c)  $H_2SO_3$  (d)  $K_2Cr_2O_7$
15. The Hinsberg's method is used for  
 (a) Preparation of primary amines  
 (b) Preparation of secondary amines  
 (c) Preparation of tertiary amines  
 (d) Separation of amine mixtures
16. Which one of the following compound gives a secondary amine on reduction  
 (a) Nitromethane (b) Nitrobenzene  
 (c) Methyl isocyanide (d) Methyl cyanide
17. Chloropicrin is manufactured by the reaction between  $Cl_2$ ,  $NaOH$  and  
 (a) Nitromethane (b) Nitroethane  
 (c) Nitrophenol (d) Nitrostyrene
18. In the reaction  

$$R-\overset{\overset{O}{||}}{C}-OH \xleftarrow{H_3O^+} X \xrightarrow{[H]} RCH_2NH_2; 'X' \text{ is}$$
 [MP PMT 1990]  
 (a) Isonitrile (b) Nitrile  
 (c) Nitrite (d) Oxime
19. When ethanol is mixed with ammonia and passed over alumina the compound formed is [CBSE PMT 1990]  
 (a)  $C_2H_5NH_2$  (b)  $C_2H_4$   
 (c)  $C_2H_5OC_2H_5$  (d)  $CH_3OCH_3$
20. Which of the following reactions does not yield an amine [CPMT 1989, 93]  
 (a)  $RX + NH_3 \longrightarrow$

## 1380 Nitrogen Containing Compounds






21. Identify 'B' in the reaction



22. Which of the following gives primary amine on reduction

[MP PMT 1995]



23. Which of the following is converted into an alcohol on treatment with  $HNO_2$  [MP PET 1996; MP PMT 1999]





24. Which of the following gives  $RNC$ , when reacted with  $CHCl_3$  and  $KOH$


[MP PET 1996]






25. When aniline reacts with  $NaNO_2$  and dil.  $HCl$  at  $0^\circ - 5^\circ C$ , the product formed is [MP PMT 1996; AIIMS 1994]




26. Starting from propanoic acid, the following reactions were carried out





What is the compound Z




27. In the reaction





the final product (C) is



28. In the following reaction, X is




[CPMT 1999]






29. Which of the following reactions will not give primary amine

[CPMT 1999]




30. Carbylamine reaction is given by

[BHU 1996; EAMCET 1990]








31. The reaction





is known as

[BHU 1996]





What is Z?

[CPMT 1996]





33. Which of the following reacts with chloroform and a base to form phenyl isocyanide [AFMC 1997]






Aromatic primary amine when treated with cold  $HNO_2$  gives

[Pb. CET 2002; DCE 1999]




35. Which of the following compound is the strongest base

[BHU 1999]







36. Nitrobenzene combines with hydrogen in the presence of platinum to produce

[BHU 1999]



The product is

[RPET 2000]



[Orissa JEE 2002]





- (c) \_\_\_\_\_ (d) \_\_\_\_\_
38. Ethyl amine on heating with  $CS_2$  in presence of  $HgCl_2$  forms [MP PET 2000]  
 (a)  $C_2H_5NCS$  (b)  $(C_2H_5)_2S$   
 (c)  $(C_2H_5)_2CS$  (d)  $C_2H_5(CS)_2$
39. Which of the following reacts with  $NaNO_2 + HCl$  to give phenol [MP PMT 2000]  
 (a)  $C_6H_5CH_2NHCH_3$  (b)  $(CH_3)_2NH$   
 (c)  $CH_3NH_2$  (d)  $C_6H_5NH_2$
40. Which of the following reactions give  $RCONH_2$  [Roorkee 2000]  
 (a)  $R-C \equiv N + H_2O \xrightarrow{HCl}$   
 (b)  $RCOONH_4 \xrightarrow{\text{heat}}$   
 (c)  $R-COCl + NH_3 \longrightarrow$   
 (d)  $(RCO)_2O + NH_3 \longrightarrow$
41. When chlorobenzene is treated with  $NH_3$  in presence of  $Cu_2O$  in xylene at 570 K. The product obtained is [Pb. PMT 2000]  
 (a) Benzylamine (b) Diazonium salt  
 (c) Schiff's base (d) Aniline
42. Nitrobenzene can be prepared from benzene by using a mixture of conc.  $HNO_3$  and conc.  $H_2SO_4$ . In the nitrating mixture,  $HNO_3$  acts as a [BHU 2001]  
 (a) Base (b) Acid  
 (c) Catalyst (d) Reducing agent
43. The rate determining step for the preparation of nitrobenzene from benzene is [AIIMS 2001]  
 (a) Removal of  $NO_2^+$  (b) Removal of  $NO_3^+$   
 (c) Formation of  $NO_2^+$  (d) Formation of  $NO_3^+$
44. In this reaction  $C_6H_5NH_2 + HCl + NaNO_2 \rightarrow X$ . Product X is [RPMT 2002; AFMC 2002]  
 (a) Aniline hydrochloride  
 (b) Nitro aniline  
 (c) Benzenediazonium chloride  
 (d) None of these
45. The diazonium salts are the reaction products in presence of excess of mineral acid with nitrous acid and [MP PET 2002]  
 (a) Primary aliphatic amine  
 (b) Secondary aromatic amine  
 (c) Primary aromatic amine  
 (d) Tertiary aliphatic amine
46. In acid medium nitrobenzene is reduced to aniline as shown in the reaction  $C_6H_5-NO_2 + 6[H] \rightarrow C_6H_5-NH_2 + 2H_2O$   
 The reducing agent used in this reaction is .....  
 (a)  $LiAlH_4$  (b)  $Sn/HCl$   
 (c)  $Na$ /alcohol (d)  $H_2/Ni$
47. When aniline is treated with sodium nitrite and hydrochloric acid at  $0^\circ C$ , it gives [Orissa JEE 2003]  
 (a) Phenol and  $N_2$  (b) Diazonium salt  
 (c) Hydrazo compound (d) No reaction takes place
48.  $CH_3NO_2 \xrightarrow{Sn+HCl} CH_3X$ , the 'X' contain [CPMT 2003]  
 (a)  $-NH_2$  (b)  $-COOH$   
 (c)  $-CHO$  (d)  $(CH_3CO)_2O$
49. In the series of reaction  $C_6H_5NH_2 \xrightarrow[0-5^\circ C]{NaNO_2/HCl} X \xrightarrow[CH_2O]{HNO_2} Y + N_2 + HCl$  X and Y are respectively [EAMCET 2003]  
 (a)  $C_6H_5-N=N-C_6H_5, C_6H_5N_2^+Cl^-$   
 (b)  $C_6H_5N_2^+Cl^-, C_6H_5-N=N-C_6H_5$   
 (c)  $C_6H_5N_2^+Cl^-, C_6H_5NO_2$   
 (d)  $C_6H_5NO_2, C_6H_6$
50. Aromatic nitriles ( $ArCN$ ) are not prepared by reaction [AIIMS 2004]  
 (a)  $ArX + KCN$  (b)  $ArN_2^+ + CuCN$   
 (c)  $ArCONH_2 + P_2O_5$  (d)  $ArCONH_2 + SOCl_2$
51. An organic amino compound reacts with aqueous nitrous acid at low temperature to produce an oily nitroso amine. The compound is :  
 (a)  $CH_3NH_2$  (b)  $CH_3CH_2NH_2$   
 (c)  $CH_3CH_2NH.CH_2CH_3$  (d)  $(CH_3CH_2)_3$
52. Azo-dyes are prepared from : [BHU 2004; Pb. CET 2001]  
 (a) Aniline (b) Salicylic acid  
 (c) Benzaldehyde (d) Chlorobenzene
53. Gabriel's phthalimide synthesis is used for the preparation of [CPMT 1982; DPMT 1983]  
 (a) Primary aromatic amine (b) Secondary amine  
 (c) Primary aliphatic amine (d) Tertiary amine
54. For the preparation of *p*-nitroiodobenzene from *p*-nitroaniline, the best method is [Orissa JEE 2005]  
 (a)  $NaNO_2/HCl$  followed by  $KI$   
 (b)  $NaNO_2/HCl$  followed by  $CuCN$   
 (c)  $LiAlH_4$  followed by  $I_2$   
 (d)  $NaBH_4$  followed by  $I_2$
55.  $KCN$  reacts readily to give a cyanide with [J & K 2005]  
 (a) Ethyl alcohol (b) Ethyl bromide  
 (c) Bromobenzene (d) Chlorobenzene

### Properties of Nitrogen Containing Compounds

1. Which of the following amine will not react with nitrous acid to give nitrogen  
 (a)  $CH_3NH_2$  (b)  $CH_3-CH_2-NH_2$

## 1382 Nitrogen Containing Compounds

- (c)  $\text{CH}_3 - \text{CH} - \text{NH}_2$  (d)  $(\text{CH}_3)_3\text{N}$   
 $\quad \quad \quad |$   
 $\quad \quad \quad \text{CH}_3$
2. Which of the following compound is expected to be most basic [NCERT 1982]  
 (a) Aniline (b) Methylamine  
 (c) Hydroxylamine (d) Ethylamine
3. Which of the following compounds is an amino acid [Manipal MEE 1995]  
 (a)  $\text{CH}_3 - \text{CH}_2 - \overset{\text{O}}{\parallel} \text{C} - \text{O} - \text{NH}_4$   
 (b)  $\text{CH}_3 - \overset{\text{O}}{\parallel} \text{CH} - \text{C} - \text{OH}$   
 $\quad \quad \quad |$   
 $\quad \quad \quad \text{NH}_2$   
 (c)  $\text{CH}_3 - \text{CH}_2 - \overset{\text{O}}{\parallel} \text{C} - \text{NH}_2$   
 (d)  $\text{CH}_3 - \overset{\text{O}}{\parallel} \text{CH} - \text{C} - \text{Cl}$   
 $\quad \quad \quad |$   
 $\quad \quad \quad \text{NH}_2$
4. Nitro group in nitrobenzene is a [MNR 1986]  
 (a) Ortho director (b) Meta director  
 (c) Para director (d) Ortho and para director
5. The alkyl cyanides are  
 (a) Acidic (b) Basic  
 (c) Neutral (d) Amphoteric
6. The alkyl cyanides when hydrolysed to the corresponding acid, the gas evolved is  
 (a)  $\text{N}_2$  (b)  $\text{O}_2$   
 (c)  $\text{NH}_3$  (d)  $\text{CO}_2$
7. Aniline when treated with  $\text{HNO}_2$  and  $\text{HCl}$  at  $0^\circ\text{C}$  gives [CPMT 1982, 89; RPMT 2000]  
 (a) Phenol (b) Nitrobenzene  
 (c) A diazo compound (d) None of these
8. Nitrosobenzene can be isolated from nitrobenzene under [DPMT 1982]  
 (a) Metal and acid  
 (b)  $\text{Zn}$  dust and  $\text{NH}_4\text{Cl}$   
 (c) Alkaline sodium arsenite  
 (d) Cannot be isolated
9. Alkyl cyanides when react with Grignard reagent, the product on hydrolysis found, is  
 (a) Aldehyde (b) Ketone  
 (c) Alcohol (d) Acid
10. The product formed when benzene is nitrated by fuming nitric acid is [MP PMT 1979]  
 (a) *m*-dinitrobenzene (b) Nitrobenzene  
 (c) *sym*-trinitrobenzene (d) None of these
11. Ethyl amine undergoes oxidation in the presence of  $\text{KMnO}_4$  to form [CPMT 1985]  
 (a) An acid (b) An alcohol  
 (c) An aldehyde (d) A nitrogen oxide
12. Which of the following amines would undergo diazotisation  
 (a) Primary aliphatic amines (b)  
 (c) Both (a) and (b) (d) None of these
13. Reaction of primary amines with aldehyde yields [NCERT 1984; Manipal MEE 1995]  
 (a) Amides (b) Aldimines  
 (c) Nitriles (d) Nitro compounds
14. When acetamide is treated with  $\text{HNO}_2$ , the gas is evolved [CPMT 1993]  
 (a)  $\text{H}_2$  (b)  $\text{O}_2$   
 (c)  $\text{N}_2$  (d)  $\text{CH}_4$
15. Nitrobenzene on nitration gives [NCERT 1978; CPMT 1989]  
 (a) *o*-dinitrobenzene (b) *p*-dinitrobenzene  
 (c) *m*-dinitrobenzene (d) *o*- and *p*-nitrobenzene
16. Reduction of alkyl nitrites yields  
 (a) Alcohol (b) Base  
 (c) Amine (d) Acid
17. When primary amines are treated with  $\text{HCl}$ , the product obtained is  
 (a) An alcohol (b) A cyanide  
 (c) An amide (d) Ammonium salt
18. Which one is weakest base [BHU 1982; RPMT 2000]  
 (a) Ammonia (b) Methylamine  
 (c) Dimethylamine (d) Trimethylamine
19. Chloroform when treated with aniline and alcoholic  $\text{KOH}$  gives [CPMT 1986; EAMCET 1992; MP PMT 1997; Pb. PMT 1999]  
 (a) Phenyl cyanide (b) Phenyl isocyanide  
 (c) Chlorobenzene (d) Phenol
20. Which of following do not react with  $\text{HNO}_2$   
 (a) Primary nitroalkanes (b) Secondary nitroalkanes  
 (c) Tertiary nitroalkanes (d) All of these
21. Primary amines can be distinguished from secondary and tertiary amines by reacting with [CPMT 1986; MP PMT 1980]  
 (a) Chloroform and alcoholic  $\text{KOH}$   
 (b) Methyl iodide  
 (c) Chloroform alone  
 (d) Zinc dust
22. Which of following is not an usual method for preparation of primary amine  
 (a) Hofmann's method (b) Curtius reaction  
 (c) Schmidt reaction (d) Friedel-Craft's reaction
23. A solution of methyl amine  
 (a) Turns blue litmus red  
 (b) Turns red litmus blue  
 (c) Does not affect red or blue litmus  
 (d) Bleaches litmus













24. Mark the correct statement  
[CPMT 1974; DPMT 1983; MP PMT 1994]
- (a) Methyl amine is slightly acidic  
(b) Methyl amine is less basic than  $NH_3$   
(c) Methyl amine is stronger base than  $NH_3$   
(d) Methyl amine forms salts with alkalis
25. The product of mustard oil reaction is  
(a) Alkyl isothiocyanate (b) Dithio carbonamide  
(c) Dithio ethylacetate (d) Thioether
26. Which of the following is azo- group  
(a)  $-N =$  (b)  $-N = N -$   
(c)  $-NH -$  (d)  $-CO - NH -$
27. 'Oil of mirbane' is  
(a) Aniline (b) Nitrobenzene  
(c) *p*-nitroaniline (d) *p*-aminoazobenzene
28. The maximum number of  $-NO_2$  groups that can be introduced by nitration in benzene is usually  
(a) 4 (b) 2  
(c) 3 (d) 6
29. Nitrobenzene at room temperature is  
(a) Gas (b) Liquid  
(c) Solid (d) Solution
30. In the explosive amatol, TNT is mixed with [CPMT 1988]  
(a) Ammonium citrate (b) Ammonium nitrate  
(c) Ammonium oxalate (d) Ammonium sulphate
31. By reduction of nitrosobenzene which of the following is not obtained
- (a)  (b)   
(c)  (d) 
32. By the presence of a halogen atom in the ring, basic properties of aniline is  
(a) Increased (b) Decreased  
(c) Unchanged (d) Doubled
33. In the mustard oil reaction, an amine is treated with  
(a)  $Na / C_2H_5OH$  (b)  $Sn / HCl$   
(c)  $CS_2$  (d)  $K_2Cr_2O_7 / H_2SO_4$
34. Primary nitro compounds when react with  $HNO_2$  forms crystalline solids which on treatment with  $NaOH$  gives  
(a) Red solution (b) Blue solution  
(c) White precipitate (d) Yellow colouration
35. Secondary nitro compounds when react with  $HNO_2$  forms crystalline solids which one on treatment with  $NaOH$  gives  
(a) Red solution (b) Blue solution  
(c) White precipitate (d) Yellow colouration
36. Which of the following possess powerful mustard smell (and are called mustard oils)  
(a) Alkyl isocyanates (b) Alkyl cyanates  
(c) Alkyl isothiocyanates (d) Alkyl thiocyanates
37. On heating acetamide in presence of  $P_2O_5$ , which of the following is formed [MP PMT 1992; MP PET 1994; Kurukshetra CEE 1998]  
(a) Ammonium acetate (b) Acetonitrile  
(c)  $NH_3$  (d) Methylamines
38. When chloroform reacts with ethyl amine in presence of alcoholic  $KOH$ , the compound formed is  
[CPMT 1983; MP PMT 1993; CBSE PMT 1997; BHU 1999; AIEEE 2002]  
(a) Ethyl cyanide (b) Ethyl isocyanide  
(c) Formic acid (d) An amide
39. When methyl cyanide is hydrolysed in presence of alkali, the product is [MP PMT 1993; BCECE 2005]  
(a) Acetamide (b) Methane  
(c)  $CO_2 + H_2O$  (d) Acetic acid
40. Hofmann's hypobromite reaction affords a method of  
[MP PMT 1993]  
(a) Preparing a tertiary amine  
(b) Preparing a mixture of amines  
(c) Stepping down a series  
(d) Stepping up a series
41. The compound which on reaction with aqueous nitrous acid on  $HNO_2$  at low temperature produces an oily nitrosoamine is [IIT 1981; CPMT 1989; MP 1991; Kurukshetra CEE 1998; MP PMT 2001]  
(a) Diethylamine (b) Ethylamine  
(c) Aniline (d) Methylamine
42. Identify the product Z in the series  
 $CH_3CN \xrightarrow{Na + C_2H_5OH} X \xrightarrow{HNO_2} Y \xrightarrow[H_2SO_4]{K_2Cr_2O_7} Z$   
[AIIMS 1983; JIPMER 2001]  
(a)  $CH_3CHO$  (b)  $CH_3CONH_2$   
(c)  $CH_3COOH$  (d)  $CH_3CH_2NHOH$
43. The end product of the reactions is  
 $C_2H_5NH_2 \xrightarrow{HNO_2} A \xrightarrow{PCl_5} B \xrightarrow{H.NH_2} C$   
[CPMT 1988, 89, 93; DCE 1999; JIPMER 2000]  
(a) Ethyl cyanide (b) Ethyl amine  
(c) Methyl amine (d) Acetamide
44. Primary and secondary amines are distinguished by  
[AMU 1988; MP PMT 1996]  
(a)  $Br_2 / KOH$  (b)  $HClO_4$   
(c)  $HNO_2$  (d)  $NH_3$
45. Which one of the following will give a primary amine on hydrolysis  
[BHU 1982]  
(a) Nitroparaffin (b) Alkyl cyanide  
(c) Oxime (d) Alkyl isocyanide
46. Methyl amine reacts with  $HNO_2$  giving [RPMT 1997]  
(a)  $CH_3O - N = O$  (b)  $CH_3 - O - CH_3$   
(c)  $CH_3OH$  (d) (a) and (b) both



## 1384 Nitrogen Containing Compounds

47. Nitrobenzene on reduction by zinc and  $NH_4Cl$  gives  
 [CPMT 1989, 94; BHU 1996; Pb. PMT 1999]  
 (a) Aniline (b) Nitrosobenzene  
 (c) Hydrazobenzene (d) Phenylhydroxyl amine
48. The decreasing order of the basic character of the three amines and ammonia is [MP PET/PMT 1988; KCET 1990]  
 (a)  $NH_3 > CH_3NH_2 > C_2H_5NH_2 > C_6H_5NH_2$   
 (b)  $C_2H_5NH_2 > CH_3NH_2 > NH_3 > C_6H_5NH_2$   
 (c)  $C_6H_5NH_2 > C_2H_5NH_2 > CH_3NH_2 > NH_3$   
 (d)  $CH_3NH_2 > C_2H_5NH_2 > C_6H_5NH_2 > NH_3$
49. Correct order of increasing basicity is [CBSE PMT 1992]  
 (a)  $NH_3 < C_6H_5NH_2 < (C_2H_5)_2NH < C_2H_5NH_2 < (C_2H_5)_3N$   
 (b)  $C_6H_5NH_2 < NH_3 < (C_2H_5)_3N < (C_2H_5)_2NH < C_2H_5NH_2$   
 (c)  $C_6H_5NH_2 < NH_3 < C_2H_5NH_2 < (C_2H_5)_3N < (C_2H_5)_2NH$   
 (d)  $C_6H_5NH_2 < (C_2H_5)_3N < NH_3 < C_2H_5NH_2 < (C_2H_5)_2NH$
50. Among the following compounds nitrobenzene, benzene, aniline and phenol, the strongest basic behaviour in acid medium is exhibited by [KCET 1993]  
 (a) Phenol (b) Aniline  
 (c) Nitrobenzene (d) Benzene
51. Aniline on treatment with excess of bromine water gives  
 [AFMC 1990; MP PMT 1991; RPMT 1997]  
 (a) Aniline bromide (b) *o*-bromoaniline  
 (c) *p*-bromoaniline (d) 2, 4, 6-tribromoaniline
52. Unpleasant smelling carbylamines are formed by heating alkali and chloroform with [KCET 1987, 2000, 01]  
 (a) Any amine (b) Any aliphatic amine  
 (c) Any aromatic amine (d) Any primary amine
53. When an organic compound was treated with sodium nitrite and hydrochloric acid in the ice cold, nitrogen gas was evolved copiously. The compound is [KCET 1986]  
 (a) A nitro compound  
 (b) A primary amine  
 (c) An aliphatic primary amine  
 (d) An aromatic primary amine
54. Aniline reacts with alkyl halide to give [KCET 1984]  
 (a) Amino compound  
 (b) Tertiary compound  
 (c) Quaternary ammonium compound  
 (d) Azomethane
55. Aniline on treatment with conc.  $HNO_3$  + conc.  $H_2SO_4$  mixture yields [AIIMS 1992]  
 (a) *o*- and *p*-nitroanilines (b) *m*-nitroanilines  
 (c) A black tarry matter (d) No reaction
56. Which statement is not correct [MP PMT 1995]  
 (a) Amines form hydrogen bond  
 (b) Ethyl amine has higher boiling point than propane  
 (c) Methyl amine is more basic than ammonia  
 (d) Dimethyl amine is less basic than methyl amine
57. Which of the following is not used as an explosive [MP PET 1996]  
 (a) Trinitrotoluene (b) Trinitrobenzene  
 (c) Picric acid (d) Nitrobenzene
58. Primary amines react with nitrous acid to yield  
 (a) Insoluble nitrite salts (b) Yellow oily layer  
 (c) Nitrogen gas (d) Azo dye
59. Which of the following has the smell of bitter almonds  
 (a) Nitromethane (b) Nitroethane  
 (c) Nitrobenzene (d) Aniline
60. The reaction of  $HNO_2$  with 'A' gives quaternary ammonium salt. A is [MP PMT 1997]  
 (a) Methyl amine (b) Dimethyl amine  
 (c) Trimethyl amine (d) Aniline
61. Reaction of nitrous acid with aliphatic primary amine in the cold gives [MP PET/PMT 1998; CBSE PMT 1994]  
 (a) A diazonium salt (b) An alcohol  
 (c) A nitrite (d) A dye
62. In presence of acid, hydrolysis of methyl cyanide gives  
 [MP PET/PMT 1998]  
 (a) Acetic acid (b) Methylamine  
 (c) Methyl alcohol (d) Formic acid
63. The amine which does not react with acetyl chloride is or Which of the following cannot be acetylated  
 [MP PET 1999; MP PMT 1999]  
 (a)  $CH_3NH_2$  (b)  $(CH_3)_2NH$   
 (c)  $(CH_3)_3N$  (d) None of these
64. The fusion of sodium with amine gives mainly [MP PMT 1999; CPMT 2002]  
 (a)  $NaCN$  (b)  $NaN_3$   
 (c)  $NaSCN$  (d)  $NaNO_2$
65. Which of the following is most basic [MP PMT 1999]  
 (a)  $C_6H_5NH_2$  (b)  $(CH_3)_2NH$   
 (c)  $(CH_3)_3N$  (d)  $NH_3$
66. In reaction  


$$CH_3CN + 2H \xrightarrow[\text{Ether}]{HCl} X \xrightarrow{\text{Boiling } H_2O} Y$$
; the term Y is  
 [CBSE PMT 1999]  
 (a) Acetone (b) Ethylamine  
 (c) Acetaldehyde (d) Dimethylamine
67. The following compound can be classified as *N-N* dimethyl propanamine, *N*-methyl aniline and aniline [Bihar MEE 1996]  
 (a) Primary, secondary, tertiary  
 (b) Primary, tertiary, secondary  
 (c) Secondary, tertiary, primary

- (d) Tertiary, primary, secondary  
(e) None of these
68. Which of the following compounds does not react with  $\text{NaNO}_2$  and  $\text{HCl}$  [KCET 1996]  
(a)  $\text{C}_6\text{H}_5\text{OH}$  (b)  $\text{C}_6\text{H}_5\text{NH}_2$   
(c)  $(\text{CH}_3)_3\text{CNO}_2$  (d)  $(\text{CH}_3)_3\text{CHNO}_2$
69. In the reduction of nitrobenzene, which of the following is the intermediate  
(a)  $\text{C}_6\text{H}_5\text{N} = \text{O}$   
(b)  $\text{C}_6\text{H}_5\text{NH} - \text{NH} - \text{C}_6\text{H}_5$   
(c)  $\text{C}_6\text{H}_5 - \text{N} = \text{N} - \text{C}_6\text{H}_5$   
(d)  $\text{C}_6\text{H}_5\text{N} = \overset{\text{O}}{\underset{\uparrow}{\text{N}}} - \text{C}_6\text{H}_5$
70. Aniline when treated with conc.  $\text{HNO}_3$  gives [KCET 1996]  
(a)   (b)   
(c)  (d) 
71. Which one of the following is not a base [EAMCET 1997]  
(a)  $\text{N}_2\text{H}_4$  (b)  $\text{NH}_2\text{OH}$   
(c)  $(\text{CH}_3)_3\text{N}$  (d)  $\text{HN}_3$
72. *p*-Nitrobromobenzene can be converted to *p*-nitroaniline by using  $\text{NaNH}_2$ . The reaction proceeds through the intermediate named [Orissa JEE 2005]  
(a) Carbocation (b) Carbanion  
(c) Benzyne (d) Dianion
73. If methyl is alkyl group, then which order of basicity is correct [RPMT 1997]  
(a)  $\text{R}_2\text{NH} > \text{RNH}_2 > \text{R}_3\text{N} > \text{NH}_3$   
(b)  $\text{R}_2\text{NH} > \text{R}_3\text{N} > \text{RNH}_2 > \text{NH}_3$   
(c)  $\text{RNH}_2 > \text{NH}_3 > \text{R}_2\text{NH} > \text{R}_3\text{N}$   
(d)  $\text{NH}_3 > \text{RNH}_2 > \text{R}_2\text{NH} > \text{R}_3\text{N}$
74. Which of the following has the minimum heat of dissociation [Roorkee Qualifying 1998]  
(a)  $(\text{CH}_3)_3\text{N} \rightarrow \text{BF}_3$   
(b)  $(\text{CH}_3)_3\text{N} \rightarrow \text{B}(\text{CH}_3)\text{F}_2$   
(c)  $(\text{CH}_3)_3\text{N} \rightarrow \text{B}(\text{CH}_3)_2\text{F}$   
(d)  $(\text{CH}_3)_3\text{N} \rightarrow \text{B}(\text{CH}_3)_3$
75. The major product (70% to 80%) of the reaction between *m*-dinitrobenzene with  $\text{NH}_4\text{HS}$  is [AIIMS 1997]  
(a)  (b)   
(c)  (d) 
- (c) (d)
76. Which one is less alkaline [CPMT 1997]  
(a)   $\text{NH}_2$  (b)   $\text{NH}_2$   
(c)   $\text{NH}_2$  (d) All of these
77. In the diazotisation of aniline with sodium nitrite and hydrochloric acid, an excess of hydrochloric acid is used primarily to [Pb. PMT 1998]  
(a) Suppress the concentration of free aniline available for coupling  
(b) Suppress hydrolysis of phenol  
(c) Insure a stoichiometric amount of nitrous acid  
(d) Neutralize the base liberated
78. A primary amine can be converted to an alcohol by the action of [CET Pune 1998]  
(a) Alkali (b) Nitrous acid  
(c) Reducing agent (d) Oxidising agent
79. Arrange the following in increasing order of basicity  $\text{CH}_3\text{NH}_2$ ,  $(\text{CH}_3)_2\text{NH}$ ,  $\text{C}_6\text{H}_5\text{NH}_2$ ,  $(\text{CH}_3)_3\text{N}$  [AFMC 1997]  
(a)  $(\text{CH}_3)_3\text{N} < (\text{CH}_3)_2\text{NH} < \text{CH}_3\text{NH}_2 < \text{C}_6\text{H}_5\text{NH}_2$   
(b)  $(\text{CH}_3)_3\text{N} > (\text{CH}_3)_2\text{NH} > \text{CH}_3\text{NH}_2 > \text{C}_6\text{H}_5\text{NH}_2$   
(c)  $\text{C}_6\text{H}_5\text{NH}_2 < (\text{CH}_3)_3\text{N} < \text{CH}_3\text{NH}_2 < (\text{CH}_3)_2\text{NH}$   
(d)  $\text{C}_6\text{H}_5\text{NH}_2 > (\text{CH}_3)_3\text{N} > \text{CH}_3\text{NH}_2 > (\text{CH}_3)_2\text{NH}$
80. In the reaction  
$$\text{CH}_3\text{CN} + \text{CH}_3\text{MgI} \rightarrow \text{A} \xrightarrow{\text{H}_2\text{O}/\text{H}^+} \text{B}$$
  
The compound B is [KCET 1999]  
(a) Acetic acid (b) Acetone  
(c) Acetaldehyde (d) Ethyl alcohol
81.  $\text{CH}_3\text{CN}$  is known as acetonitrile because [AMU 1999]  
(a) It contains an aceto group  
(b) On hydrolysis it gives acetic acid  
(c) Both (a) and (b)  
(d) None of these
82. What is formed, when nitrobenzene is reduced using zinc and alkali [BHU 2000; AIIMS 2000; CBSE PMT 2000; MH CET 2003]  
(a) Phenol (b) Aniline  
(c) Nitrosobenzene (d) Hydrazobenzene
83.  $\text{RCOCl} + 2\text{Me}_2\text{NH} \rightarrow \text{A} + \text{Me}_2\text{NH}_2\text{Cl}^-$   
Here A is [RPET 2000]  
(a)  $\text{RCON} \begin{matrix} \text{Me} \\ \diagdown \\ \diagup \\ \text{Me} \end{matrix}$  (b)  $\text{RCONH}_2$   
(c)  $\text{RCONHMe}$  (d)  $(\text{RCO})_2\text{NH}$
84. Decreasing order of basicity is [RPET 2000]  
(1)  $\text{CH}_3\text{CONH}_2$  (2)  $\text{CH}_3\text{CH}_2\text{NH}_2$   
(3)  $\text{Ph}-\text{CH}_2\text{CONH}_2$   
(a)  $1 > 2 > 3$  (b)  $2 > 1 > 3$





## 1386 Nitrogen Containing Compounds

- (c)  $3 > 2 > 1$  (d) None of these
- 85.** Among the following, the strongest base is  
[UPSEAT 2000; IIT-JEE (Screening) 2000]  
(a)  $C_6H_5NH_2$  (b)  $p-NO_2C_6H_4NH_2$   
(c)  $m-NO_2-C_6H_4NH_2$  (d)  $C_6H_5CH_2NH_2$
- 86.** Aniline and methyl amine can be differentiated by  
[DPMT 2000]  
(a) Reaction with chloroform and aqueous solution of KOH  
(b) Diazotisation followed by coupling with phenol  
(c) Reaction with  $HNO_2$   
(d) None of these
- 87.** The amine which can react with  $C_6H_5-SO_2-Cl$  to form a product insoluble in alkali shall be [AMU 2000]  
(a) Primary amine  
(b) Secondary amine  
(c) Tertiary amine  
(d) Both primary and secondary amines
- 88.** A mixture of benzene and aniline can be separated by  
[KCET (Engg.) 2001]  
(a) Hot water (b) dil. HCl  
(c) dil. NaOH (d) Alcohol
- 89.** Nitrobenzene on further excessive nitration gives  
[AFMC 2001]  
(a) Trinitrobenzene (b)  $m$ -dinitrobenzene  
(c)  $p$ -dinitrobenzene (d) All of these
- 90.** The compound A with following sequence of reaction gave benzoic acid  
 $A \xrightarrow{NaNO_2 / HCl} B \xrightarrow{KCN} C \xrightarrow{H_3O^+} \text{benzoic acid}$ . The compound A is [AMU 2001]  
(a) Nitrobenzene (b) Aniline  
(c) Benzaldehyde (d) Amides
- 91.** Which of the following chemicals are used to manufacture methyl isocyanate that caused "Bhopal Tragedy"  
(i) Methylamine (ii) Phosgene  
(iii) Dimethylamine (iv) Phosphine  
[AIIMS 2005]  
(a) (i) and (iii) (b) (iii) and (iv)  
(c) (i) and (ii) (d) (ii) and (iv)
- 92.** An isocyanide on hydrolysis gives [AMU 2001]  
(a) An amide  
(b) A carboxylic acid and ammonia  
(c) A N-substituted amide  
(d) A  $1^\circ$ -amine and formic acid
- 93.** Methyl isocyanide on hydrolysis gives [UPSEAT 2001]  
(a)  $CH_3NH_2$  (b)  $HCOOH$   
(c)  $CH_3COOH$  (d) Both (a) and (b)
- 94.** Pure aniline is a [UPSEAT 2001]  
(a) Colourless solid  
(b) Brown coloured solid  
(c) Colourless liquid  
(d) Brown coloured liquid
- 95.** Reduction of methyl isocyanide gives [RPMT 2002]  
(a) Ethylamine (b) Methylamine  
(c) Dimethylamine (d) Trimethylamine
- 96.** Reaction of aniline with benzaldehyde is [RPMT 2002]  
(a) Polymerisation (b) Condensation  
(c) Addition (d) Substitution
- 97.** In the reaction  $C_6H_5CHO + C_6H_5NH_2 \rightarrow C_6H_5N = HCC_6H_5 + H_2O$ , the compound  $C_6H_5N = CHC_6H_5$  is known as [RPMT 2000; AIIMS 2002; AMU 2001]  
(a) Aldol (b) Schiff's reagent  
(c) Schiff's base (d) Benedict reagent
- 98.** The unshared pair of electrons on a cyanide ion can act as [Kerala (Med.) 2002]  
(a) Isocyanide centre (b) Amido centre  
(c) Cationic centre (d) Nucleophilic centre
- 99.** Electrophilic substitution of  with bromine gives [Kerala (Med.) 2002]  
(a) 1, 4, 6-tribromo aniline  
(b) 2, 4, 6-tribromo aniline  
(c) 4-bromo aniline  
(d) 3-bromo aniline
- 100.** Mustard gas is obtained by [MP PET 2002]  
(a) The action of dilute acids on mustard seeds  
(b) Treating ethylene with mustard oil  
(c) Treating sulphur chloride with ethylene  
(d) None of these
- 101.** Which of the following is capable of forming a zwitter ion [JIPMER 2002]  
(a)  $C_6H_5-OH$  (b)  $C_6H_4(NH_2)_2$   
(c)  $CH_2OH$  (d)  $H_2N-CH_2-COOH$   
|  
 $CH_2OH$
- 102.** Which one of the following reducing agents is likely to be the most effective in bringing about the following change  
 $R-\overset{O}{\parallel}C-NH_2 \rightarrow RCH_2NH_2$  [AMU 2002]  
(a)  $H_2-Ni$  (b)  $NaBH_4$   
(c)  $LiAlH_4$  (d) Na-alcohol
- 103.** During acetylation of amines what is replaced by acetyl groups [UPSEAT 2002]  
(a) Hydrogen atom attached to nitrogen atom  
(b) One or more hydrogen atoms attached to carbon atom  
(c) One or more hydrogen atoms attached to nitrogen atom

(d) Hydrogen atoms attached to either carbon atom or nitrogen atom

104. Hydrolysis of acetonitrile in acidic medium produces

[CPMT 2003; RPMT 2003]

- (a)  $CH_3CH_2OH$  (b)  $CH_3COOH$   
(c)  $CH_3NC$  (d)  $CH_3COOCH_3$

105. Which has a pyramidal structure [UPSEAT 2003]

- (a) Trimethylamine (b) Methanol  
(c) Acetylene (d) Water

106. Ethyl amine on acetylation gives [BHU 2002; BVP 2003]

- (a) N-ethyl acetamide  
(b) Acetamide  
(c) Methyl acetamide  
(d) None


107. The refluxing of  $(CH_3)_2NCOCH_3$  with acid gives

[BHU 2002; BVP 2003]




- (a)  $(CH_3)_2NH + CH_3COOH$   
(b)  $(CH_3)_2NCOOH + CH_4$   
(c)  $2CH_3OH + CH_3CONH_2$   
(d)  $2CH_3NH_2 + CH_3COOH$

108. *p*-chloro aniline and anilinium hydrogen chloride can be distinguished by [UPSEAT 2003]

- (a) Sandmaeyer reaction  
(b) Carbyl amine reaction  
(c) Hinsberg's reaction  
(d)  $AgNO_3$




Product 'A' in above reaction is

- (a)  (b)   
(c)  (d) None of these


110. Product obtained by electrolytic reduction of nitrobenzene in presence of  $H_2SO_4$  is [RPMT 2003]

- (a) *o*-amino phenol (b) *m*-amino phenol  
(c) *p*-amino phenol (d) None of these




The product 'B' is

[RPMT 2003]



112. Identify the product Z in the following reaction



[Kerala (Med.) 2003]

- (a) *p*-Bromoaniline (b) *p*-Bromoacetophenone  
(c) *o*-Bromoacetophenone (d) *o*-Bromoacetanilide


113. Benzaldehyde condenses with *N,N*-dimethylaniline in presence of anhydrous  $ZnCl_2$  to give [Kerala (Med.) 2003]

- (a) Michler's ketone (b) Azo dye  
(c) Malachite green (d) Buffer yellow


114. The correct order of reactivity towards the electrophilic substitution of the compounds aniline (I) benzene (II) and nitrobenzene (III) is [CBSE PMT 2003]

- (a) I > II > III (b) III > II > I  
(c) II > III > I (d) I < II > III

115. The final product C, obtained in this reaction



[CBSE PMT 2003]



116. The correct order of increasing basic nature for the bases  $NH_3$ ,  $CH_3NH_2$  and  $(CH_3)_2NH$  is [AIEEE 2003]

- (a)  $CH_3NH_2 < NH_3 < (CH_3)_2NH$   
(b)  $(CH_3)_2NH < NH_3 < CH_3NH_2$   
(c)  $NH_3 < CH_3NH_2 < (CH_3)_2NH$   
(d)  $CH_3NH_2 < (CH_3)_2NH < NH_3$

117. Nitrobenzene gives *N*-phenylhydroxylamine by



## 1388 Nitrogen Containing Compounds

[AIIMS 2003]

- (a)  $Sn/HCl$  (b)  $H_2/Pd-C$   
 (c)  $Zn/NaOH$  (d)  $Zn/NH_4Cl$

118. Among the following the weakest base is [AIIMS 2003]


- (a)  $C_6H_5CH_2NH_2$  (b)  $C_6H_5CH_2NHCH_3$   
 (c)  $O_2NCH_2NH_2$  (d)  $CH_3NHCHO$

119. The correct order of basicity of amines in water is :

[Pb. CET 2003]

- (a)  $(CH_3)_2NH > (CH_3)_3N > CH_3NH_2$   
 (b)  $CH_3NH_2 > (CH_3)_2NH > (CH_3)_3N$   
 (c)  $(CH_3)_3N > (CH_3)_2NH > CH_3NH_2$   
 (d)  $(CH_3)_3N > CH_3NH_2 > (CH_3)_2NH$

120. Complete the following reaction :



- (a)  $[R-NH_3]^+ HSO_4^-$  (b)  $[R-NH_3]_2^+ SO_4^{2-}$   
 (c)  $R-NH_2 \cdot H_2SO_4$  (d) No reaction

121. Which of the following compound reacts with chloroform and a base to form phenyl isocyanide ? [MH CET 2003]

- (a) Phenol (b) Aniline  
 (c) Benzene (d) Nitro benzene

122. Which one doesn't liberate  $NH_3$  when undergoes hydrolysis [Orissa JEE 2005]

- (a) Acetanilide (b) Acetonitrile  
 (c) Acetamide (d) Phenyl isocyanide

123. A nitrogen containing organic compound gave an oily liquid on heating with bromine and potassium hydroxide solution. On shaking the product with acetic anhydride, an antipyretic drug was obtained. The reactions indicate that the starting compound is [KCET 2004]

- (a) Aniline (b) Benzamide  
 (c) Acetamide (d) Nitrobenzene

124. Benzamide on reaction with  $POCl_3$  gives [IIT-JEE 2004]

- (a) Aniline (b) Chlorobenzene  
 (c) Benzyl amine (d) Benzonitrile

125. Among the following which one does not act as an intermediate in Hofmann rearrangement [AIIMS 2005]

- (a)  $RNCO$  (b)  $RCO\ddot{N}$   
 (c)  $RCO\ddot{N}HBr$  (d)  $RNC$

126. Aniline reacts with which of these to form Schiff base

[AFMC 2004]

- (a) Acetic acid (b) Benzaldehyde  
 (c) Acetone (d)  $NH_3$

127. Which of the following does not reduce Tollen's reagent


[Kerala PMT 2004]

- (a)  $CH_3CHO$  (b)  $C_6H_5NHOH$   
 (c)  $HCOOH$  (d)  $C_6H_5NO_2$

(e) None of these

128. Which one of the following compound is most basic ?

[UPSEAT 2004]




- (A) (a) (A) (b) (B)  
 (c) (C) (d) All are equally basic

129. Which one of the following methods is neither meant for the synthesis nor for separation of amines [AIEEE 2005]

- (a) Hinsberg method (b) Hofmann method  
 (c) Wurtz reaction (d) Curtius reaction

130. Aniline in a set of reactions yielded a product D.



The structure of product D would be [CBSE PMT 2005]

- (a)  $C_6H_5CH_2NH_2$  (b)  $C_6H_5NHCH_2CH_3$   
 (c)  $C_6H_5NHOH$  (d)  $C_6H_5CH_2OH$

131. Electrolytic reduction of nitrobenzene in weakly acidic medium gives [CBSE PMT 2005]

- (a) Aniline (b) Nitrosobenzene  
 (c) *N*-Phenylhydroxylamine (d) *p*-Hydroxylaniline

132. Among the following compounds  $C_3H_7NH_2$ ,  $NH_3$ ,  $CH_3NH_2$ ,  $C_2H_5NH_2$  and  $C_6H_5NH_2$ , the least basic compound is

- (a)  $C_3H_7NH_2$  (b)  $NH_3$   
 (c)  $CH_3NH_2$  (d)  $C_6H_5NH_2$   
 (e)  $C_2H_5NH_2$


133. The reduction of which of the following compound would yield secondary amine ?

- (a) Alkyl nitrite  
 (b) Carbylamine  
 (c) Primary amine  
 (d) Secondary nitro compound

134. Azo dye is prepared by the coupling of phenol and :

[Pb. CET 2000]

- (a) Diazonium chloride  
 (b) *o*-nitro aniline  
 (c) Benzoic acid  
 (d) Chlorobenzene



Z is identified as :

[Pb. PMT 2004]


- (a)  $C_6H_5-NH-CH_3$   
 (b)  $C_6H_5-COOH$   
 (c)  $C_6H_5-CH_2-NH_2$

(d)  $C_6H_5-CH_2-COOH$

136. When acetamide reacts with  $Br_2$  and caustic soda, then we get :

- (a) Acetic acid                      (b) Bromoacetic acid  
(c) Methyl amine                    (d) Ethyl amine

137. In the reaction



the term  $Y$  is: [BHU 2004]

- (a) Acetone                              (b) Ethyl amine  
(c) Acetaldehyde                      (d) Dimethyl amine

138. Reaction of cyclohexanone with dimethylamine in the presence of catalytic amount of an acid forms a compounds if water during the reaction is continuously removed. The compound formed is generally known as [AIIEE 2005]

- (a) A Schiff's base                    (b) An enamine  
(c) An imine                            (d) An amine

139.  $R-NH-COH \xrightarrow[\text{pyridine}]{POCl_3}$  product

In the given reaction what will be the product [BHU 2005]

- (a)  $R-N=C=O$                       (b)  $R-\overset{+}{N} \equiv C^-$   
(c)  $R-C \equiv N$                         (d) None of these.

140. Which of the following is secondary pollutant.


[BHU 2005]

- (a)  $CO_2$                                 (b)  $N_2O$   
(c) PAN                                  (d)  $SO_2$

141. Nitration of aniline also gives  $m$ -nitro aniline, in strong acidic medium because

- (a) In electrophilic substitution reaction amino group is meta directive  
(b) In spite of substituents nitro group always goes to  $m$ - position  
(c) In strong acidic medium, nitration of aniline is a nucleophilic substitution reaction  
(d) In strong acidic medium aniline present as anilinium ion  
(e) Strong acid, gives nitrate anion, which attacks at  $m$ -position

142. Identify the product in following order



- (a) 3, 4,5 -Tribromobenzene  
(b) 1, 2, 3 - Tribromobenzene  
(c) 2, 4, 6 - Tribromobenzene  
(d) 3, 4, 5 - Tribromo nitro benzene  
(e) 3, 4, 5 - Tribromo phenol

143. The correct order of basicity in amines

- (i)  $C_4H_5NH_2$                       (ii)  $CH_3NH_2$   
(iii)  $(CH_3)_2NH$                     (iv)  $(CH_3)_3N$

[Kerala CET 2005]

- (a) (i) < (iv) < (ii) < (iii) (b) (iv) < (iii) < (ii) < (i)  
(c) (i) < (ii) < (iii) < (iv) (d) (ii) < (iii) < (iv) < (i)  
(e) (iv) < (iii) < (ii) < (i)

### Tests for Nitrogen Containing Compounds

[CPMT 2004]

1. When acetamide reacts with  $Br_2$  and caustic soda, then we get

[DPMT 1983; BHU 1997; Orissa JEE 2002;

CPMT 1971, 78, 79, 81, 85, 2000, 03;

MP PMT 1989; MP PET 1995, 2002]

- (a) Acetic acid                      (b) Bromoacetic acid  
(c) Methyl amine                    (d) Ethane

2. In organic compounds, nitrogen is tested in Lassaigne's test as

- (a)  $NaNH_2$                             (b)  $NaCN$   
(c)  $NaNO_2$                         (d)  $NaNO_3$

3. Liebermann's nitroso reaction is used for testing

- (a) Primary amines                  (b) Secondary amines  
(c) Tertiary amines                  (d) All the above

4. A nauseating smell in the carbylamine test for primary amines is due to the formation of [MP PET 1993]

- (a) Isocyanide                        (b) Chloroform  
(c) Cyanide                            (d) DDT

5. A positive carbylamine test is given by [IIT-JEE 1999]

- (a)  $N, N$ -dimethylaniline  
(b) 2, 4-dimethylaniline  
(c)  $N$ -methyl- $o$ -methylaniline  
(d)  $p$ -methylbenzylamine

6. The colour of  $p$ -amino azobenzene is [BHU 1997]

- (a) Orange [Kerala CET 2005]      (b) Congo red  
(c) Bismark brown                    (d) Indigo

7. When primary amine is heated with  $CS_2$  in presence of excess mercuric chloride, it gives isothiocyanate. This reaction is called [KCET 1998; CPMT 1999]

- (a) Hofmann bromide reaction  
(b) Hofmann mustard oil reaction  
(c) Carbylamine reaction  
(d) Perkin reaction

8. Diazo-coupling is useful to prepare some [CPMT 1999]

- (a) Dyes                                (b) Proteins  
(c) Pesticides                        (d) Vitamins

9. Carbylamine test is used in the detection of [DCE 1999]

- (a) Aliphatic  $2^\circ$  amine  
(b) Aromatic  $1^\circ$  amine  
(c) Aliphatic  $1^\circ$  amine  
(d) Both aliphatic and aromatic  $1^\circ$  amines

10. Which of the following substance does not give iodoform test

- (a)  $C_6H_5CN$                         (b)  $RNH_2$   
(c)  $CH_3OH$                             (d) All

## 1390 Nitrogen Containing Compounds

11. Which one of the following compounds when heated with  $KOH$  and a primary amine gives carbylamine test


[Orissa JEE 2005]

- (a)  $CHCl_3$  (b)  $CH_3Cl$   
(c)  $CH_3OH$  (d)  $CH_3CN$

## Critical Thinking

### Objective Questions


1. The compound



forms nitroso amines when the substituents are




[Roorkee 1999]

- (a)  $R_1 = CH_3, R_2 = R_3 = H$   
(b)  $R_1 = R_2 = H, R_3 = C_2H_5$   
(c)  $R_1 = H, R_2 = R_3 = CH_3$   
(d)  $R_1 = CH_3, R_2 = C_2H_5, R_3 = H$
2. The action of nitrous acid on ethyl amine gives  
[DPMT 1982; CPMT 1971, 89, 94; MP PET 1993, 2001; RPMT 1997; Pb. PMT 1999]
- (a) Ethane (b) Ammonia  
(c) Ethyl alcohol (d) Nitroethane
3. Aniline when diazotized in cold and then treated with dimethyl aniline gives a coloured product. Its structure would be  
[CBSE PMT 2004]







4. Indicate which nitrogen compound amongst the following would undergo Hofmann's reaction (i.e. reaction with  $Br_2$  and strong  $KOH$ ) to furnish the primary amine ( $R-NH_2$ )



[CBSE PMT 1989]



5.   
[BHU 1995]




6. The correct order of basicities of the following compounds is



[IIT-JEE (Screening) 2001]


- (a)  $2 > 1 > 3 > 4$  (b)  $1 > 3 > 2 > 4$   
(c)  $3 > 1 > 2 > 4$  (d)  $1 > 2 > 3 > 4$
7. Which of the following would be most reactive towards nitration  
[AMU 2000; UPSEAT 2002]
- (a) Benzene (b) Nitro benzene  
(c) Toluene (d) Chloro benzene
8. Aniline reacts with acetaldehyde to form  
[MHCET 2004; AFMC 2004]
- (a) Schiff's base (b) Carbylamine  
(c) Imine (d) None of these
9. *p*-chloroaniline and anilinium hydrochloride can be distinguished by  
[IIT-JEE 1998]
- (a) Sandmeyer reaction (b)  $NaHCO_3$   
(c)  $AgNO_3$  (d) Carbylamine test



In the above reaction 'X' stands for

[CPMT 1986, 2001; MP PET 1992; KCET (Engg./Med.) 2000]

- (a)  $NH_2$  (b)  $SnCl_2$   
(c)  $Cl$  (d)  $NH_4^+Cl^-$



[BHU 2000; Pb. PMT 2000; Kerala 2003]



- (a) Phenyl isocyanide (b) Benzyl amine  
(c) Benzyl chloride (d) None of these
12. The order of basic strength among the following amines in benzene solution is [AIIMS 1991; RPMT 2002]
- (a)  $CH_3NH_2 > (CH_3)_3N > (CH_3)_2NH$   
(b)  $(CH_3)_2NH > CH_3NH_2 > (CH_3)_3N$   
(c)  $CH_3NH_2 > (CH_3)_2NH > (CH_3)_3N$   
(d)  $(CH_3)_3N > CH_3NH_2 > (CH_3)_2NH$


13. The refluxing of  $(CH_3)_2NCOCH_3$  with acid gives [KCET 1996]

- (a)  $2CH_3NH_2 + CH_3COOH$   
(b)  $2CH_3OH + CH_3COOH$   
(c)  $(CH_3)_2NH + CH_3COOH$   
(d)  $(CH_3)_2NCOOH + CH_4$

14. Order of basicity of ethyl amines is [MP PMT/PET 1988]

- (a) Secondary > Primary > Tertiary  
(b) Primary > Secondary > Tertiary  
(c) Secondary > Tertiary > Primary  
(d) Tertiary > Primary > Secondary

15. The following reaction is [KCET 1996]




- (a) Nucleophilic substitution  
(b) Electrophilic substitution  
(c) Free radical substitution  
(d) None of these
16.  $RNH_2$  reacts with  $C_6H_5SO_2Cl$  in aqueous  $KOH$  to give a clear solution. On acidification a precipitate is obtained which is due to the formation of [Roorkee 2000]

- (a)  $R-\overset{\overset{H}{|}}{N^+}-SO_2C_6H_5OH^-$   
(b)  $R-N^-SO_2C_6H_5K^+$   
(c)  $R-NHSO_2C_6H_5$   
(d)  $C_6H_5SO_2NH_2$



17. If  $N$  and  $S$  are present in an organic compound during Lassaigne test, then both changes into [CPMT 1997]

- (a)  $Na_2S$  and  $NaCN$   
(b)  $NaSCN$   
(c)  $Na_2SO_3$  and  $NaCN$   
(d)  $Na_2S$  and  $NaCNO$

18. The strongest base among the following is [AIIMS 2004; BHU 2004]



- (a) (b)







19. Nitroso amines ( $R_2N-N=O$ ) are soluble in water. On heating them with concentrated  $H_2SO_4$  they give secondary amines. The reaction is called [AFMC 1998; AIIMS]

- (a) Perkin's reaction  
(b) Fittig's reaction  
(c) Sandmeyer's reaction  
(d) Liebermann's nitroso reaction





20. A primary amine is formed an amide by the treatment of bromine and alkali. The primary amine has : [BHU 2004]

- (a) 1 carbon atom less than amide  
(b) 1 carbon atom more than amide  
(c) 1 hydrogen atom less than amide  
(d) 1 hydrogen atom more than amide

21. The structural formula of Indigo dye is : [DPMT 2004]







- Which of the following is the strongest base ? [AIEEE 2004]



## 1392 Nitrogen Containing Compounds

23. The following compound on hydrolysis in aqueous acetone will give



- (a) Mixture of (K) and (L)    (b) Mixture of (K) and (M)  
(c) Only (M)    (d) Only (K)

## Assertion & Reason

For AIIMS Aspirants

Read the assertion and reason carefully to mark the correct option out of the options given below :

- (a) If both assertion and reason are true and the reason is the correct explanation of the assertion.  
(b) If both assertion and reason are true but reason is not the correct explanation of the assertion.  
(c) If assertion is true but reason is false.  
(d) If the assertion and reason both are false.  
(e) If assertion is false but reason is true.

1. Assertion : Benzene diazonium chloride does not give tests for nitrogen.  
Reason :  $N_2$  gas loss takes place during heating  
[AIIMS 1999]
2. Assertion : Amines are basic in nature.  
Reason : Presence of lone pair of electron on nitrogen atom. [AIIMS 1999]

3. Assertion : Methyl isocyanide reacts with ozone to form methyl isocyanate.  
Reason : Methyl isocyanate was responsible for Bhopal tragedy.
4. Assertion : Alkyl cyanide can be prepared by carbylamine reaction  
Reason : Ethyl amine when heated with chloroform in presence of alcoholic KOH, cyanide is formed.
5. Assertion :  $CN^-$  ion is an ambident nucleophile.  
Reason : Nucleophiles are electron rich species.
6. Assertion : Sulphanilic acid exists as dipolar ion whereas *p*-aminobenzoic acid does not.  
Reason : Carboxyl group being more acidic than  $-SO_3H$  group can easily transfer a  $H^+$  to the amino group.
7. Assertion : Nitrating mixture used for carrying out nitration of benzene consists of conc.  $HNO_3$  + conc.  $H_2SO_4$ .  
Reason : In presence of  $H_2SO_4$ ,  $HNO_3$  acts as a base and produces  $NO_2^+$  ions.
8. Assertion : In order to convert  $R-Cl$  to pure  $R-NH_2$ , Gabriel phthalimide synthesis can be used.  
Reason : With proper choice of alkyl halides, phthalimide synthesis can be used to prepare  $1^\circ$ ,  $2^\circ$  or  $3^\circ$  amines.
9. Assertion : Ammonolysis of alkyl halides involves the reaction between alkyl halides and alcoholic ammonia.  
Reason : Reaction can be used to prepare only  $2^\circ$  amines.
10. Assertion : Nitroalkanes, but not nitroarenes can be distilled at normal atmospheric pressure.  
Reason : Nitroalkanes are sparingly soluble in water while nitroarenes are insoluble.
11. Assertion : In Hofmann bromide reaction, the amine formed has one carbon atom less than the parent  $1^\circ$  amide.  
Reason : *N*-methyl acetamide undergoes Hofmann bromamide reaction.
12. Assertion : Nitrobenzene does not undergo Friedel Craft alkylation.  
Reason : Nitrobenzene is used as solvent in laboratory and industry.
13. Assertion : Ammonia is less basic than water.  
Reason : Nitrogen is less electronegative than oxygen.
14. Assertion : The reaction between a diazo salt and an aromatic amine or a phenol,

## Nitrogen Containing Compounds 1393

giving an aminoazo or hydroxyazo compounds is called coupling reaction.

Reason : Condensation of diazonium salt with phenol is carried out in weakly acidic medium.

15. Assertion : Carbylamine reaction involves the reaction between  $1^\circ$  amine and chloroform in basic medium.

Reason : In carbylamine reaction,  $-NH_2$  group is converted into  $-NC$  group.

16. Assertion :  $Me_3N$  reacts with  $BF_3$  whereas  $Ph_3N$  does not.

Reason : The electron pair on nitrogen atom in  $Ph_3N$  is delocalised in the benzene ring and is not available to boron in  $BF_3$ .

17. Assertion : *p*-Anisidine is weaker base than aniline.

Reason :  $-OCH_3$  group in anisidine exerts  $-R$  effect.

18. Assertion : Lower aldehydes and ketones are soluble in water but the solubility decreases as the molecular mass increases.

Reason : Distinction between aldehydes and ketones can be made by Tollen's test.

[AIIMS 1999]

19. Assertion : Aniline hydrogen sulphate on heating forms a mixture of ortho and para aminobenzene sulphonic acids.

Reason : The sulphonic acid group is electron withdrawing. [AIIMS 1996]

20. Assertion :  $p-O_2N-C_6H_5COCH_3$  is prepared by Friedel Crafts acylation of nitrobenzene.

Reason : Nitrobenzene easily undergoes electrophilic substitution reaction. [AIIMS 2005]

21. Assertion : Alkyl isocyanides in acidified water give alkyl formamides.

Reason : In isocyanides, carbon first acts as a nucleophile and then as an electrophile.

[AIIMS 2005]

1	a	2	c	3	a	4	d	5	c
6	d	7	c	8	d	9	c	10	a
11	a	12	b	13	b	14	a	15	b

### Preparation of Nitrogen Containing Compounds

1	c	2	d	3	b	4	b	5	b
6	b	7	c	8	a	9	c	10	a
11	d	12	d	13	b	14	b	15	d
16	c	17	a	18	b	19	a	20	c
21	b	22	a	23	a	24	a	25	b
26	b	27	c	28	d	29	c	30	a
31	a	32	a	33	a	34	d	35	c
36	c	37	b	38	a	39	d	40	b,c
41	d	42	b	43	c	44	c	45	c
46	b	47	b	48	a	49	c	50	a
51	c	52	a	53	c	54	a	55	b

### Properties of Nitrogen Containing Compounds

1	d	2	d	3	b	4	b	5	a
6	c	7	c	8	d	9	b	10	c
11	c	12	b	13	b	14	c	15	c
16	a	17	d	18	a	19	b	20	c
21	a	22	d	23	b	24	c	25	a
26	b	27	b	28	c	29	b	30	b
31	d	32	a	33	c	34	a	35	b
36	c	37	b	38	b	39	d	40	c
41	a	42	c	43	b	44	c	45	d
46	d	47	d	48	b	49	d	50	b
51	d	52	d	53	c	54	c	55	c
56	d	57	d	58	c	59	c	60	c
61	b	62	a	63	c	64	a	65	b
66	c	67	e	68	c	69	a	70	c
71	d	72	c	73	a	74	b	75	b
76	a	77	a	78	b	79	c	80	b
81	b	82	d	83	a	84	b	85	d
86	b	87	b	88	b	89	b	90	b
91	c	92	d	93	d	94	c	95	c
96	b	97	c	98	c	99	b	100	c
101	d	102	a	103	c	104	b	105	a
106	a	107	a	108	d	109	b	110	c
111	c	112	a	113	c	114	a	115	d
116	c	117	d	118	b	119	a	120	b
121	b	122	d	123	b	124	d	125	d
126	b	127	d	128	b	129	c	130	d

# Answers

## Introduction of Nitrogen Containing Compounds



## 1394 Nitrogen Containing Compounds

131	a	132	d	133	b	134	a	135	b
136	c	137	c	138	b	139	b	140	c
141	d	142	b	143	a				

## Tests for Nitrogen Containing Compounds

1	c	2	b	3	b	4	a	5	b
6	a	7	b	8	a	9	d	10	d
11	a								

## Critical Thinking Questions

1	c	2	c	3	a	4	c	5	b
6	b	7	b	8	a	9	bc	10	a
11	a	12	b	13	c	14	a	15	a
16	c	17	b	18	c	19	d	20	a
21	c	22	d	23	a				



Assertion & Reason

1	a	2	a	3	b	4	d	5	b
6	c	7	a	8	c	9	c	10	b
11	c	12	b	13	e	14	c	15	a
16	a	17	d	18	b	19	d	20	d
21	a								


# AS Answers and Solutions

## Introduction of Nitrogen Containing Compounds



4. (d)  $C_3H_9N$  can form all the 3 amines.  
 $CH_3CH_2CH_2-NH_2$ ,  $CH_3-CH_2-NH-CH_3$   
 $1^\circ$  amine  $2^\circ$  amine
- $CH_3-\overset{\text{N}}{\underset{\text{CH}_3}{|}}-CH_3$   
 $3^\circ$  amine
5. (c)  $(CH_3)_2C=O + H.CH_2-COCH_3 + NH_3$   
 $(CH_3)_2C-\underset{\text{NH}_2}{\text{CH}_2}-COCH_3$   
 diacetone amine
8. (d) Allyl isocyanide.  $CH=CH-CH_2-N\equiv C$
9. (c)
- 
12. (b)  $CH_3CH_2-O-N=O$  is a nitrite derivative, hence it is not a nitro derivative.
13. (b)  $CH_3CN$  is called acetonitrile....
15. (b) Four  $1^\circ$  amines are possible  
 $CH_3CH_2CH_2CH_2NH_2$ ,  $(CH_3)_2CHCH_2NH_2$ ,  
 $CH_3CH(NH_2)CH_2CH_3$ ,  $(CH_3)_3CNH_2$

## Preparation of Nitrogen Containing Compounds








1. (c) Hofmann's bromamide reaction  
 $CH_3-CO-NH_2 + Br_2 + 4KOH \xrightarrow{H_2O}$   
 Acetamide
- $CH_3NH_2 + K_2CO_3 + 2KBr + 2H_2O$   
 Methyl amine
2. (d)  $CH_3CONH_2 \xrightarrow{NaOBr} CH_3NH_2$ .
3. (b)  $CH_3CONH_2 + Br_2 + 4NaOH \rightarrow$   
 Acetamide










4. (b)  $CH_3-CO-NH_2 \xrightarrow[2c]{NaOH / Br_2} CH_3-NH_2$   
 Hofmann's bromamide (1c)
5. (b)  $CH_3C\equiv N + 4[H] \xrightarrow{\text{Reduction}} CH_3CH_2NH_2$   
 Reduction
6. (b)  $CH_3-CH_2-CO-NH_2 + Br_2 + 4KOH \rightarrow$   
 Propionamide  
 $CH_3CH_2NH_2 + K_2CO_3 + 2KBr + 2H_2O$
7. (c)  $C_2H_5I + NH_3 \rightarrow HI + C_2H_5-NH_2$   
 $C_2H_5OH + NH_3 \rightarrow H_2O + C_2H_5-NH_2$
9. (c)  $CH_3-CH_2-NO_2 + 6[H] \xrightarrow{Sn / HCl}$   
 Nitro ethane  
 $CH_3-CH_2-NH_2 + 2H_2O$   
 Ethyl amine
11. (d)  $CH_3I \xrightarrow{\Delta} CH_3NH_2 \xrightarrow{CH_3I} (CH_3)_2NH$   
 Methyl amine Dimethyl amine  
 $\xrightarrow{CH_3I} (CH_3)_3N$   
 Trimethyl amine
12. (d)
- 
- Aniline + Acetic anhydride  $\rightarrow$  Acetanilide +  $CH_3COOH$
16. (c)  $CH_3-N\equiv C + 4[H] \xrightarrow{\text{Reduction}} CH_3-NH-CH_3$   
 $2^\circ$  amine
17. (a)  $CH_3NO_2 + 3Cl_2 + 3NaOH \rightarrow$   
 Nitromethane  
 $CCl_3-NO_2 + 3NaCl + 3H_2O$   
 (chloropicrin)
18. (b)
- 
19. (a)  $C_2H_5OH + NH_3 \xrightarrow{\text{alumina}} C_2H_5NH_2 + H_2O$ .
20. (c)  $R-CN + H_2O \xrightarrow{H_2O / H^+} RCOOH + NH_3$   
 It yield amine when reduced as -  
 $R-CN + H_2 \rightarrow R-CH_2-NH_2$
21. (b)  $CH_3CONH_2 \xrightarrow{P_2O_5} CH_3CN \xrightarrow{4H} CH_3CH_2NH_2$
22. (a)  $CH_3-CH_2-\overset{O}{\parallel}{N} \rightarrow O + 3H_2 \rightarrow CH_3CH_2NH_2 + 2H_2O$ .
23. (a)  $CH_3NH_2 \xrightarrow{NO_2} CH_3OH + N_2 + H_2O$   
 Methyl amine
24. (a)  $R-NH_2 + CHCl_3 + 3KOH \rightarrow R-NC + 3KCl + 3HO$   
 $1^\circ$ -amine






## 1394 Nitrogen Containing Compounds




35. (c) Methyl amine is the strongest base.





50. (a) Halogen have  $-I$  and  $+M$  effect by which its electron delocalized in benzene ring by resonance & due to its  $-I$  effect its bonded with benzene ring and cannot be substitute by  $CN^-$  & show the inertness against  $KCN$  while other option gives Aromatic nitrile





51. (c) Secondary amines gives oily nitrosamine with nitrous acid.



52. (a) When aniline is treated with  $HNO_2$  at  $0-5^\circ C$  then diazonium salt is formed and by the coupling of diazonium salt and phenol azo dyes are prepared.




54. (a)  $p$ -nitrobenzene from  $p$ -nitroaniline.




## Properties of Nitrogen Containing Compounds








- (d) Tertiary amine does not react with nitrous acid because in it  $\alpha$ -H atom is absent.
- (d) Due to  $+ve$  I.E. of alkyl group,  $N$ -atom of amines acquires partial  $-ve$  charge and thus electron pair is easily donated.







The compounds in which both amino ( $-NH_2$ ) as well as acidic ( $-COOH$ ) group is present is called amino acid.





Presence of  $-NO_2$  group decreases electron density at  $o$ - and  $p$ - positions. Hence, incoming electrophile goes to  $m$  position. Therefore it is  $m$ -directing group.






12. (b) Only primary aromatic amines can undergo diazotisation.



$-NO_2$  group is meta directing group.




Amines are basic in nature they react with acid to form salt.



20. (c) Because in tertiary nitroalkanes  $\alpha-H$  atom is absent.

21. (a) Primary amine reacts with  $CHCl_3$  and alc.  $KOH$  to form isocyanide while secondary and tertiary amines do not react.




22. (d) Friedel-craft's reaction is used for the preparation of alkyl benzene or acetophenone. It is not a method to prepare amine.












This litmus paper test shows basic nature of amine.


24. (c) Presence of alkyl group increases electron density on nitrogen atom due to +I effect. Thus basic nature increases.

25. (a) Mustard oil reaction









3- nitro group can be introduced.











### 1396 Nitrogen Containing Compounds



50. (b) Because the N atom in aniline has a lone pair to donate and also due to +I effect of  $-NH_2$  group.



Nitration of aniline without protecting the amino group is not possible because  $HNO_3$  is a strong oxidising agent which oxidises aniline.






In methyl amine only one electron releasing group is present but in dimethyl amine two electron releasing groups are present which increase the basicity higher in diethyl amine.

57. (d) Nitro compounds are not explosive but stable compound.





60. (c)  $R_3N + HONO \rightarrow R_3N \cdot HONO$  called as Quaternary ammonium salt.






63. (c) 3° amine cannot be Acetylated because replaceable H-atom is absent.


67. (e) Because N-N dimethyl propanimine



68. (c) Replaceable  $H^-$  is absent.



73. (a)  $R_2NH > RNH_2 > R_3N > NH_3$ .




76. (a) Because of presence of electron withdrawing group -NO<sub>2</sub>.

77. (a) To suppress the concentration of the aniline available for coupling other rise coupling occurs.

78. (b)  $R-NH_2 + HNO_2 \rightarrow R-OH + N_2 + H_2O$   
alcohol

79. (c)  $C_6H_5NH_2 < (CH_3)_3N < CH_3NH_2 < (CH_3)_2NH$

80. (b)  $CH_3CN + CH_3MgI \rightarrow (CH_3)_2CNMgI \xrightarrow[-NH_3]{H_2O / H^+}$



82. (d)  $C_6H_5-NO_2 \xrightarrow[Zn / NaOH]{10[H]} C_6H_5NH-NHC_6H_5 + 4H_2O$   
Hydrazo benzene


83. (a)  $RCOCl + 2Me_2NH \rightarrow RCON \begin{matrix} < \\ < \end{matrix} \begin{matrix} Me \\ Me \end{matrix} + Me_2 + NH_2Cl^-$

Me = Methyl.


86. (b) Phenol react with aniline to give diazonium salt by coupling but Methyl amine not react with phenol.

87. (b)  $C_6H_5SO_2Cl$  is called Hinsberg's reagent they react with sec amine to form a product in soluble in alkalis. This reaction used to separate 1<sup>o</sup>, 2<sup>o</sup> and 3<sup>o</sup> amine from their mixture.


88. (b) A mixture of benzene and aniline can be separated by dil. HCl.



90. (b)




91. (c) Methyl isocyanate is industrially prepared by the action of methyl amine with phosgene.



92. (d)  $R-NC + 2H_2O \rightarrow RNH_2 + HCOOH$ .

93. (d)  $CH_3-NC + 2H_2O \rightarrow CH_3NH_2 + HCOOH$

95. (c)  $CH_3NC + 4H \xrightarrow[ether]{LiAlH_4} (CH_3)_2NH$ .





102. (a)  $R-\overset{O}{\parallel}C-NH_2 \xrightarrow{H_2-Ni} R-CH_2-NH_2$

104. (b)  $CH_3CN + 2H_2O \xrightarrow{HCl} CH_3COOH + NH_3$



106. (a)  $CH_3CH_2NH_2 + CH_3COCl \rightarrow CH_3CH_2NHCOCH_3 + HCl$   
N-Ethylacetanilide

107. (a)  $\begin{matrix} CH_3 \\ | \\ CH_3 \end{matrix} > NH + CH_3COOH \rightarrow \begin{matrix} CH_3 \\ | \\ CH_3 \end{matrix} > N-\overset{O}{\parallel}C-CH_3$

108. (d) Anilinium hydrogen chloride produces chloride ion which gives white precipitate with AgNO<sub>3</sub>. In fact anilium chloride is a part of aniline.




Nitrobenzene                      Phenyl hydroxyl amine                      p-amino phenol




119. (a) Basicity of amines increase with increase in number of -CH<sub>3</sub> groups (or any group which cause +I effect), due to increase in electron density on N atom. As a rule, the basicity of t-amine should be more than that of s-amine, but actually it is found to be lesser than s-amines. This is due to steric hinderence of bulkier alkyl groups, which decreases the availability of lone pair of electron on the N atom of the amino group. Hence the correct order of basicity is :


## 1398 Nitrogen Containing Compounds



120. (b) Amines are basic in nature, hence form salts with acid.










121. (b) We know that




Thus in this reaction phenyl isocyanide is produced. this is called carbylamine reaction.






122. (d) Isocyanides on hydrolysis forms primary amines not ammonia




126. (b) Aniline reacts with benzaldehyde and forms Schiff's base (benzal aniline) or anils.



Benzylidene aniline




132. (d)  $C_6H_5NH_2$  is least basic compound due to resonance by which the Lone pair of nitrogen takes part in resonance & due to unavailability








of lone pair on N Aniline become less basic. The Lone pair of N is delocalized into benzene ring by resonance

133. (b) Carbylamine (or isocyanides) give secondary amine on reduction.







134. (a) Azo dye is prepared by the coupling of phenol and diazonium chloride.




Thus product Z is identified as  $C_6H_5 - COOH$

136. (c) This is Hofman-bromide reaction. In this reaction one carbon less amines are formed from amides.




139. (b) N-alkyl formamides when dehydrated with  $POCl_3$  in presence of pyridine give isocyanides.

140. (c) Pollutants which are formed by reaction amongst the primary pollutants (persist in the environment in the form they are passed into it) are called as secondary pollutants. e.g. peroxyacyl nitrates (PAN) are formed through reaction between nitrogen oxides and hydrocarbons in the presence of sunlight.






The reason for this is that, in acidic condition protonation of  $-\text{NH}_2$  group gives anilinium ion ( $+\text{NH}_3$ ), which is of deactivating nature and of *m*-directive nature.




143. (a) Basicity order is
- $C_4H_5NH_2 < (CH_3)_3N < CH_3NH_2 < (CH_3)_2NH$
- 
- $(CH_3)_3N$
- is less basic due to steric effect while
- $C_4H_5NH_2$
- is less basic due to resonance.


### Tests for Nitrogen Containing Compounds




5. (b)
- 
- is a type of
- $1^\circ$
- amine and hence gives +ve




8. (a) Diazo-coupling is useful to prepare some dyes.



11. (a)
- $CHCl_3$
- gives carbylamine test.





1. (c)
- $R_1 = H$
- and
- $R_2 = R_3 = CH_3$





Sec. amine reacts with Nitrous acid to form nitroso amine yellow liquid.

2. (c)
- $CH_3CH_2NH_2 + HNO_2 \rightarrow CH_3CH_2OH + N_2 + H_2O$
- 
- Ethylamine Ethylalcohol




4. (c) Hofmann degradation of amide




6. (b) The relative basic character of
- $1^\circ$
- ,
- $2^\circ$
- and
- $3^\circ$
- amines also depends upon the nature of the alkyl group.

R	Relative basic strength
$-CH_3$	$R_2NH > R-NH_2 > R_3N > NH_3$
$-C_2H_5$	$R_2NH > R-NH_2 > NH_3 > R_3N$
$-CHMe_2$	$R-NH_2 > NH_3 > R_2NH > R_3N$
$-CMe_3$	$NH_3 > R-NH_2 > R_2NH > R_3N$

7. (b) The nitrogroup is very firmly linked to the benzene nucleus and does not undergo any displacement reaction. Nitro group deactivates the benzene nucleus.



9. (bc) (1) With
- $\text{NaHCO}_3 \rightarrow$




Anilinium hydrochloride is an acid salt and liberates  $\text{CO}_2$  from  $\text{NaHCO}_3$ .

### Critical Thinking Questions







## 1400 Nitrogen Containing Compounds


But p-chloro aniline is basic not acidic it does not liberate  $CO_2$ .




p-chloro aniline does not contain ionic chlorine to it does not give white ppt with  $AgNO_3$




14. (a) Order of basicity of amines  
(i)  $2^\circ > 1^\circ > 3^\circ$   
(ii)  $R_2NH > RNH_2 > ArCH_2-NH_2 > NH_3 > ArNH-R > ArNH_2 > ArNH-Ar$




Because  $OH^-$  is nucleophile.




17. (b) When sulphur and nitrogen both are present in organic compound during Lassaigne's Test, both changes into "sodium thiocyanate". ( $NaSCN$ ) which gives a blood red colouration with Ferric ion.




18. (c)  does not have aromaticity by which the Lone pair of electron of Nitrogen does not delocalised in benzene ring so it will be strong base on other hand rest 3 have aromaticity i.e., they follow the huckel rule so the electron pair of Nitrogen delocalised in ring by resonance &

19. (d) Liebermann's Nitroso reaction.




21. (c) The class of Indigo dye is Indigoid or vat dye. Indigo dyes are insoluble in water.  
22. (d) Lone pair of  $\ddot{N}$  are not taking part in conjugation whereas in other parts lone pairs are taking part in conjugation



## Assertion and Reason

- (a) It is true that benzene diazonium chloride does not respond Lassaigne test of nitrogen because benzene diazonium chloride losses  $N_2$  on Slight heat and thus it can't react with sodium metal.
- (a) Amines are basic due to the presence of a lone pair of electrons on nitrogen atom. The lone pair can be easily donated.
- (b) The reason being that the terminal carbon atom in isocyanide has electron-deficient carbon having a sextet of electrons and hence undergoes addition reactions with ozone.
- (d) When primary amines are heated with chloroform in the presence of alcoholic  $KOH$ , isocyanides are formed. This reaction is known as carbylamine reaction. eg. ethyl amine gives ethyl isocyanide on treatment with  $CHCl_3$  and alcoholic  $KOH$ .



5. (b) Nucleophilic species which have more than one site of reaction are called ambident nucleophiles.  
 $:\bar{C} = N : \leftrightarrow : C = \bar{N} :$
6. (c)  $-SO_3H$  group being more acidic than  $-CO_2H$  group can easily transfer a proton to the amino group.
7. (a)  $HNO_3 + 2H_2SO_4 \rightleftharpoons 2HSO_4^- + NO_2^+ + H_3O^+$
8. (c) Only primary aliphatic amines can be prepared by Gabriel phthalimide reaction.
9. (c) Reaction can be used to prepare  $1^\circ$ ,  $2^\circ$ ,  $3^\circ$  amines and finally quarternary ammonium salts.
10. (b) Nitroarene cannot be distilled under normal atmospheric pressure. This is because either they decompose or they explode on strong heating.
11. (c) Only  $1^\circ$  amines undergo Hofmann bromamide reaction. Since  $CH_3CONHCH_3$  is a  $2^\circ$  amine therefore, it does not undergo Hofmann bromamide reaction.
12. (b) Nitrobenzene does not undergo Friedel Craft reaction because nitro group deactivate the ring towards electrophilic substitution and drastic conditions are needed to carry out the electrophilic substitution reactions.
13. (e) Ammonia is more basic than water. It is because nitrogen being less electronegative than oxygen has a greater tendency to donate electrons.
14. (c) Condensation of diazonium salt with phenol is carried out in weakly alkaline medium, ( $pH = 9$ ). This is due to the fact that in strongly acid medium the  $-OH$  of a phenol remains unionised, and an amine forms a salt. Phenol exists as phenoxide ion and the latter is readily substituted by electrophiles than phenol itself. Thus, in phenol, coupling is carried out in alkaline medium.
15. (a)  $RNH_2 + CHCl_3 + 3KOH (alc) \rightarrow R-N \equiv C + 3KCl + 3H_2O$
17. (d) *p*-Anisidine is a stronger base than aniline.  
 $-OCH_3$  group in anisidine exerts +R- effect.
18. (b) Solubility of aldehydes and ketones decrease as the molecular mass increase.
20. (d) The nitro group strongly deactivates the benzene ring towards electrophilic substitution.  
 Nitrobenzene does not undergo Friedel-Craft acylation reaction.
21. (a) In an isocyanide, first an electrophile and then a nucleophile add at the carbon to form a species which usually undergoes further transformations.

