

Ordinary Thinking

Objective Questions

Carbohydrates

- The change in optical rotation, with time, of freshly prepared solution of sugar is known as [CPMT 1982, 85; BHU 1997]
 - Rotatory motion
 - Inversion
 - Specific rotation
 - Mutarotation
- Gun-cotton is
 - Nitrosucrose
 - Nitrocellulose
 - Nitroglucose
 - Nitropicrin
- Which of the following monosaccharide is a pentose [CPMT 1982, 87, 89, 93]
 - Galactose
 - Glucose
 - Fructose
 - Arabinose
- Amide group is present in
 - Lipids
 - Carbohydrates
 - Amino acids
 - Proteins
- Which of the following is a carbohydrate
 - Leucine
 - Albumin
 - Inulin
 - Maltase
- General formula for carbohydrates is
 - $C_nH_{2n}O_{2n+2}$
 - $C_x(H_2O)_{2x}$
 - $C_x(H_2O)_y$
 - None of these
- Benedict solution provides [CPMT 1983]
 - Ag^+
 - Li^+
 - Cu^{+2}
 - Ba^{+2}
- Glucose gives silver mirror with Tollen's reagent. It shows the presence of [MNR 1981; CPMT 1974, 81; MP PMT 1994]
 - An acidic group
 - An alcoholic group
 - A ketonic group
 - An aldehydic group
- A certain compound gives negative test with ninhydrin and positive test with Benedict's solution. The compound is [NCERT 1978; KCET 2000]
 - A protein
 - A monosaccharide
 - A lipid
 - An amino acid
- An organic compound answers Molisch's test as well as Benedict's test. But it does not answer Seliwanoff's test. Most probably, it is [KCET 2003]
 - Sucrose
 - Protein
 - Fructose
 - Maltose
- Glucose when heated with CH_3OH in presence of dry HCl gas gives α and β -methyl glucosides because it contains [CPMT 1982, 85]
 - An aldehyde group
 - A $-CH_2OH$ group
 - A ring structure
 - Five hydroxyl groups
- Which one is a disaccharide [CPMT 1981, 83]
 - Glucose
 - Fructose
 - Xylose
 - Sucrose
- Molecular formula $C_6H_{12}O_6$ is of
 - Glucose
 - Fructose
 - Both (a) and (b)
 - None of these
- Hydrolysis of sucrose is called [BHU 1979, 83; Pb. PMT 1999; Pb. CET 2000]
 - Esterification
 - Saponification
 - Inversion
 - Hydration
- In the 'glycolipids', the two sugars known to occur are glucose and
 - Fructose
 - Lactose
 - Galactose
 - Sucrose
- The 'epimerisation' involves
 - Change of configuration
 - Addition of one more 'C'
 - Substraction of a 'C'
 - Conversion of $-CHO$ to $-C=O$
- The compound which does not contain an asymmetric carbon atom is
 - Glycolaldehyde
 - Glyceraldehyde
 - Glucose
 - Galactose
- Which of the following sign indicate that the sugar is actually 'dextrorotatory'
 -
 - +
 - R -
 - All of these
- The standard compound for determination of configuration in the 'sugar chemistry' is
 - Glycolaldehyde
 - Glyceraldehyde
 - Glucose
 - Fructose
- Sugars are
 - Optically active polyhydroxy aldehydes
 - Optically active polyhydroxy ketones
 - Optically active polyhydroxy aldehydes or ketones
 - Polyhydroxy aldehydes or ketones which may or may not be optically active
- Molecular formula of pentahydroxy acid obtained when glucose is oxidised with Br_2 water is
 - $C_6H_{12}O_7$
 - $C_6H_{12}O_8$
 - $C_6H_{12}O_6$
 - $C_6H_{10}O_6$
- The 'phosphoglycerides' occur in
 - The brain and the spinal chord
 - Nails and hairs
 - Oils and fats
 - Waxes
- Sucrose is a [CPMT 1983]
 - Monosaccharide
 - Disaccharide



- (c) Trisaccharide (d) Polysaccharide
24. The commonest disaccharide has the molecular formula
[CPMT 1982; Manipal MEE 1995; MP PET 1999; AIIMS 1999]
(a) $C_{10}H_{18}O_9$ (b) $C_{10}H_{20}O_{10}$
(c) $C_{18}H_{22}O_{11}$ (d) $C_{12}H_{22}O_{11}$
25. On complete hydrolysis of starch, we finally get
[MNR 1982; DPMT 1979; CBSE PMT 1991; MP PMT 1987; MP PET 1993]
(a) Glucose (b) Fructose
(c) Glucose and fructose (d) Sucrose
26. Which is monosaccharide
(a) Glucose (b) Fructose
(c) Galactose (d) All of these
27. Which is polysaccharide
(a) Starch (b) Cellulose
(c) Glycogen (d) All of these
28. The calorific values of fats, carbohydrates and proteins vary in the order
(a) Fats > Carbohydrates > Proteins
(b) Fats > Proteins > Carbohydrates
(c) Carbohydrates > Proteins > Fats
(d) Proteins > Carbohydrates > Fats
29. Gun-cotton is obtained when conc. nitric acid reacts with
(a) Glycerine (b) Glycol
(c) Cellulose (d) Starch
30. A carbohydrate consists of [NCERT 1971]
(a) C and O (b) C, H and O
(c) C, H, N and O (d) C and H
31. Glucose forms many derivatives. The derivative which will help to prove the furanose structure is
[AIIMS 1980; DPMT 1985]
(a) Acetyl (b) Benzoyl
(c) Osazone (d) Isopropylidene
32. Glucose and fructose form [MP PMT 1986]
(a) Same osazone
(b) Same acid on oxidation
(c) Same alcohol when reduced
(d) Different osazone
33. On heating with conc. H_2SO_4 , sucrose gives [DPMT 1984]
(a) CO and CO_2 (b) CO and SO_2
(c) CO, CO_2 and SO_2 (d) None of these
34. The letter 'D' in carbohydrates represents
(a) Its direct synthesis (b) Its dextrorotation
(c) Its mutarotation (d) Its configuration
35. Starch can be used as an indicator for the detection of traces of [CPMT 1986]
(a) Glucose in aqueous solution
(b) Protein in blood
(c) Iodine in aqueous solution
(d) Urea in blood
36. It is best to carry out reactions with sugars in neutral or acid medium and not in alkaline medium. This is because in alkaline medium sugars undergo one of the following changes
(a) Racemisation (b) Decomposition
(c) Inversion (d) Rearrangement
37. Which one of the following compounds is found abundantly in nature [BHU 1983; Manipal MEE 1995; DCE 2002]
(a) Fructose (b) Starch
(c) Glucose (d) Cellulose
38. The substance that forms the plant cell walls is or Which carbohydrates is an essential constituents of plant cells
[KCET 1984; MP PET 1999; CPMT 2002]
(a) Cellulose (b) Sucrose
(c) Vitamins (d) Starch
39. Sugar can be tested in urine by
(a) Molisch test (b) Dunstan's test
(c) Benedict's test (d) Legal's test
40. When sucrose is heated with conc. HNO_3 the product is [CPMT 1979]
(a) Sucrose nitrate (b) Formic acid
(c) Oxalic acid (d) Citric acid
41. Amylopectin is [KCET 2005]
(a) Water soluble
(b) Water insoluble
(c) Forms colloidal solution with water
(d) Both (b) and (c)
42. Which of the following statements about ribose is incorrect [CPMT 1985]
(a) It is a polyhydroxy compound
(b) It is an aldehyde sugar
(c) It has six carbon atoms
(d) It exhibits optical activity
43. Maltose contains how many oxygen atoms
(a) 6 (b) 10
(c) 11 (d) 22
44. The correct name of 'sucrose' is
(a) α -D-glucopyranosyl- β -D-fructofuranoside
(b) β -D-glucopyranosyl- β -D-fructofuranoside
(c) α -D-glucopyranosyl- α -D-fructofuranoside
(d) β -D-glucopyranosyl- α -L-fructofuranoside
45. Sucrose is
(a) Laevorotatory (b) Dextrorotatory
(c) Racemic mixture (d) Optically inactive



1448 Biomolecules

46. The hydrolysis of sucrose produces a mixture which is
(a) Laevorotatory
(b) Dextrorotatory
(c) Equally both (+) and (-) rotatory
(d) Optically inactive
47. Sucrose is
(a) A reducing sugar
(b) Not a reducing sugar
(c) Partial reducing sugar
(d) Mixed sugar
48. Sucrose contains which of the following groups
(a) $-CHO$ (b) $>C=O$
(c) Both (a) and (b) (d) None of these
49. The fructose molecule in sucrose exists as
(a) Furanose (b) Pyranose
(c) Open chain (d) All
50. Which one of the following is laevorotatory [DPMT 1989]
(a) Glucose (b) Sucrose
(c) Fructose (d) None of these
51. Chemically 'digestion' is [NCERT 1978]
(a) Hydrolysis (b) Change in bacteria
(c) Hydrogenation (d) Dehydrogenation
52. Which one of the following is the reagent used to identify glucose [MP PMT 1993]
(a) Neutral ferric chloride
(b) Chloroform and alcoholic KOH
(c) Ammoniacal silver nitrate
(d) Sodium ethoxide
53. Sucrose on hydrolysis gives [MP PMT 1993; Bihar MEE 1997]
(a) Two molecules of glucose
(b) Two molecules of fructose
(c) One molecule each of glucose and fructose
(d) One molecule each of glucose and mannose
54. Which of the following is a disaccharide [CPMT 1990, 94]
(a) Lactose (b) Starch
(c) Cellulose (d) Glucose
55. Glucose cannot be classified as [CPMT 1989]
(a) A hexose (b) A carbohydrate
(c) An oligosaccharide (d) An aldose
56. The reagent which forms crystalline osazone derivative when reacted with glucose, is [CPMT 1990]
(a) Fehling solution (b) Phenylhydrazine
(c) Benedict solution (d) Hydroxylamine
57. An enzyme which brings about the conversion of starch into maltose is known as [BHU 1979]
(a) Maltase (b) Zymase
(c) Invertase (d) Diastase
58. Canesugar on hydrolysis gives [MADT Bihar 1984; NCERT 1977; AMU 1985]
(a) Glucose and maltose (b) Glucose and lactose
(c) Glucose and fructose (d) Only glucose
59. Glucose is a [CPMT 1984]
(a) Monosaccharide (b) Disaccharide
(c) Trisaccharide (d) Polysaccharide
60. Which carbohydrate is used in silvering of mirrors [BHU 1973; CPMT 1991]
(a) Sucrose (b) Starch
(c) Glucose (d) Fructose
61. A carbohydrate that cannot be hydrolysed to simpler forms is called
(a) Disaccharide (b) Monosaccharide
(c) Polysaccharide (d) Trisaccharide
62. If monosaccharide contains an aldehyde group, it is known as
(a) Epimer (b) Osone
(c) Osazone (d) Aldose
63. If a monosaccharide contains a ketogroup, it is known as
(a) Ketose (b) Osone
(c) Epimer (d) Osazone
64. The aqueous solution of a carbohydrate gives dark blue colour with iodine. It is
(a) Glucose (b) Fructose
(c) Sucrose (d) Starch
65. Which of the following carbohydrates is a disaccharide
(a) Glucose (b) Fructose
(c) Raffinose (d) Maltose
66. Optical activity is shown by
(a) Glucose (b) Fructose
(c) Sucrose (d) All of these
67. Which is a reducing sugar
(a) Glucose (b) Fructose
(c) Galactose (d) All of these
68. The ultimate product of oxidation of most of hydrogen and carbon in foodstuffs are [CPMT 1981]
(a) H_2O alone (b) CO_2 alone

- (c) H_2O and CO_2 (d) None of these
69. Osazone formation involves only 2 carbon atoms of glucose because of [MP PMT 1986]
 (a) Chelation (b) Oxidation
 (c) Reduction (d) Hydrolysis
70. Glucose will show mutarotation when solvent is [MP PMT 1986]
 (a) Acidic (b) Basic
 (c) Neutral (d) Amphoteric
71. Glucose contains [CPMT 1982]
 (a) One $-CHO$ group
 (b) Five $-OH$ groups
 (c) One primary alcoholic group
 (d) Four secondary alcoholic groups
 (e) All are correct
72. Carbohydrates are stored in human body as [MP PMT 1999; Kerala PMT 2004]
 (a) Glucose (b) Glycogen
 (c) Starch (d) Fructose
73. An example of a disaccharide made up of two units of the same monosaccharides is [KCET 1989; MP PET 1996; AFMC 2005]
 (a) Sucrose (b) Maltose
 (c) Lactose (d) None of these
74. The sugar present in fruits is [KCET 1984]
 (a) Fructose (b) Glucose
 (c) Sucrose (d) Galactose
75. Carbohydrates are [MADT Bihar 1983]
 (a) Hydrates of carbon
 (b) Polyhydroxy aldehydes or ketones
 (c) Polyhydroxy acid compounds
 (d) None of these
76. Glucose and fructose are [Bihar MADT 1982]
 (a) Isotopes (b) Isotones
 (c) Isomers (d) Homologues of each other
77. Hydrolytic conversion of sucrose into glucose and fructose is known as [BHU 1979, 97]
 (a) Induction (b) Saponification
 (c) Inversion (d) Esterification
78. Starch is a polymer of [DPMT 1982; CPMT 1975, 80; MP PMT 1994]
 (a) Glucose (b) Fructose
 (c) Both (a) and (b) (d) None of these
79. To become a carbohydrate a compound must contain at least [AFMC 1991]
 (a) 2 carbons (b) 3 carbons
 (c) 4 carbons (d) 6 carbons
80. Lactose on hydrolysis gives [KCET 1983]
 (a) Two glucose molecules
 (b) Two galactose molecules
 (c) A galactose molecule and a fructose molecule
 (d) A galactose molecule and a glucose molecule
81. An example of non-reducing sugar is [KCET 1988]
 (a) Cane sugar (b) Fructose
 (c) Lactose (d) Cellobiose
82. Cellulose is a polymer of [KCET 1984]
 (a) L-fructose (b) D-mannose
 (c) D-glucose (d) Amylose
83. The intermediate compound formed in the conversion of starch to glucose is [KCET 1984]
 (a) Lactose (b) Sucrose
 (c) Maltose (d) Fructose
84. Invertase brings about the conversion of [KCET 1986]
 (a) Starch to glucose
 (b) Sucrose to glucose and fructose
 (c) Maltose to glucose
 (d) Glucose to C_2H_5OH and CO_2
85. Which of the following pentoses will be optically active [MP PET 1994]
- | | | |
|----------|----------|----------|
| CHO | CHO | CHO |
| | | |
| $HCOH$ | $HCOH$ | $HCOH$ |
| | | |
| $HOCH$ | $HCOH$ | $HCOH$ |
| | | |
| $HCOH$ | $HOCH$ | $HCOH$ |
| | | |
| CH_2OH | CH_2OH | CH_2OH |
| I | II | III |
- (a) All (b) II and III
 (c) I (d) II
86. α -D-glucose and β -D-glucose differ from each other due to difference in one of the carbons with respect to its [CBSE PMT 1995; AFMC 1999]
 (a) Size of hemiacetal ring (b) Number of OH groups
 (c) Configuration (d) Conformation
87. Which carbohydrates has highest abundance in human blood [MP PET 1995]
 (a) d-fructose (b) d-glucose
 (c) Sucrose (d) Lactose
88. Formation of silver mirror by glucose shows that it is a/an

1450 Biomolecules

- (a) Oxidising agent (b) Acid [CPMT 1982, 87, 91; MP PET 2001]
 (c) Reducing agent (d) A salt of silver
89. Which of the following statements is right
 (a) Cellulose are linear polymers of β -glucose molecules with β -1,4-linkages
 (b) Starches are polymers of α -glucose molecules with β -1,4-linkages and some β -1,6-cross-linkages
 (c) Proteins are polyamides of β -amino acids
 (d) The structural information about their biosynthesis is contained in a class of compounds called nucleic acids, e.g. RNA and DNA
90. The number of atoms in the cyclic structure of D-fructose is [MP PMT 1997]
 (a) 5 (b) 6
 (c) 4 (d) 7
91. Which is used in motion picture films
 (a) Cellulose acetate (b) Glucose acetate
 (c) Starch acetate (d) Sucrose acetate
92. Glucose reacts with acetic anhydride to form [KCET 1996]
 (a) Mono-acetate (b) Tetra-acetate
 (c) Penta-acetate (d) Hexa-acetate
93. Which of the following does not show any reducing test of aldehyde [CPMT 1996; Orissa JEE 2004]
 (a) Sucrose (b) Fructose
 (c) Maltose (d) Lactose
94. When amylases catalyse the hydrolysis of starch, the final product obtained is chiefly [Pb. PMT 1998]
 (a) Cellobiose (b) Glucose
 (c) Maltose (d) Sucrose
95. Galactose is converted into glucose in [AFMC 1998]
 (a) Mouth (b) Stomach
 (c) Liver (d) Intestine
96. Which among the following is the simplest [CPMT 1999]
 (a) Glucose (b) Cellulose
 (c) Starch (d) None of these
97. Indigestible carbohydrate, which is also a constituent of our diet, is [Kerala (Med.) 1999]
 (a) Cellulose (b) Galactose
 (c) Maltose (d) Starch
98. Starch is converted into maltose by the [DPMT 1979; CPMT 1982; BHU 1999]
 (a) Maltase (b) Invertase
 (c) Zymase (d) Diastase
99. The disaccharide present in milk is
 (a) Maltose (b) Lactose
 (c) Sucrose (d) Cellobiose
100. Carbohydrates are used by body mainly [DCE 1999]
 (a) For obtaining vitamins
 (b) As source of energy
 (c) For all its developmental needs
 (d) For building muscles
101. In the viscose process the solvent for cellulose consists of [JIPMER 1999]
 (a) Ether and alcohol
 (b) Copper sulphate and ammonia
 (c) Sodium hydroxide and carbon disulphide
 (d) Acetic acid and acetic anhydride
102. Which of the following does not reduce Benedict's solution [KCET 2000]
 (a) Sucrose (b) Aldehyde
 (c) Glucose (d) Fructose
103. In polysaccharides the linkage connecting monosaccharide units is called
 (a) Glycoside linkage (b) Nucleoside linkage
 (c) Glycogen linkage (d) Peptide linkage
104. Blood sugar is the same as [DPMT 2000]
 (a) Glucose (b) Galactose
 (c) Glycogen (d) Fructose
105. Glucose has functional group [MH CET 2000]
 (a) Aldehydic
 (b) Aldehydic and alcoholic
 (c) Alcoholic
 (d) Ketonic and alcoholic
106. Which of the following is an aldohexose [KCET (Engg.) 2001]
 (a) Cellulose (b) Sucrose
 (c) Glucose (d) Raffinose
107. The calorific value is maximum in case of [Kerala (Med.) 2000]
 (a) Milk (b) Proteins
 (c) Minerals (d) Carbohydrates
108. An invert sugar is [AFMC 2000]
 (a) Isorotatory (b) Dextrorotatory
 (c) Laevorotatory (d) Optically inactive
109. The change in optical rotation with time of freshly prepared solutions of sugar is known as [JIPMER 2000]
 (a) Maturation (b) Rotatory motion
 (c) Inversion (d) Specific rotation

110. Yeast cell derive their energy from glucose by
[AIIMS 2001]
(a) Glycolysis formation (b) Respiration
(c) Formation (d) None of these
111. Which of the following is correct statement
[CBSE PMT 2001]
(a) Troleins are amino acid
(b) α -hydrogen is present in fructose
(c) Starch is polymer of α -glucose
(d) Amylose is compound of cellulose
112. Which of the following is a aldohexose [KCET 2001]
(a) Cellulose (b) Sucrose
(c) Galactose (d) Raffinose
113. The ultimate product of the hydrolysis of starch is
[DPMT 2001]
(a) Fructose (b) Glucose
(c) Sucrose (d) None of these
114. Raffinose is [Pb. PMT 2001]
(a) Trisaccharide (b) Monosaccharide
(c) Disaccharide (d) None of these
115. A sugar, that is not a disaccharide, among the following is
[KCET (Med./Engg.) 2002]
(a) Lactose (b) Galactose
(c) Sucrose (d) Maltose
116. To detect the reducing and non reducing sugars, which of the following test is used [MH CET 2002]
(a) Molisch test (b) Biuret test
(c) Fehling's test (d) Millions test
117. Which of the following is a disaccharide [MH CET 2002]
(a) Glucose (b) Ribulose
(c) Lactose (d) Arabinose
118. On heating glucose with Fehling's solution we get a precipitate whose colour is [CPMT 1979; CBSE PMT 1988; KCET 1992; DPMT 1983, 86; MP PMT 1996]
(a) Yellow (b) Red
(c) Black (d) White
119. Glycolysis is [CBSE PMT 2003]
(a) Conversion of glucose to haem
(b) Oxidation of glucose to glutamate
(c) Conversion of pyruvate to citrate
(d) Oxidation of glucose to pyruvate
120. Which of the following is an example of ketohexose
[Orissa JEE 2003]
(a) Mannose (b) Galactose
(c) Maltose (d) Fructose
121. The safest and the most common alternative of sugar is
[MP PMT 2003]
(a) Glucose (b) Aspartame
(c) Saccharin (d) Cyclodextrin
122. The specific rotation of equilibrium mixture of α -D-glucose and β -D-glucose, is
(a) $+19^\circ$ (b) $+112^\circ$
(c) $+52^\circ$ (d) $+100^\circ$
123. The charring of sugar, when treated with conc. H_2SO_4 , is due to [Pb. CET 2002]
(a) Oxidation (b) Reduction
(c) Dehydration (d) Hydrolysis
124. Which among the following is the simplest sugar
[Pb. CET 2002]
(a) Glucose (b) Cellulose
(c) Starch (d) Glycogen
125. Glucose and mannose are
(a) Epimers (b) Anomers
(c) Ketohexoses (d) Disaccharides
126. On hydrolysis, which produces only glucose [BVP 2004]
(a) Galactose (b) Maltose
(c) Sucrose (d) None
127. Pick out the one which does not belong to the family
[KCET 2004]
(a) Pepsin (b) Cellulose
(c) Ptyalin (d) Lipase
128. Which of the following is the sweetest sugar
[MP PMT 1997; CBSE PMT 1999; AIIMS 2000 Manipal MEE 1995; CPMT 1996; BHU 1997;]
(a) Glucose (b) Fructose
(c) Lactose (d) Sucrose
129. Oxidation of glucose is one of the most important reactions in a living cell. What is the number of ATP molecules generated in cells from one molecule of glucose
[CBSE PMT 1995]
(a) 38 (b) 12
(c) 18 (d) 28
130. Glucose has difference from fructose in that it [BHU 2005]
(a) Does not undergo hydrolysis
(b) Gives silver mirror with Tollen's reagent
(c) Monosaccharide
(d) None of these
131. In fructose, the possible optical isomers are
[Orissa JEE 2005]

1452 Biomolecules

- (a) 12 (b) 8
(c) 16 (d) 4
132. If an aqueous solution of glucose is allowed to freeze then crystal of which will be separated out first [DPMT 2005]
(a) Glucose (b) Water
(c) Both of these (d) None of these
133. Which is false [J & K 2005]
(a) Glucose is a disaccharide
(b) Starch is a polysaccharide
(c) Glucose and fructose are not anomers
(d) Invert sugar consists of glucose and fructose

Proteins, Amino Acids and Enzymes

1. Insulin is [CBSE PMT 1991]
(a) An amino acid (b) Protein
(c) A carbohydrate (d) A lipid
2. Peptides are
(a) Esters (b) Salts
(c) Amides (d) Ketones
3. The proteins which are insoluble in water are
(a) Fibrous proteins (b) Globular proteins
(c) Both (a) and (b) (d) None of these
4. Irreversible precipitation of proteins is called
(a) Denaturation (b) Hydrolysis
(c) Rearrangement (d) Electrophoresis
5. The proteins with a prosthetic group are called
(a) Pseudo proteins (b) Complex proteins
(c) Conjugated proteins (d) Polypeptides
6. The prosthetic group of haemoglobin is
(a) Porphin (b) Haem
(c) Globin (d) Globulin
7. When collagen is boiled with water, it forms
(a) Precipitate (b) Solution
(c) Gelatin (d) Complex collagen
8. Which of the following is not essential amino acid
(a) Valine (b) Lysine
(c) Histidine (d) Glycine
9. Amino acids are
(a) Liquids
(b) Volatile solids
(c) Non-volatile crystalline compounds
(d) Mixture of amines and acids
10. Isoelectric point is a
(a) Specific temperature
(b) Suitable concentration of amino acid
(c) Hydrogen ion concentration that does not allow migration of amino acid under electric field
(d) Melting point of an amino acid under the influence of electric field
11. Proteins are hydrolysed by enzymes into [CPMT 1981; BHU 1987; MP PMT 1994, 2002]
(a) Dicarboxylic acids (b) Hydroxy acids
(c) Amino acids (d) Aromatic acids
12. Proteins when heated with conc. HNO_3 give a yellow colour. This is [CPMT 1989]
(a) Oxidising test (b) Xanthoprotic test
(c) Hoppe's test (d) Acid-base test
13. Enzymes are [DPMT 1980; MP PMT 1993, 96]
(a) Proteins (b) Minerals
(c) Oils (d) Fatty acids
14. Proteins are built up of [CPMT 1981, 99; BHU 1987; CBSE PMT 2001; MP PMT 1987, 96; KCET 1984]
(a) Dicarboxylic acids (b) Amino acids
(c) Alcohols (d) Hydroxy acids
15. The main structural feature of proteins is [MNR 1987; MP PET 1993, 97, 2004]
(a) The ester linkage (b) The ether linkage
(c) The peptide linkage (d) All of these
16. Pepsin enzyme hydrolyses [NCERT 1984; MP PET 1999; MP PMT 2001]
(a) Proteins to amino acids
(b) Fats to fatty acids
(c) Glucose to ethyl alcohol
(d) Polysaccharides to monosaccharides
17. Which one of the following proteins transports oxygen in the blood stream
(a) Myoglobin (b) Insulin
(c) Albumin (d) Haemoglobin
18. Enzymes are [MP PET 1993]
(a) Living organisms
(b) Dead organisms
(c) Complex nitrogenous substances produced in living cells
(d) None of these
19. Which is an essential constituent of diet [AFMC 1980]
(a) Starch (b) Glucose
(c) Carbohydrate (d) Protein
20. Proteins can be used
(a) As food (b) In textile
(c) As enzyme (d) All of these

21. Which of the following foodstuffs contains nitrogen
[DPMT 1986; MH CET 2002]
(a) Carbohydrates (b) Fats
(c) Proteins (d) None of these
22. pH in stomach is approximately
(a) 7 (b) 2.0
(c) 6.5 (d) 10
23. The helical structure of proteins is established by
[CPMT 1988]
(a) Peptide bonds (b) Dipeptide bond
(c) Hydrogen bond (d) Vander Waal's forces
24. Natural silk is a
(a) Polyester (b) Polyamide
(c) Polyacid (d) Polysaccharide
25. Protein contains [CPMT 1975; MP PMT 2002]
(a) C, H, O and N (b) Only C and H
(c) Cl, H and O (d) All of these
26. The end product of protein digestion is
[CPMT 1981; KCET 1984]
(a) Amino acid (b) Glucose
(c) Glycerol (d) Oxalic acid
27. Protein can be most easily removed from [MNR 1988]
(a) Alkanes (b) Alkenes
(c) Alkynes (d) Benzene
28. Which of the following contains the highest percentage of protein [CPMT 1984]
(a) Groundnut (b) Cow's milk
(c) Egg (d) Wheat
29. The enzyme ptylin used for the digestion of food is present in [CPMT 1981; Pb. PMT 2004]
(a) Saliva (b) Blood
(c) Intestines (d) Adrenal glands
30. Which one of the following is an amino acid [KCET 1984]
(a) CH_3CONH_2 (b) $CH_3CONHCH_3$
(c) CH_3NHCHO (d) $NH_2CH_2.COOH$
31. Biuret test is used for the detection of [KCET 1993]
(a) Saturated oils (b) Sugars
(c) Proteins (d) Fats
32. Out of the following the best category of proteins is
[SCRA 1991]
(a) Polyamides (b) Polythioethers
(c) Glycerides (d) Polysaccharides
33. The molecular weight of protein is [KCET 1984]
(a) < 10000 (b) > 10000
(c) > 1000 (d) > 1000 and < 10000
34. Which of the following is not a classification of proteins
[KCET 1984]
(a) Enzymes (b) Antibodies
(c) Antigens (d) Hormones
35. The protein that is a structural material is [KCET 1984]
(a) Albumin (b) Oxytocin
(c) Haemoglobin (d) Keratin
36. For α -amino acids having the structure
$$R - \underset{\substack{| \\ NH_2}}{CH} - CO_2H$$

Which of the following statements are true
(A) Water solubility is maximum at a pH when concentrations of anions and cations are equal
(B) They give ninhydrin test
(C) On reacting with nitrous acid give off N_2
[MP PET 1994]
(a) All (b) B and C
(c) A and B (d) A
37. Which of the following reacts with haemoglobin in the blood to form carboxyhaemoglobin [Manipal MEE 1995]
(a) CO (b) CO_2
(c) HCOOH (d) H_2CO_3
38. Secondary structure of a protein refers to [CBSE PMT 1995]
(a) Mainly denatured proteins and structures of prosthetic groups
(b) Three dimensional structure, specially the bond between amino acid residues that are distant from each other in the polypeptide chain
(c) Linear sequence of amino acid residues in the polypeptide chain
(d) Regular folding patterns of continuous portions of the polypeptide chain
39. Of the following statements about enzymes which ones are true
(i) Enzymes lack in nucleophilic groups
(ii) Enzymes are highly specific both in binding chiral substrates and in catalyzing their reactions
(iii) Enzymes catalyse chemical reactions by lowering the activation energy
(iv) Pepsin is a proteolytic enzyme
(a) (i) and (iv) (b) (i) and (iii)
(c) (ii), (iii) and (iv) (d) (i)
40. Proteins are composed of [MP PMT 1995; J & K 2005]
(a) α -amino acids (b) Carbohydrates
(c) Vitamins (d) Mineral salts

1454 Biomolecules

41. Read the following statements carefully
 (A) Albumin is a simple protein
 (B) The amino acid alanine contains an acidic side chain
 (C) Insulin is a hormone
 (D) Muscles contain the protein keratin
 Point out the wrong statements in the above set of statements
 (a) A, B (b) C, D
 (c) A, C (d) B, D
42. Enzymes in the living systems [CPMT 1999; AIIMS 2000; CBSE PMT 1997; MP PET 1999;]
 (a) Provide energy
 (b) Provide immunity
 (c) Transport oxygen
 (d) Catalyse biological processes
43. Which of the following statements about proteins is not true [MP PET 2001]
 (a) Amino acid residues join together to make a protein molecule
 (b) Proteins are polymers with formula $(C_6H_{10}O_5)_n$
 (c) Eggs are rich in protein
 (d) Pulses are good source of proteins
44. Enzymes [AIIMS 1996]
 (a) Accelerate biochemical reactions
 (b) Have optimum activity at body temperature
 (c) Consist of amino acids
 (d) Have all these properties
45. The functional group which is found in amino acid is [AFMC 1998; AIEEE 2002]
 (a) $-COOH$ group (b) $-NH_2$ group
 (c) $-CH_3$ group (d) Both (a) and (b)
46. Amino acids are produced on hydrolysis of [AIIMS 1996]
 (a) Nucleic acid (b) Carbohydrates
 (c) Fats (d) Proteins
47. Enzymes belong to which class of compounds [KCET 1996]
 (a) Polysaccharides
 (b) Polypeptides
 (c) Polynitrogen heterocyclic compounds
 (d) Hydrocarbons
48. By the action of enzymes, the rate of biochemical reaction [CBSE PMT 1994]
 (a) Decreases (b) Increases
 (c) Does not change (d) Either (a) or (c)
49. Metal present in blood is [CPMT 1997]
 (a) Al (b) Mg
 (c) Cu (d) Fe
50. Which compound can exist in a dipolar (zwitter ion) state [Pb. PMT 1998]
 (a) $C_6H_5CH_2CH(N=CH_2)COOH$
 (b) $(CH_3)_2CH.CH(NH_2)COOH$
 (c) $C_6H_5CONHCH_2COOH$
 (d) $HOOC.CH_2CH_2COCOHOH$
51. What is the monomer of polypeptide [KCET 1998; JIPMER 1999; Pb. CET 2002]
 (a) Amino acid (b) Glucose
 (c) Nucleoside (d) Nucleotide
52. Which of the following enzymes is not useful in the digestion of proteins [KCET 1998]
 (a) Chymotrypsin (b) Pepsin
 (c) Trypsin (d) Lipase
53. Haemoglobin is [CBSE PMT 1997; BHU 2004]
 (a) An enzyme (b) A globular protein
 (c) A vitamin (d) A carbohydrate
54. Albumin proteins are most abundant in [BHU 1998]
 (a) Meat (b) Milk
 (c) Egg (d) Soyabean
55. Dialysis can separate [BHU 1998]
 (a) Glucose and fructose
 (b) Glucose and sucrose
 (c) Glucose and NaCl
 (d) Glucose and proteins
56. Which one of the following is an example of a globular protein [Orissa JEE 1997]
 (a) Keratin (b) Insulin
 (c) Collagen (d) Myoglobin
57. Leucine amino acids is the [Pb. PMT 1999]
 (a) Essential (b) Non-essential
 (c) Aromatic (d) Basic
58. Which of the following tests is not used for testing proteins [Kerala PMT 1999; KCET 1999]
 (a) Millon's test (b) Molisch's test
 (c) Biuret test (d) Ninhydrin test
59. Amino acids usually exist in the form of Zwitter ions. This means that it consists of [KCET 2000]
 (a) The basic group $-NH_2$ and the acidic group $-COOH$



- (b) The basic group $-NH_3^+$ and the acidic group $-CO_2^-$
- (c) The basic group $-CO_2^-$ and the acidic group NH_3^+
- (d) No acidic or basic group
60. The most important energy carrier in all the living cells is
[MP PET 2000; KCET 2000]
- (a) AMP (b) ATP
(c) ADP (d) UDP
61. The 10% energy transfer law of food chain was given by
[BHU 2000]
- (a) Stanley (b) Weismann
(c) Lindemann (d) Tansley
62. Which of the following is a conjugated protein[BHU 2000]
- (a) Glycoprotein (b) Phosphoprotein
(c) Chromoprotein (d) All of these
63. The number of essential amino acids in man is
[CBSE PMT 2000]
- (a) 8 (b) 10
(c) 18 (d) 20
64. Pick out wrong combination [DCE 2000]
- (a) $Fe^{+2} \rightarrow$ Haemoglobin
(b) $Mg^{2+} \rightarrow$ Photosynthesis
(c) $Se^{2+} \rightarrow$ Krebs Cycle
(d) $CO^{+2} \rightarrow$ Vitamin B-12
65. The decomposition of complex organic compounds into simpler compound with the help of enzyme is known as
[Pb. PMT 2000]
- (a) Catabolism (b) Anabolism
(c) Fermentation (d) Metabolism
66. A biological catalyst is essentially
[Pb. PMT 2000; BHU 2004]
- (a) A carbohydrates (b) An amino acids
(c) A nitrogen molecule (d) Fats
67. The test used for identifying peptide linkage in proteins is
[KCET (Engg.) 2001]
- (a) Borsche's test (b) Molisch's test
(c) Ninhydrin test (d) Biuret test
68. Which of the following is not a function of proteins
[MP PMT 2001]
- (a) Nails formation (b) Skin formation
(c) Muscle formation (d) Providing energy for metabolism
69. The helical structure of proteins is stabilized by
[MP PMT 2001]
- (a) Peptide bonds (b) Dipeptide bond
(c) Hydrogen bond (d) Vander Waal's forces
70. The optically inactive amino acid is
[MP PMT 2001; BHU 2005]
- (a) Lysine (b) Glycine
(c) Arginine (d) Alanine
71. Which α amino acid can cross link peptide chains
[AIIMS 2001]
- (a) Serine (b) Cysteine
(c) Glutamine (d) Tyrosine
72. Amino acids are the building blocks of [MH CET 2001]
- (a) Fat (b) Vitamin
(c) Protein (d) Carbohydrate
73. Which of the following protein destroys the antigen when it enters in body cell[AIIMS 2001; Pb. PMT 2000]
- (a) Antibodies (b) Insulin
(c) Chromoprotein (d) Phosphoprotein
74. An antibiotic with a broad spectrum [AFMC 2001]
- (a) Kills the antibodies
(b) Acts on a specific antigen
(c) Acts on different antigens
(d) Acts on both the antigens and antibodies
75. Antibodies are [CBSE PMT 2001]
- (a) Carbohydrate (b) Globular protein
(c) Immunoglobulins (d) Cellulose compounds
76. Excess of Na^+ ions in our system causes[BHU 2001]
- (a) High B.P. (b) Low B.P.
(c) Diabetes (d) Anaemia
77. The example of a protein is [MP PET 2003]
- (a) Narvone (b) Lacithin
(c) Cellulose (d) Insulin
78. Enzymes are made up of [CBSE PMT 2002]
- (a) Carbohydrates
(b) Edible proteins
(c) Nitrogen containing carbohydrates
(d) Proteins with specific structure
79. Chlorophyll contains [RPMT 2002]
- (a) Fe (b) Na
(c) Mg (d) Zn
80. Which one of the following biomolecules is insoluble in water [AIIMS 2005]
- (a) α - Keratin (b) Haemoglobin



1456 Biomolecules

- (c) Ribonuclease (d) Adenine
81. A nanopeptide contains peptide linkages [KCET 2005]
 (a) 10 (b) 8
 (c) 9 (d) 18
82. Identify the incorrect statement [Kerala (Med.) 2003]
 (a) An octa deca peptide contains 18 amino acid residues and 17 peptide bonds
 (b) Addition of an inert gas into a system in thermodynamic equilibrium for the dissociation of PCl_5 shifts the equilibrium to the left.
 (c) When gold dissolves in aquaregia the complex formed is chloroauric acid
 (d) In the extraction of aluminium purified bauxite is dissolved in molten fluorospar
 (e) If the pH value of a solution is to be decreased from 5 to 3, the hydrogen ion concentration must be increased twice the initial value.
83. α -helix is found in [Kerala (Engg.) 2002]
 (a) DNA (b) RNA
 (c) Lipid (d) Protein
84. The main structural of protein is [UPSEAT 2000, 02]
 (a) The ester linkage (b) The ether linkage
 (c) The peptide linkage (d) All of these
85. Among the following, the achiral amino acid is [AIIMS 2003]
 (a) 2-Ethylalanine
 (b) 2-Methylglycine
 (c) 2-Hydroxymethyl serine
 (d) Tryptophan
86. Which of the following could act as a propellant or rockets [CBSE PMT 2003]
 (a) Liquid hydrogen + liquid nitrogen
 (b) Liquid oxygen + liquid argon
 (c) Liquid hydrogen + liquid oxygen
 (d) Liquid nitrogen + liquid oxygen
87. Which amino acid has aromatic ring [CPMT 2003]
 (a) Alanine (b) Glycine
 (c) Tyrosine (d) Lysine
88. The pH value of the solution in which a particular amino acid does not migrate under the influence of an electric field is called the [Kerala (Med.) 2003]
 (a) Eutectic point (b) Yielding point
 (c) Neutralisation point (d) Effusion
 (e) Isoelectric point
89. Which part of the protein molecule is responsible for function and activity of the proteins [AMU 2002]
 (a) Secondary structure (b) Peptide bond
 (c) Primary structure (d) Binding sites
90. The Structural formula of an amino acid, isoleucine is [MP PMT 2003]
- (a) $CH_3 - \overset{NH_2}{\underset{|}{CH}} - COOH$
- (b) $\begin{array}{c} CH_3 \\ \diagdown \\ CH \\ \diagup \\ CH_3 \end{array} - \overset{NH_2}{\underset{|}{CH}} - COOH$
- (c) $\begin{array}{c} CH_3 \\ \diagdown \\ CH \\ \diagup \\ C_2H_5 \end{array} - \overset{NH_2}{\underset{|}{CH}} - COOH$
- (d) $\begin{array}{c} C_2H_5 \\ \diagdown \\ CH \\ \diagup \\ C_2H_5 \end{array} - \overset{NH_2}{\underset{|}{CH}} - COOH$
91. The process by which synthesis of protein takes place based on the genetic information present in m-RNA is called [KCET 2003; Kerala CET 2005]
 (a) Translation (b) Transcription
 (c) Replication (d) Messenger hypothesis
92. Which of the following is used in our body as a fuel for muscles and nerves and to build and repair body tissues? [DCE 2003]
 (a) Cane sugar (b) Fructose
 (c) Proteins (d) Glucose
93. Which enzyme convert glucose into alcohol [Pb. CET 2003]
 (a) Invertase (b) Zymase
 (c) Maltase (d) Diastase
94. Which one of the following structures represents the peptide chain [CBSE PMT 2004; CPMT 2003; DCE 2002; MP PET 1994; Bihar MEE 1997; Orissa JEE 1997]
- (a) $\begin{array}{ccccccc} H & & H & & H & & O \\ | & & | & & | & & || \\ -N & -C & -C & -N & -C & -C & -N & -C & -C- \\ & & || & & || & & & & \\ & & O & & O & & & & \end{array}$
- (b) $\begin{array}{ccccccc} H & & O & & H & & \\ | & & || & & | & & \\ -N & -C & -C & -C & -N & -C & -C & -N & -C & -C & -C- \\ & & & & | & & | & & || & & \\ & & & & H & & O & & & & \end{array}$





95. The correct statement in respect of protein haemoglobin is that it [CBSE PMT 2004]
- Acts as an oxygen carrier in the blood
 - Forms antibodies and offers resistance to diseases
 - Functions as a catalyst for biological reactions
 - Maintains blood sugar level
96. Identify the correct statement regarding enzymes [AIIEEE 2004]
- Enzymes are specific biological catalysts that cannot be poisoned
 - Enzymes are normally heterogeneous catalysts that are very specific in their action
 - Enzymes are specific biological catalysts that can normally function at very high temperature ($T \sim 1000K$)
 - Enzymes are specific biological catalysts that possess well-defined active sites
97. A biological catalyst is essentially
- An enzyme
 - A carbohydrate
 - An amino acid
 - A nitrogen compound
98. Which synthesis was done by Stainley Millar [CPMT 1979]
- Amino acid
 - Protein
 - Virus
 - Vitamin
99. The bond that determines the secondary structure of proteins is or secondary structure of protein is due to [NCERT 1984; MP PET 1996; MP PMT 1997]
- Coordinate bond
 - Covalent bond
 - Hydrogen bond
 - Peptide bond
- (a) Monocarboxylic acids (b) Monohydric alcohols
(c) Monohaloalkanes (d) Enzymes
3. Which of the following is not a lipid
- Oils
 - Fats
 - Waxes
 - Proteins
4. The 'acid value' of an oil or fat is measured in terms of weight of
- NH_4OH
 - $NaOH$
 - KOH
 - CH_3COOH
5. The 'saponification value' of an oil or fat is measured in terms of
- NH_4OH
 - $NaOH$
 - KOH
 - C_6H_5OH
6. The 'iodine value' of an oil indicates
- Its boiling point
 - Inflammability
 - Unsaturation present in acid contents
 - Solubility of salt in oils
7. Hardening of oils is caused by
- H_2
 - N_2
 - O_2
 - CO_2
8. Which of the following is obtained when an oil is hydrolysed with alkali [BHU 2004]
- Fat
 - Wax
 - Soap
 - Vitamin
9. Which of the following indicates the number of free $-OH$ groups in an oil or fat
- Iodine value
 - Acid value
 - Acetyl value
 - Saponification value
10. Which of the following is not glyceride
- Lipids (simple)
 - Phospholipids
 - Sphingolipids
 - All
11. The most important food reserves of animals and plants are [MP PET 1993]
- Carbohydrates
 - Proteins
 - Vitamins
 - Fats
12. Which of the following gives maximum energy in metabolic processes [CPMT 1991; MP PET 1999]
- Proteins
 - Carbohydrates
 - Lipids
 - Vitamins
13. The energy change produced by the combustion of food is called the 'calorific value'. The highest calorific value is given by [NCERT 1984; AFMC 1988]

Fats and Lipids

- Tripalmitin is
 - A protein
 - An enzyme
 - A lipid
 - A carbohydrate
- On hydrolysis, all lipids yield



1458 Biomolecules

14. Cell membrane contains
 (a) Proteins (b) Fats
 (c) Carbohydrates (d) Vitamins
15. Which of the following compounds do not belong to lipids
 (a) Alternate layers of phospholipid and coline
 (b) Double layers of phospholipid
 (c) Double layers of phospholipid with polar ends projected outside
 (d) Double layers of phospholipid with polar ends projected inside
16. Which is not a macromolecule
 (a) DNA (b) Starch
 (c) Palmitate (d) Insulin
17. A distinctive and characteristic functional group of fats is
 (a) An ester group
 (b) A peptide group
 (c) A ketonic group
 (d) An alcoholic group
18. The waxes are long chain compounds of fatty acids, which belong to the class of
 (a) Esters (b) Ethers
 (c) Alcohols (d) Acetic acid
19. Hydrolytic reaction of fats, with caustic soda, is known as
 (a) Acetylation (b) Carboxylation
 (c) Saponification (d) Esterification
20. Fat consists of
 (a) Monohydroxy carboxylic acid
 (b) Monohydroxy aliphatic carboxylic acid
 (c) Monohydroxy aliphatic, saturated carboxylic acid
 (d) Dihydroxy aliphatic carboxylic acid
21. The alcohol obtained by the hydrolysis of oils and fats is
 (a) Glycol (b) Glycerol
 (c) Propanol (d) Pentanol
22. Iodine value is related to
 (a) Fats and oils (b) Alcohols
 (c) Esters (d) Hydrocarbons
23. Phospholipids are esters of glycerol with
 (a) Three phosphate groups
 (b) Three carboxylic acid residues
 (c) Two carboxylic acid residues and one phosphate group
 (d) One carboxylic acid residue and two phosphate groups
24. Oils and fats are jointly called
 (a) Lipids (b) Soaps
 (c) Proteins (d) Polymer
25.
$$\begin{array}{c} \text{CH}_2\text{OOCR}' \\ | \\ \text{CHOOCR}'' \\ | \\ \text{CH}_2\text{OOCR}''' \end{array} \xrightarrow[\text{Hydrolysis}]{\text{Enzyme}} \begin{array}{c} \text{CH}_2\text{OH} \quad \text{R}'\text{COOH} \\ | \quad \quad \quad | \\ \text{CHOH} + \text{R}''\text{COOH} \\ | \quad \quad \quad | \\ \text{CH}_2\text{OH} \quad \text{R}'''\text{COOH} \end{array}$$
- The enzyme used in the above reaction is
 (a) Amylase (b) Lactase
 (c) Lipase (d) Invertase
26. Oleic, stearic and palmitic acids are
 (a) Fatty acid (b) Amino acid
 (c) Nucleic acid (d) Essential acid
27. An example for a saturated fatty acid, present in nature is
 (a) Oleic acid (b) linoleic acid
 (c) Linolenic acid (d) Palmitic acid

Vitamin, Hormone and Nucleic acid

- A nucleotide consists of
 (a) Base and sugar (b) Base and phosphate
 (c) Sugar and phosphate (d) Base, sugar and phosphate
2. Which of the following is responsible for heredity character
 (a) DNA (b) RNA
 (c) Proteins (d) Hormones
3. The base adenine occurs in
 (a) DNA only (b) RNA only
 (c) DNA and RNA both (d) Protein
4. The protein which maintains blood sugar level in the human body
 (a) Haemoglobin (b) Oxytocin
 (c) Insulin (d) Ptyalin
5. Which of the following statements about the assembly of nucleotides in a molecule of deoxyribose nucleic acid (DNA) is correct
 (a) A pentose of one unit connects to a pentose of another
 (b) A pentose of one unit connects to the base of another
 (c) A phosphate of one unit connects to a pentose of another

- (d) A phosphate of one unit connects to the base of another
6. Vitamin A is present in [MP PET 1995, 2000]
 (a) Cod liver oil (b) Carrot
 (c) Milk (d) In all of these
7. Ascorbic acid is a [Bihar CEE 1995; MP PET 1995]
 (a) Vitamin (b) Enzyme
 (c) Protein (d) Carbohydrate
8. The chemical name of vitamin C is [J & K 2005]
 (a) Ascorbic acid (b) Folic acid
 (c) Nicotinic acid (d) Tartaric acid
9. Which of the following is not a constituent of RNA [MP PET 1996]
 (a) Ribose (b) Phosphate
 (c) Adenine (d) Pyridine
10. Which one is found in ATP ribonucleotide
 (a) Guanine (b) Uracil
 (c) Adenine (d) None of these
11. Which of the following proteins acts as a messenger in living system
 (a) Hormone (b) Enzyme
 (c) Protective protein (d) Transport protein
12. Which substance is not present in nucleic acid [MP PET/PMT 1998]
 (a) Cytosine (b) Adenine
 (c) Thymine (d) Guanidine
13. The deficiency of vitamin B_1 causes [CPMT 1994; MP PMT 1999; BHU 2000]
 (a) Beri-beri (b) Scurvy
 (c) Rickets (d) Anaemia
14. Which of the following is not present in nucleic acids [MP PMT 1999]
 (a) Uracil (b) 2-aminopyridine
 (c) Thymine (d) Adenine
15. In nucleic acids, the sequence is [AIIMS 1996]
 (a) Base-phosphate-sugar (b) Phosphate-base-sugar
 (c) Sugar-base-phosphate (d) Base-sugar-phosphate
16. The segment of DNA which acts as the instructional manual for the synthesis of the protein is [Pb. PMT 1998]
 (a) Nucleoside (b) Nucleotide
 (c) Ribose (d) Gene
17. The double helical structure of DNA was proposed by [KCET 1998]
 (a) Watson and Crick (b) Meicher
 (c) Emil Fischer (d) Khorana
18. A segment of DNA molecule which codes or specifies for one polypeptide chain is called [KCET 1998]
 (a) Phosphate group (b) Adenine
 (c) Gene (d) Amino acid
19. In DNA, the complementary bases are [CBSE PMT 1998]
 (a) Uracil and adenine; cytosine and guanine
 (b) Adenine and thymine; guanine and cytosine
 (c) Adenine and thymine; guanine and uracil
 (d) Adenine and guanine; thymine and cytosine
20. The structure of DNA is [AFMC 1999]
 (a) Linear (b) Single helix
 (c) Double helix (d) Triple helix
21. Vitamin B_1 is [MP PMT 2000]
 (a) Riboflavin (b) Cobalamin
 (c) Thiamine (d) Pyridoxine
22. A gene is a segment of a molecule of [AIIMS 1999]
 (a) DNA (b) *m*-RNA
 (c) *t*-RNA (d) Protein
23. The deficiency of vitamin-C causes [MP PMT 2000; CPMT 2000]
 (a) Scurvy (b) Rickets
 (c) Pyrohea (d) Pernicious Anaemia
24. DNA contains the sugar [MP PMT 2000]
 (a) Deoxyribose (b) Ribose
 (c) *D*-Fructose (d) *D*-glucose
25. Which of the following is not a sex hormone [MP PMT 2000]
 (a) Testosterone (b) Estrone
 (c) Estradiol (d) Cortisone
26. Acquired immune deficiency syndroms (AIDS) is characterised [AIIMS 2000]
 (a) Killer T-cells
 (b) Reduction in number of helper T-cells
 (c) An autoimmune disease
 (d) Inability of body to produce interferons
27. The base present in DNA, but not in RNA is [KCET (Engg.) 2001; NCERT 1978; Manipal MEE 1985; MP PMT 1994, MP PET 1995; DCE 2004]
 (a) Guanine (b) Adenine
 (c) Uracil (d) Thymine
28. Mutation of DNA occurs due to changes in the sequence of one of the following [MP PMT 2001]
 (a) Bases (b) Ribose units
 (c) Phosphate units (d) Sugar units
29. Which of the following is not true about vitamins [AFMC 2001]
 (a) They are vital for life
 (b) They help in digestion
 (c) They were named by "Funic"
 (d) Their deficiency causes diseases
30. Blood calcium level can be increased by the administration of [AFMC 2001]
 (a) Glucagon (b) Calcitonin
 (c) Thyroxine (d) Paratharmone



1460 Biomolecules

31. The first hormone chemically synthesised in the laboratory is [BHU 2002]
(a) Cortisone (b) Insuline
(c) Adrenaline (d) Estrone
32. Purine derivative among the following bases is [KCET (Med./Engg.) 2002; MPPET 2004]
(a) Guanine (b) Cytosine
(c) Thymine (d) Uracil
33. RNA is different from DNA because RNA contains [AIEEE 2002, 04]
(a) Ribose sugar and thymine
(b) Ribose sugar and uracil
(c) Deoxyribose sugar and thymine
(d) Deoxyribose sugar and uracil
34. Deficiency of which vitamin causes rickets [MP PET 2002]
(a) Vitamin-D (b) Vitamin-B
(c) Vitamin-A (d) Vitamin-K
35. Which do the following vitamins has isoprene units in its structure [JIPMER 2002]
(a) Vitamin A (b) Vitamin C
(c) Vitamin B₂ (d) Vitamin D
36. The reason for double helical structure of DNA is operation of [CBSE PMT 2003; DPMT 2004]
(a) Vander Waal's forces
(b) Dipole-dipole interaction
(c) Hydrogen bonding
(d) Electrostatic attractions
37. The tripeptide hormone present in most living cells is... [KCET 2003]
(a) Glutathione (b) Glutamine
(c) Oxytocin (d) Ptyalin
38. The function of DNA in an organism is [DCE 2003]
(a) To assist in the synthesis of RNA molecule
(b) To store information of heredity characteristics
(c) To assist in the synthesis of proteins and polypeptides
(d) All of these
39. The hormone that helps in the conversion of glucose to glycogen in [CBSE PMT 2004]
(a) Adrenaline (b) Insulin
(c) Cortisone (d) Bile acids
40. Insulin production and its action in human body are responsible for the level of diabetes. This compound belongs to which of the following categories [AIEEE 2004]
(a) An enzyme (b) A hormone
(c) A co-enzyme (d) An antibiotic
41. Codon is present in [Pb. PMT 2004]
(a) *t*-RNA (b) *m*-RNA
(c) *r*-RNA (c) All of these
42. Energy is stored in our body in the form of [CBSE PMT 2001; KCET 2003]
(a) ATP (b) ADP
(c) Fats (d) Carbohydrates
43. Nucleic acid is a polymer of
(a) Nucleosides (b) α -amino acids
(c) Nucleotides (d) Glucose
44. A nucleoside on hydrolysis gives
(a) A heterocyclic base and orthophosphoric acid
(b) An aldopentose, a heterocyclic base and orthophosphoric acid
(c) An aldopentose and a heterocyclic base
(d) An aldopentose and orthophosphoric acid
45. An alternation in the base sequence of nucleic acid molecule is called [Kerala PMT 2004]
(a) Replication (b) Mutation
(c) Duplication (d) Dislocation
(e) Flocculation
46. Vitamin B₆ is known as [DCE 2004]
(a) Pyridoxin (b) Thiamine
(c) Tocopherol (d) Riboflavin

Critical Thinking

Objective Questions

1. Number of chiral carbons in β -D-(+)-glucose is [CBSE PMT 2004; MHCET 2004]
(a) Three (b) Four
(c) Five (d) Six
2. The nucleic acid base having two possible binding sites is [AIIMS 2004]
(a) Thymine (b) Cytosine
(c) Guanine (d) Adenine
3. Subunits present in haemoglobin are [AIIMS 2003]
(a) 2 (b) 3
(c) 4 (d) 5
4. A sequence of how many nucleotides in messenger RNA makes a codon for an amino acid
(a) One (b) Two
(c) Three (d) Four
5. Chargaff's rule states that in an organism [CBSE PMT 2003]
(a) Amounts of all bases are equal

- (b) Amount of adenine (*A*) is equal to that of thymine (*T*) and the amount of guanine (*G*) is equal to that of cytosine (*C*)
- (c) Amount of adenine (*A*) is equal to that of guanine (*G*) and the amount of thymine (*T*) is equal to that of cytosine (*C*)
- (d) Amount of adenine (*A*) is equal to that of cytosine (*C*) and the amount of thymine (*T*) is equal to guanine (*G*)
6. DNA multiplication is called [Kerala (Med.) 2000]
 (a) Translation (b) Transduction
 (c) Transcription (d) Replication
7. Insulin is a protein which plays the role of [KCET 1986]
 (a) An antibody (b) A hormone
 (c) An enzyme (d) A transport agent
8. Proteins fulfil several functions in living systems. An example of a protein which acts as a hormone is [KCET 1985]
 (a) Casein (b) Oxytocin
 (c) Trypsin (d) Keratin
9. Pick out the unsaturated fatty acid from the following [KCET 2004; MHCET 2002]
 (a) Stearic acid (b) Lauric acid
 (c) Oleic acid (d) Palmitic acid
10. Vitamin B_{12} contains metal [Bihar MEE 1997; RPET 1999; Pb. PMT 1999; AFMC 2002; CBSE PMT 2003; CPMT 2003; MP PMT 2003]
 (a) Ca (II) (b) Zn (II)
 (c) Fe (II) (d) Co (III)
11. The number of molecules of ATP produced in the lipid metabolism of a molecule of palmitic acid is [CBSE PMT 1998]
 (a) 130 (b) 36
 (c) 56 (d) 86
12. Protein can be most easily removed from [UPSEAT 2000, 02]
 (a) Alkanes (b) Alkenes
 (c) Alkynes (d) Benzene
13. The enzyme which hydrolyses triglycerides to fatty acids and glycerol is called [UPSEAT 2000, 02]
 (a) Zymase (b) Pepsin
 (c) Maltase (d) Lipase
14. The helical structure of protein is stabilized by [CBSE PMT 2004]
 (a) Ether bonds (b) Peptide bonds
 (c) Dipeptide bonds (d) Hydrogen bonds
15. The cell membranes are mainly composed of [CBSE PMT 2005]
 (a) Carbohydrates (b) Proteins
 (c) Phospholipids (d) Fats
16. A compound of mol. wt. 180 is acetylated to give a compound of mol. wt. 390. The number of amino groups in the initial compound is [Kerala PMT 1999; KCET 1999]
 (a) 2 (b) 4
 (c) 5 (d) 6
17. Starting with three different amino acid molecules, how many different tripeptide molecules are formed [Kerala PMT 1999; KCET 1999]
 (a) 12 (b) 9
 (c) 8 (d) 6
18. Which one of the following is a polysaccharide [NDA 1999]
 (a) Nylon (b) Amylose
 (c) Ribose (d) Polyethylene
19. Which of the following is not an acidic amino acid
 (a) Lysine (b) Arginine
 (c) Aspartic acid (d) Histidine
20. Proteins do not respond to
 (a) Biuret test (b) Heller's ring test
 (c) Ninhydrin test (d) Lucas test
21. Alkyl benzene sulphonates can be conductivity used as detergents in hard water, unlike soaps, as [AMU 2002]
 (a) They are highly soluble in water
 (b) Their Ca^{++}/Mg^{++} salts are water soluble
 (c) They are non-ionic
 (d) Their Ca^{++}/Mg^{++} salts are insoluble in water
22. When glucose reacts with bromine water, the main product is [Pb. CET 2003; BHU 2004]
 (a) Acetic acid (b) Saccharic acid
 (c) Glyceraldehyde (d) Gluconic acid
23. A zwitter ion is [KCET 1989]
 (a) A positively charged ion without a metal atom in it
 (b) A negatively charged ion without metal atom in it
 (c) An ion with positive and negative charges at different points on it
 (d) A positive ion with a small charge on it
24. Ribose is an example of [KCET 1998]
 (a) Ketohexose (b) Aldopentose
 (c) Disaccharide (d) Aldohexose
25. The two forms of *D*-glucopyranose obtained from the solution of *D*-glucose are called
 (a) Isomer (b) Anomer
 (c) Epimer (d) Enantiomer
26. Sucrose molecule is made up of [KCET 2005]

- (a) A gluco pyranose and a fructo pyranose
 (b) A gluco pyranose and a fructo furanose
 (c) A gluco furanose and a fructo pyranose
 (d) A gluco furanose and a fructo furanose

Assertion & Reason

For AIIMS Aspirants

Read the assertion and reason carefully to mark the correct option out of the options given below :

- (a) If both assertion and reason are true and the reason is the correct explanation of the assertion.
 (b) If both assertion and reason are true but reason is not the correct explanation of the assertion.
 (c) If assertion is true but reason is false.
 (d) If the assertion and reason both are false.
 (e) If assertion is false but reason is true.

- Assertion : Glycine is amphoteric in nature.
Reason : Glycine contains both acid and basic groups. [AIIMS 1996]
- Assertion : Hydrolysis of sucrose is known as inversion of cane sugar.
Reason : Sucrose is a disaccharide. [AIIMS 1997]
- Assertion : Proteins on hydrolysis produce amino acids.
Reason : Amino acids contain $-NH_2$ and $-COOH$ groups. [AIIMS 1998]
- Assertion : Sucrose undergo mutarotation.
Reason : Sucrose is a disaccharide. [AIIMS 2000]
- Assertion : DNA molecules and RNA molecules are found in the nucleus of a cell.
Reason : On heating the enzyme do not lose their specific activity. [AIIMS 2002]
- Assertion : All Amino acids exist as Zwitter ions.
Reason : Amino acids have both $-NH_2$ and $-COOH$ group. [AIIMS 2002]
- Assertion : Activity of an enzyme is pH -dependent.
Reason : Change in pH affects the solubility of the enzyme in water. [AIIMS 2003]
- Assertion : Glycosides are hydrolyzed in acidic conditions.
Reason : Glycosides are acetals. [AIIMS 2003]
- Assertion : Haemoglobin is an oxygen carrier.
Reason : Oxygen binds as O_2^- to Fe of haemoglobin. [AIIMS 2003]
- Assertion : Carboxypeptidase is an exopeptidase.
Reason : It cleaves the N -terminal bond.
- Assertion : Sucrose is a non-reducing sugar.
Reason : It has glycosidic linkage. [AIIMS 2004]
- Assertion : Sucrose is a disaccharide.
Reason : Sucrose is dextro rotatory.
- Assertion : Fructose reduces Fehling's solution and Tollen's reagent.
Reason : Fructose does not contain any aldehyde group.
- Assertion : The specific rotation of a freshly prepared solution of α -glucose decreases from $+112^\circ$ to 52.7° while that of β glucose increase from $+19^\circ$ to 52.7° .
Reason : The change in specific rotation of an optically active compound with time to an equilibrium value is called mutarotation.
- Assertion : α -amino acids exist as dipolar ions or zwitter ions.
Reason : α -amino acids are the building blocks of proteins.
- Assertion : Valine is an essential amino acid.
Reason : The lack of essential amino acids in the diet causes Kwashiorkor.
- Assertion : Sequence of bases in DNA is TGAACCTT and sequence of bases in m -RNA is CATTAAACC.
Reason : In DNA nitrogenous bases have hydrogen bonds.
- Assertion : Millon's test is a test to identify carbohydrates.
Reason : Millon's reagent is solution of mercurous nitrate and mercuric nitrate in nitric acid containing little nitrous acid.
- Assertion : ATP molecules are energy rich molecules.
Reason : ATP consists of a purine base adenine, pentose sugar ribose and a string of three phosphate groups.
- Assertion : Solubilities of protein is minimum at the isoelectric point.
Reason : At isoelectric point, protein molecule behaves as a zwitter ion.
- Assertion : Amino acids are soluble in benzene and ether.
Reason : Amino acids exist as zwitter ions.
- Assertion : A solution of sucrose in water is dextrorotatory but on hydrolysis in presence of little hydrochloric acid, It becomes laevorotatory.
Reason : Sucrose on hydrolysis gives unequal amounts of glucose and fructose as.

23. Assertion : Treatment of *D*-glucose with alkali affords an equilibrium mixture consisting of *D*-mannose, *D*-fructose and starting substance *D*-glucose.

Reason : The reaction involves an intermediate in which hybridization of C_2 changes from sp^3 to sp^2 .

Answers

Carbohydrates

1	d	2	b	3	d	4	d	5	c
6	c	7	c	8	d	9	b	10	d
11	c	12	d	13	c	14	c	15	c
16	a	17	a	18	b	19	b	20	d
21	a	22	a	23	b	24	d	25	a
26	d	27	d	28	a	29	c	30	b
31	c	32	a	33	d	34	d	35	c
36	cd	37	d	38	a	39	c	40	c
41	b	42	c	43	c	44	a	45	b
46	a	47	b	48	d	49	a	50	c
51	a	52	c	53	c	54	a	55	c
56	b	57	d	58	c	59	a	60	c
61	b	62	d	63	a	64	d	65	d
66	d	67	d	68	c	69	b	70	c
71	e	72	b	73	b	74	a	75	b
76	c	77	c	78	a	79	b	80	d
81	a	82	c	83	c	84	b	85	a
86	c	87	b	88	c	89	a	90	a
91	a	92	c	93	a	94	c	95	c
96	a	97	a	98	d	99	b	100	b
101	c	102	a	103	a	104	a	105	b
106	c	107	d	108	c	109	c	110	a
111	c	112	c	113	b	114	a	115	b
116	c	117	c	118	b	119	d	120	d
121	c	122	c	123	c	124	a	125	a
126	b	127	b	128	b	129	a	130	d
131	b	132	b	133	a				



1462 Biomolecules

Proteins, Amino Acids and Enzymes

1	b	2	c	3	a	4	a	5	c
6	b	7	c	8	d	9	c	10	c
11	c	12	b	13	a	14	b	15	c
16	a	17	d	18	c	19	d	20	d
21	c	22	b	23	c	24	b	25	a
26	a	27	d	28	a	29	a	30	d
31	c	32	a	33	b	34	c	35	d
36	b	37	a	38	d	39	c	40	a
41	d	42	d	43	b	44	d	45	d
46	d	47	b	48	b	49	d	50	b
51	a	52	d	53	b	54	c	55	d
56	d	57	a	58	b	59	c	60	b
61	c	62	d	63	b	64	c	65	c
66	b	67	d	68	d	69	c	70	b
71	b	72	c	73	a	74	c	75	c
76	a	77	d	78	d	79	c	80	a
81	b	82	e	83	d	84	c	85	c
86	c	87	c	88	e	89	b	90	c
91	a	92	c	93	b	94	a	95	a
96	d	97	a	98	a	99	c		

Fats and Lipids

1	c	2	a	3	d	4	c	5	c
6	c	7	a	8	c	9	b	10	c
11	d	12	c	13	d	14	c	15	bd
16	c	17	a	18	a	19	c	20	c
21	b	22	a	23	d	24	a	25	c
26	a	27	d						

Vitamins, Hormone and Nucleic Acid

1	d	2	a	3	c	4	c	5	c
6	d	7	a	8	a	9	d	10	c
11	a	12	d	13	a	14	b	15	d
16	b	17	a	18	c	19	b	20	c
21	c	22	a	23	a	24	a	25	d
26	b	27	d	28	a	29	b	30	d
31	b	32	a	33	b	34	a	35	a
36	c	37	a	38	d	39	b	40	b
41	b	42	a	43	c	44	c	45	b
46	a								

Critical Thinking Questions

1	b	2	c	3	c	4	c	5	b
6	d	7	b	8	b	9	c	10	d
11	a	12	d	13	d	14	d	15	b
16	a	17	d	18	b	19	c	20	d
21	d	22	d	23	c	24	b	25	b
26	b								

Assertion and Reason

1	a	2	b	3	b	4	e	5	d
6	a	7	b	8	d	9	c	10	c
11	a	12	b	13	b	14	b	15	b
16	b	17	e	18	e	19	b	20	e
21	e	22	c	23	a				

AS Answers and Solutions

Carbohydrates

- (d) α -D-Glucose \rightleftharpoons Equilibrium mixture \rightleftharpoons β -D-Glucose

$\frac{[\alpha]_D = +112^\circ}{(36\%)}$
 $\frac{[\alpha]_D = +52^\circ}{(0.02\%)}$
 $\frac{[\alpha]_D = +19^\circ}{(64\%)}$


Glucose has two forms α and β . When either of these two is dissolved in water and allowed to stand, it gets converted to an equilibrium mixture of α and β forms.
- (b) Gun-cotton is a nitrocellulose or cellulose trinitrate which is used in explosive and as a binder for solid rocket propellant.
- (d) Arabinose is an aldopentose

$$HOCH_2 - (CHOH)_3 - CHO$$
- (d) In proteins amide group is present

$$\left(-NH - \underset{\substack{| \\ R}}{CH} - \underset{\substack{|| \\ O}}{C} - NH - \underset{\substack{| \\ R}}{CH} - C - \right)_n$$


Amino or peptide bond
- (c) Inulin is a carbohydrate which is stored in "Roots of Dahliya".
- (c) Carbohydrates are hydrates of carbon. Their general formula is $C_x(H_2O)_y$.
- (d) Glucose + Tollen's reagent \rightarrow Gluconic acid + Ag-mirror.
- (b) Protein gives blue-violet colour with ninhydrin

59. (c) Zwitter ion is a dipolar ion containing both a positive and negative charge in the following




The basic group CO_2^- and acidic group NH_3^+ .

63. (b) The amino acids which can't be synthesised by human body so they are essential to take from diet. They are 10 in number.
66. (b) Biological catalysts are enzymes and all enzymes are nucleic acid.
68. (d) Protein is a body building substance not energy giving substance.
76. (a) Na^+ and K^+ controls blood pressure and heart beat so excess of Na^+ ion increases B.P.
80. (a) This protein is found in hair, nail, muscle etc.
81. (b) The peptide bond is formed between two amino acids by the elimination of a water molecule. A dipeptide contains one peptide linkage. A tripeptide contains two peptide linkages. Similarly, a nanopptide contains 8 peptide linkages.
91. (a) Synthesis of polypeptide known as translation. For this process three type of RNA essential.
92. (c) Protein is used in our body as a fuel for muscles and nerves and to build and repair body tissues.
93. (b) Zymase enzyme convert glucose into alcohol. It is found in the yeast.



94. (a) In peptide linkage i.e., $-\text{CONH}-$ group, the carboxyl group of one amino acid molecules forms an amide by combination with the amino group of the next amino acid molecule with the liberation of water molecule.



95. (a) Four Fe^{2+} ions of each haemoglobin can bind with 4 molecules of O_2 and it is carried as oxyhaemoglobin.
- $$\text{Hb}_4 + 4\text{O}_2 \rightarrow \text{Hb}_4\text{O}_8$$
96. (d) Enzyme are shape selective specific biological catalyst which normally functions effectively at body temperature.
97. (a) An enzyme (protein) is a biological catalyst.
98. (a) Amino acid synthesis was done by stainley millar.

Fats and Lipids

4. (c) Acid value is the number of 1 mg of KOH required to neutralise 1 gm of the fat or oil.
5. (c) Saponification value is the number of mg of KOH required to neutralize the fatty acid resulting from the complete hydrolysis of 1gm. of oil or fat.
6. (c) Iodine number is the number of *gms* of I_2 which combine with 100 *gm* of oil or fat. It shows the degree of unsaturation of acids in fat or oil.
7. (a) Oil (unsaturated) + $\text{H}_2 \xrightarrow{\text{Ni}}$ Fat (saturated)
8. (c) Oil + NaOH (alkali) $\xrightarrow{\text{Saponification}}$ Glycerol + Soap
11. (d) Fats are called energy bank of the body. Stored below the dermis as subcutaneous fats.
12. (c)
13. (b) 1gm carbohydrate on oxidation gives 17 kJ of energy while 1 gm fat provide 37 kJ of energy.
18. (a) Waxes are esters of higher fatty acids.
19. (c) Oil/fat + Alkali $\xrightarrow{\text{Saponification}}$ Soap + Glycerol.
21. (b)
$$\begin{array}{c} \text{CH}_2\text{OOCR} \\ | \\ \text{CHOOCR} \\ | \\ \text{CH}_2\text{OOCR} \\ \text{Lipid or Triglyceride} \end{array} + \text{H}_2\text{O} \rightarrow \begin{array}{c} \text{CH}_2\text{OH} \\ | \\ \text{CHOH} \\ | \\ \text{CH}_2\text{OH} \\ \text{Glycerol or Trihydric alcohol} \end{array} + 3\text{R}-\text{COOH}$$
 Monocarboxylic acid
26. (a) Oleic acid, stearic acid and palmitic acids are produced by the hydrolysis of fats and the acid produced by hydrolysis of fats are called fatty acid.
27. (d) Oleic acid - $\text{C}_{17}\text{H}_{33}\text{COOH}$, linoleic acid - $\text{C}_{17}\text{H}_{31}\text{COOH}$, linolenic acid - $\text{C}_{17}\text{H}_{29}\text{COOH}$, palmitic acid - $\text{C}_{15}\text{H}_{31}\text{COOH}$.

Saturated monocarboxylic acids form a homologous series which has a general formula $\text{C}_n\text{H}_{2n+1}\text{COOH}$ or $\text{C}_n\text{H}_{2n}\text{O}_2$. Only palmitic acid follows this.

Vitamin, Hormone and Nucleic acid


1. (d) $\text{Nitrogen base} + \text{Sugar} + \text{Phosphate}$
Nucleoside
3. (c) Adenine is a purine base common in both RNA and DNA.
4. (c) Insulin hormone is secreted by pancreas.

8. (a)
- Ascorbic acid

1466 Biomolecules


19. (b) Adenine = Thymine, Guanine \equiv Cytosine
2 hydrogen bonds 3 hydrogen bonds
21. (c) Vitamin B_1 is thiamine. Its main source is cereals.
22. (a) Gene is a part of the DNA molecule that codes for a specific protein.
25. (d) Cortisone is not a sex hormone, it regulates metabolism of fats, carbohydrates, proteins etc.
27. (d) Thymine is present in DNA while in RNA there is Uracil.
28. (a) Mutation is a chemical change in the sequence of Nitrogenous bases along the DNA strand which can lead to the synthesis of protein with altered amino acid sequence.
39. (b) Insulin is a hormone secreted by the pancreas that lowers blood glucose level by promoting the uptake of glucose by cells and the conversion of glucose to glycogen by the liver and skeletal muscle.
40. (b) Insulin is a proteinaceous hormone secreted by β cells by islet of Langerhans of pancreas in our body.
41. (b) Codon is present in *m*-RNA, which is responsible for translation.
42. (a) Energy is stored in our body in the form of A.T.P
43. (c) Nucleic acid is a polymer of nucleotides.
44. (c) Nucleoside on hydrolysis gives an aldopentose and a heterocyclic base purine and pyrimidine.
45. (b) An alternation in the base sequence of nucleic acid molecule is called mutation which can be by radioactive ray, by adaptation etc.
46. (a) Vitamin B_6 is called pyridoxin. It is found in fruits, green-vegetables, milk, etc. Due to its deficiency, anaemia disease is caused.
5. (b) According to Chargaff's rule amount of adenine(A) is equal to that of thymine(T) and the amount of guanine(G) is equal to that of cytosine(C).
6. (d) Multiplication of DNA is called replication.
7. (b) Insulin is a hormone which decreases sugar level in the blood.
8. (b) Oxytocin hormone secreted by posterior pituitary gland plays an important role in child birth and milk ejection for feeding baby.
9. (c) Except oleic acid stearic acid, Lauric acid and Palmitic acid are saturated fatty acid. Oleic acid is unsaturated fatty acid.
10. (d) CO (III) Transition metal is present in vitamin B_{12} .
11. (a) 130 molecules of ATP produced in the lipid metabolism of a molecule of palmitic acid.
12. (d) Protein is insoluble in benzene.
13. (d)
$$\begin{array}{ccc} \begin{array}{c} H \quad | \quad OH \\ | \quad | \\ CH_2O - C - COR \\ | \quad | \\ CHO - C - COR \\ | \quad | \\ CH_2O - C - COR \end{array} & \xrightarrow[\text{Hydrolysis}]{\text{Lipase}} & \begin{array}{c} CH_2OH \\ | \\ CHOH \\ | \\ CH_2OH \\ \text{Glycerol} \end{array} + R - COOH \\ & & \text{Fatty acid} \end{array}$$
14. (d) α -helix structure is formed when the chain of α -amino acid coil as a right handed screw because of the formation of hydrogen bonds between amide groups of the same peptide chain i.e., NH group in one unit is linked to carbonyl oxygen of the third unit by hydrogen bonding. This H-bonding is responsible for holding helix in a position.
15. (b) Three types of chemicals enter the composition of all membranes proteins, lipids and carbohydrates, proteins content varies from 46-76% lipids 20-53%, while Carbohydrate content is 1-8%
16. (c) Difference in mass of compound
 $= 390 - 180 = 210$
 wt. of CH_3CO - group is = 43
 Therefore no. of $-NH_2$ group = $\frac{210}{43} = 4.88 = 5$.
17. (d) Six type of tripeptide molecules are formed.
18. (b) Amylose is a polysaccharide.
19. (c) Aspartic acid is an amino acid with acidic side chain.

Critical Thinking Questions




This structure of β -D glucose has four asymmetric carbon atom

2. (c) It is Guanine having two possible binding site.
3. (c) Four sub units are present in haemoglobin.
4. (c) The four bases in *m*-RNA : adenine, cytosine, guanine and Uracil have been shown to act in the form of triplet; each triplet behaving as a
- code for the synthesis of a particular amino acid.
5. (b) According to Chargaff's rule amount of adenine(A) is equal to that of thymine(T) and the amount of guanine(G) is equal to that of cytosine(C).
6. (d) Multiplication of DNA is called replication.
7. (b) Insulin is a hormone which decreases sugar level in the blood.
8. (b) Oxytocin hormone secreted by posterior pituitary gland plays an important role in child birth and milk ejection for feeding baby.
9. (c) Except oleic acid stearic acid, Lauric acid and Palmitic acid are saturated fatty acid. Oleic acid is unsaturated fatty acid.
10. (d) CO (III) Transition metal is present in vitamin B_{12} .
11. (a) 130 molecules of ATP produced in the lipid metabolism of a molecule of palmitic acid.
12. (d) Protein is insoluble in benzene.
13. (d)
$$\begin{array}{ccc} \begin{array}{c} H \quad | \quad OH \\ | \quad | \\ CH_2O - C - COR \\ | \quad | \\ CHO - C - COR \\ | \quad | \\ CH_2O - C - COR \end{array} & \xrightarrow[\text{Hydrolysis}]{\text{Lipase}} & \begin{array}{c} CH_2OH \\ | \\ CHOH \\ | \\ CH_2OH \\ \text{Glycerol} \end{array} + R - COOH \\ & & \text{Fatty acid} \end{array}$$
14. (d) α -helix structure is formed when the chain of α -amino acid coil as a right handed screw because of the formation of hydrogen bonds between amide groups of the same peptide chain i.e., NH group in one unit is linked to carbonyl oxygen of the third unit by hydrogen bonding. This H-bonding is responsible for holding helix in a position.
15. (b) Three types of chemicals enter the composition of all membranes proteins, lipids and carbohydrates, proteins content varies from 46-76% lipids 20-53%, while Carbohydrate content is 1-8%
16. (c) Difference in mass of compound
 $= 390 - 180 = 210$
 wt. of CH_3CO - group is = 43
 Therefore no. of $-NH_2$ group = $\frac{210}{43} = 4.88 = 5$.
17. (d) Six type of tripeptide molecules are formed.
18. (b) Amylose is a polysaccharide.
19. (c) Aspartic acid is an amino acid with acidic side chain.




Lysine, Arginine and Histidine all are basic amino acids.

20. (d) Lucas test is used for the distinction of alcohols.
21. (d) Because their Ca^{++} / Mg^{++} salts are insoluble in water.
22. (d) Glucose on reaction with bromine water followed by oxidation gives gluconic acid



23. (c) $NH_3^+ - CH_2 - COO^-$
dipolar ion
(Zwitterion or internal salt)

24. (b) Ribose is an example of aldopentose.



Two form of D-Glucopyranose are α -D(+)-Glucopyranose and β -D(+)-Glucopyranose. These are anomers (a pair of stereoisomers which differ in configuration only around C_1 are called anomers).

26. (b) Sucrose is composed of α -D-glucopyranose unit and a β -D-fructofuranose unit. These units are joined by α - β -glycosidic linkage between C-1 of the glucose unit and C-2 of the fructose unit.

Assertion and Reason

1. (a) Glycine is an amino acid, it contains both NH_2 as well as $-COOH$ groups and therefore, its aqueous solution form Zwitter ion which is amphoteric in nature.



Therefore, both assertion and reason are true.

2. (b) Hydrolysis of sucrose is known as inversion of canesugar because sucrose produce equimolecular mixture of glucose and fructose. Sucrose is dextro-rotatory which glucose and fructose mixture is laevorotatory. Sucrose is disaccharide.
3. (b) Proteins on hydrolysis gives α -amino acid because amino acids are the building block of

proteins. It is also fact that amino acids contain both $-NH_2$ and $-COOH$ group.

Here assertion and reason both are correct but reason is not a correct explanation of assertion.

4. (e) Sucrose does not undergo mutarotation. Glucose and fructose shows mutarotation because they have two forms α and β . It is fact that sucrose is a disaccharide. Therefore, assertion is false but reason is true.
5. (d) Here, both the reason and assertion are false, DNA occurs in nucleus of the cell while RNA is found mainly in cytoplasm of the cell. On heating, enzymes lose their specific activity.
6. (a) All amino acid posses amino as well as carboxylic group. $-NH_2$ group is basic while $-COOH$ group is acidic. Therefore, they behave as zwitter ion (dipolar ion). Here, both assertion and reason are true and reason is a correct explanation.
7. (b) The assertion that activity of an enzyme is pH dependent is correct. The reason that change in pH affects the solubility of enzyme in water is also true but the reason is not the correct explanation of assertion. Change in pH cause denaturation of enzyme.
8. (d) Glycosides are formed by treating glucose with methanol in presence of dry HCl gas. They cannot be hydrolysed in acidic conditions. They are not acetals but they are hemiacetals.
9. (c) The assertion is correct that haemoglobin is an oxygen carrier and the reason that oxygen binds as O_2^- to Fe of haemoglobin is incorrect, because oxygen binds as O_2 to Fe of haeme part.
10. (c) Carboxypeptidase is an exopeptidase because it breaks the peptide chain at terminal ends. Carboxypeptidase cleaves carboxy-terminal amino acids that have aromatic or branched aliphatic side chains.
11. (a) Sucrose is a non reducing sugar as it does not reduce Tollen's or Fehling's reagent, due to absence of free aldehyde of ketone group. It contains stable acetal or ketal structure which cannot be opened into a free carboxyl group.

1468 Biomolecules

Sugar is composed of α -D-glucopyranose unit and β -D-fructo furanose unit. These units are joined by α - β -glycosidic linkage between C- 1 of the glucose unit and C- 2 of the fructose unit.

12. (b) Carbohydrates which upon hydrolysis yield two molecules of the same or different monosaccharides are called disaccharides. For example, sucrose on acid hydrolysis give one molecule of glucose and fructose.
13. (b) Fructose on warming with dilute alkali, gives rise to an equilibrium mixture of glucose, fructose and mannose. The ability of fructose to reduce Fehling solution and Tollen's reagent is probably due to the isomerisation of fructose to glucose and mannose (this is called Lobry de Bruyn and Elkenstein rearrangement).
14. (b) Glucose exists in two forms, *i.e.*, α -D-glucose with a specific rotation of $+112^\circ$ and β -D-glucose with a specific rotation of $+19^\circ$. However, when either of these two forms is dissolved in water and allowed to stand. it gets converted into the same equilibrium mixture of both the α - and β -forms with a small amount of open chain form. As a result of this equilibrium, the specific rotation of a freshly prepared solution of α -glucose decreases from $+112^\circ$ to 52.7° while that of β glucose increases from $+19^\circ$ to 52.7° .
15. (b) In acidic medium $-COOH$ group acts as the base and accepts a proton. As a result, α -amino acids exist as cations in acidic medium and migrate towards cathode under the influence of an electric field. In alkaline medium NH_3^+ group acts as the acid and thus loses a proton. As a result, α -amino acids exist as anion and migrate towards anode under the influence of an electric field. However at some intermediate value of pH , the concentration of cationic form and anionic form will become equal and hence there is no net migration of α -amino acid under the influence of an electric field.
16. (b) Valine is an essential amino acid. The amino acids which the body cannot synthesize are called essential amino acid.
17. (e) Sequence of bases in DNA is TGAACCCTT. Since according to base-pairing principle, *T* in DNA faces *A* in *m*-RNA, while *G* faces *C* and *A* faces *U*. Therefore, sequence of bases in *m*-RNA is ACUUGGGAA.
18. (e) Millon's test is a test for proteins. When Millon's reagent is added to the aqueous solution of a protein, a white precipitate is formed.
19. (b) ATP has four negatively charged oxygen atoms very close to each other. So the repulsive forces between them is high. On hydrolysis of ATP, a $H_2PO_4^-$ ion is eliminated and the number of negatively charged oxygen atoms decreases. Thus, repulsive forces decreases and a large amount of energy is set free. When ATP changes to ADP, which in turn changes into AMP, energy is released at each step. This is how ATP can act as a source of energy.
20. (e) Solubility of protein is maximum at the isoelectric point.
21. (e) Amino acids are soluble in polar solvents like H_2O , $NaOH$ and HCl and insoluble in non-polar solvents like benzene, ether etc.
22. (c) Sucrose on hydrolysis gives equal amounts of glucose and fructose. Since glucose has less positive and fructose has more negative magnitude of rotation, therefore, change in the sign of rotation is observed.

