

1.1 Logarithms

1.1.1 Definition

"The Logarithm of a given number to a given base is the index of the power to which the base must be raised in order to equal the given number."

If $a > 0$ and $a \neq 1$, then logarithm of a positive number N is defined as the index x of that power of ' a ' which equals N i.e., **$\log_a N = x$ iff $a^x = N \Rightarrow a^{\log_a N} = N, a > 0, a \neq 1$ and $N > 0$**

It is also known as fundamental logarithmic identity.

The function defined by $f(x) = \log_a x, a > 0, a \neq 1$ is called logarithmic function.

Its domain is $(0, \infty)$ and range is \mathbb{R} . a is called the base of the logarithmic function.

When base is ' e ' then the logarithmic function is called natural or Napierian logarithmic function and when base is 10, then it is called common logarithmic function.

Note : The logarithm of a number is unique i.e. No number can have two different log to a given base.

$$\text{□ } \log_e a = \log_e 10 \cdot \log_{10} a \text{ or } \log_{10} a = \frac{\log_e a}{\log_e 10} = 0.434 \log_e a$$

1.1.2 Characteristic and Mantissa

(1) The integral part of a logarithm is called the characteristic and the fractional part is called mantissa.

$$\begin{array}{ccc} \log_{10} N = \text{integer} & + & \text{fraction (+ve)} \\ \downarrow & & \downarrow \\ \text{Characteristics} & & \text{Mantissa} \end{array}$$

(2) The mantissa part of log of a number is always kept positive.

(3) If the characteristics of $\log_{10} N$ be n , then the number of digits in N is $(n+1)$

(4) If the characteristics of $\log_{10} N$ be $(-n)$ then there exists $(n-1)$ number of zeros after decimal part of

N .



Example: 1 For $y = \log_a x$ to be defined 'a' must be

[IIT 1990]

- (a) Any positive real number (b) Any number
(c) $\geq e$ (d) Any positive real number $\neq 1$

Solution: (d) It is obvious (Definition).

Example: 2 Logarithm of $32\sqrt[3]{4}$ to the base $2\sqrt{2}$ is

- (a) 3.6 (b) 5 (c) 5.6 (d) None of these

Solution: (a) Let x be the required logarithm, then by definition $(2\sqrt{2})^x = 32\sqrt[3]{4}$

$$(2 \cdot 2^{1/2})^x = 2^5 \cdot 2^{2/3}; \therefore 2^{\frac{3x}{2}} = 2^{5+\frac{2}{3}}$$

$$\text{Here, by equating the indices, } \frac{3}{2}x = \frac{27}{5}, \therefore x = \frac{18}{5} = 3.6$$

1.1.3 Properties of Logarithms

Let m and n be arbitrary positive numbers such that $a > 0$, $a \neq 1$, $b > 0$, $b \neq 1$ then

- (1) $\log_a a = 1$, $\log_a 1 = 0$ (2) $\log_a b \cdot \log_b a = 1 = \log_a a = \log_b b \Rightarrow \log_a b = \frac{1}{\log_b a}$
 (3) $\log_c a = \log_b a \cdot \log_c b$ or $\log_c a = \frac{\log_b a}{\log_b c}$ (4) $\log_a (mn) = \log_a m + \log_a n$
 (5) $\log_a \left(\frac{m}{n}\right) = \log_a m - \log_a n$ (6) $\log_a m^n = n \log_a m$
 (7) $a^{\log_a m} = m$ (8) $\log_a \left(\frac{1}{n}\right) = -\log_a n$
 (9) $\log_{a^\beta} n = \frac{1}{\beta} \log_a n$ (10) $\log_{a^\beta} n^\alpha = \frac{\alpha}{\beta} \log_a n$, ($\beta \neq 0$)
 (11) $a^{\log_c b} = b^{\log_c a}$, ($a, b, c > 0$ and $c \neq 1$)

Example: 3 The number $\log_2 7$ is

[IIT 1990]

- (a) An integer (b) A rational number (c) An irrational number (d) A prime number

Solution: (c) Suppose, if possible, $\log_2 7$ is rational, say p/q where p and q are integers, prime to each other.

$$\text{Then, } \frac{p}{q} = \log_2 7 \Rightarrow 7 = 2^{p/q} \Rightarrow 2^p = 7^q,$$

Which is false since L.H.S is even and R.H.S is odd. Obviously $\log_2 7$ is not an integer and hence not a prime number

Example: 4 If $\log_7 2 = m$, then $\log_{49} 28$ is equal to

[Roorkee 1999]

4 Logarithms

- (a) $2(1+2m)$ (b) $\frac{1+2m}{2}$ (c) $\frac{2}{1+2m}$ (d) $1+m$

Solution: (b) $\log_{49} 28 = \frac{\log 28}{\log 49} = \frac{\log 7 + \log 4}{2 \log 7} = \frac{\log 7}{2 \log 7} + \frac{\log 4}{2 \log 7} = \frac{1}{2} + \frac{1}{2} \log_7 4 = \frac{1}{2} + \frac{1}{2} \cdot 2 \log_7 2 = \frac{1}{2} + \log_7 2 = \frac{1}{2} + m = \frac{1+2m}{2}$

Example: 5 If $\log_e \left(\frac{a+b}{2} \right) = \frac{1}{2} (\log_e a + \log_e b)$, then relation between a and b will be [UPSEAT 2000]

- (a) $a = b$ (b) $a = \frac{b}{2}$ (c) $2a = b$ (d) $a = \frac{b}{3}$

Solution: (a) $\log_e \left(\frac{a+b}{2} \right) = \frac{1}{2} (\log_e a + \log_e b) = \frac{1}{2} \log_e (ab) = \log_e \sqrt{ab}$
 $\Rightarrow \frac{a+b}{2} = \sqrt{ab} \Rightarrow a+b = 2\sqrt{ab} \Rightarrow (\sqrt{a} - \sqrt{b})^2 = 0 \Rightarrow \sqrt{a} - \sqrt{b} = 0 \Rightarrow a = b$

Example: 6 If $\log_{10} 3 = 0.477$, the number of digits in 3^{40} is [IIT 1992]

- (a) 18 (b) 19 (c) 20 (d) 21

Solution: (c) Let $y = 3^{40}$ is
 Taking log both the sides, $\log y = \log 3^{40} \Rightarrow \log y = 40 \log 3 \Rightarrow \log y = 19.08$
 \therefore Number of digits in $y = 19 + 1 = 20$

Example: 7 Which is the correct order for a given number α in increasing order [Roorkee 2000]

- (a) $\log_2 \alpha, \log_3 \alpha, \log_e \alpha, \log_{10} \alpha$ (b) $\log_{10} \alpha, \log_3 \alpha, \log_e \alpha, \log_2 \alpha$
 (c) $\log_{10} \alpha, \log_e \alpha, \log_2 \alpha, \log_3 \alpha$ (d) $\log_3 \alpha, \log_e \alpha, \log_2 \alpha, \log_{10} \alpha$

Solution: (b) Since 10, 3, e , 2 are in decreasing order
 Obviously, $\log_{10} \alpha, \log_3 \alpha, \log_e \alpha, \log_2 \alpha$ are in increasing order.

1.1.4 Logarithmic Inequalities

- (1) If $a > 1, p > 1 \Rightarrow \log_a p > 0$ (2) If $0 < a < 1, p > 1 \Rightarrow \log_a p < 0$
 (3) If $a > 1, 0 < p < 1 \Rightarrow \log_a p < 0$ (4) If $p > a > 1 \Rightarrow \log_a p > 1$
 (5) If $a > p > 1 \Rightarrow 0 < \log_a p < 1$ (6) If $0 < a < p < 1 \Rightarrow 0 < \log_a p < 1$
 (7) If $0 < p < a < 1 \Rightarrow \log_a p > 1$ (8) If $\log_m a > b \Rightarrow \begin{cases} a > m^b, & \text{if } m > 1 \\ a < m^b, & \text{if } 0 < m < 1 \end{cases}$
 (9) $\log_m a < b \Rightarrow \begin{cases} a < m^b, & \text{if } m > 1 \\ a > m^b, & \text{if } 0 < m < 1 \end{cases}$

(10) $\log_p a > \log_p b \Rightarrow a \geq b$ if base p is positive and > 1 or $a \leq b$ if base p is positive and < 1 i.e., $0 < p < 1$

In other words, if base is greater than 1 then inequality remains same and if base is positive but less than 1 then the sign of inequality is reversed.

Example: 8 If $x = \log_3 5$, $y = \log_{17} 25$, which one of the following is correct

[W.B. JEE 1993]

- (a) $x < y$ (b) $x = y$ (c) $x > y$ (d) None of these

Solution: (c) $y = \log_{17} 25 = 2 \log_{17} 5$

$$\therefore \frac{1}{y} = \frac{1}{2} \log_5 17$$

$$\frac{1}{x} = \log_5 3 = \frac{1}{2} \log_5 9$$

Clearly $\frac{1}{y} > \frac{1}{x}$, $\therefore x > y$

Example: 9 If $\log_{0.3}(x-1) < \log_{0.09}(x-1)$, then x lies in the interval

- (a) $(2, \infty)$ (b) $(-2, -1)$ (c) $(1, 2)$ (d) None of these

Solution: (a) $\log_{0.3}(x-1) < \log_{(0.3)^2}(x-1) = \frac{1}{2} \log_{0.3}(x-1)$

$$\therefore \frac{1}{2} \log_{0.3}(x-1) < 0$$

or $\log_{0.3}(x-1) < 0 = \log 1$ or $(x-1) > 1$ or $x > 2$

As base is less than 1, therefore the inequality is reversed, now $x > 2 \Rightarrow x$ lies in $(2, \infty)$.

