



Assignment

Solution of Quadratic equations

Basic Level

- A real root of the equation $\log_4\{\log_2(\sqrt{x+8}-\sqrt{x})\}=0$ is [AMU 1999]
(a) 1 (b) 2 (c) 3 (d) 4
- The roots of the equation $7^{\log_7(x^2-4x+5)}=x-1$ are
(a) 4, 5 (b) 2, -3 (c) 2, 3 (d) 3, 5
- The solution set of the equation $\log_x 2 \cdot \log_{2x} 2 = \log_{4x} 2$ is
(a) $\{2^{-\sqrt{2}}, 2^{\sqrt{2}}\}$ (b) $\{\frac{1}{2}, 2\}$ (c) $\{\frac{1}{4}, 2^2\}$ (d) None of these
- The solution of the equation $3^{\log_a x} + 3x^{\log_a 3} = 2$ is given by
(a) $3^{\log_2 a}$ (b) $3^{-\log_2 a}$ (c) $2^{\log_3 a}$ (d) $2^{-\log_3 a}$
- If $3^{x+1} = 6^{\log_2 3}$, then x is
(a) 3 (b) 2 (c) $\log_3 2$ (d) $\log_2 3$
- The solution of $|x/(x-1)| + |x| = x^2/|x-1|$ is
(a) $x \geq 0$ (b) $x > 0$ (c) $x \in (1, \alpha)$ (d) None of these
- If $2 \log(x+1) - \log(x^2-1) = \log 2$, then x equals
(a) 1 (b) 0 (c) 2 (d) 3
- The real roots of the equation $x^2 + 5|x| + 4 = 0$ are [MNR 1993]
(a) $\{-1, -4\}$ (b) $\{1, 4\}$ (c) $\{-4, 4\}$ (d) None of these
- If $|x^2 - x - 6| = x + 2$, then the values of x are [Roorkee 1982; Rajasthan PET 1992]
(a) -2, 2, -4 (b) -2, 2, 4 (c) 3, 2, -2 (d) 4, 4, 3
- $\{x \in R : |x-2| = x^2\} =$ [EAMCET 2000]
(a) $\{-1, 2\}$ (b) $\{1, 2\}$ (c) $\{-1, -2\}$ (d) $\{1, -2\}$
- If $ax^2 + bx + c = 0$, then x = [MP PET 1995]
(a) $\frac{b \pm \sqrt{b^2 - 4ac}}{2a}$ (b) $\frac{-b \pm \sqrt{b^2 - ac}}{2a}$ (c) $\frac{2c}{-b \pm \sqrt{b^2 - 4ac}}$ (d) None of these
- If $x^{2/3} - 7x^{1/3} + 10 = 0$, then x = [BIT Ranchi 1992]
(a) $\{125\}$ (b) $\{8\}$ (c) \emptyset (d) $\{125, 8\}$
- The roots of the given equation $(p-q)x^2 + (q-r)x + (r-p) = 0$ are [Rajasthan PET 1986; MP PET 1999]
(a) $\frac{p-q}{r-p}, 1$ (b) $\frac{q-r}{p-q}, 1$ (c) $\frac{r-p}{p-q}, 1$ (d) $1, \frac{q-r}{p-q}$
- The solution of the equation $x + \frac{1}{x} = 2$ will be [MNR 1983]



- (a) 2, -1 (b) $0, -1, -\frac{1}{5}$ (c) $-1, -\frac{1}{5}$ (d) None of these
15. One root of the following given equation $2x^5 - 14x^4 + 31x^3 - 64x^2 + 19x + 130 = 0$ is [MP PET 1985]
 (a) 1 (b) 3 (c) 5 (d) 7
16. The roots of the equation $x^4 - 4x^3 + 6x^2 - 4x + 1 = 0$ are [MP PET 1986]
 (a) 1, 1, 1, 1 (b) 2, 2, 2, 2 (c) 3, 1, 3, 1 (d) 1, 2, 1, 2
17. One root of the equation $(x+1)(x+3)(x+2)(x+4) = 120$ is [T.S. Rajendra 1991]
 (a) -1 (b) 2 (c) 1 (d) 0
18. If $9^x - 4 \times 3^{x+2} + 3^5 = 0$, then the solution pair is
 (a) (1, 2) (b) (2, 3) (c) (2, 4) (d) (1, 3)
19. In the equation $4^{x+2} = 2^{2x+3} + 48$, the value of x will be
 (a) $-\frac{3}{2}$ (b) -2 (c) -3 (d) 1
20. The roots of the equation $4^x - 3 \cdot 2^{x+3} + 128 = 0$ are [AMU 1985]
 (a) 1 and 2 (b) 2 and 3 (c) 3 and 4 (d) 4 and 5
21. The root of the equation $\sqrt{2x-2} + \sqrt{x-3} = 2$ is [Roorkee 1979]
 (a) 3 (b) 19 (c) 3, 19 (d) 3, -19
22. The solution of the equation $\sqrt{x+1} + \sqrt{x-1} = 0$ is [IIT 1978]
 (a) 1 (b) -1 (c) 5/4 (d) None of these
23. If $x = \sqrt{6 + \sqrt{6 + \sqrt{6 + \dots}}}$ to ∞ , then [Pb.CET 1999]
 (a) x is an irrational number (b) $2 < x < 3$ (c) $x = 3$ (d)
24. The real values of x which satisfy the equation $(5 + 2\sqrt{6})^{x^2-3} + (5 - 2\sqrt{6})^{x^2-3} = 10$ are [Kurukshetra CEE 1995; Karnataka CET 1993]
 (a) ± 2 (b) $\pm\sqrt{2}$ (c) $\pm 2, \pm\sqrt{2}$ (d) $2, \sqrt{2}$
25. If one root of the equation $a(b-c)x^2 + b(c-a)x + c(a-b) = 0$ is 1 then, its other roots is [Rajasthan PET 1986]
 (a) $\frac{a(b-c)}{b(c-a)}$ (b) $\frac{c(a-b)}{a(b-c)}$ (c) $\frac{b(c-a)}{a(b-c)}$ (d) None of these
26. The imaginary roots of the equation $(x^2 + 2)^2 + 8x^2 = 6x(x^2 + 2)$ are [Roorkee 1986]
 (a) $1 \pm i$ (b) $2 \pm i$ (c) $-1 \pm i$ (d) None of these
27. GM of the roots of the equation $x^2 - 18x + 9 = 0$ is [Rajasthan PET 1997]
 (a) 6 (b) 3 (c) -3 (d) ± 3
28. The solution set of the equation $(x+1)^2 + [x-1]^2 = (x-1)^2 + [x+1]^2$ is
 (a) $x \in R$ (b) $x \in N$ (c) $x \in I$ (d) $x \in Q$
29. $\left[\frac{1}{4}\right] + \left[\frac{1}{4} + \frac{1}{200}\right] + \left[\frac{1}{4} + \frac{1}{100}\right] + \dots + \left[\frac{1}{4} + \frac{199}{200}\right]$ is
 (a) 49 (b) 50 (c) 51 (d) None of these
30. The value of $x = \sqrt{2 + \sqrt{2 + \sqrt{2 + \dots}}}$ is [Karnataka CET 2001]
 (a) -1 (b) 1 (c) 2 (d) 3
31. If $x^2 - x + 1 = 0$, then value of x^{3n} is [DCE 1995]
 (a) -1, 1 (b) 1 (c) -1 (d) 0
32. For what value of a the curve $y = x^2 + ax + 25$ touches the x -axis
 (a) 0 (b) ± 5 (c) ± 10 (d) None of these
33. Let α, β be the roots of the quadratic equation $x^2 + px + p^3 = 0$ ($p \neq 0$). If (α, β) is a point on the parabola $y^2 = x$, then the roots of the quadratic equation are [MP PET 2000]

178 Quadratic Equations and Inequalities

- (a) 4, - 2 (b) - 4, - 2 (c) 4, 2 (d) - 4, 2
34. If expression $e^{\{(\sin^2 x + \sin^4 x + \sin^6 x + \dots) \ln 2\}}$ satisfies the equation $x^2 - 9x + 8 = 0$, find the value of $\frac{\cos x}{\cos x + \sin x}$, $0 < x < \frac{\pi}{2}$ [IIT 1999]
- (a) $\frac{1}{1 + \sqrt{3}}$ (b) $\frac{1}{1 - \sqrt{3}}$ (c) $\frac{2}{1 - \sqrt{2}}$ (d) None of these
35. The roots of equation $\frac{2x+31}{9} + \frac{x^2+7}{x^2-7} = \frac{2x+47}{9}$ are [Rajasthan PET 1994]
- (a) 3, - 3 (b) 5, - 5 (c) $\sqrt{3}, -\sqrt{3}$ (d) $\sqrt{5}, -\sqrt{5}$
36. If $x^2 + y^2 = 25, xy = 12$, then $x =$ [BIT Ranchi 1992]
- (a) {3, 4} (b) {3, - 3} (c) {3, 4, - 3, - 4} (d) {- 3, - 3}
37. The sum of all real roots of the equation $|x-2|^2 + |x-2| - 2 = 0$ is [IIT 1997; Himachal CET 2002]
- (a) 2 (b) 4 (c) 1 (d) None of these
38. A two digit number is four times the sum and three times the product of its digits. The number is [MP PET 1994]
- (a) 42 (b) 24 (c) 12 (d) 21
39. The number of real solutions of the equation $|x^2 + 4x + 3| + 2x + 5 = 0$ are [IIT 1988]
- (a) 1 (b) 2 (c) 3 (d) 4
40. The number of the real values of x for which the equality $|3x^2 + 12x + 6| = 5x + 16$ holds good is [AMU 1999]
- (a) 4 (b) 3 (c) 2 (d) 1
41. The number of real solutions of the equation $\sin e^x = 5^x + 5^{-x}$ is [IIT 1990, 2002]
- (a) 0 (b) 1 (c) 2 (d) Infinitely many
42. The number of the real solutions of the equation $-x^2 + x - 1 = \sin^4 x$ is
- (a) 1 (b) 2 (c) 0 (d) 4
43. The number of solutions of $\cos x = \frac{|x|}{80}$ is
- (a) 50 (b) 52 (c) 53 (d) None of these
44. The equation $\sqrt{x+1} - \sqrt{x-1} = \sqrt{4x-1}$ has [IIT 1997]
- (a) No solution (b) One solution (c) Two solutions (d) More than two solution
45. The number of real roots of $\sqrt{5x^2 - 6x + 8} - \sqrt{5x^2 - 6x - 7} = 1$ is [Roorkee 1984]
- (a) 1 (b) 2 (c) 3 (d) 4
46. The number of roots of the quadratic equation $8 \sec^2 \theta - 6 \sec \theta + 1 = 0$ is [Pb. CET 1989,94]
- (a) Infinite (b) 1 (c) 2 (d) 0
47. The number of values of x in the interval $[0, 5\pi]$ satisfying the equation $3 \sin^2 x - 7 \sin x + 2 = 0$ is [IIT 1998, MP PET 2000]
- (a) 0 (b) 5 (c) 6 (d) 10
48. The maximum number of real roots of the equation $x^{2n} - 1 = 0$, is [MP PET 2001]
- (a) 2 (b) 3 (c) n (d) $2n$
49. The equation $x + \frac{2}{1-x} = 1 + \frac{2}{1-x}$, has [IIT 1983; MNR 1998; Kurukshetra CEE 1993]
- (a) No real root (b) One real root (c) Two equal roots (d) Infinitely many roots
50. The number of real roots of equation $(x-1)^2 + (x-2)^2 + (x-3)^2 = 0$ is [IIT 1990; Karnataka CET 1998]
- (a) 2 (b) 1 (c) 0 (d) 3
51. The number of roots of the equation $\log(-2x) = 2 \log(x+1)$ are [AMU 2001]
- (a) 3 (b) 2 (c) 1 (d) None of these
52. Number of real roots of the equation $\sum_{r=1}^{10} (x-r)^3 = 0$ is
- (a) 0 (b) 1 (c) 2 (d) 3
53. The minimum value of $|x-3| + |x-2| + |x-5|$ is
- (a) 3 (b) 7 (c) 5 (d) 9

54. Rationalised denominator of $\frac{1}{\sqrt{2} + \sqrt{3} + \sqrt{5}}$ is
 (a) $\frac{2\sqrt{3} + 3\sqrt{2} - \sqrt{30}}{12}$ (b) $\frac{3\sqrt{2} - 2\sqrt{3} - \sqrt{30}}{15}$ (c) $\frac{2\sqrt{3} - 3\sqrt{2} + \sqrt{40}}{10}$ (d) $\frac{2\sqrt{3} + 3\sqrt{2} - \sqrt{20}}{15}$
55. If $x = \sqrt{7 + 4\sqrt{3}}$, then $x + \frac{1}{x} =$ [EAMCET 1994]
 (a) 4 (b) 6 (c) 3 (d) 2
56. If $\log_2 x + \log_x 2 = \frac{10}{3} = \log_2 y + \log_y 2$ and $x \neq y$, then $x + y =$ [EAMCET 1994]
 (a) 2 (b) $\frac{65}{8}$ (c) $\frac{37}{6}$ (d) None of these
57. The equation $\log_e x + \log_e(1+x) = 0$ can be written as [Kurukshetra CEE 1993; MP PET 1989]
 (a) $x^2 + x - e = 0$ (b) $x^2 + x - 1 = 0$ (c) $x^2 + x + 1 = 0$ (d) $x^2 + xe - e = 0$
58. If $f(x) = 2x^3 + mx^2 - 13x + n$ and 2, 3 are roots of the equation $f(x) = 0$, then the value of m and n are [Roorkee 1990]
 (a) -5, -30 (b) -5, 30 (c) 5, 30 (d) None of these
59. The number of real solutions of the equation $e^x = x$ is
 (a) 1 (b) 2 (c) 0 (d) None of these
60. The sum of the real roots of the equation $x^2 + |x| - 6 = 0$ is
 (a) 4 (b) 0 (c) -1 (d) None of these
61. The number of values of a for which $(a^2 - 3a + 2)x^2 + (a^2 - 5a + 6)x + a^2 - 4 = 0$ is an identity in x is
 (a) 0 (b) 2 (c) 1 (d) 3
62. The number of values of the pair (a, b) for which $a(x+1)^2 + b(x^2 - 3x - 2) + x + 1 = 0$ is an identity in x is
 (a) 0 (b) 1 (c) 2 (d) Infinite
63. If $(\sqrt{2})^x + (\sqrt{3})^x = (\sqrt{13})^{x/2}$ then the number of values of x is
 (a) 2 (b) 4 (c) 1 (d) None of these
64. The number of real solutions of the equation $\frac{6-x}{x^2-4} = 2 + \frac{x}{x+2}$ is
 (a) Two (b) One (c) Zero (d) None of these
65. The number of real solutions of $\sqrt{x^2 - 4x + 3} + \sqrt{x^2 - 9} = \sqrt{4x^2 - 14x + 6}$ is
 (a) One (b) Two (c) Three (d) None of these

Advance Level

66. If $-1 \leq x < 0$, then solution of the equation $|x+1| - |x| + 3|x-1| - |x-2| = x+2$ is [IIT 1976]
 (a) 1, 5/3 (b) 5/3 (c) 1/3 (d) None of these
67. The real roots of $|x|^3 - 3x^2 + 3|x| - 2 = 0$ are [DCE 1997]
 (a) 0, 2 (b) ± 1 (c) ± 2 (d) 1, 2
68. The number of real solutions of the equation $2^{x/2} + (\sqrt{2} + 1)^x = (5 + 2\sqrt{2})^{x/2}$ is
 (a) One (b) Two (c) Four (d) Infinite
69. The number of negative integral solutions of $x^2 \cdot 2^{x+1} + 2^{x-3} + 2 = x^2 \cdot 2^{x-3+4} + 2^{x-1}$ is [DCE 1993]
 (a) 0 (b) 1 (c) 2 (d) 4
70. The equation $e^x - x - 1 = 0$ has [Kurukshetra CEE 1998]
 (a) Only one real root $x = 0$ (b) At least two real roots (c) Exactly two real roots (d)
71. The number of real roots of the equation $e^{\sin x} - e^{-\sin x} - 4 = 0$ are [IIT 1982]
 (a) 1 (b) 2 (c) Infinite (d) None of these
72. If a, b, c are positive real numbers, then the number of real roots of the equation $ax^2 + b|x| + c = 0$ is [DCE 1998, UPSEAT 1999]

180 Quadratic Equations and Inequations

- (a) 2 (b) 4 (c) 0 (d) None of these
73. The number of real solutions of equation $\log_{10}[98 + \sqrt{[x^3 - x^2 - 12x + 36]}] = 2$ are
 (a) 4 (b) 1 (c) 2 (d) 3
74. The equation $x^{(3/4)(\log_2 x)^2 + (\log_2 x) - 5/4} = \sqrt{2}$ has [IIT 1989]
 (a) At least one real solution (b) Exactly three real solutions
 (c) Exactly one irrational solution (d) All the above
75. The number of solutions of $|[x] - 2x| = 4$, where $[x]$ is the greatest integer $\leq x$, is
 (a) 2 (b) 4 (c) 1 (d) Infinite
76. Let $f(x)$ be a function defined by $f(x) = x - [x]$, $0 \neq x \in R$, where $[x]$ is the greatest integer less than or equal to x .
 then the number of solutions of $f(x) + f\left(\frac{1}{x}\right) = 1$
 (a) 0 (b) Infinite (c) 1 (d) 2
77. If m be the number of integral solutions of equation $2x^2 - 3xy - 9y^2 - 11 = 0$ and n be the number of real solutions
 of equation $x^3 - [x] - 3 = 0$, then $m =$
 (a) n (b) $2n$ (c) $n/2$ (d) $3n$
78. The set of values of c for which $x^3 - 6x^2 + 9x - c$ is of the form $(x - \alpha)^2(x - \beta)$ (α, β real) is given by
 (a) $\{0\}$ (b) $\{4\}$ (c) $\{0, 4\}$ (d) Null set
79. If $0 < a_r < 1$ for $r = 1, 2, 3, \dots, k$ and m be the number of real solutions of equation $\sum_{r=1}^k (a_r)^x = 1$ and n be the
 number of real solution of equation $\sum_{r=1}^k (x - a_r)^{101} = 0$, then
 (a) $m = n$ (b) $m \leq n$ (c) $m \geq n$ (d) $m > n$
80. Let $P_n(x) = 1 + 2x + 3x^2 + \dots + (n+1)x^n$ be a polynomial such that n is even. Then the number of real roots of
 $P_n(x) = 0$ is [DCE 1994]
 (a) 0 (b) n (c) 1 (d) None of these
81. The number of all possible triplets (a_1, a_2, a_3) such that $a_1 + a_2 \cos 2x + a_3 \sin^2 x = 0$ for all x is [IIT 1987]
 (a) Zero (b) One (c) Three (d) Infinite
82. The solutions of the equation $2x - 2[x] = 1$, where $[x]$ = the greatest integer less than or equal to x , are
 (a) $x = n + \frac{1}{2}, n \in N$ (b) $x = n - \frac{1}{2}, n \in N$ (c) $x = n + \frac{1}{2}, n \in Z$ (d) $n < x < n+1, n \in Z$
83. The number of real solutions of $1 + |e^x - 1| = e^x(e^x - 2)$ is
 (a) 0 (b) 1 (c) 2 (d) 4
84. The equation $2 \sin^2 \frac{x}{2} \cdot \cos^2 x = x + \frac{1}{x}, 0 < x \leq \frac{\pi}{2}$ has
 (a) One real solution (b) No real solution
 (c) Infinitely many real solutions (d) None of these
85. If $y \neq 0$ then the number of values of the pair (x, y) such that $x + y + \frac{x}{y} = \frac{1}{2}$ and $(x+y)\frac{x}{y} = -\frac{1}{2}$, is
 (a) 1 (b) 2 (c) 0 (d) None of these
86. The number of real solutions of the equation $\log_{0.5} x = |x|$ is
 (a) 1 (b) 2 (c) 0 (d) None of these
87. The product of all the solutions of the equation $(x-2)^2 - 3|x-2| + 2 = 0$ is
 (a) 2 (b) -4 (c) 0 (d) None of these

88. If $0 < x < 1000$ and $\left[\frac{x}{2}\right] + \left[\frac{x}{3}\right] + \left[\frac{x}{5}\right] = \frac{31}{30}x$, where $[x]$ is the greatest integer less than or equal to x , the number of possible values of x is
 (a) 34 (b) 32 (c) 33 (d) None of these
89. The solution set of $(x)^2 + (x+1)^2 = 25$, where (x) is the least integer greater than or equal to x , is
 (a) (2, 4) (b) $(-5, -4] \cup (2, 3]$ (c) $[-4, -3) \cup (3, 4]$ (d) None of these
90. If $[x]^2 = [x+2]$, where $[x]$ = the greatest integer less than or equal to x , then x must be such that
 (a) $x = 2, -1$ (b) $x \in [2, 3)$ (c) $x \in [-1, 0)$ (d) None of these
91. The solution set of $\left|\frac{x+1}{x}\right| + |x+1| = \frac{(x+1)^2}{|x|}$ is
 (a) $\{x \mid x \geq 0\}$ (b) $\{x \mid x > 0\} \cup \{-1\}$ (c) $\{-1, 1\}$ (d) $\{x \mid x \geq 1 \text{ or } x \leq -1\}$
92. If $a \cdot 3^{\tan x} + a \cdot 3^{-\tan x} - 2 = 0$ has real solutions, $x \neq \frac{\pi}{2}, 0 \leq x \leq \pi$, then the set of possible values of the parameter a is
 (a) $[-1, 1]$ (b) $[-1, 0)$ (c) $(0, 1]$ (d) $(0, +\infty)$

Nature of roots

Basic Level

93. The roots of the quadratic equation $2x^2 + 3x + 1 = 0$, are [IIT 1983]
 (a) Irrational (b) Rational (c) Imaginary (d) None of these
94. The roots of the equation $x^2 + 2\sqrt{3}x + 3 = 0$ are [Rajasthan PET 1986]
 (a) Real and equal (b) Rational and equal (c) Irrational and equal (d) Irrational and unequal
95. If l, m, n are real and $l \neq m$, then the roots of the equation $(l-m)x^2 - 5(l+m)x - 2(l-m) = 0$ are [IIT 1979; Rajasthan PET 1983]
 (a) Complex (b) Real and distinct (c) Real and equal (d) None of these
96. If a and b are the odd integers, then the roots of the equation $2ax^2 + (2a+b)x + b = 0, a \neq 0$, will be [Pb. CET 1988]
 (a) Rational (b) Irrational (c) Non-real (d) Equal
97. If $k \in (-\infty, -2) \cup (2, \infty)$, then the roots of the equation $x^2 + 2kx + 4 = 0$ are [DCE 2002]
 (a) Complex (b) Real and unequal (c) Real and equal (d) One real and one imaginary
98. Let a, b and c be real numbers such that $4a + 2b + c = 0$ and $ab > 0$. Then the quadratic equation $ax^2 + bx + c = 0$ has [IIT 1990]
 (a) Real roots (b) Complex roots (c) Purely imaginary roots (d) Only one root
99. If $a < b < c < d$, then the roots of the equation $(x-a)(x-c) + 2(x-b)(x-d) = 0$ are [IIT 1984]
 (a) Real and distinct (b) Real and equal (c) Imaginary (d) None of these
100. If $b_1 b_2 = 2(c_1 + c_2)$, then at least one of the equations $x^2 + b_1 x + c_1 = 0$ and $x^2 + b_2 x + c_2 = 0$ has
 (a) Real roots (b) Purely imaginary roots (c) Imaginary roots (d) None of these
101. In the equation $x^3 + 3Hx + G = 0$, if G and H are real and $G^2 + 4H^3 > 0$, then the roots are [Karnataka CET 2000]
 (a) All real and equal (b) All real and distinct (c) One real and two imaginary (d)
102. The equation $(x-a)^3 + (x-b)^3 + (x-c)^3 = 0$, has
 (a) All the roots real (b) One real and two imaginary roots
 (c) Three real roots namely $x = a, x = b, x = c$ (d) None of these
103. For the equation $|x^2| + |x| - 6 = 0$, the roots are [EAMCET 1988, 93]
 (a) One and only one real number (b) Real with sum one
 (c) Real with sum zero (d) Real with product zero
104. If $a > 0, b > 0, c > 0$, then both the roots of the equation $ax^2 + bx + c = 0$ [IIT 1980]
 (a) Are real and negative (b) Have negative real parts (c) Are rational numbers (d) None of these

182 Quadratic Equations and Inequations

- 105.** Let one root of $ax^2 + bx + c = 0$, where a, b, c are integers be $3 + \sqrt{5}$, then the other root is [MNR 1982]
 (a) $3 - \sqrt{5}$ (b) 3 (c) $\sqrt{5}$ (d) None of these
- 106.** If $2 + i$ is a root of the equation $x^3 - 5x^2 + 9x - 5 = 0$, then the other roots are [Kerala (Engg.) 2002]
 (a) 1 and $2 - i$ (b) -1 and $3 + i$ (c) 0 and 1 (d) -1 and $i - 2$
- 107.** If a, b, c are nonzero, unequal rational numbers then the roots of the equation $abc^2x^2 + (3a^2 + b^2)cx - 6a^2 - ab + 2b^2 = 0$ are
 (a) Rational (b) Imaginary (c) Irrational (d) None of these
- 108.** The equation $x^2 - 6x + 8 + \lambda(x^2 - 4x + 3) = 0$, $\lambda \in R$, has
 (a) Real and unequal roots for all λ (b) Real roots for $\lambda < 0$ only
 (c) Real roots for $\lambda > 0$ only (d) Real and unequal roots for $\lambda = 0$ only
- 109.** If $a > 1$, roots of the equation $(1 - a)x^2 + 3ax - 1 = 0$ are
 (a) One positive and one negative (b) Both negative
 (c) Both positive (d) Both nonreal complex
- 110.** If the roots of the equation $ax^2 + x + b = 0$ be real, then the roots of the equation $x^2 - 4\sqrt{ab}x + 1 = 0$ will be
 (a) Rational (b) Irrational (c) Real (d) Imaginary
- 111.** If the roots of the equation $x^2 - 8x + (a^2 - 6a) = 0$ are real, then [Rajasthan PET 1987, 97]
 (a) $-2 < a < 8$ (b) $2 < a < 8$ (c) $-2 \leq a \leq 8$ (d) $2 \leq a \leq 8$
- 112.** If the roots of the given equation $(\cos p - 1)x^2 + (\cos p)x + \sin p = 0$ are real, then [IIT 1990; Rajasthan PET 1995]
 (a) $p \in (-\pi, 0)$ (b) $p \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ (c) $p \in (0, \pi)$ (d) $p \in (0, 2\pi)$
- 113.** The greatest value of a non-negative real number λ for which both the equations $2x^2 + (\lambda - 1)x + 8 = 0$ and $x^2 - 8x + \lambda + 4 = 0$ have real roots is [AMU 1990]
 (a) 9 (b) 12 (c) 15 (d) 16
- 114.** If p, q, r are positive and are in A.P., then roots of the equation $px^2 + qx + r = 0$ are real if [IIT 1995]
 (a) $\left|\frac{r}{p} - 7\right| \geq 4\sqrt{3}$ (b) $\left|\frac{p}{r} - 7\right| \geq 4\sqrt{3}$ (c) For all values of p, r (d) For no value of p, r
- 115.** Let $p, q \in \{1, 2, 3, 4\}$. The number of equations of the form $px^2 + qx + 1 = 0$ having real roots is [IIT 1994]
 (a) 15 (b) 9 (c) 7 (d) 8
- 116.** The least integer k which makes the roots of the equation $x^2 + 5x + k = 0$ imaginary is [Kerala (Engg.) 2002]
 (a) 4 (b) 5 (c) 6 (d) 7
- 117.** If $0 < a < b < c$, and the roots α, β of the equation $ax^2 + bx + c = 0$ are non-real complex numbers, then
 (a) $|\alpha| = |\beta|$ (b) $|\alpha| > 1$ (c) $|\beta| < 1$ (d) None of these
- 118.** If roots of the equation $a(b - c)x^2 + b(c - a)x + c(a - b) = 0$ are equal, then a, b, c are in [Roorkee 1993; Rajasthan PET 2001]
 (a) A.P. (b) G.P. (c) H.P. (d) None of these
- 119.** If the equation $(m - n)x^2 + (n - l)x + l - m = 0$ has equal roots, then l, m and n satisfy [DCE 2002; EAMCET 1990]
 (a) $2l = m + n$ (b) $2m = n + l$ (c) $m = n + l$ (d) $l = m + n$
- 120.** The condition for the roots of the equation $(c^2 - ab)x^2 - 2(a^2 - bc)x + (b^2 - ac) = 0$ to be equal is [TS Rajendra 1982]
 (a) $a = 0$ (b) $b = 0$ (c) $c = 0$ (d) None of these
- 121.** If the roots of the equation $(a^2 + b^2)t^2 - 2(ac + bd)t + (c^2 + d^2) = 0$ are equal, then [MP PET 1996]
 (a) $ab = dc$ (b) $ac = bd$ (c) $ad + bc = 0$ (d) $\frac{a}{b} = \frac{c}{d}$
- 122.** If one root of $x^2 + px + 12 = 0$ is 4 and roots of the equation $x^2 + px + q = 0$ are equal, then q is equal to [Rajasthan PET 1990]
 (a) $49/4$ (b) $4/49$ (c) 4 (d) None of these
- 123.** If the roots of the equation $x^2 + 2mx + m^2 - 2m + 6 = 0$ are same, then the value of m will be [MP PET 1986]

- (a) 3 (b) 0 (c) 2 (d) -1
124. If the roots of the equation $x^2 - 15 - m(2x - 8) = 0$ are equal then m is equal to [Rajasthan PET 1985]
 (a) 3, -5 (b) -3, 5 (c) 3, 5 (d) -3, -5
125. For what value of k will the equation $x^2 - (3k - 1)x + 2k^2 - 11 = 0$ have equal roots [Karnataka CET 1998]
 (a) 5 (b) 9 (c) Both the above (d) 0
126. The value of k for which the quadratic equation $kx^2 + 1 = kx + 3x - 11x^2 = 0$ has real and equal roots are [BIT Ranchi 1993]
 (a) -11, -3 (b) 5, 7 (c) 5, -7 (d) None of these
127. If the roots of $4x^2 + px + 9 = 0$ are equal, then absolute value of p is [MP PET 1995]
 (a) 144 (b) 12 (c) -12 (d) ± 12
128. The value of k for which $2x^2 - kx + x + 8 = 0$ has equal and real roots are [BIT Ranchi 1990]
 (a) -9 and -7 (b) 9 and 7 (c) -9 and 7 (d) 9 and -7
129. The roots of $4x^2 + 6px + 1 = 0$ are equal, then the value of p is [MP PET 2003]
 (a) $\frac{4}{5}$ (b) $\frac{1}{3}$ (c) $\frac{2}{3}$ (d) $\frac{4}{3}$
130. If the equation $x^2 - (2 + m)x + (m^2 - 4m + 4) = 0$ has coincident roots, then [Roorkee 1991]
 (a) $m = 0, m = 1$ (b) $m = 0, m = 2$ (c) $m = \frac{2}{3}, m = 6$ (d) $m = \frac{2}{3}, m = 1$
131. If two roots of the equation $x^3 - 3x + 2 = 0$ are same, then the roots will be [MP PET 1985]
 (a) 2, 2, 3 (b) 1, 1, -2 (c) -2, 3, 3 (d) -2, -2, 1
132. The equation $\|x - 1| + a| = 4$ can have real solutions for x if a belongs to the interval
 (a) $(-\infty, 4]$ (b) $(-\infty, -4]$ (c) $(4, \infty)$ (d) $[-4, 4]$
133. The set of values of m for which both roots of the equation $x^2 - (m + 1)x + m + 4 = 0$ are real and negative consists of all m such that [AMU 1992]
 (a) $-3 < m \leq -1$ (b) $-4 < m \leq -3$ (c) $-3 \leq m \leq 5$ (d) $-3 \geq m$ or $m \geq 5$
134. Both the roots of the given equation $(x - a)(x - b) + (x - b)(x - c) + (x - c)(x - a) = 0$ are always [MNR 1986; IIT 1980; Kurukshetra CEE 1998]
 (a) Positive (b) Negative (c) Real (d) Imaginary
135. If $P(x) = ax^2 + bx + c$ and $Q(x) = -ax^2 + dx + c$ where $ac \neq 0$, then $P(x) \cdot Q(x) = 0$, has at least [IIT 1985]
 (a) Four real roots (b) Two real roots (c) Four imaginary roots (d) None of these
136. The conditions that the equation $ax^2 + bx + c = 0$ has both the roots positive is that [SCRA 1990]
 (a) a, b and c are of the same sign (b) a and b are of the same sign
 (c) b and c have the same sign opposite to that of a (d) a and c have the same sign opposite to that of b
137. If $[x]$ denotes the integral part of x and $k = \sin^{-1} \frac{1 + t^2}{2t} > 0$, then the integral value of α for which the equation $(x - [k])(x + \alpha) - 1 = 0$ has integral roots is
 (a) 1 (b) 2 (c) 4 (d) None of these
138. If the roots of the equation $ax^2 + bx + c = 0$ are real and of the form $\frac{\alpha}{\alpha - 1}$ and $\frac{\alpha + 1}{\alpha}$, then the value of $(a + b + c)^2$ is [AMU 2000]
 (a) $b^2 - 4ac$ (b) $b^2 - 2ac$ (c) $2b^2 - ac$ (d) None of these

Advance Level

184 Quadratic Equations and Inequations

139. Equation $\frac{a^2}{x-\alpha} + \frac{b^2}{x-\beta} + \frac{c^2}{x-\gamma} = m - n^2x$ ($a, b, c, m, n \in R$) has necessarily
- (a) All the roots real (b) All the roots imaginary
(c) Two real and two imaginary roots (d) Two rational and two irrational roots
140. If $\cos \theta, \sin \phi, \sin \theta$ are in G.P. then roots of $x^2 + 2 \cot \phi x + 1 = 0$ are always
- (a) Equal (b) Real (c) Imaginary (d) Greater than 1
141. If $f(x)$ is a continuous function and attains only rational values and $f(0) = 3$, then roots of equation $f(1)x^2 + f(3)x + f(5) = 0$ are
- (a) Imaginary (b) Rational (c) Irrational (d) Real and equal
142. The roots of $ax^2 + bx + c = 0$, where $a \neq 0$ and coefficients are real, are non-real complex and $a + c < b$. Then
- (a) $4a + c > 2b$ (b) $4a + c < 2b$ (c) $4a + c = 2b$ (d) None of these
143. The equation $(a+2)x^2 + (a-3)x = 2a-1, a \neq -2$ has roots rational for
- (a) All rational values of a except $a = -2$ (b) All real values of a except $a = -2$
(c) Rational values of $a > \frac{1}{2}$ (d) None of these
144. The quadratic equation $x^2 - 2x - \lambda = 0, \lambda \neq 0$
- (a) Cannot have a real root if $\lambda < 1$
(b) Can have a rational root if λ is a perfect square
(c) Cannot have an integral root if $n^2 - 1 < \lambda < n^2 + 2n$ where $n = 0, 1, 2, 3, \dots$
(d) None of these
145. If the roots of the equation $x^2 + px + q = 0$ are α and β and roots of the equation $x^2 - xr + s = 0$ are α^4, β^4 , then the roots of the equation $x^2 - 4qx + 2q^2 - r = 0$ will be [IIT 1989]
- (a) Both negative (b) Both positive
(c) Both real (d) One negative and one positive
146. If equation $a(b-c)x^2 + b(c-a)x + c(a-b) = 0$ has equal roots, $a, b, c > 0, n \in N$, then
- (a) $a^n + c^n \geq 2b^n$ (b) $a^n + c^n > 2b^n$ (c) $a^n + c^n \leq 2b^n$ (d) $a^n + c^n < 2b^n$
147. If $\frac{\sum_{r=0}^{k-1} x^{2r}}{\sum_{r=0}^{k-1} x^r}$ is a polynomial in x for two values of p and q of k , then roots of equation $x^2 + px + q = 0$ cannot be
- (a) Real (b) Imaginary (c) Rational (d) Irrational
148. If for $x > 0, f(x) = (a - x^n)^{1/n}, g(x) = x^2 + px + q, p, q \in R$ and equation $g(x) - x = 0$ has imaginary roots, then number of real roots of equation $g(g(x)) - f(f(x)) = 0$ is
- (a) 0 (b) 2 (c) 4 (d) None of these
149. Let $p, q \in \{1, 2, 3, 4\}$. The number of equations of the form $px^2 + qx + 1 = 0$ having real and unequal roots is
- (a) 15 (b) 9 (c) 7 (d) 8
150. If α_1, α_2 and β_1, β_2 are the roots of the equations $ax^2 + bx + c = 0$ and $px^2 + qx + r = 0$ respectively and system of equations $\alpha_1 y + \alpha_2 z = 0$ and $\beta_1 y + \beta_2 z = 0$ has a non-zero solution. Then [IIT 1987]
- (a) $a^2qc = p^2br$ (b) $p^2br = q^2ac$ (c) $c^2ar = r^2pb$ (d) None of these
151. If a, b, c, d are four consecutive terms of an increasing AP then the roots of the equation $(x-a)(x-c) + 2(x-b)(x-d) = 0$ are
- (a) Real and distinct (b) Nonreal complex (c) Real and equal (d) Integers
152. If a, b, c are three distinct positive real numbers then the number of real roots of $ax^2 + 2b|x| - c = 0$ is

- (a) 4 (b) 2 (c) 0 (d) None of these
153. If $a \in R, b \in R$ then the equation $x^2 - abx - a^2 = 0$ has
 (a) One positive root and one negative root (b) Both roots positive
 (c) Both roots negative (d) Non-real roots
154. The number of integral values of a for which $x^2 - (a-1)x + 3 = 0$ has both roots positive and $x^2 + 3x + 6 - a = 0$ has both roots negative is
 (a) 0 (b) 1 (c) 2 (d) Infinite
155. The quadratic equations $x^2 + (a^2 - 2)x - 2a^2 = 0$ and $x^2 - 3x + 2 = 0$ have
 (a) No common root for all $a \in R$ (b) Exactly one common root
 for all $a \in R$
 (c) Two common roots for some $a \in R$ (d) None of these
156. If $f(x) = \frac{x^2 - 1}{x^2 + 1}$ for every real number x then the minimum value of f
 (a) Does not exist because f is unbounded (b) Is not attained even though f is bounded
 (c) Is equal to 1 (d) Is equal to -1
157. If x, y, z are real and distinct then $f(x, y) = x^2 + 4y^2 + 9z^2 - 6yz - 3zx - 2xy$ is always
 (a) Non-negative (b) Nonpositive (c) Zero (d) None of these
158. If $a \in R, b \in R$ then the factors of the expression $a(x^2 - y^2) - bxy$ are
 (a) Real and different (b) Real and identical (c) Complex (d) None of these
159. If a, b, c are in H.P. then the expression $a(b - c)x^2 + b(c - a)x + c(a - b)$
 (a) Has real and distinct factors (b) Is a perfect square
 (c) Has no real factor (d) None of these
160. If a, b, c are in G.P., where a, c are positive, then the equation $ax^2 + bx + c = 0$ has
 (a) Real roots (b) Imaginary roots
 (c) Ratio of roots = $1 : w$ where w is a nonreal cube root of unity (d) Ratio of roots = $b : ac$
161. The polynomial $(ax^2 + bx + c)(ax^2 - dx - c)$ $ac \neq 0$, has
 (a) Four real zeros (b) At least two real zeros (c) At most two real zeros (d) No real zeros

Relation between Roots and Coefficient

Basic Level

162. If α, β are roots of the equation $ax^2 + bx + c = 0$, then the value of $\alpha^3 + \beta^3$ is
 [Kurukshehra CEE 1991; BIT Ranchi 1998; MP PET 1994; Rajasthan PET 1989, 96]
 (a) $\frac{3abc + b^3}{a^3}$ (b) $\frac{a^3 + b^3}{3ab}$ (c) $\frac{3abc - b^3}{a^3}$ (d) $\frac{b^3 - 3abc}{a^3}$
163. If α, β are roots of the equation $x^2 - (1 + n^2)x + \frac{1}{2}(1 + n^2 + n^4) = 0$, then $\alpha^2 + \beta^2$ is equal to [Rajasthan PET 1996]
 (a) $2n$ (b) n^2 (c) n^3 (d) $2n^2$
164. If α and β are the roots of the equation $ax^2 + bx + c = 0$ ($a \neq 0$; a, b, c being different), then $(1 + \alpha + \alpha^2)(1 + \beta + \beta^2) =$ [DCE 2000]
 (a) Zero (b) Positive (c) Negative (d) None of these
165. If α, β are the roots of the equation $8x^2 - 3x + 27 = 0$, then the value of $\left(\frac{\alpha^2}{\beta}\right)^{\frac{1}{3}} + \left(\frac{\beta^2}{\alpha}\right)^{\frac{1}{3}}$ is [AMU 1990]
 (a) $\frac{1}{3}$ (b) $\frac{1}{4}$ (c) $\frac{7}{2}$ (d) 4
166. If α, β are the roots of the equation $x^2 + px + p^2 + q = 0$, then the value of $\alpha^2 + \alpha\beta + \beta^2 + q$ is equal to [AMU 1993]
 (a) 0 (b) 1 (c) q (d) $2q$

186 Quadratic Equations and Inequations

167. If α, β are the roots of the equation $x^2 - p(x+1) - c = 0$, then $(\alpha+1)(\beta+1) =$ [BITS Ranchi 2000; Him. CET 2001]
 (a) c (b) $c - 1$ (c) $1 - c$ (d) None of these
168. If α, β, γ are the roots of the equation $x^3 + 4x + 1 = 0$, then $(\alpha + \beta)^{-1} + (\beta + \gamma)^{-1} + (\gamma + \alpha)^{-1} =$ [EAMCET 2003]
 (a) 2 (b) 3 (c) 4 (d) 5
169. If roots of $x^2 - 7x + 6 = 0$ are α, β then $\frac{1}{\alpha} + \frac{1}{\beta} =$ [Rajasthan PET 1990, 95; MNR 1981]
 (a) $6/7$ (b) $7/6$ (c) $7/10$ (d) $8/9$
170. If α, β are the roots of $x^2 - 2x + 4 = 0$, then $\alpha^5 + \beta^5$ is equal to [EAMCET 1990]
 (a) 16 (b) 32 (c) 64 (d) None of these
171. If the roots of the equation $ax^2 + bx + c = 0$ are α, β , then the value of $\alpha\beta^2 + \alpha^2\beta + \alpha$ will be [EAMCET 1980; AMU 1984]
 (a) $\frac{c(a-b)}{a^2}$ (b) 0 (c) $-\frac{bc}{a^2}$ (d) None of these
172. If α, β be the roots of the equation $2x^2 - 35x + 2 = 0$, then the value of $(2\alpha - 35)^3 \cdot (2\beta - 35)^3$ is equal to [Bihar CEE 1994]
 (a) 1 (b) 64 (c) 8 (d) None of these
173. If α and β are roots of $ax^2 + 2bx + c = 0$, then $\sqrt{\frac{\alpha}{\beta}} + \sqrt{\frac{\beta}{\alpha}}$ is equal to [BITS Ranchi 1990]
 (a) $\frac{2b}{ac}$ (b) $\frac{2b}{\sqrt{ac}}$ (c) $-\frac{2b}{\sqrt{ac}}$ (d) $-\frac{b}{\sqrt{2}}$
174. If α, β are the roots of the equation $x^2 + 2x + 4 = 0$, then $\frac{1}{\alpha^3} + \frac{1}{\beta^3}$ is equal to [Kerala (Engg.) 2002]
 (a) $-\frac{1}{2}$ (b) $\frac{1}{2}$ (c) 32 (d) $\frac{1}{4}$
175. If α, β, γ are roots of equation $x^3 + ax^2 + bx + c = 0$, then $\alpha^{-1} + \beta^{-1} + \gamma^{-1} =$ [EAMCET 2002]
 (a) a/c (b) $-b/c$ (c) b/a (d) c/a
176. If α, β are roots of $x^2 - 3x + 1 = 0$, then the value of $\alpha^3 + \beta^3$ is [MP 1994; BIT Ranchi 1990]
 (a) 9 (b) 18 (c) -9 (d) -18
177. If A.M. of the roots of a quadratic equation is $8/5$ and A.M. of their reciprocals is $8/7$, then the equation is [AMU 2001]
 (a) $5x^2 - 16x + 7 = 0$ (b) $7x^2 - 16x + 5 = 0$ (c) $7x^2 - 16x + 8 = 0$ (d) $3x^2 - 12x + 7 = 0$
178. The quadratic in t , such that A.M. of its roots is A and G.M. is G , is [IIT 1968, 74]
 (a) $t^2 - 2At + G^2 = 0$ (b) $t^2 - 2At - G^2 = 0$ (c) $t^2 + 2At + G^2 = 0$ (d) None of these
179. In a triangle ABC , the value of $\angle A$ is given by $5 \cos A + 3 = 0$, then the equation whose roots are $\sin A$ and $\tan A$ will be [Roorkee 1972]
 (a) $15x^3 - 8x + 16 = 0$ (b) $15x^2 + 8x - 16 = 0$ (c) $15x^2 - 8\sqrt{2}x + 16 = 0$ (d) $15x^2 - 8x - 16 = 0$
180. If $x^2 + px + q = 0$ is the quadratic whose roots are $a-2$ and $b-2$ where a and b are the roots of $x^2 - 3x + 1 = 0$, then [Kerala (Engg.) 2002]
 (a) $p = 1, q = 5$ (b) $p = 1, q = -5$ (c) $p = -1, q = 1$ (d) None of these
181. The roots of the equation $x^2 + ax + b = 0$ are p and q , then the equation whose roots are p^2q and pq^2 will be [MP PET 1998]
 (a) $x^2 + abx + b^3 = 0$ (b) $x^2 - abx + b^3 = 0$ (c) $bx^2 + x + a = 0$ (d) $x^2 + ax + ab = 0$
182. The equation whose roots are $\frac{1}{3+\sqrt{2}}$ and $\frac{1}{3-\sqrt{2}}$ is [MP PET 1994]
 (a) $7x^2 - 6x + 1 = 0$ (b) $6x^2 - 7x + 1 = 0$ (c) $x^2 - 6x + 7 = 0$ (d) $x^2 - 7x + 6 = 0$
183. If α, β are the roots of the equation $lx^2 + mx + n = 0$ then the equation whose roots are $\alpha^3\beta$ and $\alpha\beta^3$ is [MP PET 1997]
 (a) $l^4x^2 - nl(m^2 - 2nl)x + n^4 = 0$ (b) $l^4x^2 + nl(m^2 - 2nl)x + n^4 = 0$
 (c) $l^4x^2 + nl(m^2 - 2nl)x - n^4 = 0$ (d) $l^4x^2 - nl(m^2 + 2nl)x + n^4 = 0$

- 184.** If α, β are the roots of $9x^2 + 6x + 1 = 0$, then the equation with the roots $\frac{1}{\alpha}, \frac{1}{\beta}$ is [EAMCET 2000]
- (a) $2x^2 + 3x + 18 = 0$ (b) $x^2 + 6x - 9 = 0$ (c) $x^2 + 6x + 9 = 0$ (d) $x^2 - 6x + 9 = 0$
- 185.** If α, β are the roots of the equation $ax^2 + bx + c = 0$, then the equation whose roots are $\alpha + \frac{1}{\beta}$ and $\beta + \frac{1}{\alpha}$, is [Rajasthan PET 2000]
- (a) $acx^2 + (a+c)bx + (a+c)^2 = 0$ (b) $abx^2 + (a+c)bx + (a+c)^2 = 0$
 (c) $acx^2 + (a+b)cx + (a+c)^2 = 0$ (d) None of these
- 186.** If α, β are the roots of $x^2 - 3x + 1 = 0$, then the equation whose roots are $\frac{1}{\alpha-2}, \frac{1}{\beta-2}$ is [Rajasthan PET 1999]
- (a) $x^2 + x - 1 = 0$ (b) $x^2 + x + 1 = 0$ (c) $x^2 - x - 1 = 0$ (d) None of these
- 187.** If α, β are the roots of $ax^2 + bx + c = 0$, then the equation whose roots are $2 + \alpha, 2 + \beta$ is [EAMCET 1994]
- (a) $ax^2 + x(4a - b) + 4a - 2b + c = 0$ (b) $ax^2 + x(4a - b) + 4a + 2b + c = 0$
 (c) $ax^2 + x(b - 4a) + 4a + 2b + c = 0$ (d) $ax^2 + x(b - 4a) + 4a - 2b + c = 0$
- 188.** If α, β are the roots of the equation $ax^2 + bx + c = 0$, then the equation with roots $1/\alpha, 1/\beta$ will be [MNR 1988; SCRA 1990; Rajasthan PET 1994]
- (a) $cx^2 - bx + a = 0$ (b) $cx^2 + bx + a = 0$ (c) $x^2 + bx + a = 0$ (d) $x^2 + bx - a = 0$
- 189.** Let α, α^2 be the roots of $x^2 + x + 1 = 0$, then the equation whose roots are α^{31}, α^{62} is [AMU 1999]
- (a) $x^2 - x + 1 = 0$ (b) $x^2 + x - 1 = 0$ (c) $x^2 + x + 1 = 0$ (d) $x^{60} + x^{30} + 1 = 0$
- 190.** If α, β are roots of the equation $x^2 - 2x \cos 2\theta + 1 = 0$ then the equation with roots $\alpha^{n/2}, \beta^{n/2}$ will be [Rajasthan PET 1998]
- (a) $x^2 - 2nx \cos \theta + 1 = 0$ (b) $x^2 + 2nx \cos n\theta + 1 = 0$ (c) $x^2 + 2x \cos n\theta + 1 = 0$ (d) $x^2 - 2x \cos n\theta + 1 = 0$
- 191.** The equation whose roots are reciprocal of the roots of the equation $3x^2 - 20x + 17 = 0$ is [DCE 2002]
- (a) $3x^2 + 20x - 17 = 0$ (b) $17x^2 - 20x + 3 = 0$ (c) $17x^2 + 20x + 3 = 0$ (d) None of these
- 192.** The sum of the roots of a equation is 2 and sum of their cubes is 98, then the equation is [MP PET 1986]
- (a) $x^2 + 2x + 15 = 0$ (b) $x^2 + 15x + 2 = 0$ (c) $2x^2 - 2x + 15 = 0$ (d) $x^2 - 2x - 15 = 0$
- 193.** Sum of roots is -1 and sum of their reciprocals is $\frac{1}{6}$, then equation is [Karnataka CET 1998]
- (a) $x^2 + x - 6 = 0$ (b) $x^2 - x + 6 = 0$ (c) $x^2 + x + 1 = 0$ (d) $x^2 - 6x + 1 = 0$
- 194.** If α, β are the roots of the quadratic equation $x^2 + bx - c = 0$, then the equation whose roots are b and c is [Pb. CET 1989]
- (a) $x^2 + \alpha x - \beta = 0$ (b) $x^2 - [(\alpha + \beta) + \alpha\beta]x - \alpha\beta(\alpha + \beta) = 0$
 (c) $x^2 + [(\alpha + \beta) + \alpha\beta]x + \alpha\beta(\alpha + \beta) = 0$ (d) $x^2 + [\alpha\beta + (\alpha + \beta)]x - \alpha\beta(\alpha + \beta) = 0$
- 195.** If α, β are roots of $x^2 - 5x - 3 = 0$, then the equation with roots $\frac{1}{2\alpha - 3}$ and $\frac{1}{2\beta - 3}$ is [Rajasthan PET 1998]
- (a) $33x^2 + 4x - 1 = 0$ (b) $33x^2 - 4x + 1 = 0$ (c) $33x^2 - 4x - 1 = 0$ (d) $33x^2 + 4x + 1 = 0$
- 196.** Given that $\tan \alpha$ and $\tan \beta$ are the roots of $x^2 - px + q = 0$, then the value of $\sin^2(\alpha + \beta) =$ [Rajasthan PET 2000]
- (a) $\frac{p^2}{p^2 + (1 - q)^2}$ (b) $\frac{p^2}{p^2 + q^2}$ (c) $\frac{q^2}{p^2 + (1 - q)^2}$ (d) $\frac{p^2}{(p + q)^2}$
- 197.** If $2 + i\sqrt{3}$ is a root of the equation $x^2 + px + q = 0$, then (p, q) is equal to [IIT 1982; MP 1997]
- (a) $(7, -4)$ (b) $(-4, 7)$ (c) $(4, 7)$ (d) $(7, 4)$
- 198.** In the equation $x^2 + px + q = 0$, the coefficient of x was taken as 17 in place of 13 and its roots were found to be -2 and -15 . The correct roots of the original equation are [Rajasthan PET 1994; IIT 1979]
- (a) $-10, -3$ (b) $10, 3$ (c) $-10, 3$ (d) $10, -3$
- 199.** Two students while solving a quadratic equation in x , one copied the constant term incorrectly and got the roots 3 and 2. The other copied the constant term and coefficient of x^2 correctly as -6 and 1 respectively. The correct roots are [EAAMCET 1991]

188 Quadratic Equations and Inequations

- (a) 3, -2 (b) -3, 2 (c) -6, -1 (d) 6, -1
- 200.** If 8, 2 are the roots of $x^2 + ax + \beta = 0$ and 3, 3 are the roots of $x^2 + \alpha x + b = 0$, then the roots of $x^2 + ax + b = 0$ are [EAMCET 2000]
- (a) 8, -1 (b) -9, 2 (c) -8, -2 (d) 9, 1
- 201.** The equation formed by decreasing each root of $ax^2 + bx + c = 0$ by 1 is $2x^2 + 8x + 2 = 0$, then [EAMCET 2000]
- (a) $a = -b$ (b) $b = -c$ (c) $c = -a$ (d) $b = a + c$
- 202.** If p and q are non-zero constants, the equation $x^2 + px + q = 0$ has roots u and v , then the equation $qx^2 + px + 1 = 0$ has roots [MNR 1988]
- (a) u and $\frac{1}{v}$ (b) $\frac{1}{u}$ and v (c) $\frac{1}{u}$ and $\frac{1}{v}$ (d) None of these
- 203.** If the sum of the roots of the equation $x^2 + px + q = 0$ is equal to the sum of their squares, then [Pb. CET 1999]
- (a) $p^2 - q^2 = 0$ (b) $p^2 + q^2 = 2q$ (c) $p^2 + p = 2q$ (d) None of these
- 204.** If the sum of the roots of the equation $x^2 + px + q = 0$ is three times their difference, then which one of the following is true [Dhanbad Engg. 1968]
- (a) $9p^2 = 2q$ (b) $2q^2 = 9p$ (c) $2p^2 = 9q$ (d) $9q^2 = 2p$
- 205.** If the sum of the roots of the quadratic equation $ax^2 + bx + c = 0$ is equal to the sum of the squares of their reciprocals, then $\frac{b^2}{ac} + \frac{bc}{a^2} =$ [BITS Ranchi 1996]
- (a) 2 (b) -2 (c) 1 (d) -1
- 206.** If the sum of the two roots of the equation $4x^3 + 16x^2 - 9x - 36 = 0$ is zero, then the roots are [MP PET 1986]
- (a) 1, 2, -2 (b) $-2, \frac{2}{3}, -\frac{2}{3}$ (c) $-3, \frac{3}{2}, -\frac{3}{2}$ (d) $-4, \frac{3}{2}, -\frac{3}{2}$
- 207.** If the roots of the equation $ax^2 + bx + c = 0$ are l and $2l$, then [MP PET 1986]
- (a) $b^2 = 9ac$ (b) $2b^2 = 9ac$ (c) $b^2 = -4ac$ (d) $a^2 = c^2$
- 208.** If α, β are the roots of the equation $x^2 - px + 36 = 0$ and $\alpha^2 + \beta^2 = 9$, then the value of p are [AMU 1991]
- (a) ± 3 (b) ± 6 (c) ± 8 (d) ± 9
- 209.** If α, β, γ are the roots of $2x^3 - 2x - 1 = 0$, then $(\sum \alpha\beta)^2 =$ [EAMCET 2002]
- (a) -1 (b) 3 (c) 2 (d) 1
- 210.** If α, β be the roots of $x^2 + px + q = 0$ and $\alpha + h, \beta + h$ are the roots of $x^2 + rx + s = 0$, then [AMU 2001]
- (a) $\frac{p}{r} = \frac{q}{s}$ (b) $2h = \left[\frac{p}{q} + \frac{r}{s} \right]$ (c) $p^2 - 4q = r^2 - 4s$ (d) $pr^2 = qs^2$
- 211.** The quadratic equation with real coefficients whose one root is $7 + 5i$ will be [Kerala (Engg.) 2001, 02; Rajasthan PET 1999]
- (a) $x^2 - 14x - 74 = 0$ (b) $x^2 + 14x + 74 = 0$ (c) $x^2 + 14x - 74 = 0$ (d) $x^2 - 14x + 74 = 0$
- 212.** The quadratic equation with one root as the square root of $-47 + 8\sqrt{-3}$ is [IIT 1995]
- (a) $x^2 + 2x + 49 = 0$ (b) $x^2 - 2x + 49 = 0$ (c) $x^2 \pm 2x + 49 = 0$ (d) $x^2 \pm 2x - 49 = 0$
- 213.** The quadratic equation whose one root is $\frac{1}{2 + \sqrt{5}}$ will be [Rajasthan PET 1987]
- (a) $x^2 + 4x - 1 = 0$ (b) $x^2 - 4x - 1 = 0$ (c) $x^2 + 4x + 1 = 0$ (d) None of these
- 214.** The quadratic equation with one root $2 - \sqrt{3}$ is [Rajasthan PET 1985]
- (a) $x^2 - 4x + 1 = 0$ (b) $x^2 - 4x - 1 = 0$ (c) $x^2 + 4x + 1 = 0$ (d) $x^2 + 4x - 1 = 0$
- 215.** The quadratic equation whose roots are three times the roots of the equation $3ax^2 + 3bx + c = 0$ is [AMU 1990]
- (a) $ax^2 + bx + c = 0$ (b) $ax^2 + 3bx + c = 0$ (c) $ax^2 + bx + 3c = 0$ (d) $ax^2 + 3bx + 3c = 0$
- 216.** If α, β are the roots of $x^2 + px + q = 0$ then $-\frac{1}{\alpha}, -\frac{1}{\beta}$ are the roots of the equation [TS Rajendra 1991]

- (a) $qx^2 - px + 1 = 0$ (b) $qx^2 + px + 1 = 0$ (c) $x^2 + px + q = 0$ (d) $x^2 - px + q = 0$
- 217.** If a root of the equation $ax^2 + bx + c = 0$ be reciprocal of a root of the equation $a'x^2 + b'x + c' = 0$, then [IIT 1968]
 (a) $(cc' - aa')^2 = (ba' - cb')(ab' - bc')$ (b) $(bb' - aa')^2 = (ca' - bc')(ab' - bc')$
 (c) $(cc' - aa')^2 = (ba' + cb')(ab' + bc')$ (d) None of these
- 218.** One root of $ax^2 + bx + c = 0$ is reciprocal of other root if [Rajasthan PET 1985]
 (a) $a + c = 0$ (b) $b + c = 0$ (c) $b - c = 0$ (d) $a - c = 0$
- 219.** If the roots of the equation $5x^2 + 13x + k = 0$ be reciprocals of each other, then k is equal to [MNR 1980; Rajasthan PET 1990]
 (a) 0 (b) 5 (c) $1/6$ (d) 6
- 220.** If one root of the equation $x^2 = px + q$ is reciprocal of the other, then the correct relationship is [AMU 1987, 89]
 (a) $q = -1$ (b) $q = 1$ (c) $pq = -1$ (d) $pq = 1$
- 221.** If the roots of the quadratic equation $\frac{x-m}{mx+1} = \frac{x+n}{nx+1}$ are reciprocal to each other, then [MP PET 2001]
 (a) $n = 0$ (b) $m = n$ (c) $m + n = 1$ (d) $m^2 + n^2 = 1$
- 222.** The roots of the quadratic equation $ax^2 + bx + c = 0$ will be reciprocal to each other if
 (a) $a = \frac{1}{c}$ (b) $a = c$ (c) $b = ac$ (d) $a = b$
- 223.** If the absolute difference between two roots of the equation $x^2 + px + 3 = 0$ is \sqrt{p} , then p equals [Bihar CEE 1998]
 (a) -3, 4 (b) 4 (c) -3 (d) None of these
- 224.** If the roots of equation $x^2 - px + q = 0$ differ by 1, then [MP PET 1999]
 (a) $p^2 = 4q$ (b) $p^2 = 4q + 1$ (c) $p^2 = 4q - 1$ (d) None of these
- 225.** The numerical difference of the roots of $x^2 - 7x - 9 = 0$ is
 (a) 5 (b) $2\sqrt{85}$ (c) $9\sqrt{7}$ (d) $\sqrt{85}$
- 226.** If the difference of the roots of $x^2 - px + 8 = 0$ be 2, then the value of p is [Roorkee 1992]
 (a) ± 2 (b) ± 4 (c) ± 6 (d) ± 8



227. If the difference of the roots of the equation $x^2 - bx + c = 0$ be 1, then [Rajasthan PET 1991]
 (a) $b^2 - 4c - 1 = 0$ (b) $b^2 - 4c = 0$ (c) $b^2 - 4c + 1 = 0$ (d) $b^2 + 4c - 1 = 0$
228. If the roots of the equations $x^2 - bx + c = 0$ and $x^2 - cx + b = 0$ differ by the same quantity, then $b + c$ is equal to [BIT Ranchi 1969; MP PET 1993]
 (a) 4 (b) 1 (c) 0 (d) -4
229. If the roots of $x^2 - bx + c = 0$ are two consecutive integers, then $b^2 - 4c$ is [Kurukshetra CEE 1998]
 (a) 1 (b) 2 (c) 3 (d) 4
230. If α, β are the roots of $x^2 - 3x + a = 0, a \in R$ and $\alpha < 1 < \beta$ then
 (a) $a \in (-\infty, 2)$ (b) $a \in \left(-\infty, \frac{9}{4}\right)$ (c) $a \in \left(2, \frac{9}{4}\right)$ (d) None of these
231. If α, β be the roots of $4x^2 - 16x + \lambda = 0, \lambda \in R$ such that $1 < \alpha < 2$ and $2 < \beta < 3$ then the number of integral solutions of λ is
 (a) 5 (b) 6 (c) 2 (d) 3
232. If X denotes the set of real numbers p for which the equation $x^2 = p(x + p)$ has its roots greater than p then X is equal to
 (a) $\left(-2, -\frac{1}{2}\right)$ (b) $\left(-\frac{1}{2}, \frac{1}{4}\right)$ (c) Null set (d) $(-\infty, 0)$
233. If one root of the quadratic equation $ax^2 + bx + c = 0$ is equal to the n^{th} power of the other root, then the value of $(ac^n)^{\frac{1}{n+1}} + (a^n c)^{\frac{1}{n+1}} =$ [IIT 1983]
 (a) b (b) $-b$ (c) $b^{\frac{1}{n+1}}$ (d) $-b^{\frac{1}{n+1}}$
234. If one root of the equation $ax^2 - bx + c = 0$ is square of the other, then [Rajasthan PET 1998]
 (a) $a^2c + ac^2 + 3abc - b^3 = 0$ (b) $a^2c + ac^2 - 3abc + b^3 = 0$ (c) $a^3 + b^3 = 3abc$ (d) $(a+b)^3 = 3abc$
235. For the equation $3x^2 + px + 3, p > 0$ if one of the root is square of the other, then p is equal to [IIT Screening 2000]
 (a) $\frac{1}{3}$ (b) 1 (c) 3 (d) $\frac{2}{3}$
236. If one root of equation $px^2 - qx + r = 0$ is double of the other, then
 (a) $9q^2 = 2pr$ (b) $2q^2 = 9pr$ (c) $3q^2 = 4pr$ (d) $4q^2 = 3pr$
237. The value of k for which one of the roots of $x^2 - x + 3k = 0$ is double of one of the roots of $x^2 - x + k = 0$ is [UPSEAT 2000]
 (a) 1 (b) -2 (c) 2 (d) None of these
238. The function $f(x) = ax^2 + 2x + 1$ has one double root if [AMU 1989]
 (a) $a = 0$ (b) $a = -1$ (c) $a = 1$ (d) $a = 2$
239. If $\sin \alpha, \cos \alpha$ are the roots of the equation $ax^2 + bx + c = 0$, then [MP PET 1993]
 (a) $a^2 - b^2 + 2ac = 0$ (b) $(a - c)^2 = b^2 + c^2$ (c) $a^2 + b^2 - 2ac = 0$ (d) $a^2 + b^2 + 2ac = 0$
240. If the roots of $ax^2 + bx + c = 0$ are α, β and root of $Ax^2 + Bx + c = 0$ are $\alpha - k, \beta - k$, then $\frac{B^2 - 4AC}{b^2 - 4ac}$ is equal to [Rajasthan PET 1999]
 (a) $\frac{a}{A}$ (b) $\frac{A}{a}$ (c) $\left(\frac{a}{A}\right)^2$ (d) $\left(\frac{A}{a}\right)^2$
241. If the product of roots of the equation $x^2 - 3kx + 2e^{2 \log k} - 1 = 0$ is 7, then its roots will real when [Pb. CET 1990; IIT 1984]
 (a) $k = 1$ (b) $k = 2$ (c) $k = 3$ (d) None of these



190 Quadratic Equations and Inequalities

- 242.** If a and b are rational and b is not a perfect square then the quadratic equation with rational coefficients whose one root is $\frac{1}{a + \sqrt{b}}$ is
 (a) $x^2 - 2ax + (a^2 - b) = 0$ (b) $(a^2 - b)x^2 - 2ax + 1 = 0$ (c) $(a^2 - b)x^2 - 2bx + 1 = 0$ (d) None of these
- 243.** If $\frac{1}{4 - 3i}$ is a root of $ax^2 + bx + 1 = 0$, where a, b are real, then
 (a) $a = 25, b = -8$ (b) $a = 25, b = 8$ (c) $a = 5, b = 4$ (d) None of these
- 244.** If α, β, γ be the roots of the equation $x(1 + x^2) + x^2(6 + x) + 2 = 0$ then the value of $\alpha^{-1} + \beta^{-1} + \gamma^{-1}$ is
 (a) -3 (b) $\frac{1}{2}$ (c) $-\frac{1}{2}$ (d) None of these
- 245.** If the roots of $x^3 - 12x^2 + 39x - 28 = 0$ are in A.P. then their common difference is
 (a) ± 1 (b) ± 2 (c) ± 3 (d) ± 4
- 246.** The roots of the equation $x^3 + 14x^2 - 84x - 216 = 0$ are in
 (a) A.P. (b) G.P. (c) H.P. (d) None of these
- 247.** If 3 and $1 + \sqrt{2}$ are two roots of a cubic equation with rational coefficients, then the equation is
 (a) $x^3 - 5x^2 + 9x - 9 = 0$ (b) $x^3 - 3x^2 - 4x + 12 = 0$ (c) $x^3 - 5x^2 + 7x + 3 = 0$ (d) None of these
- 248.** What is the sum of the squares of roots of $x^2 - 3x + 1 = 0$ [Karnataka CET 1993]
 (a) 5 (b) 7 (c) 9 (d) 10
- 249.** If $\alpha + \beta = 3$ and $\alpha^3 + \beta^3 = 27$, then α and β are the roots of
 (a) $3x^2 + 9x + 7 = 0$ (b) $9x^2 - 27x + 20 = 0$ (c) $2x^2 - 6x + 15 = 0$ (d) None of these
- 250.** For what value of λ the sum of the squares of the roots of $x^2 + (2 + \lambda)x - \frac{1}{2}(1 + \lambda) = 0$ is minimum [AMU 1999]
 (a) $3/2$ (b) 1 (c) $1/2$ (d) $11/4$
- 251.** The value of $a (a \geq 3)$ for which the sum of the cubes of the roots of $x^2 - (a - 2)x + (a - 3) = 0$, assumes the least value is [Orissa JEE 2002]
 (a) 3 (b) 4 (c) 5 (d) None of these
- 252.** Let α, β be the roots of $x^2 + (3 - \lambda)x - \lambda = 0$. The value of λ for which $\alpha^2 + \beta^2$ is minimum, is [AMU 2000]
 (a) 0 (b) 1 (c) 2 (d) 3
- 253.** If the sum of squares of the roots of the equation $x^2 - (a - 2)x - (a + 1) = 0$ is least, then the value of a is [Rajasthan PET 2000. Pb. CET 2002]
 (a) 0 (b) 2 (c) -1 (d) 1
- 254.** If α, β are roots of $Ax^2 + Bx + C = 0$ and α^2, β^2 are roots of $x^2 + px + q = 0$, then p is equal to [Rajasthan PET 1986]
 (a) $(B^2 - 2AC)/A^2$ (b) $(2AC - B^2)/A^2$ (c) $(B^2 - 4AC)/A^2$ (d) $(4AC - B^2)/A^2$
- 255.** If α, β are roots of the equation $x^2 + x + 1 = 0$ and $\frac{\alpha}{\beta}, \frac{\beta}{\alpha}$ are roots of the equation $x^2 + px + q = 0$, then p equals [Rajasthan PET 1987, 93]
 (a) -1 (b) 1 (c) -2 (d) 2
- 256.** If α, β are real and α^2, β^2 are the roots of the equation $a^2x^2 + x + 1 - a^2 = 0 (a > 1)$, then $\beta^2 =$ [EAMCET 1999]
 (a) a^2 (b) $1 - \frac{1}{a^2}$ (c) $1 - a^2$ (d) $1 + a^2$
- 257.** The H.M. of the roots of the equation $x^2 - 8x + 4 = 0$ is [Rajasthan PET 1988]
 (a) 1 (b) 2 (c) 3 (d) None of these
- 258.** If α, β are the roots of the equation $x^2 + x\sqrt{\alpha} + \beta = 0$, then the value of α and β are [AMU 1990, 92]
 (a) $\alpha = 1$ and $\beta = -1$ (b) $\alpha = 1$ and $\beta = -2$ (c) $\alpha = 2$ and $\beta = 1$ (d) $\alpha = 2$ and $\beta = -2$
- 259.** If p and q are the roots of $x^2 + px + q = 0$, then [IIT 1995, AIEEE 2002]
 (a) $p = 1$ (b) $p = -2$ (c) $p = 1$ or 0 (d) $p = -2$ or 0

- 260.** If roots of the equation $2x^2 - (a^2 + 8a + 1)x + a^2 - 4a = 0$ are in opposite sign, then [AMU 1998]
 (a) $0 < a < 4$ (b) $a > 0$ (c) $a < 8$ (d) $-4 < a < 0$
- 261.** Which of the following equation has 1 and -2 as the roots [SCRA 1999]
 (a) $x^2 - x - 2 = 0$ (b) $x^2 + x - 2 = 0$ (c) $x^2 - x + 2 = 0$ (d) $x^2 + x + 2 = 0$
- 262.** If the roots of the equation $x^2 + x + 1 = 0$ are in the ratio $m : n$ then [Rajasthan PET 1994]
 (a) $\sqrt{\frac{m}{n}} + \sqrt{\frac{n}{m}} + 1 = 0$ (b) $\sqrt{m} + \sqrt{n} + 1 = 0$ (c) $\frac{m}{n} + \frac{n}{m} + 1 = 0$ (d) $m + n + 1 = 0$
- 263.** If the roots of the equation $lx^2 + nx + n = 0$ are in the ratio $p : q$ then $\sqrt{\frac{p}{q}} + \sqrt{\frac{q}{p}}$ is equal to [Rajasthan PET 1997; BIT Ranchi 2002]
 (a) $\sqrt{n/l}$ (b) $\sqrt{l/n}$ (c) $\pm \sqrt{n/l}$ (d) $-\sqrt{l/n}$
- 264.** If the roots of the equation $12x^2 - mx + 5 = 0$ are in the ratio $2 : 3$, then $m =$ [Rajasthan PET 2002]
 (a) $5\sqrt{10}$ (b) $3\sqrt{10}$ (c) $2\sqrt{10}$ (d) None of these
- 265.** If the ratio of the roots of the equation $ax^2 + bx + c = 0$ be $p : q$, then [Pb. CET 1994]
 (a) $pqb^2 + (P+q)^2ac = 0$ (b) $pqb^2 - (P+q)^2ac = 0$ (c) $pqa^2 - (P+q)^2bc = 0$ (d) None of these
- 266.** The two roots of an equation $x^3 - 9x^2 + 14x + 24 = 0$ are in the ratio $3 : 2$. The roots will be [UPSEAT 1999]
 (a) 6, 4, -1 (b) 6, 4, 1 (c) -6, 4, 1 (d) -6, -4, 1
- 267.** The condition that one root of the equation $ax^2 + bx + c = 0$ is three times the other is [DCE 2002]
 (a) $b^2 = 8ac$ (b) $3b^2 + 16ac = 0$ (c) $3b^2 = 16ac$ (d) $b^2 + 3ac = 0$
- 268.** If the roots of the equation $\frac{x^2 - bx}{ax - c} = \frac{\lambda - 1}{\lambda + 1}$ are such that $\alpha + \beta = 0$, then the value of λ is [Kurukhstra CEE 1995; MP PET 1996, 2002; Rajasthan PET 2001]
 (a) $\frac{a-b}{a+b}$ (b) c (c) $\frac{1}{c}$ (d) $\frac{a+b}{a-b}$
- 269.** For the equation $\frac{1}{x+a} - \frac{1}{x+b} = \frac{1}{x+c}$, if the product of the roots is zero, then the sum of the roots is [AMU 1992]
 (a) 0 (b) $\frac{2ab}{b+c}$ (c) $\frac{2bc}{b+c}$ (d) $-\frac{2bc}{b+c}$
- 270.** If the sum of two of the roots of $x^3 + px^2 + qx + r = 0$ is zero, then $pr =$ [EAMCET 2003]
 (a) $-r$ (b) r (c) $2r$ (d) $-2r$
- 271.** If the roots of the equation $\frac{1}{x+p} + \frac{1}{x+q} = \frac{1}{r}$ are equal in magnitude but opposite in sign, then the product of the roots will be [IIT 1967]
 (a) $\frac{p^2 + q^2}{2}$ (b) $-\frac{(p^2 + q^2)}{2}$ (c) $\frac{p^2 - q^2}{2}$ (d) $-\frac{(p^2 - q^2)}{2}$
- 272.** The value of m for which the equation $x^3 - mx^2 + 3x - 2 = 0$ has two roots equal in magnitude but opposite in sign, is [Kurukhstra CEE 1996]
 (a) $1/2$ (b) $2/3$ (c) $3/4$ (d) $4/5$
- 273.** If $ax^2 + bx + c = a(x - \alpha)(x - \beta)$, then $a(\alpha x + 1)(\beta x + 1)$ is equal to [AMU 1986]
 (a) $ax^2 + bx + c$ (b) $cx^2 - bx + a$ (c) $cx^2 - bx - a$ (d) $cx^2 + bx + a$
- 274.** If α, β are the roots of the equation $ax^2 + bx + c = 0$ ($a \neq 0$) and $\alpha + \delta, \beta + \delta$ are the roots of $Ax^2 + Bx + C = 0$ ($A \neq 0$) for some constant, then [IIT 2000]
 (a) $\frac{b^2 - 4ac}{a^2} = \frac{B^2 - 4AC}{A^2}$ (b) $\frac{b^2 - 2ac}{a^2} = \frac{B^2 - 2AC}{A^2}$ (c) $\frac{b^2 - 8ac}{a^2} = \frac{B^2 - 8AC}{A^2}$ (d) None of these
- 275.** In a triangle PQR , $\angle R = \frac{\pi}{2}$. If $\tan\left(\frac{p}{2}\right)$ and $\tan\left(\frac{q}{2}\right)$ are the roots of the equation $ax^2 + bx + c = 0$ ($a \neq 0$), then [IIT 1999]
 (a) $a + b = c$ (b) $b + c = 0$ (c) $a + c = b$ (d) $b = c$

192 Quadratic Equations and Inequalities

- 276.** The product of all real roots of the equation $x^2 - |x| - 6 = 0$ is [Roorkee 2000]
 (a) -9 (b) 6 (c) 9 (d) 36
- 277.** If the sum of the roots of the equation $ax^2 + bx + c = 0$ is equal to the sum of the squares of their reciprocals then bc^2, ca^2, ab^2 are in [IIT 1976]
 (a) A.P. (b) G.P. (c) H.P. (d) None of these
- 278.** The roots of the equation $x^2 - 2x + A = 0$ are p, q and the roots of the equation $x^2 - 18x + B = 0$ are r, s . If $p < q < r < s$ are in A.P., then [IIT 1997]
 (a) $A = 3, B = 77$ (b) $A = -3, B = 77$ (c) $A = 3, B = -77$ (d) $A = -3, B = -77$
- 279.** If the roots of the equation $x^2 + bx + c = 0$ and $x^2 + qx + r = 0$ are in the same ratio, then [EAMCET 1994]
 (a) $r^2c = qb^2$ (b) $r^2b = qc^2$ (c) $c^2r = q^2b$ (d) $b^2r = q^2c$
- 280.** If one root of the equation $x^2 + px + q = 0$ is $2 + \sqrt{3}$, then values of p and q are [UPSEAT 2002]
 (a) -4, 1 (b) 4, -1 (c) $2, \sqrt{3}$ (d) $-2, -\sqrt{3}$
- 281.** If $1 - i$ is a root of the equation $x^2 - ax + b = 0$, then $b =$ [EAMCET 2002]
 (a) -2 (b) -1 (c) 1 (d) 2

Advance Level

- 282.** If α, β are the roots of $x^2 + px + 1 = 0$ and γ, δ are the roots of $x^2 + qx + 1 = 0$, then $q^2 - p^2 =$ [IIT 1978; DCE 2000]
 (a) $(\alpha - \gamma)(\beta - \gamma)(\alpha + \delta)(\beta + \delta)$ (b) $(\alpha + \gamma)(\beta + \gamma)(\alpha - \delta)(\beta + \delta)$
 (c) $(\alpha + \gamma)(\beta + \gamma)(\alpha + \delta)(\beta + \delta)$ (d) None of these
- 283.** If α, β be the roots of $x^2 - px + q = 0$ and α', β' be the roots of $x^2 - p'x + q' = 0$, then the value of $(\alpha - \alpha')^2 + (\beta - \alpha')^2 + (\alpha - \beta')^2 + (\beta - \beta')^2$ is
 (a) $2\{p^2 - 2q + p'^2 - 2q' - pp'\}$ (b) $2\{p^2 - 2q + p'^2 - 2q' - qq'\}$
 (c) $2\{p^2 - 2q - p'^2 - 2q' - pp'\}$ (d) $2\{p^2 - 2q - p'^2 - 2q' - qq'\}$
- 284.** If α and β are the roots of the equation $x^2 - ax + b = 0$ and $A_n = \alpha^n + \beta^n$, then which of the following is true [Karnataka CET 2000]
 (a) $A_{n+1} = aA_n + bA_{n-1}$ (b) $A_{n+1} = bA_n + aA_{n-1}$ (c) $A_{n+1} = aA_n - bA_{n-1}$ (d) $A_{n+1} = bA_n - aA_{n-1}$
- 285.** If roots of an equation $x^n - 1 = 0$ are $1, a_1, a_2, \dots, a_{n-1}$, then the value of $(1 - a_1)(1 - a_2)(1 - a_3) \dots (1 - a_{n-1})$ will be [UPSEAT 1999]
 (a) n (b) n^2 (c) n^n (d) 0
- 286.** If α and β are the roots of $6x^2 - 6x + 1 = 0$, then the value of $\frac{1}{2}[a + b\alpha + c\alpha^2 + d\alpha^3] + \frac{1}{2}[a + b\beta + c\beta^2 + d\beta^3]$ is [Rajasthan PET 2000]
 (a) $\frac{1}{4}(a + b + c + d)$ (b) $\frac{a}{1} + \frac{b}{2} + \frac{c}{3} + \frac{d}{4}$ (c) $\frac{a}{1} - \frac{b}{2} + \frac{c}{3} - \frac{d}{4}$ (d) None of these
- 287.** If α_1, α_2 are the roots of equation $x^2 - px + 1 = 0$ and β_1, β_2 be those of equation $x^2 - qx + 1 = 0$ and vector $\alpha_1\hat{i} + \beta_1\hat{j}$ is parallel to $\alpha_2\hat{i} + \beta_2\hat{j}$, then
 (a) $p = \pm q$ (b) $p = \pm 2q$ (c) $p = 2q$ (d) None of these
- 288.** If the roots of $a_1x^2 + b_1x + c_1 = 0$ are α_1 and β_1 and those of $a_2x^2 + b_2x + c_2 = 0$ are α_2 and β_2 such that $\alpha_1\alpha_2 = \beta_1\beta_2 = 1$, then
 (a) $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$ (b) $\frac{a_1}{c_2} = \frac{b_1}{b_2} = \frac{c_1}{a_2}$ (c) $a_1a_2 = b_1b_2 = c_1c_2$ (d) None of these
- 289.** If the sum of the roots of the equation $qx^2 + 2x + 3q = 0$ is equal to their product, then the value of q is equal to

- (a) $-\frac{2}{3}$ (b) $\frac{3}{2}$ (c) 3 (d) -6
290. If $x = (\beta - \gamma)(\alpha - \delta)$, $y = (\gamma - \alpha)(\beta - \delta)$, $z = (\alpha - \beta)(\gamma - \delta)$, then the value of $x^3 + y^3 + z^3 - 3xyz$ is
 (a) 0 (b) $\alpha^6 + \beta^6 + \gamma^6 + \delta^6$ (c) $\alpha^6 \beta^6 \gamma^6 \delta^6$ (d) None of these
291. If α, β, γ are the roots of the equation $x^3 + px^2 + qx + r = 0$, then $(1 - \alpha^2)(1 - \beta^2)(1 - \gamma^2)$ is equal to
 (a) $(1 + q)^2 - (p + r)^2$ (b) $(1 + q)^2 + (p + r)^2$ (c) $(1 - q)^2 + (p - r)^2$ (d) None of these
292. If α, β, γ are the roots of the equation $x^3 + ax + b = 0$, then $\frac{\alpha^3 + \beta^3 + \gamma^3}{\alpha^2 + \beta^2 + \gamma^2} =$
 (a) $\frac{3b}{2a}$ (b) $\frac{-3b}{2a}$ (c) $3b$ (d) $2a$
293. If α, β are the roots of $6x^2 - 2x + 1 = 0$ and $s_x = \alpha^n + \beta^n$, then $\lim_{n \rightarrow \infty} \sum_{r=1}^n S_r$ is
 (a) $\frac{5}{17}$ (b) 0 (c) $\frac{3}{37}$ (d) None of these
294. Let α, β be the roots of the equation $ax^2 + bx + c = 0$ and let $\alpha^n + \beta^n = S_n$ for $n \geq 1$. Then the value of the determinant

$$\begin{vmatrix} 3 & 1 + S_1 & 1 + S_2 \\ 1 + S_1 & 1 + S_2 & 1 + S_3 \\ 1 + S_2 & 1 + S_3 & 1 + S_4 \end{vmatrix}$$
 is
 (a) $\frac{b^2 - 4ac}{a^4}$ (b) $\frac{(a + b + c)(b^2 + 4ac)}{a^4}$ (c) $\frac{(a + b + c)(b^2 - 4ac)}{a^4}$ (d) $\frac{(a + b + c)^2(b^2 - 4ac)}{a^4}$
295. If α, β are roots of the equation $2x^2 + 6x + b = 0$ ($b < 0$), then $\frac{\alpha}{\beta} + \frac{\beta}{\alpha}$ is less than
 (a) 2 (b) -2 (c) 18 (d) None of these
296. If α, β are roots of the equation $ax^2 + 3x + 2 = 0$ ($a < 0$), then $\frac{\alpha^2}{\beta} + \frac{\beta^2}{\alpha}$ is greater than
 (a) 0 (b) 1 (c) 2 (d) None of these
297. If $\alpha, \beta, \gamma, \sigma$ are the roots of the equation $x^4 + 4x^3 - 6x^2 + 7x - 9 = 0$, then the value of $(1 + \alpha^2)(1 + \beta^2)(1 + \gamma^2)(1 + \sigma^2)$ is
 (a) 5 (b) 9 (c) 11 (d) 13
298. If α and β are the roots of the equation $x^2 - p(x + 1) - q = 0$, then the value of $\frac{\alpha^2 + 2\alpha + 1}{\alpha^2 + 2\alpha + q} + \frac{\beta^2 + 2\beta + 1}{\beta^2 + 2\beta + q}$ is
 (a) 2 (b) 3 (c) 0 (d) 1
299. If A, G, H be respectively, the A.M., G.M. and H.M. of three positive number a, b, c then the equation whose roots are these number is given by
 (a) $x^3 - 3Ax^2 + G^3(3x - 1) = 0$ (b) $x^3 - 3Ax^2 + 3(G^3 / H)x - G^3 = 0$
 (c) $x^3 + 3Ax^2 + 3(G^3 / H)x - G^3 = 0$ (d) $x^3 - 3Ax^2 - 3(G^3 / H)x + G^3 = 0$
300. Let $a = \cos \frac{2\pi}{7} + i \sin \frac{2\pi}{7}$, $A = a + a^2 + a^4$ and $B = a^3 + a^5 + a^6$ then A and B are roots of the equation [Rajasthan PET 2000]
 (a) $x^2 - x + 2 = 0$ (b) $x^2 - x - 2 = 0$ (c) $x^2 + x + 2 = 0$ (d) None of these
301. If α, β are the roots of the equation $x^2 - px + q = 0$, then the quadratic equation whose roots are $(\alpha^2 - \beta^2)(\alpha^3 - \beta^3)$ and $\alpha^3 \beta^2 + \alpha^2 \beta^3$ is [Roorkee 1994]
 (a) $x^2 - Sx + P = 0$ (b) $x^2 + Sx + P = 0$ (c) $x^2 + Sx - P = 0$ (d) None of these
 [Where $S = p(p^4 - 5p^2q + 5q^2)$ and $P = p^2q^2(p^4 - 5p^2q + 4q^2)$]
302. Let A, G and H are the A.M., G.M. and H.M. respectively of two unequal positive integers. Then the equation $Ax^2 - |G|x - H = 0$ has
 (a) Both roots as fractions (b) At least one root which is a negative fraction
 (c) Exactly one positive root (d) At least one root which is an integer



194 Quadratic Equations and Inequalities

- 303.** Let $x^2 - px + q = 0$, where $p \in R, q \in R$, have the roots α, β such that $\alpha + 2\beta = 0$ then
 (a) $2p^2 + q = 0$ (b) $2q^2 + p = 0$ (c) $q < 0$ (d) None of these
- 304.** The cubic equation whose roots are the A.M., G.M. and H.M. of the roots of $x^2 - 2px + q^2 = 0$ is
 (a) $(x - p)(x - q)(x - p - q) = 0$ (b) $(x - p)(x - |q|)(px - q^2) = 0$
 (c) $x^3 - \left(p + |q| + \frac{q^2}{p}\right)x^2 + \left(p|q| + q^2 + \frac{|q|^3}{p}\right)x - |q|^3 = 0$ (d) None of these
- 305.** If α, β are the roots of $x^2 + px + q = 0$ and also of $x^{2n} + p^n x^n + q^n = 0$ and if $\frac{\alpha}{\beta}, \frac{\beta}{\alpha}$ are the roots of $x^n + 1 + (x + 1)^n = 0$, then n is
 (a) An odd integer (b) An even integer (c) Any integer (d) None of these
- 306.** If $\cos^4 x + \sin^2 x - p = 0, p \in R$ has real solutions then
 (a) $p \leq 1$ (b) $\frac{3}{4} \leq p \leq 1$ (c) $p \geq \frac{3}{4}$ (d) None of these
- 307.** If the ratio of the roots of $\lambda x^2 + \mu x + \nu = 0$ is equal to the ratio of the roots of $x^2 + x + 1 = 0$ then λ, μ, ν are in
 (a) A.P. (b) G.P. (c) H.P. (d) None of these
- 308.** P, q, r and s are integers. If the A.M. of the roots of $x^2 - px + q^2 = 0$ and G.M. of the roots of $x^2 - rx + s^2 = 0$ are equal then
 (a) q is an odd integer (b) r is an even integer (c) p is an even integer (d) s is an odd integer
- 309.** If the roots of $4x^2 + 5k = (5k + 1)x$ differ by unity then the negative value of k is
 (a) -3 (b) $-\frac{1}{5}$ (c) $-\frac{3}{5}$ (d) None of these
- 310.** The harmonic mean of the roots of the equation $(5 + \sqrt{2})x^2 - (4 + \sqrt{5})x + 8 + 2\sqrt{5} = 0$ is
 (a) 2 (b) 4 (c) 6 (d) 8
- 311.** If α, β are the roots of $ax^2 + c = bx$ then the equation $(a + cy)^2 = b^2 y$ in y has the roots
 (a) α^{-1}, β^{-1} (b) α^2, β^2 (c) $\alpha\beta^{-1}, \alpha^{-1}\beta$ (d) α^{-2}, β^{-2}
- 312.** If the roots of $ax^2 - bx - c = 0$ change by the same quantity then the expression in a, b, c that does not change is
 (a) $\frac{b^2 - 4ac}{a^2}$ (b) $\frac{b - 4c}{a}$ (c) $\frac{b^2 + 4ac}{a^2}$ (d) None of these
- 313.** If α, β are the roots of $x^2 - px + q = 0$ then the product of the roots of the quadratic equation whose roots are $\alpha^2 - \beta^2$ and $\alpha^3 - \beta^3$ is
 (a) $p(p^2 - q)^2$ (b) $p(p^2 - q)(p^2 - 4q)$ (c) $p(p^2 - 4q)(p^2 + q)$ (d) None of these
- 314.** The quadratic equation whose roots are the A.M. and H.M. of the roots of the equation $x^2 + 7x - 1 = 0$ is
 (a) $14x^2 + 14x - 45 = 0$ (b) $45x^2 - 14x + 14 = 0$ (c) $14x^2 + 45x - 14 = 0$ (d) None of these
- 315.** If $z_0 = \alpha + i\beta, i = \sqrt{-1}$, then the roots of the cubic equation $x^3 - 2(1 + \alpha)x^2 + (4\alpha + \alpha^2 + \beta^2)x + 2(\alpha^2 + \beta^2) = 0$ are
 (a) $2, z_0, \bar{z}_0$ (b) $1, z_0, -z_0$ (c) $2, z_0, -\bar{z}_0$ (d) $2, -z_0, \bar{z}_0$
- 316.** Let a, b, c be real numbers and $a \neq 0$. If α is a root of $a^2x^2 + bx + c = 0$, β is a root of $a^2x^2 - bx - c = 0$, and $0 < \alpha < \beta$ then the equation $a^2x^2 + 2bx + 2c = 0$ has a root γ that always satisfies
 (a) $\gamma = \frac{1}{2}(\alpha + \beta)$ (b) $\gamma = \alpha + \frac{\beta}{2}$ (c) $\gamma = \alpha$ (d) $\alpha < \gamma < \beta$
- 317.** If $(\lambda^2 + \lambda - 2)x^2 + (\lambda + 2)x < 1$ for all $x \in R$, then λ belongs to the interval
 (a) $(-2, 1)$ (b) $\left(-2, \frac{2}{5}\right)$ (c) $\left(\frac{2}{5}, 1\right)$ (d) None of these
- 318.** The least integral value of k for which $(k - 2)x^2 + 8x + k + 4 > 0$ for all $x \in R$, is
 (a) 5 (b) 4 (c) 3 (d) None of these

- 319.** The set of possible values of λ for which $x^2 - (\lambda^2 - 5\lambda + 5)x + (2\lambda^2 - 3\lambda - 4) = 0$ has roots whose sum and product are both less than 1 is
 (a) $\left(-1, \frac{5}{2}\right)$ (b) (1, 4) (c) $\left[1, \frac{5}{2}\right]$ (d) $\left(1, \frac{5}{2}\right)$
- 320.** The set of the possible values of x such that $5^x + (2\sqrt{3})^{2x} - 169$ is always positive is
 (a) $[3, +\infty)$ (b) $[2, +\infty)$ (c) $(2, +\infty)$ (d) None of these
- 321.** If all real value of x obtained from the equation $4^x - (a-3)2^x + a - 4 = 0$ are nonpositive then
 (a) $a \in (4, 5]$ (b) $a \in (0, 4)$ (c) $a \in (4, +\infty)$ (d) None of these
- 322.** If $ax^2 + bx + 6 = 0$ does not have two distinct real roots $a \in R, b \in R$, then the least value of $3a + b$ is
 (a) 4 (b) -1 (c) 1 (d) -2
- 323.** If $ab = 2a + 3b, a > 0, b > 0$ then the minimum value of ab is
 (a) 12 (b) 24 (c) $\frac{1}{4}$ (d) None of these
- 324.** The number of values of k for which $\{x^2 - (k-2)x + k^2\}\{x^2 + kx + (2k-1)\}$ is a perfect square is
 (a) 1 (b) 2 (c) 0 (d) None of these
- 325.** If $x^2 - bx + c = 0$ has equal integral roots then
 (a) b and c are integers
 (b) b and c are even integers
 (c) b is an even integer and c is a perfect square of a positive integer
 (d) None of these
- 326.** Let A, G and H be the A.M., G.M. and H.M. of two positive number a and b . The quadratic equation whose roots are A and H is
 (a) $Ax^2 - (A^2 + G^2)x + AG^2 = 0$ (b) $Ax^2 - (A^2 + H^2)x + AH^2 = 0$
 (c) $Hx^2 - (H^2 + G^2)x + HG^2 = 0$ (d) None of these
- 327.** If $x^2 + y^2 + z^2 = 1$, then the value of $xy + yz + zx$ lies in the interval
 (a) $\left[\frac{1}{2}, 2\right]$ (b) $[-1, 2]$ (c) $\left[-\frac{1}{2}, 1\right]$ (d) $\left[-1, \frac{1}{2}\right]$
- 328.** If $px^2 + qx + r = 0$ has no real roots and p, q, r are real such that $p + r > 0$, then
 (a) $p - q + r < 0$ (b) $p - q + r > 0$ (c) $p + r = q$ (d) All of these
- 329.** The quadratic equation $x^2 - 2x - \lambda = 0, \lambda \neq 0$
 (a) Cannot have a real root if $\lambda < -1$
 (b) Can have a rational root if λ is a perfect square
 (c) Cannot have an integral root if $n^2 - 1 < \lambda < n^2 + 2n$ where $n = 0, 1, 2, 3, \dots$
 (d) None of these
- 330.** A quadratic equation whose roots are $\left(\frac{\gamma}{\alpha}\right)^2$ and $\left(\frac{\beta}{\alpha}\right)^2$, where α, β, γ are the roots of $x^3 + 27 = 0$, is
 (a) $x^2 - x + 1 = 0$ (b) $x^2 + 3x + 9 = 0$ (c) $x^2 + x + 1 = 0$ (d) $x^2 - 3x + 9 = 0$
- 331.** If a, b are the real roots of $x^2 + px + 1 = 0$ and c, d are the real roots of $x^2 + qx + 1 = 0$, then $(a-c)(b-c)(a+d)(b+d)$ is divisible by
 (a) $a+b+c+d$ (b) $a+b-c-d$ (c) $a-b+c-d$ (d) $a-b-c-d$
- 332.** If $0 < a < 5, 0 < b < 5$ and $\frac{x^2+5}{2} = x - 2\cos(a+bx)$ is satisfied for at least one real x then the greatest value of $a+b$ is
 (a) π (b) $\frac{\pi}{2}$ (c) 3π (d) 4π
- 333.** $a(x^2 - y^2) + \lambda\{x(y+1)+1\}$ can be resolved into linear rational factors. Then

196 Quadratic Equations and Inequalities

- (a) $\lambda = 1$ (b) $\lambda = \frac{4a^2}{a-1}, a \neq 1$ (c) $\lambda = 0, a = 1$ (d) None of these
- 334.** If α, β are the roots of the equation $x^2 + x + 3 = 0$ then equation $3x^2 + 5x + 3 = 0$ has a root
 (a) $\frac{\alpha}{\beta}$ (b) $\frac{\beta}{\alpha}$ (c) $\frac{\alpha}{\beta} + \frac{\beta}{\alpha}$ (d) None of these
- 335.** If α, β are the roots of $x^2 - 2ax + b^2 = 0$ and γ, δ are the roots of $x^2 - 2bx + a^2 = 0$, then
 (a) A.M. of $\alpha, \beta =$ G.M. of γ, δ (b) G.M. of $\alpha, \beta =$ A.M. of γ, δ
 (c) $\alpha, \beta, \gamma, \delta$ are in A.P. (d) $\alpha, \beta, \gamma, \delta$ are in G.P.
- 336.** If the roots of the equation $ax^2 - 4x + a^2 = 0$ are imaginary and the sum of the roots is equal to their product then a is
 (a) -2 (b) 4 (c) 2 (d) None of these

Condition for common roots

Basic Level

- 337.** If equations $x^2 + bx + a = 0$ and $x^2 + ax + b = 0$ have one root common and $a \neq b$, then
 (a) $a + b = 1$ (b) $a - b = 1$ (c) $a + b = -1$ (d) $a + b = 0$ [Rajasthan PET 1992; IIT 1986]
- 338.** If equations $x^2 + 2x + 3\lambda = 0$ and $2x^2 + 3x + 5\lambda = 0$ have one non-zero root common, then λ is equal to [Rajasthan PET 1992]
 (a) 2 (b) -1 (c) 1 (d) 3
- 339.** If $x^2 + ax + 10 = 0$ and $x^2 + bx - 10 = 0$ have a common root, then $a^2 - b^2$ is equal to [Kerala (Engg.) 2002]
 (a) 10 (b) 20 (c) 30 (d) 40
- 340.** If two equations $a_1x^2 + b_1x + c_1 = 0$ and $a_2x^2 + b_2x + c_2 = 0$ have a common root, then the value of $(a_1b_2 - a_2b_1)(b_1c_2 - c_1b_2)$ is [Roorkee 1992]
 (a) $-(a_1c_2 - a_2c_1)^2$ (b) $(a_1a_2 - c_1c_2)^2$ (c) $(a_1c_1 - a_2c_2)^2$ (d) $(a_1c_2 - c_1a_2)^2$
- 341.** If the roots of $a_1x^2 + b_1x + c_1 = 0$ and $a_2x^2 + b_2x + c_2 = 0$ are the same, then [Kurukshetra CEE 1995]
 (a) $a_1 = a_2, b_1 = b_2, c_1 = c_2$ (b) $c_1 = c_2 = 0$
 (c) $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$ (d) $a_1 = b_1 = c_1; a_2 = b_2 = c_2$
- 342.** If one root of the equation $(k^2 + 1)x^2 + 13x + 4k = 0$ is reciprocal of the other then k has the value
 (a) $-2 + \sqrt{3}$ (b) $2 - \sqrt{3}$ (c) 1 (d) None of these
- 343.** If the product of the roots of the equation $x^2 - 5x + 4^{\log_2 \lambda} = 0$ is 8 then λ is
 (a) $\pm 2\sqrt{2}$ (b) $2\sqrt{2}$ (c) 3 (d) None of these
- 344.** If the absolute value of the difference of roots of the equation $x^2 + px + 1 = 0$ exceeds $\sqrt{3p}$ then
 (a) $p < -1$ or $p > 4$ (b) $p > 4$ (c) $-1 < p < 4$ (d) $0 \leq p < 4$
- 345.** If α, β are roots of $x^2 + px + q = 0$ and γ, δ are the roots of $x^2 + px - r = 0$, then $(\alpha - \gamma)(\alpha - \delta)$ is equal to
 (a) $q + r$ (b) $q - r$ (c) $-(q + r)$ (d) $-(p + q + r)$
- 346.** If the equation $2x^2 + 3x + 5\lambda = 0$ and $x^2 + 2x + 3\lambda = 0$ have a common root, then $\lambda =$ [Rajasthan PET 1989]
 (a) 0 (b) -1 (c) 0, -1 (d) 2, -1
- 347.** If a root of the equations $x^2 + px + q = 0$ and $x^2 + ax + \beta = 0$ is common, then its value will be (where $p \neq \alpha$ and $q \neq \beta$) [IIT 1974, 76; Rajasthan PET 1997]
 (a) $\frac{q - \beta}{\alpha - p}$ (b) $\frac{p\beta - \alpha q}{q - \beta}$ (c) $\frac{q - \beta}{\alpha - p}$ or $\frac{p\beta - \alpha q}{q - \beta}$ (d) None of these



348. If $ax^2 + bx + c = 0$ and $bx^2 + cx + a = 0$ have a common root and $a \neq 0$, then $\frac{a^3 + b^3 + c^3}{abc} =$ [IIT 1982; MNR 1983]
 (a) 1 (b) 2 (c) 3 (d) None of these
349. If the equation $x^2 + px + q = 0$ and $x^2 + qx + p = 0$, have a common root, then $p + q + 1 =$ [Orissa JEE 2002]
 (a) 0 (b) 1 (c) 2 (d) -1

Advance Level

350. If every pair from among the equation $x^2 + px + qr = 0$, $x^2 + qx + rp = 0$ and $x^2 + rx + pq = 0$ has a common root, then the product of three common roots is
 (a) pqr (b) $2pqr$ (c) $p^2q^2r^2$ (d) None of these
351. If the equation $x^2 + px + qr = 0$ and $x^2 + qx + pr = 0$ have a common root, then the sum and product of their other roots are respectively
 (a) r, pq (b) $-r, pq$ (c) pq, r (d) $-pq, r$
352. The value of 'a' for which the equations $x^3 + ax + 1 = 0$ and $x^4 + ax^2 + 1 = 0$ have a common root is
 (a) 2 (b) -2 (c) 0 (d) None of these
353. If the equations $ax^2 + bx + c = 0$ and $cx^2 + bx + a = 0$, $a \neq c$ have a negative common root then the value of $a - b + c$ is
 (a) 0 (b) 2 (c) 1 (d) None of these
354. If $x^2 + ax + b = 0$ and $x^2 + bx + a = 0$, $a \neq b$, have a common root α then
 (a) $a + b = 1$ (b) $\alpha + 1 = 0$ (c) $\alpha = 1$ (d) $a + b + 1 = 0$
355. If α is a root of the equation $2x(2x + 1) = 1$ then the other root is
 (a) $3\alpha^3 - 4\alpha$ (b) $-2\alpha(\alpha + 1)$ (c) $4\alpha^3 - 3\alpha$ (d) None of these
356. The common roots of the equations $x^3 + 2x^2 + 2x + 1 = 0$ and $1 + x^{130} + x^{1988} = 0$ are (where ω is a nonreal cube root of unity)
 (a) ω (b) ω^2 (c) -1 (d) $\omega - \omega^2$
357. If a, b, c are rational and no two of them are equal then the equations $(b - c)x^2 + (c - a)x + a - b = 0$ and $a(b - c)x^2 + b(c - a)x + c(a - b) = 0$
 (a) Have rational roots (b) Will be such at least one has rational roots
 (c) Have exactly one root common (d) Have at least one root common
358. If the equations $ax^2 + bx + c = 0$ and $x^3 + 3x^2 + 3x + 2 = 0$ have two common roots, then
 (a) $a = b \neq c$ (b) $a = -b = c$ (c) $a = b = c$ (d) None of these
359. The equations $ax^2 + bx + a = 0$ and $x^3 - 2x^2 + 2x - 1 = 0$ have 2 roots in common. Then $a + b$ must be equal to
 (a) 1 (b) -1 (c) 0 (d) None of these
360. If a, b, c are in G.P. then the equations $ax^2 + 2bx + c = 0$ and $dx^2 + 2ex + f = 0$ have a common root if $\frac{d}{a}, \frac{e}{b}, \frac{f}{c}$ are in
 [IIT 1985; Pb. CET 2000; DCE 2000]
 (a) A.P. (b) G.P. (c) H.P. (d) None of these
361. If the equations $x^2 + ix + a = 0$, $x^2 - 2x + ia = 0$, $a \neq 0$ have a common root then
 (a) a is real (b) $a = \frac{1}{2} + i$
 (c) $a = \frac{1}{2} - i$ (d) The other root is also common
362. If $x^2 - 2r.p_r x + r = 0$; $r = 1, 2, 3$ are three quadratic equations of which each pair has exactly one root common then the number of solutions of the triplet (p_1, p_2, p_3) is
 (a) 2 (b) 1 (c) 9 (d) 27
363. If x, y, z are three consecutive terms of a G.P., where $x > 0$ and the common ratio is r , then the inequality $z + 3x > 4y$ holds for


198 Quadratic Equations and Inequalities

- (a) $r \in (-\infty, 1)$ (b) $r = \frac{24}{5}$ (c) $r \in (3, +\infty)$ (d) $r = \frac{1}{2}$
364. If x is real, then the value of $x^2 - 6x + 13$ will not be less than [Rajasthan PET 1986]
 (a) 4 (b) 6 (c) 7 (d) 8
365. If x be real, the least value of $x^2 - 6x + 10$ is [Kurukshetra CEE 1998]
 (a) 1 (b) 2 (c) 3 (d) 10
366. The smallest value of $x^2 - 3x + 3$ in the interval $(-3, 3/2)$ is [EAMCET 1991]
 (a) $3/4$ (b) 5 (c) -15 (d) -20
367. If $x = 2 + 2^{1/3} + 2^{2/3}$, then $x^3 - 6x^2 + 6x$ equals [Rajasthan PET 1995; MNR 1985]
 (a) 2 (b) -2 (c) 0 (d) 1
368. If x be real, then the minimum value of $x^2 - 8x + 17$ is [MNR 1980]
 (a) -1 (b) 0 (c) 1 (d) 2
369. If x be real, then the maximum value of $5 + 4x - 4x^2$ will be equal to [MNR 1979]
 (a) 5 (b) 6 (c) 1 (d) 2
370. The expression $ax^2 + bx + c$ has the same sign as of 'a' of [Kurukshetra CEE 1995]
 (a) $b^2 - 4ac > 0$ (b) $b^2 - 4ac = 0$
 (c) $b^2 - 4ac \leq 0$ (d) b and c have the same sign as a .
371. The value of $x^2 + 2bx + c$ is positive if [Roorkee 1995]
 (a) $b^2 - 4c > 0$ (b) $b^2 - 4c < 0$ (c) $c^2 < b$ (d) $b^2 < c$
372. The values of 'a' for which $(a^2 - 1)x^2 + 2(a - 1)x + 2$ is positive for any x are [UPSEAT 2001]
 (a) $a \geq 1$ (b) $a \leq 1$ (c) $a > -3$ (d) $a < -3$ or $a > 1$

Quadratic Expressions

Basic Level

373. If x is real, then the maximum and minimum values of the expression $\frac{x^2 - 3x + 4}{x^2 + 3x + 4}$ will be [IIT 1984]
 (a) 2, 1 (b) $5, \frac{1}{5}$ (c) $7, \frac{1}{7}$ (d) None of these
374. If x is real, then the value of $\frac{x^2 + 34x - 71}{x^2 + 2x - 7}$ does not lie between [Roorkee 1983, 89]
 (a) -9 and -5 (b) -5 and 9 (c) 0 and 9 (d) 5 and 9
375. The adjoining figure shows the graph of $y = ax^2 + bx + c$. Then



- (a) $a < 0$ (b) $b^2 < 4ac$
 (c) $c > 0$ (d) a and b are of opposite signs
376. If $x + 2$ is a common factor of $px^2 + qx + r$ and $qx^2 + px + r$, then
 (a) $p = q = r$ (b) $p = q$ or $p + q + r = 0$ (c) $p = r$ or $p + q + r = 0$ (d) $q = r$ or $p + q + r = 0$
377. $x^2 - 11x + a$ and $x^2 - 14x + 2a$ will have a common factor, if $a =$ [Roorkee 1981]

- (a) 24 (b) 0, 24 (c) 3, 24 (d) 0, 3
378. If $x^2 - 3x + 2$ is a factor of $x^4 - px^2 + q$, then [IIT 1974; MP PET 1995]
 (a) $p = 4, q = 5$ (b) $p = 5, q = 4$ (c) $p = -5, q = -4$ (d) None of these
379. If $x + 1$ is a factor of $x^4 - (p - 3)x^3 - (3p - 5)x^2 + (2p - 7)x + 6$, then p is equal to [IIT 1975]
 (a) -4 (b) 4 (c) -1 (d) 1
380. If $x^2 + px + 1$ is a factor of the expression $ax^3 + bx + c$, then [IIT 1980]
 (a) $a^2 + c^2 = -ab$ (b) $a^2 - c^2 = -ab$ (c) $a^2 - c^2 = ab$ (d) None of these
381. The condition that $x^3 - 3px + 2q$ may be divisible by a factor of the form $x^2 + 2ax + a^2$ is [AMU 2002]
 (a) $3p = 2q$ (b) $3p + 2q = 0$ (c) $p^3 = q^2$ (d) $27p^3 = 4q^2$
382. If x be real then $\frac{(x-a)(x-b)}{x-c}$ will take all real values when [IIT 1984; Karnataka CET 2002]
 (a) $a < b < c$ (b) $a > b > c$ (c) $a < c < b$ (d) Always
383. Let $y = \sqrt{\frac{(x+1)(x-3)}{(x-2)}}$, then all real values of x for which y takes real values, are [IIT 1980]
 (a) $-1 \leq x < 2$ or $x \geq 3$ (b) $-1 \leq x < 3$ or $x > 2$ (c) $1 \leq x < 2$ or $x \geq 3$ (d) None of these
384. The graph of the curve $x^2 = 3x - y - 2$ is
 (a) Between the lines $x = 1$ and $x = \frac{3}{2}$ (b) Between the lines $x = 1$ and $x = 2$
 (c) Strictly below the line $4y = 1$ (d) None of these
385. If $x^2 + px + 1$ is a factor of the expression $ax^3 + bx + c$ then
 (a) $a^2 + c^2 = -ab$ (b) $a^2 - c^2 = -ab$ (c) $a^2 - c^2 = ab$ (d) None of these
386. If $x + \lambda y - 2$ and $x - \mu y + 1$ are factors of the expression $6x^2 - xy - y^2 - 6x + 8y - 12$, then
 (a) $\lambda = \frac{1}{3}, \mu = \frac{1}{2}$ (b) $\lambda = 2, \mu = 3$ (c) $\lambda = \frac{1}{3}, \mu = -\frac{1}{2}$ (d) None of these

Advance Level

387. Given that, for all real x , the expression $\frac{x^2 - 2x + 4}{x^2 + 2x + 4}$ lies between $\frac{1}{3}$ and 3. The values between which the expression $\frac{9 \cdot 3^{2x} + 6 \cdot 3^x + 4}{9 \cdot 3^{2x} - 6 \cdot 3^x + 4}$ lies are [Karnataka CET 1998]
 (a) $\frac{1}{3}$ and 3 (b) -2 and 0 (c) -1 and 1 (d) 0 and 2
388. If x, y, z are real and distinct, then $u = x^2 + 4y^2 + 9z^2 - 6yz - 3zx - 2xy$ is always [IIT 1979]
 (a) Non-negative (b) Non-positive (c) Zero (d) None of these
389. If $x + y$ and $y + 3x$ are two factors of the expression $\lambda x^3 - \mu x^2 y + xy^2 + y^3$, then the third factor is
 (a) $y + 3x$ (b) $y - 3x$ (c) $y - x$ (d) None of these
390. If $\log_{10} x + \log_{10} y \geq 2$ then the smallest possible value of $x + y$ is
 (a) 10 (b) 30 (c) 20 (d) None of these
391. If α be the number of solutions of equation $[\sin x] = x$, where $[x]$ denote the integral part of x and m be the greatest value of $\cos(x^2 + xe^x - [x])$ on the interval $[-1, 1]$, then
 (a) $\alpha = m$ (b) $\alpha < m$ (c) $\alpha > m$ (d) $\alpha \neq m$
392. If $f(x) = 3^x + 4^x + 5^x - 6^x$, then $f(x) < f(3)$ for
 (a) Only one value of x (b) No value of x (c) Only two values of x (d) Infinitely many values of x

200 Quadratic Equations and Inequalities

393. If $f(x) = \sum_{r=0}^{100} a_r x^r$ and $f(0)$ and $f(1)$ are odd numbers, then for any integer x
- (a) $f(x)$ is odd or even according as x is odd or even (b) $f(x)$ is even or odd according as x is odd or even
 (c) $f(x)$ is even for all integral values of x (d) $f(x)$ is odd for all integral values of x
394. If $x \in [2, 4]$ then for the expression $x^2 - 6x + 5 = 0$
- (a) The least value = -4 (b) The greatest value = 4 (c) The least value = 3 (d) The greatest value = -3
395. The value of 'a' for which $(a^2 - 1)x^2 + 2(a - 1)x + 2$ is positive for any x are
- (a) $a \geq 1$ (b) $a \leq 1$ (c) $a \geq -3$ (d) $a \leq -3$ or $a \geq 1$
396. Let $f(x)$ be a quadratic expression which is positive for all real values of x , then for all real x , $10[f(x) + f(-x)]$ is
- (a) > 0 (b) ≥ 0 (c) < 0 (d) ≤ 0
397. The constant term of the quadratic expression $\sum_{k=1}^n \left(x - \frac{1}{k+1}\right) \left(x - \frac{1}{k}\right)$ as $n \rightarrow \infty$ is
- (a) -1 (b) 0 (c) 1 (d) None of these
398. Let $f(x) = (1 + b^2)x^2 + 2bx + 1$ and let $m(b)$ be the minimum value of $f(x)$. As b varies, the range of $m(b)$ is
- (a) $[0, 1]$ (b) $\left[0, \frac{1}{2}\right]$ (c) $\left[\frac{1}{2}, 1\right]$ (d) $(0, 1]$
399. If $p(x)$ be a polynomial satisfying the identity $p(x^2) + 2x^2 + 10x = 2xp(x+1) + 3$, then $p(x)$ is given by
- (a) $2x + 3$ (b) $3x - 4$ (c) $3x + 2$ (d) $2x - 3$
400. Let $y = \frac{\sin x \cos 3x}{\cos x \sin 3x}$, then
- (a) y may be equal to $\frac{1}{3}$ (b) y may be equal to 3
 (c) Set of possible value of y is $\left(-\infty, \frac{1}{3}\right) \cup (3, \infty)$ (d) Set of possible values of y is $\left(-\infty, \frac{1}{3}\right] \cup (3, \infty)$
401. If $a = \frac{x^2 - 2x + 4}{x^2 + 2x + 4}$, and equation of lines AB and CD be $3y = x$ and $y = 3x$ respectively, then for all real x , point $P(a, a^2)$
- (a) Lies in the acute angle between lines AB and CD (b) Lies in the obtuse angle between lines AB and CD
 (c) Cannot be in the acute angle between lines AB and CD (d) Cannot lie in the obtuse angle between lines AB and CD

Position of roots

Basic Level

402. If a, b, c are real numbers such that $a + b + c = 0$, then the quadratic equation $3ax^2 + 2bx + c = 0$ has [MNR 1992; DCE 1995]
- (a) At least one root in $[0, 1]$ (b) At least one root in $[1, 2]$
 (c) At least one root in $[-1, 0]$ (d) None of these
403. The number of values of k for which the equation $x^2 - 3x + k = 0$ has two real and distinct roots lying in the interval $(0, 1)$, are [UPSEAT 2001; Kurukshetra CEET 2002]
- (a) 0 (b) 2 (c) 3 (d) Infinitely many
404. The value of k for which the equation $(k - 2)x^2 + 8x + k + 4 = 0$ has both real, distinct and negative is [Orissa JEE 2002]
- (a) 0 (b) 2 (c) 3 (d) -4

Advance Level

- 405.** Let a, b, c be real number $a \neq 0$. If α is a root of $a^2x^2 + bx + c = 0$; β is a root of $a^2x^2 - bx - c = 0$ and $0 < \alpha < \beta$; then the equation $a^2x^2 + 2bx + 2c = 0$ has a root γ which always satisfies [IIT 1989]
- (a) $\gamma = \frac{\alpha + \beta}{2}$ (b) $\gamma = \alpha + \frac{\beta}{2}$ (c) $\gamma = \alpha$ (d) $\alpha < \gamma < \beta$
- 406.** Let a, b, c be non-zero real numbers such that $\int_0^1 (1 + \cos^8 x)(ax^2 + bx + c) dx = \int_0^2 (1 + \cos^8 x)(ax^2 + bx + c) dx$, then the quadratic equation $ax^2 + bx + c = 0$ has
- (a) No root in $(0, 2)$ (b) At least one root in $(0, 1)$ (c) A double root in $(0, 2)$ (d) Two imaginary roots
- 407.** For the equation $2x^2 + 6\sqrt{2}x + 1 = 0$
- (a) Roots are rational (b) If one root is $p + \sqrt{q}$ then the other is $-p + \sqrt{q}$
 (c) Roots are irrational (d) If one root is $p + \sqrt{q}$ then the other is $p - \sqrt{q}$
- 408.** The values of a for which both roots of the equation $(1 - a^2)x^2 + 2ax - 1 = 0$ lie between 0 and 1 are given by
- (a) $a > 2$ (b) $1 < a < 2$ (c) $-\infty < a < \infty$ (d) None of these
- 409.** If p, q be non-zero real numbers and $f(x) \neq 0$ in $[0, 2]$ and $\int_0^1 f(x).(x^2 + px + q) dx = \int_0^2 f(x).(x^2 + px + q) dx = 0$ then equation $x^2 + px + q = 0$ has
- (a) Two imaginary roots (b) No root in $(0, 2)$
 (c) One root in $(0, 1)$ and other in $(1, 2)$ (d) One root in $(-\infty, 0)$ and other in $(2, \infty)$
- 410.** If $a, b, c \in R, a \neq 0$ and $(b - 1)^2 < 4ac$, then the number of real roots of the system of equation (in three unknowns x_1, x_2, x_3)
 $ax_1^2 + bx_1 + c = x_2, ax_2^2 + bx_2 + c = x_3, ax_3^2 + bx_3 + c = x_1$ is
- (a) 0 (b) 1 (c) 2 (d) 3
- 411.** If $0 < \alpha < \frac{\pi}{4}$, equation $(x - \sin \alpha)(x - \cos \alpha) - 2 = 0$ has
- (a) Both roots in $(\sin \alpha, \cos \alpha)$ (b) Both roots in $(\cos \alpha, \sin \alpha)$
 (c) One root in $(-\infty, \cos \alpha)$ and other in $(\sin \alpha, \infty)$ (d) One root in $(-\infty, \sin \alpha)$ and other in $(\cos \alpha, \infty)$
- 412.** For equation $x^3 - 6x^2 + 9x + k = 0$ to have exactly one root in $(1, 3)$, the set of values of k is
- (a) $(-4, 0)$ (b) $(1, 3)$ (c) $(0, 4)$ (d) None of these
- 413.** Let $f(x) = x^3 - 6x^2 + 3(1 + \pi)x + 7, p > q > r$, then $\frac{\{x - f(p)\}\{x - f(r)\}}{x - f(q)}$ has no value in
- (a) (p, q) (b) (q, r) (c) (r, ∞) (d) None of these
- 414.** If $a + b + 2c = 0, c \neq 0$, then equation $ax^2 + bx + c = 0$ has
- (a) At least one root in $(0, 1)$ (b) At least one root in $(0, 2)$
 (c) At least one root in $(-1, 1)$ (d) None of these
- 415.** If $ax^2 - bx + c = 0$ has two distinct real roots in $(0, 1)$, where $a, b, c \in N$, then $16c(a - b + c)$
- (a) $= a^2$ (b) $< a^2$ (c) $> a^2$ (d) $\geq a^2$

Solution of Quadratic inequations

Basic Level

- 416.** If $a < b$, then the solution of $x^2 + (a + b)x + ab < 0$, is given by
- (a) $a < x < b$ (b) $x < a$ or $x > b$ (c) $-b < x < -a$ (d) $x < -b$ or $x < -a$
- 417.** The solution of $6 + x - x^2 > 0$ is [DCE 2000; Kurakshetra CEE 1999]
- (a) $-1 < x < 2$ (b) $-2 < x < 3$ (c) $-2 < x < -1$ (d) None of these

202 Quadratic Equations and Inequalities

418. For all $x \in R$, if $mx^2 - 9mx + 5m + 1 > 0$, then m lies in the interval [AMU 1989]
- (a) $\left(-\frac{4}{61}, 0\right)$ (b) $\left[0, \frac{4}{61}\right)$ (c) $\left(\frac{4}{61}, \frac{61}{4}\right)$ (d) $\left[-\frac{61}{4}, 0\right]$
419. If $x^2 - 1$ is a factor of $x^4 + ax^3 + 3x - b$, then
- (a) $a = 3, b = -1$ (b) $a = -3, b = 1$ (c) $a = 3, b = 1$ (d) None of these
420. If $(x - 1)^3$ is factor of $x^4 + ax^3 + bx^2 + cx - 1$ then the other factor is
- (a) $x - 3$ (b) $x + 1$ (c) $x + 2$ (d) None of these
421. The set of values of x which satisfy $5x + 2 < 3x + 8$ and $\frac{x+2}{x-1} < 4$, is [EAMCET 1989]
- (a) (2, 3) (b) $(-\infty, 1) \cup (2, 3)$ (c) $(-\infty, 1)$ (d) (1, 3)
422. The solution of the equation $2x^2 + 3x - 9 \leq 0$ is given by [Kurukshetra CEE 1998]
- (a) $\frac{3}{2} \leq x \leq 3$ (b) $-3 \leq x \leq \frac{3}{2}$ (c) $-3 \leq x \leq 3$ (d) $\frac{3}{2} \leq x \leq 2$
423. The complete solution of the inequation $x^2 - 4x < 12$ is [AMU 1999]
- (a) $x < -2$ or $x > 6$ (b) $-6 < x < 2$ (c) $2 < x < 6$ (d) $-2 < x < 6$
424. If x is real and satisfies $x + 2 > \sqrt{x + 4}$, then [AMU 1999]
- (a) $x < -2$ (b) $x > 0$ (c) $-3 < x < 0$ (d) $-3 < x < 4$
425. If $a < 0$ then the inequality $ax^2 - 2x + 4 > 0$ has the solution represented by [AMU 2001]
- (a) $\frac{1 + \sqrt{1 - 4a}}{a} > x > \frac{1 - \sqrt{1 - 4a}}{a}$ (b) $x < \frac{1 - \sqrt{1 - 4a}}{a}$
- (c) $x < 2$ (d) $2 > x > \frac{1 + \sqrt{1 - 4a}}{a}$

Advance Level

426. If x satisfies $|x - 1| + |x - 2| + |x - 3| \geq 6$, then
- (a) $0 \leq x \leq 4$ (b) $x \leq -2$ or $x \geq 4$ (c) $x \leq 0$ (d) None of these
427. The number of positive integral solutions of $\frac{x^2(3x - 4)^3(x - 2)^4}{(x - 5)^5(2x - 7)^6} \leq 0$ is
- (a) 4 (b) 3 (c) 2 (d) 1
428. If $5^x + (2\sqrt{3})2^x \geq 13^x$, then the solution set for x is
- (a) $[2, \infty)$ (b) $\{2\}$ (c) $(-\infty, 2]$ (d) $[0, 2]$
429. The inequality $|2x - 3| < 1$ is valid when x lies in [IIT 1993]
- (a) (3, 4) (b) (1, 2) (c) (-1, 2) (d) (-4, 3)
430. The graph of the function $y = 16x^2 + 8(a + 5)x - 7a - 5$ is strictly above the x -axis, then 'a' must satisfy the inequality
- (a) $-15 < a < -2$ (b) $-2 < a < -1$ (c) $5 < a < 7$ (d) None of these
431. If x is a real number such that $x(x^2 + 1), (-1/2)x^2, 6$ are three consecutive terms of an A.P. then the next two consecutive term of the A.P. are
- (a) 14, 6 (b) -2, -10 (c) 14, 22 (d) None of these
432. If x, y are rational numbers such that $x + y + (x - 2y)\sqrt{2} = 2x - y + (x - y - 1)\sqrt{6}$, then
- (a) x and y cannot be determined (b) $x = 2, y = 1$
- (c) $x = 5, y = 1$ (d) None of these

433. If $[x]$ = the greatest interger less than or equal to x , and (x) = the least interger greatest than or equal to x and $[x]^2 + (x)^2 > 25$ then x belongs to
 (a) $[3, 4]$ (b) $(-\infty, -4]$ (c) $[4, +\infty)$ (d) $(-\infty, -4] \cup [4, +\infty)$
434. The set of real values of x satisfying $|x-1| \leq 3$ and $|x-1| \geq 1$ is
 (a) $[2, 4]$ (b) $(-\infty, 2] \cup [4, +\infty)$ (c) $[-2, 0] \cup [2, 4]$ (d) None of these
435. The set of real values of x satisfying $||x-1|-1| \leq 1$ is
 (a) $[-1, 3]$ (b) $[0, 2]$ (c) $[-1, 1]$ (d) None of these
436. If $x \in Z$ (the set of integers) such that $x^2 - 3x < 4$ then the number of possible values of x is
 (a) 3 (b) 4 (c) 6 (d) None of these
437. If x is an interger satisfying $x^2 - 6x + 5 \leq 0$ and $x^2 - 2x > 0$ then the number of possible values of x is
 (a) 3 (b) 4 (c) 2 (d) Infinite
438. The solution set of the ineuation $\log_{1/3}(x^2 + x + 1) + 1 > 0$ is
 (a) $(-\infty, -2) \cup (1, +\infty)$ (b) $[-1, 2]$ (c) $(-2, 1)$ (d) $(-\infty, +\infty)$
439. If $3^{x/2} + 2^x > 25$ then the solution set is
 (a) R (b) $(2, +\infty)$ (c) $(4, +\infty)$ (d) None of these
440. The solution set of $\frac{x^2 - 3x + 4}{x + 1} > 1, x \in R$, is
 (a) $(3, +\infty)$ (b) $(-1, 1) \cup (3, +\infty)$ (c) $[-1, 1] \cup [3, +\infty)$ (d) None of these
441. The equation $|x+1||x-1| = a^2 - 2a - 3$ can have real solutions for x if a belongs to
 (a) $(-\infty, -1] \cup [3, +\infty)$ (b) $[1 - \sqrt{5}, 1 + \sqrt{5}]$ (c) $[1 - \sqrt{5}, -1] \cup [3, 1 + \sqrt{5}]$ (d) None of these

Miscellaneous Problems

Basic Level

442. If $x^2 + 2x + 2xy + my - 3$ has two rational factors, then the value of m will be [Rajasthan PET 1990]
 (a) $-6, -2$ (b) $-6, 2$ (c) $6, -2$ (d) $6, 2$
443. If $x^2 - hx - 21 = 0, x^2 - 3hx + 35 = 0$ ($h > 0$) has a common root, then the value of h is equal to [EAMCET 1986]
 (a) 1 (b) 2 (c) 3 (d) 4
444. Minimum value of $(a+b+c+d)\left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d}\right)$ is
 (a) 4 (b) 9 (c) 16 (d) 25
445. Let $f(x) = ax^3 + 5x^2 - bx + 1$. If $f(x)$ when divided by $2x + 1$ leaves 5 as remainder, and $f'(x)$ is divisible by $3x - 1$ then
 (a) $a = 26, b = 10$ (b) $a = 24, b = 11$ (c) $a = 26, b = 12$ (d) None of these
446. $x^{3^n} + y^{3^n}$ is divisible by $x + y$ if
 (a) n is any integer ≥ 0 (b) n is an odd positive integer
 (c) n is an even positive integer (d) n is a rational number
447. The number of solution of the equation $|x| = \cos x$ is
 (a) One (b) Two (c) Three (d) Zero
448. The line $y + 14 = 0$ cuts the curve whose equation is $x(x^2 + x + 1) + y = 0$ at
 (a) Three real points (b) One real point (c) At least one real point (d) No real point

204 Quadratic Equations and Inequalities

449. Let R = the set of real numbers, \mathbb{J} = the set of integers, N = the set of natural numbers. If S be the solution set of the equation $(x)^2 + [x]^2 = (x-1)^2 + [x+1]^2$, where (x) = the least integer greater than or equal to x and $[x]$ = the greatest integer less than or equal to x , then
- (a) $S = R$ (b) $S = R - Z$ (c) $S = R - N$ (d) None of these
450. The number of real roots of $x^8 - x^5 + x^2 - x + 1 = 0$ is equal to
- (a) 0 (b) 2 (c) 4 (d) 6
451. The number of positive real roots of $x^4 - 4x - 1 = 0$ is
- (a) 3 (b) 2 (c) 1 (d) 0
452. The number of negative real roots of $x^4 - 4x - 1 = 0$ is
- (a) 3 (b) 2 (c) 1 (d) 0
453. The number of complex roots of the equation $x^4 - 4x - 1 = 0$ is
- (a) 3 (b) 2 (c) 1 (d) 0
454. $x^2 - 4$ is a factor of $f(x) = (a_1x^2 + b_1x + c_1)(a_2x^2 + b_2x + c_2)$ if
- (a) $b_1 = 0, c_1 + 4a_1 = 0$ (b) $b_2 = 0, c_2 + 4a_2 = 0$
(c) $4a_1 + 2b_1 + c_1 = 0, 4a_2 + c_2 = 2b_2$ (d) $4a_1 + c_1 = 2b_1, 4a_2 + 2b_2 + c_2 = 0$





Answer Sheet

Quadratic Equations and Inequalities **Assignment (Basic and Advance Level)**

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
a	c	a	d	d	c	d	d	b	d	d	d	c	d	c	a	c	b	d	c
21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40
c	d	c	c	b	c	d	c	b	c	b	c	a	a	b	c	b	b	b	c
41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60
a	c	a	a	d	d	c	a	a	c	b	a	a	a	a	d	b	b	c	b
61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80
c	a	c	b	a	d	c	a	a	a	d	c	b	d	b	b	b	c	b	a
81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
d	c	b	b	b	a	c	c	b	d	b	c	b	c	b	a	b	a	a	a
101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120
c	b	c	b	a	a	a	a	c	d	c	c	b	a	c	d	b	c	b	a
121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140
d	a	a	c	c	c	d	d	c	c	b	b	b	c	b	d	d	a	a	b
141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160
a	b	a	a,c	c	b	c	a	c	b	a	b	a	d	b	d	a	a	b	b
161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180
b	c	b	b	b	a	c	c	b	b	a	b	c	d	b	b	a	a	b	d
181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200
a	a	a	c	a	c	d	b	c	d	b	d	a	c	a	a	b	a	d	d
201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220
b	c	c	c	a	d	b	d	d	c	d	c	a	a	d	a	a	d	b	a
221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240
a	b	b	b	d	c	a	d	a	a	d	c	b	a	c	b	b	c	a	d
241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260
b	b	a	c	c	b	d	b	d	c	a	c	d	b	b	b	a	b	a	a
261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280
b	a	c	a	b	a	c	a	d	b	b	b	b	a	a	a	a	b	d	a
281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300
d	a	a	c	a	b	a	b	a	a	a	a	b	d	b	d	d	d	b	c
301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320
a	b,c	a,c	b,c	b	b	b	c	b	b	d	c	b	c	a	d	b	a	d	c
321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340
a	d	b	a	c	a,c	c	b	a,c	c	a,b	c	c	a	a,b	c	c	c	d	d
341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360
c	b	b	b	c	c	c	c	a	a	b	b	a	c,d	b,c	a,b	a,c	c	c	a
361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380
c	a	a,b,c,d	a	a	a	a	c	b	c	d	d	c	d	d	b	b	b	b	c
381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400
c	c	a	c	c	a	a	a	b	c	a	d	d	a,d	d	a	c	d	a	c
401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420
d	a	a	c	d	b	b,c	a	c	a	d	a	d	a,b,c	b	c	b	b	b	b
421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440
b	b	d	b	a	c	b	c	b	a	c	b	d	a	a	a,b	a	c	c	b

441	442	443	444	445	446	447	448	449	450	451	452	453	454
a,c	c	d	c	c	a	b	b	b	a	c	c	b	a,b,c,d

