



Assignment

Significant Digit and Rounding off Numbers, Fundamental concepts

Basic Level

- The number 3.14150 rounded to 3 decimals is [MP PET 2000]
 (a) 3.14 (b) 3.141 (c) 3.142 (d) None of these
- The number of significant digits in 0.003050 is
 (a) 7 (b) 6 (c) 4 (d) None of these
- The number of significant digits in 20.035 is
 (a) 3 (b) 5 (c) 4 (d) None of these
- The number of significant digits in 20340 is
 (a) 4 (b) 5 (c) 3 (d) None of these
- The number 0.0008857 when rounded off to three significant digits yields
 (a) 0.001 (b) 0.000886 (c) 0.000885 (d) None of these
- The number 3.68451 when rounded off to three decimal places becomes
 (a) 3.68 (b) 3.684 (c) 3.685 (d) None of these
- The number of significant digits in the number 0.00452000 is
 (a) 3 (b) 5 (c) 8 (d) None of these
- When a number is approximated to n decimal places by chopping off the extra digits, then the absolute value of the relative error does not exceed
 (a) 10^{-n} (b) 10^{-n+1} (c) $0.5 \times 10^{-n+1}$ (d) None of these
- When the number 6.878652 is rounded off to five significant figures, then the round off error is
 (a) -0.000048 (b) -0.00048 (c) 0.000048 (d) 0.00048
- The number 0.0009845 when rounded off to three significant digits yields [DCE 1998]



- (a) 0.001 (b) 0.000987 (c) 0.000985 (d) None of these
11. A decimal number is chopped off to four decimal places, then the absolute value of the relative error is not greater than [DCE 1996]
 (a) 10^{-2} (b) 10^{-3} (c) 10^{-4} (d) None of these
12. If e_1 and e_2 are absolute errors in two numbers n_1 and n_2 respectively due to rounding or truncation, then $\left| \frac{e_1}{n_1} + \frac{e_2}{n_2} \right|$
 (a) Is equal to $e_1 + e_2$ (b) Is less than $e_1 + e_2$
 (c) Is less than or equal to $e_1 + e_2$ (d) Is greater than or equal to $e_1 + e_2$
13. In general the ratio of the truncation error to that of round off error is
 (a) 1 : 2 (b) 2 : 1 (c) 1 : 1 (d) None of these
14. The equation $e^{-2x} - \sin x + 1 = 0$ is of the form
 (a) Algebraic (b) Linear (c) Quadratic (d) Transcendental
15. The root of the equation $x^3 - 6x + 1 = 0$ lies in the interval
 (a) (2, 3) (b) (3, 4) (c) (3, 5) (d) (4, 6)
16. The root of the equation $x^3 - 3x - 5 = 0$ in the interval (1, 2) is
 (a) 1.13 (b) 1.98
 (c) 1.54 (d) No root lies in the interval (1, 2)
17. The equation $f(x) = 0$ has repeated root $a \in (x_1, x_2)$, if
 (a) $f'(a) < 0$ (b) $f'(a) > 0$ (c) $f'(a) = 0$ (d) None of these
18. The root of the equation $2x - \log_{10} x = 7$ lies between
 (a) 3 and 3.5 (b) 2 and 3 (c) 3.5 and 4 (d) None of these
19. For the equation $f(x) = 0$, if $f(a) < 0$, $f(b) > 0$, $f(c) > 0$ and $b > c$ then we will discard the value of the function $f(x)$ at the point
 (a) a (b) b (c) c (d) Anyone out of a, b, c
20. The positive root of the equation $e^x + x - 3 = 0$ lies in the interval
 (a) (0, 1) (b) (1, 2) (c) (2, 3) (d) (2, 4)
21. The positive root of the equation $x^3 - 2x - 5 = 0$ lies in the interval
 (a) (0, 1) (b) (1, 2) (c) (2, 3) (d) (3, 4)
22. One real root of the equation $x^3 - 5x + 1 = 0$ must lie in the interval
 (a) (0, 1) (b) (1, 2) (c) (-1, 0) (d) (-2, 0)
23. The number of positive roots of the equation $x^3 - 3x + 5 = 0$ is [MP PET 1998]



- (a) 1 (b) 2 (c) 3 (d) None of these

24. Let $f(x) = 0$ be an equation and x_1, x_2 be two real numbers such that $f(x_1)f(x_2) < 0$, then [MP PET 1989, 1997]
- (a) At least one root of the equation lies in the interval (x_1, x_2)
 (b) No root of the equation lies in the interval (x_1, x_2)
 (c) Either no root or more than one root of the equation lies the interval (x_1, x_2)
 (d) None of these
25. Let $f(x) = 0$ be an equation let x_1, x_2 be two real numbers such that $f(x_1)f(x_2) > 0$, then
- (a) At least one root of the equation lies in (x_1, x_2)
 (b) No root of the equation lies in (x_1, x_2)
 (c) Either no root or an even number of roots lie in (x_1, x_2)
 (d) None of these
26. If for $f(x) = 0$, $f(a) < 0$ and $f(b) > 0$, then one root of $f(x) = 0$ is
- (a) Between a and b (b) One of from a and b
 (c) Less than a and greater than b (d) None of these
27. If $f(a)f(b) < 0$, then an approximate value of a real root of $f(x) = 0$ lying between a and b is given by
- (a) $\frac{af(b) - bf(a)}{b - a}$ (b) $\frac{bf(a) - af(b)}{b - a}$
 (c) $\frac{af(b) - bf(a)}{f(b) - f(a)}$ (d) None of these

Successive bisection method

Basic Level

28. One root of $x^3 - x - 4 = 0$ lies in $(1, 2)$. In bisection method, after first iteration the root lies in the interval
- (a) $(1, 1.5)$ (b) $(1.5, 2.0)$ (c) $(1.25, 1.75)$ (d) $(1.75, 2)$
29. A root of the equation $x^3 - x - 1 = 0$ lies between 1 and 2. Its approximate value as obtained by applying bisection method 3 times is [MP PET 1993]
- (a) 1.375 (b) 1.625 (c) 1.125 (d) 1.25
30. A root of the equation $x^3 - x - 4 = 0$ lies between 1 and 2. Its approximate value, as obtained by applying bisection method 3 times, is

- (a) 1.375 (b) 1.750 (c) 1.975 (d) 1.875
31. Performing 3 iterations of bisection method, the smallest positive approximate root of equation $x^3 - 5x + 1 = 0$ is [MP PET 1996]
 (a) 0.25 (b) 0.125 (c) 0.50 (d) 0.1875
32. A root of the equation $x^3 - 3x - 5 = 0$ lies between 2 and 2.5. Its approximate value, by applying bisection method 3 times is
 (a) 2.0625 (b) 2.3125 (c) 2.3725 (d) 2.4225
33. If for the function $f(x) = 0$, $f(a) < 0$ and $f(b) > 0$, then the value of x in first iteration is
 (a) $\frac{a+b}{2}$ (b) $\frac{b-a}{2}$ (c) $\frac{2a-b}{2}$ (d) $\frac{2b-a}{2}$
34. Using successive bisection method, a root of the equation $x^3 - 4x + 1 = 0$ lies between 1 and 2, at the end of first interaction, it lies between [DCE 1996]
 (a) 1.62 and 1.75 (b) 1.5 and 1.75 (c) 1.75 and 1.87 (d) None of these
35. The nearest real root of the equation $xe^x - 2 = 0$ correct to two decimal places, is
 (a) 1.08 (b) 0.92 (c) 0.85 (d) 0.80

*Regula-Falsi method**Basic Level*

36. By the false position method, the root of the equation $x^3 - 9x + 1 = 0$ lies in interval (2, 4) after first iteration. It is
 (a) 3 (b) 2.5 (c) 3.57 (d) 2.47
37. The formula [where $f(x_{n-1})$ and $f(x_n)$ have opposite sign at each step $n \geq 1$] of method of False position of successive approximation to find the approximate value of a root of the equation $f(x) = 0$ is [MP PET 1995, 97]
 (a) $x_{n+1} = x_n - \frac{f(x_n) - f(x_{n-1})}{f(x_n)}(x_n - x_{n-1})$ (b) $x_{n+1} = x_n - \frac{f(x_n)}{f(x_n) - f(x_{n-1})}(x_n - x_{n-1})$
 (c) $x_{n+1} = x_n + \frac{f(x_n) + f(x_{n-1})}{f(x_n)}(x_n - x_{n-1})$ (d) $x_{n+1} = x_n + \frac{f(x_n)}{f(x_n) + f(x_{n-1})}(x_n - x_{n-1})$
38. By false positioning, the second approximation of a root of equation $f(x) = 0$ is (where x_0, x_1 are initial and first approximations respectively) [MP PET 1996; DCE 2001]
 (a) $x_0 - \frac{f(x_0)}{f(x_1) - f(x_0)}$ (b) $\frac{x_0 f(x_1) - x_1 f(x_0)}{f(x_1) - f(x_0)}$ (c) $\frac{x_0 f(x_0) - x_1 f(x_1)}{f(x_1) - f(x_0)}$ (d) $x_0 - \frac{f(x_0)}{f(x_1) - f(x_0)}$
39. A root of the equation $x^3 - 18 = 0$ lies between 2 and 3. The value of the root by the method of false position is
 (a) 2.526 (b) 2.536 (c) 2.546 (d) 2.556



40. The equation $x^3 - 3x + 4 = 0$ has only one real root. What is its first approximate value as obtained by the method of false position in $(-3, -2)$ [MP PET 1999]
- (a) -2.125 (b) 2.125 (c) -2.812 (d) 2.812
41. A root of equation $x^3 + 2x - 5 = 0$ lies between 1 and 1.5. its value as obtained by applying the method of false position only once is [MP PET 1993]
- (a) $\frac{4}{3}$ (b) $\frac{35}{27}$ (c) $\frac{23}{25}$ (d) $\frac{5}{4}$

Newton-Raphson method
Basic Level

42. If successive approximations are given by $x_1, x_2, x_3, \dots, x_n, x_{n+1}$, then Newton-Raphson formula is given as [MP PET 1993, 95]
- (a) $x_{n+1} = x_n + \frac{f(x_{n+1})}{f'(x)}$ (b) $x_{n+1} = x_n + \frac{f(x_n)}{f'(x_n)}$
- (c) $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$ (d) $x_{n+1} = x_n - \frac{f'(x_n)}{f(x_n)}$
43. Newton-Raphson method is applicable only when
- (a) $f(x) \neq 0$ in the neighbourhood of actual root $x = \alpha$ (b) $f'(x) \neq 0$ in the neighbourhood of actual root $x = \alpha$
- (c) $f''(x) \neq 0$ in the neighbourhood of actual root $x = \alpha$ (d) None of these
44. Newton-Raphson processes has a
- (a) Linear convergence (b) Quadratic convergence (c) Cubic convergence (d) None of these
45. The condition for convergence of the Newton-Raphson method to a root α is [MP PET 2001]
- (a) $\frac{1}{2} \frac{f'(\alpha)}{f''(\alpha)} < 1$ (b) $\frac{f'(\alpha)}{f''(\alpha)} < 1$
- (c) $\frac{1}{2} \frac{f'(\alpha)}{f''(\alpha)} > 1$ (d) None of these
46. The real root of the equation $x^3 - x - 5 = 0$ lying between -1 and 2 after first iteration by Newton-Raphson method is
- (a) 1.909 (b) 1.904 (c) 1.921 (d) 1.940
47. A root of the equation $x^3 - 4x + 1 = 0$ lies between 1 and 2. Its value as obtained by using Newton-Raphson method is
- (a) 1.775 (b) 1.850 (c) 1.875 (d) 1.950
48. The value of x_0 (the initial value of x) to get the solution in interval $(0.5, 0.75)$ of the equation $x^3 - 5x + 3 = 0$ by Newton-Raphson method, is
- (a) 0.5 (b) 0.75 (c) 0.625 (d) None of these



49. If a and $a + h$ are two consecutive approximate roots of the equation $f(x) = 0$ as obtained by Newton's method, then h is equal to [MP PET 1999]
- (a) $f(a)/f'(a)$ (b) $f'(a)/f(a)$ (c) $-f'(a)/f(a)$ (d) $-f(a)/f'(a)$
50. The Newton-Raphson method converges fast if $f'(\alpha)$ is (α is the exact value of the root) [DCE 1998]
- (a) Small (b) Large (c) 0 (d) None of these

Advance Level

51. If one root of the equation $f(x) = 0$ is near to x_0 then the first approximation of this root as calculated by Newton-Raphson method is the abscissa of the point where the following straight line intersects the x -axis [MP PET 1998]
- (a) Normal to the curve $y = f(x)$ at the point $(x_0, f(x_0))$
- (b) Tangent to the curve $y = f(x)$ at the point $(x_0, f(x_0))$
- (c) The straight line through the point $(x_0, f(x_0))$ having the gradient $\frac{1}{f'(x_0)}$
- (d) The ordinate through the point $(x_0, f(x_0))$
52. A root of the equation $x^3 - 3x - 5 = 0$ lies between 2 and 2.5. Its value as obtained by using Newton-Raphson method, is
- (a) 2.25 (b) 2.33 (c) 2.35 (d) 2.45
53. After second iteration of Newton-Raphson method, the positive root of equation $x^2 = 3$ is (taking initial approximation $\frac{3}{2}$) [MP PET 1996]
- (a) $\frac{3}{2}$ (b) $\frac{7}{4}$ (c) $\frac{97}{56}$ (d) $\frac{347}{200}$
54. If one root of the equation $x^3 + x^2 - 1 = 0$ is near to 1.0, then by Newton-Raphson method the first calculated approximate value of this root is [MP PET 1998]
- (a) 0.9 (b) 0.6 (c) 1.2 (d) 0.8
55. The approximate value of a root of the equation $x^3 - 3x - 5 = 0$ at the end of the second iteration by taking the initial value of the roots as 2, and by using Newton-Raphson method, is [AI CBSE 1990]
- (a) 2.2806 (b) 2.2701 (c) 2.3333 (d) None of these
56. Newton-Raphson method is used to calculate $\sqrt[3]{65}$ by solving $x^3 = 65$. If $x_0 = 4$ is taken as initial approximation then the first approximation x_1 is [AMU 1999]
- (a) $65/16$ (b) $131/32$ (c) $191/48$ (d) $193/48$
57. Starting with $x_0 = 1$, the next approximation x_1 to $2^{1/3}$ obtained by Newton's method is [DCE 1997]

(a) $\frac{5}{3}$

(b) $\frac{4}{3}$

(c) $\frac{5}{4}$

(d) $\frac{5}{6}$

Trapezoidal rule

Basic Level

58. Approximate value of $\int_{x_0}^{x_0+nh} y dx$ by Trapezoidal rule, is [MP PET 1993, 97]

[Where $y(x_i) = y_i$, $x_{i+1} - x_i = h$, $i = 0, 1, 2, \dots, n$]

(a) $\frac{h}{2} [y_0 + y_n + 2(y_1 + y_2 + y_3 + \dots + y_{n-1})]$

(b) $\frac{h}{3} [y_0 + y_n + 4(y_1 + y_3 + y_5 + \dots + y_{n-1}) + 2(y_2 + y_4 + \dots + y_{n-2})]$

(c) $\frac{h}{4} [y_0 + y_n + 2(y_1 + y_3 + y_5 + \dots + y_{n-1}) + 4(y_2 + y_4 + \dots + y_{n-2})]$

(d) $\frac{h}{2} [y_0 + y_2 + y_4 + \dots + y_n] + 2(y_1 + y_3 + y_5 + \dots + y_{n-1})]$

59. Trapezoidal rule for evaluation of $\int_a^b f(x) dx$ requires the interval (a, b) to be divided into [DCE 1994; MP PET 1996]

(a) $2n$ sub-intervals of equal width(b) $2n + 1$ sub-intervals of equal width

(c) Any number of sub-intervals of equal width

(d) $3n$ sub-intervals of equal width

60. The value of $f(x)$ is given only at $x = 0, \frac{1}{3}, \frac{2}{3}, 1$. Which of the following can be used to evaluate $\int_0^1 f(x) dx$ approximately [MP PET 1999]

(a) Trapezoidal rule

(b) Simpson rule

(c) Trapezoidal as well as Simpson rule

(d) None of these

61. A river is 80 metre wide. Its depth d metre and corresponding distance x metre from one bank is given below in table

x :	0	10	20	30	40	50	60	70	80
y :	0	4	7	9	12	15	14	8	3

Then the approximate area of cross-section of river by Trapezoidal rule, is [MP PET 1994]

(a) 710 sq.m

(b) 730 sq.m

(c) 705 sq.m

(d) 750 sq.m

62. A curve passes through the points given by the following table

x :	1	2	3	4	5
y :	10	50	70	80	100

By Trapezoidal rule, the area bounded by the curve, the x -axis and the lines $x = 1$, $x = 5$, is

(a) 310

(b) 255

(c) 305

(d) 275

63. From the following table, using Trapezoidal rule, the area bounded by the curve, the x -axis and the lines $x = 7.47$, $x = 7.52$, is

x :	7.47	7.48	7.49	7.50	7.51	7.52
-------	------	------	------	------	------	------



$f(x)$: 1.93 1.95 1.98 2.01 2.03 2.06

- (a) 0.0996 (b) 0.0896 (c) 0.1096 (d) 0.0776

64. Let $f(0) = 1$, $f(1) = 2.72$, then the trapezoidal rule gives approximate value of $\int_0^1 f(x) dx$ [MP PET 1999; DCE 2001]

- (a) 3.72 (b) 1.86 (c) 1.72 (d) 0.86

65. By Trapezoidal rule, the value of $\int_0^1 x^3 dx$ considering five sub-intervals, is

- (a) 0.21 (b) 0.23 (c) 0.24 (d) 0.26

Advance Level

66. The approximate value of $\int_1^9 x^2 dx$ by using Trapezoidal rule with 4 equal intervals is [EAMCET 2002]

- (a) 243 (b) 248 (c) 242.8 (d) 242.5

67. Taking $n = 4$, by trapezoidal rule, the value of $\int_0^2 \frac{dx}{1+x}$ is [DCE 1999, 2000]

- (a) 1.1125 (b) 1.1176 (c) 1.118 (d) None of these

68. With the help of trapezoidal rule for numerical integration and the following table

x :	0	0.25	0.50	0.75	1
$f(x)$:	0	0.0625	0.2500	0.5625	1

The value of $\int_0^1 f(x) dx$ is [MP PET 1996]

- (a) 0.35342 (b) 0.34375 (c) 0.34457 (d) 0.33334

69. If for $n = 3$, the integral $\int_1^{10} x^3 dx$ is approximately evaluated by Trapezoidal rule $\int_1^{10} x^3 dx = 3 \left[\frac{1+10^3}{2} + \alpha + 7^3 \right]$, then $\alpha =$

- (a) 3^3 (b) 4^3 (c) 5^3 (d) 6^3

[MP PET 2000]

70. By trapezoidal rule, the value of $\int_1^2 \frac{1}{x} dx$, (using five ordinates) is nearly [DCE 1994]

- (a) 0.216 (b) 0.697 (c) 0.921 (d) None of these

Simpson's one third rule

Basic Level



71. The value of $\int_{x_0}^{x_0+nh} y dx$, n is even number, by Simpson's one-third rule is [MP PET 1995]
- (a) $\frac{h}{3} [(y_0 + y_n) + 2(y_1 + y_3 + \dots + y_{n-1}) + 4(y_2 + y_4 + \dots + y_{n-2})]$ (b) $\frac{h}{3} [(y_0 + y_n) + 4(y_1 + y_3 + \dots + y_{n-1}) + 2(y_2 + y_4 + \dots + y_{n-2})]$
- (c) $\frac{h}{3} [(y_0 + y_n) - 2(y_1 + y_3 + \dots + y_{n-1}) + 4(y_2 + y_4 + \dots + y_{n-2})]$ (d) None of these
72. Simpson's one-third rule for evaluation $\int_a^b f(x) dx$ requires the interval $[a, b]$ to be divided into [DCE 1999]
- (a) An even number of sub-intervals of equal width (b) Any number of sub-intervals
- (c) Any number of sub-intervals of equal width (d) An odd number of sub-intervals of equal width
73. Simpson rule for evaluation of $\int_a^b f(x) dx$ requires the interval (a, b) to be divided into [Haryana CEE 1993; DCE 1994]
- (a) $3n$ intervals (b) $2n + 1$ intervals (c) $2n$ intervals (d) Any number of intervals
74. To calculate approximate value of π by Simpson's rule, the approximate formula is [MP PET 2000]
- (a) $\int_0^1 \left(\frac{1}{1+x^2} \right) dx, n = 16$ (b) $\int_0^1 \left(\frac{1}{1+x^2} \right) dx, n = 9$ (c) $\int_0^1 \left(\frac{1}{1+x} \right) dx, n = 11$ (d) $\int_0^1 \left(\frac{1}{1+x} \right) dx, n = 9$
75. In Simpson's one-third rule, the curve $y = f(x)$ is assumed to be a [MP PET 2001]
- (a) Circle (b) Parabola (c) Hyperbola (d) None of these
76. A river is 80 feet wide. The depth d (in feet) of the river at a distance of x feet from one bank is given by the following table
- | | | | | | | | | | |
|-------|---|----|----|----|----|----|----|----|----|
| x : | 0 | 10 | 20 | 30 | 40 | 50 | 60 | 70 | 80 |
| y : | 0 | 4 | 7 | 9 | 12 | 15 | 14 | 8 | 3 |
- By Simpson's rule, the area of the cross-section of the river is
- (a) 705 sq. feet (b) 690 sq. feet (c) 710 sq. feet (d) 715 sq. feet
77. A curve passes through the points given by the following table
- | | | | | | | | |
|-------|---|-----|-----|-----|---|-----|-----|
| x : | 1 | 1.5 | 2 | 2.5 | 3 | 3.5 | 4 |
| y : | 2 | 2.4 | 2.7 | 2.8 | 3 | 2.6 | 2.1 |
- By Simpson's rule, the area bounded by the curve, the x -axis and the lines $x = 1, x = 4$, is
- (a) 7.583 (b) 6.783
- (c) 7.783 (d) 7.275
78. Using Simpson's $\frac{1}{3}$ rule, the value of $\int_1^3 f(x) dx$ for the following data, is
- | | | | | | |
|----------|-----|-----|-----|-----|---|
| x : | 1 | 1.5 | 2 | 2.5 | 3 |
| $f(x)$: | 2.1 | 2.4 | 2.2 | 2.8 | 3 |



[MP PET 1993]

- (a) 55.5 (b) 11.1 (c) 5.05 (d) 4.975

79. By the application of Simpson's one-third rule for numerical integration, with two subintervals, the value of $\int_0^1 \frac{dx}{1+x}$ is [MP PET 1996]

- (a) $\frac{17}{24}$ (b) $\frac{17}{36}$ (c) $\frac{25}{35}$ (d) $\frac{17}{25}$

80. By Simpson's rule, the value of $\int_{-3}^3 x^4 dx$ by taking 6 sub-intervals, is

- (a) 98 (b) 96 (c) 100 (d) 99

81. If $\int_a^b f(x) dx$ is numerically integrated by Simpson's rule, then in any pair of consecutive sub-intervals by which of the following curves, the curve $y = f(x)$ is approximated [MP PET 1998]

- (a) Straight line (b) Parabola (c) Circle (d) Ellipse

82. If by Simpson's rule $\int_0^1 \frac{1}{1+x^2} dx = \frac{1}{12} [3.1 + 4(a+b)]$ when the interval $[0, 1]$ is divided into 4 sub-intervals and a and b are the values of $\frac{1}{1+x^2}$ at two of its division points, then the values of a and b are the following [MP PET 1998]

- (a) $a = \frac{1}{1.0625}, b = \frac{1}{1.25}$ (b) $a = \frac{1}{1.0625}, b = \frac{1}{1.5625}$ (c) $a = \frac{1}{1.25}, b = 1$ (d) $a = \frac{1}{1.5625}, b = \frac{1}{1.25}$

83. If $e^0 = 1, e^1 = 2.72, e^2 = 7.39, e^3 = 20.09$ and $e^4 = 54.60$, then by Simpson's rule, the value of $\int_0^4 e^x dx$ is

[MP PET 1994, 95, 2001, 02]

- (a) 5.387 (b) 53.87 (c) 52.78 (d) 53.17

84. If $(2, 6)$ is divided into four intervals of equal region, then the approximate value of $\int_2^6 \frac{1}{x^2-x} dx$ using Simpson's rule, is

[EAMCET 2002]

- (a) 0.3222 (b) 0.2333 (c) 0.5222 (d) 0.2555

85. If $h = 1$ in Simpson's rule, the value of $\int_1^5 \frac{dx}{x}$ is

- (a) 1.62 (b) 1.43 (c) 1.48 (d) 1.56

* * *





Answer Sheet

Numerical Methods

Assignment (Basic and Advance Level)

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
c	c	b	a	b	c	d	b	a	c	b	c	b	d	a	d	c	c	b	a
21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40
c	a	d	a	c	a	c	b	a	d	d	b	a	d	c	d	b	b	a	a
41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60
b	c	b	b	c	a	c	b	d	b	a	b	c	d	a	d	b	a	c	a
61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80
c	b	a	b	d	b	a	b	b	b	b	a	c	a	b	c	c	c	c	a
81	82	83	84	85															
b	b	b	c	a															

