

Topics : Method of Differentiation, Complex Number, Continuity & Derivability,
Application of Derivatives, Sequence & Series, Straight Line

Type of Questions		M.M., Min.
Single choice Objective (no negative marking) Q.1,2,3,4,5	(3 marks, 3 min.)	[15, 15]
Subjective Questions (no negative marking) Q.6,7,8	(4 marks, 5 min.)	[12, 15]

- Sum to infinite terms of the series $\frac{1}{1.3} + \frac{2}{1.3.5} + \frac{3}{1.3.5.7} + \frac{4}{1.3.5.7.9} + \dots$ is
 (A) 1 (B) $\frac{1}{2}$ (C) $\frac{3}{2}$ (D) none of these
- Consider the function $f(x) = x - |x - x^2|$, $-1 \leq x \leq 2$. Then point of discontinuities of $f(x)$ for $x \in [-1, 2]$ are
 (A) $x = 0, 1$ (B) $x = 1, 2$ (C) $x = 0, \frac{1}{2}, 1$ (D) None of these
- Given that f is a real valued differentiable function such that $f(x) f'(x) < 0$ for all real x , it follows that
 (A) $f(x)$ is an increasing function (B) $f(x)$ is a decreasing function
 (C) $|f(x)|$ is an increasing function (D) $|f(x)|$ is a decreasing function
- If $f'(1) = -2\sqrt{2}$ and $g'(\sqrt{2}) = 4$, then the derivative of $f(\tan x)$ with respect to $g(\sec x)$ at $x = \frac{\pi}{4}$, is
 (A) 1 (B) -1 (C) 2 (D) 4
- If $y = (\sqrt{x})^{x^{\dots \dots \dots}}$, then $\frac{dy}{dx}$ is equal to
 (A) $\frac{y^3}{2x(1-y^2 \ln x)}$ (B) $\frac{y^2}{2x(1+y^2 \ln x)}$ (C) $\frac{y^2}{2x(1-y^2 \ln x)}$ (D) $\frac{y^3}{2x(1+y^2 \ln x)}$
- If $Y = \left(\frac{ax+b}{x^2+c}\right)$, then show that $(2xy' + y) y'' = 3(xy'' + y')y'$, where a, b, c are constants
- If the lines $L_1 : 2x - 3y - 6 = 0$, $L_2 : x + y - 4 = 0$ and $L_3 : x + 2 = 0$ taken pair wise in order constitute the angles A, B and C respectively of $\triangle ABC$, then find the equation whose roots are $\tan A$, $\tan B$ and $\tan C$.
- Sketch the region given by $|z| \leq 4$ & $\text{Arg}(z - i - 1) > \pi/4$



Answers Key

1. (B) 2. (D) 3. (D) 4. (B)
5. (A) 7. $2x^3 - 15x^2 + 28x - 15 = 0$

