

Topic : Vector

Type of Questions

M.M. Min.

Single choice Objective ('-1' negative marking) Q.1,2,3,4,5,6,7,8,9 (3 marks, 3 min.) [27, 27]

Subjective Questions (no negative marking) Q.10 (4 marks, 5 min.) [4, 5]

- If  $\vec{a} + 2\vec{b} + 3\vec{c} = 0$  and  $|\vec{a}| = 6$ ,  $|\vec{b}| = 3$  and  $|\vec{c}| = 2$ , then angle between  $\vec{a}$  and  $\vec{b}$  is

(A)  $\pi + \cos^{-1}\left(\frac{3}{4}\right)$       (B)  $\sin^{-1}\left(\frac{4}{5}\right)$       (C)  $\pi - \cos^{-1}\left(\frac{3}{4}\right)$       (D) None of these
- The value of  $\lambda$  for which the vector  $\vec{r} = (\lambda^2 - 9)\hat{i} + 2\hat{j} - (\lambda^2 - 16)\hat{k}$  makes acute angle with the positive direction of coordinate axis.

(A)  $(-\infty, -3) \cup (3, \infty)$       (B)  $(4, 4)$       (C)  $(-4, -3) \cup (3, 4)$       (D) None of these
- The set of all values of  $\lambda$  for which the vectors  $\vec{a} = (\lambda \log_2 x)\hat{i} - 6\hat{j} + 3\hat{k}$  and  $\vec{b} = (\log_2 x)\hat{i} + 2\hat{j} + (2\lambda \log_2 x)\hat{k}$  make an obtuse angle for any  $x \in (0, \infty)$

(A)  $\left(0, \frac{4}{3}\right)$       (B)  $\left(-\frac{4}{3}, 0\right)$       (C)  $\left(\frac{4}{3}, \infty\right)$       (D)  $\left(-\frac{4}{3}, 0\right]$
- $(\vec{r} \cdot \hat{i})(\hat{i} \times \vec{r}) + (\vec{r} \cdot \hat{j})(\hat{j} \times \vec{r}) + (\vec{r} \cdot \hat{k})(\hat{k} \times \vec{r}) =$

(A) 0      (B)  $\vec{r}$       (C)  $2\vec{r}$       (D)  $3\vec{r}$
- If  $|\vec{a}| = 3$ ,  $|\vec{b}| = 4$ , then a value of  $\lambda$  for which  $\vec{a} + \lambda\vec{b}$  is perpendicular to  $\vec{a} - \lambda\vec{b}$ , is :-

(A)  $\frac{9}{16}$       (B)  $\frac{3}{4}$       (C)  $\frac{3}{2}$       (D)  $\frac{4}{3}$
- If ABCDEF is regular hexagon, then  $\vec{AD} + \vec{EB} + \vec{FC}$  is equal to

(A) 0      (B)  $2\vec{AB}$       (C)  $3\vec{AB}$       (D)  $4\vec{AB}$



7. If  $\vec{a} = \hat{i} + \hat{j}$ ,  $\vec{b} = 2\hat{j} - \hat{k}$  and  $\vec{r} \times \vec{a} = \vec{b} \times \vec{a}$ ,  $\vec{r} \times \vec{b} = \vec{a} \times \vec{b}$ , then  $\frac{\vec{r}}{|\vec{r}|}$  is equal to
- (A)  $\frac{1}{\sqrt{11}}(\hat{i} + 3\hat{j} - \hat{k})$       (B)  $\frac{1}{\sqrt{11}}(\hat{i} - 3\hat{j} + \hat{k})$       (C)  $\frac{1}{\sqrt{3}}(\hat{i} - \hat{j} + \hat{k})$       (D) none of these
8. If  $\vec{a}$ ,  $\vec{b}$  are nonzero and noncollinear vectors, then  $[\vec{a} \ \vec{b} \ \vec{i}] \vec{i} + [\vec{a} \ \vec{b} \ \vec{j}] \vec{j} + [\vec{a} \ \vec{b} \ \vec{k}] \vec{k} =$
- (A)  $\vec{a} + \vec{b}$       (B)  $\vec{a} \times \vec{b}$       (C)  $\vec{a} - \vec{b}$       (D)  $\vec{b} \times \vec{a}$
9. A vector  $\vec{c}$  of magnitude  $20\sqrt{6}$  parallel to the bisector of the angle between  $\vec{a} = 7\hat{i} - 4\hat{j} - 4\hat{k}$  and  $\vec{b} = -2\hat{i} - \hat{j} + 2\hat{k}$  is
- (A)  $\pm \frac{20}{3}(2\hat{i} + 7\hat{j} + \hat{k})$       (B)  $\pm \frac{3}{20}(\hat{i} + 7\hat{j} + 2\hat{k})$
- (C)  $\pm \frac{20}{3}(\hat{i} - 2\hat{j} + 7\hat{k})$       (D)  $\pm \frac{20}{3}(\hat{i} - 7\hat{j} + 2\hat{k})$
10. In a triangle OAB,  $\angle AOB = 90^\circ$  where O is origin. If P and Q are point of trisection of AB then prove that  $OP^2 + OQ^2 = \frac{5}{9} AB^2$

## Answers Key

1. (D)      2. (C)      3. (D)      4. (A)
5. (B)      6. (D)      7. (A)      8. (B)      9. (D)

