



Series : A4BAB/3

SET-2

प्रश्न-पत्र कोड
Q.P. Code

55/3/2

रोल नं.

Roll No.

--	--	--	--	--	--	--	--

परीक्षार्थी प्रश्न-पत्र कोड को उत्तर-पुस्तिका के मुख-पृष्ठ पर अवश्य लिखें।

Candidates must write the Q.P. Code on the title page of the answer-book.

नोट

- (I) कृपया जाँच कर लें कि इस प्रश्न-पत्र में मुद्रित पृष्ठ 12 हैं।
- (II) प्रश्न-पत्र में दाहिने हाथ की ओर दिए गए प्रश्न-पत्र कोड को छात्र उत्तर-पुस्तिका के मुख-पृष्ठ पर लिखें।
- (III) कृपया जाँच कर लें कि इस प्रश्न-पत्र में 12 प्रश्न हैं।
- (IV) कृपया प्रश्न का उत्तर लिखना शुरू करने से पहले, प्रश्न का क्रमांक अवश्य लिखें।
- (V) इस प्रश्न-पत्र को पढ़ने के लिए 15 मिनट का समय दिया गया है। प्रश्न-पत्र का वितरण पूर्वाह्न में 10.15 बजे किया जाएगा। 10.15 बजे से 10.30 बजे तक छात्र केवल प्रश्न-पत्र को पढ़ेंगे और इस अवधि के दौरान वे उत्तर-पुस्तिका पर कोई उत्तर नहीं लिखेंगे।

NOTE

- (I) Please check that this question paper contains 12 printed pages.
- (II) Q.P. Code given on the right hand side of the question paper should be written on the title page of the answer-book by the candidate.
- (III) Please check that this question paper contains 12 questions.
- (IV) Please write down the Serial Number of the question in the answer-book before attempting it.
- (V) 15 minute time has been allotted to read this question paper. The question paper will be distributed at 10.15 a.m. From 10.15 a.m. to 10.30 a.m., the candidates will read the question paper only and will not write any answer on the answer-book during this period. *

भौतिक विज्ञान (सैद्धान्तिक)

PHYSICS (Theory)

निर्धारित समय : 2 घण्टे

Time allowed : 2 hours

अधिकतम अंक : 35

Maximum Marks : 35

55/3/2

257 B

1

P.T.O.





सामान्य निर्देश :

निम्नलिखित निर्देशों को बहुत सावधानी से पढ़िए और उनका पालन कीजिए :

- (i) इस प्रश्न-पत्र में कुल 12 प्रश्न हैं। सभी प्रश्न अनिवार्य हैं।
- (ii) यह प्रश्न पत्र तीन खंडों में विभाजित है – खंड क, ख और ग।
- (iii) खण्ड-क : प्रश्न संख्या 1 से 3 तक प्रत्येक प्रश्न 2 अंक का है।
- (iv) खण्ड-ख : प्रश्न संख्या 4 से 11 तक प्रत्येक प्रश्न 3 अंक का है।
- (v) खण्ड-ग : प्रश्न संख्या 12 प्रकरण अध्ययन आधारित प्रश्न है, यह प्रश्न 5 अंक का है।
- (vi) प्रश्न पत्र में कोई समग्र विकल्प नहीं है। हालाँकि कुछ प्रश्नों में आंतरिक विकल्प प्रदान किए गए हैं। इनमें से केवल एक ही प्रश्न का उत्तर लिखिए।
- (vii) लॉग टेबल का उपयोग कर सकते हैं, यदि आवश्यक हो, लेकिन कैल्कुलेटर के उपयोग की अनुमति नहीं है।

$$c = 3 \times 10^8 \text{ m/s}$$

$$h = 6.63 \times 10^{-34} \text{ Js}$$

$$e = 1.6 \times 10^{-19} \text{ C}$$

$$\mu_0 = 4\pi \times 10^{-7} \text{ T m A}^{-1}$$

$$\epsilon_0 = 8.854 \times 10^{-12} \text{ C}^2 \text{ N}^{-1} \text{ m}^{-2}$$

$$\frac{1}{4\pi\epsilon_0} = 9 \times 10^9 \text{ N m}^2 \text{ C}^{-2}$$

$$\text{इलेक्ट्रॉन का द्रव्यमान (m}_e\text{)} = 9.1 \times 10^{-31} \text{ kg}$$

$$\text{न्यूट्रॉन का द्रव्यमान} = 1.675 \times 10^{-27} \text{ kg}$$

$$\text{प्रोटॉन का द्रव्यमान} = 1.673 \times 10^{-27} \text{ kg}$$

$$\text{आवोगाद्रो संख्या} = 6.023 \times 10^{23} \text{ प्रति ग्राम मोल}$$

$$\text{बोल्जमान नियतांक} = 1.38 \times 10^{-23} \text{ JK}^{-1}$$

.55/3/2

2





General Instructions :

Read the following instructions very carefully and strictly follow them :

- (i) *This question paper contains 12 questions. All questions are compulsory.*
- (ii) *This question paper is divided into three sections – Section A, B and C.*
- (iii) **Section A :** *Q. Nos. 1 to 3 are of 2 marks each.*
- (iv) **Section B :** *Q. Nos. 4 to 11 are of 3 marks each.*
- (v) **Section C :** *Q. No. 12 is a case study based questions of 5 marks.*
- (vi) *There is no overall choice in the question paper. However, internal choice has been provided in some of the questions. Attempt any one of the alternatives in such questions.*
- (vii) *Use of log tables is permitted, if necessary, but use of calculator is not permitted.*

$$c = 3 \times 10^8 \text{ m/s}$$

$$h = 6.63 \times 10^{-34} \text{ Js}$$

$$e = 1.6 \times 10^{-19} \text{ C}$$

$$\mu_0 = 4\pi \times 10^{-7} \text{ T m A}^{-1}$$

$$\epsilon_0 = 8.854 \times 10^{-12} \text{ C}^2 \text{ N}^{-1} \text{ m}^{-2}$$

$$\frac{1}{4\pi\epsilon_0} = 9 \times 10^9 \text{ N m}^2 \text{ C}^{-2}$$

$$\text{Mass of electron (} m_e \text{)} = 9.1 \times 10^{-31} \text{ kg}$$

$$\text{Mass of neutron} = 1.675 \times 10^{-27} \text{ kg}$$

$$\text{Mass of proton} = 1.673 \times 10^{-27} \text{ kg}$$

$$\text{Avogadro's number} = 6.023 \times 10^{23} \text{ per gram mole}$$

$$\text{Boltzmann constant} = 1.38 \times 10^{-23} \text{ JK}^{-1}$$



खण्ड – क

1. किसी p-n संधि में रोधिका विभव बनने की व्याख्या कीजिए। 2
2. (a) किसी प्रकाशित फोटोडायोड का परिपथ आरेख और उसका I-V अभिलाक्षणिक खींचिए। 1
(b) किसी फोटोडायोड का उपयोग प्रकाश की तीव्रता के मापन में किस प्रकार किया जा सकता है ? 1
3. (a) (i) समस्थानिकों और समभारिकों के बीच विभेदन कीजिए। 2
(ii) दो नाभिकों के विभिन्न परमाणु द्रव्यमान A_1 और A_2 हैं। क्या ये नाभिक आवश्यक रूप से समान तत्व के समस्थानिक हैं ? व्याख्या कीजिए।

अथवा

- (b) (i) उन कारकों का नाम लिखिए जिन पर किसी पृष्ठ से प्रकाश-विद्युत उत्सर्जन निर्भर करता है।
(ii) किसी प्रकाश सुग्राही पदार्थ के लिए देहली आवृत्ति की परिभाषा लिखिए।

खण्ड – ख

4. (a) किसी फोटॉन, जिसकी ऊर्जा $6.5 \times 10^{-19} \text{ J}$ है, की आवृत्ति परिकलित कीजिए।
(b) क्या यह फोटॉन 2.14 eV कार्यफलन के Cs के पृष्ठ से इलेक्ट्रॉन का उत्सर्जन कर सकता है ? यदि हाँ, तो फोटोइलेक्ट्रॉन की अधिकतम गतिज ऊर्जा क्या होगी ? 3
5. 600 nm का कोई एकवर्णी प्रकाश वायु से जल के पृष्ठ पर आपतन करता है। जल का अपवर्तनांक 1.33 है। परावर्तित और अपवर्तित प्रकाश की (i) तरंगदैर्घ्य (ii) आवृत्ति और (iii) चाल ज्ञात कीजिए। 3
6. (a) λ_1 , λ_2 और λ_3 तरंगदैर्घ्य की विद्युत-चुम्बकीय तरंगों का उपयोग क्रमशः रेडार निकायों में, जल शोधकों में और टीवी के सुदूर स्विचों में किया जाता है।
(i) इन विद्युत-चुम्बकीय तरंगों को पहचानिए, तथा
(ii) इनमें प्रत्येक के लिए एक-एक स्रोत लिखिए। 3

अथवा

- (b) (i) दो प्रकाश स्रोतों के कलासंबद्ध होने के लिए दो शर्तों का उल्लेख कीजिए।
(ii) दो झिर्रियों के कारण उत्पन्न व्यतिकरण पैटर्न और एकल झिर्री के कारण विवर्तन पैटर्न के बीच दो अन्तर लिखिए।





SECTION – A

1. Explain the formation of the barrier potential in a p-n junction. 2
2. (a) Draw the circuit diagram of an illuminated photodiode and its I-V characteristics. 1
(b) How can a photodiode be used to measure the light intensity ? 1
3. (a) (i) Distinguish between isotopes and isobars. 2
(ii) Two nuclei have different mass numbers A_1 and A_2 . Are these nuclei necessarily the isotopes of the same element ? Explain.

OR

- (b) (i) Name the factors on which photoelectric emission from a surface depends.
(ii) Define the term 'threshold frequency' for a photosensitive material.

SECTION – B

4. (a) Calculate the frequency of a photon of energy 6.5×10^{-19} J.
(b) Can this photon cause emission of an electron from the surface of Cs of work function 2.14 eV ? If yes, what will be maximum kinetic energy of the photoelectron ? 3
5. Monochromatic light of wavelength 600 nm is incident from air on a water surface. The refractive index of water is 1.33. Find the (i) wavelength, (ii) frequency and (iii) speed, of reflected and refracted light. 3
6. (a) Electromagnetic waves of wavelengths λ_1 , λ_2 and λ_3 are used in radar systems, in water purifiers and in remote switches of TV, respectively.
(i) Identify the electromagnetic waves, and
(ii) Write one source of each of them. 3

OR

- (b) (i) State two conditions for two light sources to be coherent.
(ii) Give two points of difference between an interference pattern due to a double – slit and a diffraction pattern due to a single slit.

.55/3/2


5

P.T.O.





7. किसी हाइड्रोजन परमाणु में स्थायी कक्षाओं की व्याख्या के लिए बोर के अभिगृहीत का उल्लेख कीजिए। यह सत्यापित कीजिए कि n वीं कक्षा में परिक्रमा करने वाले किसी इलेक्ट्रॉन की चाल $(1/n)$ के आनुपातिक होती है। 3
8. चालकों, अर्धचालकों और विद्युतरोधियों के ऊर्जा बैंड आरेखित कीजिए। कौन सा बैंड ठोसों की वैद्युत चालकता निर्धारित करता है? ताप में वृद्धि होने पर किसी अर्धचालक की वैद्युत चालकता किस प्रकार प्रभावित होती है? व्याख्या कीजिए। 3
9. कोई प्रोटॉनों का पतला प्रकाश पुंज जिसमें प्रत्येक प्रोटॉन की ऊर्जा 4.1 MeV है लैड ($Z = 82$) की किसी चादर की ओर उपगमन कर रहा है। परिकल्पित कीजिए :
(i) पुंज में किसी प्रोटॉन की चाल, और
(ii) उसके उपगमन की समीपस्थ दूरी 3
10. द्वि-झिरी प्रयोग में फ्रिंजों के पृथक्करण पर क्या प्रभाव पड़ेगा यदि :
(i) झिरीयों के पृथक्करण में वृद्धि कर दी जाए,
(ii) उपयोग किए गए प्रकाश का वर्ण लाल से नीले में परिवर्तित कर दिया जाए,
(iii) सारे उपकरण को अपवर्तनांक 1.2 के किसी तेल में डुबो दिया जाए ?
प्रत्येक प्रकरण में अपने उत्तर की पुष्टि कीजिए। 3
11. (a) पूर्ण आन्तरिक परावर्तन के लिए दो आवश्यक शर्तें लिखिए।
(b) आरेख में दर्शाए अनुसार दो प्रिज्मों ABC और DBC को व्यवस्थित किया गया है। वायु के सापेक्ष इन दोनों प्रिज्मों के क्रांतिक कोण क्रमशः 41.1° और 45° हैं। प्रिज्मों के संयोजन से प्रकाश किरण के पथ को आरेखित कीजिए। 3




अथवा

6

.55/3/2



7. State Bohr's postulate to explain stable orbits in a hydrogen atom. Prove that the speed with which the electron revolves in n^{th} orbit is proportional to $(1/n)$. 3
8. Draw the energy band diagrams for conductors, semiconductors and insulators. Which band determines the electrical conductivity of a solid? How is the electrical conductivity of a semiconductor affected with rise in its temperature? Explain. 3
9. A narrow beam of protons, each having 4.1 MeV energy is approaching a sheet of lead ($Z = 82$). Calculate :
(i) the speed of a proton in the beam, and
(ii) the distance of its closest approach 3
10. How is the spacing between fringes in a double slit experiment affected if :
(i) the slits separation is increased,
(ii) the colour of light used is changed from red to blue,
(iii) the whole apparatus is submerged in a oil of refractive index 1.2 ?
Justify your answer in each case. 3
11. (a) Write two necessary conditions for total internal reflection.
(b) Two prisms ABC and DBC are arranged as shown in figure.



The critical angles for the two prisms with respect to air are 41.1° and 45° respectively. Trace the path of the ray through the combination. 3

OR


.55/3/2

7

P.T.O.



- (a) कोई बिम्ब किसी अभिसारी लेंस के सामने स्थित है। वह शर्ते प्राप्त कीजिए जिसमें इस लेंस द्वारा उत्पन्न आवर्धन (i) ऋणात्मक और (ii) धनात्मक होता है।
- (b) कोई बिन्दुकित बिम्ब किसी काँच के गोले के सामने आरेख में दर्शाए अनुसार O पर स्थित है। इस गोले द्वारा प्रतिबिम्ब बनना दर्शाइए।



खण्ड – ग

प्रकरण अध्ययन

12. संयुक्त सूक्ष्मदर्शी दो अभिसारी लेंसों से मिलकर बनता है। जिनमें एक लेंस जिसका द्वारक छोटा और फोकस दूरी कम होती है उसे अभिदृश्यक कहते हैं तथा दूसरे लेंस को जिसका द्वारक कुछ बड़ा होता है और फोकस दूरी भी कुछ अधिक होती है उसे नेत्रिका कहते हैं। दोनों लेंसों को किसी नलिका में इस प्रकार व्यवस्थित किया जाता है कि इन दोनों लेंसों के बीच की दूरी को परिवर्तित किया जा सके। किसी लघु बिम्ब को अभिदृश्यक के सामने इसकी फोकस दूरी से कुछ अधिक दूरी पर रखा जाता है। अभिदृश्यक इस बिम्ब का प्रतिबिम्ब बनाता है, जो नेत्रिका के लिए बिम्ब की भांति कार्य करता है। नेत्रिका फिर बिम्ब का अंतिम आवर्धित प्रतिबिम्ब बना देती हैं।

$$1 \times 5 = 5$$

- I. किसी संयुक्त सूक्ष्मदर्शी में अभिदृश्यक और नेत्रिका द्वारा बनाए गए प्रतिबिम्ब होते हैं क्रमशः


- (A) आभासी, वास्तविक (B) वास्तविक, आभासी
(C) आभासी, आभासी (D) वास्तविक, वास्तविक

- II. किसी संयुक्त सूक्ष्मदर्शी के कारण आवर्धन निम्नलिखित में से किस पर निर्भर नहीं करता है ?

- (A) अभिदृश्यक और नेत्रिका के द्वारक
(B) अभिदृश्यक और नेत्रिका की फोकस दूरी
(C) नलिका की लम्बाई
(D) उपयोग किया गया प्रकाश



- (a) An object is placed in front of a converging lens. Obtain the conditions under which the magnification produced by the lens is (i) negative and (ii) positive.
- (b) A point object is placed at O in front of a glass sphere as shown in figure.



Show the formation of image by the sphere.

SECTION – C

CASE STUDY

12. A compound microscope consists of two converging lenses. One of them, of smaller aperture and smaller focal length is called objective and the other of slightly larger aperture and slightly larger focal length is called eye-piece. Both the lenses are fitted in a tube with an arrangement to vary the distance between them. A tiny object is placed in front of the objective at a distance slightly greater than its focal length. The objective produces the image of the object which acts as an object for the eye-piece. The eye piece, in turn produces the final magnified image. $1 \times 5 = 5$

- I. In a compound microscope the images formed by the objective and the eye-piece are respectively
- (A) virtual, real (B) real, virtual
(C) virtual, virtual (D) real, real
- II. The magnification due to a compound microscope *does not* depend upon
- (A) the aperture of the objective and the eye-piece
(B) the focal length of the objective and the eye-piece
(C) the length of the tube
(D) the colour of the light used

.55/3/2

9

P.T.O.





- III. संयुक्त सूक्ष्मदर्शी के संदर्भ में कौन सा कथन *सही नहीं* है ?
- (A) दोनों लेंस कम फोकस दूरी के होते हैं ।
(B) दोनों लेंसों की फोकस दूरी कम करने पर आवर्धन क्षमता बढ़ जाती है ।
(C) दोनों लेंसों के बीच की फोकस दूरी ($f_o + f_e$) से अधिक होती है ।
(D) दोनों लेंसों की अदला-बदली करके इस सूक्ष्मदर्शी का उपयोग दूरदर्शक के रूप में किया जा सकता है ।
- IV. किसी संयुक्त सूक्ष्मदर्शी में अभिदृश्यक 10X का और नेत्रिका 20X की है । इस सूक्ष्मदर्शी के कारण आवर्धन होगा
- (A) 2 (B) 10
(C) 30 (D) 200
- V. किसी संयुक्त सूक्ष्मदर्शी के अभिदृश्यक और नेत्रिका की फोकस दूरियाँ क्रमशः 1.2 cm और 3.0 cm हैं । बिम्ब अभिदृश्यक से 1.25 cm दूरी पर स्थित है । यदि अन्तिम प्रतिबिम्ब अनन्त पर बनता है, तो सूक्ष्मदर्शी की आवर्धन क्षमता होगी
- (A) 100 (B) 150
(C) 200 (D) 250
-





- III. Which of the following is *not correct* in the context of a compound microscope ?
- (A) Both the lenses are of short focal lengths.
 - (B) The magnifying power increases by decreasing the focal lengths of the two lenses.
 - (C) The distance between the two lenses is more than $(f_o + f_e)$.
 - (D) The microscope can be used as a telescope by interchanging the two lenses.
- IV. A compound microscope consists of an objective of 10X and an eye-piece of 20X. The magnification due to the microscope would be
- (A) 2
 - (B) 10
 - (C) 30
 - (D) 200
- V. The focal lengths of objective and eye-piece of a compound microscope are 1.2 cm and 3.0 cm respectively. The object is placed at a distance of 1.25 cm from the objective. If the final image is formed at infinity, the magnifying power of the microscope would be
- (A) 100
 - (B) 150
 - (C) 200
 - (D) 250
-





*

.55/3/2

257 B

12

Strictly Confidential: (For Internal and Restricted use only)

SeniorSecondary School ,Term II Examination2022

Marking Scheme – PHYSICS (SUBJECT CODE — 042)

(PAPER CODE — 55/3/2)

General Instructions: -

1. You are aware that evaluation is the most important process in the actual and correct assessment of the candidates. A small mistake in evaluation may lead to serious problems which may affect the future of the candidates, education system and teaching profession. To avoid mistakes, it is requested that before starting evaluation, you must read and understand the spot evaluation guidelines carefully.
2. **“Evaluation policy is a confidential policy as it is related to the confidentiality of the examinations conducted, Evaluation done and several other aspects. Its’ leakage to public in any manner could lead to derailment of the examination system and affect the life and future of millions of candidates. Sharing this policy/document to anyone, publishing in any magazine and printing in News Paper/Website etc may invite action under IPC.”**
3. Evaluation is to be done as per instructions provided in the Marking Scheme. It should not be done according to one’s own interpretation or any other consideration. Marking Scheme should be strictly adhered to and religiously followed. **However, while evaluating, answers which are based on latest information or knowledge and/or are innovative, they may be assessed for their correctness otherwise and marks be awarded to them. In class-X, while evaluating two competency based questions, please try to understand given answer and even if reply is not from marking scheme but correct competency is enumerated by the candidate, marks should be awarded.**
4. The Head-Examiner must go through the first five answer books evaluated by each evaluator on the first day, to ensure that evaluation has been carried out as per the instructions given in the Marking Scheme. The remaining answer books meant for evaluation shall be given only after ensuring that there is no significant variation in the marking of individual evaluators.
5. Evaluators will mark(\surd) wherever answer is correct. For wrong answer ‘X’ be marked. Evaluators will not put right kind of mark while evaluating which gives an impression that answer is correct and no marks are awarded. **This is most common mistake which evaluators are committing.**
6. If a question has parts, please award marks on the right-hand side for each part. Marks awarded for different parts of the question should then be totalled up and written in the left-hand margin and encircled. This may be followed strictly.
7. If a question does not have any parts, marks must be awarded in the left-hand margin and encircled. This may also be followed strictly.
8. If a student has attempted an extra question, answer of the question deserving more marks should be retained and the other answer scored out.
9. No marks to be deducted for the cumulative effect of an error. It should be penalized only once.



10. A full scale of marks ____35 ____ (example 0-40 marks as given in Question Paper) has to be used. Please do not hesitate to award full marks if the answer deserves it.
11. Every examiner has to necessarily do evaluation work for full working hours i.e. 8 hours every day and evaluate 30 answer books per day in main subjects and 35 answer books per day in other subjects (Details are given in Spot Guidelines). This is in view of the .of questions in question paper reduced syllabus and number
12. Ensure that you do not make the following common types of errors committed by the Examiner in the past:-
 - Leaving answer or part thereof un assessed in an answer book.
 - Giving more marks for an answer than assigned to it.
 - Wrong totalling of marks awarded on a reply.
 - Wrong transfer of marks from the inside pages of the answer book to the title page.
 - Wrong question wise totalling on the title page.
 - Wrong totalling of marks of the two columns on the title page.
 - Wrong grand total.
 - Marks in words and figures not tallying.
 - Wrong transfer of marks from the answer book to online award list.
 - Answers marked as correct, but marks not awarded. (Ensure that the right tick mark is correctly and clearly indicated. It should merely be a line. Same is with the X for incorrect answer.)
 - Half or a part of answer marked correct and the rest as wrong, but no marks awarded.
13. While evaluating the answer books if the answer is found to be totally incorrect, it should be marked as cross (X) and awarded zero (0)Marks.
14. Any un assessed portion, non-carrying over of marks to the title page, or totalling error detected by the candidate shall damage the prestige of all the personnel engaged in the evaluation work as also of the Board. Hence, in order to uphold the prestige of all concerned, it is again reiterated that the instructions be followed meticulously and judiciously.
15. The Examiners should acquaint themselves with the guidelines given in the Guidelines for spot Evaluation before starting the actual evaluation.
16. Every Examiner shall also ensure that all the answers are evaluated, marks carried over to the title page, correctly totalled and written in figures and words.
17. The Board permits candidates to obtain photocopy of the Answer Book on request in an RTI application and also separately as a part of the re-evaluation process on payment of the processing charges.





MARKING SCHEME

Senior Secondary School Examination TERM–II, 2022

PHYSICS (Subject Code — 042)

[Paper Code — 55/3/2]

Maximum Marks : 35


Q. No.	EXPECTED ANSWER / VALUE POINTS	Marks	Total Marks				
SECTION—A							
1.	<table border="1"> <tr> <td>Explanation of movement of charge carriers / diffusion</td> <td>1</td> </tr> <tr> <td>Formation of the barrier potential</td> <td>1</td> </tr> </table> <p>The diffusion of electrons from n-region to p-region and that of the holes from p-region to n-region creates positive charge on the n-side and negative charge on the p-side which causes a difference of potential across the junction.</p> <p>This potential, setup across the junction tends to prevent the movement of electrons from the n-region to p-region .This is called barrier potential.</p>	Explanation of movement of charge carriers / diffusion	1	Formation of the barrier potential	1	1 1	2
Explanation of movement of charge carriers / diffusion	1						
Formation of the barrier potential	1						
2.	<table border="1"> <tr> <td>(a) Circuit diagram and I- V characteristic</td> <td>1</td> </tr> <tr> <td>(b) To measure the light intensity using photodiode</td> <td>1</td> </tr> </table> <p>(a)</p>  <p>(b)</p>  <p>(b) Photo current flows in an illuminated photodiode under reverse bias. The magnitude of photocurrent varies linearly as the intensity of incident light. This fact is used to measure the light intensity.</p>	(a) Circuit diagram and I - V characteristic	1	(b) To measure the light intensity using photodiode	1	1/2 1/2	2
(a) Circuit diagram and I - V characteristic	1						
(b) To measure the light intensity using photodiode	1						
3.	<table border="1"> <tr> <td>Distinction between isotopes and isobars</td> <td>1</td> </tr> <tr> <td>Explanation</td> <td>1</td> </tr> </table>	Distinction between isotopes and isobars	1	Explanation	1		
Distinction between isotopes and isobars	1						
Explanation	1						




	<p>(i) Isotopes – These are the atoms having same atomic number (Z) but different atomic mass(A). Isobars – The atoms of different element having same atomic masses.</p> <p>(ii) No The mass number of a nucleus is the sum of number of proton(Z) and number of neutrons (N) / $A = Z + N$ / Two nuclei with different mass numbers A_1 and A_2, may have, have different Z .</p> <p style="text-align: center;">OR</p> <p>b) <table border="1" style="display: inline-table; vertical-align: middle;"> <tr> <td>Two factors</td> <td style="text-align: right;">1</td> </tr> <tr> <td>Definition of threshold frequency</td> <td style="text-align: right;">1</td> </tr> </table></p> <p>(i) Factors (a) Frequency of incident radiation (b) Work function of the surface</p> <p>(ii) The minimum frequency of the incident radiation below which photoelectric emission does not take place.</p>	Two factors	1	Definition of threshold frequency	1	<p>½</p> <p>½</p> <p>½</p> <p>½</p> <p>1</p>	<p>2</p> <p>2</p>
Two factors	1						
Definition of threshold frequency	1						
	SECTION B						
4.	<table border="1" style="display: inline-table; vertical-align: middle;"> <tr> <td>a) Calculation of frequency</td> <td style="text-align: right;">1</td> </tr> <tr> <td>b) Determination of energy for electron emission</td> <td style="text-align: right;">2</td> </tr> </table> <p>$E = h\nu$</p> $\nu = \frac{E}{h} = \frac{6.5 \times 10^{-19}}{6.63 \times 10^{-34}}$ $= 9.8 \times 10^{14} \text{ Hz}$ <p>b) Energy of photon in eV $E = 6.5 \times 10^{-19} \text{ J} = 4.06 \text{ eV}$ $\therefore E > \phi_0$ There will be photoelectric emission. K.E of photoelectron $E_k = E - \phi_0$ $= 4.06 - 2.14$ $= 1.92 \text{ eV}$</p>	a) Calculation of frequency	1	b) Determination of energy for electron emission	2	<p>½</p> <p>½</p> <p>½</p> <p>½</p> <p>½</p> <p>½</p>	<p>3</p>
a) Calculation of frequency	1						
b) Determination of energy for electron emission	2						
5.	<table border="1" style="display: inline-table; vertical-align: middle;"> <tr> <td>(i) Finding wavelength ,frequency and speed of reflected light</td> <td style="text-align: right;">1 ½</td> </tr> <tr> <td>(ii) Finding wavelength ,frequency and speed of refracted light</td> <td style="text-align: right;">1 ½</td> </tr> </table> <p>(a) Reflected high $\lambda' = \lambda = 600 \text{ nm.}$ <i>No change in wavelength</i> $c = 3 \times 10^8 \text{ m/s in air}$</p>	(i) Finding wavelength ,frequency and speed of reflected light	1 ½	(ii) Finding wavelength ,frequency and speed of refracted light	1 ½	<p>½</p> <p>½</p>	
(i) Finding wavelength ,frequency and speed of reflected light	1 ½						
(ii) Finding wavelength ,frequency and speed of refracted light	1 ½						



	$v = \frac{c}{\lambda} = \frac{3 \times 10^8}{5 \times 10^{14}} \text{ Hz}$ <p>Refracted light</p> $\lambda' = \frac{\lambda}{\mu} = \frac{600}{1.33} = 451.1 \text{ nm}$ $\approx 450 \text{ nm}$ $v = \frac{c}{\mu} = \frac{3 \times 10^8}{1.33}$ $= 2.26 \times 10^8 \text{ m/s}$ $= 5 \times 10^{14} \text{ Hz}$	<p>½</p> <p>½</p> <p>½</p> <p>½</p>	<p>3</p>																						
<p>6.</p>	<p>a)</p> <table border="1" style="width: 100%;"> <tr> <td>(i) Identification of the three em waves</td> <td>1 ½</td> </tr> <tr> <td>(ii) Sources of the three em waves</td> <td>1 ½</td> </tr> </table> <table style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 30%; text-align: center;">EM Waves</th> <th style="width: 70%; text-align: center;">Sources</th> </tr> </thead> <tbody> <tr> <td>$\lambda_1 \rightarrow$ Microwaves</td> <td>magnetron valve / klystron valve / gun diodes</td> </tr> <tr> <td>$\lambda_2 \rightarrow$ Ultra Violet</td> <td>high voltage gas discharge tube</td> </tr> <tr> <td>$\lambda_3 \rightarrow$ Infra red</td> <td>hot bodies and molecules</td> </tr> </tbody> </table> <p style="text-align: center;">OR</p> <p>b)</p> <table border="1" style="width: 100%;"> <tr> <td>(i) Two conditions</td> <td>1</td> </tr> <tr> <td>(ii) Two points of difference</td> <td>2</td> </tr> </table> <p>i) The two sources should emit the waves in the same phase or with a constant phase difference . The two sources must continuously emit light wave of the same wavelength.</p> <p>ii)</p> <table border="1" style="width: 100%;"> <thead> <tr> <th style="width: 50%; text-align: center;">Interference</th> <th style="width: 50%; text-align: center;">Diffraction</th> </tr> </thead> <tbody> <tr> <td>1) Intensity of all bright bands are nearly same</td> <td>1) Intensity of bright bands decreases rapidly on both sides of central maxima.</td> </tr> <tr> <td>2) All fringes are of same width.</td> <td>2) All the fringes are not of same width / the width of central maximum is double that of the width of secondary maxima.</td> </tr> </tbody> </table> <p>Or any other two differences.</p>	(i) Identification of the three em waves	1 ½	(ii) Sources of the three em waves	1 ½	EM Waves	Sources	$\lambda_1 \rightarrow$ Microwaves	magnetron valve / klystron valve / gun diodes	$\lambda_2 \rightarrow$ Ultra Violet	high voltage gas discharge tube	$\lambda_3 \rightarrow$ Infra red	hot bodies and molecules	(i) Two conditions	1	(ii) Two points of difference	2	Interference	Diffraction	1) Intensity of all bright bands are nearly same	1) Intensity of bright bands decreases rapidly on both sides of central maxima.	2) All fringes are of same width.	2) All the fringes are not of same width / the width of central maximum is double that of the width of secondary maxima.	<p>½+ ½</p> <p>½+ ½</p> <p>½+ ½</p> <p>½</p> <p>½</p> <p>1+1</p>	<p>3</p> <p>3</p>
(i) Identification of the three em waves	1 ½																								
(ii) Sources of the three em waves	1 ½																								
EM Waves	Sources																								
$\lambda_1 \rightarrow$ Microwaves	magnetron valve / klystron valve / gun diodes																								
$\lambda_2 \rightarrow$ Ultra Violet	high voltage gas discharge tube																								
$\lambda_3 \rightarrow$ Infra red	hot bodies and molecules																								
(i) Two conditions	1																								
(ii) Two points of difference	2																								
Interference	Diffraction																								
1) Intensity of all bright bands are nearly same	1) Intensity of bright bands decreases rapidly on both sides of central maxima.																								
2) All fringes are of same width.	2) All the fringes are not of same width / the width of central maximum is double that of the width of secondary maxima.																								

7.	<div style="border: 1px solid black; padding: 5px; margin-bottom: 10px;"> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="padding: 2px;">Statement of Bohr's 2nd postulate</td> <td style="text-align: right; padding: 2px;">1</td> </tr> <tr> <td style="padding: 2px;">Derivation of speed</td> <td style="text-align: right; padding: 2px;">2</td> </tr> </table> </div> <p>(i) An electron can revolve around the nucleus in an orbit in which its angular momentum is an integral multiple of $\frac{h}{2\pi}$.</p> <p>(ii) Proof</p> $\frac{mv^2}{r} = \frac{1}{4\pi\epsilon_0} \frac{e^2}{r^2}$ $mvr = \frac{nh}{2\pi}$ <p>Eliminating r we get</p> $v = \frac{e^2}{2\epsilon_0 h} \cdot \frac{1}{n}$ $\therefore v \propto \frac{1}{n}$	Statement of Bohr's 2 nd postulate	1	Derivation of speed	2	1 1/2 1/2 1/2 1/2	3		
Statement of Bohr's 2 nd postulate	1								
Derivation of speed	2								
8	<div style="border: 1px solid black; padding: 5px; margin-bottom: 10px;"> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="padding: 2px;">Energy band diagrams for three cases</td> <td style="text-align: right; padding: 2px;">1 1/2</td> </tr> <tr> <td style="padding: 2px;">Bandwidth determination for conductivity</td> <td style="text-align: right; padding: 2px;">1/2</td> </tr> <tr> <td style="padding: 2px;">Effect of temperature</td> <td style="text-align: right; padding: 2px;">1</td> </tr> </table> </div>  <p>Conduction band determines electrical conductivity</p> <p>As temperature of a semiconductor rises, the carrier concentration (electron-hole pair) increases due to breaking of covalent bonds and the conductivity of the semiconductor increases.</p>	Energy band diagrams for three cases	1 1/2	Bandwidth determination for conductivity	1/2	Effect of temperature	1	1/2 1/2 + 1/2 1/2 1	3
Energy band diagrams for three cases	1 1/2								
Bandwidth determination for conductivity	1/2								
Effect of temperature	1								



<p>9.</p>	<div style="border: 1px solid black; padding: 5px; margin-bottom: 10px;"> (i) Calculation of speed 1 ½ (ii) Calculation of the distance of closest approach 1 ½ </div> <p>(i) $\frac{1}{2}mv^2 = 4 \cdot 1 \times 1.6 \times 10^{-13} \text{ J}$</p> $v = \sqrt{\frac{2 \times 4 \cdot 1 \times 1.6 \times 10^{-13}}{1.673 \times 10^{-27}}}$ $= 2.8 \times 10^7 \text{ m/s}$ <p>(ii) $d = \frac{Ze^2}{4\pi\epsilon_0 \times E_k}$</p> $= \frac{9 \times 10^9 \times 82 \times 1.6 \times 10^{-19} \times 1.6 \times 10^{-19}}{4 \cdot 1 \times 1.6 \times 10^{-13}}$ $= 2.88 \times 10^{-14} \text{ m}$	<p>½</p> <p>½</p> <p>½</p> <hr style="width: 50%; margin-left: 0;"/> <p>½</p> <p>½</p> <p>½</p>	<p>3</p>
<p>10.</p>	<div style="border: 1px solid black; padding: 5px; margin-bottom: 10px;"> Effect on spacing between fringes 1 ½ Justification 1 ½ </div> <p>Spacing between fringes in Young's double slit experiment</p> $\beta = \frac{\lambda D}{d}$ <p>1) β decreases as d increases as $\beta \propto \frac{1}{d}$</p> <p>2) $\lambda_{\text{blue}} < \lambda_{\text{red}}$ as $\beta \propto \lambda \therefore$ fringe width decreases.</p> <p>3) $\beta' = \frac{\beta}{\mu} \therefore \beta$ decreases as $\mu > 1$ [$\therefore \mu = 1.2$]</p>	<p>½ + ½</p> <p>½ + ½</p> <p>½ + ½</p>	<p>3</p>
<p>11.</p>	<p>a) (a) Two necessary conditions 2 (b) Tracing the path of the ray 1</p> <p>a) Two conditions</p> <p>(i) The light must travel from an optically denser medium to a rarer medium.</p> <p>(ii) Angle of incidence should be greater than the critical angle.</p> <p>(b) </p>	<p>1</p> <p>1</p> <p>1</p>	<p>3</p>

<u>OR</u>							
b)	<table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>(a) Obtaining conditions for two cases</td> <td style="text-align: center;">2</td> </tr> <tr> <td>(b) Formation of image</td> <td style="text-align: center;">1</td> </tr> </table>	(a) Obtaining conditions for two cases	2	(b) Formation of image	1		
(a) Obtaining conditions for two cases	2						
(b) Formation of image	1						
(a)	Two conditions $m = \frac{h'}{h} = \frac{v}{u}$						
(i)	For real images / when object is placed beyond F, As u is negative and v is positive..	$\frac{1}{2} + \frac{1}{2}$					
(ii)	for virtual image / when object is kept between F and the optical centre of the lens. As u and v both are negative.	$\frac{1}{2} + \frac{1}{2}$					
		1					
			3				
SECTION—C							
12.	(I) (B) real , virtual	1					
	(II) (A) The aperture of the objective and the eye piece	1					
	(III) (D) The microscope can be used as a telescope by interchanging the two lenses.	1					
	(IV) (D) 200	1					
	(V) (C) 200	1	5				

* * *