

## Syllabus

*Vectors and Scalars, Magnitude and direction of a vector. Direction cosines and direction ratios of a vector, "types of vectors" equal, unit, zero, parallel and collinear vectors, position vector of a "point, negative of a vector", components of a vector, addition of vectors (properties of addition, laws of addition), Multiplication of a vector by a scalar, position vector of a point dividing a line segment in a given ratio. Definition, Geometrical interpretation, properties and application of Scalar (dot) product of vectors, Vector (cross) product of vectors, scalar triple product of vectors.*

## Chapter Analysis

TOPIC	2016		2017		2018
	Delhi	OD	Delhi	OD	Delhi/OD
Properties	2 Q. (1 Mark)	-	-	-	-
Angle between vectors	-	-	1 Q. (4 Marks)	-	2 Q. (1 Mark) 2 Q. (2 Marks)
Dot Product	-	-	-	-	-
Cross product	-	2 Q. (1 Mark)	-	-	1 Q. (4 Marks)
Area of triangle	-	1 Q. (4 Marks)	-	1 Q. (4 Marks)	-
Coplanarity	1 Q. (4 Marks)	-	1 Q. (4 Marks)	1 Q. (4 Marks)	-



### TOPIC-1 Basic Algebra of Vectors

TOPIC - 1 Page 366  
Basic Algebra of Vectors

TOPIC - 2 Page 378  
Dot Product of Vectors

TOPIC - 3 Page 389  
Cross Product

TOPIC - 4 Page 403  
Scalar Triple Product

## Revision Notes

### 1. Vector : Basic Introduction :

- A quantity having magnitude as well as the direction is called a vector. It is denoted as  $\vec{AB}$  or  $\vec{a}$ . Its magnitude (or modulus) is  $|\vec{AB}|$  or  $|\vec{a}|$  otherwise, simply  $AB$  or  $a$ .
- Vectors are denoted by symbols such as  $\vec{a}$ . [Pictorial representation of vector]

### 2. Initial and Terminal Points :

The initial and terminal points means that point from which the vector originates and terminates respectively.

**3. Position Vector :**

The position vector of a point say  $P(x, y, z)$  is  $\vec{OP} = \vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$  and the magnitude is  $r = \sqrt{x^2 + y^2 + z^2}$ .

The vector  $\vec{OP} = \vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$  is said to be in its **component form**. Here  $x, y, z$  are called the scalar components or rectangular components of  $\vec{r}$  and  $x\hat{i}, y\hat{j}, z\hat{k}$  are the vector components of  $\vec{r}$  along  $x, y, z$ -axis respectively.

- Also,  $\vec{r} = (\text{Position Vector of } B) - (\text{Position Vector of } A)$ . For example, let  $A(x_1, y_1, z_1)$  and  $B(x_2, y_2, z_2)$ . Then,  $\vec{r} = (x_2\hat{i} + y_2\hat{j} + z_2\hat{k}) - (x_1\hat{i} + y_1\hat{j} + z_1\hat{k})$ .
- Here  $\hat{i}, \hat{j}, \hat{k}$  are the unit vectors along the axes  $OX, OY$  and  $OZ$  respectively (The discussion about unit vectors is given later under 'types of vectors').

**4. Direction Ratios and Direction Cosines :**

If  $r = xi + yj + zk$ , then coefficient of  $i, j, k$  in  $\vec{r}$  i.e.,  $x, y, z$  are called the direction ratios (abbreviated as d.r.'s) of vector  $\vec{r}$ . These are denoted by  $a, b, c$  (i.e.,  $a = x, b = y, c = z$ ; in a manner we can say that scalar components of vector  $\vec{r}$  and its d.r.'s both are the same).

Also, the coefficients of  $i, j, k$  in  $\frac{\vec{r}}{r}$  (which is the unit vector of  $\vec{r}$ ) i.e.,  $\frac{x}{\sqrt{x^2 + y^2 + z^2}}, \frac{y}{\sqrt{x^2 + y^2 + z^2}}, \frac{z}{\sqrt{x^2 + y^2 + z^2}}$  are called direction cosines (which is abbreviated as d.c.'s) of vector  $\vec{r}$ .

- These direction cosines are denoted by  $l, m, n$  such that  $l = \cos \alpha, m = \cos \beta, n = \cos \gamma$  and  $l^2 + m^2 + n^2 = 1 \Rightarrow \cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$ .
- It can be easily concluded that  $\frac{x}{r} = l = \cos \alpha, \frac{y}{r} = m = \cos \beta, \frac{z}{r} = n = \cos \gamma$ .

Therefore,  $r = lri + mrj + nrk = r(\cos \alpha i + \cos \beta j + \cos \gamma k)$ . [Here  $r = |\vec{r}|$ ].

**5. Addition of vectors**

(a) **Triangular law** : If two adjacent sides (say sides  $AB$  and  $BC$ ) of a triangle  $ABC$  are represented by  $\vec{AB}$  and  $\vec{BC}$  taken in same order, then the third side of the triangle taken in the reverse order gives the sum of vectors  $\vec{AB} + \vec{BC} = \vec{AC}$  and  $\vec{AC} = -\vec{CA} \Rightarrow \vec{AB} + \vec{BC} = -\vec{CA}$ .

Also since  $\vec{AC} = -\vec{CA} \Rightarrow \vec{AB} + \vec{BC} = -\vec{CA}$ .

And  $\vec{AB} + \vec{BC} - \vec{AC} = \vec{AB} + \vec{BC} + \vec{CA} = \vec{0}$ .

(b) **Parallelogram law** : If two vectors  $\vec{OA}$  and  $\vec{OB}$  are represented in magnitude and the direction by the two adjacent sides (say  $AB$  and  $AD$ ) of a parallelogram  $ABCD$ , then their sum is given by that diagonal of parallelogram which is co-initial with  $\vec{OA}$  and  $\vec{OB}$  i.e.,  $\vec{OC} = \vec{OA} + \vec{OB}$ .

**6. Properties of Vector Addition**

(a) **Commutative property** :

Consider  $a = a_1i + a_2j + a_3k$  and  $b = b_1i + b_2j + b_3k$  be any two given vectors,

then  $a + b = (a_1 + b_1)i + (a_2 + b_2)j + (a_3 + b_3)k = b + a$ .

(b) **Associative property** :  $(a + b) + c = a + (b + c)$ .

(c) **Additive identity property** :

(d) **Additive inverse property** :  $a + (-a) = 0 = (-a) + a$ .

**Note : Multiplication of a vector by a scalar**

Let  $\vec{a}$  be any vector and  $k$  be any scalar. Then the product  $k\vec{a}$  is defined as a vector whose magnitude is  $|k|$  times that of  $\vec{a}$  and the direction is

- (i) same as that of  $\vec{a}$  if  $k$  is positive, and (ii) opposite as that of  $\vec{a}$  if  $k$  is negative.



## Know the Terms

### Types of Vectors :

- (a) **Zero or Null vector** : It is that vector whose initial and terminal points are coincident. It is denoted by  $\vec{0}$ . Ofcourse its magnitude is 0 (zero).
- Any non-zero vector is called a **proper vector**.
- (b) **Co-initial vectors** : Those vectors (two or more) having the same starting point are called the co-initial vectors.
- (c) **Co-terminus vectors** : Those vectors (two or more) having the same terminal point are called the co-terminus vectors.
- (d) **Negative of a vector** : The vector which has the same magnitude as the  $\vec{a}$  but opposite direction. It is denoted by  $-\vec{a}$ . Hence if,  $\vec{AB} = \vec{a}$  then  $-\vec{a} = \vec{BA}$  i.e.,  $\vec{AB} = -\vec{BA}$ ,  $PQ = -QP$  etc.
- (e) **Unit vector** : It is a vector with the unit magnitude. The unit vector in the direction of vector  $\vec{a}$  is given by  $\hat{a} = \frac{\vec{a}}{|\vec{a}|}$  such that  $|\hat{a}| = 1$ , so, if  $\vec{a} = x\hat{i} + y\hat{j} + z\hat{k}$  then its unit vector is :

$$\hat{a} = \frac{x}{\sqrt{x^2 + y^2 + z^2}}\hat{i} + \frac{y}{\sqrt{x^2 + y^2 + z^2}}\hat{j} + \frac{z}{\sqrt{x^2 + y^2 + z^2}}\hat{k}.$$

- Unit vector perpendicular to the plane  $\vec{a}$  and  $\vec{b}$  is :  $\pm \frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|}$ .
- (f) **Reciprocal of a vector** : It is a vector which has the same direction as the vector  $\vec{a}$  but magnitude equal to the reciprocal of the magnitude of  $\vec{a}$ . It is denoted as  $\vec{a}^{-1}$ . Hence  $|\vec{a}^{-1}| = \frac{1}{|\vec{a}|}$ .
- (g) **Equal vectors** : Two vectors are said to be equal if they have the same magnitude as well as direction, regardless of the position of their initial points.

$$\text{Thus } \vec{a} = \vec{b} \Leftrightarrow \begin{cases} |\vec{a}| = |\vec{b}| \\ \vec{a} \text{ and } \vec{b} \text{ have same direction} \end{cases}$$

Also, if  $\vec{a} = \vec{b} \Rightarrow a_1\hat{i} + a_2\hat{j} + a_3\hat{k} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k} \Rightarrow a_1 = b_1, a_2 = b_2, a_3 = b_3$ .

- (h) **Collinear or Parallel vector** : Two vectors  $\vec{a}$  and  $\vec{b}$  are collinear or parallel if there exists a non-zero scalar  $\lambda$  such that  $\vec{a} = \lambda\vec{b}$ .
- It is important to note that the respective coefficients of  $\hat{i}, \hat{j}, \hat{k}$  in  $\vec{a}$  and  $\vec{b}$  are proportional provided they are parallel or collinear to each other.
  - The d.r.'s of parallel vectors are same (or are in proportion).
  - The vectors  $\vec{a}$  and  $\vec{b}$  will have same or opposite direction as  $\lambda$  is positive or negative respectively.
  - The vectors  $\vec{a}$  and  $\vec{b}$  are collinear if  $\vec{a} \times \vec{b} = \vec{0}$ .
- (i) **Free vectors** : The vectors which can undergo parallel displacement without changing its magnitude and direction are called free vectors.

## Know the Formulae

The position vector of a point say  $P$  dividing a line segment joining the points  $A$  and  $B$  whose position vectors are  $\vec{a}$  and  $\vec{b}$  respectively, in the ratio  $m : n$ .

(a) Internally,  $\vec{OP} = \frac{m\vec{b} + n\vec{a}}{m+n}$

(b) Externally,  $\vec{OP} = \frac{m\vec{b} - n\vec{a}}{m-n}$

- Also if point  $P$  is the mid-point of line segment  $AB$ , then  $\vec{OP} = \frac{\vec{a} + \vec{b}}{2}$ .

## Objective Type Questions

(1 mark each)

Q.1. Area of a rectangle having vertices A, B, C and

... (i)

D with position vectors

$$\vec{AB} \cdot \vec{BC} = -\vec{CA}$$

... (ii)

and respectively is

∴ The equation given in alternative (a) is true.

- (a)  $\frac{1}{2}$
- (b) 1
- (c) 2
- (d) 4

$$\vec{AB} + \vec{BC} = \vec{AC}$$

[NCERT Ex.]

∴ The equation given in alternative (b) is true.

Ans. Correct option : (c)

From equation (ii), we have

Explanation :

The position vectors of vertices A, B, C and D of rectangle ABCD are given as :

The equation given in alternative (d) is true.

$$\vec{OA} = -\hat{i} - \hat{j} + \hat{k}, \vec{OB} = \hat{i} - \hat{j} + 4\hat{k}$$

Now, consider the equation given in alternative (c) :

$$= \hat{i} - \frac{1}{2}\hat{j} + 4\hat{k}, \vec{OD} = -\hat{i} - \frac{1}{2}\hat{j} + 4\hat{k}$$

$$\vec{AB} + \vec{BC} - \vec{CA} = 0$$

... (iii)

The adjacent sides and of the given rectangle are given as :

For equations (i) and (iii), we have :

$$\vec{AB} = (1+1)\hat{i} + \left(\frac{1}{2} - \frac{1}{2}\right)\hat{j} + (4-4)\hat{k} = 2\hat{i}$$

$$\vec{AC} = \vec{CA}$$

$$\vec{BC} = (1-1)\hat{i} + \left(-\frac{1}{2} - \frac{1}{2}\right)\hat{j} + (4-4)\hat{k} = -\hat{j}$$

$$\Rightarrow \vec{AC} + \vec{AC} =$$

$$\vec{AB} \cdot \vec{BC} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 0 & 0 \\ 0 & -1 & 0 \end{vmatrix}$$

, which is not true.

So that, the equation given in alternative (c) is incorrect.

$$= (-) = -2$$

$$|\vec{AB} \cdot \vec{BC}| = 2$$

Q.3. If  $\vec{a}$  is a non-zero vector of magnitude 'a' and  $\lambda$  a

non-zero scalar, then  $\lambda\vec{a}$  is unit vector if

- (a)  $\lambda = 1$
- (b)  $\lambda = -1$
- (c)  $a = |\lambda|$
- (d)  $a = 1/|\lambda|$

[NCERT Ex.]

Ans. Correct option : (d)

Explanation :

Vector  $\vec{a}$  is a unit vector if  $|\vec{a}| = 1$ .

Now,

$$|\lambda\vec{a}| = 1$$

$$\lambda a = 1$$


$$\vec{a} = \frac{1}{\lambda} \quad [\lambda \neq 0]$$

$$\vec{a} = \frac{1}{\lambda} \vec{a} \quad [1 = a]$$

So that, vector  $\vec{a}$  is a unit vector if  $a = \frac{1}{|\lambda|}$ .

Q.2. In triangle ABC (Figure), which of the following is not true :


- (a)  $\vec{AB} + \vec{BC} = \vec{AC}$
- (b)  $\vec{AB} + \vec{AC} = \vec{BC}$
- (c)  $\vec{AB} + \vec{BC} + \vec{CA} = \vec{0}$
- (d)  $\vec{AB} + \vec{BC} + \vec{AC} = \vec{0}$



[NCERT Ex.]

Ans. Correct option : (c)

Explanation :



Applying the triangle law of addition in the above triangle, we have

Q.4. If  $\vec{a}$  and  $\vec{b}$  are two collinear vectors, then which of the following are incorrect :

- (a)  $\vec{a} = \lambda\vec{b}$ , for some scalar  $\lambda$
- (b)  $\vec{a} \cdot \vec{b} = |\vec{a}||\vec{b}|$
- (c) the respective components of  $\vec{a}$  and  $\vec{b}$  are not proportional
- (d) both the vectors  $\vec{a}$  and  $\vec{b}$  have same direction, but different magnitudes

[NCERT Ex.]

Ans. Correct option : (d)

**Explanation :**

If  $\vec{a}$  and  $\vec{b}$  are two collinear vectors, then they are parallel.

Therefore, we have

(For some scalar  $\lambda$ )

If  $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$  and  $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$  then,

$$b_1\hat{i} + b_2\hat{j} + b_3\hat{k} = \lambda (a_1\hat{i} + a_2\hat{j} + a_3\hat{k})$$

$$b_1\hat{i} + b_2\hat{j} + b_3\hat{k} = (\lambda a_1)\hat{i} + (\lambda a_2)\hat{j} + (\lambda a_3)\hat{k}$$

$$b_1 = \lambda a_1, b_2 = \lambda a_2, b_3 = \lambda a_3$$

$$\frac{b_1}{a_1} = \frac{b_2}{a_2} = \frac{b_3}{a_3} = \lambda$$

So that, the respective components of  $\vec{a}$  and  $\vec{b}$  are proportional. However, vectors  $\vec{a}$  and  $\vec{b}$  can have different directions. Hence, the statement given in option (d) is incorrect.

**Q. 5. The vector in the direction of the vector  $i - j + k$  that has magnitude 9 is**

(a)  $i - j + k$                       (b)  $\frac{i - j + k}{3}$

(c)  $3i - 2j + 2k$                       (d)  $9i - 2j + 2k$

[NCERT Exemp.]

**Ans. Correct option : (c)**

**Explanation :**

Let  $\vec{a} = i - j + k$

Any vector in the direction of a vector  $\vec{a}$  is given by  $\frac{\vec{a}}{|\vec{a}|}$ .

$$= \frac{i - j + k}{\sqrt{1^2 + (-1)^2 + 1^2}} = \frac{i - j + k}{\sqrt{3}}$$

$\therefore$  Vector in the direction of  $\vec{a}$  with magnitude 9.

$$= 9 \frac{i - j + k}{\sqrt{3}}$$

$$= 3i - 2j + 2k$$

**Q. 6. The position vector of the point which divides the join of points  $P$  and  $Q$  in the ratio 3 : 1 is :**

(a)  $\frac{3\vec{p} + \vec{q}}{4}$                       (b)  $\frac{3\vec{q} + \vec{p}}{4}$

(c)  $\frac{3\vec{p} + 3\vec{q}}{4}$                       (d)  $\frac{3\vec{p} + \vec{q}}{4}$

[NCERT Exemp.]

**Ans. Correct option : (d)**

**Explanation :**

Let the position vector of the  $R$  divides the join of points  $P$  and  $Q$ .

$$\therefore \text{Position vector, } R = \frac{3(a+b)+1}{4}$$

Since, the position vector of a point  $R$  dividing the line segments joining the points  $P$  and  $Q$ , whose position vectors are  $p$  and  $q$  in the ration  $m : n$

internally, is given by  $\frac{mq + np}{m+n}$ .

$$\therefore R = \frac{5a}{4}$$

## Very Short Answer Type Questions

(1 mark each)

**Q. 1. Find a vector in the direction of  $\vec{a} = i - j$  that has magnitude 7 units.** [R&U] [NCERT] [Delhi Set I, II, III Comptt. 2015]

**Sol.**  $\vec{a} = \frac{1}{\sqrt{5}}\hat{i} - \frac{2}{\sqrt{5}}\hat{j}$                        $\frac{\vec{a}}{|\vec{a}|} = \frac{i - j}{\sqrt{2}}$                        $\frac{1}{2}$

then  $\vec{b} = \frac{7}{\sqrt{2}}\hat{i} - \frac{14}{\sqrt{2}}\hat{j}$                        $\frac{1}{2}$

[CBSE Marking Scheme 2015]

**Q. 2. Write a vector in the direction of the vector  $\hat{i} - 2\hat{j} + 2\hat{k}$  that has magnitude 9 units.**

[R&U] [Delhi Set I Comptt. 2014]

**Sol.** Let  $\vec{a} = i - 2j + 2k$

The vector in the direction of  $\vec{a}$  with magnitude 9 is  $\frac{9\vec{a}}{|\vec{a}|}$ .

$$\therefore \text{Required vector} = 9 \times \frac{i - 2j + 2k}{\sqrt{1^2 + (-2)^2 + 2^2}}$$

$$= 9 \times \frac{i - j + k}{\sqrt{2}}$$

$$= 3i - 6j + 6k \quad 1$$

**Q. 3. Write a unit vector in the direction of the sum of vectors  $\vec{a} = 2\hat{i} + 2\hat{j} - 5\hat{k}$  and  $\vec{b} = 2\hat{i} + \hat{j} - 7\hat{k}$ .**

[R&U] [Delhi Set III, 2014]

**Sol.** Let  $\vec{c} = \vec{a} + \vec{b} = 4\hat{i} + 3\hat{j} - 12\hat{k}$

$$r = \sqrt{16 + 9 + 144} = \sqrt{169} = 13 \quad \frac{1}{2}$$

So, unit vector

$$\hat{c} = \frac{\vec{c}}{r} = \frac{4\hat{i} + 3\hat{j} - 12\hat{k}}{13}$$

$$= \frac{4}{13}\hat{i} + \frac{3}{13}\hat{j} - \frac{12}{13}\hat{k} \quad \frac{1}{2}$$

**Q. 4. Find a vector in the direction of vector  $2\hat{i} - 3\hat{j} + 6\hat{k}$  which has magnitude 21 units.**

[R&U] [Foreign Set I, II, III, 2014]

**Sol.** Let  $\vec{a} = 2\hat{i} - 3\hat{j} + 6\hat{k}$

The vector in the direction of  $\vec{a}$  with a magnitude 21 is  $21 \times \hat{a}$ .

$$\begin{aligned} \therefore \text{Required vector} &= 21 \times \frac{2i-3j+6k}{\sqrt{2^2+(-3)^2+6^2}} \\ &= 21 \times \frac{2i-3j+6k}{7} \\ &= 6i-9j+18k \quad 1 \end{aligned}$$

**Q. 5.** If  $\vec{a} = x\hat{i} + 2\hat{j} - z\hat{k}$  and  $\vec{b} = 3\hat{i} - y\hat{j} + k$  are two equal vectors, then write the value of  $x + y + z$ . R&U [Delhi Set I, 2013]

**Sol.**

$$\begin{aligned} \vec{a} &= x\hat{i} + 2\hat{j} - z\hat{k} \\ \text{and } \vec{b} &= 3\hat{i} - y\hat{j} + k \\ \text{are equal vectors} \\ \text{So, } & x = 3, y = -2, z = -1 \\ \text{or } x\hat{i} + 2\hat{j} - z\hat{k} &= 3\hat{i} - y\hat{j} + k \quad \frac{1}{2} \\ \therefore x = 3, y = -2, z = -1 & \quad \frac{1}{2} \\ \therefore x + y + z = 3 - 2 - 1 = 0. & \quad \frac{1}{2} \end{aligned}$$

**Q. 6.** Write a unit vector in the direction of the sum of vectors :

$$\vec{a} = 2\hat{i} - \hat{j} + 2\hat{k} \quad \text{and} \quad \vec{b} = -\hat{i} + \hat{j} + 3\hat{k}$$

R&U [NCERT] [Delhi Set III, 2013]

**Sol.** Given,

$$\begin{aligned} \vec{a} &= 2\hat{i} - \hat{j} + 2\hat{k} \\ \text{and } \vec{b} &= -\hat{i} + \hat{j} + 3\hat{k} \\ \text{Let, } \vec{r} &= \vec{a} + \vec{b} \\ &= (2\hat{i} - \hat{j} + 2\hat{k}) + (-\hat{i} + \hat{j} + 3\hat{k}) \\ &= \hat{i} + 5\hat{k} \\ |\vec{r}| &= \sqrt{1^2 + 5^2} = \sqrt{26} \quad \frac{1}{2} \end{aligned}$$

So, required unit vector

$$\begin{aligned} r &= \frac{\vec{r}}{|\vec{r}|} = \frac{\hat{i} + 5\hat{k}}{\sqrt{26}} \\ &= \frac{1}{\sqrt{26}}\hat{i} + \frac{5}{\sqrt{26}}\hat{k} \quad \frac{1}{2} \end{aligned}$$

**Commonly Made Error**

- Generally students commit errors in finding the unit vector as they don't get the result in required vector form.

**Q. 7.** Find a unit vector parallel to the sum of vectors  $\hat{i} + \hat{j} + \hat{k}$  and  $2\hat{i} - 3\hat{j} + 5\hat{k}$ .

U [Delhi Set I Comptt. 2012]

**Sol.** Sum of given two vectors is given as

$$\begin{aligned} &(i + j + k) + (2i - 3j + 5k) \\ &= (1+2)i + (1-3)j + (1+5)k \\ &= 3i - 2j + 6k = \quad \text{(say) } \vec{A} \end{aligned}$$

A unit vector parallel to this vector

$$\begin{aligned} &= \frac{3i - 2j + 6k}{|\vec{A}|} \\ &= \frac{3i - 2j + 6k}{\sqrt{3^2 + (-2)^2 + 6^2}} = \frac{3i - 2j + 6k}{\sqrt{49}} \\ &= \frac{3}{7}i - \frac{2}{7}j + \frac{6}{7}k. \quad \frac{1}{2} \end{aligned}$$

**Q. 8.** Find a unit vector in the direction of

$$\vec{A} = 3\hat{i} - 2\hat{j} + \hat{k}$$

U [O.D. Set I Comptt. 2012]

**Sol.** A unit vector in the direction of vector  $\vec{A}$  is given by

$$\begin{aligned} &= \frac{\vec{A}}{|\vec{A}|} \\ \vec{A} &= \sqrt{3^2 + (-2)^2 + 1^2} = 4 \\ &= \frac{3i - 2j + k}{4} \end{aligned}$$

A unit vector in the direction of

$$= \frac{3}{4}i - \frac{2}{4}j + \frac{1}{4}k. \quad 1$$

**Q. 9.** Find the sum of the vectors  $\vec{a} = \hat{i} - 2\hat{j} + \hat{k}$ ,  $\vec{b} = -2\hat{i} + 4\hat{j} + 5\hat{k}$  and  $\vec{c} = \hat{i} - 6\hat{j} - \hat{k}$

U [NCERT] [Delhi Set I, 2012]

**Sol.**  $\vec{a} + \vec{b} + \vec{c}$

$$\begin{aligned} &= (i - 2j + k) + (-2i + 4j + 5k) + (i - 6j - k) \\ &= (1 - 2 + 1)i + (-2 + 4 - 6)j + (1 + 5 - 1)k \\ &= 0i - 4j - 1k = -4j - k. \quad 1 \end{aligned}$$

**Q.10.** Find the sum of the vectors :

$$a = i - 2j, b = 2i - 3j, c = 2i + k$$

U [Delhi Set II, 2012]

**Sol.**  $\vec{a} + \vec{b} + \vec{c}$

$$\begin{aligned} &= (i - 2j) + (2i - 3j) + (2i + k) \\ &= (1 + 2 + 2)i + (-2 - 3)j + k \\ &= 5i - 5j + k. \quad 1 \end{aligned}$$

**Q. 11. Find the sum of the vectors :**

$$\vec{a} = \hat{i} - 3\hat{k}, \vec{b} = 2\hat{j} - \hat{k}, \vec{c} = 2\hat{i} - 3\hat{j} + \hat{k}$$

**R&I** [Delhi Set III, 2012]

**Sol.**  $\vec{a} + \vec{b} + \vec{c} = (i - 3k) + (2j - k) + (2i - 3j + k)$   
 $= (1+2)i + (2-3)j + (-3-1+k)$   
 $= i - j - k.$  1

**Q. 12. Write the number of vectors of unit length**

**perpendicular to both the vector  $\vec{a} = 2\hat{i} + \hat{j} + 2\hat{k}$  and  $\vec{b} = \hat{j} + \hat{k}$ .** **O.D.** [Set I, II, III 2016]

**Sol.** There are two such vectors of unit length perpendicular to both the given vectors  $\vec{a}$  and  $\vec{b}$

and vectors are  $\pm \frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|}$ .

**Q. 13. Find the position vector of a point which divides the join of points with position vectors**


**$(a - 2b)$  and  $(2a + b)$  externally in the ratio 2 : 1.** **R&I** [Delhi Set I, II, III 2016]

**Sol.** Required vector =  $\frac{1(a - 2b) - 2(2a + b)}{1 - 2}$   
 $= \frac{(a - 2b) - 4a - 2b}{-1}$   
 $= 3a + 4b$  1/2

**R&I Q. 14. The two vectors  $\hat{i} + \hat{j}$  and  $\hat{i} - \hat{j} + 4\hat{k}$  represent the two sides  $AB$  and  $AC$ , respectively of  $\Delta ABC$ . Find the length of the median through  $A$ .**

**O.D.** [Delhi Set I, II, III 2016] [Foreign 2015]

**Sol.**  $\vec{AB} = \hat{i} + \hat{j}$  and  $\vec{AC} = \hat{i} - \hat{j} + 4\hat{k}$



Now  $ABEC$  represent a parallelogram with  $AE$  as the diagonal.

$$= AB + AC \quad 1/2$$

$$= (j+k) + 3i - j + 4k = i + k$$

Now,  $AE = \sqrt{(3)^2 + (5)^2} = \sqrt{9+25} = \sqrt{34}$

$\therefore \vec{AD} = \frac{1}{\sqrt{34}}$  units 1/2

**Q. 15. Write the position vector of the point which divides the join of points with position vectors**

**$(a - 2b)$  and  $(2a + b)$  in the ratio 2:1.** **R&I**

**Sol.** Let

$$\vec{OQ} =$$

The position vector of the point  $R$  dividing the join of  $P$  and  $Q$  internally in the ratio 2 : 1 is

$$\vec{r} = \frac{2(2a + b) + (a - 2b)}{2 + 1}$$
 1/2

$$= \frac{4a + 2b + a - 2b}{3}$$

$$= \frac{5a}{3} + \frac{2b}{3}$$
 1/2

**Q. 16. Write the value of  $p$  for which the vectors**

**$3\hat{i} + 2\hat{j} + 9\hat{k}$  and  $\hat{i} - 2p\hat{j} + 3\hat{k}$  are parallel vectors.** **O.D.** [Set I, 2014]

**Sol.**  $\frac{1}{a_1} = \frac{1}{b_1} = \frac{1}{c_1} = \frac{1}{a_2} = \frac{1}{b_2} = \frac{1}{c_2}$  [For parallel vectors]

or  $\frac{3}{1} = \frac{2}{-2p} = \frac{9}{3}$

or  $p = -1$  1

**Q. 17. Find a vector  $\vec{r}$  of magnitude  $\sqrt{14}$  making an angle of  $\frac{\pi}{4}$  with  $x$ -axis,  $\frac{\pi}{3}$  with  $y$ -axis and an acute angle  $\theta$  with  $z$ -axis.** **O.D.** [Set II 2014]

**Sol.** Let  $\hat{i}$  be the unit vector in the direction of  $x$ -axis.

Since vector  $\vec{r}$  makes an angle of  $\frac{\pi}{4}$  with  $x$ -axis,

$\frac{\pi}{3}$  with  $y$ -axis and an acute angle  $\theta$  with  $z$ -axis

therefore

$$\cos \frac{\pi}{4} = \frac{r_x}{\sqrt{14}}, \cos \frac{\pi}{3} = \frac{r_y}{\sqrt{14}}, \cos \theta = \frac{r_z}{\sqrt{14}}$$

[Using  $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$ ]

or  $\cos^2\theta = \frac{1}{2}$

or  $\cos \theta = \frac{1}{\sqrt{2}}$

or  $\theta = \frac{\pi}{4}$

Therefore,  $\vec{a} = 5\sqrt{2}a$   
 $= 5\sqrt{2}\left(\cos\frac{\pi}{4}\hat{i} + \cos\frac{\pi}{4}\hat{j} + \cos\frac{\pi}{4}\hat{k}\right)$   
 $= 5\sqrt{2}\left(\frac{1}{\sqrt{2}}\hat{i} + \frac{1}{\sqrt{2}}\hat{j} + \frac{1}{\sqrt{2}}\hat{k}\right)$   
 $= 5(\hat{i} + \hat{j} + \hat{k})$

**Q. 18.** If a unit vector  $\vec{a}$  makes an angle  $\frac{\pi}{4}$  with  $\hat{i}$ ,  $\frac{\pi}{4}$  with  $\hat{j}$  and an acute angle  $\theta$  with  $\hat{k}$ , then find the value of  $\theta$ . □ [Delhi Set I, 2013]

**Sol.** We know that if a vector  $\vec{a}$  makes angle  $\alpha$ ,  $\beta$  &  $\gamma$  with  $\hat{i}$ ,  $\hat{j}$  and  $\hat{k}$  respectively, then

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$$

Here, we have

$$\alpha = \frac{\pi}{4}, \beta = \frac{\pi}{4} \text{ and } \gamma = \theta, \text{ an acute angle}$$

$$\therefore \cos^2 \frac{\pi}{4} + \cos^2 \frac{\pi}{4} + \cos^2 \theta = 1$$

or  $\frac{1}{2} + \frac{1}{2} + \cos^2 \theta = 1$

or  $\cos^2 \theta = \frac{1}{2}$

or  $\cos \theta = \pm \frac{1}{\sqrt{2}}$  or  $\theta = \frac{\pi}{4}$

**Q. 19.**  $P$  and  $Q$  are two points with position vectors  $\vec{p}$  and  $\vec{q}$  respectively. Write the position vector of a point  $R$  which divides the line segment  $PQ$  externally in the ratio  $2 : 1$ .

□ [NCERT] [O.D. Set I, 2013]

**Sol.** Consider two points  $P$  and  $Q$  with position vectors  $\vec{p}$  and  $\vec{q}$

and  $OQ = 2OP$ , then position vector of the point  $R$  dividing the join of  $P$  and  $Q$  externally in the ratio  $2 : 1$  is

$$\vec{r} = \frac{2\vec{q} - \vec{p}}{2 - 1}$$

**Q. 20.**  $L$  and  $M$  are two points with position vectors  $\vec{l}$  and  $\vec{m}$  respectively. Write the position vectors of a point  $N$  which divides the line segment  $LM$  in the ratio  $2:1$  externally.

□ [O.D. Set I, 2013]

**Sol.** If  $\vec{n}$  is the position vector of  $N$ , then by section formula

$$\vec{n} = \frac{2(\vec{a} + 2\vec{b}) - 1 \cdot 2\vec{a}}{2 - 1}$$

$$= \frac{2\vec{a} + 4\vec{b} - 2\vec{a}}{1}$$

$$= 4\vec{b}$$

**Q.21.** Find the scalar components of the vector  $\vec{AB}$  with initial point  $A(2, 1)$  and terminal point  $B(-5, 7)$ .

□ [O.D. Set I, II, III, 2012]

**Sol.**  $\vec{AB}$  = Position vector of  $B$  - Position vector of  $A$

$$= (-5\hat{i} + 7\hat{j}) - (2\hat{i} + \hat{j})$$

$$= (-5 - 2)\hat{i} + (7 - 1)\hat{j}$$

$$= -7\hat{i} + 6\hat{j}$$

$\therefore$  The scalar components are  $(-7, 6)$ .

**Q. 22.** If a line has direction ratios  $2, -1, -2$ , then what are its direction cosines?

□ [Delhi Set I, II, III, 2012]

**Sol.** Here direction ratios of line are  $2, -1, -2$ .

$$\therefore \text{Direction cosines of line are } \frac{2}{\sqrt{2^2 + (-1)^2 + (-2)^2}}, \frac{-1}{\sqrt{2^2 + (-1)^2 + (-2)^2}}, \frac{-2}{\sqrt{2^2 + (-1)^2 + (-2)^2}}$$

$$\text{i.e., } \frac{2}{3}, \frac{-1}{3}, \frac{-2}{3}$$

[Note : If  $a, b, c$  are the direction ratios of a line, the direction cosines are  $\frac{a}{\sqrt{a^2 + b^2 + c^2}}, \frac{b}{\sqrt{a^2 + b^2 + c^2}}, \frac{c}{\sqrt{a^2 + b^2 + c^2}}$ ]

**Q. 23.** If  $\vec{a}$  and  $\vec{b}$  denote the position vectors of points  $A$  and  $B$  respectively and  $C$  is a point on  $AB$  such that  $AC = 2CB$ , then write the position vector of  $C$ . □ [Outside Delhi Set I, II, III comptt. 2016]

**Sol.**  $AC : CB = 2 : 1$   
 Position vector of  $C$   
 $= \frac{2\vec{b} + \vec{a}}{2 + 1}$

1

[CBSE Marking Scheme 2016]

**Q. 24.** If  $\vec{a} = 4\hat{i} - \hat{j} + \hat{k}$  and  $\vec{b} = 2\hat{i} - 2\hat{j} + \hat{k}$ , then find a unit vector parallel to the vector  $\vec{a} + \vec{b}$ .

□ [Outside Delhi Set I, II, III comptt. 2016]



**Sol.**  $\vec{a} = 6i - 3j + 2k$   $\frac{1}{7}$   
 Unit vector parallel to  $\vec{a}$  is  $\frac{1}{7}(6\hat{i} - 3\hat{j} + 2\hat{k})$   $\frac{1}{7}$   
**[CBSE Marking Scheme 2016]**

**Q. 25.** Give an example of vectors  $\vec{a}$  and  $\vec{b}$  such that  $\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}|$  but  $\vec{a} \neq \vec{b}$ . **[R&U] [SQP 2017-18]**

**Sol.**  $\vec{a} = \hat{i}, \vec{b} = \hat{j}$  [or any other correct answer] **1**  
**[CBSE Marking Scheme, 2017-18]**

**Q. 26.** Write a unit vector in the direction of vector  $\vec{PQ}$  where  $\vec{P}$  and  $\vec{Q}$  are the points (1, 3, 0) and (4, 5, 6), respectively. **[U] [Foreign 2014]**

**Sol.** Given points are  $\vec{P}(1, 3, 0)$  and  $\vec{Q}(4, 5, 6)$ .  
 Here,  $x_1 = 1, y_1 = 3, z_1 = 0$   
 and  $x_2 = 4, y_2 = 5, z_2 = 6$   
 So, vector  $\vec{PQ} = (x_2 - x_1)\hat{i} + (y_2 - y_1)\hat{j} + (z_2 - z_1)\hat{k}$   
 $= (4 - 1)\hat{i} + (5 - 3)\hat{j} + (6 - 0)\hat{k}$   
 $= 3\hat{i} + 2\hat{j} + 6\hat{k}$   $\frac{1}{7}$

$\therefore$  Magnitude of given vector  
 $|\vec{PQ}| = \sqrt{3^2 + 2^2 + 6^2} = \sqrt{49} = 7$  units  
 Hence, the unit vector in the direction of  $\vec{PQ}$  is  
 $\frac{\vec{PQ}}{|\vec{PQ}|} = \frac{3\hat{i} + 2\hat{j} + 6\hat{k}}{7} = \frac{3}{7}\hat{i} + \frac{2}{7}\hat{j} + \frac{6}{7}\hat{k}$   $\frac{1}{7}$

**Q. 27.** For what values of  $k$ , the vectors  $2\hat{i} - 3\hat{j} + 4\hat{k}$  and  $k\hat{i} + \hat{j} - \hat{k}$  are collinear?  
**[R&U] [HOTS; Delhi 2011]**

**Sol.** Let given vectors are  $\vec{a} = 2\hat{i} - 3\hat{j} + 4\hat{k}$  and  $\vec{b} = k\hat{i} + \hat{j} - \hat{k}$   
 vectors  $\vec{a}$  and  $\vec{b}$  are said to be collinear, if  
 $\vec{a} = k\vec{b}$ , where  $k$  is a scalar.  
 $\therefore 2\hat{i} - 3\hat{j} + 4\hat{k} = k(k\hat{i} + \hat{j} - \hat{k})$

On comparing the coefficients of  $\hat{i}$  and  $\hat{j}$ , we get  
 $2 = ka$  and  $-3 = 6k$  or  $k = -\frac{1}{2}$   
 $\therefore 2 = -\frac{1}{2}a$  or  $a = -4$  **1**

**Answering Tips**  
 • Clarify the concept of collinearity of two vectors.

**Q. 28.** If  $A, B$  and  $C$  are the vertices of a  $\triangle ABC$ , then what is the value of  $\vec{AB} + \vec{BC} + \vec{CA}$ ? **[U] [Delhi 2011C]**  
**Sol.** Let  $\triangle ABC$  be the given triangle.

Now, by triangle law of vector addition, we have  
 $\vec{AB} + \vec{BC} = \vec{AC}$   
 or  $\vec{AB} + \vec{BC} + \vec{CA} = \vec{AC} + \vec{CA}$  [adding  $\vec{CA}$  on both sides]  
 or  $\vec{AB} + \vec{BC} + \vec{CA} = \vec{0}$   $\therefore AC = -CA$   
 $\therefore \vec{AB} + \vec{BC} + \vec{CA} = \vec{0}$  **1**

**Q. 29.** Find the unit vector in the direction of the sum of vectors  $2\hat{i} + 3\hat{j} - \hat{k}$  and  $4\hat{i} - 3\hat{j} - 2\hat{k}$ .  
**[U] [Foreign 2015]**

**Sol.** Try Yourself  
**Q. 30.**  $A$  and  $B$  are two points with position vectors  $\vec{a}$  and  $\vec{b}$  respectively. Write the position vector of a point  $P$  which divides the line segment  $AB$  internally in the ratio 1 : 2.  
**[U] [All India 2013]**

**Sol.** Try Yourself  
**Q. 31.** Write the position vector of mid-point of the vector joining points  $P(2, 3, 4)$  and  $Q(4, 1, -2)$ .  
**[U] [Foreign 2011]**

**Sol.** Try Yourself  
**Q. 32.** Write a unit vector in the direction of vector  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$  **[R&U] [All India 2011; Delhi 2009]**

**Sol.** Try Yourself  
**Q. 33.** Find the magnitude of the vector  $\vec{a} = 3\hat{i} + 2\hat{j} + 6\hat{k}$ .  
**[U] [All India 2011C; Delhi 2008]**

**Sol.** Try Yourself

Q. 34. Find a unit vector in the direction of vector  
 $\vec{a} = 2\hat{i} + 3\hat{j} + 6\hat{k}$  R& [All India 2011C]

Sol. Try Yourself

Q. 35. Find a unit vector in the direction of vector  
 $\vec{a} = 2\hat{i} - 3\hat{j} + 6\hat{k}$  U [Delhi 2011C]

Sol. Try Yourself

Q. 36. Find a vector in the direction of  $\vec{a} = \hat{i} - \hat{j} + \hat{k}$ ,  
 which has magnitude 6 units. U [Delhi 2010C]

Sol. Try Yourself

Q. 37. Find a unit vector in the direction of vector  
 $\vec{a} = 6\hat{i} - 2\hat{j} + 3\hat{k}$ . R& [All India 2009C]

Sol. Try Yourself

Q. 38. Find a unit vector in the direction of vector  
 $\vec{a} = 2\hat{i} - 6\hat{j} + 3\hat{k}$ . R& [Delhi 2009]

Sol. Try Yourself

Q. 39. Find the position vector of mid-point of the line  
 segment AB, where A is point (3, 4, -2) and B is  
 point (1, 2, 4). R& [Delhi 2010]

Sol. Try Yourself

Q. 40. Write a vector of magnitude 9 units in the direction  
 of vector  $-\hat{i} + \hat{j} + \hat{k}$ . U [All India 2010]

Sol. Try Yourself

Q. 41. Write a vector of magnitude 15 units in the  
 direction of vector  $\hat{i} - \hat{j} + \hat{k}$ . R& [Delhi 2010]

Sol. Try yourself

## Short Answer Type Questions

(2 marks each)

Q. 1. Find the area of the parallelogram whose diagonals  
 are represented by the vectors  $\vec{a} = 2\hat{i} - 3\hat{j} + 4\hat{k}$

and  $\vec{b} = \hat{i} - \hat{j} + \hat{k}$  R& [SQP 2018-19]

Sol.

$$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & -3 & 4 \\ 2 & -1 & 2 \end{vmatrix}$$

$$= -2\hat{i} + 4\hat{j} + 4\hat{k} \quad 1$$

$$|\vec{a} \times \vec{b}| = \sqrt{4+16+16} = 6 \quad \frac{1}{2}$$

$$\text{Area of the parallelogram} = \frac{|\vec{a} \times \vec{b}|}{2} = 3 \text{ sq units. } \frac{1}{2}$$

[CBSE Marking Scheme 2018-19]

### Answering Tip

- Clarify the concept of finding area of parallelogram whose diagonal all vectors.

Q. 2. Find the angle between the vectors

$\vec{a} = \hat{i} + \hat{j} - \hat{k}$  and  $\vec{b} = \hat{i} - \hat{j} + \hat{k}$  U [SQP 2018-19]

Sol. The angle  $\theta$  between the vectors  $\vec{a}$  and  $\vec{b}$  is given  
 by

$$\cos\theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} \quad 1$$

$$\text{i.e., } \cos\theta = \frac{(\hat{i} + \hat{j} - \hat{k}) \cdot (\hat{i} - \hat{j} + \hat{k})}{\sqrt{(1)^2 + (1)^2 + (-1)^2} \cdot \sqrt{(1)^2 + (-1)^2 + (1)^2}}$$

$$\text{i.e., } \cos\theta = \frac{1-1-1}{\sqrt{3}\sqrt{3}}$$

$$\text{i.e., } \cos\theta = -\frac{1}{3}$$

$$\Rightarrow \theta = \cos^{-1}\left(-\frac{1}{3}\right) \quad 1$$

[CBSE Marking Scheme 2018-19]

### Answering Tip

- Concept of Angle between two vectors should be revised thoroughly.

Q. 3. Show that each of three vectors is a unit vector :

$$\frac{1}{7}(2\hat{i} + 3\hat{j} + 6\hat{k}), \frac{1}{7}(6\hat{i} - 3\hat{j} + 2\hat{k}), \frac{1}{7}(3\hat{i} - 6\hat{j} + \hat{k})$$

Sol.  $\vec{a} = \frac{1}{7}(2\hat{i} + 3\hat{j} + 6\hat{k})$

$$= \frac{1}{7}(6\hat{i} - 3\hat{j} + 2\hat{k})$$

and  $\vec{b} = \frac{1}{7}(3\hat{i} - 6\hat{j} + 2\hat{k})$

$$\vec{a} = \frac{1}{7}\sqrt{4+9+36}$$

$$= \frac{1}{7}\sqrt{49}$$

$$= \frac{7}{7} = 1 \quad \frac{1}{2}$$

$$\vec{b} = \frac{1}{7}\sqrt{36+9+4}$$



$$= \frac{1}{\sqrt{49}}$$

$$= \frac{1}{7} = 1 \quad \frac{1}{2}$$

and  $\vec{c} = \frac{1}{\sqrt{9+3}} = \frac{1}{\sqrt{49}}$

$$= \frac{1}{7} = 1 \quad \frac{1}{2}$$

$$a = b = c = 1 \quad \frac{1}{2}$$

Hence, vectors are unit vectors.

**Q.4.** If  $\vec{a} = 3\hat{i} + 7\hat{j} + 2\hat{k}$ ,  $\vec{b} = 7\hat{i} - 2\hat{j} + 3\hat{k}$ . Is

$|\vec{a}| = |\vec{b}|$ . Can we say  $\vec{a} = \vec{b}$ ? Give reason.

**Sol.**  $\vec{a} = 3\hat{i} + 7\hat{j} + 2\hat{k}$

$$|\vec{a}| = \sqrt{3^2 + 7^2 + 2^2} = \sqrt{58}$$

and

$$\vec{b} = 7\hat{i} - 2\hat{j} + 3\hat{k}$$

$$|\vec{b}| = \sqrt{7^2 + 4 + 9} = \sqrt{62}$$

Thus,  $|\vec{a}| \neq |\vec{b}|$

but  $\vec{a} \neq \vec{b}$  because their corresponding components are different.

**Q.5.** Find the vector of magnitude of 9 units in the direction of  $\vec{a} = 3\hat{i} - 2\hat{j} + 3\hat{k}$  and

$$\vec{b} = \hat{i} - 4\hat{j}$$

**Sol.** Let,  $\vec{c} = \lambda(3\hat{i} - 2\hat{j} + 3\hat{k}) + \mu(\hat{i} - 4\hat{j})$

$$= (3\lambda + \mu)\hat{i} - (2\lambda + 4\mu)\hat{j} + 3\lambda\hat{k}$$

or

$$= 2\hat{i} + 2\hat{j} + 4\hat{k}$$

$$= \frac{\vec{c}}{|\vec{c}|} = \frac{2\hat{i} + 2\hat{j} + 4\hat{k}}{\sqrt{4 + 4 + 16}}$$

$$= \frac{2\hat{i} + 2\hat{j} + 4\hat{k}}{\sqrt{24}}$$

$$= \frac{\hat{i} + \hat{j} + 2\hat{k}}{\sqrt{6}}$$

$$= \frac{\hat{i} + \hat{j} + 2\hat{k}}{\sqrt{6}}$$

$\therefore$  Required vector  $= 9\hat{c} = \frac{9\hat{i} + 9\hat{j} + 18\hat{k}}{\sqrt{6}}$  1

**Q.6.** If  $|\vec{a} + \vec{b}| = 60$ ,  $|\vec{a} - \vec{b}| = 40$  and  $|\vec{a}| = 22$ , then

find  $|\vec{b}|$ . [R&] [S.Q.P. 2016-17]

**Sol.**  $|\vec{a} + \vec{b}|^2 + |\vec{a} - \vec{b}|^2 = 2(|\vec{a}|^2 + |\vec{b}|^2)$  1

or  $60^2 + 40^2 = 2(22^2 + |\vec{b}|^2)$  1/2

or  $|\vec{b}| = 46$  1/2

[CBSE Marking Scheme 2016]

**Q.7.** The position vectors of points A, B and C are  $\hat{i} + 3\hat{j}$ ,  $12\hat{i} + 3\hat{j}$  and  $11\hat{i} - 3\hat{j}$  respectively. If C divides the line segment joining A and B in the ratio 3 : 1, find the values of  $\lambda$  and  $\mu$ .

[R&] [Delhi Comptt. 2017]

**Sol.**  $11\hat{i} - 3\hat{j} = \frac{3(\hat{i} + 3\hat{j}) + \mu(12\hat{i} + 3\hat{j})}{3 + \mu}$  1

$$44 = 36 + \lambda, -12 = 3\mu + 3$$

$$\lambda = 8, \mu = -5 \quad \frac{1}{2} + \frac{1}{2}$$

[CBSE Marking Scheme, 2017]

## Long Answer Type Questions-I

(4 marks each)

**Q.1.** Find a vector of magnitude 5 units and parallel to the resultant of  $\vec{a} = 2\hat{i} + 3\hat{j} - \hat{k}$  and  $\vec{b} = \hat{i} - 2\hat{j} + \hat{k}$ . [R&] [Delhi 2011]

**Sol.** Given,  $\vec{a} = 2\hat{i} + 3\hat{j} - \hat{k}$  and  $\vec{b} = \hat{i} - 2\hat{j} + \hat{k}$

Now, resultant of above vectors =

$$= (2\hat{i} + 3\hat{j} - \hat{k}) + (\hat{i} - 2\hat{j} + \hat{k}) = 3\hat{i} + \hat{j}$$
 1

Let  $\vec{c} = \lambda(3\hat{i} + \hat{j})$

$$\therefore \vec{c} = 3\lambda\hat{i} + \lambda\hat{j}$$

Now unit vector  $\hat{c}$  in the direction of  $\vec{c}$  is  $\frac{\vec{c}}{|\vec{c}|}$

$$= \frac{3\lambda\hat{i} + \lambda\hat{j}}{\sqrt{(3\lambda)^2 + (\lambda)^2}}$$
 1

$$= \frac{3 + \hat{i}}{\sqrt{10}} = \frac{3}{\sqrt{10}} + \frac{1}{\sqrt{10}}\hat{i}$$
 1

Hence, vector of magnitude 5 units and parallel to resultant of  $\vec{a}$  and  $\vec{b}$  is.

$$5\left(\frac{3}{\sqrt{10}} + \frac{1}{\sqrt{10}}\right) \text{ or } \frac{15}{\sqrt{10}} + \frac{5}{\sqrt{10}} \quad 1$$

**Q. 2.** Let  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$ ,  $\vec{b} = 4\hat{i} - 2\hat{j} + 3\hat{k}$ , and  $\vec{c} = \hat{i} - 2\hat{j} + \hat{k}$ . Find a vector of magnitude 6 units, which is parallel to the vector  $\vec{a} + \vec{b} + \vec{c}$ .

**U [All India 2010]**

**Sol.** Given,  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$ ,  $\vec{b} = 4\hat{i} - 2\hat{j} + 3\hat{k}$

and  $\vec{c} = \hat{i} - 2\hat{j} + \hat{k}$

$\therefore \vec{a} + \vec{b} + \vec{c} =$

$$= 2(\hat{i} + \hat{j} + \hat{k}) - (4\hat{i} - 2\hat{j} + 3\hat{k}) + 3(\hat{i} - 2\hat{j} + \hat{k})$$

$$= 2\hat{i} + 2\hat{j} + 2\hat{k} - 4\hat{i} + 2\hat{j} - 3\hat{k} + 3\hat{i} - 6\hat{j} + 3\hat{k}$$

or  $\vec{a} + \vec{b} + \vec{c} = \hat{i} - 2\hat{j} + \hat{k} \quad 1\frac{1}{2}$

Now, a unit vector in the direction of vector

$$\begin{aligned} \vec{u} &= \frac{\hat{i} - 2\hat{j} + \hat{k}}{\sqrt{(1)^2 + (-2)^2 + (1)^2}} \\ &= \frac{\hat{i} - 2\hat{j} + \hat{k}}{\sqrt{6}} \\ &= \frac{1}{\sqrt{6}}(\hat{i} - 2\hat{j} + \hat{k}) \quad 1\frac{1}{2} \end{aligned}$$


Hence, vector of magnitude 6 units parallel to the

$$\begin{aligned} \text{vector } \vec{v} &= 6\left(\frac{1}{\sqrt{6}}(\hat{i} - 2\hat{j} + \hat{k})\right) \\ &= 2\hat{i} - 4\hat{j} + 2\hat{k} \quad 1 \end{aligned}$$

**Q. 3.** Find the position vector of a point R, which divides the line joining two points P and Q whose position vectors are  $\vec{a}$  and  $\vec{b}$  respectively, externally in the ratio 1 : 2. Also, show that P is the mid-point, of line segment RQ.

**R&U [HOTS; Delhi 2010]**

**Sol.** Given,  $\vec{a}$  = Position vector of P =  $\vec{a}$  and  $\vec{b}$  = Position vector of Q =  $\vec{b}$ . Let  $\vec{r}$  be the position vector of point R, which divides PQ in the ratio 1 : 2 externally.



$$\therefore \vec{r} = \frac{1(\vec{a} - 3\vec{b}) - 2\vec{a} + 2\vec{b}}{1 - 2} \quad 1$$

$$\left[ \therefore \vec{OR} = \frac{m(\vec{OQ}) - n(\vec{OP})}{m - n} \text{ Here, } m = 1, n = 2 \right]$$

$$= \frac{\vec{a} - 3\vec{b} - 2\vec{a} + 2\vec{b}}{-1}$$

$$= -\vec{a} + \vec{b} = \vec{b} - \vec{a}$$

Hence,  $\vec{r} = \vec{b} - \vec{a} \quad 1\frac{1}{2}$

Now, we have to show that P is the mid-point of RQ,

$$\text{i.e. } \vec{r} = \frac{\vec{OR} + \vec{OQ}}{2}$$

We have,  $\vec{r} = \vec{b} - \vec{a}$ ,  $\vec{OQ} = \vec{b}$

$$\therefore \frac{\vec{OR} + \vec{OQ}}{2} = \frac{(3\vec{a} + 5\vec{b}) + \vec{b} - 3\vec{a}}{2}$$

$$= \frac{\vec{a} + \vec{b}}{2} = \frac{2\vec{a} + 2\vec{b}}{2}$$

$$= \vec{a} + \vec{b} = \vec{a} + \vec{b} \quad \therefore OP = \frac{OR + OQ}{2}$$

Hence, P is the mid-point of line segment RQ.  $1\frac{1}{2}$

**Q. 4.** Show that the points  $A(-2\hat{i} + 3\hat{j} + 5\hat{k})$ ,  $B(\hat{i} + 2\hat{j} + 3\hat{k})$  and  $C(7 - \hat{i})$  are collinear. **[Delhi 2009C]**

**Sol.** Try yourself


## TOPIC-2

### Dot Product of Vectors

#### Revision Notes

##### 1. Products of Two Vectors and Projection of Vectors

- (a) **Scalar Product or Dot Product** : The dot product of two vectors  $\vec{a}$  and  $\vec{b}$  is defined by,  $\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta$  where  $\theta$  is the angle between  $\vec{a}$  and  $\vec{b}$ ,  $0 \leq \theta \leq \pi$ .



Consider  $\vec{a} = a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k}$ ,  $\vec{b} = b_1 \hat{i} + b_2 \hat{j} + b_3 \hat{k}$ , then  $\vec{a} \cdot \vec{b} = a_1 b_1 + a_2 b_2 + a_3 b_3$ .

⇒ **Projection of a vector**  $\vec{a}$  on the other vector say  $\vec{b}$  is given as  $\left( \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} \right) \frac{\vec{b}}{|\vec{b}|}$ .

⇒ **Projection of a vector**  $\vec{a}$  on the other vector say  $\vec{a}$  is given as  $\left( \frac{\vec{a} \cdot \vec{a}}{|\vec{a}|} \right) \frac{\vec{a}}{|\vec{a}|}$ .

#### Know the Properties (Dot Product)

- **Properties/Observations of Dot product**

⇒  $\hat{i} \cdot \hat{i} = |\hat{i}| |\hat{i}| \cos 0 = 1$  or  $\hat{i} \cdot \hat{i} = 1 = \hat{j} \cdot \hat{j} = \hat{k} \cdot \hat{k}$

⇒  $\hat{i} \cdot \hat{j} = |\hat{i}| |\hat{j}| \cos \frac{\pi}{2} = 0$  or  $\hat{i} \cdot \hat{j} = 0 = \hat{j} \cdot \hat{k} = \hat{k} \cdot \hat{i}$

⇒  $\vec{a} \cdot \vec{a} \in \mathbb{R}$ , where  $\mathbb{R}$  is real number i.e., any scalar.

⇒  $\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{a}$  (Commutative property of dot product).

⇒  $\vec{a} \cdot \vec{b} = 0 \Leftrightarrow \vec{a} \perp \vec{b}$  or  $|\vec{a}| = 0$  or  $|\vec{b}| = 0$ .

⇒ If  $\theta = 0$ , then  $\vec{a} \cdot \vec{a} = |\vec{a}|^2$ . Also  $\vec{a} \cdot \vec{a} = |\vec{a}|^2 \cos \theta$  as  $\theta$  in this case is 0.

Moreover if  $\theta = \pi$ , then  $\vec{a} \cdot \vec{a} = -|\vec{a}|^2$ .

⇒  $\vec{a} \cdot (\vec{b} + \vec{c}) = \vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c}$  (Distributive property of dot product).

⇒  $\vec{a} \cdot (-\vec{b}) = -(\vec{a} \cdot \vec{b}) = (-\vec{a}) \cdot \vec{b}$

#### Know the Formulae

- ⇒ **Angle between two vectors**  $\vec{a}$  and  $\vec{b}$  can be found by the expression given below :

$$\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} \text{ or } \theta = \cos^{-1} \left( \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} \right)$$

## Objective Type Questions

(1 mark each)

Q. 1. If  $\theta$  is the angle between two vectors  $\vec{a}$  and  $\vec{b}$ , then  $\vec{a} \cdot \vec{b} = 0$  only when

- (a)  $0 < \theta < \frac{\pi}{2}$                       (b)  $0 \leq \theta \leq \frac{\pi}{2}$   
 (c)  $0 < \theta < \pi$                       (d)  $0 \leq \theta \leq \pi$

[NCERT Misc. Ex. Q. 16, Page 459]

Ans. Correct option : (b)

Explanation :

Let  $\theta$  be the angle between two vectors  $\vec{a}$  and  $\vec{b}$ . Then, without loss of generality,  $\vec{a}$  and  $\vec{b}$  are non-zero vectors so that  $|\vec{a}|$  and  $|\vec{b}|$  are positive.

It is known that,  $\vec{a} \cdot \vec{b} = |\vec{a}||\vec{b}|\cos\theta$ .

$$\therefore 0 = |\vec{a}||\vec{b}|\cos\theta$$

$$\Rightarrow \cos\theta = 0$$

$$\left[ \because |\vec{a}| \text{ and } |\vec{b}| \text{ are positive.} \right]$$

$$\Rightarrow 0 \leq \theta \leq \pi$$

Q. 2. Let  $\vec{a}$  and  $\vec{b}$  be two-unit vectors and  $\theta$  is the angle between them. Then  $\vec{a} + \vec{b}$  is a unit vector if

- (a)  $\theta = \frac{\pi}{2}$                       (b)  $\theta = \frac{\pi}{3}$   
 (c)  $\theta = \frac{\pi}{4}$                       (d)  $\theta = \frac{2\pi}{3}$

[NCERT Misc. Ex. Q. 17, Page 459]

Ans. Correct option : (d)

Explanation :

Let  $\vec{a}$  and  $\vec{b}$  be two-unit vectors and  $\theta$  be the angle between them.

$$\text{Then, } |\vec{a} + \vec{b}| = |\vec{a}| + |\vec{b}| = 1.$$

Now,  $\vec{a} + \vec{b}$  is a unit vector if  $|\vec{a} + \vec{b}| = 1$ .

$$|\vec{a} + \vec{b}| = 1$$

$$(\vec{a} + \vec{b}) \cdot (\vec{a} + \vec{b}) = 1$$

$$\Rightarrow (\vec{a} + \vec{b}) \cdot (\vec{a} + \vec{b}) = 1$$

$$\Rightarrow \vec{a} \cdot \vec{a} + \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{a} + \vec{b} \cdot \vec{b} = 1$$

$$\Rightarrow |\vec{a}|^2 + \vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{b} + |\vec{b}|^2 = 1$$

$$\Rightarrow 1 + 2|\vec{a}||\vec{b}|\cos\theta + 1 = 1$$

$$\cos\theta = -\frac{1}{2}$$

$$\Rightarrow \theta = \frac{2\pi}{3}$$

So that,  $\vec{a} + \vec{b}$  is a unit vector if  $\theta = \frac{2\pi}{3}$ .

Q. 3. The angle between two vectors  $\vec{a}$  and  $\vec{b}$  with magnitudes  $\sqrt{3}$  and 4, respectively, and  $\vec{a} \cdot \vec{b} = \sqrt{3}$  is:

- (a)  $\frac{\pi}{6}$                       (b)  $\frac{\pi}{3}$   
 (c)  $\frac{\pi}{2}$                       (d)  $\frac{5\pi}{6}$

[NCERT Exemp. Ex. 10.3, Q. 22, Page 217]

Ans. Correct option : (b)

Explanation :

Here,  $|\vec{a}| = \sqrt{3}$ ,  $|\vec{b}| = 4$  and  $\vec{a} \cdot \vec{b} = \sqrt{3}$  Given]

We know that,

$$\vec{a} \cdot \vec{b} = |\vec{a}||\vec{b}|\cos\theta$$

$$\Rightarrow \sqrt{3} = \sqrt{3} \cdot 4 \cdot \cos\theta$$

$$\Rightarrow \cos\theta = \frac{\sqrt{3}}{4\sqrt{3}} = \frac{1}{4}$$

$$\therefore \theta = \frac{\pi}{3}$$

Q. 4. Find the value of  $\lambda$  such that the vectors  $\vec{a} = 2\hat{i} + \lambda\hat{j} + \hat{k}$  and  $\vec{b} = \hat{i} + \hat{j} + \hat{k}$  are orthogonal.

- (a) 0                      (b) 1  
 (c)  $\frac{3}{2}$                       (d)  $\frac{5}{2}$

[NCERT Exemp. Ex. 10.3, Q. 23, Page 217]

Ans. Correct option : (d)

Explanation :

Since, two non-zero vectors  $\vec{a}$  and  $\vec{b}$  are orthogonal, i.e.,  $\vec{a} \cdot \vec{b} = 0$

$$\therefore (2\hat{i} + \lambda\hat{j} + \hat{k}) \cdot (\hat{i} + \hat{j} + \hat{k}) = 0$$

$$2 + 2\lambda + 1 = 0$$

$$\lambda = -\frac{5}{2}$$

Q. 5. The value of  $\lambda$  for which the vectors  $3\hat{i} + 6\hat{j} + \hat{k}$  and  $2\hat{i} - 4\hat{j} + \lambda\hat{k}$  are parallel is

- (a)  $\frac{2}{3}$                       (b)  $\frac{3}{2}$   
 (c)  $\frac{5}{2}$                       (d)  $\frac{2}{5}$

[NCERT Exemp. Ex. 10.3, Q. 24, Page 217]

Ans. Correct option : (a)

Explanation :

Let  $\vec{a} = 3\hat{i} - 6\hat{j} + \hat{k}$  and  $\vec{b} = 2\hat{i} - 4\hat{j} + \lambda\hat{k}$

Since,  $\vec{a} \parallel \vec{b}$

$$\frac{3}{2} = \frac{-6}{-4} = \frac{1}{\lambda}$$

$$\lambda = \frac{2}{5}$$

Q. 6. If  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are unit vectors such that  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ , then the value of  $\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}$  is



- (a) 1
- (b) 3
- (c) -3/2
- (d) None of these

[NCERT Exemp. Ex. 10.3, Q. 29, Page 218]

Ans. Correct option : (c)

Explanation :

We have,  $a^2 + b^2 + c^2 = 1 + 1 + 1 = 3$  and  $a \cdot b + b \cdot c + c \cdot a = 0$

$$a^2 + b^2 + c^2 + 2(a \cdot b + b \cdot c + c \cdot a) = 3 + 2(0) = 3$$

$$a^2 + b^2 + c^2 + 2(a \cdot b + b \cdot c + c \cdot a) = 3$$

$$[ \because a \cdot b = b \cdot a, b \cdot c = c \cdot b \text{ and } c \cdot a = a \cdot c ]$$

$$1 + 1 + 1 + 2(a \cdot b + b \cdot c + c \cdot a) = 0$$

$$a \cdot b + b \cdot c + c \cdot a = -\frac{3}{2}$$

Q. 7. The projection vector of  $\vec{a}$  on  $\vec{b}$  is

- (a)  $\left( \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} \right) \vec{b}$
- (b)  $\vec{b}$
- (c)  $\vec{a}$
- (d)  $\left( \frac{\vec{a} \cdot \vec{b}}{|\vec{a}|} \right) \vec{b}$

[NCERT Exemp. Ex. 10.3, Q. 30, Page 218]

Ans. Correct option : (a)

Explanation :

Projection vector of  $\vec{a}$  on  $\vec{b}$  is given by,

$$\vec{a} \cdot \frac{\vec{b}}{|\vec{b}|}$$

$$\left( \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} \right) \vec{b}$$

Q. 8. If  $\vec{a}, \vec{b}$  and  $\vec{c}$  are three vectors such that  $a = 2, |b| = 3$  and  $c = 4$ , then the value of  $(\vec{a} + \vec{b} + \vec{c}) \cdot (\vec{a} + \vec{b} + \vec{c})$  is

- (a) 0
- (b) 1
- (c) -19
- (d) 38

[NCERT Exemp. Ex. 10.3, Q. 31, Page 218]

Ans. Correct option : (c)

Explanation :

Here,  $a = 2, b = 3$  and  $c = 4$

$$\therefore (\vec{a} + \vec{b} + \vec{c}) \cdot (\vec{a} + \vec{b} + \vec{c}) = 0$$

$$\Rightarrow a^2 + b^2 + c^2 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = 2^2 + 3^2 + 4^2 + 2(0)$$

$$\Rightarrow 4 + 9 + 16 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = 0$$

$$\Rightarrow 29 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = 0$$

$$\Rightarrow \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a} = -\frac{29}{2} = -14.5$$

## Very Short Answer Type Questions

(1 mark each)

Q. 1. If  $\vec{a}$  and  $\vec{b}$  are unit vectors, then what is the angle between  $\vec{a}$  and  $\vec{b}$ , so that  $\frac{\vec{a} + \vec{b}}{\sqrt{2}}$  is a unit vector? [Delhi Set I, II, III Comptt. 2015]

Sol.  $\frac{\vec{a} + \vec{b}}{\sqrt{2}}$  is a unit vector

$$2|\vec{a}|^2 + 2|\vec{b}|^2 + 2\vec{a} \cdot \vec{b} = 2$$

$$2 + 2 + 2\vec{a} \cdot \vec{b} = 2$$

$$4 + 2\vec{a} \cdot \vec{b} = 2$$

$$2\vec{a} \cdot \vec{b} = -2$$

$$\vec{a} \cdot \vec{b} = -1$$

$$\cos \theta = -1$$

$$1.1. \cos \theta = \frac{1}{\sqrt{2}}$$

$$\theta = \frac{\pi}{4}$$

[CBSE Marking Scheme 2015]

Q. 2. Find the projection of vector  $\vec{a} = 2\hat{i} + 3\hat{j} + 2\hat{k}$  on

the vector  $\vec{b} = 2\hat{i} + 2\hat{j} + \hat{k}$  [R&U] [NCERT]

[O.D. Set I, II, III Comptt. 2015]

Sol.  $(2i + 3j + 2k) \cdot (2i + 2j + k) = 12$  1/2

$$p = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} \text{ or } p = \frac{12}{\sqrt{2^2 + 2^2 + 1^2}}$$

$$= \frac{12}{3} = 4$$
 1/2

[CBSE Marking Scheme 2015]

Q. 3. Write the projection of the vector  $\vec{a} = 2\hat{i} - \hat{j} + \hat{k}$  on the vector  $\vec{b} = \hat{i} + 2\hat{j} + 2\hat{k}$ .

[R&U] [Delhi Set I, II, III Comptt. 2014]

Sol. Projection of a vector  $\vec{a}$  on the vector  $\vec{b}$  is given by

$$= \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|}$$

$$= \frac{(2i - j + k) \cdot (i + 2j + 2k)}{\sqrt{1^2 + 2^2 + 2^2}}$$

$$= \frac{2 + 2 - 2 + 2}{3} = \frac{4}{3}$$

[CBSE Marking Scheme 2014]

**Commonly Made Error**

- Most of the candidates calculate dot product instead of applying projection formula.

**Answering Tip**

- Clarify the concept of scalar projection of vector thoroughly.

**Q. 4.** Find the projection of vector  $\hat{i} + 3\hat{j} + 7\hat{k}$  on the vector  $2\hat{i} - 3\hat{j} + 6\hat{k}$ . R&U [Delhi Set II, III 2014]

**Sol.** Required projection :

$$= \frac{(\hat{i} + 3\hat{j} + 7\hat{k}) \cdot (2\hat{i} - 3\hat{j} + 6\hat{k})}{\sqrt{2^2 + (-3)^2 + 6^2}}$$

$$= \frac{2 - 9 + 42}{7} = 5 \quad 1$$

**Q. 5.** Write the projection of the vector  $\hat{i} + \hat{j} + \hat{k}$  along the vector  $\hat{j}$ . □ U [Foreign Set I, II, III 2014]

**Sol.** We know that the projection of a vector  $\vec{a}$  on the vector  $\vec{b}$  is given by  $\frac{\vec{a} \cdot \vec{b}}{|\vec{b}|}$ .

∴ The projection of the vector  $\hat{i} + \hat{j} + \hat{k}$  along the vector  $\hat{j}$  is,

$$\left( \hat{i} + \hat{j} + \hat{k} \right) \cdot \left( \frac{\hat{j}}{\sqrt{0^2 + 1^2 + 0^2}} \right) = 1 \quad 1$$

**Q. 6.** Write the projection of the vector  $7\hat{i} + \hat{j} - 4\hat{k}$  on the vector  $2\hat{i} + 6\hat{j} + 3\hat{k}$ . R&□ [Delhi 2015]  
[Delhi Set I, II, III Comptt. 2013]

**Sol.** Projection of  $\vec{a}$  on  $\vec{b} = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|}$

$$= \frac{(7\hat{i} + \hat{j} - 4\hat{k}) \cdot (2\hat{i} + 6\hat{j} + 3\hat{k})}{\sqrt{2^2 + 6^2 + 3^2}}$$

$$= \frac{14 + 6 - 12}{7} = 2 \quad 1$$

[CBSE Marking Scheme 2013]

**Q.7.** Write the projection of  $\vec{a} + \vec{c}$  on  $\vec{b}$ , where  $\vec{a} = 2\hat{i} + 2\hat{j} + \hat{k}$ ,  $\vec{b} = \hat{i} + 2\hat{j} + 2\hat{k}$  and  $\vec{c} = 2\hat{i} - \hat{j} + 4\hat{k}$ .

U [O.D. Set I, II, III Comptt. 2013]

**Sol.**

$$\vec{a} + \vec{c} = 2\hat{i} + 2\hat{j} + \hat{k} + 2\hat{i} - \hat{j} + 4\hat{k}$$

$$= 4\hat{i} + \hat{j} + 5\hat{k}$$

and  $\vec{b} = \hat{i} + 2\hat{j} + 2\hat{k}$

$$+ \vec{c} = 2\hat{i} - \hat{j} + 4\hat{k}$$

$$= 4\hat{i} + \hat{j} + 5\hat{k} + 2\hat{i} - \hat{j} + 4\hat{k} = 6\hat{i} + 9\hat{k}$$

Projection of  $\vec{a} + \vec{c}$  on  $\vec{b}$

$$= \frac{(\vec{a} + \vec{c}) \cdot \vec{b}}{|\vec{b}|} = \frac{(6\hat{i} + 9\hat{k}) \cdot (\hat{i} + 2\hat{j} + 2\hat{k})}{\sqrt{1^2 + 2^2 + 2^2}}$$

$$= \frac{6 + 18}{3} = 8 \quad \frac{1}{2}$$

[CBSE Marking Scheme 2013]

**Q. 8.** Find 'λ' when the projection of  $\vec{a} = \lambda\hat{i} + \hat{j} + 4\hat{k}$  on  $\vec{b} = 2\hat{i} + 6\hat{j} + 3\hat{k}$  is 4 units. R& [Delhi Set I, II, III 2012]

**Sol.** Projection of  $\vec{a}$  on  $\vec{b} = 4$ , (given)

$$\frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} = 4 \quad \frac{1}{2}$$

∴

$$\frac{(\lambda\hat{i} + \hat{j} + 4\hat{k}) \cdot (2\hat{i} + 6\hat{j} + 3\hat{k})}{\sqrt{2^2 + 6^2 + 3^2}} = 4$$

or  $2\lambda + 6 + 12 = 7 \times 4$

or  $2\lambda = 28 - 18 = 10$

or  $\lambda = \frac{10}{2} = 5 \quad \frac{1}{2}$

[CBSE Marking Scheme 2012]

**Q. 9.** If  $\vec{a}$  and  $\vec{b}$  are perpendicular vectors,  $|\vec{a} + \vec{b}| = 13$  and  $|\vec{a}| = 5$ , find the value of  $|\vec{b}|$ . □ [O.D. Set III 2014]



**Sol.**  $a + b^2 = |a|^2 + |b|^2 + 2 \dots$   
 or  $(13)^2 = (\dots) |\vec{b}|$   
 $\{\therefore \dots \Rightarrow \dots \cdot |\cos \theta = 0 \text{ as } \theta = 90^\circ\}$   
 or  $(169 - 25) = \vec{b}^2$   
 or  $b = 12$  1  
**[CBSE Marking Scheme 2014]**

**Q. 10.** Write the value of  $\lambda$  so that the vectors  $a = 2i - \lambda j + k$  and  $b = i - 2j + 3k$  are perpendicular to each other?

[Delhi Set I, II, III Comptt. 2013]  
 [O.D. Set I, II, III Comptt. 2012]

**Sol.**  $a = 2i - \lambda j + k$   
 and  $b = i - 2j + 3k$   
 For perpendicular :  
 $\dots = 0$   
 or  $2 \times 1 + \lambda(-2) + 1 \times 3 = 0$   $\frac{1}{2}$   
 or  $2\lambda = 5$   
 or  $\lambda = -\dots$   $\frac{1}{2}$

**Q. 11.** For what value of  $\lambda$  are the vectors  $2i + j - 3k$  and  $i + 2\lambda j + k$  perpendicular?

R& [O.D. Set I, II, III Comptt. 2013, 2011]  
 [Delhi Set I, II, III Comptt. 2012]

**Sol.** For two vectors to be perpendicular, their product should be zero.

$\therefore (i + 2\lambda j + k) \cdot (2i + j - 3k) = 0$   
 or  $1 \times 2 + 2\lambda \times 1 + 1 \times (-3) = 0$   
 $\lambda = -\dots$  1

**Q. 12.** Find  $x$ , if for a unit vector  $a$  and  $(x + a) \cdot (x - a) = 15$ .  [O.D. Set I 2013]

**Sol.**  $a = 1$   
 Given  $(x + a) \cdot (x - a) = 15$   
 or  $\dots = 15$   
 or  $x^2 - 1 = 15$   
 $\{\therefore a \text{ is a unit vector } |a| = 1\}$   
 $x^2 = 16$  or  $x = \pm 4$   
 As magnitude of a vector is non-negative.  
 So  $x = 4$ . 1  
**[CBSE Marking Scheme 2013]**

**Q. 13.** If  $\vec{a}$  and  $\vec{b}$  are unit vectors, then what is the angle between  $\vec{a} + \vec{b}$  and  $\vec{a} - \vec{b}$  for  $-\sqrt{b}$  to be unit vectors?  [O.D. Set-II, 2016]

**Sol.**

**[Topper's Answer 2016]**

**Q. 14.** If  $\hat{a}, \hat{b}$  and  $\hat{c}$  are mutually perpendicular unit vectors, then find value of  $2\hat{a} + \hat{b} + \hat{c}$ .  [All India 2015]

**Sol.** Given  $\hat{a}, \hat{b}$  and  $\hat{c}$  are mutually perpendicular unit vectors, i.e.,  
 $\dots = \dots = 0$  ...(i)

and  $\hat{a} = \hat{b} = \hat{c} = 1$  ...(ii)  
 Now,  $2\hat{a} + \hat{b} + \hat{c} = (\hat{a} + \hat{b} + \hat{c}) \cdot (\hat{a} + \hat{b} + \hat{c})$   
 $= 4(\hat{a} \cdot \hat{a}) + (\hat{a} \cdot \hat{b}) + (\hat{a} \cdot \hat{c}) + (\hat{b} \cdot \hat{a}) + (\hat{b} \cdot \hat{b}) + (\hat{b} \cdot \hat{c}) + 2(\hat{c} \cdot \hat{a}) + (\hat{c} \cdot \hat{b}) + (\hat{c} \cdot \hat{c})$   
 $[\therefore \vec{a} \cdot \vec{a} = |\vec{a}|^2]$

[∵ dot product is distributive over addition]  $\frac{1}{2}$

$$= 4(|\vec{a}| + 2(0) + 2(0) + 2(|\vec{c}|)) \cdot \frac{1}{2}$$

$$= 4(1 + 0 + 0 + 1) = 8$$

from Eq. (i) and  $\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{a}$

$$= 4(1) + 1 + 1 = 4 + 1 + 1 = 6$$

∴  $2a + b + c = \sqrt{6}$   $\frac{1}{2}$

[∵ length cannot be negative]  
[CBSE Marking Scheme 2015]

Q. 15. If  $\vec{a}$  and  $\vec{b}$  are two unit vectors such that  $\vec{a} + \vec{b}$  is also a unit vector, then find the angle between  $\vec{a}$  and  $\vec{b}$ . [R&U] [Delhi 2014]

Sol. Given,  $|\vec{a}| = 1, |\vec{b}| = 1$  and  $|\vec{a} + \vec{b}| = 1$

Now,  $(\vec{a} + \vec{b}) \cdot (\vec{a} + \vec{b}) = |\vec{a} + \vec{b}|^2$

$$= (\vec{a} \cdot \vec{a}) + (\vec{b} \cdot \vec{b}) + 2(\vec{a} \cdot \vec{b})$$

or  $1 + 1 + 2\vec{a} \cdot \vec{b} = 1$

$$2\vec{a} \cdot \vec{b} = 1 - 2 = -1$$

or  $\vec{a} \cdot \vec{b} = -\frac{1}{2}$  [∵  $\vec{a} \cdot \vec{b} = |\vec{a}||\vec{b}|\cos\theta$ ]

or  $\cos\theta = -\frac{1}{2}$  [∵  $|\vec{a}| = |\vec{b}| = 1$ ]

or  $\cos\theta = \cos\frac{2\pi}{3}$  or  $\theta = \frac{2\pi}{3}$

Hence, the angle between  $\vec{a}$  and  $\vec{b}$  is  $\frac{2\pi}{3}$ .  $\frac{1}{2}$

[CBSE Marking Scheme 2014]

Q. 16. Find  $\vec{a} \cdot \vec{b}$  if  $\vec{a} = -\hat{i} + \hat{j} - 2\hat{k}$  and  $\vec{b} = 2\hat{i} + 3\hat{j} - \hat{k}$ . [R&U] [All India 2009C]

Sol. Given,  $\vec{a} = -\hat{i} + \hat{j} - 2\hat{k}$  and  $\vec{b} = 2\hat{i} + 3\hat{j} - \hat{k}$

Then,  $\vec{a} \cdot \vec{b} = (-\hat{i} + \hat{j} - 2\hat{k}) \cdot (2\hat{i} + 3\hat{j} - \hat{k})$

$$= -2 + 3 + 2 = 3$$

[CBSE Marking Scheme 2019]

Q. 17. If  $|\vec{a}| = 2, |\vec{b}| = 3$  and  $|\vec{a} + \vec{b}| = 3$ , then find the projection of  $\vec{a}$  on  $\vec{b}$ . [R&U] [All India 2010C]

Sol. Given,  $|\vec{a}| = 2, |\vec{b}| = 3$  and  $|\vec{a} + \vec{b}| = 3$

∴ Projection of  $\vec{a}$  on  $\vec{b}$

$$= \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|} = \frac{\vec{a} \cdot (\vec{a} + \vec{b})}{|\vec{b}|}$$

$$= \frac{|\vec{a}|^2 + \vec{a} \cdot \vec{b}}{|\vec{b}|}$$

[∵  $\vec{a} \cdot \vec{a} = |\vec{a}|^2$  and  $|\vec{a}| = 2$ ]

Q. 18. If  $\vec{a}, \vec{b}, \vec{c}$  are unit vectors such that  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ , then write the value of  $\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}$ . [Foreign 2015]

Sol. Try Yourself [1]

Q. 19. If  $\vec{a}$  and  $\vec{b}$  are unit vectors, then find the angle between  $\vec{a} + \vec{b}$  and  $\vec{a} - \vec{b}$ , given that  $(\sqrt{3} - 1)$  is a unit vector. [R&U] [Delhi 2014C][NCERT Exemplar]

Sol. Try Yourself [1]

Q. 20. Write the projection of vector  $\vec{a} - \vec{b}$  on the vector  $\vec{a} + \vec{b}$ . [All India 2011]

Sol. Try Yourself [1]

Q. 21. If  $\hat{p}$  is a unit vector and  $(\vec{x} - \hat{p}) \cdot \hat{p} = 80$ , then find  $|\vec{x}|$ . [U] [All India 2009]

Sol. Try Yourself [1]

Q. 22. Write the angle between vectors  $\vec{a}$  and  $\vec{b}$  with magnitudes  $\sqrt{3}$  and 2 respectively, having  $|\vec{a} + \vec{b}| = \sqrt{6}$ . [U] [All India 2011]

Sol. Try Yourself [1]

Q. 23. If  $|\vec{a}| = \sqrt{3}, |\vec{b}| = 2$  and angle between  $\vec{a}$  and  $\vec{b}$  is  $60^\circ$ , then find  $|\vec{a} + \vec{b}|$ . [U] [Delhi 2011C]

Sol. Try Yourself [1]

Q. 24. Find  $\vec{a} \cdot \vec{b}$  if  $\vec{a} = \hat{i} - \hat{j}$  and  $\vec{b} = 2\hat{i} + 3\hat{j} - 3\hat{k}$ . [U] [Delhi 2009C]

Sol. Try Yourself [1]

Q. 25. Find the value of  $\lambda$ , if the vectors  $\hat{i} + \lambda\hat{j} + \hat{k}$  and  $3\hat{i} + 2\hat{j} - 4\hat{k}$  are perpendicular to each other.

[R&U] [All India 2010C]  
Sol. Try Yourself [1]

**Q. 26.** Find the projection of  $\vec{a}$  on  $\vec{b}$ , if  $|\vec{a}| = 8$  and

$$\vec{a} = 2\hat{i} + 6\hat{j} + 3\hat{k} \quad \text{R\&U [Delhi 2009]}$$

**Sol.** Try Yourself 1

**Q. 27.** Find the magnitude of each of the vectors  $\vec{a}$  and  $\vec{b}$ , having the same magnitude such that the angle between them is  $60^\circ$  and their scalar product is  $\frac{9}{2}$ .

R&U [Delhi/O.D. 2018]

**Sol.**  $|\vec{a}| = |\vec{b}| = 3$   $\frac{1}{2} + \frac{1}{2}$   
 [CBSE Marking Scheme, 2018]

**Detailed Solution :**

Given, angle  $\theta = 60^\circ$  and  $\vec{a} \cdot \vec{b} = \frac{9}{2}$

$$\therefore \cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|}$$

$$\Rightarrow \cos 60^\circ = \frac{\frac{9}{2}}{|\vec{a}| |\vec{b}|} \quad (\because |\vec{a}| = |\vec{b}|)$$

$$\Rightarrow \frac{1}{2} |\vec{a}|^2 = \frac{9}{2} \Rightarrow |\vec{a}| = 3.$$

$$\therefore |\vec{a}| = |\vec{b}| = 3. \quad \frac{1}{2}$$

## Short Answer Type Questions

(2 marks each)

**Q. 1.** If either vector  $\vec{a}$  or  $\vec{b}$  then  $\vec{a} \cdot \vec{b} = 0$ . But the converse need not be true. Justify your answer.

□ [NCERT] [Foreign 2011]

**Sol.** Let  $\vec{a} \cdot \vec{b} = 0$

$$\text{or } |\vec{a}| |\vec{b}| \cos \theta = 0$$

$$\text{as } \cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} \quad \text{1}$$

$$\text{Either } \vec{a} = 0 \text{ or } \vec{b} = 0$$

$$\text{or } \cos \theta = 0$$

$$\theta = 90^\circ \quad \text{1}$$

thus, it is clear that the dot product of two non-zero perpendicular vector is always zero. This shows that the converse is not true.

**Q. 2.** If  $\vec{a}$  and  $\vec{b}$  are two vectors such that  $|\vec{a} + \vec{b}| = |\vec{a}|$ ,

then prove that  $\vec{b}$  is perpendicular to  $\vec{a}$ .

□ [Delhi Set I 2013]

**Sol.** Given,  $|\vec{a} + \vec{b}| = |\vec{a}|$

$$|\vec{a} + \vec{b}|^2 = |\vec{a}|^2$$

$$\text{or } |\vec{a}|^2 + |\vec{b}|^2 + 2\vec{a} \cdot \vec{b} = |\vec{a}|^2$$

$$|\vec{b}|^2 + 2\vec{a} \cdot \vec{b} = 0 \quad \dots(i)$$

Now,  $2\vec{a} + \vec{b} \cdot \vec{b} = 2\vec{a} \cdot \vec{b} + |\vec{b}|^2$  1

$$\vec{b} \cdot (2\vec{a} + \vec{b}) = 0 \quad \text{1}$$

From eqn. (i),  $2\vec{a} + \vec{b}$  is  $\perp$  to  $\vec{b}$

[CBSE Marking Scheme, 2013]

**Q. 3.** Find the projection (vector) of  $2\hat{i} - \hat{j} + \hat{k}$  on

$$\hat{i} - 2\hat{j} + \hat{k}. \quad \text{R\&U [SQP 2017-18]}$$

**Sol.**  $\vec{a} = 2\hat{i} - \hat{j} + \hat{k}$   $\vec{b} = \hat{i} - 2\hat{j} + \hat{k}$   $\therefore |\vec{a}| = 5, |\vec{b}| = \sqrt{6}$   $\frac{1}{2}$

The required projection (vector) of  $\vec{a}$  on  $\vec{b}$

$$= \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|^2} \vec{b} \quad \text{1}$$

$$= \frac{5}{6} (\hat{i} - 2\hat{j} + \hat{k}) \quad \frac{1}{2}$$

[CBSE Marking Scheme, 2017]

**Q. 4.** If  $\theta$  is the angle between two vectors  $\hat{i} - \hat{j} + \hat{k}$

and  $3\hat{i} - 2\hat{j} + \hat{k}$  find  $\sin \theta$ . R&U [Delhi/O.D.-2018]

**Sol.**  $\sin \theta = \frac{|\hat{i} - 2\hat{j} + 3\hat{k} \times 3\hat{i} - 2\hat{j} + \hat{k}|}{|\hat{i} - 2\hat{j} + \hat{k}| |3\hat{i} - 2\hat{j} + \hat{k}|} \quad \frac{1}{2}$

$$\hat{i} - 2\hat{j} + 3\hat{k} \times 3\hat{i} - 2\hat{j} + \hat{k} = 4\hat{i} + 8\hat{j} + \hat{k} = \sqrt{6}$$

$$\sin \theta = \frac{4\sqrt{6}}{\sqrt{6} \cdot \sqrt{6}} = \frac{4}{3} \quad \frac{1}{2}$$

[CBSE Marking Scheme, 2018]

Detailed Answer :

$$\begin{aligned} \text{Let } \vec{a} &= \hat{i} - 2\hat{j} + 3\hat{k} \Rightarrow a = \sqrt{1 + (2)^2 + 3^2} = \sqrt{14} \\ \vec{b} &= 3\hat{i} - 2\hat{j} + \hat{k} \Rightarrow b = \sqrt{3^2 + (2)^2 + 1^2} = \sqrt{14} \\ \therefore \cos \theta &= \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} \\ &= \frac{(\hat{i} - 2\hat{j} + 3\hat{k}) \cdot (3\hat{i} - 2\hat{j} + \hat{k})}{\sqrt{14} \sqrt{14}} = 1 \end{aligned}$$

$$= \frac{3 \times 3 + (-2) \times (-2) + 3 \times 1}{14} = \frac{10}{14} = \frac{5}{7}$$

$$\therefore \sin \theta = \sqrt{1 - \cos^2 \theta}$$

$$\sin \theta = \sqrt{1 - \frac{25}{49}}$$

$$= \frac{\sqrt{24}}{7} = \frac{\sqrt{6}}{7}$$

1

### Long Answer Type Questions-I

(4 marks each)

Q.1. Find the vector  $\vec{p}$  which is perpendicular to both

$$\vec{\alpha} = 4\hat{i} + 5\hat{j} - k \text{ and } \vec{\beta} = \hat{i} - 4\hat{j} + 5\hat{k} \text{ and } \vec{p} \cdot \vec{q} = 21,$$

$$\text{where } \vec{q} = 3\hat{i} - \hat{j} - \hat{k}$$

[O.D. Set I, II, III Comptt. 2014]

Sol. Any vector perpendicular to both  $\vec{\alpha}$  and  $\vec{\beta}$

Parallel to  $(\vec{\alpha} \times \vec{\beta})$

$$\therefore \vec{p} = \lambda (\vec{\alpha} \times \vec{\beta})$$

$$= \lambda \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 4 & 5 & -1 \\ 1 & -4 & 5 \end{vmatrix}$$

$$= \lambda [i(25 - 4) - j(20 + 1) + k(-16 - 5)]$$

$$= \lambda (21\hat{i} - 21\hat{j} - 21\hat{k})$$

$$= 21 \quad \text{(Given)}$$

$$\lambda (21\hat{i} - 21\hat{j} - 21\hat{k}) \cdot (3\hat{i} - \hat{j} - \hat{k}) = 21$$

$$\lambda (63 - 21 + 21) = 21$$

$$\lambda = \frac{1}{7}$$

$$\therefore \vec{p} = \frac{1}{7} (21\hat{i} - 21\hat{j} - 21\hat{k})$$

$$\vec{p} = 3\hat{i} - 3\hat{j} - 3\hat{k}$$

$$\vec{p} = 3(\hat{i} - \hat{j} - \hat{k})$$

[CBSE Marking Scheme 2014]

Q.2. Find the unit vector perpendicular to the plane ABC where the position vectors A, B and C are

$$2\hat{i} - \hat{j} + \hat{k}, \hat{i} + 2\hat{j} + \hat{k} \text{ and } 2\hat{i} + \hat{j} + \hat{k}$$

[O.D. Set I, II, III Comptt. 2014]

Sol. Required vector =  $\frac{AB \times AC}{|AB \times AC|}$  1

$$\text{or } = (\text{Position vector of } B) - (\text{Position vector of } A)$$

$$\text{or } = -\hat{i} + 2\hat{j} + \hat{k}$$

Similarly

$$\vec{p} = \hat{i} + \hat{j} + \hat{k} \quad 1$$

$$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -1 & 2 & 1 \\ 0 & 1 & 2 \end{vmatrix}$$

$$= i(4 - 1) - j(-2 - 0) + k(-2 - 0)$$

$$AB \times AC = 3\hat{i} - 2\hat{j} - 2\hat{k} \quad 1$$

\therefore Required unit vector

$$= \frac{1}{\sqrt{17}} (3\hat{i} - 2\hat{j} - 2\hat{k}) \quad 1$$

[CBSE Marking Scheme 2014]

#### Commonly Made Error

- Sometimes, candidates use cross product without evaluating  $\vec{a} \times \vec{b}$  and  $|\vec{a} \times \vec{b}|$ . Some candidates make mistakes while evaluating the unit vector in the final answer.

#### Answering Tip

- Vector algebra in finding unit vector need to be understood by the students.

Q.3. If  $\vec{a} = 7\hat{i} - \hat{j} + \hat{k}$  and  $\vec{b} = 5\hat{i} - \hat{j} + \lambda\hat{k}$  then find

the value of  $\lambda$  so that  $\vec{a}$  and  $\vec{b}$  are perpendicular vectors. [O.D. Set I 2013]



**Sol.** Given ,

$$\vec{a} = i - j + 7k$$

and  $\vec{b} = 5i - j + \lambda k$

$$\vec{a} + \vec{b} = (\hat{i} - \hat{j} + 7\hat{k}) + 5\hat{i} - \hat{j} + \lambda\hat{k} \quad 1$$

$$= 6\hat{i} - 2\hat{j} + 7\hat{k} + \lambda\hat{k}$$

and  $\vec{a} - \vec{b} = (\hat{i} - \hat{j} + 7\hat{k}) - 5\hat{i} + \hat{j} - \lambda\hat{k}$

$$= -4\hat{i} + 7\hat{k} - \lambda\hat{k} \quad 1$$

Since  $(\vec{a} + \vec{b})$  and  $(\vec{a} - \vec{b})$  are perpendicular vectors,

$$\therefore (\vec{a} + \vec{b}) \cdot (\vec{a} - \vec{b}) = 0$$

$$\{6\hat{i} - 2\hat{j} + (7 + \lambda)\hat{k}\} \cdot \{-4\hat{i} + (7 - \lambda)\hat{k}\} = 0$$

$$\text{or } -24 + (7 + \lambda)(7 - \lambda) = 0 \quad 1$$

$$\text{or } 49 - \lambda^2 - 24 = 0$$

$$\text{or } \lambda^2 = 49 - 24 = 25$$

$$\text{or } \lambda = \pm 5 \text{ units} \quad 1$$

[CBSE Marking Scheme, 2013]

**Q. 4.** Vectors  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are such that  $\vec{a} + \vec{b} + \vec{c}$

and  $|\vec{a}| = 3, |\vec{b}| = 5$  and  $|\vec{c}| = 7$ . Find the angle between  $\vec{a}$  and  $\vec{b}$ . [R&U] [Delhi Set I, II, III, 2014]

**Sol.**  $\vec{a} + \vec{b} + \vec{c} = \vec{0} \Rightarrow \vec{a} + \vec{b} = -\vec{c}$   $\therefore |\vec{a} + \vec{b}| = |\vec{c}|$   $\frac{1}{2}$

$$\text{or } \sqrt{3^2 + 5^2 + 2|\vec{a}||\vec{b}|\cos\theta} = 7 \quad \frac{1}{2}$$

$$\text{or } 9 + 25 + 2|\vec{a}||\vec{b}|\cos\theta = 49 \quad 1$$

$\theta$  being angle between  $\vec{a}$  and  $\vec{b}$ ,  $\frac{1}{2}$

$$\therefore \cos\theta = \frac{49 - 34}{2 \times 3 \times 5} = -\frac{1}{5} \text{ or } \theta = \cos^{-1}\left(-\frac{1}{5}\right) \quad 1$$

[CBSE Marking Scheme 2014]

**Q. 5.** The scalar product of the vector  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$  with a unit vector along the sum of vectors  $\vec{b} = 2\hat{i} + 4\hat{j} - 5\hat{k}$  and  $\vec{c} = \lambda\hat{i} + 2\hat{j} + 3\hat{k}$  is equal to one. Find the value of  $\lambda$  and hence find the unit vector along  $\vec{b} \cdot \vec{c}$ . [U] [OD 2009]

[O.D. Set I, II, III, 2014]

**Sol.** Given that

$$\vec{a} \cdot \frac{\vec{b} + \vec{c}}{|\vec{b} + \vec{c}|} = 1 \quad \frac{1}{2}$$

or  $\vec{a} \cdot (\vec{b} + \vec{c}) = |\vec{b} + \vec{c}| \quad \frac{1}{2}$

$$(\hat{i} + \hat{j} + \hat{k}) \cdot (2\hat{i} + 4\hat{j} - 5\hat{k}) + (\lambda\hat{i} + 2\hat{j} + 3\hat{k}) \cdot (\lambda\hat{i} + 2\hat{j} + 3\hat{k}) = \sqrt{(2 + \lambda)^2 + (4 + 2)^2 + (-5 + 3)^2}$$

$$= \sqrt{(\lambda + 2)^2 + 6^2 + (-2)^2} \quad \frac{1}{2}$$

$$\text{or } (2 + \lambda + 5) + (\lambda + 2 + 3) = \sqrt{(\lambda + 2)^2 + 36 + 4} \quad 1$$

$$\therefore (\lambda + 6)^2 = (\lambda + 2)^2 + 40 \text{ or } \lambda = 1 \quad \frac{1}{2}$$

Hence  $\frac{\vec{b} + \vec{c}}{|\vec{b} + \vec{c}|} = \frac{3\hat{i} + 6\hat{j} - 2\hat{k}}{\sqrt{3^2 + 6^2 + (-2)^2}}$

$$= \frac{3\hat{i} + 6\hat{j} - 2\hat{k}}{7} \quad 1$$

[CBSE Marking Scheme 2014]

**Alternative Method :**

Let  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$

and  $\vec{b} + \vec{c} = (2\hat{i} + 4\hat{j} - 5\hat{k}) + (\lambda\hat{i} + 2\hat{j} + 3\hat{k})$

$$= (\lambda + 2)\hat{i} + 6\hat{j} - 2\hat{k}$$

So,  $\frac{\vec{b} + \vec{c}}{|\vec{b} + \vec{c}|} = \frac{(\lambda + 2)\hat{i} + 6\hat{j} - 2\hat{k}}{\sqrt{(\lambda + 2)^2 + 6^2 + (-2)^2}}$

$$= \frac{(\lambda + 2)\hat{i} + 6\hat{j} - 2\hat{k}}{\sqrt{\lambda^2 + 4\lambda + 44}} = r, \text{ (say) } \dots(i) \quad \frac{1}{2}$$

$\therefore$  Unit vector along  $\vec{b} + \vec{c}$  is given by :

$$\frac{\vec{b} + \vec{c}}{|\vec{b} + \vec{c}|} = \frac{(\lambda + 2)\hat{i} + 6\hat{j} - 2\hat{k}}{\sqrt{\lambda^2 + 4\lambda + 44}} \quad \frac{1}{2}$$

Since,  $\left(\frac{\vec{b} + \vec{c}}{|\vec{b} + \vec{c}|}\right)^2 = 1$

$$\text{or } (\hat{i} + \hat{j} + \hat{k}) \cdot \left(\frac{(\lambda + 2)\hat{i} + 6\hat{j} - 2\hat{k}}{\sqrt{\lambda^2 + 4\lambda + 44}}\right) = 1 \quad \frac{1}{2}$$

$$\text{or } 1\left(\frac{\lambda + 2}{\sqrt{\lambda^2 + 4\lambda + 44}}\right) + 1\left(\frac{6}{\sqrt{\lambda^2 + 4\lambda + 44}}\right) + 1\left(\frac{-2}{\sqrt{\lambda^2 + 4\lambda + 44}}\right) = 1 \quad \frac{1}{2}$$

or  $\frac{\lambda + 2 + 6 - 2}{\sqrt{\lambda^2 + 4\lambda + 44}} = 1 \Rightarrow \lambda + 6 = \sqrt{\lambda^2 + 4\lambda + 44}$

or  $\lambda + 6 = \sqrt{\lambda^2 + 4\lambda + 44}$ , from (i)  $\frac{1}{2}$

or  $\lambda^2 + 12\lambda + 36 = \lambda^2 + 4\lambda + 44$

or  $8\lambda = 8$

$\therefore \lambda = 1 \Rightarrow r = 7 \quad \frac{1}{2}$

Hence,  $\frac{\vec{b} + \vec{c}}{|\vec{b} + \vec{c}|} = \frac{3\hat{i} + 6\hat{j} - 2\hat{k}}{7}$

$$= \frac{3\hat{i} + 6\hat{j} - 2\hat{k}}{7} \quad 1$$

**Q. 6.** Dot product of a vector with vectors  $\hat{i} + \hat{j} + \hat{k}$ ,  $2\hat{i} + 3\hat{j} + \hat{k}$  and  $\hat{i} + \hat{j} + \hat{k}$  are respectively 4, 0 and 2. Find the vector.

[Delhi Set I, II, III Comptt. 2013]

**Sol.** Let the required vector be  $\vec{r} = xi + yj + zk$  1/2

Also, let  $\vec{r} = \hat{i} - \hat{j} + \hat{k}$ ,

$\vec{r} = \hat{i} + \hat{j} - \hat{k}$

and  $\vec{r} = \hat{i} + \hat{j} + \hat{k}$

Given,  $x - y + z = 4$  ... (i) 1/2

$2x + y - 3z = 0$  ... (ii) 1/2

and  $x + y + z = 2$  ... (iii) 1/2

By solving eqns. (i), (ii), & (iii), we get  $x = 2, y = -1, z = 1$  1 1/2

$\therefore$  The req. vector is  $\vec{r} = 2\hat{i} - \hat{j} + \hat{k}$  1/2

[CBSE Marking Scheme 2013]

**Answering Tip**

- Generally students commit errors in simplifying equation which leads to get the wrong result.

**Q. 7.** Find the values of  $\lambda$  for which the angle between the vectors  $\vec{a} = 2\hat{i} + 4\hat{j} + \lambda\hat{k}$  and  $\vec{b} = 7\hat{i} - 2\hat{j} + \lambda\hat{k}$  is obtuse. [O.D. Set I, II, III Comptt. 2013]

**Sol.** Here,  $\vec{a} = 2\hat{i} + 4\hat{j} + \lambda\hat{k}$

and  $\vec{b} = 7\hat{i} - 2\hat{j} + \lambda\hat{k}$

If  $\theta$  is the angle between the vectors  $\vec{a}$  and  $\vec{b}$ , then

$$\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta$$

Or  $\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|}$  1/2

for  $\theta$  to be obtuse

$$\cos \theta < 0 \text{ or } \vec{a} \cdot \vec{b} < 0$$

or  $2 \cdot 7 + 4 \cdot (-2) + \lambda \cdot \lambda < 0$  1

or  $14\lambda^2 - 8\lambda + \lambda < 0$

or  $14\lambda^2 - 7\lambda < 0$

or  $2\lambda^2 - \lambda < 0$

or  $\lambda(2\lambda - 1) < 0$  1/2

$\therefore \lambda \in \left(0, \frac{1}{2}\right)$  1

[CBSE Marking Scheme 2013]

**Q. 8.** If the sum of two unit vectors  $\vec{a}$  and  $\vec{b}$  is a unit vector, then show that the magnitude of their difference is  $\sqrt{3}$ . [Delhi Set I, II, III Comptt. 2012]

**Sol.** Let,  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$

or  $|\vec{c}| = \sqrt{|a|^2 + |b|^2 + 2|a||b|\cos\theta}$  1

Given that  $\vec{a}$  and  $\vec{b}$  are unit vectors.

So,  $|a| = |b| = |c| = 1$

or  $2|a||b|\cos\theta = -1$  1

or  $\cos\theta = -\frac{1}{2}$

$|\vec{d}| = \sqrt{|a|^2 + |b|^2 - 2|a||b|\cos\theta}$

$= \sqrt{1 + 1 - 2 \cdot 1 \cdot 1 \cdot (-\frac{1}{2})}$  1

$\therefore |\vec{d}| = \sqrt{3}$  1

[CBSE Marking Scheme 2012]

**Q. 9.** If  $a, b, c$  are three vectors such that  $|a| = 5, |b| = 12$  and  $|c| = 13$  and  $a + b + c = 0$ , find the value of  $\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}$ . [Delhi Set I, II, III, 2012]

**Sol.**  $\vec{a} + \vec{b} + \vec{c} = 0$

or  $(\vec{a} + \vec{b} + \vec{c}) \cdot (\vec{a} + \vec{b} + \vec{c}) = 0$  1/2

or  $|\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = 0$  1

or  $25 + 144 + 169 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = 0$  1

$\therefore \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a} = -\frac{1}{2} [25 + 144 + 169]$  1/2

$= -169$  1

[CBSE Marking Scheme 2012]

**Q. 10.** If  $\vec{a}, \vec{b}$  and  $\vec{c}$  are three vectors such that  $|\vec{a}| = 3, |\vec{b}| = 4$  and  $|\vec{c}| = 5$  and each one of them is perpendicular to the sum of the other two, then find  $\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}$ . [O.D. Comptt. 2011, 2010]  
[O.D. Set I, II, III Comptt. 2013]

**Sol.** Since  $\vec{a} \perp (\vec{b} + \vec{c})$ , therefore  $\vec{a} \cdot (\vec{b} + \vec{c}) = 0$

or  $\vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c} = 0$

or  $\vec{a} \cdot (\vec{a} + \vec{b} + \vec{c}) = 3^2 = 9$  ... (i)  $\frac{1}{2}$

or  $\vec{b} \cdot (\vec{a} + \vec{b} + \vec{c}) = 4^2 = 16$  ... (ii)  $\frac{1}{2}$

Similarly,  $\vec{c} \cdot (\vec{a} + \vec{b} + \vec{c}) = 5^2 = 25$  ... (iii)  $\frac{1}{2}$

Adding eqn. (i), (ii) and (iii), we get

$$\vec{a} \cdot (\vec{a} + \vec{b} + \vec{c}) + \vec{b} \cdot (\vec{a} + \vec{b} + \vec{c}) + \vec{c} \cdot (\vec{a} + \vec{b} + \vec{c}) = 50$$

or  $(\vec{a} + \vec{b} + \vec{c}) \cdot (\vec{a} + \vec{b} + \vec{c}) = 50$   $\frac{1}{2}$

or  $|\vec{a} + \vec{b} + \vec{c}|^2 = 50$   $\frac{1}{2}$

or  $|\vec{a} + \vec{b} + \vec{c}| = \sqrt{50} = 5\sqrt{2}$   $\frac{1}{2}$

[CBSE Marking Scheme 2013]

**Q. 11.** Find the angle between the vectors  $\vec{a} - \vec{b}$  and  $\vec{a} + \vec{b}$ .

[R&U] [NCERT Exemplar]

[Outside Delhi Set-II, 2015]

**Sol.**

$$\vec{a} - \vec{b} = \sqrt{2}(\hat{i} - \hat{j})$$

$$\vec{a} + \vec{b} = \sqrt{2}(\hat{i} + \hat{j})$$

$$\therefore \cos \theta = \frac{(\vec{a} - \vec{b}) \cdot (\vec{a} + \vec{b})}{|\vec{a} - \vec{b}| |\vec{a} + \vec{b}|}$$

$$= \frac{(\hat{i} - \hat{j}) \cdot (\hat{i} + \hat{j})}{\sqrt{2} \times \sqrt{2}}$$

$$= \frac{1 - 1}{2} = 0$$

$$\therefore \theta = \frac{\pi}{2}$$

$\therefore$  angle between vectors is  $\frac{\pi}{2}$

[CBSE Marking Scheme 2015]

**Q.12.** If  $a, b, c$  are mutually perpendicular vectors

of equal magnitudes, show that the vector  $\vec{a} + \vec{b} + \vec{c}$  is equally inclined to  $a, b$  and  $c$ . Also, find the angle which  $\vec{a} + \vec{b} + \vec{c}$  makes with  $\vec{a}$  or  $\vec{b}$  or  $\vec{c}$ .

[Delhi 2017]

**Sol.**  $\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{c} = \vec{c} \cdot \vec{a} = 0$  ... (i)

Let  $\alpha, \beta$  and  $\gamma$  be the angles made by  $\vec{a} + \vec{b} + \vec{c}$  with  $\vec{a}, \vec{b}$  and  $\vec{c}$  respectively

$$\cos \alpha = \frac{(\vec{a} + \vec{b} + \vec{c}) \cdot \vec{a}}{|\vec{a} + \vec{b} + \vec{c}| |\vec{a}|}$$

or  $\alpha = \cos^{-1} \left( \frac{|\vec{a}|}{|\vec{a} + \vec{b} + \vec{c}|} \right)$

Similarly,  $\beta = \cos^{-1} \left( \frac{|\vec{b}|}{|\vec{a} + \vec{b} + \vec{c}|} \right)$  and  $\gamma = \cos^{-1} \left( \frac{|\vec{c}|}{|\vec{a} + \vec{b} + \vec{c}|} \right)$  **1**

using (i), we get  $\alpha = \beta = \gamma$   $\frac{1}{2}$

Now

$$|\vec{a} + \vec{b} + \vec{c}|^2 = |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a})$$

or  $|\vec{a} + \vec{b} + \vec{c}|^2 = 3|\vec{a}|^2$  (using (i)) **1**

or  $|\vec{a} + \vec{b} + \vec{c}| = \sqrt{3} |\vec{a}|$

$\therefore \alpha = \cos^{-1} \left( \frac{1}{\sqrt{3}} \right) = \beta = \gamma$   $\frac{1}{2}$

[CBSE Marking Scheme, 2017]

**Q. 13.** If  $a, b$  and  $c$  are mutually perpendicular vectors of equal magnitudes, find the angles which the vector  $\vec{a} + \vec{b} + \vec{c}$  makes with the vectors  $\vec{a}, \vec{b}$  and  $\vec{c}$ .

[O.D. Comptt. 2017]

**Sol.** Let the vector  $\vec{r} = \vec{a} + \vec{b} + \vec{c}$  makes angles  $\alpha, \beta, \gamma$  respectively with the vector  $a, b, c$

Given that  $\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{c} = \vec{c} \cdot \vec{a} = 0$  **1**

$$\cos \alpha = \frac{(\vec{a} + \vec{b} + \vec{c}) \cdot \vec{a}}{|\vec{a} + \vec{b} + \vec{c}| |\vec{a}|}$$

$$= \frac{|\vec{a}|^2}{3|\vec{a}| |\vec{a}|} = \frac{1}{3}$$

or  $\alpha = \cos^{-1} \frac{1}{3}$  **1**

$$\cos \beta = \frac{(\vec{a} + \vec{b} + \vec{c}) \cdot \vec{b}}{|\vec{a} + \vec{b} + \vec{c}| |\vec{b}|} = \frac{|\vec{b}|^2}{3|\vec{b}| |\vec{b}|} = \frac{1}{3}$$

or  $\beta = \cos^{-1} \frac{1}{3}$  **1**

$$\cos \gamma = \frac{(\vec{a} + \vec{b} + \vec{c}) \cdot \vec{c}}{|\vec{a} + \vec{b} + \vec{c}| |\vec{c}|} = \frac{|\vec{c}|^2}{3|\vec{c}| |\vec{c}|} = \frac{1}{3}$$

or  $\gamma = \cos^{-1} \frac{1}{3}$  **1**

$$= \frac{2|\vec{c}|^2}{3|\vec{c}|} = \frac{2}{3}$$

or  $\gamma = \cos^{-1} \frac{2}{3}$  1/2

[CBSE Marking Scheme, 2017]

**Q. 14.** If  $\vec{a}$  and  $\vec{b}$  are two vectors, such that  $|\vec{a}| = 2$ ,  $|\vec{b}| = 1$  and  $\vec{a} \cdot \vec{b} = 1$ , then find  $(3\vec{a} - 5\vec{b}) \cdot (3\vec{a} + 5\vec{b})$ . R& [Delhi 2011]

**Sol.** Given,  $|\vec{a}| = 2$ ,  $|\vec{b}| = 1$  and  $\vec{a} \cdot \vec{b} = 1$

Now,  $(3\vec{a} - 5\vec{b}) \cdot (3\vec{a} + 5\vec{b})$

$$= 6\vec{a} \cdot \vec{a} + 21\vec{a} \cdot \vec{b} - 15\vec{b} \cdot \vec{a} - 25\vec{b} \cdot \vec{b}$$

$$= 6|\vec{a}|^2 + 21\vec{a} \cdot \vec{b} - 15\vec{b} \cdot \vec{a} - 25$$

1

$[\because \vec{x} \cdot \vec{x} = |\vec{x}|^2 \text{ and } \vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{a}]$

$$= 6(2)^2 + 11\vec{a} \cdot \vec{b} - 35$$

$$= 6(2)^2 + 11(1) - 35(1)^2$$

$$= 24 + 11 - 35 = 0$$

1

$[\because |\vec{a}| = 2 \text{ and } |\vec{b}| = 1]$

Hence,  $(3\vec{a} - 5\vec{b}) \cdot (3\vec{a} + 5\vec{b}) = 0$  2

[CBSE Marking Scheme 2011]

**Q. 15.** If vectors  $\vec{c} = 2\hat{i} + 2\hat{j} + 3\hat{k}$ ,  $\vec{d} = -\hat{i} + 2\hat{j} + \hat{k}$ , and  $\vec{c} + \lambda\vec{d}$  is perpendicular to  $\vec{c}$ , then find the value of  $\lambda$ . U [Foreign 2011; All India 2009C]

**Sol.** Given,  $\vec{c} = 2\hat{i} + 2\hat{j} + 3\hat{k}$ ,  $\vec{d} = -\hat{i} + 2\hat{j} + \hat{k}$

and  $\vec{c} + \lambda\vec{d}$  is perpendicular to  $\vec{c}$

Also,  $(\vec{c} + \lambda\vec{d}) \cdot \vec{c} = 0$  ...(i) 1

$[\because \text{when } \vec{u} \perp \vec{v}, \text{ then } \vec{u} \cdot \vec{v} = 0]$

Now,  $(2\hat{i} + 2\hat{j} + 3\hat{k} + \lambda(-\hat{i} + 2\hat{j} + \hat{k})) \cdot (2\hat{i} + 2\hat{j} + 3\hat{k}) = 0$

or  $(2 - \lambda)\hat{i} + (2 + 2\lambda)\hat{j} + (3 + \lambda)\hat{k} \cdot (2\hat{i} + 2\hat{j} + 3\hat{k}) = 0$  1

Then from Eq. (i), we get

$$[2(2 - \lambda) + 2(2 + 2\lambda) + 3(3 + \lambda)] = 0$$

1

or  $3(2 - \lambda) + 1(2 + 2\lambda) = 0$

or  $8 - \lambda = 0$

$\therefore \lambda = 8$  1

**Q. 16.** If  $\vec{p} = 5\hat{i} + \lambda\hat{j} - 3\hat{k}$  and  $\vec{q} = \hat{i} + 3\hat{j} - 5\hat{k}$ , then find the value of  $\lambda$ , so that  $\vec{p}$  and  $\vec{q}$  are perpendicular vectors. U [All India 2013]

**Sol.** Try Yourself Like Q.3 LATQ-I.

## TOPIC-3 Cross Product


### Revision Notes

1. The cross product of two vectors  $\vec{a}$  and  $\vec{b}$  is defined by,

$$\vec{a} \times \vec{b} = |\vec{a}| |\vec{b}| \sin \theta \hat{n}$$

where  $\theta$  is the angle between the vectors  $\vec{a}$  and  $\vec{b}$ ,  $0 \leq \theta \leq \pi$  and  $\hat{n}$  is a unit vector

perpendicular to both  $\vec{a}$  and  $\vec{b}$ . For better illustration, see figure.



Consider  $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$   $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$

$$\text{then, } \vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} = (a_2b_3 - a_3b_2)\hat{i} - (a_1b_3 - a_3b_1)\hat{j} + (a_1b_2 - a_2b_1)\hat{k}$$



• **Properties/Observations of Cross Product**

$$\Rightarrow \hat{i} \times \hat{i} = |\hat{i}| |\hat{i}| \sin 0 = \vec{0} \text{ or } \hat{i} \times \hat{i} = \vec{0} = \hat{j} \times \hat{j} = \vec{0}$$

$$\Rightarrow \hat{i} \times \hat{j} = |\hat{i}| |\hat{j}| \sin \frac{\pi}{2} = \hat{k} \text{ or } \hat{i} \times \hat{j} = \hat{k}, \hat{j} \times \hat{k} = \hat{i}, \hat{k} \times \hat{i} = \hat{j}$$

$\Rightarrow \vec{a} \times \vec{b}$  is a vector  $\vec{c}$  (say) then this vector  $\vec{c}$  is perpendicular to both the vectors  $\vec{a}$  and  $\vec{b}$ .

$$\Rightarrow \vec{a} \times \vec{b} = \vec{0} \text{ if } \vec{a} \parallel \vec{b} \text{ or } \vec{a} = \vec{0}, \vec{b} = \vec{0}$$

$$\Rightarrow \vec{a} \times \vec{a} = \vec{0}$$

$\Rightarrow \vec{a} \times \vec{b} \neq \vec{b} \times \vec{a}$  (Commutative property does not hold for cross product).

$$\Rightarrow \vec{a} \times (\vec{b} + \vec{c}) = \vec{a} \times \vec{b} + \vec{a} \times \vec{c} \text{ (Left distributive)}$$

$$\Rightarrow (\vec{b} + \vec{c}) \times \vec{a} = \vec{b} \times \vec{a} + \vec{c} \times \vec{a} \text{ (Right distributive)}$$

(Distributive property of the vector product or cross product)

**2. Relationship between Vector product and Scalar product [Lagrange's Identity]**

$$\text{or } |\vec{a} \times \vec{b}|^2 + (\vec{a} \cdot \vec{b})^2 = |\vec{a}|^2 |\vec{b}|^2$$

**3. Cauchy-Schwarz inequality :**

For any two vectors  $\vec{a}$  and  $\vec{b}$ , always have  $|\vec{a} \cdot \vec{b}| \leq |\vec{a}| |\vec{b}|$ .

**Note :**

- If  $\vec{a}$  and  $\vec{b}$  represent the adjacent sides of a triangle, then the area of triangle can be obtained by evaluating  $\frac{1}{2} |\vec{a} \times \vec{b}|$ .
- If  $\vec{a}$  and  $\vec{b}$  represent the adjacent sides of a parallelogram, then the area of parallelogram can be obtained by evaluating  $|\vec{a} \times \vec{b}|$ .
- The area of the parallelogram with diagonals  $\vec{a} + \vec{b}$  and  $\vec{a} - \vec{b}$  is  $|\vec{a} \times \vec{b}|$ .

**Know the Formulae**

$\Rightarrow$  Angle between two vectors  $\vec{a}$  and  $\vec{b}$  in terms of cross-product can be found by the expression given here :

$$\sin \theta = \frac{|\vec{a} \times \vec{b}|}{|\vec{a}| |\vec{b}|} \text{ or } \theta = \sin^{-1} \left( \frac{|\vec{a} \times \vec{b}|}{|\vec{a}| |\vec{b}|} \right)$$

**Theorem**

**Triangle Inequality**

For any two vectors  $\vec{a}$  and  $\vec{b}$ , we always have  $|\vec{a} + \vec{b}| \leq |\vec{a}| + |\vec{b}|$ .

**Proof :** The given inequality holds trivially when either  $\vec{a} = \vec{0}$  or  $\vec{b} = \vec{0}$ , i.e., in such a case  $|\vec{a} + \vec{b}| = |\vec{a}| + |\vec{b}|$ .

So, let us check it for  $\vec{a} \neq \vec{0}$  and  $\vec{b} \neq \vec{0}$ .

Then consider  $|\vec{a} + \vec{b}|^2 = |\vec{a}|^2 + |\vec{b}|^2 + 2 \vec{a} \cdot \vec{b}$ .

or 
$$\vec{a} + \vec{b} \cdot \vec{a} + \vec{b} \cdot \vec{b} = \vec{a} \cdot \vec{a} + \vec{b} \cdot \vec{b} + 2|\vec{a}||\vec{b}|\cos\theta$$

For  $\cos\theta \leq 1$ , we have :  $|\vec{a}||\vec{b}|\cos\theta \leq |\vec{a}||\vec{b}|$

or 
$$\vec{a} \cdot \vec{a} + \vec{b} \cdot \vec{b} + |\vec{a}||\vec{b}|\cos\theta \leq \vec{a} \cdot \vec{a} + \vec{b} \cdot \vec{b} + |\vec{a}||\vec{b}|$$

or 
$$\vec{a} + \vec{b} \cdot \vec{a} + \vec{b} \cdot \vec{b} \leq (|\vec{a}| + |\vec{b}|)^2$$

or 
$$|\vec{a} + \vec{b}| \leq |\vec{a}| + |\vec{b}|$$

Hence proved

## Objective Type Questions

(1 mark each)

Q. 1. The value of  $i.(j \times k) + j.(i \times k) + k.(i \times j)$  is

- (a) 0
- (b) -1
- (c) 1
- (d) 3

[NCERT Misc.]

Ans. Correct option : (c)

Explanation :

$$\begin{aligned} & i.(j \times k) + j.(i \times k) + k.(i \times j) \\ &= \hat{i}\hat{i} + \hat{j}(-\hat{j}) + \hat{k}\hat{k} \\ &= 1 - \hat{j} \cdot \hat{j} + 1 \\ &= 1 - 1 + 1 \end{aligned}$$

Q. 2. If  $\theta$  is the angle between any two vectors  $\vec{a}$  and  $\vec{b}$ , then  $|\vec{a} \cdot \vec{b}|$  is equal to  $|\vec{a}||\vec{b}|\cos\theta$  when  $\theta$  is equal to

- (a) 0
- (b)  $\pi/4$
- (c)  $\pi/2$
- (d)  $\pi$

[NCERT Misc.]

Ans. Correct option : (b)

Explanation :

Let  $\theta$  be the angle between two vectors  $\vec{a}$  and  $\vec{b}$ . Then, without loss of generality,  $\vec{a}$  and  $\vec{b}$  are non-zero vectors, so that  $|\vec{a}|$  and  $|\vec{b}|$  are positive.

$$\begin{aligned} |\vec{a} \cdot \vec{b}| &= |\vec{a}||\vec{b}|\cos\theta \\ \Rightarrow |\vec{a}||\vec{b}|\cos\theta &= |\vec{a}||\vec{b}|\sin\theta \\ \Rightarrow \cos\theta &= \sin\theta \quad [\because |\vec{a}| \text{ and } |\vec{b}| \text{ are positive.}] \\ \Rightarrow \tan\theta &= 1 \\ \Rightarrow \theta &= \frac{\pi}{4} \end{aligned}$$

So that,  $|\vec{a} \cdot \vec{b}| = |\vec{a}||\vec{b}|\cos\theta$  when  $\theta$  is equal to  $\frac{\pi}{4}$ .

Q. 3. Let the vectors  $\vec{a}$  and  $\vec{b}$  be such that  $|\vec{a}| = 3$  and  $|\vec{b}| = \frac{\sqrt{2}}{3}$ , then  $\frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|}$  is a unit vector, if the angle between  $\vec{a}$  and  $\vec{b}$  is

- (a)  $\pi/6$
- (b)  $\pi/4$
- (c)  $\pi/3$
- (d)  $\pi/2$

[NCERT Ex.]

Ans. Correct option : (b)

Explanation :

It is given that  $|\vec{a}| = 3$  and  $|\vec{b}| = \frac{\sqrt{2}}{3}$ .

We know that  $\vec{a} \times \vec{b} = |\vec{a}||\vec{b}|\sin\theta \hat{n}$ , where  $\hat{n}$  is a unit vector perpendicular to both  $\vec{a}$  and  $\vec{b}$  and  $\theta$  is the angle between  $\vec{a}$  and  $\vec{b}$ .

Now,  $\frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|}$  is a unit vector if  $|\vec{a} \times \vec{b}| = |\vec{a}||\vec{b}|\sin\theta$ .

$$\begin{aligned} |\vec{a} \times \vec{b}| &= 1 \\ |\vec{a}||\vec{b}|\sin\theta &= 1 \\ |\vec{a}||\vec{b}|\sin\theta &= 1 \\ \Rightarrow 3 \times \frac{\sqrt{2}}{3} \sin\theta &= 1 \\ \sin\theta &= \frac{1}{\sqrt{2}} \\ \theta &= \frac{\pi}{4} \end{aligned}$$

So that,  $\frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|}$  is a unit vector if the angle between  $\vec{a}$  and  $\vec{b}$  is  $\frac{\pi}{4}$ .

Q. 4. The vectors from origin to the points A and B are  $\vec{a}$  and  $\vec{b}$  respectively, then the area of triangle OAB is

- (a) 340
- (b)  $\sqrt{10}$
- (c)  $\sqrt{5}$
- (d)  $-\sqrt{5}$

[NCERT Exemp.]

Ans. Correct option : (d)

Explanation :

$$\begin{aligned} \text{Area of } \Delta OAB &= \frac{1}{2} |\vec{OA} \times \vec{OB}| \\ &= \frac{1}{2} |(2\hat{i} - 3\hat{j} + 2\hat{k}) \times (2\hat{i} + 3\hat{j} + \hat{k})| \\ &= \frac{1}{2} \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & -3 & 2 \\ 2 & 3 & 1 \end{vmatrix} \\ &= \frac{1}{2} [(2(-3-6) - \hat{j}(2-4) + \hat{k}(6+6))] \\ &= \frac{1}{2} |-9\hat{i} + 2\hat{j} + 12\hat{k}| \end{aligned}$$



Area of  $\Delta OAB$   
 $\frac{1}{2}\sqrt{81 + 144}$   
 $\frac{1}{2}\sqrt{225}$

Q. 5. For any vector  $\vec{a}$ , the value of  $(\vec{a} \times \hat{i})^2 + (\vec{a} \times \hat{j})^2 + (\vec{a} \times \hat{k})^2$

is equal to

- (a)  $a^2$  (b)  $2a^2$   
 (c)  $3a^2$  (d)  $4a^2$

[NCERT Exemp.]

Ans. Correct option : (d)

Explanation :

Let  $a = xi + yj + zk$

$$\vec{a} \times \hat{i} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ x & y & z \\ 1 & 0 & 0 \end{vmatrix}$$

$$= \hat{i}[0] - \hat{j}[-z] + \hat{k}[-y]$$

$$= z\hat{j} - y\hat{k}$$

$$(\vec{a} \times \hat{i})^2 = (z\hat{j} - y\hat{k})^2 = z^2 + y^2$$

Similarly,  $(\vec{a} \times \hat{j})^2 = x^2 + z^2$  and  $(\vec{a} \times \hat{k})^2 = x^2 + y^2$   
 $a^2 \times i^2 + (a \times j)^2 + (a \times k)^2 = y^2 + z^2 + x^2 + z^2 + x^2 + y^2 = 2a^2$

Q. 6. If  $|\vec{a}| = 10$ ,  $|\vec{b}| = 2$  and  $|\vec{a} \times \vec{b}| = 16$ , then the value of  $|\vec{a} \cdot \vec{b}|$  is

- (a) 5 (b) 10  
 (c) 14 (d) 16

[NCERT Exemp.]

Ans. Correct option : (d)

Explanation :

Here,  $a = 10$ ,  $b = 2$  and  $a \cdot b = 2$  [Given]

$$a \cdot b = |a||b|\cos\theta$$

$$12 = 10 \times 2 \cos\theta$$

$$\cos\theta = \frac{12}{20} = \frac{3}{5}$$

$$\sin\theta = \sqrt{1 - \cos^2\theta} = \sqrt{1 - \frac{9}{25}}$$

$$\sin\theta = \pm \frac{4}{5}$$

$$|\vec{a} \times \vec{b}| = |a||b|\sin\theta$$

$$= 10 \times 2 \times \frac{4}{5}$$

$$= 16$$

Q. 7. The vectors  $\lambda\hat{i} + \hat{j} + 2\hat{k}$ ,  $\hat{i} + \lambda\hat{j} + \hat{k}$  and  $2\hat{i} - \hat{j} + \lambda\hat{k}$  are coplanar, if

- (a)  $\lambda = -2$  (b)  $\lambda = 0$   
 (c)  $\lambda = 1$  (d)  $\lambda = -1$

[NCERT Exemp.]

Ans. Correct option : (a)

Explanation :

Let  $\vec{a} = \lambda\hat{i} + \hat{j} + 2\hat{k}$ ,  $\vec{b} = \hat{i} + \lambda\hat{j} + \hat{k}$  and  $\vec{c} = 2\hat{i} - \hat{j} + \lambda\hat{k}$

For  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  to be coplanar,

$$\lambda \begin{vmatrix} 1 & 1 & 2 \\ 1 & \lambda & 1 \\ 2 & -1 & \lambda \end{vmatrix} = 0$$

$$\lambda^3 - \lambda - \lambda - 2 - 2 - 4\lambda = 0$$

$$\lambda^3 - 6\lambda - 4 = 0$$

$$(\lambda + 2)(\lambda^2 - 2\lambda - 2) = 0$$

$$\lambda = -2 \text{ or } \lambda = \frac{2 \pm \sqrt{12}}{2}$$

$$\lambda = -2 \text{ or } \lambda = \frac{2 \pm 2\sqrt{3}}{2} = \pm\sqrt{3}$$

## Very Short Answer Type Questions

(1 mark each)

Q. 1. Write the value of  $(\hat{i} \times \hat{j}) \cdot \hat{k} + \hat{i} \cdot \hat{j}$ .

[O.D. Set I, 2012]

Sol.  $(\hat{i} \times \hat{j}) \cdot \hat{k} + \hat{i} \cdot \hat{j} = \hat{k} \cdot \hat{k} + 0$   
 $= 1 + 0 = 1$   
 [CBSE Marking Scheme 2012]

Q. 2. Write the value of  $(\hat{k} \times \hat{j}) \cdot \hat{i} + \hat{j} \cdot \hat{k}$ .

[O.D. Set II, 2012]

Sol.  $(\hat{k} \times \hat{j}) \cdot \hat{i} + \hat{j} \cdot \hat{k} = -\hat{i} \cdot \hat{i} + 0$   
 $= -1 + 0 = -1$   
 [CBSE Marking Scheme 2012]

Q. 3. Write the value of  $(\hat{k} \times \hat{i}) \cdot \hat{j} + \hat{i} \cdot \hat{k}$ .

[O.D. Set III, 2012]

Sol.  $(\hat{k} \times \hat{i}) \cdot \hat{j} + \hat{i} \cdot \hat{k} = \hat{j} \cdot \hat{j} + 0$   
 $= 1 + 0 = 1$   
 [CBSE Marking Scheme 2012]

Q. 4. If  $\vec{a}$  and  $\vec{b}$  are two vectors of magnitude 3 and 4 respectively such that  $\vec{a} \times \vec{b}$  is a unit vector, write the angle between  $\vec{a}$  &  $\vec{b}$ .

[O.D. 2010] [Delhi Set II, 2014] [S.Q.P. 2012]

Sol. We know  $|\vec{a} \times \vec{b}| = |a||b|\sin\theta$ , where  $\hat{n} = 1$

$$\begin{aligned} \times &= 1 \\ a &= 3 \quad \text{[Given]} \\ \text{and } b &= - \\ \therefore 1 &= 3 \times - \sin \theta \\ \text{or } \sin \theta &= \frac{1}{3} \\ \text{or } \theta &= \dots \quad 1 \end{aligned}$$

[CBSE Marking Scheme 2012]

Q. 5. Vectors  $\vec{a}$  and  $\vec{b}$  are such that  $|\vec{a}| = \sqrt{3}$ ,  $|\vec{b}| = 2$  and  $\vec{a} \times \vec{b}$  is a unit vector. Write the angle between  $\vec{a}$  &  $\vec{b}$ .  [Delhi Set II, 2014]

Sol. Since  $\vec{a} \times \vec{b}$  is a unit vector, therefore

$$|\vec{a} \times \vec{b}| = 1$$

or  $|\vec{a}| |\vec{b}| \sin \theta = 1$

or  $(\sqrt{3}) \left(\frac{2}{3}\right) \sin \theta = 1$

or  $\sin \theta = \frac{\sqrt{3}}{2}$

$\therefore \theta = \dots$  1

[CBSE Marking Scheme 2014]

Q. 6. If  $|\vec{a}| = 8$ ,  $|\vec{b}| = 3$  and  $|\vec{a} \times \vec{b}| = 12$ , find the angle between  $\vec{a}$  and  $\vec{b}$ .  [O.D. Set I, II, III Comptt. 2014]

Sol. We know

$$|\vec{a} \times \vec{b}| = |\vec{a}| |\vec{b}| \sin \theta$$

or  $\sin \theta = \frac{|\vec{a} \times \vec{b}|}{|\vec{a}| |\vec{b}|} = \frac{12}{8 \times 3} = \frac{1}{2}$

$\therefore \theta = \dots$  1

[CBSE Marking Scheme 2014]

Q. 7. For vector  $\vec{a}$ , if  $|\vec{a}| = a$ , then write the value of :

$$\left(\vec{a} \times \hat{i}\right)^2 + \left(\vec{a} \times \hat{j}\right)^2 + \left(\vec{a} \times \hat{k}\right)^2$$

R& [NCERT Exemplar] [Delhi Set I, II, III Comptt. 2016]

Sol. Let  $\vec{a} = xi + yj + zk$

then  $x^2 + y^2 + z^2 = a^2$  (as,  $|\vec{a}| = a$ )

$$\vec{a} \times \hat{i} = -yk + zj,$$

$$\vec{a} \times \hat{j} = -xz + yi$$

and  $\vec{a} \times \hat{k} = -xy + zi$

$$\therefore (\vec{a} \times \hat{i})^2 + (\vec{a} \times \hat{j})^2 + (\vec{a} \times \hat{k})^2 = 2(x^2 + y^2 + z^2) = 2a^2 \quad 1$$

[CBSE Marking Scheme 2016]

Q. 8. Find the direction cosines of the vector joining the points  $A(1, 2, -3)$  and  $B(-1, -2, 1)$  directed from  $B$  to  $A$ .  [Outside Delhi Set I, II, III Comptt. 2016]

Sol.  $\vec{AB} = 2i + 4j - 4k$

or d-ratios of  $\vec{AB}$  are 2, 4, -4

$\therefore$  Direction cosines are :  $\frac{1}{3}, \frac{2}{3}, -\frac{2}{3}$  1

[CBSE Marking Scheme 2016]

Q. 9. Find the angle between two vectors  $\vec{a}$  and  $\vec{b}$  having the same length  $\sqrt{2}$  and their vector product is  $-\hat{i} - \hat{j} + \hat{k}$ .  [Outside Delhi Set I, II, III Comptt. 2016]

Sol.  $\sin \theta = \frac{-i - j + k}{\sqrt{2} \cdot \sqrt{2}}$

$$\sin \theta = \frac{\sqrt{3}}{2}$$

or  $\theta = 60^\circ$

or  $\theta = \dots$  1

[CBSE Marking Scheme 2016]

Q. 10. Find  $\lambda$  and  $\mu$  if  $(\hat{i} + 3\hat{j} + 9\hat{k}) \times (\hat{i} - \lambda\hat{j} + \mu\hat{k}) = 0$ .  [O.D. 2016]

Sol. Getting  $\lambda = -9$

and  $\mu = 27$  1

[CBSE Marking Scheme 2016]

Detailed Solution :

$$(i + 3j + 9k) \times (i - \lambda j + \mu k) = 0$$

or  $\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 3 & 9 \\ 1 & -\lambda & \mu \end{vmatrix} = 0$

$$\begin{aligned} \text{or } \hat{i}(3\mu + 9\lambda) - \hat{j}(\mu - 27) + \hat{k}(-\lambda - 9) &= 0 \\ \text{or } 3\mu + 9\lambda &= 0 \quad \dots(i) \\ \text{or } \mu - 27 &= 0 \quad \dots(ii) \\ \text{or } -\lambda - 9 &= 0 \quad \dots(iii) \end{aligned}$$

from eqn. (ii) and (iii),

$$\begin{aligned} \mu &= 27 \\ \text{and } \lambda &= -9 \end{aligned}$$

**Q. 11.** If  $\vec{a}$  and  $\vec{b}$  are two non-zero vectors such that  $|\vec{a} \times \vec{b}| = a \cdot b$ , then find the angle between  $\vec{a}$  and  $\vec{b}$ . R&U [O.D. 2010] [S.Q.P. 2016]

**Sol.**

$$\begin{aligned} \sin \theta &= \cos \theta \\ \text{or } \theta &= 45^\circ \end{aligned}$$

**1**  
[CBSE Marking Scheme 2016]

**Q. 12.** If vectors  $\vec{a}$  and  $\vec{b}$  are such that  $|\vec{a}| = \frac{1}{2}$ ,  $|\vec{b}| = \frac{4}{\sqrt{3}}$  and  $|\vec{a} \times \vec{b}| = \frac{1}{\sqrt{3}}$ , then find  $|\vec{a} \cdot \vec{b}|$ . R&U [O.D. Set II 2016]

**Sol.**

$$|\vec{a}| = \frac{1}{2} \quad |\vec{b}| = \frac{4}{\sqrt{3}} \quad |\vec{a} \times \vec{b}| = \frac{1}{\sqrt{3}} \quad |\vec{a} \cdot \vec{b}| = ?$$

$$|\vec{a} \times \vec{b}| = |\vec{a}| |\vec{b}| \sin \theta \quad (\theta: \text{Angle between vectors } \vec{a} \text{ \& } \vec{b})$$

$$\frac{1}{\sqrt{3}} = \frac{1}{2} \times \frac{4}{\sqrt{3}} \sin \theta$$

$$\sin \theta = \frac{1}{2} \quad \theta = 30^\circ$$

$$|\vec{a} \cdot \vec{b}| = |\vec{a}| |\vec{b}| \cos \theta$$

$$= \frac{1}{2} \times \frac{4}{\sqrt{3}} \times \frac{\sqrt{3}}{2} = 1$$

$$\therefore |\vec{a} \cdot \vec{b}| = 1$$

**1**  
[Topper's Answer 2016]

**Q. 13.** Write the angle between the vectors  $\vec{a} \times \vec{b}$  and  $\vec{b} \times \vec{a}$ . R&U [Delhi Comptt. 2017]

**Sol.** Angle between  $\vec{a} \times \vec{b}$  and  $\vec{b} \times \vec{a}$  is  $\pi$ . **1**  
[CBSE Marking Scheme 2017]

**Q. 14.** If  $(\vec{a} \times \vec{b})^2 + (\vec{a} \cdot \vec{b})^2 = 225$  and  $|\vec{a}| = 5$ , then write the value of  $|\vec{b}|$ . R&U [O.D. Comptt. 2017]

**Sol.**

$$(\vec{a} \times \vec{b})^2 + (\vec{a} \cdot \vec{b})^2 = 225$$

$$\text{or } |\vec{a}|^2 |\vec{b}|^2 (\sin^2 \theta + \cos^2 \theta) = 225$$

$$\text{or } (5)^2 |\vec{b}|^2 = 225 \text{ or } |\vec{b}| = 3 \quad \mathbf{1}$$

[CBSE Marking Scheme 2017]

**Q. 15.** Write the value of the following.

$$\hat{i} \times (\hat{j} + \hat{k}) + \hat{j} \times (\hat{k} + \hat{i}) + \hat{k} \times (\hat{i} + \hat{j})$$

R&U [Foreign 2014]

**Sol.**

$$\hat{i} \times (\hat{j} + \hat{k}) + \hat{j} \times (\hat{k} + \hat{i}) + \hat{k} \times (\hat{i} + \hat{j})$$

$$= \hat{i} \times \hat{j} + \hat{i} \times \hat{k} + \hat{j} \times \hat{k} + \hat{j} \times \hat{i} + \hat{k} \times \hat{i} + \hat{k} \times \hat{j}$$

$$= \hat{k} - \hat{j} + \hat{i} - \hat{k} + \hat{j} - \hat{i} = \vec{0}$$

$$[\because \hat{i} \times \hat{j} = \hat{k}, \hat{i} \times \hat{k} = -\hat{j}, \hat{j} \times \hat{k} = \hat{i}, \hat{j} \times \hat{i} = -\hat{k}, \hat{k} \times \hat{i} = \hat{j}, \hat{k} \times \hat{j} = -\hat{i}] \quad \mathbf{1}$$

[CBSE Marking Scheme 2014]

**Answering Tip**

- Practice of calculation of vectors should be done properly.

**Q. 16.** Find the angle between  $\vec{a}$  and  $\vec{b}$  with magnitudes 1 and 2 respectively, when  $|\vec{a} \times \vec{b}| = \sqrt{3}$ .

R&U [Delhi 2009]

**Sol.** Given,  $|\vec{a}| = 1, |\vec{b}| = 2$  and  $|\vec{a} \times \vec{b}| = \sqrt{3}$

$$\begin{aligned} \text{or } |\vec{a}||\vec{b}|\sin\theta &= \sqrt{3} \\ [\therefore \vec{a} \times \vec{b} &= |\vec{a}||\vec{b}|\sin\theta \hat{n} \text{ and } |\hat{n}|=1] \\ \text{or } 1 \times 2 \times \sin\theta &= \sqrt{3} \\ \text{or } \sin\theta &= \frac{\sqrt{3}}{2} = \sin\frac{\pi}{3} \text{ or } \theta = \frac{\pi}{3} \end{aligned}$$

Hence, the angle between  $\vec{a}$  and  $\vec{b}$  is  $\frac{\pi}{3}$ . 1

[CBSE Marking Scheme 2009]

**Q. 17.** Write the value of  $p$ , for which  $\vec{a} = 3\hat{i} + 2\hat{j} + 9\hat{k}$  and  $\vec{b} = \hat{i} + p\hat{j} + 3\hat{k}$  are parallel vectors.

[R&U] [Delhi 2009]

**Sol.** Given vectors are  $\vec{a} = 3\hat{i} + 2\hat{j} + 9\hat{k}$  and  $\vec{b} = \hat{i} + p\hat{j} + 3\hat{k}$

Also  $\vec{a}$  and  $\vec{b}$  are parallel vectors.

So,  $\vec{a} \times \vec{b} = 0$

$$\text{or } \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & 2 & 9 \\ 1 & p & 3 \end{vmatrix} = \vec{0}$$

or  $\hat{i}(6-9p) - \hat{j}(9-9) + \hat{k}(3p-2) = \vec{0}$

or  $\hat{i}(6-9p) + \hat{k}(3p-2) = 0\hat{i} + 0\hat{j} + 0\hat{k}$

On comparing the coefficients of  $\hat{i}$  or  $\hat{k}$  form both sides, we get

$$\begin{aligned} 6-9p &= 0 \\ \therefore p &= \frac{2}{3} \end{aligned}$$

1

[CBSE Marking Scheme 2009]

**Alternate Method :**

Since  $\vec{a}$  and  $\vec{b}$  are parallel,  $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$

$$\Rightarrow \frac{3}{1} = \frac{2}{p} = \frac{9}{3} \Rightarrow p = \frac{2}{3}$$

**Q. 18.** If  $|\vec{a} \times \vec{b}|^2 + |\vec{a} \cdot \vec{b}|^2 = 400$  and  $|\vec{a}| = 5$ , then write the value of  $|\vec{b}|$ . [R&U] [Foreign 2016]

**Sol.** Try Yourself

**Q. 19.** Find  $\lambda$ , if

$$(2\hat{i} + 6\hat{j} + 14\hat{k}) \times (\hat{i} - \lambda\hat{j} + 7\hat{k}) = 0$$

[R&U] [All India 2010]

**Sol.** Try Yourself

**Q. 20.** Find the value of  $p$ ,

$$\text{if } (2\hat{i} + 6\hat{j} + 27\hat{k}) \times (\hat{i} + 3\hat{j} + p\hat{k}) = \vec{0}$$

[R&U] [All India 2009]

**Sol.** Try Yourself

#### Commonly Made Error

- Some candidates make mistakes while calculating dot & cross product as vector  $i \cdot i = 1$  and  $i \times i = 0$  is right method but candidates like  $i \times i = 1$  and  $i \cdot i = 0$  which is wrong.

## ? Short Answer Type Questions

(2 marks each)

**Q. 1.** Find  $|\vec{a} \times \vec{b}|$ , if  $\vec{a} = \hat{i} + 2\hat{j} - \hat{k}$ ,  $\vec{b} = 3\hat{i} + \hat{j} - \hat{k}$ .

[R&U]

**Sol.**

$$\begin{aligned} \vec{a} \times \vec{b} &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 2 & -1 \\ 3 & 1 & -1 \end{vmatrix} \\ &= \hat{i}(-2+1) - \hat{j}(-1+3) + \hat{k}(1-6) \\ &= -\hat{i} - 2\hat{j} - 5\hat{k} \end{aligned}$$

$$\begin{aligned} |\vec{a} \times \vec{b}| &= \sqrt{1^2 + 2^2 + 5^2} \\ &= \sqrt{1+4+25} = \sqrt{30} \end{aligned}$$

1

**Q. 2.** Find the area of parallelogram whose adjacent sides are determined by the vector  $\vec{a} = \hat{i} - \hat{j} + 2\hat{k}$  and  $\vec{b} = 2\hat{i} - \hat{j} - \hat{k}$ .

[R&U]

**Sol.**

$$\begin{aligned} \vec{a} \times \vec{b} &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -1 & 2 \\ 2 & -1 & -1 \end{vmatrix} \\ &= \hat{i}(1+2) - \hat{j}(-1-4) + \hat{k}(-1+2) \\ &= 3\hat{i} + 5\hat{j} + \hat{k} \end{aligned}$$

Area of  $|\vec{a} \times \vec{b}| = \sqrt{9+25+1} = \sqrt{35}$  sq. units 1

**Q. 3.** Find  $|\vec{a} \times \vec{b}|$  if  $|\vec{a}| = 10$ ,  $|\vec{b}| = 2$  and  $\vec{a} \cdot \vec{b} = 12$ .

[R&U]

**Sol.**  $\therefore \cos\theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}||\vec{b}|} = \frac{12}{10 \times 2} = \frac{3}{5}$

$$\cos\theta = \frac{3}{5}$$

$$\sin \theta = \sqrt{1 - \cos^2 \theta} = \sqrt{1 - \frac{9}{25}} = \frac{4}{5} \quad 1$$

$$\sin \theta = \frac{|\vec{a} \times \vec{b}|}{|\vec{a}| |\vec{b}|}$$

$$\text{or } |\vec{a} \times \vec{b}| = |\vec{a}| |\vec{b}| \sin \theta$$

$$\text{or } |\vec{a} \times \vec{b}| = 10 \times 2 \times \frac{4}{5} = 16 \quad 1$$

**Q. 4.** Using vectors, find the area of triangle ABC, with vertices A(1, 2, 3), B(2, -1, 4) and C(4, 5, -1).

**R&U** [Foreign 2017]

$$\text{Sol. } \vec{AB} = \hat{i} - 3\hat{j} + \hat{k}, \quad \vec{AC} = 3\hat{i} + 3\hat{j} - 4\hat{k} \quad \frac{1}{2} + \frac{1}{2}$$

$$\text{Area of } \triangle ABC = \frac{1}{2} |\vec{AB} \times \vec{AC}|$$

$$= \frac{1}{2} \text{ magnitude of } \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -3 & 1 \\ 3 & 3 & -4 \end{vmatrix} \quad 1$$

$$= \frac{\sqrt{274}}{2} \text{ sq. units}$$

[CBSE Marking Scheme 2017]

### Answering Tip

- Learn the concept of area of triangle in terms of vector algebra.

**Q. 5.** If  $\vec{a} \times \vec{b} = \vec{c} \times \vec{d}$  and  $\vec{a} \times \vec{c} = \vec{b} \times \vec{d}$ , then show that

$(\vec{a} - \vec{d})$  is parallel to  $(\vec{b} - \vec{c})$ , it is being given that  $\vec{a} \neq \vec{d}$  and  $\vec{b} \neq \vec{c}$ . [Foreign 2016] [Delhi 2009]

**Sol.** Given,

$$\vec{a} \times \vec{b} = \vec{c} \times \vec{d} \quad \text{and} \quad \vec{a} \times \vec{c} = \vec{b} \times \vec{d}$$

$$\text{or } \vec{a} \times \vec{b} - \vec{a} \times \vec{c} = \vec{c} \times \vec{d} - \vec{b} \times \vec{d}$$

$$\text{or } \vec{a} \times (\vec{b} - \vec{c}) + (\vec{b} - \vec{c}) \times \vec{d} = \vec{0} \quad 1$$

[By left and right distributive law]

$$\text{or } \vec{a} \times (\vec{b} - \vec{c}) - \vec{d} \times (\vec{b} - \vec{c}) = \vec{0} \quad [\because \vec{a} \times \vec{b} = -\vec{b} \times \vec{a}]$$

$$\text{or } (\vec{a} - \vec{d}) \times (\vec{b} - \vec{c}) = \vec{0}$$

$$\text{or } (\vec{a} - \vec{d}) \parallel (\vec{b} - \vec{c}) \quad 1$$

[CBSE Marking Scheme 2009]

## ? Long Answer Type Questions-I

(4 marks each)

**Q. 1.** The two adjacent sides of a parallelogram are  $2\hat{i} - 4\hat{j} - 5\hat{k}$  and  $2\hat{i} + 2\hat{j} + 3\hat{k}$ . Find the two unit vectors parallel to its diagonals. Using the diagonal vectors, find the area of the parallelogram.


**R&U** [O.D. Set I, II, III 2016]

$$\text{Sol. } \vec{a} = 2\hat{i} - 4\hat{j} - 5\hat{k}$$

and  $\vec{b} = 2\hat{i} + 2\hat{j} + 3\hat{k}$

$$\therefore \vec{a} + \vec{b} = 4\hat{i} - 2\hat{j} - 2\hat{k}$$

and  $\vec{b} - \vec{a} = 6\hat{j} + 8\hat{k} \quad \frac{1}{2}$



Unit vector parallel to  $\vec{d}_1 = \vec{a} + \vec{b}$  is

$$= \frac{\vec{a} + \vec{b}}{|\vec{a} + \vec{b}|} \quad \frac{1}{2}$$

$$= \frac{4\hat{i} - 2\hat{j} - 2\hat{k}}{\sqrt{16 + 4 + 4}}$$

$$= \frac{4}{\sqrt{24}}\hat{i} - \frac{2}{\sqrt{24}}\hat{j} - \frac{2}{\sqrt{24}}\hat{k}$$

$$= \frac{2}{\sqrt{6}}\hat{i} - \frac{1}{\sqrt{6}}\hat{j} - \frac{1}{\sqrt{6}}\hat{k} \quad \frac{1}{2}$$

Unit vector parallel to  $\vec{d}_2 = \vec{b} - \vec{a}$  is

$$= \frac{\vec{b} - \vec{a}}{|\vec{b} - \vec{a}|} = \frac{6\hat{j} + 8\hat{k}}{\sqrt{36 + 64}}$$

$$= \frac{6}{10}\hat{j} + \frac{8}{10}\hat{k} = \frac{3}{5}\hat{j} + \frac{4}{5}\hat{k} \quad \frac{1}{2}$$

Area of parallelogram  $\frac{1}{2} |\vec{d}_1 \times \vec{d}_2|$

$$\therefore \vec{d}_1 \times \vec{d}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 4 & -2 & -2 \\ 0 & 6 & 8 \end{vmatrix}$$

$$= \hat{i}(-16+12) - \hat{j}(32-0) + \hat{k}(24-0)$$

$$= -4\hat{i} - 32\hat{j} + 24\hat{k}$$

$$|\vec{d}_1 \times \vec{d}_2| = \sqrt{16+1024+576}$$

$$= \sqrt{1,616} \quad \frac{1}{2}$$

∴ Area of parallelogram

$$= \frac{1}{2} |\vec{d}_1 \times \vec{d}_2| \quad \frac{1}{2}$$

$$= \frac{1}{2} \sqrt{1616}$$

$$= \frac{1}{2} \times 4\sqrt{101}$$

$$= 2\sqrt{101}$$

$$= 20.09 \text{ or } 20.1 \text{ sq. units} \quad 1$$

[CBSE Marking Scheme 2016]

Q. 2. If  $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$  find  $(\vec{r} \times \hat{i}) \cdot (\vec{r} \times \hat{j}) + xy$ .

[R&U] [Delhi, 2015]

Sol.  $\vec{r} \times \hat{i} = (x\hat{i} + y\hat{j} + z\hat{k}) \times \hat{i}$   
 $= -y\hat{k} + z\hat{j} \quad 1$

$$\vec{r} \times \hat{j} = (x\hat{i} + y\hat{j} + z\hat{k}) \times \hat{j}$$

$$= x\hat{k} - z\hat{i} \quad 1$$

$$\therefore (\vec{r} \times \hat{i}) \cdot (\vec{r} \times \hat{j})$$

$$= (0\hat{i} + z\hat{j} - y\hat{k}) \cdot (-z\hat{i} + 0\hat{j} + x\hat{k})$$

$$= -xy \quad 1$$

$$\therefore (\vec{r} \times \hat{i}) \cdot (\vec{r} \times \hat{j}) + xy$$

$$= -xy + xy = 0 \quad 1$$

[CBSE Marking Scheme 2015]

Q. 3. If  $\vec{a} = 2\hat{i} - 3\hat{j} + \hat{k}$ ,  $\vec{b} = -\hat{i} + \hat{k}$ ,  $\vec{c} = 2\hat{j} - \hat{k}$  are three vectors, find the area of the parallelogram having diagonals  $\vec{a} + \vec{b}$  and  $\vec{b} + \vec{c}$ .

[R&U] [Delhi Set I Comptt. 2014]

Sol.  $\vec{a} + \vec{b} = \hat{i} - 3\hat{j} + 2\hat{k}$ ;  $\vec{b} + \vec{c} = -\hat{i} + 2\hat{j}$  1

$$(\vec{a} + \vec{b}) \times (\vec{b} + \vec{c})$$

$$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -3 & 2 \\ -1 & 2 & 0 \end{vmatrix} = -4\hat{i} + 2\hat{j} - \hat{k} \quad 1\frac{1}{2}$$

Area of parallelogram

$$= \frac{1}{2} |(\vec{a} + \vec{b}) \times (\vec{b} + \vec{c})|$$

$$= \frac{\sqrt{21}}{2} \text{ sq. units} \quad 1\frac{1}{2}$$

[CBSE Marking Scheme 2014]

Q. 4. Find the unit vector perpendicular to both the vectors  $\vec{a} + \vec{b}$  and  $\vec{a} - \vec{b}$ , where  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$  and  $\vec{b} = \hat{i} + 2\hat{j} + 3\hat{k}$ .

[R&U] [Foreign Set I, II, III 2014]

Sol.  $\vec{a} + \vec{b} = (2\hat{i} + 3\hat{j} + 4\hat{k})$ , 1

and  $\vec{a} - \vec{b} = -\hat{j} - 2\hat{k}$  1/2

Let  $\vec{c} = (\vec{a} + \vec{b}) \times (\vec{a} - \vec{b})$  1

$$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 3 & 4 \\ 0 & -1 & -2 \end{vmatrix} \quad \frac{1}{2}$$

or  $\vec{c} = -2\hat{i} + 4\hat{j} - 2\hat{k}$

or  $\hat{c} = -\frac{1}{\sqrt{6}}\hat{i} + \frac{2}{\sqrt{6}}\hat{j} - \frac{1}{\sqrt{6}}\hat{k}$  1

[CBSE Marking Scheme 2014]

Q. 5. If  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$  and  $\vec{b} = \hat{j} - \hat{k}$ , find a vector  $\vec{c}$ , such that  $\vec{a} \times \vec{c} = \vec{b}$  and  $\vec{a} \cdot \vec{c} = 3$ .

[A] [Delhi Set II, 2013]

Sol. Let  $\vec{c} = x\hat{i} + y\hat{j} + z\hat{k}$

Given  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$

$$\vec{b} = \hat{j} - \hat{k}$$

According to the question,

$$\vec{a} \cdot \vec{c} = 3$$

or  $(\hat{i} + \hat{j} + \hat{k}) \cdot (x\hat{i} + y\hat{j} + z\hat{k}) = 3$  1

or  $x + y + z = 3$  ...(i)

and  $\vec{a} \times \vec{c} = \vec{b}$





$$\text{or } \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & 1 \\ x & y & z \end{vmatrix} = \vec{b} = \hat{j} - \hat{k} \quad 1$$

$$\text{or } (z-y)\hat{i} + (x-z)\hat{j} + (y-x)\hat{k} = \hat{j} - \hat{k}$$

On equating the coefficients of like terms, we get

$$z-y=0, \text{ or } y=z \quad \dots(\text{ii})$$

$$x-z=1 \quad \dots(\text{iii})$$

$$\text{and } y-x=-1 \quad \dots(\text{iv}) \quad 1$$

Solving eqns. (i), (ii), (iii) and (iv), we get

$$x=5/3$$

$$\text{and } y=2/3=z$$

$$\text{Hence, } \vec{c} = \frac{5}{3}\hat{i} + \frac{2}{3}\hat{j} + \frac{2}{3}\hat{k} \quad 1$$

[CBSE Marking Scheme 2013]

**Q. 6.** Using vectors, find the area of the triangle whose vertices are  $A(1, 2, 3)$ ,  $B(2, -1, 4)$  and  $C(4, 5, -1)$ . **R&U** [Delhi 2017] [Delhi Set III, 2013]

OR

Using vectors find the area of triangle ABC with vertices  $A(1, 2, 3)$ ,  $B(2, -1, 4)$  and  $C(4, 5, -1)$ .

**R&U** [Delhi 2017]

**Sol.** Given,  $\vec{OA} = \hat{i} + 2\hat{j} + 3\hat{k}$

$$\vec{OB} = 2\hat{i} - \hat{j} + 4\hat{k}$$

and  $\vec{OC} = 4\hat{i} + 5\hat{j} - \hat{k}$

Now,  $\vec{AB} = \vec{OB} - \vec{OA}$

$$= (2\hat{i} - \hat{j} + 4\hat{k}) - (\hat{i} + 2\hat{j} + 3\hat{k})$$

$$= \hat{i} - 3\hat{j} + \hat{k} \quad \frac{1}{2}$$

and  $\vec{AC} = \vec{OC} - \vec{OA}$

$$= (4\hat{i} + 5\hat{j} - \hat{k}) - (\hat{i} + 2\hat{j} + 3\hat{k})$$

$$= 3\hat{i} + 3\hat{j} - 4\hat{k} \quad \frac{1}{2}$$

$\therefore$  The area of the given triangle

$$= \frac{1}{2} |\vec{AB} \times \vec{AC}| \quad 1$$

Now,  $\vec{AB} \times \vec{AC} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -3 & 1 \\ 3 & 3 & -4 \end{vmatrix}$

$$= \hat{i}(12-3) + \hat{j}(3+4) + \hat{k}(3+9)$$

$$= 9\hat{i} + 7\hat{j} + 12\hat{k} \quad 1$$

Therefore,

$$|\vec{AB} \times \vec{AC}| = \sqrt{(9)^2 + (7)^2 + (12)^2}$$

$$= \sqrt{81 + 49 + 144}$$

$$= \sqrt{274}$$

Hence,

$$\text{required area} = \frac{1}{2} \sqrt{274} \text{ unit}^2 \quad 1$$

[CBSE Marking Scheme 2013]

**Q. 7.** Find the unit vector perpendicular to the plane of  $\triangle ABC$  whose vertices are  $A(3, -1, 2)$ ,  $B(1, -1, -3)$  and  $C(4, -3, 1)$  **R&U** [S.Q.P., 2013]

**Sol.** A vector  $\perp$ , to the plane of  $\triangle ABC$ ,

$$\vec{OA} = 3\hat{i} - \hat{j} + 2\hat{k},$$

$$\vec{OB} = \hat{i} - \hat{j} - 3\hat{k},$$

$$\vec{OC} = 4\hat{i} - 3\hat{j} + \hat{k}$$

$$\vec{AB} = \vec{OB} - \vec{OA} = -2\hat{i} + 0\hat{j} - 5\hat{k}$$

$$\vec{BC} = \vec{OC} - \vec{OB} = 3\hat{i} - 2\hat{j} + 4\hat{k}$$

$$\text{Unit vector perpendicular to plane} = \frac{\vec{AB} \times \vec{BC}}{|\vec{AB} \times \vec{BC}|}$$

$$\vec{AB} \times \vec{BC} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -2 & 0 & -5 \\ 3 & -2 & 4 \end{vmatrix} = -10\hat{i} + 7\hat{j} + 4\hat{k} \quad 1$$

$$|\vec{AB} \times \vec{BC}| = \sqrt{100 + 49 + 16} = \sqrt{165} \quad 1$$

$\therefore$  Unit vector  $\perp$ , to the plane

$$= \frac{1}{\sqrt{165}} (-10\hat{i} + 7\hat{j} + 4\hat{k}). \quad 1$$

[CBSE Marking Scheme 2013]

**Q. 8.** Let  $\vec{a} = \hat{i} + 4\hat{j} + 2\hat{k}$ ,  $\vec{b} = 3\hat{i} - 2\hat{j} + 7\hat{k}$  and  $\vec{c} = 2\hat{i} - \hat{j} + 4\hat{k}$ . Find a vector  $\vec{p}$  which is perpendicular to both  $\vec{a}$  and  $\vec{b}$  and  $\vec{p} \cdot \vec{c} = 18$ .

**R&U** [O.D. Set I, II, III, 2012]

**Sol.**  $\vec{p}$  is  $\perp$  to both  $\vec{a}$  and  $\vec{b}$

$$\text{or } \vec{p} = \lambda(\vec{a} \times \vec{b}) \quad 1\frac{1}{2}$$

$$\text{Now, } \vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 4 & 2 \\ 3 & -2 & 7 \end{vmatrix}$$

$$= 32\hat{i} - \hat{j} - 14\hat{k} \quad 1$$

Given that  $\vec{p} \cdot \vec{c} = 18$

$$\text{or } \lambda(32\hat{i} - \hat{j} - 14\hat{k}) \cdot (2\hat{i} - \hat{j} + 4\hat{k}) = 18 \quad 1$$

$$\text{or } \lambda(64 + 1 - 56) = 18$$

$$\lambda = 2$$

$$\therefore \vec{p} = 64\hat{i} - 2\hat{j} - 28\hat{k} \quad \frac{1}{2}$$

[CBSE Marking Scheme 2012]

Q. 9. If  $\vec{a} = 2\hat{i} + \hat{j} + 3\hat{k}$  and  $\vec{b} = 3\hat{i} + \hat{j} - 2\hat{k}$ , then find  $|\vec{a} \times \vec{b}|$ . [R&U] [Outside Delhi Set-II, 2015]

Sol.

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 1 & 3 \\ 3 & 5 & -2 \end{vmatrix}$$

$$\vec{a} \times \vec{b} = \hat{i}(-2 - 15) - \hat{j}(-4 - 9) + \hat{k}(10 - 3)$$

$$|\vec{a} \times \vec{b}| = \sqrt{(-17)^2 + (13)^2 + (7)^2}$$

$$= \sqrt{289 + 169 + 49}$$

$$= \sqrt{507}$$

$$= \sqrt{3 \times 169}$$

$$|\vec{a} \times \vec{b}| = 13\sqrt{3}$$

[CBSE Marking Scheme 2015]

Q. 10. If  $\vec{a} = 3\hat{i} - \hat{j}$  and  $\vec{b} = 2\hat{i} + \hat{j} - 3\hat{k}$ , then express  $\vec{b}$  in the form of  $\vec{b} = b_1 + b_2$ , where  $b_1 \parallel \vec{a}$  and  $b_2$  perpendicular to  $\vec{a}$ .

[A] [NCERT] [O.D. Set I, II, III, 2013]

Sol. Here  $\vec{a} = 3\hat{i} - \hat{j}$

and  $\vec{b} = 2\hat{i} + \hat{j} - 3\hat{k}$

To express  $\vec{b} = \vec{b}_1 + \vec{b}_2$ ,

where  $\vec{b}_1 \parallel \vec{a}$  or  $\vec{b}_1 = \lambda \vec{a}$

$$\vec{b}_1 = \lambda(3\hat{i} - \hat{j}) \quad \frac{1}{2}$$

Let  $\vec{b}_2 = x\hat{i} + y\hat{j} + z\hat{k}$

Now,  $\vec{b}_2 \perp \vec{a}$  or  $\vec{b}_2 \cdot \vec{a} = 0$   $\frac{1}{2}$

$$3x - y = 0 \quad \dots(i)$$

Now,  $\vec{b} = \vec{b}_1 + \vec{b}_2$

$$\text{or } 2\hat{i} + \hat{j} - 3\hat{k} = (3\lambda + x)\hat{i} + (y - \lambda)\hat{j} + z\hat{k}$$

Comparing the corresponding components

$$2 = 3\lambda + x \quad \dots(ii) \frac{1}{2}$$

$$1 = -\lambda + y$$

$$\text{or } \lambda = y - 1 \quad \dots(iii) \frac{1}{2}$$

$$-3 = z \quad \dots(iv) \frac{1}{2}$$

From eqn. (ii),  $2 = 3(y - 1) + x$

$$\text{or } 2 = 3y - 3 + x$$

$$\text{or } x + 3y = 5 \quad \dots(v) \frac{1}{2}$$

Solving eqn. (i) & (v),  $x = \frac{1}{2}, y = \frac{3}{2}$

$\therefore$  From eqn. (iii),  $\lambda = \frac{1}{2}$

$$\therefore \vec{b}_1 = \frac{3}{2}\hat{i} - \frac{1}{2}\hat{j} \quad \frac{1}{2}$$

and  $\vec{b}_2 = \frac{1}{2}\hat{i} + \frac{3}{2}\hat{j} - 3\hat{k} \quad \frac{1}{2}$

[CBSE Marking Scheme 2013]

#### Commonly Made Error

- Generally students do not able to find the vector  $\vec{b}$  in the form of  $\vec{b} = \vec{b}_1 + \vec{b}_2$ , instead they add vector  $\vec{a}$  and  $\vec{b}$  which leads incorrect result.

#### Answering Tip

- Read and understand the question carefully to avoid such errors.

Q. 11. Find a unit vector perpendicular to both of the vectors  $3\vec{a} + 2\vec{b}$  and  $3\vec{a} - 2\vec{b}$ , where  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$  and  $\vec{b} = \hat{i} + 2\hat{j} + 3\hat{k}$ .

[R&U] [Delhi Set I, II, III, Comptt. 2016]

Sol. Here  $3\vec{a} + 2\vec{b} = 5\hat{i} + 7\hat{j} + 9\hat{k}$

and  $3\vec{a} - 2\vec{b} = \hat{i} - \hat{j} - 3\hat{k} \quad 1 + 1$

Let  $\vec{c}$  be the vector perpendicular to both  $(3\vec{a} + 2\vec{b})$  &  $(3\vec{a} - 2\vec{b})$ .

Then,  $\vec{c} = (3\vec{a} + 2\vec{b}) \times (3\vec{a} - 2\vec{b})$

$$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 5 & 7 & 9 \\ 1 & -1 & -3 \end{vmatrix}$$

$$= -12\hat{i} + 24\hat{j} - 12\hat{k} \quad 2$$

[CBSE Marking Scheme 2016]

**Detailed Solution :**

$$\text{Given } \vec{a} = \hat{i} + \hat{j} + \hat{k}$$

$$\vec{b} = \hat{i} + 2\hat{j} + 3\hat{k}$$

$$3\vec{a} = 3\hat{i} + 3\hat{j} + 3\hat{k}$$

$$\text{and } 2\vec{b} = 2\hat{i} + 4\hat{j} + 6\hat{k} \quad 1$$

Let  $\vec{c}$  be the vector perpendicular to both  $(3\vec{a} + 2\vec{b})$

and  $(3\vec{a} - 2\vec{b})$ .

$$\therefore 3\vec{a} + 2\vec{b} = 5\hat{i} + 7\hat{j} + 9\hat{k} \quad 1$$

$$\text{and } 3\vec{a} - 2\vec{b} = \hat{i} - \hat{j} - 3\hat{k} \quad 1$$

$$\therefore \vec{c} = (3\vec{a} + 2\vec{b}) \times (3\vec{a} - 2\vec{b})$$

$$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 5 & 7 & 9 \\ 1 & -1 & -3 \end{vmatrix}$$

$$= \hat{i}(-21+9) - \hat{j}(-15-9) + \hat{i}(-5-7)$$

$$= -12\hat{i} + 24\hat{j} - 12\hat{k} \quad 1$$

**Q. 12.** Show that the points  $A, B, C$  with position vectors  $2\hat{i} - \hat{j} + \hat{k}, \hat{i} - 3\hat{j} - 5\hat{k}$  and  $3\hat{i} - 4\hat{j} - 4\hat{k}$  respectively, are the vertices of a right-angled triangle, hence find the area of the triangle. R&U [O.D. Set-I, 2017]

**Sol.**  $\vec{AB} = -\hat{i} - 2\hat{j} - 6\hat{k}, \vec{BC} = 2\hat{i} - \hat{j} + \hat{k}, \vec{CA} = -\hat{i} + 3\hat{j} + 5\hat{k}$  1

Since  $\vec{AB}, \vec{BC}, \vec{CA}$  are not parallel vectors, and  $\vec{AB} + \vec{BC} + \vec{CA} = \vec{0} \therefore A, B, C$  form a triangle 1

Also  $\vec{BC} \cdot \vec{CA} = 0 \therefore A, B, C$  form a right triangle 1

Area of  $\Delta = \frac{1}{2} |\vec{AB} \times \vec{BC}| = \frac{1}{2} \sqrt{210}$  [CBSE Marking Scheme 2017] 1

OR

Handwritten solution for Q. 12:

$$\vec{OA} = 2\hat{i} - \hat{j} + \hat{k}$$

$$\vec{OB} = \hat{i} - 3\hat{j} - 5\hat{k}$$

$$\vec{OC} = 3\hat{i} - 4\hat{j} - 4\hat{k}$$

$$\vec{AB} = \vec{OB} - \vec{OA} = -\hat{i} - 2\hat{j} - 6\hat{k}$$

$$|\vec{AB}| = \sqrt{(-1)^2 + (-2)^2 + (-6)^2} = \sqrt{41} \text{ units}$$

$$\vec{AC} = \vec{OC} - \vec{OA} = \hat{i} - 3\hat{j} - 5\hat{k}$$

$$|\vec{AC}| = \sqrt{1^2 + (-3)^2 + (-5)^2} = \sqrt{35} \text{ units}$$

$$\vec{BC} = \vec{OC} - \vec{OB} = 2\hat{i} - \hat{j} + \hat{k}$$

$$|\vec{BC}| = \sqrt{(2)^2 + (-1)^2 + (1)^2} = \sqrt{6} \text{ units}$$

$$\vec{AC} \cdot \vec{BC} = 2 + 3 - 5 = 0 \text{ Hence } \vec{AC} \perp \vec{BC} \text{ Hence } \angle C = 90^\circ$$

$$|\vec{BC}|^2 + |\vec{AC}|^2 = |\vec{AB}|^2$$

Hence  $\Delta ABC$  is right angled at  $C$ .

$$\text{area} = \frac{1}{2} |\vec{AC} \times \vec{BC}|$$

$$= \frac{1}{2} \left| \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -3 & -5 \\ 2 & -1 & 1 \end{vmatrix} \right| = \frac{1}{2} | \hat{i}(-8) - \hat{j}(11) + \hat{k}(5) |$$

$$= \frac{1}{2} \sqrt{(-8)^2 + (-11)^2 + (5)^2} = \frac{1}{2} \sqrt{209} \text{ sq. units}$$

$$= \frac{1}{2} \sqrt{210} \text{ sq. units}$$

$$= \sqrt{\frac{210}{4}} = \sqrt{52.5} \text{ sq. units}$$

4  
[Topper's Answer 2017]

**Q. 13.** If  $\vec{a} = 2\hat{i} - \hat{j} - 2\hat{k}$  and  $\vec{b} = 7\hat{i} + 2\hat{j} - 3\hat{k}$  then express  $\vec{b}$  in the form of  $\vec{b} = \vec{b}_1 + \vec{b}_2$ , where  $\vec{b}_1$  is parallel to  $\vec{a}$  and  $\vec{b}_2$  is perpendicular to  $\vec{a}$ .

[R&U] [O.D. 2017]

Sol.  $\vec{b}_1 \parallel \vec{a}$  or let  $\vec{b}_1 = \lambda(2\hat{i} - \hat{j} - 2\hat{k})$  1/2  
 $\vec{b}_2 = \vec{b} - \vec{b}_1$   
 $= (7\hat{i} + 2\hat{j} - 3\hat{k}) - (2\lambda\hat{i} - \lambda\hat{j} - 2\lambda\hat{k})$  1/2  
 $= (7 - 2\lambda)\hat{i} + (2 + \lambda)\hat{j} - (3 - 2\lambda)\hat{k}$  1

$\vec{b}_2 \perp \vec{a}$  or  $2(7 - 2\lambda) - 1(2 + \lambda) + 2(3 - 2\lambda) = 0$   
 or  $\lambda = 2$   
 $\therefore \vec{b}_1 = 4\hat{i} - 2\hat{j} - 4\hat{k}$  1/2  
 and  $\vec{b}_2 = 3\hat{i} + 4\hat{j} + \hat{k}$  1/2  
 or  $(7\hat{i} + 2\hat{j} - 3\hat{k}) = (4\hat{i} - 2\hat{j} - 4\hat{k}) + (3\hat{i} + 4\hat{j} + \hat{k})$  1

[CBSE Marking Scheme 2017]


**Answering Tip**

- Clarify the concept of scalar projection of vector thoroughly.

**Q. 14.** Given that vectors  $\vec{a}, \vec{b}, \vec{c}$  form a triangle such that  $\vec{a} = \vec{b} + \vec{c}$ . Find  $p, q, r, s$  such that area of triangle is  $5\sqrt{6}$  where  $\vec{a} = p\hat{i} + q\hat{j} + r\hat{k}, \vec{b} = s\hat{i} + 3\hat{j} + 4\hat{k}$  and  $\vec{c} = 3\hat{i} + \hat{j} - 2\hat{k}$ .

[K&A] [O.D. Set II 2016]

Sol.  $\vec{a} = \vec{b} + \vec{c}$  Hence  $\vec{a} = p\hat{i} + q\hat{j} + r\hat{k}$   
 $\vec{b} = s\hat{i} + 3\hat{j} + 4\hat{k}$   
 $\vec{c} = 3\hat{i} + \hat{j} - 2\hat{k}$



$\vec{a} = \vec{b} + \vec{c}$   
 $p\hat{i} + q\hat{j} + r\hat{k} = s\hat{i} + 3\hat{j} + 4\hat{k} + 3\hat{i} + \hat{j} - 2\hat{k}$   
 $p\hat{i} + q\hat{j} + r\hat{k} = (s+3)\hat{i} + 4\hat{j} + 2\hat{k}$   
 Equating components,  
 $p = s + 3$   
 $q = 4$   
 $r = 2$

Area of triangle =  $5\sqrt{6}$   
 But Area of triangle =  $\frac{1}{2} |\vec{a} \times \vec{b}|$

$$5\sqrt{6} = \frac{1}{2} \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ p & 4 & 2 \\ s+3 & 4 & 2 \end{vmatrix} = \frac{1}{2} |10\hat{i} - (2p+6)\hat{j} + (2-p)\hat{k}|$$

$$600 = 100 + (2p+6)^2 + (2-p)^2$$

$$500 = 4p^2 + 36 + 24p + 14 + p^2 - 24p \Rightarrow 5p^2 = 320$$

$p^2 = 64$   
 $p = \pm 8$   
 If  $p = 8, s = p - 3 = 5$   
 If  $p = -8, s = p - 3 = -11$


[Topper's Answer 2016]

**Q. 15.** Find the area of a parallelogram  $ABCD$  whose side  $AB$  and the diagonal  $DB$  are given by the vectors  $5\hat{i} + 7\hat{k}$  and  $2\hat{i} + 2\hat{j} + 3\hat{k}$  respectively.

[K] [Foreign 2017]

**Sol.**  $\vec{AD} = \vec{AB} - \vec{DB} = 3\hat{i} - 2\hat{j} + 4\hat{k}$  1

Area =  $|\vec{AB} \times \vec{AD}|$



= magnitude of  $\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 5 & 0 & 7 \\ 3 & -2 & 4 \end{vmatrix}$  1

=  $|14\hat{i} + \hat{j} - 10\hat{k}|$  1


=  $\sqrt{297}$  sq. units or  $3\sqrt{33}$  sq. units 1

**Q. 16.** Find the area of a parallelogram  $ABCD$  whose side  $AB$  and the diagonal  $AC$  are given by the vectors  $3\hat{i} + \hat{j} + 4\hat{k}$  and  $4\hat{i} + 5\hat{k}$  respectively.

[R&U] [Foreign 2017]

**Sol.**  $\vec{BC} = \vec{AC} - \vec{AB} = \hat{i} - \hat{j} + \hat{k}$  1

Area =  $|\vec{AB} \times \vec{BC}|$



= magnitude of  $\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & 1 & 4 \\ 1 & -1 & 1 \end{vmatrix}$  1

=  $|5\hat{i} + \hat{j} - 4\hat{k}|$  1

=  $\sqrt{42}$  sq. units 1

[CBSE Marking Scheme 2017]

**Q. 17.** If  $\vec{a} = 2\hat{i} + \hat{j} - \hat{k}$ ,  $\vec{b} = 4\hat{i} - 7\hat{j} + \hat{k}$ , find a vector  $\vec{c}$  such that  $\vec{a} \times \vec{c} = \vec{b}$  and  $\vec{a} \cdot \vec{c} = 6$ .

[K&U] [Foreign 2017]

**Sol.** Let  $\vec{c} = x\hat{i} + y\hat{j} + z\hat{k}$ ;  $\vec{a} \cdot \vec{c} = 6$  or  $2x + y - z = 6$

Now,  $\vec{a} \times \vec{c} = \vec{b}$

or  $\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 1 & -1 \\ x & y & z \end{vmatrix} = 4\hat{i} - 7\hat{j} + \hat{k}$  1½

or  $\hat{i}(z + y) - \hat{j}(2z + x) + \hat{k}(2y - x) = 4\hat{i} - 7\hat{j} + \hat{k}$

or  $z + y = 4, 2z + x = 7, 2y - x = 1$  1

Solving and getting  $x = 3, y = 2, z = 2$

$\vec{c} = 3\hat{i} + 2\hat{j} + 2\hat{k}$  1½

[CBSE Marking Scheme 2017]

**Q. 18.** If  $\vec{a} = \hat{i} + 2\hat{j} + \hat{k}$ ,  $\vec{b} = 2\hat{i} + \hat{j}$  and  $\vec{c} = 3\hat{i} - 4\hat{j} - 5\hat{k}$ , then find a unit vector perpendicular to both of the

vectors  $(\vec{a} - \vec{b})$  and  $(\vec{c} - \vec{b})$ . [R&U] [All India 2015]

**Sol.** Try Yourself

**Q. 19.** Find a unit vector perpendicular to both of the vectors  $\vec{a} + \vec{b}$  and  $\vec{a} - \vec{b}$  where,  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$  and

$\vec{b} = \hat{i} + 2\hat{j} + 3\hat{k}$ . [R&U] [Foreign 2014]

**Sol.** Try Yourself

**Q. 20.** Find a unit vector perpendicular to each of the vectors  $\vec{a} + \vec{b}$  and  $\vec{a} - \vec{b}$ , where  $\vec{a} = 3\hat{i} + 2\hat{j} + 2\hat{k}$

and  $\vec{b} = \hat{i} + 2\hat{j} - 2\hat{k}$ . [R&U] [Delhi 2011]

**Sol.** Try Yourself

**Q. 21.** Using vectors, find the area of triangle with vertices  $A(1, 1, 2)$ ,  $B(2, 3, 5)$  and  $C(1, 5, 5)$ .

[R&U] [All India 2011]

**Sol.** Try Yourself

**Q. 22.** Using vectors, find the area of triangle with vertices  $A(2, 3, 5)$ ,  $B(3, 5, 8)$  and  $C(2, 7, 8)$ .

[R&U] [Delhi 2010C]

**Sol.** Try Yourself





## TOPIC-4 Scalar Triple Product

### Revision Notes

#### 1. Scalar Triple Product

If  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are any three vectors, then the scalar product of  $\vec{a} \times \vec{b}$  with  $\vec{c}$  is called scalar triple product of  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$ .

Thus,  $(\vec{a} \times \vec{b}) \cdot \vec{c}$  is called the scalar triple product of  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$ .

- **Notation for scalar triple product** : The scalar triple product of  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  is denoted by  $[\vec{a} \vec{b} \vec{c}]$  i.e.,  $(\vec{a} \times \vec{b}) \cdot \vec{c} = [\vec{a} \vec{b} \vec{c}]$ .

- **Properties/Observations of Scalar Triple Product**

⊕  $(\vec{a} \times \vec{b}) \cdot \vec{c} = \vec{a} \cdot (\vec{b} \times \vec{c})$  i.e., the position of dot and cross can be interchanged without change in the value of the scalar triple product (provided their cyclic order remains the same).

⊕  $[\vec{a} \vec{b} \vec{c}] = [\vec{b} \vec{c} \vec{a}] = [\vec{c} \vec{a} \vec{b}]$ , i.e., the value of scalar triple product doesn't change when cyclic order of the vectors is maintained.

Also,  $[\vec{a} \vec{b} \vec{c}] = -[\vec{b} \vec{a} \vec{c}]$ ;  $[\vec{b} \vec{c} \vec{a}] = -[\vec{b} \vec{a} \vec{c}]$ . i.e., the value of scalar triple product remains the same in magnitude but changes the sign when cyclic order of the vectors is altered.

⊕ For any three vectors  $\vec{a}$ ,  $\vec{b}$ ,  $\vec{c}$  and scalar  $\lambda$ , we have  $[\lambda \vec{a} \vec{b} \vec{c}] = \lambda [\vec{a} \vec{b} \vec{c}]$ .

⊕ The value of scalar triple product is zero if any two of the three vectors are identical. That is,  $[\vec{a} \vec{a} \vec{c}] = 0 = [\vec{a} \vec{b} \vec{b}] = [\vec{a} \vec{b} \vec{a}]$  etc.

⊕ Value of scalar triple product is zero if any two of the three vectors are parallel or collinear.

⊕ Scalar triple product of  $\hat{i}$ ,  $\hat{j}$  and  $\hat{k}$  is 1 (unity) i.e.,  $[\hat{i} \hat{j} \hat{k}] = 1$

⊕ If  $[\vec{a} \vec{b} \vec{c}] = 0$ , then the non-parallel and non-zero vectors  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are **coplanar**.

### Know the Formulae

#### Volume of Parallelepiped

- If  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  represent the three co-terminus edges of a parallelepiped, then its volume can be obtained by :

$$[\vec{a} \vec{b} \vec{c}] = (\vec{a} \times \vec{b}) \cdot \vec{c} \text{ i.e.,}$$

$$(\vec{a} \times \vec{b}) \cdot \vec{c} = \text{Base area of Parallelepiped} \times \text{Height of Parallelepiped on this base}$$

#### Note :

- If for any three vectors  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$ , we have  $[\vec{a} \vec{b} \vec{c}] = 0$ , then volume of parallelepiped with the co-terminus edges as  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$ , is zero. This is possible only if the vectors  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are co-planar.



## ? Very Short Answer Type Questions

(1 mark each)

Q. 1. Find  $\vec{a} \cdot (\vec{b} \times \vec{c})$ , if

$$\vec{a} = 2\hat{i} + \hat{j} + 3\hat{k}, \vec{b} = -\hat{i} + 2\hat{j} + \hat{k} \text{ and}$$

$$\vec{c} = 3\hat{i} + \hat{j} + 2\hat{k}. \quad \text{R\&U [O.D. Set I, II, III, 2014]}$$

Sol. 
$$\vec{b} \times \vec{c} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -1 & 2 & 1 \\ 3 & 1 & 2 \end{vmatrix}$$

$$= \hat{i}(4-1) - \hat{j}(-2-3) + \hat{k}(-1-6)$$

$$= 3\hat{i} + 5\hat{j} - 7\hat{k}$$

$$\therefore \vec{a} \cdot (\vec{b} \times \vec{c}) = (2\hat{i} + \hat{j} + 3\hat{k}) \cdot (3\hat{i} + 5\hat{j} - 7\hat{k})$$

$$= 6 + 5 - 21 = -10 \quad 1$$

[CBSE Marking Scheme 2014]

Q. 2. Find  $\lambda$ , if the vectors

$$\vec{a} = \hat{i} + 3\hat{j} + \hat{k}, \vec{b} = 2\hat{i} - \hat{j} - \hat{k} \text{ and}$$

$$\vec{c} = \lambda\hat{j} + 3\hat{k} \text{ are coplanar.} \quad \text{R\&U [Delhi, 2015]}$$

Sol. 
$$\begin{vmatrix} 1 & 3 & 1 \\ 2 & -1 & -1 \\ 0 & \lambda & 3 \end{vmatrix} = 0 \quad \frac{1}{2}$$

or 
$$\lambda = 7 \quad \frac{1}{2}$$

[CBSE Marking Scheme 2015]

## ? Short Answer Type Questions

(2 marks each)

Q. 1. If the vectors  $\hat{i} - \hat{j} + \hat{k}$ ,  $3\hat{i} + \hat{j} + 2\hat{k}$  and  $\hat{i} + \lambda\hat{j} - 3\hat{k}$  are coplanar, then find the value of  $\lambda$ .

R&amp;U [O.D. Comptt, 2017]

Sol. For three vectors to be coplanar

$$\begin{vmatrix} 1 & -1 & 1 \\ 3 & 1 & 2 \\ 1 & \lambda & -3 \end{vmatrix} = 0 \quad 1$$

or 
$$\lambda = 15 \quad 1$$

[CBSE Marking Scheme 2017]

Q. 2. If the vectors  $\vec{a} = \hat{i} + 3\hat{j} + \hat{k}$ ,  $\vec{b} = \lambda\hat{i} + 7\hat{j} + 3\hat{k}$  and  $\vec{c} = 2\hat{i} - \hat{j} - \hat{k}$  are coplanar, then find the value of  $\lambda$ .

R&amp;U [O.D. Comptt, 2017]

Sol. For coplanarity of vectors 
$$\begin{vmatrix} 1 & 3 & 1 \\ 2 & -1 & -1 \\ \lambda & 7 & 3 \end{vmatrix} = 0 \quad 1$$

$$4 - 18 + 3\lambda + 14 + \lambda = 0$$

Solving to get  $\lambda = 0 \quad 1$

[CBSE Marking Scheme 2017]

## ? Long Answer Type Questions-I

(4 marks each)

Q. 1. Prove that :

$$\vec{a} \cdot \{(\vec{b} + \vec{c}) \times (\vec{a} + 2\vec{b} + 3\vec{c})\} = [\vec{a} \ \vec{b} \ \vec{c}].$$

R&amp;U [S.Q.P. 2016-17]

Sol. LHS = 
$$\vec{a} \cdot (\vec{b} \times \vec{a} + 2\vec{b} \times \vec{b} + 3\vec{b} \times \vec{c} + \vec{c} \times \vec{a} + 2\vec{c} \times \vec{b} + 3\vec{c} \times \vec{c}) \quad 1$$

$$= \vec{a} \cdot (\vec{b} \times \vec{a}) + 3\vec{a} \cdot (\vec{b} \times \vec{c}) + \vec{a} \cdot (\vec{c} \times \vec{a}) + 2\vec{a} \cdot (\vec{c} \times \vec{b})$$

as 
$$\vec{b} \times \vec{b} = \vec{c} \times \vec{c} = 0 \quad 1$$

$$= 3[\vec{a} \ \vec{b} \ \vec{c}] + 2[\vec{a} \ \vec{c} \ \vec{b}] \quad 1$$

$$= 3[\vec{a} \ \vec{b} \ \vec{c}] - 2[\vec{a} \ \vec{b} \ \vec{c}]$$

$$= [\vec{a} \ \vec{b} \ \vec{c}] \quad 1$$

[CBSE Marking Scheme 2016]

### Commonly Made Error

- Many candidates make errors while simplifying the scalar triple product.

### Answering Tip

- Scalar triple product and its applications need to be practiced with the help of practical examples.

Q. 2. Show that the four points  $A(4, 5, 1)$ ,  $B(0, -1, -1)$ ,  $C(3, 9, 4)$  and  $D(-4, 4, 4)$  are coplanar.

R&amp;U [CBSE Outside Delhi, 2016]



**Sol.**  $\vec{AB} = -4\hat{i} - 6\hat{j} - 2\hat{k}$ ,  
 $\vec{AC} = -\hat{i} + 4\hat{j} + 3\hat{k}$ ,  
and  $\vec{AD} = -8\hat{i} - \hat{j} + 3\hat{k}$  1½

For 4 points to be coplanar,  
 $|\vec{AB} \ \vec{AC} \ \vec{AD}| = 0$

i.e.,  $\begin{vmatrix} -4 & -6 & -2 \\ -1 & 4 & 3 \\ -8 & -1 & 3 \end{vmatrix} = 0$  1½

$= -4(12 + 3) + 6(-3 + 24) - 2(1 + 32)$   
 $= -60 + 126 - 66 = 0$ , which is true  
Hence, points are coplanar. 1

**[CBSE Marking Scheme 2016]**

**Alternative Method :**

If four points  $A, B, C, D$  are coplanar, then vector

$\vec{AB}, \vec{AC}$  and  $\vec{AD}$  will be coplanar and so

$$|\vec{AB} \ \vec{AC} \ \vec{AD}| = 0$$

$$\begin{aligned} A &= (4, 5, 1) \\ B &= (0, -1, -1) \\ C &= (3, 9, 4) \\ D &= (-4, 4, 4) \end{aligned}$$

By considering  $O = (0, 0, 0)$  as initial point

$$\vec{OA} = 4\hat{i} + 5\hat{j} + \hat{k}$$

$$\vec{OB} = -\hat{j} - \hat{k},$$

$$\vec{OC} = 3\hat{i} + 9\hat{j} + 4\hat{k}$$

and  $\vec{OD} = -4\hat{i} + 4\hat{j} + 4\hat{k}$  ½

$\therefore \vec{AB} = \vec{OB} - \vec{OA}$   
 $= -\hat{j} - \hat{k} - (4\hat{i} + 5\hat{j} + \hat{k})$   
 $= -4\hat{i} - 6\hat{j} - 2\hat{k}$  ½

$\vec{AC} = \vec{OC} - \vec{OA}$   
 $= 3\hat{i} + 9\hat{j} + 4\hat{k} - (4\hat{i} + 5\hat{j} + \hat{k})$   
 $= -\hat{i} + 4\hat{j} + 3\hat{k}$  ½

and  $\vec{AD} = \vec{OD} - \vec{OA}$   
 $= -4\hat{i} + 4\hat{j} + 4\hat{k} - (4\hat{i} + 5\hat{j} + \hat{k})$   
 $= -8\hat{i} - \hat{j} + 3\hat{k}$  ½

Now,

$$|\vec{AB} \ \vec{AC} \ \vec{AD}| = \begin{vmatrix} -4 & -6 & -2 \\ -1 & 4 & 3 \\ -8 & -1 & 3 \end{vmatrix}$$

$$\begin{aligned} &= -4(12 + 3) + 6(-3 + 24) + (-2)(1 + 32) \\ &= -60 + 126 - 66 \\ &= 0 \end{aligned}$$

1

$\therefore \vec{AB}, \vec{AC}, \vec{AD}$  are coplanar and these three vectors are co-initial vectors. So, points  $A, B, C, D$  are coplanar. 1

**Q. 3. Prove that**

$$\vec{a} \ \vec{b} + \vec{c} \ \vec{d} = \vec{a} \ \vec{b} \ \vec{d} + \vec{a} \ \vec{c} \ \vec{d}.$$

**[A] [O.D. Set I, II, III Comptt. 2015]**

**Sol.** Taking LHS =  $\vec{a} \cdot \{(\vec{b} + \vec{c}) \times \vec{d}\}$  1 + 1

$$= \vec{a} \cdot \{(\vec{b} \times \vec{d}) + (\vec{c} \times \vec{d})\}$$
 1

$$= \vec{a} \cdot (\vec{b} \times \vec{d}) + \vec{a} \cdot (\vec{c} \times \vec{d})$$
 1

$$= [\vec{a} \ \vec{b} \ \vec{d}] + [\vec{a} \ \vec{c} \ \vec{d}]$$

**[CBSE Marking Scheme 2015]**

**[AI] Q. 4. If the vectors  $\vec{a}, \vec{b}$  and  $\vec{c}$  are coplanar, prove that the vectors  $\vec{a} + \vec{b}, \vec{b} + \vec{c}$  and  $\vec{c} + \vec{a}$  are also coplanar.** **[R&U] [Delhi Set I Comptt. 2014]**

**[Delhi Set III Comptt. 2013]**

**[Foreign Set I, II, III, 2014] [Delhi Set I, II, III, 2016]**

**Sol.** Here,  $(\vec{a} + \vec{b}), (\vec{b} + \vec{c}), (\vec{c} + \vec{a})$  are coplanar, 1

$$\therefore (\vec{a} + \vec{b}) \cdot \{(\vec{b} + \vec{c}) \times (\vec{c} + \vec{a})\} = 0$$
 1

or  $(\vec{a} + \vec{b}) \cdot (\vec{b} \times \vec{c} + \vec{b} \times \vec{a} + \vec{c} \times \vec{c} + \vec{c} \times \vec{a}) = 0$  1

or  $\vec{a} \cdot (\vec{b} \times \vec{c}) + \vec{a} \cdot (\vec{b} \times \vec{a}) + \vec{a} \cdot (\vec{c} \times \vec{a})$   
 $+ \vec{a} \cdot (\vec{c} \times \vec{c}) + \vec{b} \cdot (\vec{b} \times \vec{c}) + \vec{b} \cdot (\vec{b} \times \vec{a})$   
 $+ \vec{b} \cdot (\vec{c} \times \vec{c}) + \vec{b} \cdot (\vec{c} \times \vec{a}) = 0$  ½

or  $2\{ \vec{a} \cdot (\vec{b} \times \vec{c}) \} = 0$

$\therefore \vec{a} \cdot (\vec{b} \times \vec{c}) = 0$  ½

$\therefore \vec{a}, \vec{b}, \vec{c}$  are coplanar.

Similarly converse part can also be proved.

**[CBSE Marking Scheme 2014]**

**Q. 5. Find the value of  $\lambda$ , if the points with position vectors  $3\hat{i} - 2\hat{j} - \hat{k}, 2\hat{i} + 3\hat{j} - 4\hat{k}, -\hat{i} + \hat{j} + 2\hat{k}$  and  $4\hat{i} + 5\hat{j} + \lambda\hat{k}$  are coplanar.** **[R&U] [S.Q.P. 2013]**

**Sol.** Let the points be  $A(3, -2, -1), B(2, 3, -4), C(-1, 1, 2)$  and  $D(4, 5, \lambda)$



$$\begin{aligned}\vec{AB} &= (\text{Position vector of } B) \\ &\quad - (\text{Position vector of } A)\end{aligned}$$

$$\therefore \vec{AB} = -\hat{i} + 5\hat{j} - 3\hat{k},$$

Similarly

$$\vec{AC} = -4\hat{i} + 3\hat{j} + 3\hat{k}$$

$$\text{and } \vec{AD} = \hat{i} + 7\hat{j} + (\lambda + 1)\hat{k} \quad 1\frac{1}{2}$$

$$A, B, C, D \text{ are coplanar if } [\vec{AB} \ \vec{AC} \ \vec{AD}] = 0 \quad \frac{1}{2}$$

$$[\vec{AB} \ \vec{AC} \ \vec{AD}] = \begin{vmatrix} -1 & 5 & -3 \\ -4 & 3 & 3 \\ 1 & 7 & \lambda + 1 \end{vmatrix} = 0 \quad \frac{1}{2}$$

$$\therefore 1(15 + 9) - 7(-3 - 12) + (\lambda + 1)(-3 + 20) = 0 \quad 1$$

$$24 + 105 + 17\lambda + 17 = 0$$

$$\text{or } \lambda = -\frac{146}{17} \quad \frac{1}{2}$$

#### Commonly Made Error

- Some candidates fail to apply condition of coplanarity.

#### Answering Tip

- Scalar triple product and its applications need to be practiced with the help of practical examples.

Q. 6. If  $\vec{a} = 2\hat{i} - 3\hat{j} + 4\hat{k}$ ,  $\vec{b} = \hat{i} + 2\hat{j} - 3\hat{k}$  and

$$\vec{c} = 3\hat{i} + 4\hat{j} - \hat{k}, \text{ then find}$$

$$\vec{a} \cdot (\vec{b} \times \vec{c}) \text{ and } (\vec{a} \times \vec{b}) \cdot \vec{c}. \text{ Is,}$$

$$\vec{a} \cdot (\vec{b} \times \vec{c}) = (\vec{a} \times \vec{b}) \cdot \vec{c} ? \quad \text{R\&U}$$

$$\text{Sol. } \vec{a} \cdot (\vec{b} \times \vec{c}) = \begin{vmatrix} 2 & -3 & 4 \\ 1 & 2 & -3 \\ 3 & 4 & -1 \end{vmatrix} \quad \frac{1}{2}$$

$$\begin{aligned}&= 2(-2 + 12) + 3(-1 + 9) + 4(4 - 6) \\ &= 20 + 24 - 8 \\ &= 36 \quad 1\end{aligned}$$

$$\text{and } (\vec{a} \times \vec{b}) \cdot \vec{c} = \vec{c} \cdot (\vec{a} \times \vec{b})$$

$$= \begin{vmatrix} 3 & 4 & -1 \\ 2 & -3 & 4 \\ 1 & 2 & -3 \end{vmatrix} \quad \frac{1}{2}$$

$$\begin{aligned}&= 3(9 - 8) - 4(-6 - 4) - 1(4 + 3) \\ &= 3 + 40 - 7 = 36 \quad 1\end{aligned}$$

$$\therefore \vec{a} \cdot (\vec{b} \times \vec{c}) = (\vec{a} \times \vec{b}) \cdot \vec{c}. \quad 1$$

[CBSE Marking Scheme 2013]

Q. 7. If  $\vec{p} = \hat{i} + \hat{j} + \hat{k}$  and  $\vec{q} = \hat{i} - 2\hat{j} + \hat{k}$ , find a vector of magnitude  $5\sqrt{3}$  units perpendicular to the vector  $\vec{q}$  and coplanar with vectors  $\vec{p}$  and  $\vec{q}$ .

[R&U] (SQP 2018-19)

Sol. Let  $\vec{r} = a\hat{i} + b\hat{j} + c\hat{k}$  be the required vector.

$$\text{Since, } \vec{r} \perp \vec{q}$$

$$\text{therefore, } 1a - 2b + 1c = 0 \quad \dots(1) \quad 1$$

Also,  $\vec{p}$ ,  $\vec{q}$  and  $\vec{r}$  are coplanar.

$$\therefore \begin{vmatrix} \vec{p} & \vec{q} & \vec{r} \end{vmatrix} = 0$$

$$\begin{vmatrix} 1 & 1 & 1 \\ 1 & -2 & 1 \\ a & b & c \end{vmatrix} = 0 \Rightarrow a - c = 0 \quad \dots(2) \quad 1$$

Solving equation (1) and (2)

$$\frac{a}{2-0} = \frac{b}{1+1} = \frac{c}{0+2}$$

$$\Rightarrow \frac{a}{2} = \frac{b}{2} = \frac{c}{2}$$

$$\text{i.e., } \frac{a}{1} = \frac{b}{1} = \frac{c}{1}$$

$$\therefore \vec{r} = 1\hat{i} + 1\hat{j} + 1\hat{k}$$

$$|\vec{r}| = \sqrt{3}$$

$$\therefore \text{Unit vector } \hat{r} = \frac{\vec{r}}{|\vec{r}|} = \frac{\hat{i} + \hat{j} + \hat{k}}{\sqrt{3}} \quad 1$$

$$\therefore \text{Required vector} = 5\sqrt{3}\hat{r} = 5(\hat{i} + \hat{j} + \hat{k}) \quad 1$$

[CBSE Marking Scheme 2018-19]

Q. 8. Find  $x$  such that the four points  $A(4, 1, 2)$ ,  $B(5, x, 6)$ ,  $C(5, 1, -1)$  and  $D(7, 4, 0)$  are coplanar.

[A] [Outside Delhi Set-II, 2015]

Sol. Here,  $\vec{AB}$ ,  $\vec{BC}$ ,  $\vec{CD}$  are coplanar

$$\text{So } \vec{AB} \cdot (\vec{BC} \times \vec{CD}) = 0 \quad 1$$

triple product is 0.

$$\vec{AB} = 1\hat{i} + (x-1)\hat{j} + 4\hat{k}$$

$$\vec{BC} = 0\hat{i} + (1-x)\hat{j} - 7\hat{k}$$

$$\vec{CD} = 2\hat{i} + 3\hat{j} + \hat{k} \quad 1\frac{1}{2}$$

$$\vec{AB} \cdot (\vec{BC} \times \vec{CD}) = \begin{vmatrix} 1 & x-1 & 4 \\ 0 & 1-x & -7 \\ 2 & 3 & 1 \end{vmatrix} = 0$$

expanding by  $R_1$

$$1(1-x+21) - (x-1)14 + 4(2(x-1)) = 0$$

$$22 - x - 14x + 14 + 8x - 8 = 0$$

$$-7x = -28$$

$$x = 4$$

1½

[CBSE Marking Scheme 2015]

**Q. 9.** If the vector  $\vec{p} = a\hat{i} + \hat{j} + \hat{k}$ ,  $\vec{q} = \hat{i} + b\hat{j} + \hat{k}$  and  $\vec{r} = \hat{i} + \hat{j} + c\hat{k}$  are coplanar, then for  $a, b, c \neq 1$ , then show that.

$$\frac{1}{1-a} + \frac{1}{1-b} + \frac{1}{1-c} = 1$$

[R&amp;U] [Outside Delhi Set I, II, III, Comptt. 2016]

[SQP Dec. 2016-17]

**Sol.** Since the vector  $\vec{p}$ ,  $\vec{q}$  and  $\vec{r}$  are coplanar,

$$\therefore [\vec{p}, \vec{q}, \vec{r}] = 0 \quad 1$$

$$\text{i.e.,} \quad \begin{vmatrix} a & 1 & 1 \\ 1 & b & 1 \\ 1 & 1 & c \end{vmatrix} = 0$$

$$\text{and} \quad \begin{matrix} R_2 \rightarrow R_2 - R_1 \\ R_3 \rightarrow R_3 - R_1 \end{matrix} \quad 1$$

$$\text{or} \quad \begin{vmatrix} a & 1 & 1 \\ 1-a & b-1 & 0 \\ 1-a & 0 & c-1 \end{vmatrix} = 0$$

$$\text{Or } a(b-1)(c-1) - 1(1-a)(c-1) - 1(1-a)(b-1) = 0$$

$$\text{i.e., } a(1-b)(1-c) + (1-a)(1-c) + (1-a)(1-b) = 0$$

1

Dividing both the sides by  $(1-a)(1-b)(1-c)$ , we get

$$\frac{a}{1-a} + \frac{1}{1-b} + \frac{1}{1-c} = 0$$

$$\text{i.e.,} \quad -1 + \frac{1}{1-a} + \frac{1}{1-b} + \frac{1}{1-c} = 0$$

$$\text{i.e.,} \quad \frac{1}{1-a} + \frac{1}{1-b} + \frac{1}{1-c} = 1 \quad 1$$

[CBSE Marking Scheme 2016]

**Commonly Made Error**

• Some candidates do mistake while doing scalar triple product.

• The right product is  $\begin{vmatrix} \vec{a} & \vec{b} & \vec{c} \end{vmatrix} = \vec{a} \cdot (\vec{b} \times \vec{c})$

• but student do mistake  $\begin{vmatrix} \vec{a} & \vec{b} & \vec{c} \end{vmatrix} = \begin{vmatrix} \vec{a} & \vec{b} \end{vmatrix} \times \vec{c}$  or

$$\begin{vmatrix} \vec{a} & \vec{b} & \vec{c} \end{vmatrix} = \begin{vmatrix} \vec{a} & \vec{c} \end{vmatrix} \times \vec{b}$$

**Q. 10.** If  $\vec{a}, \vec{b}, \vec{c}$  are three vectors such that  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$

then prove that  $\vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{c} \times \vec{a}$ , and hence

show that  $[\vec{a} \ \vec{b} \ \vec{c}] = 0$ . [R&U] [SQP 2017-18]

$$\text{Sol.} \quad \vec{a} \times (\vec{a} + \vec{b} + \vec{c}) = \vec{a} \times \vec{0} \quad 1$$

$$\text{or} \quad \vec{a} \times \vec{b} + \vec{a} \times \vec{c} = \vec{0} \quad \frac{1}{2}$$

$$\text{or} \quad \vec{a} \times \vec{b} = \vec{c} \times \vec{a} \quad \frac{1}{2}$$

$$\vec{b} \times (\vec{a} + \vec{b} + \vec{c}) = \vec{b} \times \vec{0} \quad \frac{1}{2}$$

$$\text{or} \quad \vec{b} \times \vec{c} = \vec{a} \times \vec{b} \quad \frac{1}{2}$$

$$\therefore \vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{c} \times \vec{a} \quad \frac{1}{2}$$

$$[\vec{a} \ \vec{b} \ \vec{c}] = \vec{a} \cdot (\vec{b} \times \vec{c}) = \vec{c} \cdot (\vec{a} \times \vec{b}) = 0$$

[As the scalar triple product of three vectors is zero if any two of them are equal.] ½

[CBSE Marking Scheme 2017-18]

**Q. 11.** Find the value of  $\lambda$ , if four points with position vectors  $3\hat{i} + 6\hat{j} + 9\hat{k}$ ,  $\hat{i} + 2\hat{j} + 3\hat{k}$ ,  $2\hat{i} + 3\hat{j} + \hat{k}$  and  $4\hat{i} + 6\hat{j} + \lambda\hat{k}$  are coplanar. [R&U] [O.D. Set I 2017]

**Sol.** Given points,  $A, B, C, D$  are coplanar, if the vectors  $\vec{AB}, \vec{AC}$  and  $\vec{AD}$  are coplanar, i.e.,

$$\vec{AB} = -2\hat{i} - 4\hat{j} - 6\hat{k}, \vec{AC} = -\hat{i} - 3\hat{j} - 8\hat{k}, \vec{AD} = \hat{i} + (\lambda - 9)\hat{k} \text{ are coplanar} \quad 1\frac{1}{2}$$

$$\text{i.e.,} \quad \begin{vmatrix} -2 & -4 & -6 \\ -1 & -3 & -8 \\ 1 & 0 & \lambda - 9 \end{vmatrix} = 0$$

$$\text{or} \quad -2[-3\lambda + 27] + 4[-\lambda + 17] - 6(3) = 0 \quad 1+1$$

$$\text{or} \quad \lambda = 2. \quad \text{[CBSE Marking Scheme 2017] } \frac{1}{2}$$



OR

$\vec{OA} = 3\hat{i} + 6\hat{j} + 9\hat{k}$   
 $\vec{OB} = \hat{i} + 2\hat{j} + 3\hat{k}$   
 $\vec{OC} = 2\hat{i} + 3\hat{j} + \hat{k}$   
 $\vec{OD} = 4\hat{i} + 6\hat{j} + \lambda\hat{k}$

$\vec{AB} = \vec{OB} - \vec{OA} = -2\hat{i} - 4\hat{j} - 6\hat{k}$   
 $\vec{AC} = \vec{OC} - \vec{OA} = -\hat{i} - 3\hat{j} - 8\hat{k}$   
 $\vec{AD} = \vec{OD} - \vec{OA} = \hat{i} + 0\hat{j} + (\lambda - 9)\hat{k}$

Scalar triple product  $[\vec{a} \ \vec{b} \ \vec{c}] = 0 \quad \therefore \vec{a}, \vec{b}, \vec{c}$  are coplanar.

$$\begin{vmatrix} -2 & -4 & -6 \\ -1 & -3 & -8 \\ 1 & 0 & \lambda - 9 \end{vmatrix} = 0$$

$(-2)(-3\lambda + 27) + 4(-\lambda + 9 + 8) - 6(3) = 0$   
 $6\lambda - 54 - 4\lambda + 68 - 18 = 0$   
 $2\lambda = 54 + 18 - 68$   
 $2\lambda = 72 - 68 = 4$   
 $\lambda = 2$

4  
[Topper's Answer 2017]

Q. 12. Find the value of  $x$  such that the points  $A(3, 2, 1)$ ,  $B(4, x, 5)$ ,  $C(4, 2, -2)$  and  $D(6, 5, -1)$  are coplanar. **R&U [O.D. 2017]**

Sol. Points  $A, B, C$  and  $D$  are coplanar, then the vectors  $\vec{AB}, \vec{AC}$ , and  $\vec{AD}$  must be coplanar.

$\vec{AB} = \hat{i} + (x-2)\hat{j} + 4\hat{k}; \vec{AC} = \hat{i} - 3\hat{k}$

$\vec{AD} = 3\hat{i} + 3\hat{j} - 2\hat{k}$  1½

i.e.,  $\begin{vmatrix} 1 & x-2 & 4 \\ 1 & 0 & -3 \\ 3 & 3 & -2 \end{vmatrix} = 0$  1

or  $1(9) - (x-2)(7) + 4(3) = 0$  or  $x = 5$ . 1½  
[CBSE Marking Scheme 2017]

Q. 13. Let  $\vec{a} = \hat{i} + \hat{j} + \hat{k}, \vec{b} = \hat{i}$  and  $\vec{c} = c_1\hat{i} + c_2\hat{j} + c_3\hat{k}$ , then

(i) Let  $c_1 = 1$  and  $c_2 = 2$ , find  $c_3$  which makes  $\vec{a}, \vec{b}$  and  $\vec{c}$  coplanar.

(ii) If  $c_2 = -1$  and  $c_3 = 1$ , show that no value of  $c_1$  can make  $\vec{a}, \vec{b}$  and  $\vec{c}$  coplanar. **R&U [Delhi 2017]**

Sol.  $[\vec{a} \ \vec{b} \ \vec{c}] = \begin{vmatrix} 1 & 1 & 1 \\ 1 & 0 & 0 \\ c_1 & c_2 & c_3 \end{vmatrix} = c_2 - c_3$  1

(i)  $c_1 = 1, c_2 = 2$

$[\vec{a} \ \vec{b} \ \vec{c}] = 2 - c_3$  1

$\therefore \vec{a}, \vec{b}, \vec{c}$  are coplanar  $[\vec{a} \ \vec{b} \ \vec{c}] = 0$  or  $c_3 = 2$  1

(ii)  $c_2 = -1, c_3 = 1$

$[\vec{a} \ \vec{b} \ \vec{c}] = c_2 - c_3 = -2 \neq 0$

or No value of  $c_1$  can make  $\vec{a}, \vec{b}, \vec{c}$  coplanar 1

[CBSE Marking Scheme 2017]

Q. 14. If four points  $A, B, C$  and  $D$  with position vectors  $4\hat{i} + 3\hat{j} + 3\hat{k}, 5\hat{i} + x\hat{j} + 7\hat{k}, 5\hat{i} + 3\hat{j}$  and  $7\hat{i} + 6\hat{j} + \hat{k}$  respectively are coplanar, then find the value of  $x$ . **R&U [Delhi Comptt. 2017]**

Sol.  $\vec{AB} = \hat{i} + (x-3)\hat{j} + 4\hat{k}$

$\vec{AC} = \hat{i} - 3\hat{k}$

$\vec{AD} = 3\hat{i} + 3\hat{j} - 2\hat{k}$  1½

As  $A, B, C$  and  $D$  are coplanar

$\therefore \vec{AB} \cdot (\vec{AC} \times \vec{AD}) = 0$

i.e.,  $\begin{vmatrix} 1 & x-3 & 4 \\ 1 & 0 & -3 \\ 3 & 3 & -2 \end{vmatrix} = 0$  1½

$9 - (x-3)(7) + 12 = 0$   
which gives

$x = 6$  1  
[CBSE Marking Scheme 2017]

## ? Long Answer Type Question-II

(6 marks each)

Q.1. If  $\vec{a}, \vec{b}, \vec{c}$  are unit vectors such that  $\vec{a} \cdot \vec{b} = \vec{a} \cdot \vec{c} = 0$  and the angle between  $\vec{b}$  and  $\vec{c}$

is  $\frac{\pi}{6}$ , then prove that :

(i)  $\vec{a} = \pm 2(\vec{b} \times \vec{c})$

(ii)  $[\vec{a} + \vec{b} \ \vec{b} + \vec{c} \ \vec{c} + \vec{a}] = \pm 1$ . [A] [S.Q.P. 2015-16]

Sol. (i) As given

$$\vec{a} \cdot \vec{b} = 0, \vec{a} \cdot \vec{c} = 0 \Rightarrow \vec{a} \perp \text{both } \vec{b} \text{ and } \vec{c}$$

(as  $\vec{a}, \vec{b}, \vec{c}$  are non-zero vectors) 1

or  $\vec{a} \parallel (\vec{b} \times \vec{c})$

Let  $\vec{a} = \lambda(\vec{b} \times \vec{c})$ ,

then  $|\vec{a}| = |\lambda| |(\vec{b} \times \vec{c})|$

or  $\frac{|\vec{a}|}{|(\vec{b} \times \vec{c})|} = |\lambda|$

or  $|\lambda| = \frac{1}{\sin \frac{\pi}{6}} = 2$

$\therefore \lambda = \pm 2$

$\therefore \vec{a} = \pm 2(\vec{b} \times \vec{c})$

Hence proved. 2

(ii) Now  $[\vec{a} + \vec{b} \ \vec{b} + \vec{c} \ \vec{c} + \vec{a}]$

$$= [(\vec{a} + \vec{b}) \times (\vec{b} + \vec{c})] \cdot (\vec{c} + \vec{a})$$

$$= (\vec{a} \times \vec{b}) \cdot \vec{c} + (\vec{b} \times \vec{c}) \cdot \vec{a} \quad 1$$

(As the scalar triple product = 0, if any two vectors are equal)

Hence,

$$\vec{a} \cdot (\vec{b} \times \vec{c}) + (\vec{a} \times \vec{b}) \cdot \vec{c} = \vec{a} \cdot (\vec{b} \times \vec{c}) + \vec{c} \cdot (\vec{b} \times \vec{c})$$

$$= 2 \vec{a} \cdot (\vec{b} \times \vec{c}) \quad 1 + \frac{1}{2}$$

$$= 2 \vec{a} \cdot \left( \pm \frac{1}{2} \vec{a} \right)$$

$$= \pm 1 \quad \frac{1}{2}$$

Hence proved.

[CBSE Marking Scheme 2015]

Q.2. If  $\hat{a}$  and  $\hat{b}$  are unit vectors inclined an angle  $\theta$ ,

then prove that  $\tan \frac{\theta}{2} = \frac{|\vec{a} - \vec{b}|}{|\vec{a} + \vec{b}|}$ .

Sol. Given  $|\vec{a}| = |\vec{b}| = 1$  &  $\theta$  is angle between  $\vec{a}$  and  $\vec{b}$

$$(\vec{a} - \vec{b}) \cdot (\vec{a} - \vec{b}) = \vec{a} \cdot \vec{a} - \vec{a} \cdot \vec{b} - \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{b}$$

$$= |\vec{a}|^2 - 2\vec{a} \cdot \vec{b} + |\vec{b}|^2$$

$$= 1 - 2\vec{a} \cdot \vec{b} + 1 \quad 1$$

$$|\vec{a} - \vec{b}|^2 = 2 - 2\vec{a} \cdot \vec{b}$$

$$= 2 - 2|a||b| \cos \theta$$

$$|\vec{a} - \vec{b}|^2 = 2(1 - \cos \theta) = 2 \left( 2 \sin^2 \frac{\theta}{2} \right) \quad 1$$

$$|\vec{a} - \vec{b}|^2 = 4 \sec^2 \frac{\theta}{2}$$

$$2 \sin \frac{\theta}{2} = |\vec{a} - \vec{b}|$$

$$\sin \frac{\theta}{2} = \frac{1}{2} |\vec{a} - \vec{b}| \quad 1$$

Now  $(\vec{a} + \vec{b}) \cdot (\vec{a} + \vec{b}) = \vec{a} \cdot \vec{a} + \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{a} + \vec{b} \cdot \vec{b}$

$$|\vec{a} + \vec{b}|^2 = |\vec{a}|^2 + 2\vec{a} \cdot \vec{b} + |\vec{b}|^2$$

$$|\vec{a} + \vec{b}|^2 = 1 + 2\vec{a} \cdot \vec{b} + 1 \quad \frac{1}{2}$$

$$|\vec{a} + \vec{b}|^2 = 2 + 2|a||b| \cos \theta$$

$$|\vec{a} + \vec{b}|^2 = 2(1 + \cos \theta)$$

$$|\vec{a} + \vec{b}|^2 = 4 \cos^2 \frac{\theta}{2} \quad \frac{1}{2}$$

$$|\vec{a} + \vec{b}|^2 = 2 \cos \frac{\theta}{2}$$

$$\cos \frac{\theta}{2} = \frac{1}{2} |\vec{a} + \vec{b}|$$

$$\tan \frac{\theta}{2} = \frac{|\vec{a} - \vec{b}|}{|\vec{a} + \vec{b}|} \quad \text{Hence Proved. 2}$$

Q.3. If with reference to right handed system of mutually perpendicular unit vectors  $\hat{i}, \hat{j}$  and  $\hat{k}$

and  $\vec{\alpha} = 3\hat{i} - \hat{j}$ ,  $\vec{\beta} = 2\hat{i} + \hat{j} - 3\hat{k}$ , then express in

the form  $\vec{\beta} = \vec{\beta}_1 + \vec{\beta}_2$ , where  $\vec{\beta}_1$  is parallel to  $\vec{\alpha}$

and  $\vec{\beta}_2$  is perpendicular to  $\vec{\alpha}$ .

Sol. Given  $\vec{\beta}_1 \parallel \vec{\alpha}$

$$\therefore \vec{\beta}_1 = \lambda \vec{\alpha}$$

$$\vec{\beta}_1 = \lambda(3\hat{i} - \hat{j}) \quad 2$$

$\vec{\beta}_2$  is  $\perp$  to  $\vec{\alpha}$

$$\therefore \vec{\beta}_2 \cdot \vec{\alpha} = 0$$

$$\text{Given } \vec{\beta} = \vec{\beta}_1 + \vec{\beta}_2 \Rightarrow \vec{\beta}_2 = (2\hat{i} + \hat{j} - 3\hat{k}) - \lambda(3\hat{i} - \hat{j}) \quad 2$$

$$\vec{\beta}_2 = (2 - 3\lambda)\hat{i} + (1 + \lambda)\hat{j} - 3\hat{k}$$

$$\vec{\beta}_2 \cdot \vec{\alpha} = [(2 - 3\lambda)\hat{i} + (1 + \lambda)\hat{j} - 3\hat{k}] \cdot [3\hat{i} - \hat{j}] = 0$$

$$3(2 - 3\lambda) - (1 + \lambda) = 0$$

$$5 - 10\lambda = 0$$

$$\lambda = \frac{1}{2}$$

$$\vec{\beta}_1 = \frac{1}{2}(3\hat{i} - \hat{j}) \quad \& \quad \vec{\beta}_2 = \frac{1}{2}\hat{i} + \frac{3}{2}\hat{j} - 3\hat{k} \quad 2$$

#### Commonly Made Errors

- Some students do dot product first then cross product which is wrong.
- The right method is to do the cross product first then dot product in scalar triple product.

 <b>OSWAAL LEARNING TOOLS</b>	
<b>For Suggested Online Videos</b>	
Visit : <a href="https://qr.go.page.link/7gJLR">https://qr.go.page.link/7gJLR</a>	Visit : <a href="https://qr.go.page.link/amYe3">https://qr.go.page.link/amYe3</a>
 Or Scan the Code	 Or Scan the Code

