DAY THIRTY SIX

Biodiversity, its Conservation and Environmental Issues

Learning & Revision for the Day

Concept of Biodiversity

Loss of Biodiversity

- Biodiversity Conservation
- Patterns of Biodiversity
 Red Data Book
 - Important Wildlife Projects of India
- Pollution
- Deforestation
- Case studies

Concept of Biodiversity

- Biodiversity is the heterogeneity which exists at different levels of biological organisations in our biosphere.
- The term **biodiversity** was popularised by the sociobiologist Edward Wilson.
- The scientific estimation of total number of species made by **Robert Mayer** is about 7 million.
- More than 70% of all the species recorded are animals, while plants (algae, fungi, bryophytes, gymnosperms and angiosperms) comprise not more than 22% of the total.
- Among animals, insects are the most species-rich taxonomic group making up more than 70% of the total.
- Fungi species in the world are more than the combined total of the species of fishes, amphibians, reptiles and mammals, i.e. they represent maximum number of species among global biodiversity.
- Maximum nutritional diversity is found in the group Monera.
- India shares 8.1% of global species diversity and it makes India one of the 12 mega diversity countries of the world.
- The biological diversity includes three hierarchical levels
 - (i) **Genetic diversity** is the diversity in the number and types of genes present in different species and the genetic variations in the same species.





For example,

- (a) Genetic variation in medicinal plant *Rauwolfia vomitoria* growing in different Himalayan ranges occurs in terms of the potency and concentration of reserpine (active chemical).
- (b) In India, rice shows maximum genetic diversity with more than 50,000 different strains.
- (ii) Species diversity is the variety of species within a region. It indicates the species richness in any habitat. For example, the Western Ghats have a greater amphibian species diversity than the Eastern Ghats.
- (iii) Ecological diversity is the diversity at ecosystem level. Diversity at the level of community and ecosystem has three perspectives
 - α-diversity also called as local diversity, is the diversity within a community.
 - β-diversity is the diversity between two communities.
 - γ-diversity, also called as regional diversity, represents the total richness of species in all the habitats found within a geographical region or landscape.

Patterns of Biodiversity

Species diversity decreases as we move away from the equator towards the poles.

The main reasons for more diversity in tropics than temperate regions are

- (i) Unlike temperate regions subjected to frequent glaciation for much time period in the past, tropics remained relatively undisturbed for millions of years and favours species diversification.
- (ii) Tropical environments are less seasonal, relatively more constant and predictable, which promote niche specialisation and lead to a greater species diversity.
- (iii) There is more solar energy available in the tropics, which contributes to higher productivity and inturn greater diversity.

Therefore, according to non-uniform diversity on the earth, the biodiversity is divided into following patterns as

- (i) In latitudinal gradients, tropics (latitudinal range of 23.5°N to 23.5°S) harbour more species than temperate or polar areas.
 - India with much of its land area in the tropical latitudes, has more than 1200 species of birds.
 - The **tropical Amazonian** rainforest in South America has the greatest biodiversity on earth.
- (ii) Species-area relationship was first described by Alexander von Humboldt. He observed that within a region, species richness increase with increasing explored areas, but only upto a limit.
 - On a logarithmic scale, the relationship is a straight line described by the equation



Loss of Biodiversity

Area

IUCN (International Union for Conservation of Nature and Natural Resources), Red List (2004) documented the extinction of 784 species (338 vertebrates, 359 invertebrates and 87 plants) in the last 500 years.

The major causes of biodiversity loss (The Evil Quartet) are

- (i) Habitat loss and fragmentation It results due to over population, urbanisation, industrialisation, etc.,
 e.g. 90% of wetlands of New Zealand have been destroyed by European settlers.
- (ii) Overexploitation Excessive exploitation of a species reduces its population size, so that it becomes vulnerable to extinction, e.g. Dodo, passenger pigeon, three subspecies of Tiger (Bali, Javan and Caspian) and Stellar's sea cow have become extinct due to overexploitation by humans.
- (iii) Alien species invasion Non-native or alien species often become invasive and drive away the local population.

For example,

CLICK HERE

- *Eichhornia crassipes* killed several aquatic plants and animals.
- Nile perch killed small cichlid fishes in lake Victoria of South Africa.
- *Eupatorium hysterophorus* has reduced the population of *Tectona grandis* in North-East.
- *Parthenium hysterophorus* has pushed several herbs and shrubs from the plains.
- African catfish, *Clarias gariepinus* has threatened native catfishes, *Clarias bacterachus* in Indian rivers.
- (iv) **Coextinction** Extinction of one species causes extinction of the other due to obligatory mutualistic relation between them, e.g. *Pronuba yuccaselles* and *Yucca*.

Biodiversity Conservation

Conservation means protection, upliftment and scientific management of biodiversity, so as to maintain it at its optimum level and derive sustainable benefits for the present as well as future generations

🕀 www.studentbro.in

The following are the three major reasons to conserve biodiversity

- 1. Narrowly utilitarian The useful human products like food, fibres, drugs and medicines are obtained from biodiversity.
- 2. **Broadly utilitarian** Biodiversity provides ecosystem services like providing oxygen, pollinating crops and controlling floods, erosions, etc.
- 3. Ethical utilitarian Every living species has an intrinsic value, through it may not have direct economic value and also every species has right to live.
- The Convention of Biodiversity (CBD) came into force on 29th December, 1993. It has three objectives conservation, sustainable use and equitable sharing of benefits of biodiversity.
- Biodiversity Act of India was passed by the Parliament in 2002.
- Insularization is the reduction of species diversity in small patches.
- World Wildlife Fund (WWF) was established in 1961 at Switzerland and giant panda (*Ailuropoda melanoleuca*) was selected as its symbol. It aims at protection and preservation of wild plants and animals.
- World Wildlife week is celebrated in first week of October.

There are two major methods of biodiversity conservation, as discussed below

In Situ Conservation

It is the conservation of living resources through their maintenance within the natural ecosystem in which they occur. These include hotspots and Protected Area Network (PAN). PAN includes sacred lands, biosphere reserves, national parks and wildlife sanctuaries.

- (i) Hotspots The concept of hotspot was given by Norman Myers in 1988. Hotspots are the areas that are extremely rich in species diversity, have high endemism, lesser interspecific competition and are under constant threat.
 - Among the 34 hotspots (cover less than 2% of earth land area) of the world, two are found in India extending into neighbouring countries. These two are the **Western Ghats/Sri Lanka** and the **Indo-Burma Region** (covering the Eastern Himalayas also known as cradle of speciation).

The two hotspots in India are as follows

• **Eastern himalayas** It extends to the North-Eastern India and Bhutan. The temperate forests are found at altitudes of 1780-3500 metres.

Many deep and semi-isolated valleys found in this region are exceptionally rich in endemic plant species.

Besides being an active centre of evolution and rich diversity of flowering plants, the numerous primitive angiosperm families, e.g. Magnoliaceae and Winteraceae and primitive genera of plants, like *Magnolia* and *Betula* are found in Eastern Himalayas.

- Western Ghats This region lies parallel to the Western coast of Indian Peninsula for almost 1600 km in Maharashtra, Karnataka, Tamil Nadu and Kerala. These regions are rich in amphibians. The forests at low elevation (500 m above mean sea level) are mostly evergreen, while those found at 500-1500 m height are generally semi-evergreen forests.
- (ii) Biosphere reserves were introduced under MAB (Man And Biosphere) programme of UNESCO. Biosphere reserve programme was started in India in 1986. Total biosphere reserves in India are 14.
 - The first biosphere reserve established in 1986 was Nilgiri Biosphere Reserve.
 - A biosphere reserve is made of core, buffer and manipulation zone.
 - In the core zone, no human activity is allowed and hence, the area remains undisturbed and legally protected.
 - In the buffer zone, limited human activities are involved for resource use strategies, research and education.
 - In the manipulation zone, active cooperation is present between reserve management and local people for cropping, settlements, etc.

Some Biosphere Re	eserves in	India
-------------------	------------	-------

Site	Location (State)	Year	Area in km ²
Nilgiri	Karnataka, Kerala and Tamil Nadu	1986	5520
Nanda Devi	Uttarakhand	1988	5860
Norek	Meghalaya	1988	820
Manas	Assom	1989	2837
Sunderbans	West Bengal	1989	9630

- (iii) **National parks** are an area strictly reserved for the protection and welfare of wildlife.
 - In India, there are 96 national parks (April 2007) covering an area of 1.16% of India's total surface area.
 - The first national park in the world, the **Yellowstone National Park**, was founded in 1872 in USA.
 - In 1935, the first national park of India was established in the foothills of the Himalayas (Hailey National Park) presently known as **Corbett National Park.**
 - A total of 166 National Parks have been authorised.



Some National Parks in India

Name	State	Famous for
Bandipur National Park	Karnataka	Elephant and tiger
Corbett National Park	Uttarakhand	Tiger
Dachigam National Park	Jammu and Kashmir	Hangul (Kashmir stag)
Gir National Park	Gujarat	Asiatic lion
Kanha National Park	Madhya Pradesh	Tiger
Kaziranga National Park	Assom	One-horned rhinoceros
Kanchenjunga National Park	Sikkim	Tiger and elephants
Madhav National Park	Madhya Pradesh	Chital, deer
Mahavir Harina Vanasthali National Park	Rajasthan	Deer
Manas National Park	Assom	Wild water buffalo
Keibul Lamjao National Park	Manipur	Brow antlered deer
Periyar National Park	Kerala	Elephant
Rajaji National Park	Uttarakhand	Elephant
Sariska National Park	Rajasthan	Tiger
Silent Valley National Park	Kerala	Lion tailed macaque
Sundarbans National Park	West Bengal	Royal Bengal tiger
Tadoba National Park	Maharashtra	Tiger

(iv) Sanctuaries are notified for the protection of wild animal and fauna. In India, there are over 500 sanctuaries in different states.

Among these, the 28 tiger reserves are governed by Project Tiger, one of special significance in the conservation of the tiger. Some wildlife sanctuaries are specifically named bird sanctuary.

Some Important Sanctuaries of India

Name and Location	Area (in sq km)	Key Vertebrate Species being Protected
Chilka Lake (Odisha)	990	Flamingoes, sandpipers, ducks, water fowls, cranes, golden plovers and ospreys.
Keoladeo Ghana Bird Sanctuary (Rajasthan)	29	Siberian crane, spoon bill, herons, egrets and variety of other local birds. Blue bull, wild boar, black buck and spotted deer.

Name and Location	Area (in sq km)	Key Vertebrate Species being Protected
Mudumalai Wildlife Sanctuary, Nilgiri (Tamil Nadu)	520	Flying squirrel, porcupine, elephant, sambhar, cheetal, barking deer, mouse, deer, four-horned antelope, giant squirrel flying lizard and monitor lizard.
Manas Wildlife Sanctuary, Kamrup (Assom)	_	Tiger, wild boar, sambhar, golden langoor, one-horned rhino, panther, swamp deer, wild dog and wild buffalo.
Periyar Sanctuary (Kerala)	777	Elephants, leopard, black langoor, sambhar, gaur, bison, Egret and horn bills.
Sultanpur Lake Bird Sanctuary (Uttar Pradesh)	12	Cranes, duck, green pigeon, drake and spot bill, <i>Python</i> and crocodile.

(v) Sacred forests are undisturbed forests without any human intervention and highly surrounded by degraded lanseales. These forests contain number of rare, endangered and endemic species. Such sacred groves are found in

- Khasi and Jaintia Hills in Meghalaya
- Aravali Hills of Rajasthan
- Western Ghat regions of Karnataka and Maharashtra
- Sarguja, Chanda and Bastar areas of Madhya Pradesh

Ex Situ Conservation (Off-site Conservation)

CLICK HERE

It means conservation outside the habitats by perpetuating sample population in genetic resource centres.

- **Botanical garden** is a collection of living plants maintained for both pure and applied studies.
- Wildlife safari parks are used for *ex situ* conservation of threatened animals and plants.
- In **seed banks**, germplasm is stored as seeds of various accessions. Under suitable conditions, seeds of many species can be stored for upto 50-100 year.
- **Pollen storage** is considerably important in the conservation of genetic diversity. The life of pollen has been reported 3 years in some species. They can be stored for several years in liquid nitrogen having temperature -196°C.
- **Tissue culture** can be extended to endangered species as well as those which may otherwise require very rapid climatic condition and can be maintained at one place in aseptic cultures.

🕀 www.studentbro.in

• **Zoo** is a place, where wild animals are kept for public showing. They have recorded success with captive breeding of animals.

Some Zoos in India

Name	City	State
Indira Gandhi Zoological Park	Visakhapattnam	Andhra Pradesh
Nehru Zoological Park	Hyderabad	Andhra Pradesh
Arignar Anna Zoological Park	Chennai	Tamil Nadu
Sri Chamarajendra Zoological Park	Mysore	Karnataka

Extinction of Species

The total elimination or dying out of a particular species from the earth leads to extinction of that species. Population traits, which make a species susceptible to extinction are

- (i) Large body size, e.g. elephant, rhinoceros and lion.
- (ii) Small population size.
- (iii) Low reproductive potential, e.g. blue whale, giant panda.
- (iv) Higher states of tropic level, e.g. Bengal tiger, bald eagle.

Red Data Book

A **Red Data Book** or **Red List** is a catalogue of taxa facing risk of extinction.

IUCN is International Union for Conservation of Nature and Natural Resources, which is now called World Conservation Union (WCU) maintains it.

The main objective of IUCN is to promote and support the conservation of wildlife and natural resources. WCU has its headquarter at **Morgan**, **Switzerland**.

Red List has following categories of species

- (i) **Threatened species** liable to be extinct if not allowed to realise it's full biotic potential by providing protection from exotic species, e.g. mountain gorilla, Giant panda, rhinoceros, etc.
- (ii) A taxon is **extinct**, when there is no reasonable doubt that it's last individual has died, e.g. passenger pigeon, Dodo, etc.
- (iii) A taxon is extinct in the wild, when it is known only to survive in cultivation, in capacity or as a naturalised population, well outside the past range, e.g. Hawain crow, Pinta Island tortoise, etc.
- (iv) A taxon is critically endangered, when it is facing an extremely high risk of extinction in the wild in the immediate future (925 animals and 1014 plants), e.g. Great Indian Bustard, Sumatran orangutan, etc.
- (v) A taxon is **endangered**, when it is not critically endangered but facing a very high risk of extinction in the

wild in the near future, e.g. Blue whale, black winged Indonesian parrot, etc.

- (vi) A taxon is vulnerable, when it is not critically endangered or endangered but is facing a high risk of extinction in the wild in the medium term future, e.g. Madagascar frog (Dyscophus antongilii), black-buck (Antilope cervicapra).
- (vii) Taxa that do not currently qualify as critically endangered, or vulnerable and may be classified as **conservation dependent**.
- (viii) A taxon is **data deficient**, when there is inadequate information to make a direct or indirect assessment of it's risk of extinction based on it's distribution or population status.
- (ix) A taxon is under the category **non-evaluated** when it has not yet been assessed against any criteria.

Important Wildlife Organisations of the World

- CITES Convention on International Trade in Endangered Species of Wild Fauna and Flora
- IBWL Indian Board for Wildlife
- IUCN International Union for Conservation of Nature and Natural Resources
- NWAP National Wildlife Action Plan
- UNCED United Nations Conference on Environment and Development
- WPSI Wildlife Preservation Society of India
- WWF World Wildlife Fund

CLICK HERE

• NEERI National Environment Engineering Research Institute

Important Wildlife Projects of India

- **Project tiger** (*Panthera tigris*) The project was started in 1973 in order to check depletion in population of tiger. Initially, it was undertaken in 17 National Parks. But recently the project has been extended to more National Parks (a total of 23).
- **Project lion** (*Panthera leo persica*) The project was started in 1972. It is located in Gir National Park, Junagarh (Gujarat).
- **Project snow leopard** (*Panthera uncia*) Throughout Himalayas, e.g. Khangchendzonga National Park (Gangtok).
- **Project musk deer** (*Moschus moschiferus*) Kedarnath Sanctuary (Uttarakhand), Manali Sanctuary (HP) and Shikari Devi Sanctuary (HP).
- **Project elephant** It was launched in 1992 and it covers both wild and domestic elephants.

Environmental issues include the aspects which adversely affect our biophysical environment. Pollution, global warming, deforestation, etc., are the topics of major concern in current perspective.

🕀 www.studentbro.in

Pollution

- Any undesirable change in the physical, chemical or biological characteristics of the atmosphere (air), lithosphere (land) and hydrosphere (water), which is harmful to living organisms directly or indirectly is called **pollution**.
- Pollutants are divided into many types on the basis of several categories as given below
 - (i) On the basis of natural degradation or disposal, pollutants are of two types
 - Biodegradable Degrades after sometime either automatically, e.g. by heat or through the agency of microorganisms, e.g. sewage, domestic wastes, etc.
 - Non-biodegradable Not degraded by living organisms. These are the most harmful environmental pollutants, e.g. DDT, glass, plastic, pesticides, radioactive substances, heavy metals like mercury, lead, cadmium, etc.
 - (ii) On the basis of persistance, pollutants are of two types
 - Primary pollutant Persist in the atmosphere in same form in which they are released, e.g. CO, DDT, plastic ware, etc.
 - Secondary pollutant Formed by the interaction between primary pollutants and are more toxic than the primary pollutants, e.g. PAN, ozone, HNO₃, H₂SO₄, etc.

Types of Pollution

On the basis of location, reservoir or the area where the pollutants are present. The pollution can be of air pollution, water pollution, soil pollution, etc.

Air Pollution and its Control

It is undesirable change in the natural characteristics of the atmosphere due to contamination by any chemical, biological or physical agent.

Sources of Air Pollution

Various air pollutants, their origin and effects are described below

• **Particulate matter** These are solid or liquid particles found in air. Settleable particulate, (about $\geq 10 \,\mu$ m) settle out in less than a day. Suspended particulate (about $\leq 10 \,\mu$ m) remain suspended for weeks, e.g. aerosol, dust, mist, etc. Particles of 2.5 μ m or less in diameter cause greater harm to humans (as per CPCB–Central Pollution Control Board).

These cause breathing and respiratory problems, irritation, inflammation and damage to lungs.

• **Fluorocarbon** Chemicals released with force in the form of mist or vapour, into the atmosphere. Released by jet planes and also called as **aerosols**.

- **Smog** It is a mixture of fog and smoke which occurs in some busy industrical cities. It can be of two types
- Photochemical/Los Angeles Smog Nitrogen oxides, hydrocarbons and O₂ interact in the presence of sunlight to form ozone (corrodes the heritage building surfaces and marble statues) and Peroxy Acetyl Nitrate (PAN) which inhibits ETC as well as damage chloroplast.
- Classical/London Smog Formed from reaction of smoke, dust, H₂S and SO₂. Also known as sulphur smog, it causes stone cancer.
- **Smoke** It is obtained by incomplete combustion of carboneous material and smoke stacks of thermal power plants.
- **Carbon Monoxide** (CO) It is formed due to incomplete combustion of fuels from motor vehicles and industries. It Combines with haemoglobin to form carboxyhaemoglobin (COHb) which reduces O₂ carrying capacity of blood.
- Sulphur Dioxide (SO₂) Gaseous pollutant released by combustion of sulphur containing fossil fuels, smelting of ore and from oil refineries. It causes membrane damage, inhibits electron

transportation in plants and leads to respiratory problems in organisms.

• **Nitrogen Oxide** (NO₂) Released by combustion of fossil fuels at high temperature in automobile engines. Forms brown air that leads to various heart and lung problems.

Methods of Controlling Air Pollution

The air pollutants must be separated out before releasing the harmless gases into the atmosphere, which can be done by adopting these methods

- Electrostatic Precipitator (ESP) It is the most commonly used method for the removal of particulate matter. About 99% of particulate matter are removed from the exhaust of thermal power plant.
- **Scrubber** A scrubber can remove gases like sulphur dioxide from the industrial exhaust. The exhaust is passed through a spray of water or lime which reacts with sulphur dioxide to form a precipitate of calcium sulphate and sulphide.
- **Catalytic converters** They have expensive metals namely platinum, palladium and rhodium as the catalysts and are fitted into automobiles for reducing emission of poisonous gases.

As the exhaust passes through the catalytic converter, unburnt hydrocarbons are converted into CO_2 and water and carbon monoxide and nitric oxide are changed to carbon dioxide and nitrogen, oxygen gas respectively. Motor vehicles equipped with catalytic converter should use unleaded petrol because lead in the petrol inactivates the catalyst.

🕀 www.studentbro.in

CLICK HERE

• **Government's norms for emission** In the line of world standard, Government of India also formulate new fuel policy time to time. These fuel policies with their applicable regions are given below

Fuel policies

	1 401	Policios	
Standard	Reference	Year	Region
India 2000	Euro 1	2000 - 01	NCR*, Mumbai, Kolkata, Chennai
Bharat Stage II	Euro 2	2003 - 04	NCR*, 10 cities +
		2005 - 04	Nationwide
Bharat stage III	Euro 3	2005 - 04	NCR*, 10, cities +
		2010 - 05	Nationwide
Bharat stage IV	Euro 4	2010 - 04	NCR*, 10 cities +

* National Capital Region (Delhi).

+ Mumbai, Kolkata, Chennai, Bengaluru, Hyderabad, Ahmedabad, Pune, Surat, Kanpur and Agra.

Effects of Air Pollution on Environment

Air pollution plays a major role indamaging environment and climate in the following ways:

Acid Rain (pH ≤ 5.65)

- The term acid rain was first proposed by **Robert August**. Precipitation of oxides of sulphur and nitrogen along with rain is termed as acid rain.
- It takes place when SO₂ and oxides of nitrogen such as N₂O (nitrous oxide) and NO (nitric oxide) present in air dissolve in rain water to form sulphuric acid (H₂SO₄) and nitric acid (HNO₂), respectively.
- Effects of acid rain are given below:
 - Effect on aquatic life Acidic deposition adversely affect the aquatic life by making water acidic. The water body in which the biodiversity is reduced at significant level, is called **Biologically dead** (completely eutrophic).
 - Effects on forest Acid deposition affect the forests negatively, e.g. the red spruce forest in tropical areas are killed severely.
 - Effects on building and monuments The oldest building and monuments all over the world are destroyed by atmospheric acid at an alarming rate, e.g. Taj Mahal.

Greenhouse Effect and Global Warming

The term greenhouse effect was coined by **Arrhenius**. It is a naturally occurring phenomenon that is responsible for heating of Earth's surface and atmosphere due to the presence of certain gases in the atmosphere.

Brief description of chief Greenhouse Gases (GHGs) with their sources and effects are given below

- (i) Carbon Dioxide (CO₂) Present level in atmosphere is 380 ppm (parts per million). Atmospheric, lifetime is 5-200 yr. Its amount in atmosphere is increasing due to fossil fuel's burning, deforestation and change in land use. High concentration may cause effects such as increase in rate of photosynthesis and growth of plants, decrease in stomatal conductance and transpiration rate, etc.
- (ii) Nitrous Oxide (N_2O) Present atmospheric concentration is 316 ppb (parts per billion). Major sources are agriculture, biomass burning, nylon industries, nitrogen rich fertilisers and fuels.
- (iii) **Methane** (CH_4) Present level in atmosphere is 1750 ppb (parts per billion). **Methanogen** bacteria increase greenhouse effect by producing methane. The major sources are freshwater wetlands, enteric fermentation in cattle. Flooded rice fields along with biomass burning.
- (iv) Chlorofluorocarbons (CFCs) Present atmospheric concentration is 282 ppt (part per trillion). Atmospheric lifetime is 45-260 yr. Major sources are leakage from air conditioners, refrigeration units, evaporation of industrial solvents, production of plastic foams and propellants in aerosol, spray cans.
- The gradual continuous increase in average temperature of surface of the earth as a result of increase in concentration of greenhouse gases is termed as global warming.
- During the past century, the temperature of earh has increased by 0.6°C, most of it during the last three decades.
- Effects of global warming are as follows
 - Earth's temperature has increased by 0.6°C in last three decades. This causes change in precipitation patterns.
 - The rise in temperature leads to harmful effects in environment leading to odd climatic changes, e.g. El Nino effect.
 - The high temperature will result in melting of polar ice caps, which will lead to rise in sea level and many coastal areas will be submerged.
 - The high levels of temperature lead to increased weed growth, eruption of diseases and pests. Thus, the crop productivity will decrease.
 - Reducing use of fossile fuels, improving efficiency of energy usage, reducing deforestation, planting trees and slowing down the growth of human population can control the global warming.

Ozone Depletion

CLICK HERE

• Ozone layer present in stratosphere protects us from harmful ultraviolet radiations coming from sun. Ozone gas is continuously formed by the action of UV rays on molecular

🕀 www.studentbro.in

oxygen. The thickness of ozone is measured in **Dobson Units** (DU). Ozone can be

- **Bad ozone** Formed in troposphere and is harmful to plants and animals.
- **Good ozone** Present in stratosphere and acts as a shield, absorbing harmful UV radiations from the sun.
- This ozone layer is being destroyed by some pollution like CFCs. Harmful effects of ozone depletion are as follows
 - UV-B damages DNA, causing mutation.
 - Ageing of skin, damage to skin cells and various types of skin cancer.
 - High dose of UV-B causes inflammation of cornea. This is called snow-blindness, cataract, etc. Such exposure may permanently damage the cornea.
 - The depletion of ozone is particularly marked over the Antarctic region.
 - Ozone is commonly called as chemical weed.
 - Montreal Protocol, was signed at Montreal (Canada) in 1987 (effective in 1989) to control the emission of ozone depleting substances.
 - Earth Summit (1992), to reduce greenhouse gases.
 - **Kyoto Protocol** (1997), convention on climate change, to reduce **greenhouse emission**.

Water Pollution and its Control

Water pollution is the undesirable presence of some organic, inorganic, biological or physical substances in water which makes it unfit for use.

Sources of Water Pollution

Various water pollutants are described below

- **Toxic Metals** Pb, Zn, Ar, Cu, Cd, Hg, Ni from electroplating, chemical and copper pickling industries inhibit self-purification of water body and lead to chromosomal damage and thus, interfere with heredity.
- Acids HNO₃, H₂SO₄ lowers the pH of water which is lethal to fishes, etc.
- Alkalis pH shocks, due to addition of alkaline water causes asphyxiation of fishes.
- **Coal** It interferes with self-purification of river water.
- **Dyes** These increase the BOD and change colour of water.
- Gaseous pollutants Major gaseous pollutants include
 - NH₃ produced in fertiliser industry causes irritation, pulmonary oedema.
 - Cl Causes pulmonary oedema, corrosive, fatal for fishes.
 - = H_2S Produces noxious smell, irritation, respiratory depressant.

- Industrial agents Major industrial agents include
 - Paper and pulp Produces free chlorine
 - Textile Minor acids, fats, oils and grease.
 - Food Processing produces starch.
 - **Chemicals** Like mineral acids OH, NH₃, tartaric acid and nitrocompounds P, S, F.
 - **Metals** Like fluorides, cyanogen and limestone are called nuisance.
- **Fertilisers** Phosphatic, nitrogenous fertilisers and potash when mixed with water through run off causes water pollution.
- **Pesticides** DDT, 2, 4-D, TEPP, Aldrin, BHC parathion do not degrade (biomagnification).

Effects of Water Pollution

The pollutants of water owing to their toxic effects have diverse effects on humans, environment and aquatic life, etc.

- Major toxic elements found in water and their various effects are as follows
 - Aluminium Interferes with phosphate metabolism, inhibit absorption of fluorides, Ca and iron compounds.
 - **Arsenic** Loss of appetite, copious secretion of mucus in respiratory tract, black foot disease.
 - Cadmium Itai-itai disease (Japan), kidney damage.
 - **Fluorine** Fluorosis, about 5-12 ppm is toxic, enamel becomes brittle, bones lose their elasticity and are prone to fractures, impairs glycolysis, knock-knee disease.
 - Lead Anaemia and mental retardation due to degenerative changes in motor nerves.
 - Mercury Minamata disease, main site of injury is CNS leading to tremors inability to coordinate, impairment of vision and loss of hearing.

Two major episodes of mercury poisoning have occurred in Japan, in Minamata bay and Niigata. Mercury was absorbed, bioaccumulated and biomagnified to high levels. Fish collected from this bay had 10-12 mg of Hg per kg of their flesh and bones. The largest mercury epidemic occurred in 1971-72 in Iraq when 6000 people were affected and 500 died also. It causes infertility in human.

- Some important effect of water pollution on water bodies are as follows
 - Dissolved Oxygen (DO) It decreases due to increase in domestic sewage, toxic pollutants and microbes, thus threatening aquatic life. It decreases with rise in temperature.
 - **Biological Oxygen Demand** (BOD) It is defined as amount of oxygen (in mg) required to decompose organic matter present in one litre of water. It is expressed in parts per million (ppm). Its value increases when sewage is mixed with water bodies.

🕀 www.studentbro.in

CLICK HERE

- Chemical Oxygen Demand (COD) It indicates the total oxygen required by organic matter (pollutants, etc) in a sample of water for its oxidation by a strong chemical oxidant. Its value is higher than BOD.
- **Eutrophication** Eutrophic (*eu* + *trophic* = truely nourished) water are rich in organisms and organic materials. Eutrophication is an increase in nutrient level and productivity. Along with BOD, eutrophication often results from nutrient enrichment. Sewage, fertiliser, run off and other human activities causing increase in biological productivity is called **cultural eutrophication**.

Methods of Controlling Water Pollution

Water pollution can be controlled through various measures. Some of them are discussed here

- Proper maintenance of water bodies.
- Reduced use of pesticides and chemical fertilisers in agriculture.
- Avoid the disposal of waste into water.
- Proper sewage treatment before disposal into large water bodies.
- Control of disposal of industrial waste into water.

Soil Pollution

It is alteration in soil caused by removal or addition of substances and factors which decrease its productivity, quality of plants and ground water.

Main Soil Pollutants, their Characteristics and Effects

Type of Pollutant	Characteristic and Effect
Pesticides (broad spectrum)	Insecticides (kill insects), fungicides (kill fungi), algicides (kill algal bloom), rodenticides (kill rodents), etc., reduce pest population but also other biota. Degraded products of pesticides are bioaccumulated by plants and passed on to food chain/web.
Fertilisers	Excessive use of fertilisers kills useful microbes, increases salinity, reduces soil productivity.
Manures	Excreta of livestock/human sewage, sludge pollutes/contaminates soil and crops from such areas if consumed can cause important health hazards.
Radioactive waste	Of nuclear plants, laboratories using them and from mining activities gets into soil and causes mutations/genomic changes, acid rain; weathering of rocks pollute soil.

Agro-Chemicals and Their Effects

- The increasing amount of agricultural chemicals like fertilisers can create biological magnification conditions in aquatic and terrestial ecosystems and destroy non-target organisms.
- **Biomagnification** occurs due to persistent pesticides such as DDT which have a long lifetime in the environment.
- They are fat soluble and generally non-biodegradable, therefore they can get incorporated into the food chain and ultimately deposited in the fatty tissues of animals and humans.
- The magnification of these pesticides in successive higher trophic levels is known as **biological magnification**.
- As a result of this, a decline in bird population in a region is observed.
- Agrochemical pollution can be controlled by adapting organic farming. It is a cyclic, zero-waste procedure, where waste products from one process are used as nutrients for other processes. This allows the maximum utilisation of resource and increases the efficiency of production.

Solid Waste Management

Solid waste means everything that goes out in trash, i.e.

- (i) Municipal solid waste includes wastes from homes, offices, schools, etc., that are collected and disposed by the municipality.
- (ii) Fly ash generated by thermal power plants, which is composed of oxides of silica, iron and aluminium.
- (iii) Hospital wastes include hazardous wastes.

 $(iv) \ Industrial wastes include paper, rubber, pesticides, dye, etc.$

Disposal of solid wastes is done by

- Burning the municipal waste to reduce volume.
- Sanitary landfills as open dumps.
- Fly ash is used in construction of industry or buried in landfills.
- E-wastes are buried in landfills.

CLICK HERE

Radioactive Waste Management

The release of radioactive material into environment is called radioactive pollution. Radioactivity is the property of certain elements (radium, thorium, etc) to spontaneously emit alpha (α) particles, beta (β) particles and gamma (γ) rays by disintegration of their atomic nuclei, Nuclear energy is now considered as the most potent pollutant. Earlier, it was assumed to be a natural, non-polluting way for electricity generation.

🕀 www.studentbro.in

- Radioactive pollution is caused due to leakage of radioactive material from thermal power plants or due to unsafe disposal of radioactive wastes.
- It cause mutations at a very high rate. At high doses, nuclear radiation is lethal but a lower doses, it causes various disorders and cancer.
- Nuclear and radioactive waste must be deal with utmost caution, Nuclear waste should be pre-treated and stored in shielded containers and then buried about 500 m deep within the rocks.

Deforestation

- It is the conversion of forested areas to non-forested ones. In **India** at the beginning of 20th century, forests covered about 30% of total land but at the end of the century, it had shrunk to 19.4%.
- National Forest Policy (1988) of India has recommended 33% forest cover for plains and 67% for hills. Cutting of trees for timber, firewood, cattle ranching and Jhum cultivation are responsible for deforestation.
- Slash and burn agriculture, commonly called as Jhum cultivation is common in North-Eastern states of India. In Jhum cultivation, the farmers cut down the trees of the forest and burn the plant remains.

The ash produced is used as a fertiliser and the land is then used for farming or cattle grazing.

- After cultivation, the area is left for several years so as to allow its recovery. The farmers move on to other areas and repeat this process.
- **Reforestation** is the process of restoring a forest that once existed but was removed in the past.

Case Studies

Three case studies addressing environmental issues are given below

- 1. People's Participation in Conservation of Forest
- A Bishnoi woman **Amrita Devi** showed exemplary courage by protecting trees from the men cutting them. She sacrified her

life along with her three daughters, while hugging the trees to protect them from axemen.

- The Government of India has recently instituted the **Amrita Devi Bishnoi Wildlife Protection Award** for the individuals or communities from rural areas that have shown extraordinary courage and dedication in protecting wildlife.
- In 1973, the Chipko Movement was launched by Chandi Prasad Bhatt and Sunder Lal Bahuguna in Chamoli district of Garhwal Himalayas.
- During this movement, local women showed enormous bravery in protecting trees from the axemen of contractor by hugging the trees.
- The Government of India has introduced **Joint Forest Management** (JFM) so as to work closely with the local communities for protecting and managing forests. In return, the communities get benefit of various forest products.

2. Remedy of Plastic Waste

- Ahmed Khan a plastic sack manufacturer in Bengaluru has found and ideal solution to deal with the problem of plastic bags. His company developed polyblend, a fine powder of recycled modified plastic.
- Mixing this powder with bitumen, a blend was prepared, which when laid on the roads enhanced the bitumen's water repellant properties and helped to increase the life of the road three times. Using Khan's technique, by the year 2002, more than 40 km of road in Bengaluru has already been laid.

3. Controlling Vehicular Air Pollution : Delhi

• Delhi leads the country for its high level of air pollution. In 1990, It was the most polluted city of the world. After seeing the serious problem of air pollution, Public Interest Litigatin (PIT) was filed in supreme court of India and the strong steps controlled it to some extent. The result was that the entire fleet of public transport (buses etc.) were converted to run on CNG.



(DAY PRACTICE SESSION 1)

FOUNDATION QUESTIONS EXERCISE

1 Which of the following represents maximum number of species among global biodiversity? → NEET 2018

(a) Algae	(b) Lichens
(c) Fungi	(d) Mosses and ferns

2 Which one of the following has the highest number of species in nature?

(a) Insects	(b) Birds
(c) Angiosperms	(d) Fungi

- **3** Maximum nutritional diversity is found in the group (a) Fungi (b) Animalia (c) Monera (d) Plantae
- **4** Which one of the following shows maximum genetic diversity in India?
 - (a) Rice (b) Maize (c) Mango (d) Groundnut
- 5 Genetic diversity is related to
 - (a) types of species within a community
 - (b) types of community in an area
 - (c) gene based diversity
 - (d) diversity and evolution of species with a genus
- **6** In India, we find mangoes with different flavours, colours, fibre content, sugar content and even shelf life. The large variation is an account of
 - (a) species diversity (b (c) genetic diversity (d
 - (b) induced mutations (d) hybridisation
- 7 α -diversity is biodiversity present
 - (a) within community (b) between community (c) ranges of community (d) All of these
- 8 Rate of replacement of species along a gradient of habitats/communities is called
 - (a) α -diversity (b) β -diversity (c) γ -diversity (d) ω -diversity
- 9 Which of the following is called β-diversity?
 (a) Habitat diversity
 (b) Differentiation diversity
 (c) Resource diversity
 (d) Species diversity
- 10 Biodiversity is determined by
 - (a) number of individuals in an area
 - (b) species richness
 - (c) evenness
 - (d) Both (b) and (c)
- 11 Species diversity increases as one proceeds from
 - (a) high altitude to low altitude and high latitude to low latitude
 - (b) low altitude to high altitude and high latitude to low latitude
 - (c) low altitude to high altitude and low latitude to high latitude
 - (d) high altitude to low altitude and low latitude to high latitude

- **12** Which of the below mentioned regions exhibit less seasonal variations?
 - (a) Tropics (c) Alpines

(b) Temperates (d) Both (a) and (b)

- **13** Which of the following regions has the greatest biodiversity on earth?
 - (a) Moist deciduous forests(b) Northern boreal forest(c) Amazon rainforest(d) None of these
- 14 Alexander von Humboldt described for the first time → NEET 2017
 - (a) ecological biodiversity(b) law of limiting factor(c) species-area relationships(d) population growth equation
- **15** Susceptibility to extinction is due to
 - (a) large body size(b) small population(c) high trophic level(d) All of these
- 16 The cause of extinction of blue whale is
 - (a) small population size and low reproductive rate
 - (b) crushing of bones under body weight
 - (c) incapability to breath
 - (d) cracking of skin
- **17** Which of the following is the most important cause of animals and plants being driven to extinction?
 - → NEET-I 2016

🕀 www.studentbro.in

- (a) Alien species invasion
- (b) Habitat loss and fragmentation
- (c) Coextinctions
- (d) Overexploitation
- 18 Which of the following is most dangerous to wildlife?
 - (a) Overexploitation
 - (b) Man-made forest
 - (c) Habitat destruction
 - (d) Introduction of foreign species
- 19 Which of the following statements is correct?
 - (a) Steller's sea cow is an extinct animal
 - (b) Lantana is popularly known as carrot grass
 - (c) Parthenium is an endemic species of our country
 - (d) African catfish is not a threat to indigenous catfishes
- 20 The extinction of passenger pigeon was due to
 - (a) increased number of predatory birds
 - (b) over-exploitation by humans
 - (c) non-availability of the food
 - (d) bird flu virus infection
- **21** Which one of the following pairs of organisms is exotic species introduced in India?
 - (a) Ficus religiosa, Lantana camara
 - (b) Lantana camara, Water hyacinth
 - (c) Water hyacinth, Prosopis, Cinereria
 - (d) Nile perch, Ficus religiosa

CLICK HERE

22 Which of the following is not an alien species?

(a) Parthenium	(b) Eichhornia

(c) Clarias gariepinus (d) Podophyllum

23 Which of the following is correctly matched?

→ NEET-II 2016

(a) Aerenchyma	— Opuntia
(b) Age pyramid	— Biome
(c) Parthenium hysteropho	orus — Threat to biodiversity
(d) Stratification	— Population

24 The alien species introduced into Lake Victoria that was responsible for the extinction of cichlid fishes is

(a) Murrels	(b) Nile perch
(c) Carrot grass	(d) African catfis

25 Which of the following is not an invasive alien species in the Indian context?

(a) Lantana (b) Cynodon (c) Parthenium (d) Eichhornia

26 Match the following columns.

	Column I (Animals)				Со	lumn	II (Loo	cation))		
	Α.	Dodo			1.	A	frica				
	В.	Quage	ja		2.	R	lussia				
	C.	Thylac	ine		3.	N	1auriti	us			
	D.	Stellar	's sea	COW	4.	Α	ustra	lia			
	Code	es									
	A	A B	С	D			А	В	С	D	
	(a) 3	3 1	2	4		(b)	3	1	4	2	
	(c) 1	3	2	4		(d)	4	3	1	2	
27	How	many h	otspo	ots of I	biodiv	ers	ity in	the v	vorld	have	
	been	Identifi	ea tii	l date	DY NO	orma	an ivi	yers	→ NI	EET-II	2016
	(a) 1	7	(b)	25		(c)	34		(d) 4	3	2010
20	The e	, noning	(D)	inod to		(O) (ulor	ragio		l not f	ound
20	elsew	here is	term	ineu il ied as	σαμα	u uc	ulai	iegioi ⇒ (BSF-4		2015
	(a) ke	evstone		iou uo		(b) ;	alien				2010
	(c) e	ndemic				(d) i	rare				
29	Whick	n one c	f the	follow	ing ar	eas	s in Ir	ndia, i	s a h	otspo	t of
	biodiv	versity?)		0			→ (BSE-A	NPMT	2012
	(a) E	astern (Ghats			(b) (Gang	jetic p	olain		
	(c) S	underb	ans			(d) \	West	ern G	hats		
30	Selec	t the co	orrect	t state	ment	abc	out bi	odive	ersity.		
	(a) Tl	he dese	ert are	eas of I	Rajast	han	and	Gujar	at ha	ve a v	ery
	h	ligh leve	el ot c	lesert	anıma	l sp	ecies	s as w	ell as	nume	rous
	(b) L	arde sc	ale p	lantino	of Bt	cot	ton h	as no	adve	rse ef	fect
	Ó	n biodi	versit	y Ű							
	(c) W	/estern	Ghate	s have	a ver	y hię	gh de	egree	of spe	ecies	
	ri (d) C	Chness	and	enderr of bioc	nism Iivorsi	tv ic	inet	a fad	nurei	ind hy	, tha
	(u) U d	levelop	ed co	untries	aivei si S	ty is	just	a iau	puisc	ieu by	/ 110
31	The re	eaion c	of hio	sohere	rese	rve	whi	ch is	legall	V	
5.	prote	cted ar	nd wh	iere no	b hum	an	activ	ity is	allow	, ed, is	
	know	n as						<i>,</i>	→	NEET	2017
	(a) c	ore zon	е			(b) l	buffe	r zone	Э		

(c) transition zone (d) restoration zone

- 32 Buffer zone of biosphere reserve is where
 - (a) no human activity is allowed
 - (b) human activity is limited
 - (c) free human activity is allowed
 - (d) wild animals are absent
- 33 The first biosphere reserve established in India for conserving the gene pool of flora and fauna and the life style of tribals is
 - (a) Nilgiri biosphere reserve
 - (b) Nandadevi biosphere reserve
 - (c) Uttarakhand biosphere reserve
 - (d) Great Nicobar biosphere reserve
- 34 Which one of these is an in situ method of conservation? (a) National park (b) Botanical garden
 - (c) Tissue culture (d) Genetic engineering
- 35 Which of the following will help most in conservation of wildlife?
 - (a) Making stringent laws
 - (b) Making numerous zoos
 - (c) Making numerous sanctuaries
 - (d) All of the above
- 36 First National Park of India was

(a) Jim Corbett National Park (b) Kaziranga National Park (c) Panna National Park (d) Gir National Park

37 Simlipal is

CLICK HERE

(a) sanctuary	(b) biosphere reserve
(c) only national park	(d) zoo

- 38 Asiatic lion is protected in (a) Kaziranga National Park (b) Gir National Park (c) Kanha National Park (d) Desert National Park
- **39** Identify the odd combination of the habitat and the
- particular animal concerned. (a) Dachigam National Park — Snow leopard
 - (b) Sunderbans Bengal tiger
 - (c) Periyar Elephant
 - Wild ass (d) Rann of Kutch
- 40 Tiger is not a resident in which one of the following national parks? → CBSE-AIPMT 2009 (a) Ranthambhor (b) Sunderbans (c) Gir (d) Jim Corbett
- 41 Which of the following national parks is home to the famous musk deer or hangul? → NEET-II 2016
 - (a) Keibul Lamjao National Park, Manipur
 - (b) Bandhavgarh Naitonal Park, Madhya Pradesh
 - (c) Eaglenest Wildlife Sancturay, Arunachal Pradesh
 - (d) Dachigam Naitonal Park, Jammu and Kashmir
- 42 Brow antiered deer is one of the rarest mammal found in India. It is found in
 - (a) Nanda Devi Himachal Pradesh
 - (b) Keibul Lamjao National Park Manipur
 - (c) Dudhwa National Park Uttar Pradesh

🕀 www.studentbro.in

(d) Periyar Wildlife Sanctuary Kerala

- 43 Biosphere reserves differ from National Parks and Wildlife Sanctuaries because in the former.
 - (a) human beings are not allowed to enter
 - (b) people are an integral part of the system
 - (c) plants are paid greater attention than the animals
 - (d) living organisms are brought from all over the world and preserved for posterity

44 Bandipur (Karnataka) National Park is the site of

- (a) Deer project (b) Peacock project
- (c) Elephant project (d) Tiger project
- 45 Hoolock Gibbon (India's only ape) is found in
 - (a) Kaziranga Bird Sanctuary (b) Hazaribagh National Park (c) Corbett National Park (d) Gir National Park
- 46 Rajaji National Park is situated in
 - (a) Tamil Nadu (b) Karnataka (c) Uttarakhand (d) Rajasthan
- 47 What is true of National Park?
 - (a) Tourism is allowed in buffer zone
 - (b) No human activity is allowed
 - (c) Cattle grazing is allowed in buffer zone
 - (d) Hunting is allowed in core zone
- 48 Biosphere reserves are different from National Parks as
 - (a) plants and animals are protected in biosphere reserves
 - (b) humans are integral part of biosphere reserves
 - (c) humans are not involved in biosphere reserves
 - (d) None of the above
- 49 Kanha National Park is located in
 - (a) Assom (b) Rajasthan (c) Uttar Pradesh (d) Madhya Pradesh
- 50 Rare animal Hispid hare (Caprologus hispidus) is found in
 - (a) Dachigam National Park (b) Kaziranga National Park
 - (c) Manas National Park (d) Kanha National Park
- 51 National park and sanctuary have the common characteristic as
 - (a) boundries are circumscribed by state legislation
 - (b) there is biotic interference
 - (c) tourism is permissible
 - (d) research and scientific management is possible
- 52 'Hangul Project' was started by government to save Hangul (Cervus hangul) in 1970. The sanctuary, where it is started is (a) National Chambal Sanctuary
 - (b) Dachigam Sanctuary
 - (c) Corbett National Park
 - (d) Bandipur National Park
- 53 Ranganathitu sanctuary (Mysore) is known for population of

(a) bison	(b) tigers	(c) goats	(d) birds
-----------	------------	-----------	-----------

54 Sanctuary which has maximum number of rare animals is (a) Manas National Park (b) Kaziranga National Park (c) Dudhwa National Park (d) Corbett National Park

- 55 The one-horned rhinoceros is specific to which of the following sanctuary?
 - (a) Bhitar Kanika (b) Bandipur
 - (d) Corbett Park (c) Kaziranga
- 56 Which of the following is correctly matched?
 - Sanctuary Animal (a) Gir Lion (b) Kaziranga Musk deer (c) Sunderbans Rhino (d) NE Himalayan region Sambar
- 57 Which among the following is a sacred lake?
 - (a) Dal lake
 - (b) Khecheopalri lake of Sikkim
 - (c) Surajkund lake
 - (d) Chilka lake
- 58 Rare endangered and endemic taxa can be found intact and flourishing in
 - (a) sacred groves (b) buffer zone (c) tropical forests
 - (d) temperate forests
- 59 Significant wetlands of India have been declared as
 - (a) Bastic sites (b) Gaston sites (c) Ramsar sites (d) Spicer sites
- 60 Breeding place of flamingo (hansawar) in India is (a) Chilka Lake (b) Sambhar Lake (c) Rann of Kutch (d) Ghana Vihar
- 61 All of the following are included in ex situ conservation → NEET-II 2016 except (a) botanical gardens (b) sacred groves
 - (c) wildlife safari parks (d) seed banks
- 62 The first white tiger safari in the world is in
 - (a) Nagarjuna sagar Andhra Pradesh
 - (b) Kalkad Mundan thurai Tamil Nadu
 - Kerala (c) Perivar
 - (d) Nandan Kanan Zoo Park Odisha
- 63 In which of the following both pairs has correct combination? → CBSE-AIPMT 2015
 - (a) In situ conservation/National park Ex situ conservation/Botanical garden
 - (b) In situ conservation/Cryopreservation Ex situ conservation/Wildlife sanctuary
 - (c) In situ conservation/Seed bank Ex situ conservation/National park
 - (d) In situ conservation Tissue culture
 - Ex situ conservation/Sacred groves
- 64 Which one of the following is related to ex situ conservation of threatened animals and plants?

→ NEET 2017

- (a) Wildlife safari parks (b) Biodiversity hotspots (c) Amazon rainforest (d) Himalayan region
- 65 Which one of the following is not used for ex situ plant conservation? → NEET 2013
 - (a) Field gene banks (c) Shifting cultivation

(b) Seed banks

(d) Botanical gardens





66 Pollen grains can be stored for several years in liquid nitrogen having temperature of → NEET 2018
 (a) -196°C
 (b) -80°C

(a) -190 C	(U) - 00 C
(c) -120°C	(d) −160°C

67 Cryopreservation of gametes of threatened species in viable and fertile condition can be referred to as

→ CBSE-AIPMT 2015

- (a) *in situ* conservation of biodiversity(b) advanced *ex situ* conservation of biodiversity
- (c) *in situ* conservation by sacred groves
- (d) *in situ* cryoconservation of biodiversity
- 68 A collection of plants and seeds having diverse alleles of all the genes of a crop is called → CBSE-AIPMT 2011
 (a) germplasm
 (b) gene library
 (c) genome
 (d) herbarium
- 69 The organisation which publishes the Red List of species is → CBSE-AIPMT 2014

(a) ICFRE	(b) IUCN
(c) UNED	(d) WWF

70 IUCN, now called World Conservation Union (WCU) has its headquarter at

(a) South Africa	(b) America
(c) India	(d) Switzerland

- 71 A species facing extremely high risk of extinction in the immediate future is called → CBSE-AIPMT 2014
 - (a) vulnerable
 - (b) endemic
 - (c) critically endangered
 - (d) extinct
- 72 Red List contains data or information on

→ NEET-II 2016

- (a) all economically important plants
- (b) plants whose products are in international trade
- (c) threatened species
- (d) marine vertebrates only
- **73** Amongst the animal groups given below, which one has the highest percentage of endangered species?

(a) Insects	(b) Reptiles
(c) Mammals	(d) Amphibians

- **74** Which one of the following is an endangered plant species of India?
 - (a) Santalum album (Sandal wood)
 - (b) Rauwolfia serpentina
 - (c) Cycas beddonei
 - (d) All of the above
- **75** Which one of the following sets consists entirely of endangered wildlife species of India?
 - (a) Egret, Black boar, Bison, Spotted deer
 - (b) Himalayan musk deer, Black buck, Indian lion, Rhino, Sambar
 - (c) Himalayan musk deer, Indian lion, Rhino, Wild buffalo, Golden cat
 - (d) Himalayan musk deer, Great Indian bustard, Snow leopard, Kashmir stag, Wild ass

- **76** Which of the following pairs of an animal and a plant represents endangered organisms in India?
 - (a) Cinchona and Leopard
 - (b) Banyan and Black buck
 - (c) Tamarind and Rhesus monkey
 - (d) Bentinckia nicobarica and Red panda
- 77 Which endangered animal is the source of the world's finest, lightest, warmest and most expensive wool-the shahtoosh?(a) Nilgai(b) Cheetal
 - (c) Kashmiri goat (d) Chiru
- **78** The endangered largest living lemur *Idri idri* is inhabitant of
 - (a) Madagascar (b) Mauritius (c) Sri Lanka (d) India
- **79** The historic convention on Biological Diversity held in Rio de Janeiro in 1992 is known as
 - (a) G-16 Summit (b) CITES Convention (c) The Earth Summit (d) MAB Programme
- 80 Earth summit at Rio de Janeiro was related to

 (a) soil fertility
 (b) conservation of environment
 (c) afforestation
 (d) natural resources
- 81 Which animal is the symbol of the World Wildlife Fund?
 (a) Tiger
 (b) Hornbill
 (c) Giant panda
 (d) White bear
- 82 Match the following columns.

→ NEET-II 2016

🕀 www.studentbro.in

	Column I					Colun	nn ll	
A.	Bio	osphe	ere res	serves	1.	Wild	animal	S
В.	MA	٨B			2.	1992		
C.	Sa	nctua	aries		3.	1994		
D.	Pro	oject	eleph	ant	4.	1986		
					5.	1971		
Cod	es							
	А	В	С	D		А	В	С
(a)	4	5	1	2	(b)	1	2	3
(C)	5	4	2	1	(d)	1	4	5

- 83 In 1984, the Bhopal gas tragedy occurred because gas Methyl Isocyanate reacted (MIC) chemically with
 (a) DDT
 (b) ammonia
 (c) CO₂
 (d) water
- 84 Photochemical smog is formed due to emission of primary air pollutants
 - (a) SO_2 and CO (b) O_3 and PAN (c) NO_2 and hydrocarbons (d) CO_2
- 85 Nitrogen oxides produced from the emission of automobiles and power plants are the source of fine air borne particles which lead to
 - (a) photochemical smog (b) dry acid deposition (c) wet acid deposition (d) Both (b) and (c)
- **86** Suspended particles such as ash, soot smoke may affect human body as they cause

(a) eye irritation	(b) emphysema
(c) mutations	(d) Both (a) and (b)

CLICK HERE

- 87 Which one of the following statements is not valid for aerosols? → NEET 2017
 - (a) They are harmful to human health
 - (b) They alter rainfall and monsoon patterns
 - (c) They cause increased agricultural productivity
 - (d) They have negative impact on agricultural land
- 88 In almost all Indian metropolitan cities like Delhi, the major atmospheric pollutant(s) is/are
 - (a) Suspended Particulate Matter (SPM)
 - (b) oxides of sulphur
 - (c) carbon dioxide and carbon monoxide
 - (d) oxides of nitrogen
- 89 Burning sensation of throat, eyes and nausea is associated with present in air.
 - (a) hydrogen sulphide (b) sulphur (c) hydrogen cyanide (d) arsenic
- 90 Carbon dioxide, methane, nitrogen oxide and chlorofluorocarbons are called greenhouse gases because they can absorb
 - (a) ultraviolet radiations (c) γ -rays radiations
- (b) visible light radiations (d) long wave infrared radiations
- 91 If there was no carbon dioxide on the earth's atmosphere, the temperature of the earth's surface would be
 - (a) same as the present level
 - (b) more than the present level
 - (c) less than the present level
 - (d) dependent on the oxygen content in the atmosphere
- **92** At present, the concentration of CO₂ in the atmosphere is about

(a) 100 ppm	(b) 240 ppm
(c) 380 ppm	(d) 520 pmm

- 93 Greenhouse effect with respect to global climate refers to (a) cooling and moist condition
 - (b) warming effect
 - (c) increase rainfall and greenery
 - (d) desertification
- 94 Which one of the following is the correct percentage of the two (out of the total of four) greenhouse gases that contribute to the total global warming? → CBSE-AIPMT 2008 (a) CFCs 14% and CH $_{\rm 4}$ 20% (b) CO $_2$ 40% and CFCs 30% (c) N_2O 6% and CO_2 86% (d) CH_4 20% and N_2O 18%
- 95 Carbon dioxide is called greenhouse gas because it is
 - (a) used in greenhouse to increase plant growth
 - (b) transparent to heat but traps sunlight
 - (c) transparent to sunlight but traps heat
 - (d) transparent to both sunlight and heat
- 96 According to Kyoto protocol, the major nations abide to reduce concentration of greenhouse gases by

(a) 2008	(b) 2010
(c) 2012	(d) 2018

consequence of rising CO₂ levels? (a) C_4 crops, such as corn, being replaced by more C_3 plants such as wheat and soybean (b) Rising global temperature (c) Increased breakdown of atmosphere ozone (d) Increased vegetative productivity **98** Global warming can be controlled by → NEET 2013 (a) reducing deforestation, cutting down use of fossil fuel

97 Which of the following is not thought to be a possible

- (b) reducing reforestation, increasing the use of fossil fuel.
- (c) increasing deforestation, slowing down the growth of human population
- (d) increasing deforestation, reducing efficiency of energy usade
- 99 The UN conference of Parties on climate change in the year 2012 was held at → NEET 2015 (a) Durban
 - (b) Doha (d) Warsaw
- 100 World Ozone day is celebrated on → NEET 2018 (a) 16th September (b) 21th April (c) 5th June (d) 22nd April
- **101** Depletion of which gas in the atmosphere can lead to an increased incidence of skin cancers? → NEET 2016 (b) Ammonia (a) Ozone
 - (c) Methane (d) Nitrous oxide
- 102 The destruction of the ozone layer is most directly linked to the
 - (a) destruction of tropical rainforests
 - (b) release of Chlorofluorocarbons (CFCs) into the atmosphere
 - (c) decrease in the production of ozone precursors by chemical production plants
 - (d) global warming

(c) Lima

- **103** In stratosphere, which one of the following elements acts as catalyst in degradation of ozone and release of molecular oxygen? → NEET 2018
 - (b) Cl (c) Carbon (d) Oxygen
- 104 Ozone layer is found in two atmospheric layers of atmosphere, identify the option which associates with good ozone.
 - (a) Troposphere

(a) Fe

- (b) Mesosphere
- (c) Stratosphere
- (d) Thermosphere
- 105 Ozone layer thickness is measured in
 - (a) decibels (b) millimetrer
 - (c) Dobson units (d) centimetre
- 106 In 1975, atmospheric scientists first discovered the formation of ozone hole, maximum over
 - (a) Arctic (b) Antarctica (d) Pakistan (c) India





- 107 Global agreement in specific control strategies to reduce the release of ozone depleting substances, was adopted by → CBSE-AIPMT 2009
 - (a) Rio de Janeiro Conference
 - (b) Montreal Protocol
 - (c) Kyoto Protocol
 - (d) Vienna Convention
- 108 Which of the following is not correct?
 - (a) Greenhouse effect is related to global warming
 - (b) First earth summit held at Rio de Janeiro (Brazil) in 1992
 - (c) Kyoto protocol held in Japan
 - (d) Montreal protocol is related to greenhouse gases
- **109** Which of the following is not one of prime health risks associated with greater UV radiation through the atmosphere due to the depletion of stratospheric zone?

→ CBSE-AIPMT 2015

- (a) Increased skin cancer (c) Damage to eyes
- cer (b) Reduced immune system (d) Increased liver cancer
- **110** The molecular action of ultraviolet light is mainly reflected through
 - (a) destruction of hydrogen bonds between DNA strands
 - (b) photodynamic action
 - (c) formation of pyrimidine
 - (d) formation of sticky metaphase
- **111** A substantial fall in CO₂ and SO₂ level has been found in Delhi between 1997 and 2005 due to
 - (a) use of purified, unleaded petrol
 - (b) use of purified, unleaded diesel
 - (c) use of Compressed Natural Gas (CNG) in public transports
 - (d) use of Liquified Petroleum Gas (LPG) in public transports
- A chemical industrial plant is releasing large amount of exhaust in the atmosphere. As a control method, scrubber in the exhaust was advised because it removes
 → CBSE-AIPMT 2015
 - (a) gases like sulphur dioxide
 - (b) particulate matter of the size 5 micrometre or above
 - (c) gases like ozone and methane
 - (d) particulate matter of the size 2.5 micrometre or less
- 113 The Air Prevention and Control of Pollution Act came into force in → NEET 2013
 (a) 1975 (b) 1981 (c) 1985 (d) 1990
- 114 Catalytic converters are fitted into automobiles to reduce emission of harmful gases. Catalytic converters change
- emission of harmful gases. Catalytic converters change unburnt hydrocarbons into

(a) carbon dioxide and water	(b) carbon monooxide
(c) methane	(d) carbon dioxide and methane

115 Match the following columns.

	Column I		Column II
Α.	Electrostatic precipitator	1.	Removes gases like SO_2
В.	Scrubber	2.	Reduces automobile emission
C.	Catalytic converter	3.	Removes particulate matter

Codes

	А	В	С		А	В	С
(a)	3	2	1	(b)	1	2	3
(c)	3	1	2	(d)	1	3	2

- **116** Steps taken by the Government of India to control air pollution include → CBSE-AIPMT 2009
 - (a) compulsory mixing of 20% ethyl alcohol with petrol and 20% biodiesel with diesel
 - (b) compulsory Pollution Under Control (PUC) certification of petrol driven vehicles, which tests for carbon monoxide and hydrocarbons
 - (c) permission to use only pure diesel with a maximum of 500 ppm sulphur as fuel for vehicles
 - (d) use of non-polluting Compressed Natural Gas (CNG) only as fuel by all buses and trucks
- **117** Which one of the following statements pertaining to water pollutants is correct?
 - (a) DDT is a non-biodegradable pollutant
 - (b) Excess fluoride in drinking water causes osteoporosis
 - (c) Excess cadmium in drinking water causes black foot disease
 - (d) Methylmercury in water may cause 'itai-itai' disease
- **118** Addition of phosphate and nitrates rich fertiliser into water bodies which ultimately lead into lake, first affects the
 - (a) growth of aquatic organisms in lake
 - (b) eutrophication rate of lake
 - (c) environment of lake
 - (d) organic remains deposited on the bottom of lake
- **119** The presence of large amounts of nutrients in water causes excessive growth of planktonic algae, called (a) algal bloom
 - (b) eutrophication
 - (c) biomagnification
 - (d) acidification
- **120** A lake which is rich in organic waste may results in

→ NEET-II 2016

- (a) increased population of aquatic organisms due to minerals
- (b) drying of the lake due to algal bloom
- (c) increased population of fish due to lots of nutrients
- (d) mortality of fish due to lack of oxygen
- 121 The green scum seen in the freshwater bodies is
 - (a) blue-green algae
 - (b) red algae
 - (c) green algae

CLICK HERE

- (d) Both (a) and (c)
- **122** Eutrophication of water bodies leading to killing of fishes is mainly due to non-availability of

(г	ι) food		(b)	light
(c) esse	ential minerals	(d)	oxvaen

- **123** The highest DDT concentration in aquatic food chain shall occur in
 - (a) phytoplankton (b) seagull (c) crab (d) eel



- **124** If we continuously add sewage water in a river, the BOD of river will
 - (a) decrease continuously
 - (b) increase continuously
 - (c) be higher before addition of sewage water
 - (d) be lower after addition of sewage water
- 125 BOD is
 - (a) amount of oxygen utilised by microorganisms
 - (b) amount of oxygen utilised by organisms in water
 - (c) total amount of oxygen present in water
 - (d) All of the above
- **126** A river with an inflow of domestic sewage rich in organic → NEET-I 2016 waste may result in
 - (a) an increased population of aquatic food web organisms
 - (b) an increased production of fish due to biodegradable nutrients
 - (c) death of fish due to lack of oxygen
 - (d) drying of the river very soon due to algal bloom
- **127** A water body is polluted, this can be confirmed by the presence of
 - (a) Lemna pancicostata
 - (b) Eichhornia crassipes
 - (c) Escherichia coli
 - (d) Entamoeba histolytica
- **128** Sewage water can be purified for recycling with the action of

(a) microorganisms	(b) penicillin
(c) fishes	(d) aquatic plants

- **129** A sewage treatment process in which a portion of the decomposer bacteria present in the waste is recycled into the beginning of the process, is called
 - (a) cvclic treatment
 - (b) primary treatment
 - (c) activated sludge treatment
 - (d) tertiary treatment
- 130 DDT has been a major pollutant because it
 - (a) kills aquatic animals
 - (b) kills pests
 - (c) destroys many valuable species
 - (d) is non-biodegradable
- 131 Some possible properties of modern insecticides are listed below. When these insecticides are used, which property helps to keep environmental pollution at the lowest level?
 - (a) Accumulates in the bodies of predators
 - (b) Broken down by soil bacteria
 - (c) Easily washed into lakes and rivers
 - (d) Taken up by the plant roots
- **132** Formation of non-functional methaemoglobin causes blue-baby syndrome. This is due to
 - (a) excess of arsenic concentration in drinking water
 - (b) excess of nitrates in drinking water
 - (c) deficiency of iron in food
 - (d) increased methane content in the atmosphere

- 133 Itai-itai disease which results in kidney damage is caused by the poisoning of
 - (a) mercury (b) lead (c) chromium (d) cadmium
- 134 In a polluted environment, the maximum pollutant will occur in
 - (a) primary producers (b) tertiary producers (c) secondary producers (d) primary consumers
- 135 Bad taste and odour of the water bodies of tanks, ponds, etc. is due to
 - (a) acids and grease (b) free chlorine (c) fats (d) starch
- **136** Aluminium in drinking water may influence metabolism. (a) nitrate (b) phosphate
 - (d) fluorine
 - (c) calcium
- **137** Eutrophic lakes are highly (a) productive (b) enriched with phosphates (c) Both (a) and (b) (d) None of these
- 138 The index of pollution in a polluted lake is (b) BOD (a) Daphnia (c) Stoneflies
 - (d) All of these
- **139** During sewage treatment, biogases are produced which include → NEET 2013
 - (a) methane, oxygen, hydrogen sulphide
 - (b) hydrogen sulphide, methane, sulphur dioxide
 - (c) hydrogen sulphide, nitrogen, methane
 - (d) methane, hydrogen sulphide, carbon dioxide
- 140 Lead, a toxic chemical is mainly considered as (a) air pollutant (b) water pollutant (c) soil pollutant (d) noise pollutant
- 141 Which of the following is/are responsible for increased soil pollution?
 - (a) Excreta of human and other living beings
 - (b) Radioactive wastes as C¹⁴
 - (c) Fertilisers, pesticides, etc
 - (d) All of the above
- 142 Soil pollution more often results in
 - (a) changing the chemistry of soil
 - (b) decreased soil fertility
 - (c) increased crop yield
 - (d) Both (a) and (b)
- **143** The chemicals released due to unsustainable agricultural practices and causing soil pollution are
 - (a) pathogens (b) chemical fertilisers
 - (c) strontium-90 (d) None of these
- 144 Organic farming is an excellent substitute for increased chemical use in agriculture. Select an option which justifies the above statement.
 - (a) Waste of one process is cycled as nutrients in other processes

🕀 www.studentbro.in

- (b) It is a zero waste procedure
- (c) Both (a) and (b)

CLICK HERE

(d) It is carried out at high temperatures

145 Percentage of forest area recommended by the national forest policy in plains is

(a) 33%	(b) 67%
(c) 30%	(d) 10%

- **146** Chipko movement, a first of its kind public campaign first occurred in
 - (a) Sundarban area in Bengal (b) Tehri Garhwal of UK
 - (c) Rajasthan (d) None of these
- **147** The practice of restoring a forest cover over an area where one existed previously but was removed at some time, in the past is
 - (a) afforestation
 - (b) reforestation
 - (c) desertification
 - (d) deforestation
- **148** Polyblend is a fine powder of recycled modified plastic which
 - (a) enhance the bitumen's water repellant properties
 - (b) helps to increase the life of road
 - (c) Both (a) and (b)
 - (d) is a type of magnet and helps to improve blood circulation when applied on human body
- **149** Steps taken by the Government of india to control air pollution includes
 - (a) complulsory mixing of 20% ethyl alcohol with petrol and 20% biodiesel with diesel
 - (b) compulsory PUC (Pollution Under Control) certification of petrol driven vehicles, which tests for carbon monoxide and hydrocarbons
 - (c) permission to use only pure diesel with a miximum of 500 ppm sulphur as fuel for vehicles
 - (d) use of non-polluting compressed Natural Gas (CNG) only as fuel by all buses and trucks

150 Match the following columns.

		C	olumi	٦l			Column	ı II
Α.	Bis	hnoi (Comm	nunity		1.	1988	
В.	. Chipko movement					2.	1980	
C.	C. Joint Forest Management					3.	1974	
D.	The National Forest Policy				icy	4.	1731	
Coc	les							
	А	В	С	D				
(a)	1	4	3	2				
(b)	4	3	2	1				
(C)	3	2	1	4				
(d)	4	1	2	3				

Directions (Q. Nos. 151-158) In each of the following questions a statement of Assertion is given followed by a corresponding statement of Reason. Of the statements, mark the correct answer as

- (a) If both Assertion and Reason are true and Reason is the correct explanation of the Assertion
- (b) If both Assertion and Reason are true but Reason is not the correct explanation of Assertion
- (c) If Assertion is true but Reason is false
- (d) If both Assertion and Reason are false
- **151** Assertion Garden is an example of *ex situ* conservation. **Reason** Garden is an artificial habitat resembling to the natural habitat of organisms.
- **152** Assertion A Suspended Particulate Matter (SPM) is an important pollutant released by diesel vehicles.Reason Catalytic converters greatly reduce pollution caused by automobiles.
- **153** Assertion Agricultural output increased several times after introduction of DDT.**Reason** DDT was the first insecticide used on a wide scale.
- **154** Assertion A concentration of methane in the atmosphere has more than doubled in the last 250 years.**Reason** Wetlands and rice fields are the major sources of methane.
- **155** Assertion Methane component of greenhouse gases contribution to global warming is about 20%. **Reason** Introduction of multipoint fuel injection engines in automobiles has decreased methane content in the exhausts.
- **156** Assertion Presently, the global atmosphere is warming up. **Reason** The depletion of stratospheric ozone layer has resulted in increase in ultraviolet radiations reaching the earth.
- **157** Assertion UV radiation causes photodissociation of ozone into O₂ and O, thus causing damage to the stratospheric ozone layer. **Reason** Ozone hole is resulting in global warming and climate change.
- **158** Assertion Deforestation is one of the main factors contributing to global warming.**Reason** Besides CO₂, two other gases methane and CFCs are also included under greenhouse gases.

🕀 www.studentbro.in



DAY PRACTICE SESSION 2

PROGRESSIVE QUESTIONS EXERCISE

- 1 Name the medicinal plant growing in different Himalayan ranges. (a) Plantago ovata (b) Atropa belladona (c) Rauwolfia vomitoria (d) Cinchona officinalis 2 Hotspots of biodiversity mean (a) species in particular niche/area (b) species diversity at particular area (c) areas of the earth that contain many endemic species (d) species serves as proxy for entire communities in particular area 3 Oran is a (a) sacred grove (b) sacred landscape (c) sacred animal (d) endangered animal 4 Which of the following animals has become extinct from India? (a) Snow leopard (b) Hippopotamus (c) Wolf (d) Cheetah 5 Maximum species diversity is seen in latitudinal range of (b) 23.5° N to 23.5°S (a) 23.5° N to 66.5° N (d) 66.5° N to 90° N (c) 23.5° S to 66.5° N 6 Which one of the following is not included under in situ conservation? (a) Sanctuary (b) Botanical garden (c) Biosphere reserve (d) National park 7 Plant species on verge of extinction due to overexploitation is (a) Centella (b) Podophyllum (c) Gloriosa (d) All of these 8 The venue and year of the Earth Summit Conservation of **Biodiversity** was (a) Ramsar, 1974 (b) Stockholm, 1974 (c) South Africa, 2002 (d) Rio de Janeiro, 1992 9 'Evil Quartet' is related with (a) loss of biodiversity (b) loss of standing crop (c) loss of alien species (d) loss of climax community 10 Which of the following is ex situ conservation? (a) Protecting fishing in Bhitar kanika (b) Banning of Akhard Sikar in Simlipal (c) Breeding of animals in Nandankanan (d) Protecting migration of birds in Chilka lake
- 11 Which of the following has the greatest biodiversity on earth?
 - (a) Tropical Amazonian rainforests in South America
 - (b) Eastern Ghats and Western Himalayas
 - (c) Sahara deserts
 - (d) Savanna forests

- 12 Which one of the following is a pair of endangered species?
 - (a) Garden lizard and Mexican poppy
 - (b) Rhesus monkey and Sal tree
 - (c) Indian peacock and Carrot grass
 - (d) Hornbill and Indian aconite
- 13 An institution, where valuable plant material likely to become irretrievably lost in the wild or in cultivation is preserved in a viable condition is known as
 - (a) genome (b) gene library (d) herbarium (c) gene bank
- 14 Which one of the following is not a wildlife conservation project?
 - (a) Project Dodo (b) Project Tiger (d) Project Indian Bustard (c) Project Hangul
- 15 What are large undisturbed areas where wildlife is protected in its natural habitat?
 - (a) Biosphere reserves
 - (b) National parks
 - (c) Sacred landscapes
 - (d) Wildlife sanctuaries
- 16 'Project Tiger' was launched following the recommendations of

(a) IBWL	(b) BNHS
(c) CITES	(d) NWAP

- 17 Endangered or threatened animals are protected from extinction by ex situ conservation in
 - (a) national parks (b) zoological parks (c) wildlife sanctuary
 - (d) biosphere reserves
- **18** Which one of the following is not observed in biodiversity hotspot?
 - (a) Endemism

CLICK HERE

- (b) Species richness
- (c) Accelerated species loss
- (d) Lesser interspecific competition
- 19 Most recently notified biosphere reserve in India is (a) Cold desert (b) Seshachalam
 - (c) Dehang Debang (d) Agasthyamalai
- 20 Which of the following statements is not true? (a) Robert May places the global species diversity at about
 - 7 million
 - (b) More than 70 per cent of all the species recorded are animals
 - (c) Plants comprise not more than 22 per cent of the total recorded global species

🕀 www.studentbro.in

(d) Eastern Ghats have a greater amphibian species diversity than the Western Ghats

- 21 Select the incorrect statement.
 - (a) Stellar's sea cow and passenger pigeon got extinct due to overexploitation by man
 - (b) Lantana and Eichhornia are invasive weed species in India
 - (c) Species diversity increases as we move away from the equator towards the poles
 - (d) The historic convention on biological diversity was held in 1992
- 22 What is common to Lantana. Eichhornia and African catfish?
 - (a) All are endangered species of India
 - (b) All the species are neither threatened nor indigenous species of India
 - (c) All are keystone species
 - (d) All are mammals found in India
- 23 India now has
 - (a) 10 Biosphere Reserves, 50 National Parks and 400 Wildlife Sanctuaries
 - (b) 14 Biosphere Reserves, 50 National Parks and 400 Wildlife Sanctuaries
 - (c) 10 Biosphere Reserves, 90 National Parks and 448 Wildlife Sanctuaries
 - (d) 17 Biosphere Reserves, 96 National Parks and 441 Wildlife Sanctuaries
- 24 Match the following columns.

		Co (Ca	lumn tegory	 √)		Colu (Exar	mn ll nple)		
А.	Ex	tinct			1.	Polar	bear		
В.	Er	ndang	ered		2.	Tiger	shark	<	
C.	Vu	Inera	ble		3.	Mami	noth		
D.	Ne	ear thi	reater	ned	4.	Giant	panc	la	
Co	des								
	А	В	С	D		А	В	С	D
(a)	3	1	4	2	(b) 1	2	3	4
(c)	3	4	1	2	(d) 3	2	1	4

25 From the point of view of rhinoceros reserve which one of the following is correct?

(a) Corbett		Punjab
(b) Palamou	_	Odisha
(c) Kaziranga	_	Assom
(d) Nandan Kanan	—	Rajasthan

- 26 Which one of the following is the correctly matched pair of an endangered animal and a National Park?
 - (a) Lion - Corbett National Park — Kaziranga National Park (b) Rhinoceros
 - (c) Wild ass — Dudhwa National Park
 - (d) Great Indian bustard Keoladeo National Park

27 Choose the incorrect matched pair.

(a) Carrot grass

(b) Wildlife safari parks

- Ex situ conservation

— Lantana

- (c) Amazon rainforest
- (d) Khasi and Jaintia hills
- Lungs of the planet

CLICK HERE

- Meghalaya

28 Match the following columns.

$\begin{tabular}{ c c c c } \hline Column I & C \\ (Reserve area) & (Reserve ar$	olumn II (State) Asom Chhattisgarh Rajasthan Uttar Pradesh			
A.Keoladeo Bird Sanctuary1.AB.Dudhwa National Park2.CC.Kanha National Park3.FD.Kaziranga National Park4.CCodesABCDABC(a) 1243(b) 342(c) 2314(d) 41329Which of the following is a secondary poll(a) SO2(b) CO2(c) CO(d) O3	Asom Chhattisgarh Rajasthan Uttar Pradesh			
B. Dudhwa National Park 2. 0 C. Kanha National Park 3. F D. Kaziranga National Park 4. U Codes A B C D A B C (a) 1 2 4 3 (b) 3 4 2 (c) 2 3 1 4 (d) 4 1 3 29 Which of the following is a secondary poll (a) SO2 (b) CO2 (c) CO (d) O3 4	Chhattisgarh Rajasthan Uttar Pradesh			
C. Kanha National Park 3. F D. Kaziranga National Park 4. U Codes A B C D A B C (a) 1 2 4 3 (b) 3 4 2 (c) 2 3 1 4 (d) 4 1 3 29 Which of the following is a secondary poll (a) SO2 (b) CO2 (c) CO (d) O3 4	Rajasthan Uttar Pradesh			
D. Kaziranga National Park 4. U Codes A B C D A B C (a) 1 2 4 3 (b) 3 4 2 (c) 2 3 1 4 (d) 4 1 3 29 Which of the following is a secondary poll (a) SO2 (b) CO2 (c) (c) CO (d) 0 (d) (d) (d) (d) (d) (d)	Jttar Pradesh			
Codes A B C D A B C (a) 1 2 4 3 (b) 3 4 2 (c) 2 3 1 4 (d) 4 1 3 29 Which of the following is a secondary poll (a) SO ₂ (b) CO ₂ (c) CO (c) CO ₂ (c) CO ₂ (c) CO ₂	D			
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	D			
(a) 1 2 4 3 (b) 3 4 2 (c) 2 3 1 4 (d) 4 1 3 29 Which of the following is a secondary poll (a) SO_2 (b) CO_2 (c) CO (d) O_3				
29 Which of the following is a secondary poll (a) SO_2 (b) CO_2 (c) CO (d) O_3	1			
(a) SO_2 (b) CO_2 (c) CO (d) O_3	utant?			
30 Anxiety and stress reactions are common exposed to increased levels of	in people			
(a) air pollution (b) noise pollu (c) water pollution (d) nuclear po	ution ollution			
 According to the Central Pollution Control particles that are responsible for causing human health are of diameter (a) 2.50 micrometres (b) 5.00 micro 	billution Control Board, le for causing great harm to ter			
(c) 10.00 micrometres (d) 7.5 micror	netres			
 32 Carbon monoxide combines with haemog increased rate to form (a) carboxyhaemoglobin (b) leghaemoglobin (c) haemocyanin (d) methaemoglobin 	s with naemoglobin at an(b) leghaemoglobin(d) methaemoglobin			
33 Ecosanitation is a sustainable system for l reduce its level in water bodies.	handling t			
(a) soil(b) human exit(c) water(d) radioactive	creta e materials			
34 An effective method used for disposing of and hazardous hospital wastes is	ff the infections			
(a) recycling(b) open dum(c) burning(d) None of th	iping iese			
 World's most problematic aquatic weed re killing aquatic biodiversity by suffocating (a) Azolla (b) Wolffia (c) Eichhornia 	esponsible for them. a (d) <i>Trapa</i>			
36 Joint Forest Management (JFM) concept in India during	was introduced			
(a) 1970s (b) 1980s (c) 1990s (d) 1960s				
 Acid rain is caused by increase in the atm concentration of (a) SO₂ and NO₂ (b) SO₃ and CO (c) CO and CO (d) CO and CO 	nospheric CO			
38 The two gases making highest relative co	ncentration to			
the greenhouse gases are(a) CO_2 and CH_4 (b) CH_4 and N(c) CFCs and N2O(d) CO_2 and N	10 ₂ 1 ₂ 0			

🕀 www.studentbro.in

39 In a coal fired power plant, electrostatic precipitators are installed to control emission of

(a)	SO ₂	(b) NO _x
(c)	SPM	(d) CO

- **40** Indicator plants, which can be used to indicate atmospheric pollution by SO₂ are
 - (a) grassland like Deschampsia
 - (b) moss like Sphagnum
 - (c) lichens like *Usnea*
 - (d) climbers like Cucurbita
- **41** Accodring to Euro II norms (1993), content of sulphur in diesel should be

(a) 350 ppm	(b) 150 ppm
(c) 25 ppm	(d) 5ppm

- 42 The presence of water blooms in a lake indicates
 - (a) excessive nutrient
 - (b) nutrient deficiency
 - (c) oxygen deficiency
 - (d) the absence of herbivores
- **43** Limit of BOD prescribed by Central Pollution Control Board for the discharge of industrial and municipal waste water into natural surface water, is

(a) < 3.0 ppm	(b) < 10 ppm
(c) < 100 ppm	(d) < 30 ppm

44 Biochemical Oxygen Demand (BOD) may not be a good index for pollution in water bodies receiving effluents from

(a) domestic sewage	(b
(c) petroleum industry	(d

- (b) dairy industry (d) sugar industry
- **45** 'Chipko Movement' launched in Uttarakhand hills is connected with
 - (a) project tiger
 - (b) plant breeding
 - (c) plant/forest conservation
 - (d) conservation of natural resources
- **46** DDT residues are rapidly passed through food chain causing biomagnification because DDT is
 - (a) liposoluble
 - (b) moderately toxic
 - (c) non-toxic to aquatic animals
 - (d) water soluble
- **47** UV rays are non-ionising type and are lethal due to inactivation of

(a) proteins	(b) pigments
(c) nucleic acid	(d) All of these

- 48 Which of the following is correct?
 - (a) PAN (Peroxy Acetyl Nitrate) is a secondary pollutant
 - (b) Photochemical smog was first observed in Los Angeles (c) Carbon monoxide has about 200 times greater affinity
 - for haemoglobin as compared to oxygen
 - (d) All of the above

- **49** Recent reports of acid rains in big industrial cities are due to the effect of atmospheric pollution by
 - (a) more release of NO_2 and SO_2 by burning of fossil fuels
 - (b) more release of CO₂ by burning of coal/wood cutting of forests and increasing populations
 - (c) excessive release of NH₃ by coal gas/industries
 - (d) excessive release of CO by incomplete combustion of carbonaceous fuels
- **50** Pollution from animal excreta and organic waste from kitchen can be most profitably minimised by
 - (a) stroring them in underground storage tanks
 - (b) using them for producing biogas
 - (c) vermiculture
 - (d) using them directly as biofertilisers
- 51 Greenhouse effect is due to the presence of
 - (a) ozone layer in the atmosphere
 - (b) moisture layer in the atmosphere
 - (c) CO₂ layer in the atmosphere
 - (d) infrared light reaching the earth
- 52 Minamata disease was caused due to the consumption of
 - (a) sea food containing lot of cadmium
 - (b) fish contaminated with mercury
 - (c) oysters with lot of pesticide
 - (d) sea food contaminated with sellenium
- **53** Global warming can be controlled by
 - (a) reducing reforestation, increasing the use of fossil fuel
 - (b) increasing deforestation, slowing down the growth of human population
 - (c) increasing deforestation, reducing efficiency of energy usage
 - (d) reducing deforestation, cutting down use of fossil fuel
- 54 Choose the incorrect statement.
 - (a) The Montreal Protocol is associated with the control of emission of ozone depleting substance
 - (b) Methane and carbon dioxide are greenhouse gases
 - (c) Dobson units are used to measure oxygen content
 - (d) Use of incinerators is crucial to disposal of hospital wastes
- **55** 'Kyoto Protocol' is multinational international treaty for (a) phasing out greenhouse gases
 - (b) controlling ozone destroying substances
 - (c) management of hazardous wastes
 - (d) conservation of biodiversity
- 56 Match the following columns.

CLICK HERE

				0								
		C	olum	Column II								
Α.	DD	Г			1.	Snow blindness						
В.	Biod	degra	dable	e orga	2.	Algal bloom						
C.	UV					З.	BOD					
D.	Pho	phate	es an	d nitra	4.	Biomagnification						
Codes												
	А	В	С	D			А	В	С	D		
(a)	1	2	3	4		(b)	2	1	4	3		
(c)	4	3	1	2		(d)	1	4	3	2		

🕀 www.studentbro.in

Directions (Q. Nos. 57-59) In each of the following questions a statement of Assertion is given followed by the corresponding statement of Reason. Of the statements, mark the correct answer as

- (a) If both Assertion and Reason are true and Reason is the correct explanation of Assertion
- (b) If both Assertion and Reason are true but Reason is not the correct explanation of Assertion
- (c) If Assertion is true but Reason is false
- (d) If both Assertion and Reason are false

57 Assertion In tropical rainforests, O-horizon and A-horizon of soil profile are shallow and nutrient poor.
 Reason Excessive growth of microorganisms in the soil depletes its organic content.

58 Assertion Increase in ozone concentration near the earth's surface affects crop yield.

Reason The crop yield reduces because UV rays reach the earth's surface unfiltered.

59 Assertion Kyoto protocol an international conference was initiated to discuss global warming.
 Reason According to this protocol, the major nations abide to reduce concentration of greenhouse gases by 2012.

ANSWERS

(SESSION 1)	1	(c)	2	(a)	3	(c)	4	(a)	5	(c)	6	(c)	7	(a)	8	(b)	9	(b)	10	(d)
	11	(a)	12	(a)	13	(c)	14	(c)	15	(d)	16	(a)	17	(b)	18	(c)	19	(a)	20	(b)
	21	(b)	22	(d)	23	(c)	24	(b)	25	(b)	26	(b)	27	(c)	28	(c)	29	(d)	30	(c)
	31	(a)	32	(b)	33	(a)	34	(a)	35	(d)	36	(a)	37	(b)	38	(b)	39	(a)	40	(c)
	41	(d)	42	(b)	43	(b)	44	(c)	45	(b)	46	(c)	47	(b)	48	(b)	49	(d)	50	(c)
	51	(c)	52	(b)	53	(d)	54	(a)	55	(c)	56	(a)	57	(b)	58	(a)	59	(c)	60	(a)
	61	(b)	62	(d)	63	(a)	64	(a)	65	(c)	66	(a)	67	(b)	68	(a)	69	(b)	70	(d)
	71	(c)	72	(c)	73	(d)	74	(c)	75	(d)	76	(d)	77	(d)	78	(c)	79	(c)	80	(d)
	81	(c)	82	(d)	83	(d)	84	(b)	85	(a)	86	(d)	87	(c)	88	(a)	89	(a)	90	(d)
	91	(c)	92	(c)	93	(b)	94	(a)	95	(c)	96	(c)	97	(c)	98	(a)	99	(b)	100	(a)
	101	(a)	102	(b)	103	(b)	104	(c)	105	(c)	106	(b)	107	(b)	108	(d)	109	(b)	110	(a)
	111	(c)	112	(b)	113	(a)	114	(a)	115	(c)	116	(d)	117	(a)	118	(a)	119	(a)	120	(d)
	121	(d)	122	(d)	123	(d)	124	(b)	125	(a)	126	(c)	127	(b)	128	(a)	129	(c)	130	(d)
	131	(b)	132	(b)	133	(d)	134	(c)	135	(a)	136	(d)	137	(c)	138	(a)	139	(d)	140	(c)
	141	(d)	142	(d)	143	(b)	144	(c)	145	(a)	146	(b)	147	(b)	148	(c)	149	(d)	150	(b)
	151	(a)	152	(b)	153	(a)	154	(a)	155	(b)	156	(b)	157	(d)	158	(b)				
(SESSION 2)	1	(c)	2	(c)	3	(a)	4	(d)	5	(b)	6	(b)	7	(a)	8	(d)	9	(a)	10	(c)
	11	(a)	12	(d)	13	(c)	14	(a)	15	(a)	16	(a)	17	(b)	18	(d)	19	(b)	20	(d)
	21	(c)	22	(b)	23	(d)	24	(c)	25	(c)	26	(b)	27	(a)	28	(b)	29	(d)	30	(b)
	31	(a)	32	(d)	33	(b)	34	(c)	35	(c)	36	(b)	37	(b)	38	(a)	39	(c)	40	(c)
	41	(a)	42	(a)	43	(d)	44	(C)	45	(C)	46	(a)	47	(a)	48	(a)	49	(a)	50	(D)
	ЭТ	(C)	52	(a)	53	(d)	54	(C)	55	(a)	50	(C)	57	(a)	58	(a)	59	(a)		

