

CO-ORDINATE GEOMETRY

Single Correct Answer Type

1. In ΔABC , $a^2(\cos^2 B - \cos^2 C) + b^2(\cos^2 C - \cos^2 A) + c^2(\cos^2 A - \cos^2 B)$ is equal to
 a) 0 b) 1 c) $a^2 + b^2 + c^2$ d) $2(a^2 + b^2 + c^2)$
2. If $\sin A : \sin B : \sin C = 3 : 4 : 5$, then $\cos A : \cos B$ is equal to
 a) 4:3 b) 5:3 c) 3:4 d) 3:5
3. If A, B, C are the angles of a triangle, then $\cot \frac{A}{2} + \cot \frac{B}{2} + \cot \frac{C}{2}$ is equal to
 a) $\frac{s}{R}$ b) $\frac{R}{s}$ c) $\frac{\Delta}{s^2}$ d) $\frac{s^2}{\Delta}$
4. Coordinates of the foot of the perpendicular drawn from $(0, 0)$ to the line joining $(a \cos \alpha, a \sin \alpha)$ and $(a \cos \beta, a \sin \beta)$ are
 a) $(\frac{a}{2}, \frac{b}{2})$ b) $(\frac{a}{2}(\cos \alpha + \cos \beta), \frac{a}{2}(\sin \alpha + \sin \beta))$
 c) $(\cos \frac{\alpha + \beta}{2}, \sin \frac{\alpha + \beta}{2})$ d) $(0, \frac{b}{2})$
5. Three points are $A(6, 3), B(-3, 5), C(4, -2)$ and $P(x, y)$ is a point, then the ratio of area of ΔPBC and ΔABC is
 a) $|\frac{x + y - 2}{7}|$ b) $|\frac{x - y + 2}{2}|$ c) $|\frac{x - y - 2}{7}|$ d) None of these
6. Two vertical poles 20 m and 80 m stands apart on a horizontal plane. The height of the point of intersection of the lines joining the top of each pole to the foot of the other is
 a) 15 m b) 16 m c) 18 m d) 50 m
7. A person on a ship sailing north sees two lighthouses which are 6 km apart, in a line due west. After an hour's sailing one of them bears south west and the other southern south west. The ship is travelling at a rate of
 a) 12 km/hr b) 6 km/hr c) $3\sqrt{2}$ km/hr d) $(6 + 3\sqrt{2})$ km/hr
8. If α, β, γ are the real roots of the equation $x^3 - 3px^2 + 3qx - 1 = 0$,
 Then the centroid of the triangle whose vertices are $(\alpha, \frac{1}{\alpha}), (\beta, \frac{1}{\beta})$ and $(\gamma, \frac{1}{\gamma})$, is
 a) (p, q) b) (q, p) c) $(-p, q)$ d) $(q, -p)$
9. If two vertices of a triangle are $(-2, 3)$ and $(5, -1)$. Orthocentre lies at the origin and centroid on the line $x + y = 7$, then the third vertex lies at
 a) $(7, 4)$ b) $(8, 14)$ c) $(12, 21)$ d) None of these
10. What is the equation of the locus of a point which moves such that 4 times its distance from the x -axis is the square of its distance from the origin?
 a) $x^2 + y^2 - 4y = 0$ b) $x^2 + y^2 - 4|y| = 0$ c) $x^2 + y^2 - 4x = 0$ d) $x^2 + y^2 - 4|x| = 0$
11. If $a^2 + b^2 = c^2$, then $s(s - a)(s - b)(s - c)$ is equal to
 a) a^2b^2 b) $\frac{1}{4}a^2b^2$ c) $\frac{1}{2}a^2b^2$ d) $\frac{1}{2}ab$
12. The harmonic conjugate of $(4, -2)$ with respect to $(2, -4)$ and $(7, 1)$ is
 a) $(-8, -14)$ b) $(2, 3)$ c) $(-2, -3)$ d) $(13, -5)$

13. If O is the origin and $P(x_1, y_1), Q(x_2, y_2)$ are two points, then $OP \cdot OQ \sin \angle POQ =$
 a) $x_1x_2 + y_1y_2$ b) $x_1y_2 + x_2y_1$ c) $|x_1y_2 - x_2y_1|$ d) None of these
14. If ΔABC , if $a = 3, b = 4, c = 5$, then the value of $\sin 2B$ is
 a) $4/5$ b) $3/20$ c) $24/25$ d) $1/50$
15. From an aeroplane vertically over a straight horizontal road, the angles of depression of two consecutive milestones on opposite sides of the aeroplane are observed to be α and β . The height of the aeroplane above the road is
 a) $\frac{\tan \alpha + \tan \beta}{\tan \alpha \tan \beta}$ b) $\frac{\tan \alpha \tan \beta}{\tan \alpha + \tan \beta}$ c) $\frac{\cot \alpha \cot \beta}{\cot \alpha + \cot \beta}$ d) None of these
16. In ΔABC , if $\angle A = 45^\circ, \angle B = 75^\circ$, then $a + c\sqrt{2}$ is equal to
 a) 0 b) 1 c) b d) $2b$
17. Three vertical poles of heights h_1, h_2 and h_3 at the vertices A, B and C of a ΔABC subtend angles α, β and γ respectively at the circumcentre of the triangle. If $\cot \alpha, \cot \beta$ and $\cot \gamma$ are in AP, then h_1, h_2, h_3 are in
 a) AP b) GP c) HP d) None of these
18. The area enclosed within the curve $|x| + |y| = 1$ is
 a) 1 sq unit b) $2\sqrt{2}$ sq units c) $\sqrt{2}$ sq units d) 2 sq units
19. P is a point on the segment joining the feet of two vertical poles of height a and b . The angles of elevation of the top of the poles from P are 45° each. Then, the square of the distance between the top of the poles is
 a) $\frac{a^2 + b^2}{2}$ b) $a^2 + b^2$ c) $2(a^2 + b^2)$ d) $4(a^2 + b^2)$
20. By rotating the coordinates axes through 30° in anticlockwise sense the equation $x^2 + 2\sqrt{3}xy - y^2 = 2a^2$ changes to
 a) $X^2 - Y^2 = 3a^2$ b) $X^2 - Y^2 = a^2$ c) $X^2 - Y^2 = 2a^2$ d) None of these
21. The x -coordinate of the incentre of the triangle where the mid points of the sides are $(0, 1), (1, 1)$ and $(1, 0)$ is
 a) $2 + \sqrt{2}$ b) $1 + \sqrt{2}$ c) $2 - \sqrt{2}$ d) $1 - \sqrt{2}$
22. Let $A(2, -3)$ and $B(-2, 1)$ be vertices of a triangle ABC . If the centroid of this triangle moves on the line $2x + 3y = 1$, then the locus of the vertex C is the line
 a) $2x + 3y = 9$ b) $2x - 3y = 7$ c) $3x + 2y = 5$ d) $3x - 2y = 3$
23. The angle of elevation of the top of a tower at a point on the ground is 30° . If on walking 20 m toward the tower the angle of elevation becomes 60° , then the height of the tower is
 a) 10 m b) $\frac{10}{\sqrt{3}}$ m c) $10\sqrt{3}$ m d) None of these
24. In a ΔABC , if $2s = a + b + c$ and $(s - b)(s - c) = x \sin^2 \frac{A}{2}$, then the value of x is
 a) bc b) ca c) ab d) abc
25. If p_1, p_2 denote the length of the perpendiculars from the origin on the lines $x \sec \alpha + y \operatorname{cosec} \alpha = 2a$ and $x \cos \alpha + y \sin \alpha = a \cos 2\alpha$ respectively, then $\left(\frac{p_1}{p_2}, \frac{p_2}{p_1}\right)^2$ is equal to
 a) $4 \sin^2 4\alpha$ b) $4 \cos^2 4\alpha$ c) $4 \operatorname{cosec}^2 4\alpha$ d) $4 \sec^2 4\alpha$
26. The equation $\sqrt{(x - 2)^2 + (y - 1)^2} + \sqrt{(x + 2)^2 + (y - 4)^2} = 5$ represents
 a) Circle b) Ellipse c) Line segment d) None of these
27. The value of $\frac{1}{r_1^2} + \frac{1}{r_2^2} + \frac{1}{r_3^2} + \frac{1}{r^2}$ is
 a) 0 b) $\frac{a^2 + b^2 + c^2}{\Delta^2}$ c) $\frac{\Delta^2}{a^2 + b^2 + c^2}$ d) $\frac{a^2 + b^2 + c^2}{\Delta}$
28. The sides of a triangle are 4cm, 5cm and 6cm. the area of the triangle is equal to
 a) $\frac{15}{4} \text{ cm}^2$ b) $\frac{15}{4}\sqrt{7} \text{ cm}^2$ c) $\frac{4}{15}\sqrt{7} \text{ cm}^2$ d) None of these

29. A vertical lamp-post, 6 m high, stands at a distance of 2 m from a wall, 4 m high. A 1.5 m tall man starts to walk away from the wall on the other side of the wall, in line with the lamp-post the maximum distance to which the man can walk remaining in the shadow is
 a) $\frac{5}{2}$ m b) $\frac{3}{2}$ m c) 4 m d) None of these
30. A tower subtends an angle α at a point A in the plane of its base and the angle of depression of the foot of the tower at a point b feet just above A is β . Then, the height of the tower is
 a) $b \tan \alpha \cot \beta$ b) $b \cot \alpha \tan \beta$ c) $b \cot \alpha \cot \beta$ d) $b \tan \alpha \tan \beta$
31. In a ΔABC , if $b = 2$, $\angle B = 30^\circ$, then the area of the circumcircle of ΔABC in square unit is
 a) π b) 2π c) 4π d) 6π
32. The base of a cliff is circular. From the extremities of a diameter of the base of angle of elevation of the top of the cliff are 30° and 60° . If the height of the cliff be 500 m, then the diameter of the base of the cliff is
 a) $1000\sqrt{3}$ m b) $\frac{2000}{\sqrt{3}}$ m c) $\frac{1000}{\sqrt{3}}$ m d) $\frac{2000}{\sqrt{2}}$ m
33. If R denotes circumradius, then in ΔABC , $\frac{b^2 - c^2}{2aR}$ is equal to
 a) $\cos(B - C)$ b) $\sin(B - C)$ c) $\cos B - \cos C$ d) None of these
34. The area between the curve $y = 1 - |x|$ and the x -axis is equal to
 a) 1 sq unit b) $\frac{1}{2}$ sq unit c) $\frac{1}{3}$ sq unit d) 2 sq units
35. Angles A, B and C of a triangle are in AP with common difference 15 degree, then angle A is equal to
 a) 45° b) 60° c) 75° d) 30°
36. In a triangle $\left(1 - \frac{r_1}{r_2}\right)\left(1 - \frac{r_1}{r_3}\right) = 2$, then the triangle is
 a) Right angled b) Equilateral c) Isosceles d) None of these
37. The angle of elevation of the sun, if the length of the shadow of a tower is $\sqrt{3}$ times the height of the pole, is
 a) 150° b) 30° c) 60° d) 45°
38. If the equation $2x^2 + y^2 - 4x - 4y = 0$ is transformed to the equation $2X^2 + Y^2 - 8X - 8Y + 18 = 0$ by shifting the origin at a point P without rotating the coordinates axes, then the coordinates of P are
 a) (1, 2) b) (1, -2) c) (-1, 2) d) (-1, -2)
39. A vertical pole PS has two marks Q and R such that the portions PQ, PR and PS subtend angles α, β, γ at a point on the ground distance x from the pole. If $PQ = a, PR = b, PS = c$ and $\alpha + \beta + \gamma = 180^\circ$ then x^2 is equal to
 a) $\frac{a}{a + b + c}$ b) $\frac{b}{a + b + c}$ c) $\frac{c}{a + b + c}$ d) $\frac{abc}{a + b + c}$
40. If in a ΔABC , $(s - a)(s - b) = s(s - c)$, then angle C is equal to
 a) 90° b) 45° c) 30° d) 75°
41. At a point on the ground the angle of elevation of a tower is such that its cotangent is $\frac{3}{5}$. On walking 32 m towards the tower the cotangent of the angle of elevation is $\frac{2}{5}$. The height of the tower is
 a) 160 m b) 120 m c) 64 m d) None of these
42. Area of quadrilateral whose vertices are (2, 3), (3, 4), (4, 5) and (5, 6), is equal to
 a) 0 b) 4 c) 6 d) None of these
43. If the area of a triangle ABC is Δ , then $a^2 \sin 2B + b^2 \sin 2A$ is equal to
 a) 3Δ b) 2Δ c) 4Δ d) -4Δ
44. Consider the following statements :
 1. If in a ΔABC , $\frac{\sin A}{\sin C} = \frac{\sin(A-B)}{\sin(B-C)}$, then a^2, b^2, c^2 are in AP
 2. If exradius r_1, r_2 and r_3 of a ΔABC are in HP, then the sides a, b, c are in AP
 Which of these is/are correct?
 a) Only (1) b) Only (2) c) Both (1) and (2) d) None of these

45. If the sides of the triangle are $p, q, \sqrt{p^2 + q^2 + pq}$, then the greatest angle is
 a) $\frac{\pi}{2}$ b) $\frac{5\pi}{4}$ c) $\frac{2\pi}{3}$ d) $\frac{7\pi}{4}$
46. If x, y, z are perpendicular drawn from the vertices of triangle having sides a, b and c , then the value of $\frac{bx}{c} + \frac{cy}{a} + \frac{az}{b}$ will be
 a) $\frac{a^2 + b^2 + c^2}{2R}$ b) $\frac{a^2 + b^2 + c^2}{R}$ c) $\frac{a^2 + b^2 + c^2}{4R}$ d) $\frac{2(a^2 + b^2 + c^2)}{R}$
47. A balloon is observed simultaneously from three points A, B and C on a straight road directly under it. The angular elevation at B is twice and at C is thrice that of A . If the distance between A and B is 200 m and the distance between B and C is 100 m, then the height of balloon is given by
 a) 50 m b) $50\sqrt{3}$ m c) $50\sqrt{2}$ m d) None of these
48. If the distance of any point P from the points $A(a + b, a - b)$ and $B(a - b, a + b)$ are equal, then the locus of P is
 a) $x - y = 0$ b) $ax + by = 0$ c) $bx - ay = 0$ d) $x + y = 0$
49. The length of altitude through A of the ΔABC , where $A \equiv (-3, 0), B \equiv (4, -1), C \equiv (5, 2)$, is
 a) $\frac{2}{\sqrt{10}}$ b) $\frac{4}{\sqrt{10}}$ c) $\frac{11}{\sqrt{10}}$ d) $\frac{22}{\sqrt{10}}$
50. Triangle ABC has vertices $(0, 0), (11, 60)$ and $(91, 0)$. If the line $y = kx$ cuts the triangle into two triangles of equal area, then k is equal to
 a) $\frac{30}{51}$ b) $\frac{4}{7}$ c) $\frac{7}{4}$ d) $\frac{30}{91}$
51. A pole stands at the centre of a rectangular field and it subtends angles of 15° and 45° at the mid points of the side of the field. If the length of its diagonal is 1200 m, then the height of flag staff is
 a) 400 m b) 200 m c) $300\sqrt{2 + \sqrt{3}}$ m d) $300\sqrt{2 - \sqrt{3}}$ m
52. What is the equation of the locus a point which moves such that 4 times its distance from the x -axis is the square of its distance from the origin?
 a) $x^2 - y^2 - 4y = 0$ b) $x^2 + y^2 - 4|y| = 0$ c) $x^2 + y^2 - 4x = 0$ d) $x^2 + y^2 - 4|x| = 0$
53. A person standing on the bank of a river, observe that the angle of elevation of the top of a tree on the opposite bank of the river is 60° and when he retries 40m a way from the tree the angle of elevation become 30° . The breadth of the river is
 a) 20 m b) 30 m c) 40 m d) 60 m
54. There exist a ΔABC satisfying
 a) $\tan A + \tan B + \tan C = 0$ b) $\frac{\sin A}{2} = \frac{\sin B}{3} = \frac{\sin C}{1}$
 c) $\sin A + \sin B = -\left(\frac{\sqrt{3} + 1}{2\sqrt{2}}\right) \cos A \cos B$ d) $(a + b)^2 = c^2 + ab$ and $\sqrt{2}(\sin A + \cos A) = \sqrt{3}$
 $= \frac{\sqrt{3}}{4} = \sin A \sin B$
55. From a point a meters above a lake the angle of elevation of a cloud is α and the angle of depression of its reflection is β . The height of the cloud is
 a) $\frac{a \sin(\alpha + \beta)}{\sin(\alpha + \beta)}$ m b) $\frac{a \sin(\alpha + \beta)}{\sin(\beta - \alpha)}$ m c) $\frac{a \sin(\beta - \alpha)}{\sin(\alpha + \beta)}$ d) None of these
56. The orthocentre of the triangle formed by $(0, 0), (8, 0), (4, 6)$ is
 a) $\left(4, \frac{8}{3}\right)$ b) $(3, 4)$ c) $(4, 3)$ d) $(-3, 4)$
57. The x -coordinate of the incentre of the triangle where the mid point of the sides are $(0, 1), (1, 1)$ and $(1, 0)$, is
 a) $2 + \sqrt{2}$ b) $1 + \sqrt{2}$ c) $2 - \sqrt{2}$ d) $1 - \sqrt{2}$
58. The locus of the point (x, y) which is equidistant from the points $(a + b, b - a)$ and $(a - b, a + b)$ is

- a) $ax = by$ b) $ax + by = 0$ c) $bx + ay = 0$ d) $bx - ay = 0$
59. If the sum of the distances from two perpendicular lines in a plane is 1, then its locus is
a) A square b) A circle
c) A straight line d) Two intersecting lines
60. A tower of x metres high, has a flagstaff at its top. The tower and the flagstaff subtend equal angles at a point distant y metres from the foot of the tower. Then the length of the flagstaff (in meters), is
a) $\frac{y(x^2 - y^2)}{(x^2 + y^2)}$ b) $\frac{x(y^2 + x^2)}{(y^2 - x^2)}$ c) $\frac{x(x^2 + y^2)}{(x^2 - y^2)}$ d) $\frac{x(x^2 - y^2)}{(x^2 + y^2)}$
61. In a ΔABC , $2ac \sin \frac{A-B+C}{2}$ is equal to
a) $a^2 + b^2 - c^2$ b) $c^2 + a^2 - b^2$ c) $b^2 - a^2 - c^2$ d) $c^2 - a^2 - b^2$
62. If $P = (1, 0)$, $Q = (-1, 0)$ and $R = (2, 0)$ are three given points, then the locus of the point $S(x, y)$ satisfying the relation $SQ^2 + SR^2 = 2SP^2$ is
a) A straight line parallel to x -axis b) A circle through the origin
c) A circle with centre at the origin d) A straight line parallel to y -axis
63. If orthocenter and circumcentre of a triangle are respectively $(1, 1)$ and $(3, 2)$, then the coordinates of its centroid are
a) $(\frac{7}{3}, \frac{5}{3})$ b) $(\frac{5}{3}, \frac{7}{3})$ c) $(7, 5)$ d) None of these
64. The locus of the point of intersection of the lines $x \cot \theta + y \operatorname{cosec} \theta = 2$ and $x \operatorname{cosec} \theta + y \cot \theta = 6$ is
a) A straight line b) Circle c) A hyperbola d) An ellipse
65. In ΔABC , if $\cot A, \cot B, \cot C$ be in AP, then a^2, b^2, c^2 are in
a) HP b) GP c) AP d) None of these
66. The angles of elevation of the cloud at a point 2500 m high from the lake is 15° and the angle of depression of its reflection to the lake is 45° . Then the height of cloud from the foot of lake is
a) $2500\sqrt{3}$ m b) 2500 m c) $500\sqrt{3}$ m d) None of these
67. ABC is a triangular park with $AB = AC = 100$ m. A clock tower is situated at the mid point of BC . The angle of elevation, if the top of the tower at A and B are $\cot^{-1} 3.2$ and $\operatorname{cosec}^{-1} 2.6$ respectively. The height of the tower is
a) 16 m b) 25 m c) 50 m d) None of these
68. In ΔABC , $b = \sqrt{3}$, $c = 1$ and $\angle A = 30^\circ$, then the largest angle of the triangle is
a) 60° b) 135° c) 90° d) 120°
69. In an equilateral triangle, $R:r:r_1$ is equal to
a) 1:1:1 b) 1:2:3 c) 2:1:3 d) 3:2:4
70. In a triangle, if $r_1 = 2r_2 = 3r_3$, then $\frac{a}{b} + \frac{b}{c} + \frac{c}{a}$ is equal to
a) $\frac{75}{60}$ b) $\frac{155}{60}$ c) $\frac{176}{60}$ d) $\frac{191}{60}$
71. In a triangle ABC, $a : b : c = 4 : 5 : 6$. The ratio of the radius of the circumcircle to that of the incircle is
a) $\frac{15}{4}$ b) $\frac{11}{5}$ c) $\frac{16}{7}$ d) $\frac{16}{3}$
72. An aeroplane flying with uniform speed horizontally one kilometer above the ground is observed at an elevation of 60° . After 10 s, if the elevation is observed to be 30° , then the speed of the plane (in km/h) is
a) $\frac{240}{\sqrt{3}}$ b) $200\sqrt{3}$ c) $240\sqrt{3}$ d) $\frac{120}{\sqrt{3}}$
73. The angle of elevation of the top of a tower standing on a horizontal plane from a point A is α . After walking a distance a towards the foot of the tower the angle of elevation is found to be β . The height of the tower is
a) $\frac{a \sin \alpha \sin \beta}{\sin(\beta - \alpha)}$ b) $\frac{a \sin \alpha \sin \beta}{\sin(\alpha - \beta)}$ c) $\frac{a \sin(\beta - \alpha)}{\sin \alpha \sin \beta}$ d) $\frac{a \sin(\alpha - \beta)}{\sin \alpha \sin \beta}$
74. If the vertices of a triangle have integral coordinates, the triangle cannot be

- a) An equilateral triangle
c) An isosceles triangle
- b) A right angled triangle
d) None of the above
75. In a ΔABC , among the following which one is true?
 a) $(b + c) \cos \frac{A}{2} = a \sin \left(\frac{B + C}{2} \right)$
 b) $(b + c) \cos \left(\frac{B + C}{2} \right) = a \sin \frac{A}{2}$
 c) $(b - c) \cos \left(\frac{B - C}{2} \right) = a \cos \left(\frac{A}{2} \right)$
 d) $(b - c) \cos \frac{A}{2} = a \sin \left(\frac{B - C}{2} \right)$
76. The upper $\left(\frac{3}{4} \right)$ th portion of a vertical pole subtends an angle $\tan^{-1} \left(\frac{3}{5} \right)$ at a point in the horizontal plane through its foot and at a distance 40 m from the foot. A possible height of the vertical pole is
 a) 20 m
b) 40 m
c) 60 m
d) 80 m
77. If C and D are the points of internal and external division of line segment AB in the same ratio, then AC, AB, AD are in
 a) AP
b) GP
c) HP
d) AGP
78. A ladder rests against a vertical wall at angle α to the horizontal. If its foot is pulled away from the wall through a distance ' a ' so that it slides a distance ' b ' down the wall making an angle β with the horizontal, then $a =$
 a) $b \tan \left(\frac{\alpha - \beta}{2} \right)$
 b) $b \tan \left(\frac{\alpha + \beta}{2} \right)$
 c) $b \cot \left(\frac{\alpha - \beta}{2} \right)$
 d) None of these
79. The angles A, B and C of a ΔABC are in A.P. If $AB = 6, BC = 7$, then $AC =$
 a) 5
b) 7
c) 8
d) None of these
80. The locus of a point whose difference of distance from points $(3, 0)$ and $(-3, 0)$ is 4, is
 a) $\frac{x^2}{4} - \frac{y^2}{5} = 1$
 b) $\frac{x^2}{5} - \frac{y^2}{4} = 1$
 c) $\frac{x^2}{2} - \frac{y^2}{3} = 1$
 d) $\frac{x^2}{3} - \frac{y^2}{2} = 1$
81. If a ΔABC , if $a^4 + b^4 + c^4 = 2c^2(a^2 + b^2)$, then $\angle C$ is equal to
 a) 60°
b) 135°
c) 90°
d) 75°
82. Given the points $A(0, 4)$ and $B(0, -4)$, then the equation of the locus of the point $P(x, y)$ such that, $|AP - BP| = 6$, is
 a) $\frac{x^2}{7} + \frac{y^2}{9} = 1$
 b) $\frac{x^2}{9} + \frac{y^2}{7} = 1$
 c) $\frac{x^2}{7} - \frac{y^2}{9} = 1$
 d) $\frac{y^2}{9} - \frac{x^2}{7} = 1$
83. If in ΔABC , $\sin \frac{A}{2} \sin \frac{C}{2} = \sin \frac{B}{2}$ and $2s$ is the perimeter of the triangle, then s is
 a) $2b$
b) b
c) $3b$
d) $4b$
84. The angle of depression of a ship from the top of a tower 30 m high is 60° . Then the distance of ship from the base of tower is
 a) 30 m
b) $30\sqrt{3}$ m
c) $10\sqrt{3}$ m
d) 10 m
85. At a distance $2h$ m from the foot of a tower of height h m the top of the tower and a pole at the top of the tower subtend equal angles. Height of the pole should be
 a) $\frac{5h}{3}$ m
b) $\frac{4h}{3}$ m
c) $\frac{7h}{5}$ m
d) $\frac{3h}{2}$ m
86. From the tower 60 m high angles of depression of the top and bottom of a house are α and β respectively. If the height of the house is $\frac{60 \sin(\beta - \alpha)}{x}$, then x is equal to
 a) $\sin \alpha \sin \beta$
b) $\cos \alpha \cos \beta$
c) $\sin \alpha \cos \beta$
d) $\cos \alpha \sin \beta$
87. In a triangle, the lengths of the two larger sides are 10 cm and 9 cm respectively. If the angles of the triangle are in AP, then the length of the third side in cm can be
 a) $5 - \sqrt{6}$ only
b) $5 + \sqrt{6}$ only
c) $5 - \sqrt{6}$ or $5 + \sqrt{6}$
d) Neither $5 - \sqrt{6}$ nor $5 + \sqrt{6}$
88. In ΔABC , if $\sin^2 \frac{A}{2}, \sin^2 \frac{B}{2}, \sin^2 \frac{C}{2}$ be in HP. Then, a, b, c will be in
 a) AP
b) GP
c) HP
d) None of these
89. The angles of elevation of the top of a tower at the top and the foot of a pole of height 10 m are 30° and 60° respectively. The height of the tower is



- a) 10 m b) 15 m c) 20 m d) None of these
90. If the points $(1, 1)$, $(-1, -1)$, $(-\sqrt{3}, \sqrt{3})$ are the vertices of a triangle, then this triangle is
a) Right angled b) Isosceles c) Equilateral d) None of these
91. The vertices of a family of triangles have integer coordinates. If two of the vertices of all the triangles are $(0, 0)$ and $(6, 8)$, then the least value of areas of the triangles is
a) 1 b) $\frac{3}{2}$ c) 2 d) $\frac{5}{2}$
92. In a ΔABC , $(\cot \frac{A}{2} + \cot \frac{B}{2})(a \sin^2 \frac{B}{2} + b \sin^2 \frac{A}{2})$ is equal to
a) $\cot C$ b) $c \cot C$ c) $\cot \frac{C}{2}$ d) $c \cot \frac{C}{2}$
93. The intercepts on the straight line $y = mx$ by the line $y = 2$ and $y = 6$ is less than 5, then m belongs to
a) $]-\frac{4}{3}, \frac{4}{3}[$ b) $]\frac{4}{3}, \frac{3}{8}[$ c) $]-\infty, -\frac{4}{3}[\cup]\frac{4}{3}, \infty[$ d) $]\frac{4}{3}, \infty[$
94. In ΔABC , $(b - c) \sin A + (c - a) \sin B + (a - b) \sin C$ is equal to
a) $ab + bc + ca$ b) $a^2 + b^2 + c^2$ c) 0 d) None of these
95. The inradius of the triangle whose sides are 3, 5, 6 is
a) $\sqrt{\frac{8}{7}}$ b) $\sqrt{8}$ c) $\sqrt{7}$ d) $\sqrt{\frac{7}{8}}$
96. In a ΔABC , if the sides are $a = 3$, $b = 5$ and $c = 4$, then $\sin \frac{B}{2} + \cos \frac{B}{2}$ is equal to
a) $\sqrt{2}$ b) $\frac{\sqrt{3} + 1}{2}$ c) $\frac{\sqrt{3} - 1}{2}$ d) 1
97. The elevation of an object on a hill is observed from a certain point in the horizontal plane through its base, to be 30° . After walking 120 m towards it on level ground the elevation is found to be 60° . Then the height of the object (in metres) is
a) 120 b) $60\sqrt{3}$ c) $120\sqrt{3}$ d) 60
98. If the area of the triangle with vertices $(x, 0)$, $(1, 1)$ and $(0, 2)$ is 4 sq unit, then the value of x is
a) -2 b) -4 c) -6 d) 8
99. At a distance 12 metres from the foot A of a tower AB of height 5 metres, a flagstaff BC on top of AB and the tower subtend the same angle. The, the height of flagstaff is
a) $\frac{1440}{119}$ metres b) $\frac{475}{119}$ metres c) $\frac{845}{119}$ metres d) None of these
100. A tower 50 m high, stands on top of a mount, from a point on the ground the angles of elevation of the top and bottom of the tower are found to be 75° and 60° respectively. the height of the mount is
a) 25 m b) $25(\sqrt{3} - 1)$ m c) $25\sqrt{3}$ m d) $25(\sqrt{3} + 1)$ m
101. Let AB is divided internally and externally at P and Q in the same ratio. Then, AP , AB , AQ are in
a) AP b) GP c) HP d) None of these
102. If the sum of the distance of a point P from two perpendicular lines in a plane is 1, then the locus of P is a
a) Rhombus b) Circle c) Straight line d) Pair of straight lines
103. A flagpole stands on a building of height 450 ft and an observer on a level ground is 300 ft from the base of the building. The angle of elevation of the bottom of the flagpole is 30° and the height of the flagpole is 50ft. If θ is the angle of elevation of the top of the flagpole, then $\tan \theta$ is equal to
a) $\frac{4}{3\sqrt{3}}$ b) $\frac{\sqrt{3}}{2}$ c) $\frac{9}{2}$ d) $\frac{\sqrt{3}}{5}$
104. In ΔABC , $2(a \sin^2 \frac{C}{2} + c \sin^2 \frac{A}{2})$ is equal to
a) $a + b - c$ b) $c + a - b$ c) $b + c - a$ d) $a + b + c$
105. Orthocenter of triangle with vertices $(0, 0)$, $(3, 4)$ and $(4, 0)$ is
a) $(3, \frac{5}{4})$ b) $(3, 12)$ c) $(3, \frac{3}{4})$ d) $(3, 9)$



106. Three vertices of a parallelogram taken in order are $(-1, -6)$, $(2, -5)$ and $(7, 2)$. The fourth vertex is
 a) $(1, 4)$ b) $(4, 1)$ c) $(1, 1)$ d) $(4, 4)$
107. If in a ΔABC , $\cos A \cos B + \sin A \sin B \sin C = 1$, then the triangle is
 a) Isosceles b) Right angled c) Isosceles right angled d) Equilateral
108. If in a ΔABC , the sides AB and AC are perpendicular, then the true equation is
 a) $\tan A + \tan B = 0$ b) $\tan B + \tan C = 0$ c) $\tan A + 2 \tan C = 0$ d) $\tan B \tan C = 1$
109. The points $(1, 1)$, $(-5, 5)$ and $(13, \lambda)$ lie on the same straight line, if λ is equal to
 a) 7 b) -7 c) ± 7 d) 0
110. Circumcentre of triangle whose vertices are $(0, 0)$, $(3, 0)$ and $(0, 4)$ is
 a) $(\frac{3}{2}, 2)$ b) $(2, \frac{3}{2})$ c) $(0, 0)$ d) None of these
111. The vertices of a triangle are $A(-1, -7)$, $B(5, 1)$ and $C(1, 4)$. The equation of the bisector of angle ABC , is
 a) $x + 7y - 2 = 0$ b) $x - 7y - 2 = 0$ c) $x - 7y + 2 = 0$ d) None of these
112. A tower subtends angles α , 2α and 3α respectively at points A , B and C , all lying on a horizontal line through the foot of the tower, then $\frac{AB}{BC}$ is equal to
 a) $\frac{\sin 3\alpha}{\sin 2\alpha}$ b) $1 + 2 \cos 2\alpha$ c) $2 \cos 2\alpha$ d) $\frac{\sin 2\alpha}{\sin \alpha}$
113. A person standing on the bank of a river finds that the angle of elevation of the top of a tower on the opposite bank is 45° , then which of the following statements is correct?
 a) Breadth of the river is twice the height of the tower
 b) Breadth of the river and the height of the tower are the same
 c) Breadth of the river is half of the height of the tower
 d) None of these
114. The angular depression of the top and the foot of the chimney as seen from the top of a second chimney which is 150 m high and standing on the same level as the first are θ and ϕ respectively. The distance between their tops when $\tan \theta = \frac{4}{3}$ and $\tan \phi = \frac{5}{2}$ is equal to
 a) 50 m b) 100 m c) 15 m d) None of these
115. A round balloon of radius r subtends an angle α at the eye of the observer, While the angle of elevation of its centre is β . The height of the center of balloon is
 a) $r \operatorname{cosec} \alpha \sin \frac{\beta}{2}$ b) $r \sin \alpha \operatorname{cosec} \frac{\beta}{2}$ c) $r \sin \frac{\alpha}{2} \operatorname{cosec} \beta$ d) $r \operatorname{cosec} \frac{\alpha}{2} \sin \beta$
116. In a ΔABC , a, c, A are given and b_1, b_2 are two values, if the third side b such that $b_2 = 2b_1$, then $\sin A$ is equal to
 a) $\frac{\sqrt{9a^2 - c^2}}{8a^2}$ b) $\sqrt{\frac{9a^2 - c^2}{8c^2}}$ c) $\frac{\sqrt{9a^2 + c^2}}{8a^2}$ d) None of these
117. If a, b, c are sides of a triangle, then
 a) $\sqrt{a} + \sqrt{b} > \sqrt{c}$ b) $|\sqrt{a} - \sqrt{b}| > \sqrt{c}$ (if c is smallest)
 c) $\sqrt{a} + \sqrt{b} < \sqrt{c}$ d) None of the above
118. ABC Is a triangle with $\angle A = 30^\circ$, $BC = 10$ cm. The area of the circumcircle of the triangle is
 a) 100π sq cm b) 5 sq cm c) 25 sq cm d) $\frac{100\pi}{3}$ sq cm
119. In a ΔABC , $a : b : c = 4 : 5 : 6$. The ratio of the radius of the circumcircle to that of the incircle is
 a) $\frac{16}{9}$ b) $\frac{16}{7}$ c) $\frac{11}{7}$ d) $\frac{7}{16}$
120. The incentre of the triangle formed by lines $x = 0$, $y = 0$ and $3x + 4y = 12$, is at
 a) $(\frac{1}{2}, \frac{1}{2})$ b) $(1, 1)$ c) $(1, \frac{1}{2})$ d) $(\frac{1}{2}, 1)$
121. Given points are $A(0, 4)$ and $B(0, -4)$, the locus of $P(x, y)$ such that $|AP - BP| = 6$, is
 a) $9x^2 - 7y^2 + 63 = 0$ b) $9x^2 + 7y^2 - 63 = 0$ c) $9x^2 + 7y^2 + 63 = 0$ d) None of these

122. The angle of elevation of the top of a tower from a point A due South of the tower is α and from a point B due East of the tower is β . If $AB = d$, then the height of the tower is
- a) $\frac{d}{\sqrt{\tan^2 \alpha - \tan^2 \beta}}$ b) $\frac{d}{\sqrt{\tan^2 \alpha + \tan^2 \beta}}$ c) $\frac{d}{\sqrt{\cot^2 \alpha + \cot^2 \beta}}$ d) $\frac{d}{\sqrt{\cot^2 \alpha - \cot^2 \beta}}$
123. Let P be the point $(1, 0)$ and Q be the point on $y^2 = 8x$. The locus of mid point of PQ is
- a) $x^2 - 4y + 2 = 0$ b) $x^2 + 4y + 2 = 0$ c) $y^2 + 4x + 2 = 0$ d) $y^2 - 4x + 2 = 0$
124. Let $A(k, 2)$ and $B(3, 5)$ are points. The point (t, t) divide \overline{AB} from A 's side in the ratio of k , then $k = \dots, k \in R - \{0, -1\}$
- a) -4 b) -2 c) 4 d) 2
125. If a, b, c the sides of a ΔABC are in AP and a is the smallest side, then $\cos A$ equals
- a) $\frac{3c - 4b}{2c}$ b) $\frac{3c - 4b}{2b}$ c) $\frac{4c - 3b}{2c}$ d) None of these
126. Area of the triangle formed by the lines $y = 2x, y = 3x$ and $y = 5$ is equal to (in square unit)
- a) $\frac{25}{6}$ b) $\frac{25}{12}$ c) $\frac{5}{6}$ d) $\frac{17}{12}$
127. The angles of depression of the top and the foot of a chimney as seen from the top of a second chimney, which is 150 m high and standing on the same level as the first are θ and ϕ respectively, then the distance between their tops when $\tan \theta = \frac{4}{3}$ and $\tan \phi = \frac{5}{2}$, is
- a) $\frac{150}{\sqrt{3}}$ m b) $100\sqrt{3}$ m c) 150 m d) 100 m
128. If one side of a triangle is double the other and the angles opposite to these sides differ by 60° , then the triangle is
- a) Obtuse angled b) Acute angled c) Isosceles d) Right angled
129. If the three points $(3q, 0), (0, 3p)$ and $(1, 1)$ are collinear then which one is true?
- a) $\frac{1}{p} + \frac{1}{q} = 0$ b) $\frac{1}{p} + \frac{1}{q} = 1$ c) $\frac{1}{p} + \frac{1}{q} = 3$ d) $\frac{1}{p} + \frac{3}{q} = 1$
130. If in a $\Delta ABC, a = 15, b = 36, c = 39$, then $\sin \frac{C}{2}$ is equal to
- a) $\frac{\sqrt{3}}{2}$ b) $\frac{1}{2}$ c) $\frac{1}{\sqrt{2}}$ d) $-\frac{1}{\sqrt{2}}$
131. In a ΔABC , let $\angle C = \frac{\pi}{2}$, if r is the inradius and R is the circumradius of the ΔABC , then $2(r + R)$ equals
- a) $c + a$ b) $a + b + c$ c) $a + b$ d) $b + c$
132. From the top of a light house 60 m high with its base at the sea level the angle of depression of a boat is 15° . The distance of the boat from the foot of light house is
- a) $\left(\frac{\sqrt{3}-1}{\sqrt{3}+1}\right) 60$ m b) $\left(\frac{\sqrt{3}+1}{\sqrt{3}-1}\right) 60$ m c) $\frac{\sqrt{3}+1}{\sqrt{3}-1}$ m d) None of these
133. If $\cos^2 A + \cos^2 C = \sin^2 B$, then ΔABC is
- a) Equilateral b) Right angled c) Isosceles d) None of these
134. The sides of triangle are in the ratio $1:\sqrt{3}:2$, then the angles of the triangle are in ratio
- a) 1:3:5 b) 2:3:1 c) 3:2:1 d) 1:2:3
135. A tower stands at the top of a hill whose height is 3 times the height of the tower. The tower is found to subtend at a point 3 km away on the horizontal through the foot of the hill, an angle θ , where $\tan \theta = \frac{1}{9}$. The height of the tower is
- a) 12 b) 3 c) $\frac{9 \pm \sqrt{33}}{8}$ d) None of these
136. Angles A, B and C of a ΔABC are in AP. If $\frac{b}{c} = \frac{\sqrt{3}}{\sqrt{2}}$, then angle A is equal to
- a) $\frac{\pi}{6}$ b) $\frac{\pi}{4}$ c) $\frac{5\pi}{12}$ d) $\frac{\pi}{2}$

137. The angle of depression of a boat in a river is 30° from the top of a tower, 87 m high and the speed of the boat is 5.8 km/h. The time taken by the boat to reach at the base of the tower is
 a) 9 min b) $\frac{9\sqrt{3}}{10}$ min c) 25 min d) 15 min
138. If the centroid of the triangle formed by the points (a, b) , (b, c) and (c, a) is at the origin, then $a^3 + b^3 + c^3 =$
 a) 0 b) abc c) $3abc$ d) $-3abc$
139. The sides of a ΔABC are $BC = 5$, $CA = 4$ and $AB = 3$. If A is at the origin and the bisector of the internal angle A meets BC in $D(12/7, 12/7)$, then the coordinates of the incentre, are
 a) $(2, 2)$ b) $(2, 3)$ c) $(3, 2)$ d) $(1, 1)$
140. If a, b and c are the sides of a triangle such that $a^4 + b^4 + c^4 = 2c^2(a^2 + b^2)$, then the angles opposite to the side C is
 a) 45° or 90° b) 30° or 135° c) 45° or 135° d) 60° or 120°
141. In radius of a circle which is inscribed in a isosceles triangle one of whose angle is $2\pi/3$, is $\sqrt{3}$, then area of triangle is
 a) $4\sqrt{3}$ b) $12 - 7\sqrt{3}$ c) $12 + 7\sqrt{3}$ d) None of these
142. A triangular park is enclosed on two sides by a fence and on the third side by a straight river bank. The two sides having fence are of same length x . The maximum area enclosed by the park is
 a) $\sqrt{\frac{x^3}{8}}$ b) $\frac{1}{2}x^2$ c) πx^2 d) $\frac{3}{2}x^2$
143. In a ΔABC , if $\tan \frac{A}{2} = \frac{5}{6}$, $\tan \frac{C}{2} = \frac{2}{5}$, then
 a) a, b, c are in AP b) a, b, c are in GP c) b, a, c are in AP d) a, b, c are in AP
144. The vertices P, Q, R of a triangle are $(2, 1)$, $(5, 2)$ and $(3, 4)$ respectively. Then, the circumcentre is
 a) $(\frac{13}{4}, -\frac{9}{4})$ b) $(-\frac{13}{4}, \frac{9}{4})$ c) $(-\frac{13}{4}, -\frac{9}{4})$ d) $(\frac{13}{4}, \frac{9}{4})$
145. In a ΔABC , $(a + b + c)(b + c - a) = kbc$, if
 a) $k < 0$ b) $k > 6$ c) $0 < k < 4$ d) $k > 4$
146. If $A(6, -3)$, $B(-3, 5)$, $C(4, -2)$, $P(\alpha, \beta)$, then the ratio of the areas of the triangles PBC, ABC is
 a) $|\alpha + \beta|$ b) $|\alpha - \beta|$ c) $|\alpha + \beta + 2|$ d) $|\alpha + \beta - 2|$
147. ABC is a triangular park with $AB = AC = 100$ m. A clock tower is situated at the mid point of BC . The angles of elevation of the top of the tower at A and B are $\cot^{-1} 3.2$ and $\operatorname{cosec}^{-1} 2.6$ respectively. The height of the tower is
 a) 50 m b) 25 m c) 40 m d) None of these
148. In a ΔABC , $a \cot A + b \cot B + c \cot C$ is equal to
 a) $r + R$ b) $r - R$ c) $2(r + R)$ d) $2(r - R)$
149. If $(1, a)$, $(2, b)$, and $(3, c)$; $a, b, c \in R$ are the vertices of a triangle, its centroid can
 a) Not be on x -axis b) Not be on y -axis c) Be on $(0, 0)$ d) None of these
150. The pair of lines $\sqrt{3}x^2 - 4xy + \sqrt{3}y^2 = 0$ are rotated about the origin by $\pi/6$ in the anti-clockwise sense. The equation of the pair in the new position is
 a) $\sqrt{3}y^2 - xy = 0$ b) $\sqrt{3}x^2 - xy = 0$ c) $x^2 - y^2 = 0$ d) $\sqrt{3}x^2 + xy = 0$
151. In triangle ABC , $a = 2$, $b = 3$ and $\sin A = \frac{2}{3}$ then B is equal to
 a) 30° b) 60° c) 90° d) 120°
152. If the sides of a right angle triangle form an AP, the 'sin' of the acute angles are
 a) $(\frac{3}{5}, \frac{4}{5})$ b) $(\sqrt{3}, \frac{1}{\sqrt{3}})$ c) $(\sqrt{\frac{\sqrt{5}-1}{2}}, \sqrt{\frac{\sqrt{5}-1}{2}})$ d) $(\sqrt{\frac{\sqrt{3}-1}{2}}, \sqrt{\frac{\sqrt{3}-1}{2}})$
153. In a ΔABC , $2a^2 + 4b^2 + c^2 = 4ab + 2ac$, then $\cos B$ is equal to

- a) 0 b) $\frac{1}{8}$ c) $\frac{3}{8}$ d) $\frac{7}{8}$

154. The line joining $A(b \cos \alpha, b \sin \alpha)$ and $B(a \cos \beta, a \sin \beta)$ is produced to the point $M(x, y)$ so that $AM:MB = b : a$, then

$x \cos\left(\frac{\alpha+\beta}{2}\right) + y \sin\left(\frac{\alpha+\beta}{2}\right)$ is

- a) -1 b) 0 c) 1 d) $a^2 + b^2$

155. A house of height 100 m subtends a right angle at the window of an opposite house. If the height of the window be 64m, then the distance between the two houses is

- a) 48 m b) 36 m c) 54 m d) 72 m

156. A vertical tower stands on a declivity which is inclined at 15° to the horizon. From the foot of the tower a man ascends the declivity for 80 ft and then, finds that the tower subtends an angle of 30° . The height of tower is

- a) $20(\sqrt{6} - \sqrt{2})$ ft b) $40(\sqrt{6} - \sqrt{2})$ ft c) $40(\sqrt{6} + \sqrt{2})$ ft d) None of these

157. $(0, -1)$ And $(0, 3)$ are two opposite vertices of a square. The other two vertices are

- a) $(0, 1), (0, -3)$ b) $(3, -1), (0, 0)$ c) $(2, 1), (-2, 1)$ d) $(2, 2), (1, 1)$

158. The points $(1, 3)$ and $(5, 1)$ are two opposite vertices of a rectangle. The other two vertices lie on the line $y = 2x + c$, are

- a) $(2, 0)$ and $(4, 4)$ b) $(2, 0)$ and $(-4, -4)$ c) $(2, 0)$ and $(-4, 4)$ d) $(-2, 0)$ and $(4, 4)$

159. If in a $\Delta ABC, r_3 = r_1 + r_2 + r$, then $\angle A + \angle B$ is equal to

- a) 120° b) 100° c) 90° d) 80°

160. In a triangle ABC , if $a = 3, b = 4, c = 5$, then the distance between its incentre and circumcentre is

- a) $\frac{1}{2}$ b) $\frac{\sqrt{3}}{2}$ c) $\frac{3}{2}$ d) $\frac{\sqrt{5}}{2}$

161. One side of length $3a$ of triangle of area a^2 square unit lies on the line $x = a$. Then, one of the lines on which the third vertex lies, is

- a) $x = -a^2$ b) $x = a^2$ c) $x = -a$ d) $x = \frac{a}{3}$

162. In a ΔABC , if D is the middle point BC and AD is perpendicular to AC , then $\cos B$ is equal to

- a) $\frac{2b}{a}$ b) $-\frac{b}{a}$ c) $\frac{b^2 + c^2}{ca}$ d) $\frac{c^2 + a^2}{ca}$

163. The angle of depression of a point situated at a distance of 70 metres from the base of a tower is 45° . The height of the tower is

- a) 70 m b) $70\sqrt{2}$ m c) $\frac{70}{\sqrt{2}}$ m d) 35 m

164. The sides of a triangle are three consecutive natural numbers and its largest angle is twice the smallest one. Then, the sides of the triangle are

- a) 1, 2, 3 b) 2, 3, 4 c) 3, 4, 5 d) 4, 5, 6

165. Consider the following statements :

1. $\frac{b^2 - c^2}{a \sin(B - C)} = 2R$

2. $a \sin(B - C) + b \sin(C - A) + c \sin(A - B) = 0$

Which of these is/are correct?

- a) Only (1) b) Only (2) c) Both (1) and (2) d) None of these

166. The four distinct point $(0, 0), (2, 0), (0, -2)$ and $(k, -2)$ are concyclic, if k is equal to

- a) -2 b) 2 c) 1 d) 0

167. If origin is shifted to $(7, -4)$, then point $(4, 5)$ shifted to

- a) $(-3, 9)$ b) $(3, 9)$ c) $(11, 1)$ d) None of these

168. In $\Delta ABC, (a + b + c) \left(\tan \frac{A}{2} + \tan \frac{B}{2}\right)$ is equal to

- a) $2c \cot \frac{C}{2}$ b) $2a \cot \frac{A}{2}$ c) $2b \cot \frac{B}{2}$ d) $\tan \frac{C}{2}$

169. In a ΔABC , sides a, b, c are in AP and $\frac{2}{1!9!} + \frac{2}{3!7!} + \frac{1}{5!5!} = \frac{8^a}{(2b)!}$, then the maximum value of $\tan A \tan B$ is equal to
- a) $\frac{1}{2}$ b) $\frac{1}{3}$ c) $\frac{1}{4}$ d) $\frac{1}{4}$
170. If the angle of elevation of two towers from the middle point of the line joining their feet be 60° and 30° respectively, then the ratio of their heights is
- a) 2:1 b) $1:\sqrt{2}$ c) 3:1 d) $1:\sqrt{3}$
171. In a ΔABC , $\angle C = 60^\circ$ then $\frac{1}{a+c} + \frac{1}{b+c}$ is equal to
- a) $\frac{1}{a+b+c}$ b) $\frac{2}{a+b+c}$ c) $\frac{3}{a+b+c}$ d) None of these
172. In ΔABC , if $(a+b+c)(a-b+c) = 3ac$, then
- a) $\angle B = 60^\circ$ b) $\angle B = 30^\circ$ c) $\angle C = 60^\circ$ d) $\angle A + \angle C = 90^\circ$
173. If a^2, b^2, c^2 are in AP, then which of the following are also in AP?
- a) $\sin A, \sin B, \sin C$ b) $\tan A, \tan B, \tan C$ c) $\cot A, \cot B, \cot C$ d) None of these
174. In a triangle ABC , if $\sin A \sin B = \frac{ab}{c^2}$, then the triangle is
- a) Equilateral b) Isosceles c) Right angled d) Obtuse angled
175. The perimeter of a ΔABC is 6 times the arithmetic mean of the sine ratios of its angles. If $a = 1$, then A is equal to
- a) $\frac{\pi}{6}$ b) $\frac{\pi}{3}$ c) $\frac{\pi}{2}$ d) $\frac{2\pi}{3}$
176. The centroid of the triangle ABC , where $A \equiv (2, 3)$, $B \equiv (8, 10)$ and $C \equiv (5, 5)$ is
- a) (5, 6) b) (6, 5) c) (6, 6) d) (15, 18)
177. The angle of elevation of the top of the tower observed from each of the tree point A, B, C on the ground forming a triangle is the same angle α . If R is the circumradius of the triangle ABC , then the height of the tower is
- a) $R \sin \alpha$ b) $R \cos \alpha$ c) $R \cot \alpha$ d) $R \tan \alpha$
178. The angle of elevation of the top of a hill from a point is α . After walking b metres towards the top up a slope inclined at an angle β to the horizon, the angle of elevation of the top becomes γ . Then, the height of the hill is
- a) $\frac{b \sin \alpha \sin(\gamma - \beta)}{\sin(\gamma - \alpha)}$ b) $\frac{b \sin \alpha \sin(\gamma - \alpha)}{\sin(\gamma - \beta)}$ c) $\frac{b \sin(\gamma - \beta)}{\sin(\gamma - \alpha)}$ d) $\frac{\sin(\gamma - \beta)}{b \sin \alpha \sin(\gamma - \alpha)}$
179. The area of the ΔABC , in which $a = 1, b = 2, \angle C = 60^\circ$, is
- a) 4 sq unit b) $\frac{1}{2}$ sq unit c) $\frac{\sqrt{3}}{2}$ sq unit d) $\sqrt{3}$ sq units
180. If t_1, t_2 and t_3 are distinct points $(t_1, 2at_1 + at_1^3), (t_2, 2at_2 + at_2^3)$ and $t_3, 2at_3 + at_3^3$ are collinear, if
- a) $t_1 t_2 t_3 = 1$ b) $t_1 + t_2 + t_3 = t_1 t_2 t_3$ c) $t_1 + t_2 + t_3 = 0$ d) $t_1 + t_2 + t_3 = -1$
181. If A and B are two points having coordinates (3, 4) and (5, -2) respectively and P is a point such that $PA = PB$ and area of triangle $PAB = 10$ sq unit, then the coordinates of P are
- a) (7, 4) and (13, 2) b) (7, 2) and (1, 0) c) (2, 7) and (4, 13) d) None of these
182. In ΔABC , $\angle A = \frac{\pi}{2}, b = 4, c = 3$, then the value of $\frac{R}{r}$ is equal to
- a) $\frac{5}{2}$ b) $\frac{7}{2}$ c) $\frac{9}{2}$ d) $\frac{35}{24}$
183. In the angles A, B and C of a triangular are in the arithmetic progression and if a, b and c denotes the lengths of the sides opposite to A, B and C respectively, then the value of the expression $\frac{a}{c} \sin 2C + \frac{c}{a} \sin 2A$ is
- a) $\frac{1}{2}$ b) $\frac{\sqrt{3}}{2}$ c) 1 d) $\sqrt{3}$

184. Two sides of a triangle are given by the roots of the equation $x^2 - 5x + 6 = 0$ and the angle between the sides is $\frac{\pi}{3}$. Then, the perimeter of the triangle is
 a) $5 + \sqrt{2}$ b) $5 + \sqrt{3}$ c) $5 + \sqrt{5}$ d) $5 + \sqrt{7}$
185. In a triangle ABC , if $\angle A = 60^\circ$, $a = 5$, $b = 4$, then c is a root of the equation
 a) $c^2 - 5c - 9 = 0$ b) $c^2 - 4c - 9 = 0$ c) $c^2 - 10c + 25 = 0$ d) $c^2 - 5c - 41 = 0$
186. The angle of elevation of the top of vertical tower from a point A on the horizontal ground is found to be $\frac{\pi}{4}$. From A , a man walks 10 m up a path sloping at a angle $\frac{\pi}{6}$. After this the slope becomes steeper and after walking up another 10 m, the man reaches the top of the tower. Distance of A from the foot of the tower is
 a) $5(1 + \sqrt{3})\text{m}$ b) $\frac{5}{2}(1 + \sqrt{3})\text{m}$ c) $5(\sqrt{3} - 1)\text{m}$ d) $\frac{5}{2}(\sqrt{3} - 1)\text{m}$
187. If the distance between the points $(a \cos \theta, a \sin \theta)$ and $(a \cos \phi, a \sin \phi)$ is $2a$, then θ is equal to
 a) $2n\pi \pm \pi + \phi, n \in Z$ b) $n\pi + \frac{\pi}{2} + \phi, n \in Z$
 c) $n\pi - \phi, n \in Z$ d) $2n\pi + \phi, n \in Z$
188. If $A(0, 0)$, $B(12, 0)$, $C(12, 2)$, $D(6, 7)$ and $E(0, 5)$ are the vertices of the pentagon $ABCDE$, then its area in square units, is
 a) 58 b) 60 c) 61 d) 63
189. A flag is standing vertically on a tower of height b . On a point at a distance a from the foot of the tower, the flag and the tower subtend equal angles. The height of the flag is
 a) $b \cdot \frac{a^2 + b^2}{a^2 - b^2}$ b) $a \cdot \frac{a^2 - b^2}{a^2 + b^2}$ c) $b \cdot \frac{a^2 - b^2}{a^2 + b^2}$ d) $a \cdot \frac{a^2 + b^2}{a^2 - b^2}$
190. A kite is flying at an inclination of 60° with the horizontal. If the length of the thread is 120 m, then the height of the kite is
 a) $60\sqrt{3}$ m b) 60 m c) $\frac{60}{\sqrt{3}}$ m d) 120 m
191. $\frac{a \cos A + b \cos B + c \cos C}{a + b + c}$ is equal to
 a) $1/r$ b) r/R c) R/r d) $1/R$
192. AB is a vertical pole. The end A is on the level ground. C is the middle point of AB . P is a point on the level ground. The portion BC subtends an angle β at P . If $AP = n AB$, then $\tan \beta =$
 a) $\frac{n}{2n^2 + 1}$ b) $\frac{n}{n^2 - 1}$ c) $\frac{n}{n^2 + 1}$ d) None of these
193. If $P(3, 7)$ is a point on the line joining $A(1, 1)$ and $B(6, 16)$, then the harmonic conjugate Q of point P has the coordinates
 a) $(9, 29)$ b) $(-9, 29)$ c) $(9, -29)$ d) $(-9, -29)$
194. The angles of a triangle are in the ratio 3:5:10. Then, the ratio of the smallest side to the greatest side is
 a) $1: \sin 10^\circ$ b) $1: 2 \sin 10^\circ$ c) $1: \cos 10^\circ$ d) $1: 2 \cos 10^\circ$
195. In ΔABC , if $\begin{vmatrix} 1 & a & b \\ 1 & c & a \\ 1 & b & c \end{vmatrix} = 0$, then $\sin^2 A + \sin^2 B + \sin^2 C$ is equal to
 a) $\frac{4}{9}$ b) $\frac{9}{4}$ c) $3\sqrt{3}$ d) 1
196. From a station A due West of a tower the angle of elevation of the top of the tower is seen to be 45° . From a station B , 10 m from A and in the direction 45° South of East of angle of elevation is 30° , the height of tower is
 a) $5\sqrt{2}(\sqrt{5} + 1)\text{m}$ b) $\frac{5(\sqrt{5} + 1)}{2}\text{m}$ c) $\frac{5\sqrt{2}(\sqrt{5} + 1)}{2}\text{m}$ d) None of these
197. A straight line with negative slope passing through the point $(1, 4)$ meets the coordinate axes at A and B . The minimum value of $OA + OB$ is equal to
 a) 5 b) 6 c) 9 d) 8

198. An observer finds that the elevation of the top of a tower is $22\frac{1}{2}^\circ$ and after walking 150 metres towards the foot of the tower he finds that the elevation of the top has increased to $67\frac{1}{2}^\circ$. The height of the tower in metres is
 a) 50 b) 75 c) 125 d) 175
199. In an isosceles ΔABC , $AB = AC$. If vertical angle A is 20° , then $a^3 + b^3$ is equal to
 a) $3a^2b$ b) $3b^2c$ c) $3c^2a$ d) abc
200. In a ΔABC , $a(\cos^2 B + \cos^2 C) + \cos A (c \cos C + b \cos B)$ is equal to
 a) a b) b c) c d) $a + b + c$
201. ABC Is a triangle with vertices $A(-1, 4)$, $B(6, -2)$ and $C(-2, 4)$. D, E And F are the points which divide each AB, BC and CA respectively in the ratio 3:1 internally. Then, the centroid of the triangle DEF is
 a) (3, 6) b) (1, 2) c) (4, 8) d) (-3, 6)
202. If in a ΔABC , $\frac{1}{a+c} + \frac{1}{b+c} = \frac{3}{a+b+c}$, then the value of the angle C is
 a) 60° b) 30° c) 45° d) None of these
203. A tower subtends an angle of 30° at a point distance d from the foot of the tower and on the same level as the foot of the tower. At a second point, h vertically above the first, the angle of depression of the foot of the tower is 60° . The height of the tower is
 a) $\frac{h}{3}$ b) $\frac{h}{3d}$ c) $3h$ d) $\frac{3h}{d}$
204. Points D, E are taken on the side BC of the ΔABC , such that $BD = DE = EC$. If $\angle BAD = x, \angle DAE = y, \angle EAC = z$, then the value of $\frac{\sin(x+y) \sin(y+z)}{\sin x \sin z}$ is equal to
 a) 1 b) 2 c) 4 d) None of these
205. $\sin A : \sin C = \sin(A - B) : \sin(B - C)$, then a^2, b^2, c^2 are in
 a) AP b) GP c) HP d) None of these
206. The point on the line $3x + 4y = 5$, which is equidistant from (1, 2) and (3, 4) is
 a) (7, -4) b) (15, -10) c) $(\frac{1}{7}, \frac{8}{7})$ d) $(0, \frac{5}{4})$
207. If A and B are two fixed points, then the locus of a point which moves in such a way that the angle, APB is a right angle is
 a) A circle b) An ellipse c) A parabola d) None of these
208. If in a ΔABC , $a \cos^2(\frac{C}{2}) + c \cos^2(\frac{A}{2}) = \frac{3b}{2}$, then the sides a, b and c
 a) Are in AP b) Are in GP c) Are in HP d) Satisfy $a + b = c$
209. For a regular polygon, let r and R be the radii of the inscribed and the circumscribed circles. A false statement among the following is
 a) There is regular polygon with $\frac{r}{R} = \frac{1}{2}$ b) There is a regular polygon $\frac{r}{R} = \frac{1}{\sqrt{2}}$
 c) There is a regular polygon with $\frac{r}{R} = \frac{2}{3}$ d) There is a regular polygon with $\frac{r}{R} = \frac{\sqrt{3}}{2}$
210. In a triangle vertex angles are A, B, C and side BC are given. The area of ΔABC is
 a) $\frac{s(s-a)(s-b)(s-c)}{2}$ b) $\frac{b^2 \sin C \sin A}{\sin B}$
 c) $ab \sin C$ d) $\frac{1}{2} \cdot \frac{a^2 \sin B \sin C}{\sin A}$
211. A flag staff in the centre of a rectangular field whose diagonal is 1200 m and subtends angle 15° and 45° at the mid point of the sides of the field. The height of the flag staff is
 a) 200 m b) $300\sqrt{2 + \sqrt{3}}$ m c) $300\sqrt{2 - \sqrt{3}}$ m d) 400 m
212. On the level ground the angle of elevation of the top of a tower is 30° . On moving 20 m nearer the tower, the angle of elevation is found to be 60° . The height of the tower is
 a) 10 m b) 20 m c) $10\sqrt{3}$ m d) None of these

- a) (1, 6) b) $(-\frac{1}{2}, 5)$ c) $(\frac{5}{6}, 6)$ d) $(-8, \frac{1}{8})$
227. Point $(\frac{1}{2}, -\frac{13}{4})$ divides the line joining the points (3, -5) and (-7, 2) in the ratio of
a) 1: 3 internally b) 3: 1 internally c) 1: 3 externally d) 3: 1 externally
228. The orthocenter of the triangle with vertices $O(0, 0), A(0, \frac{3}{2}), B(-5, 0)$ is
a) $(\frac{5}{2}, \frac{3}{4})$ b) $(-\frac{5}{2}, \frac{3}{4})$ c) $(-5, \frac{3}{2})$ d) (0, 0)
229. Area of triangle formed by the lines $x + y = 3$ and angle bisectors of the pair of straight lines $x^2 - y^2 + 2y = 1$ is
a) 2 sq units b) 4 sq units c) 6 sq units d) 8 sq units
230. $ABCD$ is a rectangular field. A vertical lamp post of height 12m stands at the corner A . If the angle of elevation of its top from B is 60° and from C is 45° , then the area of the field is
a) $48\sqrt{2}$ sq m b) $48\sqrt{3}$ sq m c) 48 sq m d) $12\sqrt{2}$ sq m
231. If two adjacent sides of a cylinder quadrilateral are 2 and 5 and the angle between them is 60° . If the third side is 3, then the remaining fourth side is
a) 2 b) 3 c) 4 d) 5
232. In ΔABC , $2R^2 \sin A \sin B \sin C$ is equal to
a) s^2 b) $ab + bc + ca$ c) Δ d) None of these
233. If in a ΔABC , the altitudes from the vertices A, B, C on opposite sides are in HP, then $\sin A, \sin B, \sin C$ are in
a) HP b) Arithmetico-Geometric Progression
c) AP d) GP
234. The middle point of the line segment joining (3, -1) and (1, 1) is shifted by two units (in the sense of increasing y) perpendicular to the line segment. Then the coordinates of the point in the new position are
a) $(2 - \sqrt{2}, 2)$ b) $(2, 2 - \sqrt{3})$ c) $(2 + \sqrt{2}, \sqrt{3})$ d) None of these
235. If area of triangle with vertices (0, 0), (0, 6) and (α, β) is 15 sq unit, then
a) $\alpha = \pm 5, \beta = 5$ b) $\alpha = \pm 10, \beta = 5$
c) $\alpha = \pm 5, \beta = 2$ d) $\alpha = \pm 5, \beta$ can take any real value
236. Area of quadrilateral whose vertices are (2, 3), (3, 4), (4, 5) and (5, 6) is equal to
a) 0 b) 4 c) 6 d) None of the above
237. If the distance between (2, 3) and (-5, 2) is equal to the distance between $(x, 2)$ and (1, 3), then the values of x are
a) -6, 8 b) 6, 8 c) -8, 6 d) -7, 7
238. A circle is inscribed in an equilateral triangle of side a . The area of the circle is
a) $3\pi a^2$ sq units b) $2a^2$ sq units c) a^2 sq units d) None of these
239. The x -axis, y -axis and a line passing through the point $A(6, 0)$ form a triangle ABC . If $\angle A = 30^\circ$ then the area of the triangle, in sq units is
a) $6\sqrt{3}$ b) $12\sqrt{3}$ c) $4\sqrt{3}$ d) $8\sqrt{3}$
240. In a ΔABC , if $\frac{\cos A}{a} = \frac{\cos B}{b} = \frac{\cos C}{c}$ and the side $a = 2$, then area of the triangle is
a) 1 sq unit b) 2 sq unit c) $\frac{\sqrt{3}}{2}$ sq unit d) $\sqrt{3}$ sq unit
241. In an ambiguous case of solving a triangle when $a = \sqrt{5}, b = 2, \angle A = \frac{\pi}{6}$, the two possible values of third side are c_1 and c_2 , then
a) $|c_1 - c_2| = 2\sqrt{6}$ b) $|c_1 - c_2| = 4\sqrt{6}$ c) $|c_1 - c_2| = 4$ d) $|c_1 - c_2| = 6$
242. In ΔABC , $(a - b)^2 \cos^2 \frac{C}{2} + (a + b)^2 \sin^2 \frac{C}{2}$ is equal to
a) a^2 b) b^2 c) c^2 d) None of these
243. In ΔABC , with usual notation, observe the two statements given below
I. $rr_1r_2r_3 = \Delta^2$

II. $r_1 r_2 + r_2 r_3 + r_3 r_1 = s^2$

Which of the following is correct?

- a) Both I and III are true b) I is true, II is false c) I is false, II is true d) Both I and II are false

244. If the angles of a triangle be in the ratio 1:2:7, then the ratio of its greatest side to the least side is
 a) 1:2 b) 2:1 c) $(\sqrt{5} + 1) : (\sqrt{5} - 1)$ d) $(\sqrt{5} - 1) : (\sqrt{5} + 1)$
245. From the top of a cliff 300 metres high, the top of a tower was observed at an angle of depression 30° and from the foot of the tower the top of the cliff was observed at an angle of elevation 45° , the height of the tower is
 a) $50(3 - \sqrt{3})$ m b) $200(3 - \sqrt{3})$ m c) $100(3 - \sqrt{3})$ m d) None of these
246. If x_1, x_2, x_3 and y_1, y_2, y_3 are both in GP with the same common ratio, then the points $(x_1, y_1), (x_2, y_2), (x_3, y_3)$
 a) Lie on a straight line b) Lie on an ellipse
 c) One vertices of a triangle d) Lie on a circle
247. The coordinates axes are rotated through an angle 135° . If the coordinates of a point P in the new system are known to be $(4, -3)$, then the coordinates of P in the original system are
 a) $(\frac{1}{\sqrt{2}}, \frac{7}{\sqrt{2}})$ b) $(\frac{1}{\sqrt{2}}, -\frac{7}{\sqrt{2}})$ c) $(-\frac{1}{\sqrt{2}}, -\frac{7}{\sqrt{2}})$ d) $(-\frac{1}{\sqrt{2}}, \frac{7}{\sqrt{2}})$
248. The coordinate axes rotated through an angle 135° . If the coordinates of a point P in the new system are known to be $(4, -3)$, then the coordinates of P in the original system are
 a) $(\frac{1}{\sqrt{2}}, \frac{7}{\sqrt{2}})$ b) $(\frac{1}{\sqrt{2}}, -\frac{7}{\sqrt{2}})$ c) $(-\frac{1}{\sqrt{2}}, -\frac{7}{\sqrt{2}})$ d) $(-\frac{1}{\sqrt{2}}, \frac{7}{\sqrt{2}})$
249. When the elevation of sun changes from 45° to 30° the shadow of a tower increases by 60 m the height of the tower is
 a) $30\sqrt{3}$ m b) $30(\sqrt{2} + 1)$ m c) $30(\sqrt{3} - 1)$ m d) $30(\sqrt{3} + 1)$ m
250. The angle of elevation of the top of a tower from a point A $\tan^{-1} 6$ and that from B due West of it, is $\tan^{-1} 7.5$. If h is the height of the tower, then $AB = \lambda h$, where λ^2 is equal to
 a) $\frac{21}{700}$ b) $\frac{42}{1300}$ c) $\frac{41}{900}$ d) None of these
251. In a triangle, if $r_1 + r_3 = k \cos^2 \frac{B}{2}$, then k is equal to
 a) R b) $2R$ c) $3R$ d) $4R$
252. AB is a vertical pole and C is its middle point. The end A is on the level ground and P is any point on the level ground other than A the portion CB subtends an angle β at P . If $AP : AB = 2 : 1$, then $\beta =$
 a) $\tan^{-1} \frac{4}{9}$ b) $\tan^{-1} \frac{1}{9}$ c) $\tan^{-1} \frac{5}{9}$ d) $\tan^{-1} \frac{2}{9}$
253. If the points $(x + 1, 2), (1, x + 2), (\frac{1}{x+1}, \frac{2}{x+1})$ are collinear, then x is
 a) 4 b) 5 c) -4 d) None of these
254. If in a ΔABC , CD is the angular bisector of the $\angle ACB$, then CD is equal to
 a) $\frac{a+b}{2ab} \cos \frac{C}{2}$ b) $\frac{a+b}{ab} \cos \frac{C}{2}$ c) $\frac{2ab}{a+b} \cos \frac{C}{2}$ d) None of these
255. A tower stands at the center of a circuit park. A and B are two points on the boundary of the park such that $AB (= a)$ subtends an angles of 60° at the foot of the tower and the angle of elevation of the top of the tower from A or B is 30° . The height of the tower is
 a) $\frac{2a}{\sqrt{3}}$ b) $2a\sqrt{3}$ c) $\frac{a}{\sqrt{3}}$ d) $\sqrt{3}$
256. Consider three points
 $P = (-\sin(\beta - \alpha), -\cos \beta)$
 $Q = (\cos(\beta - \alpha), \sin \beta)$
 And $R = (\cos(\beta - \alpha + \theta), \sin(\beta - \theta))$,
 Where $0 < \alpha, \beta, \theta < \frac{\pi}{4}$. Then,

- a) P lies on the line segment RQ b) Q lies on the line segment PR
 c) R lies on the line segment QP d) P, Q, R are non-collinear
257. In ΔABC , if $8R^2 = a^2 + b^2 + c^2$, then the triangle is
 a) Right angled b) Equilateral c) Acute angled d) Obtuse angled
258. A triangle with vertices $(4, 0), (-1, -1), (3, 5)$ is
 a) Isosceles and right angled b) Isosceles but not right angled
 c) Right angled but not isosceles d) Neither right angled nor isosceles
259. In a ΔABC , the correct formulae among the following are
 III. $r = 4R \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}$
 IV. $r_1 = (s - a) \tan \frac{A}{2}$
 V. $r_3 = \frac{\Delta}{s - c}$
 a) Only I, II b) Only II, III c) Only I, III d) I, II, III
260. The area (in square unit) of the triangle formed by the lines $x = 0, y = 0$ and $3x + 4y = 12$, is
 a) 3 b) 4 c) 6 d) 12
261. Three distinct points A, B and C given in the two dimensional coordinate plane such that the ratio of the distance of any one of them from the point $(1, 0)$ to the distance from the point $(-1, 0)$ is equal to $\frac{1}{3}$. Then, the circumcentre of the triangle ABC is at the point
 a) $(\frac{5}{4}, 0)$ b) $(\frac{5}{2}, 0)$ c) $(\frac{5}{3}, 0)$ d) $(0, 0)$
262. One possible condition for the three points $(a, b), (b, a)$ and $(a^2, -b^2)$ to be collinear, is
 a) $a - b = 2$ b) $a + b = 2$ c) $a = 1 + b$ d) $a = 1 - b$
263. Three vertical towers standing at A, B, C subtends the angle $\theta_A, \theta_B, \theta_C$ respectively at the circumcentre of the ΔABC , then $\tan \theta_A, \tan \theta_B$ and $\tan \theta_C$ are in
 a) AP b) GP c) HP d) None of these
264. The area of the triangle whose sides are $6, 5, \sqrt{13}$ (in square unit) is
 a) $5\sqrt{2}$ b) 9 c) $6\sqrt{2}$ d) 11
265. If points $A(x_1, y_1), B(x_2, y_2)$ and $C(x_3, y_3)$ are such that x_1, x_2, x_3 and y_1, y_2, y_3 are in AP, then
 a) A, B and C are concyclic points b) A, B and C are collinear points
 c) A, B and C are vertices of an equilateral triangle d) None of the above
266. Two points $P(a, 0)$ and $Q(-a, 0)$ are given, R is a variable point on one side of the line PQ such that $\angle RPQ - \angle RQP$ is 2α , then
 a) Locus of R is $x^2 - y^2 + 2xy \cot 2\alpha - a^2 = 0$ b) Locus of R is $x^2 + y^2 + 2xy \cot \alpha - a^2 = 0$
 c) Locus of R is a hyperbola, if $\alpha = \frac{\pi}{4}$ d) Locus of R is a circle, if $\alpha = \frac{\pi}{4}$
267. In ΔABC , medians, AD and BE are drawn. If $AD = 4, \angle DAB = \frac{\pi}{6}$ and $\angle ABE = \frac{\pi}{3}$, then the area of the ΔABC is
 a) $\frac{8}{3}$ sq units b) $\frac{16}{3}$ sq units c) $\frac{32}{3\sqrt{3}}$ sq units d) $\frac{64}{3}$ sq units
268. Point Q is symmetric to $P(4, -1)$ with respect to the bisector of the first quadrant. The length of PQ is
 a) $3\sqrt{2}$ b) $5\sqrt{2}$ c) $7\sqrt{2}$ d) $9\sqrt{2}$
269. The area of triangle formed by the points $(a, b + c), (b, c + a), (c, a + b)$ is equal to
 a) abc b) $a^2 + b^2 + c^2$ c) $ab + bc + ca$ d) 0
270. Let $0 \leq \theta \leq \frac{\pi}{2}$ and $x = X \cos \theta + Y \sin \theta, y = X \sin \theta - Y \cos \theta$ such that $x^2 + 4xy + y^2 = aX^2 + bY^2$, where a, b are constants, then
 a) $a = -1, b = 3, \theta = \frac{\pi}{4}$ b) $a = 1, b = -3, \theta = \frac{\pi}{3}$ c) $a = 3, b = -1, \theta = \frac{\pi}{4}$ d) $a = 3, b = -1, \theta = \frac{\pi}{3}$
271. If a point $P(4, 3)$ is shifted by a distance $\sqrt{2}$ unit parallel to the line $y = x$, then coordinates of p in new position are

- a) (5, 4) b) $(5 + \sqrt{2}, 4 + \sqrt{2})$ c) $(5 - \sqrt{2}, 4 - \sqrt{2})$ d) None of these
272. From the top of a cliff 50m high, the angles of depression of the top and bottom of a tower are observed to be 30° and 45° . The height of tower is
a) 50m b) $50\sqrt{3}$ m c) $50(\sqrt{3} - 1)$ m d) $50\left(1 - \frac{\sqrt{3}}{3}\right)$ m
273. The vertex of an equilateral triangle is $(2, -1)$ and the equation of its base is $x + 2y = 1$, the length of its sides is
a) $\frac{2}{\sqrt{15}}$ b) $\frac{4}{3\sqrt{3}}$ c) $\frac{1}{\sqrt{5}}$ d) $\frac{4}{\sqrt{15}}$
274. If the elevation of the sun is 30° , then the length of the shadow cast by a tower of 150 ft. height is
a) $75\sqrt{3}$ ft. b) $200\sqrt{3}$ ft. c) $150\sqrt{3}$ ft. d) None of these
275. The area of the triangle formed by the points $(2, 2)$, $(5, 5)$, $(6, 7)$ is equal to (in square unit)
a) $\frac{9}{2}$ b) 5 c) 10 d) $\frac{3}{2}$
276. The orthocenter of the ΔOAB , where O is the origin, $A(6, 0)$ and $B(3, 3\sqrt{3})$ is
a) $(9/2, \sqrt{3}/2)$ b) $(3, \sqrt{3})$ c) $(\sqrt{3}, 3)$ d) $(3, -\sqrt{3})$
277. In a ΔABC , $\cos A + \cos B + \cos C$ is equal to
a) $1 + \frac{r}{R}$ b) $1 - \frac{r}{R}$ c) $1 - \frac{R}{r}$ d) $1 + \frac{R}{r}$
278. If in a ΔPQR , $\sin P, \sin Q, \sin R$ are in AP, then
a) The altitudes are in AP b) The altitudes are in HP
c) The medians are in GP d) The medians are in AP
279. The angle of elevation of an object on a hill from a point on the ground is 30° . After walking 120 metres the elevation of the object is 60° . The height of the hill is
a) 120 m b) $60\sqrt{3}$ m c) $120\sqrt{3}$ m d) 60 m
280. If the median AD of ΔABC , makes an angle θ with side AB , then $\sin(A - \theta)$ is equal to
a) $\left(\frac{b}{c}\right) \operatorname{cosec} \theta$ b) $\left(\frac{b}{c}\right) \sin \theta$ c) $\left(\frac{c}{b}\right) \sin \theta$ d) $\left(\frac{c}{b}\right) \operatorname{cosec} \theta$
281. The angle of elevation of a cliff at a point A on the ground and a point B , 100 m vertically at A are α and β respectively. The height of the cliff is
a) $\frac{100 \cot \alpha}{\cot \alpha - \cot \beta}$ b) $\frac{100 \cot \beta}{\cot \alpha - \cot \beta}$ c) $\frac{100 \cot \beta}{\cot \beta - \cot \alpha}$ d) $\frac{100 \cot \beta}{\cot \beta + \cot \alpha}$
282. A quadrilateral $ABCD$ in which $AB = a, BC = b, CD = c$ and $DA = d$ is such that one circle can be inscribed in it and another circle can be circumscribed about it, then $\cos A$ is equal to
a) $\frac{ad + bc}{ad - dc}$ b) $\frac{ad - bc}{ad + bc}$ c) $\frac{ac + bd}{ac - bd}$ d) $\frac{ac - bd}{ac + bd}$
283. If λ be the perimeter of the ΔABC , then $b \cos^2\left(\frac{C}{2}\right) + c \cos^2\left(\frac{B}{2}\right)$ is equal to
a) λ b) 2λ c) $\frac{\lambda}{2}$ d) None of these
284. If in the ΔABC , $\angle B = 45^\circ$, then $a^4 + b^4 + c^4$ is equal to
a) $2a^2(b^2 + c^2)$ b) $2c^2(a^2 + b^2)$
c) $2b^2(a^2 + c^2)$ d) $2(a^2b^2 + b^2c^2 + c^2a^2)$
285. The ratio in which the line $x + y = 4$ divides the line joining the points $(1, -1)$ and $(5, 7)$ is
a) 1:2 b) 2:1 c) 1:3 d) 3:1
286. The coordinates of the orthocenter of the triangle formed by $(0, 0)$, $(8, 0)$, $(4, 6)$ is
a) $(4, 0)$ b) $(6, 3)$ c) $(6, 0)$ d) None of these
287. If $\Delta = a^2 - (b - c)^2$, where Δ is the area of ΔABC , then $\tan A$ is equal to
a) $\frac{15}{16}$ b) $\frac{8}{15}$ c) $\frac{8}{17}$ d) $\frac{1}{2}$

288. In ΔABC , $\frac{b+c}{a}$ is equal to
- a) $\frac{\cos \frac{1}{2}(B-C)}{\sin \frac{1}{2} A}$ b) $\frac{\sin \frac{1}{2}(B-C)}{\cos \frac{1}{2} A}$ c) $\frac{\cos \frac{1}{2}(B+C)}{\sin \frac{1}{2} A}$ d) $\frac{\cos \frac{1}{2}(B+C)}{\cos \frac{1}{2} A}$
289. In a cubical hall $ABCDPQRS$ with each side 10 m, G is the centre of the wall $BCRQ$ and T is the mid point of the side AB . The angle of elevation of G at the point T is
- a) $\sin^{-1}(1/\sqrt{3})$ b) $\cos^{-1}(1/\sqrt{3})$ c) $\cot^{-1}(1/\sqrt{3})$ d) None of these
290. If p_1, p_2, p_3 are respectively the perpendicular from the vertices of a triangle to the opposite sides, then p_1, p_2, p_3 is equal to
- a) $a^2 b^2 c^2$ b) $2a^2 b^2 c^2$ c) $\frac{4a^2 b^2 c^2}{R^2}$ d) $\frac{a^2 b^2 c^2}{8R^2}$
291. An observer standing on a 300 m high tower observes two boats in the same direction their angles of depression are 60° and 30° respectively. The distance between boats is
- a) 173.2 m b) 346.4 m c) 25 m d) 72 m
292. If the length of the sides of a triangle are 3, 4 and 5 unit, then R is
- a) 3.5 b) 3.0 c) 2.0 d) 2.5
293. In a triangle $\left(1 - \frac{r_1}{r_2}\right)\left(1 - \frac{r_1}{r_3}\right) = 2$, then the triangle is
- a) Right angled b) Isosceles c) Equilateral d) None of these
294. If in a ΔABC , $a \tan A + b \tan B = (a+b) \tan \frac{A+B}{2}$, then
- a) $A = B$ b) $A = -B$ c) $A = 2B$ d) $B = 2A$
295. The coordinates of the circumcentre of the triangle with vertices $(8, 6)$, $(8, -2)$ and $(2, -2)$ are
- a) $\left(6, \frac{2}{3}\right)$ b) $(8, 2)$ c) $(5, -2)$ d) $(5, 2)$
296. Points D, E are taken on the side BC of a ΔABC such that $BD = DE = EC$. If $\angle BAD = x$, $\angle DAE = y$, $\angle EAC = z$, then the value of $\frac{\sin(x+y) \sin(y+z)}{\sin x \sin z}$ is equal to
- a) 1 b) 2 c) 4 d) None of these
297. An observer on the top of tree, finds the angle of depression of a car moving towards the tree to be 30° . After 3 min this angle becomes 60° . After how much more time, the car will reach the tree?
- a) 4 min b) 4.5 min c) 1.5 min d) 2 min
298. If PQ be a vertical tower subtending angles α, β and γ at the points A, B and C respectively on the line in the horizontal plane through the foot D of tower and on the same side of it, then $BC \cot \alpha - CA \cot \alpha + AB \cot \gamma$ is equal to
- a) 0 b) 1 c) 2 d) None of these
299. The equation of the three sides of a triangle are $x = 2, y + 1 = 0$ and $x + 2y = 4$. The coordinates of the circumcentre of the triangles are
- a) $(4, 0)$ b) $(2, -1)$ c) $(0, 4)$ d) $(-1, 2)$
300. On one bank of river there is a tree. On another bank, an observer makes an angle of elevation of 60° at the top of the tree. The angle of elevation of the top of the tree at a distance 20 m away the bank is 30° . The width of the river is
- a) 20 m b) 10 m c) 5 m d) 1 m
301. Consider the following statements :
- In ΔABC , $a = \sqrt{3} + 1, \angle B = 30^\circ, \angle C = 45^\circ$, then c is equal to
 - In a triangle, if $a^2 + b^2 + c^2 = 8R^2$, then the triangle is right angled
 - In a ΔABC , $a = 2, b = 3, c = 4$, then $\cos A = \frac{7}{8}$
- Which of the statements given above is/are correct?
- a) Only (1) b) Only (2) c) Only (3) d) All of these
302. If A is the area and $2s$ the sum of three sides of a triangle, then

- a) $A \leq \frac{s^2}{3\sqrt{3}}$ b) $A \leq \frac{s^2}{2}$ c) $A > \frac{s^2}{\sqrt{3}}$ d) None of these

303. A flag staff of 5 m high stands on a building of 25 m high. At an observer at a height of 30 m, the flag staff and the building subtend equal angles. the distance of the observer from the top of the flag staff is

- a) $\frac{5\sqrt{3}}{2}$ m b) $5\sqrt{\frac{3}{2}}$ m c) $5\sqrt{\frac{2}{3}}$ m d) None of these

304. A tower of height b subtends an angle at a point O on the level of the foot of the tower and at a distance ' a ' from the foot of the tower. If the pole mounted on the tower also subtends an equal angle at O , the height of the pole is

- a) $b\left(\frac{a^2 - b^2}{a^2 + b^2}\right)$ b) $b\left(\frac{a^2 + b^2}{a^2 - b^2}\right)$ c) $a\left(\frac{a^2 - b^2}{a^2 + b^2}\right)$ d) $a\left(\frac{a^2 + b^2}{a^2 - b^2}\right)$

305. If the angles of a triangle are in the ratio 4: 1: 1, then the ratio of the longest side to the perimeter is

- a) $\sqrt{3}:(2 + \sqrt{3})$ b) 1:6 c) 1:(2 + $\sqrt{3}$) d) 2:3

306. If two angles of a triangle are 45° and $\tan^{-1}(2)$, then the third angle is

- a) 60° b) 75° c) $\tan^{-1} 3$ d) 90°

307. The angle of elevation of top of a tower from a point on the ground is 30° and it is 60° when it is viewed from a point located 40m away the initial point towards the tower. The height of the tower is

- a) $-20\sqrt{3}$ m b) $\frac{\sqrt{3}}{20}$ m c) $-\frac{\sqrt{3}}{20}$ m d) $20\sqrt{3}$ m

308. The orthocentre of the triangle formed by the points $(0, 0)$, $(4, 0)$ and $(3, 4)$ is

- a) $(2, 0)$ b) $\left(\frac{3}{2}, 2\right)$ c) $\left(\frac{3}{4}, 3\right)$ d) $\left(3, \frac{3}{4}\right)$

309. In order to remove xy -term from the equation $5x^2 + 4\sqrt{3}xy + 9y^2 - 8 = 0$ the coordinate axes must be rotated through an angle

- a) $\pi/6$ b) $\pi/4$ c) $\pi/3$ d) $\pi/2$

310. If C is a point on the line segment joining $A(-3, 4)$ and $B(2, 1)$ such that $AC = 2BC$, then the coordinate of C is

- a) $\left(\frac{1}{3}, 2\right)$ b) $\left(2, \frac{1}{3}\right)$ c) $(2, 7)$ d) $(7, 2)$

311. The image of the centre of the circle $x^2 + y^2 = a^2$ with respect to the mirror image $x + y = 1$, is

- a) $\left(\frac{1}{\sqrt{2}}, \sqrt{2}\right)$ b) $(\sqrt{2}, \sqrt{2})$ c) $(\sqrt{2}, 2\sqrt{2})$ d) None of these

312. In a ΔABC , $\operatorname{cosec} A(\sin B \cos C + \cos B \sin C)$ is equal to

- a) $\frac{c}{a}$ b) $\frac{a}{c}$ c) 1 d) $\frac{c}{ab}$

313. If $A(3, 5)$, $B(-5, -4)$, $C(7, 10)$ are the vertices of a parallelogram, taken in the order, then the coordinates of the fourth vertex are

- a) $(10, 19)$ b) $(15, 19)$ c) $(19, 10)$ d) $(19, 15)$

314. Two pillars of equal height stand on either side of a road-way which is 60 m wide. At a point in the road-way between the pillars, the elevation of the top of pillars are 60° and 30° . The height of the pillars is

- a) $15\sqrt{3}$ m b) $\frac{15}{\sqrt{3}}$ m c) 15 m d) 20 m

315. From the top of a cliff of height a , the angle of depression of the foot of a certain tower is found to be double the angle of elevation of the top of the tower of height h . If θ be the angle of elevation, then its value is

- a) $\cos^{-1} \sqrt{\frac{2h}{a}}$ b) $\sin^{-1} \sqrt{\frac{2h}{a}}$ c) $\sin^{-1} \sqrt{\frac{a}{2-h}}$ d) $\tan^{-1} \sqrt{3 - \frac{2h}{a}}$

316. The points $A(2a, 4a)$, $B(2a, 6a)$ and $C(2a + \sqrt{3}a, 5a)$, $a > 0$ are the vertices of



- a) An isosceles triangle
c) An acute angled triangle
- b) A right angled triangle
d) None of the above
317. If a vertex of a triangle is $(1, 1)$ and the mid points of two sides through the vertex are $(-1, 2)$ and $(3, 2)$, then the centroid of the triangle is
a) $(1, \frac{7}{3})$ b) $(\frac{1}{3}, \frac{7}{3})$ c) $(-\frac{1}{3}, \frac{7}{3})$ d) $(-1, \frac{7}{3})$
318. P, Q, R and S are the points on the line joining the points $P(a, x)$ and $T(b, y)$ such that $PQ = QR = RS = ST$, then $(\frac{5a+3b}{8}, \frac{5x+3y}{8})$ is the mid point of
a) PQ b) QR c) RS d) ST
319. Orthocenter of the triangle formed by the lines $x + y = 1$ and $xy = 0$ is
a) $(0, 0)$ b) $(0, 1)$ c) $(1, 0)$ d) $(-1, 1)$
320. The circumcentre of the triangle with vertices $(0, 30)$, $(4, 0)$ and $(30, 0)$ is
a) $(10, 10)$ b) $(10, 12)$ c) $(12, 12)$ d) $(17, 17)$
321. A vertical pole (more than 100 m high) consists of two portions, the lower being one third of the whole, if the upper portion subtends an angle $\tan^{-1} \frac{1}{2}$ at a point in a horizontal plane through the foot of the pole and distance 40 ft from it, then the height of the pole is
a) 100 ft b) 120 ft c) 150 ft d) None of these
322. If a point $P(4, 3)$ is rotated through an angle 45° in anti-clockwise direction about origin, then coordinates of P in new position are
a) $(\frac{1}{\sqrt{2}}, \frac{7}{\sqrt{2}})$ b) $(-\frac{7}{\sqrt{2}}, -\frac{1}{\sqrt{2}})$ c) $(-\frac{1}{\sqrt{2}}, \frac{7}{\sqrt{2}})$ d) $(\frac{1}{\sqrt{2}}, -\frac{7}{\sqrt{2}})$
323. The horizontal distance between two towers is 60m and the angle of depression of the top of the first tower as seen from the top of the second is 30° . If the height of the second tower be 150m, then the height of the first tower is
a) 90 m b) $(150 - 60\sqrt{3})$ m c) $(150 + 20\sqrt{3})$ m d) None of the above
324. In a ΔABC , if $\sin A + \sin B + \sin C)(\sin A + \sin B - \sin C) = 3 \sin A \sin B$, then the angle C is equal to
a) $\frac{\pi}{2}$ b) $\frac{\pi}{3}$ c) $\frac{\pi}{4}$ d) $\frac{\pi}{6}$
325. From the bottom of a pole of height h the angle of elevation of the top of a tower is α and the pole subtends an angle β at the top of the tower. The height of the tower is
a) $\frac{h \tan(\alpha - \beta)}{\tan(\alpha - \beta) - \tan \alpha}$ b) $\frac{h \cot(\alpha - \beta)}{\cot(\alpha - \beta) - \cot \alpha}$ c) $\frac{\cot(\alpha - \beta)}{\cot(\alpha - \beta) - \cot \alpha}$ d) None of these
326. A variable line through the point $(\frac{1}{5}, \frac{1}{5})$ cuts the coordinate axes in the points A and B . If the point P divides AB internally in the ratio 3:1, then the locus of P is
a) $3y + x = 20xy$ b) $y + 3x = 20xy$ c) $x + y = 20xy$ d) $3x + 3y = 20xy$
327. If the points $(1, 2)$ and $(3, 4)$ were to be on the same side of the line $3x - 5y + a = 0$, then
a) $1 < a < 6$ b) $7 < a < 11$ c) $a > 11$ d) $a < 7$ or $a > 11$
328. The area of the region bounded by the lines $y = |x - 2|$, $x = 1$, $x = 3$ and the x -axis is
a) 1 b) 2 c) 3 d) 4
329. In triangle ABC , the value of $\frac{\cot \frac{A}{2} \cot \frac{B}{2} - 1}{\cot \frac{A}{2} \cot \frac{B}{2}}$ is
a) $\frac{a}{a + b + c}$ b) $\frac{c}{a + b + c}$ c) $\frac{2a}{a + b + c}$ d) $\frac{2c}{a + b + c}$
330. If the coordinates of the centroid and a vertex of an equilateral triangle are $(1, 1)$ and $(1, 2)$ respectively, then the coordinates of another vertex, are
a) $(\frac{2 - \sqrt{3}}{2}, -\frac{1}{2})$ b) $(\frac{2 + 3\sqrt{3}}{2}, -\frac{1}{2})$ c) $(\frac{2 + \sqrt{3}}{2}, \frac{1}{2})$ d) None of these
331. In any triangle ABC , $c^2 \sin 2B + b^2 \sin 2C$ is equal to

- a) $\frac{\Delta}{2}$ b) Δ c) 2Δ d) 4Δ

332. A house of height 100 m subtends a right angle at the window of an opposite house. If the height of the window be 64 m, then the distance between the two houses is
 a) 48 m b) 36 m c) 54 m d) 72 m
333. The angle of elevation of the top of a vertical pole when observed from each vertex of a regular hexagon is $\frac{\pi}{3}$. If the area of the circle circumscribing the hexagon be A metre², then the area of the hexagon is
 a) $\frac{3\sqrt{3}A}{8}$ m² b) $\frac{\sqrt{3}A}{\pi}$ m² c) $\frac{3\sqrt{3}A}{4\pi}$ m² d) $\frac{3\sqrt{3}A}{2\pi}$ m²
334. The area of the segment of a circle of radius a subtending an angle of 2α at the centre is
 a) $a^2\left(\alpha + \frac{1}{2}\sin 2\alpha\right)$ b) $\frac{1}{2}a^2\sin 2\alpha$ c) $a^2\left(\alpha + \frac{1}{2}\sin 2\alpha\right)$ d) $a^2\alpha$
335. A man of height 6 ft. observes the top of a tower and the foot of the tower at angles of 45° and 30° of elevation and depression respectively. The height of the tower is
 a) 13.79 m b) 14.59 m c) 14.29 m d) None of these
336. If the sides of the triangles are $5k, 6k, 5k$ and radius of incircle is 6, the value of k is equal to
 a) 4 b) 5 c) 6 d) 7
337. At the foot of the mountain the elevation of its summit is 45° , after ascending 100 m towards the mountain up a slope of 30° inclination, the elevation is found to be 60° . The height of the mountain is
 a) $\frac{\sqrt{3}+1}{2}$ m b) $\frac{\sqrt{3}-1}{2}$ m c) $\frac{\sqrt{3}+1}{2\sqrt{3}}$ m d) None of these
338. At each end of a horizontal line of length $2a$, the angular elevation of the peak of a vertical tower is θ and that at its middle point it is ϕ . The height of the peak is
 a) $a \sin \theta \sin \phi$ b) $\frac{a \sin \theta \sin \phi}{\sqrt{\sin(\theta + \phi) \sin(\phi - \theta)}}$
 c) $\frac{a \cos \theta \cos \phi}{\sqrt{\cos(\phi + \theta) \cos(\phi - \theta)}}$ d) None of the above
339. In a triangle ABC , $\frac{\cos A}{a} = \frac{\cos B}{b} = \frac{\cos C}{c}$, if $a = \frac{1}{\sqrt{6}}$, then the area of the triangle (in square unit) is
 a) $1/24$ b) $\sqrt{3}/24$ c) $\frac{1}{8}$ d) $\frac{1}{\sqrt{3}}$
340. A ladder leans again a wall at an angle α to the horizontal. Its foot is pulled away through a distance a_1 so that it slides a distance b_1 down the wall and rests inclined at angle β with the horizontal. Its foot is further pulled away through a_2 , so that it slides a further distance b_2 down the wall and is now, inclined at an angle γ . If $a_1 a_2 = b_1 b_2$, then
 a) $\alpha + \beta + \gamma$ is greater than π b) $\alpha + \beta + \gamma$ is equal to π
 c) $\alpha + \beta + \gamma$ is less than π d) Nothing can be said about $\alpha + \beta + \gamma$
341. In a ΔABC , $\frac{(a+b+c)(b+c-a)(c+a-b)(a+b-c)}{4b^2c^2}$ equals
 a) $\cos^2 A$ b) $\cos^2 B$ c) $\sin^2 A$ d) $\sin^2 B$
342. The locus of a point P which moves such that $2PA = 3PB$, where $A(0, 0)$ and $B(4, -3)$ are points, is
 a) $5x^2 - 5y^2 - 72x + 54y + 225 = 0$ b) $5x^2 + 5y^2 - 72x + 54y + 225 = 0$
 c) $5x^2 + 5y^2 + 72x - 54y + 225 = 0$ d) $5x^2 + 5y^2 - 72x - 54y - 225 = 0$
343. The incentre of the triangle formed by $(0, 0)$, $(5, 12)$, $(16, 12)$ is
 a) $(7, 9)$ b) $(9, 7)$ c) $(-9, 7)$ d) $(-7, 9)$
344. In ΔABC , if $\tan \frac{A}{2} \tan \frac{C}{2} = \frac{1}{2}$, then a, b, c are in
 a) AP b) GP c) HP d) None of these
345. In ΔABC , if $\sin^2 \frac{A}{2}, \sin^2 \frac{B}{2}, \sin^2 \frac{C}{2}$ be in HP, then a, b, c will be in
 a) AP b) GP c) HP d) None of these
346. If O is the origin and $P(2,3)$ and $Q(4,5)$ are two points, then $OP \cdot OQ \cos \angle POQ =$

- a) 8 b) 15 c) 22 d) 23
347. If in a ΔABC , $2b^2 = a^2 + c^2$, then $\frac{\sin 3B}{\sin B}$ is equal to
a) $\frac{c^2 - a^2}{2ca}$ b) $\frac{c^2 - a^2}{ca}$ c) $\left(\frac{c^2 - a^2}{ca}\right)^2$ d) $\left(\frac{c^2 - a^2}{2ca}\right)^2$
348. A spherical balloon of radius r subtends an angle α at the eye of an observer. If the angle of elevation of the centre of the balloon be β , then height of the centre of the balloon is
a) $r \operatorname{cosec} \left(\frac{\alpha}{2}\right) \sin \beta$ b) $r \operatorname{cosec} \alpha \sin \left(\frac{\beta}{2}\right)$ c) $r \sin \left(\frac{\alpha}{2}\right) \operatorname{cosec} \beta$ d) $r \sin \alpha \operatorname{cosec} \left(\frac{\beta}{2}\right)$
349. A point $P(2, 4)$ translates to the point Q along the parallel to the positive direction of x -axis by 2 units. If O be the origin, then $\angle OPQ$ is
a) $\sin^{-1} \sqrt{\frac{399}{400}}$ b) $\cos^{-1} \left(\frac{1}{20}\right)$ c) $-\sin^{-1} \left(\sqrt{\frac{399}{400}}\right)$ d) None of these
350. From the top of a hill h meters high, the angles of depressions of the top and the bottom of a pillar are α and β respectively. The height (in meters) of the pillar is
a) $\frac{h(\tan \beta - \tan \alpha)}{\tan \beta}$ b) $\frac{h(\tan \alpha - \tan \beta)}{\tan \alpha}$ c) $\frac{h(\tan \beta + \tan \alpha)}{\tan \beta}$ d) $\frac{h(\tan \beta + \tan \alpha)}{\tan \alpha}$
351. If $G(1, 4)$ is the centroid of triangle ABC having its two vertices A and B at $(4, -3)$ and $(-9, 7)$ respectively, then area of the triangle ABC in square units, is
a) $\frac{138}{2}$ b) $\frac{319}{2}$ c) $\frac{183}{2}$ d) $\frac{381}{2}$
352. If $a = 2\sqrt{2}$, $b = 6$, $A = 45^\circ$, then
a) No triangle is possible b) One triangle is possible
c) Two triangles are possible d) Either no triangle or two triangles are possible
353. Two poles of equal height stand on either side of a 100 m wide road. At a point between the poles the angles of elevation of the tops of the poles are 30° and 60° . The height of each pole is
a) 25 m b) $25\sqrt{3}$ m c) $\frac{100}{\sqrt{3}}$ m d) None of these
354. Area (in sq unit) enclosed by $y = 1$, $2x + y = 2$ and $x + y = 2$ is
a) $\frac{1}{2}$ sq unit b) $\frac{1}{4}$ sq unit c) 1 sq unit d) 2 sq units
355. The feet of the perpendicular drawn from P to the sides of a ΔABC are collinear, then P is
a) Circumcentre of ΔABC b) Lies on the circumcircle of ΔABC
c) Excentre of ΔABC d) None of the above
356. Let $O(0, 0)$, $P(3, 4)$, $Q(6, 0)$ be the vertices of the triangle OPQ . The point R inside the triangle OPQ is such that the triangles OPR , PQR , OQR are of equal area. The coordinates of R are
a) $\left(\frac{4}{3}, 3\right)$ b) $\left(3, \frac{2}{3}\right)$ c) $\left(3, \frac{4}{3}\right)$ d) $\left(\frac{4}{3}, \frac{2}{3}\right)$
357. A tree is broken by wind, its upper part touches the ground at a point 10 metres from the foot of the tree and makes an angle of 45° with the ground. The entire length of the tree is
a) 15 metres b) 20 metres c) $10(1 + \sqrt{2})$ metres d) $10\left(1 + \frac{\sqrt{3}}{2}\right)$ metres
358. Without change of axes the origin is shifted to (h, k) , then from the equation $x^2 + y^2 - 4x + 6y - 7 = 0$ the terms containing linear powers are missing. The point (h, k) is
a) $(3, 2)$ b) $(-3, 2)$ c) $(2, -3)$ d) $(-2, -3)$
359. The horizontal distance between two towers is 60 m and the angle of depression of the top of the first tower as seen from the top of the second is 30° . If the height of the second tower be 150 m, then the height of the first tower is
a) $(150 - 60\sqrt{3})$ m b) 90 m c) $(150 - 20\sqrt{3})$ m d) None of these
360. Let $a > 0, b > 0$. The sum of the distance of the point (a, b) from the lines $\frac{x}{a} + \frac{y}{b} = 1$ and $\frac{x}{b} + \frac{y}{a} = 1$ is

- a) $\frac{a+b}{2}$ b) \sqrt{ab} c) $\frac{2ab}{a+b}$ d) $\sqrt{a^2+b^2}$
361. In ΔABC , if $\frac{1}{b+c} + \frac{1}{c+a} = \frac{3}{a+b+c}$, then C is equal to
a) 90° b) 60° c) 45° d) 30°
362. If in a ΔABC , $4 \sin A = 4 \sin B = 3 \sin C$, then $\cos C$ is equal to
a) $1/3$ b) $1/9$ c) $1/27$ d) $1/18$
363. In a ΔABC , $(b+c)(bc) \cos A + (a+c)(ac) \cos B + (a+b)(ab) \cos C$ is
a) $a^2 + b^2 + c^2$ b) $a^3 + b^3 + c^3$
c) $(a+b+c)(a^2 + b^2 + c^2)$ d) $(a+b+c)(ab + bc + ca)$
364. If a flag staff of 6 m high placed on the top of a tower throws a shadow of $2\sqrt{3}$ m along the ground, then the angle (in degrees) that the sun makes with the ground is
a) 60° b) 80° c) 75° d) None of these
365. If the angles of a ΔABC be in AP, then
a) $c^2 = a^2 + b^2 - ab$ b) $b^2 = a^2 + c^2 - ac$ c) $a^2 = b^2 + c^2 - ac$ d) $b^2 = a^2 + c^2$
366. The sides of a triangle are respectively 7 cm, $4\sqrt{3}$ cm and $\sqrt{13}$ cm, then the smallest angle of the triangle is
a) $\frac{\pi}{6}$ b) $\frac{\pi}{3}$ c) $\frac{\pi}{4}$ d) $\frac{\pi}{5}$
367. The straight lines $x + y = 0$, $3x + y - 4 = 0$ and $x + 3y - 4 = 0$ form a triangle which is
a) Right angled b) Equilateral c) Isosceles d) None of these
368. A tower of x metres height has flag staff at its top. The tower and the flag staff subtend equal angles at a point distant y metres from the foot of the tower. Then, the length of the flag staff in metres is
a) $y \left(\frac{x^2 - y^2}{x^2 + y^2} \right)$ b) $x \left(\frac{x^2 + y^2}{y^2 - x^2} \right)$ c) $x \left(\frac{x^2 + y^2}{x^2 - y^2} \right)$ d) $x \left(\frac{x^2 - y^2}{x^2 + y^2} \right)$
369. If $\begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix} = 0$, then the points (x_1, y_1) , (x_2, y_2) and (x_3, y_3) are
a) Vertices of an equilateral triangle b) Vertices of a right angled triangle
c) Vertices of an isosceles triangle d) None of the above
370. If two angles of ΔABC are 45° and 60° , then the ratio of the smallest and the greatest sides are
a) $(\sqrt{3} - 1) : 1$ b) $\sqrt{3} : \sqrt{2}$ c) $1 : \sqrt{3}$ d) $\sqrt{3} : 1$
371. An aeroplane flying horizontally 1 km above the ground is observed at an elevation of 60° and after 10 s the elevation is observed to be 30° . The uniform speed of the aeroplane (in km/h) is
a) 240 b) $240\sqrt{3}$ c) $60\sqrt{3}$ d) None of these
372. The angle of elevation of the top of a tower from the top and bottom of a building of height ' a ' are 30° and 45° respectively. If the tower and the building stand at the same level, the height of the tower is
a) $\frac{a(3 + \sqrt{3})}{2}$ b) $a(\sqrt{3} + 1)$ c) $a\sqrt{3}$ d) $a(\sqrt{3} - 1)$
373. If in a ΔABC , $2b^2 = a^2 + c^2$, then $\frac{\sin 3B}{\sin B}$ is equal to
a) $\frac{c^2 - a^2}{2ca}$ b) $\frac{c^2 - a^2}{ca}$ c) $\left(\frac{c^2 - a^2}{ca} \right)^2$ d) $\left(\frac{c^2 - a^2}{2ca} \right)^2$
374. The angle of elevation of a cloud from a point h mt. above is θ° and the angle of depression of its reflection in the lake is ϕ . Then, the height is
a) $\frac{h \sin(\phi - \theta)}{\sin(\phi + \theta)}$ b) $\frac{h \sin(\phi + \theta)}{\sin(\phi - \theta)}$ c) $\frac{h \sin(\theta + \phi)}{\sin(\theta - \phi)}$ d) None of these
375. In ΔABC , $\angle B = \frac{\pi}{3}$ and $\angle C = \frac{\pi}{4}$. If D divides BC internally in ratio 1:3, then the value of $\frac{\sin \angle BAD}{\sin \angle CAD}$ is
a) $\frac{1}{\sqrt{3}}$ b) $\frac{1}{\sqrt{6}}$ c) $\sqrt{\frac{2}{3}}$ d) $\frac{1}{3}$

376. Let equation of the side BC of a ΔABC be $x + y + 2 = 0$. If coordinates of its orthocentre and circumcentre are $(1, 1)$ and $(2, 0)$ respectively, then radius of the circumcircle of ΔABC is
 a) 3 b) $\sqrt{10}$ c) $2\sqrt{2}$ d) None of these
377. In a ΔABC , if $b + c = 2a$ and $\angle A = 60^\circ$, then ΔABC is
 a) Equilateral b) Right angled c) Isosceles d) Scalene
378. The orthocenter of the triangle with vertices $(-2, -6)$, $(-2, 4)$ and $(1, 3)$ is
 a) $(3, 1)$ b) $(1, 1/3)$ c) $(1, 3)$ d) None of these
379. If $a > 0, b > 0$ the maximum area of the triangle formed by the points $O(0, 0)$, $A(a \cos \theta, b \sin \theta)$ and $B(a \cos \theta, -b \sin \theta)$ is (in sq unit)
 a) $\frac{ab}{2}$ when $\theta = \frac{\pi}{4}$ b) $\frac{3ab}{2}$ when $\theta = \frac{\pi}{4}$ c) $\frac{ab}{2}$ when $\theta = -\frac{\pi}{4}$ d) $a^2 b^2$
380. If the coordinates of orthocentre O' and centroid G of a ΔABC are $(0, 1)$ and $(2, 3)$ respectively, then the coordinates of the circumcentre are
 a) $(3, 2)$ b) $(1, 0)$ c) $(4, 3)$ d) $(3, 4)$
381. The point P is equidistant from $A(1, 3)$, $B(-3, 5)$ and $C(5, -1)$, then PA is equal to
 a) 5 b) $5\sqrt{5}$ c) 25 d) $5\sqrt{10}$
382. The coordinates of the incentre of the triangle having sides $3x - 4y = 0$, $5x + 12y = 0$ and $y - 15 = 0$ are
 a) $-1, 8$ b) $1, -8$ c) $2, 6$ d) None of these
383. The mid point of the line joining the points $(-10, 8)$ and $(-6, 12)$ divides the line joining the points $(4, -2)$ and $(-2, 4)$ in the ratio
 a) 1: 2 internally b) 1: 2 externally c) 2: 1 internally d) 2: 1 externally
384. The centre of circle inscribed in square formed by the lines $x^2 - 8x + 12 = 0$ and $y^2 - 14y + 45 = 0$, is
 a) $(4, 7)$ b) $(7, 4)$ c) $(9, 4)$ d) $(4, 9)$
385. If A, A_1, A_2, A_3 be the areas of the incircle and excircles, then $\frac{1}{\sqrt{A_1}} + \frac{1}{\sqrt{A_2}} + \frac{1}{\sqrt{A_3}}$ is equal to
 a) $\frac{1}{\sqrt{A}}$ b) $\frac{2}{\sqrt{A}}$ c) $\frac{3}{\sqrt{A}}$ d) $\frac{4}{\sqrt{A}}$
386. If $x = X \cos \theta - Y \sin \theta$, $y = X \sin \theta + Y \cos \theta$ and $x^2 + 4xy + y^2 = AX^2 + BY^2$, $0 \leq \theta \leq \frac{\pi}{2}$, then
 a) $\theta = \frac{\pi}{6}$ b) $\theta = \frac{\pi}{4}$ c) $A = -6$ d) $B = 1$
387. If $P(1, 0)$, $Q = (-1, 0)$ and $R = (2, 0)$ are three given points, then the locus of a point S satisfying the relation $SQ^2 + SR^2 = 2SP^2$ is
 a) A straight line parallel to x -axis b) A circle through origin
 c) A circle with centre at the origin d) A straight line parallel to y -axis
388. The triangle formed by $x^2 - 3y^2 = 0$ and $x = 4$ is
 a) Isosceles b) Equilateral c) Right angled d) None of these
389. The angle of elevation of the top of an incomplete vertical pillar at a horizontal distance of 100 m from its base is 45° . If the angle of elevation of the top of the complete pillar at the same point is to be 60° , then the height of the incomplete pillar is to be increased by
 a) $50\sqrt{2}$ m b) 100 m c) $100(\sqrt{3} - 1)$ m d) $100(\sqrt{3} + 1)$ m
390. If $O(0, 0)$, $A(4, 0)$ and $B(0, 3)$ are the vertices of a triangle OAB , then the coordinates of the excentre opposite to the vertex $O(0, 0)$ are
 a) $(12, 12)$ b) $(6, 6)$ c) $(3, 3)$ d) None of these
391. A tower subtends an angle α at a point A in the plane of its base and angle of depression of the foot of the tower at a point l metres just above A is β . The height of the tower is
 a) $l \tan \beta \cot \alpha$ b) $l \tan \alpha \cot \beta$ c) $l \tan \alpha \tan \beta$ d) $l \cot \alpha \cot \beta$
392. Observe the following statements

I. In ΔABC $b \cos^2 \frac{C}{2} + c \cos^2 \frac{B}{2} = s$

II. ΔABC , $\cot \frac{A}{2} = \frac{b+c}{2} \Rightarrow B = 90^\circ$

Which of the following is correct?

- a) Both I and II are true
- b) I is true, II is false
- c) I is false, II is false
- d) Both I and II are false

393. If A and B are two points on one bank of a straight river and C, D are two other points on the other bank of river. If direction from A to B is same as that from C to D and $AB = a, \angle CAD = \alpha, \angle DAB = \beta, \angle CBA = \gamma$, then CD is equal to

- a) $\frac{a \sin \beta \sin \gamma}{\sin \alpha \sin(\alpha + \beta + \gamma)}$
- b) $\frac{a \sin \alpha \sin \gamma}{\sin \beta \sin(\alpha + \beta + \gamma)}$
- c) $\frac{a \sin \alpha \sin \beta}{\sin \gamma \sin(\alpha + \beta + \gamma)}$
- d) None of these

394. $ABCD$ is a square plot. The angle of elevation of the top of a pole standing at D from A or C is 30° and that from B is θ , then $\tan \theta$ is equal to

- a) $\sqrt{6}$
- b) $1/\sqrt{6}$
- c) $\sqrt{3}/2$
- d) $\sqrt{2/3}$


395. In a ΔABC , if $(\sqrt{3} - 1)a = 2b, A = 3B$, then $\angle C$ is

- a) 60°
- b) 120°
- c) 30°
- d) 45°

396. The angle of elevation of an object from a point on the level ground is α . Moving d meters on the ground towards the object, the angle of elevation is found to be β . Then the height (in meters) of the object is

- a) $d \tan \alpha$
- b) $d \cot \beta$
- c) $\frac{d}{\cot \alpha + \cot \beta}$
- d) $\frac{d}{\cot \alpha - \cot \beta}$

397. In a ΔPQR as shown in figure given that $x : y : z = 2 : 3 : 6$, then the value of $\angle QPR$ is



- a) $\frac{\pi}{6}$
- b) $\frac{\pi}{4}$
- c) $\frac{\pi}{3}$
- d) None of these

398. If by shifting the origin at $(1, 1)$ the coordinates of a point P become $(\cos \theta, \cos \phi)$, then the original coordinates of P were

- a) $(2 \cos^2 \theta/2, 2 \cos^2 \phi/2)$
- b) $(2 \sin^2 \theta/2, 2 \sin^2 \phi/2)$
- c) $(2 \cos \theta/2, 2 \cos \phi/2)$
- d) $(2 \sin \theta/2, 2 \sin \phi/2)$

399. In a ΔABC , if $r_1 = 2r_2 = 3r_3$, then

- a) $\frac{a}{b} = \frac{4}{5}$
- b) $\frac{a}{b} = \frac{5}{4}$
- c) $a + b - 2c = 0$
- d) $2a = b + c$

400. In a ΔABC , if a, b, c are in AP, then the value of $\frac{\sin \frac{A}{2} \sin \frac{C}{2}}{\sin \frac{B}{2}}$ is

- a) 1
- b) $\frac{1}{2}$
- c) 2
- d) -1

401. The top of a hill observed from the top and bottom of a building h is at angles of elevation p and q respectively. The height of hill is

- a) $\frac{h \cot q}{\cot q - \cot p}$
- b) $\frac{h \cot p}{\cot p - \cot q}$
- c) $\frac{h \tan p}{\tan p - \tan q}$
- d) None of these

402. In a triangle, if $b = 20, c = 21$ and $\sin A = \frac{3}{5}$, then a is equal to

- a) 12
- b) 13
- c) 14
- d) 15

403. In ΔABC , $a = 2, b = 4$ and $\angle C = 60^\circ$, then $\angle A$ and $\angle B$ are equal to

- a) $90^\circ, 30^\circ$ b) $60^\circ, 60^\circ$ c) $30^\circ, 90^\circ$ d) $60^\circ, 45^\circ$
404. If in an equilateral triangle $R = \sqrt{3}$ cm, then the length of each side of the triangle is
a) 1 cm b) 2 cm c) 3 cm d) None of these
405. In a ΔABC , $\frac{b-c \cos A}{c-b \cos A}$ is equal to
a) $\frac{\sin B}{\sin C}$ b) $\frac{\cos C}{\cos B}$ c) $\frac{\cos B}{\cos C}$ d) None of these
406. In a ΔABC , if $2s = a + b + c$, then the value of the $\frac{s(s-a)}{bc} - \frac{(s-b)(s-c)}{bc}$ is equal to
a) $\sin A$ b) $\cos A$ c) $\tan A$ d) None of these
407. The centroid of a triangle is $(2, 7)$ and two of its vertices are $(4, 8)$ and $(-2, 6)$. The third vertex is
a) $(0, 0)$ b) $(4, 7)$ c) $(7, 4)$ d) $(7, 7)$
408. Each side of a square subtends an angle of 60° at the top of a tower h metres high standing in the centre of the square. If a is the length of each side of the square, then
a) $2a^2 = h^2$ b) $2h^2 = a^2$ c) $3a^2 = 2h^2$ d) $2h^2 = 3a^2$
409. In ΔABC , $(a-b)^2 \cos^2 \frac{C}{2} + (a+b)^2 \sin^2 \frac{C}{2}$ is equal to
a) a^2 b) b^2 c) c^2 d) None of these
410. If the area of a ΔABC is given by $\Delta = a^2 - (b-c)^2$, then $\tan \left(\frac{A}{2}\right)$ is equal to
a) -1 b) 0 c) $\frac{1}{4}$ d) $\frac{1}{2}$
411. If $O'(4, 8/3)$ is the orthocentre of the triangle ABC the coordinates of whose vertices are $O(0,0), A(8,0)$ and $B(4,6)$, then the coordinates of the orthocentre of $\Delta O'AB$ are
a) $(0, 0)$ b) $(8, 0)$ c) $(4, 6)$ d) None of these
412. In a ΔABC , if $b^2 + c^2 = 3a^2$, then $\cot B + \cot C - \cot A$ is equal to
a) 1 b) $\frac{ab}{4\Delta}$ c) 0 d) $\frac{ac}{4\Delta}$
413. From the top of a tower, the angle of depression of a point on the ground is 60° . If the distance of this point from the tower is $\frac{1}{\sqrt{3}+1} m$, then the height of the tower is
a) $\frac{4\sqrt{3}}{2} m$ b) $\frac{\sqrt{3}+3}{2} m$ c) $\frac{3-\sqrt{3}}{2} m$ d) $\frac{\sqrt{3}}{2} m$
414. If in a ΔABC , $a = 6$ cm, $b = 8$ cm, $c = 10$ cm, then the value of $\sin 2A$ is
a) $6/25$ b) $8/25$ c) $10/25$ d) $24/25$
415. In ΔABC , $\frac{1+\cos(A-B)\cos C}{1+\cos(A-C)\cos B}$ is equal to
a) $\frac{a-b}{a+b}$ b) $\frac{a+b}{a+c}$ c) $\frac{a^2-b^2}{a^2-c^2}$ d) $\frac{a^2+b^2}{a^2+c^2}$
416. The base of a cliff is circular. From the extremities of a diameter of the base angles of elevation of the top of the cliff are 30° and 60° . If the height of the cliff be 500 m, then the diameter of the base of the cliff is
a) $\frac{2000}{\sqrt{3}} m$ b) $\frac{1000}{\sqrt{3}} m$ c) $\frac{2000}{\sqrt{2}} m$ d) $1000\sqrt{3} m$
417. The sides BC, CA and AB of a triangle ABC are of lengths a, b and c respectively. If D is the mid point of BC and AD is perpendicular to AC , then the value of $\cos A \cos C$ is
a) $\frac{3(a^2-c^2)}{2ac}$ b) $\frac{2(a^2-c^2)}{3bc}$ c) $\frac{(a^2-c^2)}{3ac}$ d) $\frac{2(c^2-a^2)}{3ac}$
418. The mid points of the sides of a triangle are $D(6, 1), E(3, 5)$ and $F(-1, -2)$, then the vertex opposite to D is
a) $(-4, 2)$ b) $(-4, 5)$ c) $(2, 5)$ d) $(10, 8)$
419. The locus of a points which moves such that the sum of the squares of its distance from three vertices of the triangle is constant is a/an
a) Circle b) Straight line c) Ellipse d) None of the above
420. The angles A, B and C of a ΔABC are in AP. if $b:c = \sqrt{3}:\sqrt{2}$, then the angle A is
a) 30° b) 15° c) 75° d) 45°


421. If the three points $(0, 1)$, $(0, -1)$ and $(x, 0)$ are vertices of an equilateral triangle, then the values of x are
 a) $\sqrt{3}, \sqrt{2}$ b) $\sqrt{3}, -\sqrt{3}$ c) $-\sqrt{5}, \sqrt{3}$ d) $\sqrt{2}, -\sqrt{2}$
422. The sides of a triangle are $\sin \alpha \cos \alpha$ and $\sqrt{1 + \sin \alpha \cos \alpha}$ for some $0 < \alpha < \frac{\pi}{2}$. Then, the greatest angle of the triangle is
 a) 60° b) 90° c) 120° d) 150°
423. From an aeroplane flying, vertically above a horizontal road, the angles of depression of two consecutive stones on the same side of the aeroplane are observed to be 30° and 60° respectively. The height at which the aeroplane is flying in km, is
 a) $\frac{4}{\sqrt{3}}$ b) $\frac{\sqrt{3}}{2}$ c) $\frac{2}{\sqrt{3}}$ d) 2
424. A flag staff 20 m long standing on a wall 10 m high subtends an angle whose tangent is 0.5 at a point on the ground. If θ is the angle subtended by the wall at this point, then
 a) $\tan \theta = 1$ b) $\tan \theta = 3$ c) $\tan \theta = \frac{1}{2}$ d) None of these
425. The angle of elevation of the top of a vertical tower from two points distance a and b from the base and in the same line with it, are complimentary. If θ is the angle subtended at the top of the tower by the line joining these points then $\sin \theta =$
 a) $\frac{a-b}{\sqrt{2}(a+b)}$ b) $\frac{a+b}{a-b}$ c) $\frac{a-b}{a+b}$ d) None of these
426. In a triangle with one angle of 120° , the length of the sides forms an AP. If the length of the greatest side is 7 cm, then area of triangle is
 a) $\frac{3\sqrt{15}}{4} \text{ cm}^2$ b) $\frac{15\sqrt{3}}{4} \text{ cm}^2$ c) $\frac{15}{4} \text{ cm}^2$ d) $\frac{3\sqrt{3}}{4} \text{ cm}^2$
427. If in a ΔABC , $\frac{b+c}{11} = \frac{c+a}{12} = \frac{a+b}{13}$, then $\cos A$ is equal to
 a) $\frac{1}{5}$ b) $\frac{5}{7}$ c) $\frac{19}{35}$ d) None of these
428. If two vertices of an equilateral triangle are $(0, 0)$ and $(3, 3\sqrt{3})$, then the third vertex lies at
 a) $(3, -3)$ b) $(-3, 3)$ c) $(-3, 3\sqrt{3})$ d) None of these
429. In an isosceles right angled ΔABC , $\angle B = 90^\circ$, AD is the median, then $\frac{\sin \angle BAD}{\sin \angle CAD}$ is
 a) $\frac{1}{\sqrt{2}}$ b) $\sqrt{2}$ c) 1 d) None of these
430. If $P(1, 2)$, $Q(4, 6)$, $R(5, 7)$ and $S(a, b)$ are the vertices of a parallelogram $PQRS$, then
 a) $a = 2, b = 4$ b) $a = b, b = 4$ c) $a = 2, b = 3$ d) $a = 3, b = 5$
431. Angles of a triangle are in the ratio 4: 1: 1. The ratio between its greatest side and perimeter is
 a) $\frac{3}{2 + \sqrt{3}}$ b) $\frac{1}{2 + \sqrt{3}}$ c) $\frac{\sqrt{3}}{\sqrt{3} + 2}$ d) $\frac{2}{2 + \sqrt{3}}$
432. From the top of a cliff h metres above sea level an observer notices that angles of depression of an object A and its image B are complementary. If the angle of depression at A is θ . The height of A above sea level is
 a) $h \sin \theta$ b) $h \cos \theta$ c) $h \sin 2\theta$ d) $h \cos 2\theta$
433. The radius of the incircle of triangle when sides are 18, 24 and 30 cm is
 a) 2 cm b) 4 cm c) 6 cm d) 9 cm
434. The points $(1, 3)$ and $(5, 1)$ are the opposite vertices of a rectangle. The other two vertices lie on the line $y = 2x + c$, then the value of c will be
 a) 4 b) -4 c) 2 d) -2
435. If the vertices of a triangle at $O(0,0)$, $A(a, 0)$ and $B(0, a)$. Then, the distance between its circumcentre and orthocentre is
 a) $\frac{a}{2}$ b) $\frac{a}{\sqrt{2}}$ c) $\sqrt{2} a$ d) $\frac{a}{4}$
436. The straight lines $x = y$, $x - 2y = 3$ and $x + 2y = -3$ form a triangle, which is

- a) Isosceles b) Equilateral c) Right angled d) None of these
437. The vertices of a triangle are $(6, 0)$, $(0, 6)$ and $(6, 6)$. The distance between its circumcentre and centroid is
a) 2 b) $\sqrt{2}$ c) 1 d) $2\sqrt{2}$
438. In a ΔABC , $a = 5$, $b = 7$ and $\sin A = \frac{3}{4}$, then the number of possible triangles are
a) 1 b) 0 c) 2 d) Infinite
439. The points $(k, 2 - 2k)$, $(-k + 1, 2k)$, $(-4 - k, 6 - 2k)$ are collinear, then k is equal to
a) 2, 3 b) 1, 0 c) $\frac{1}{2}, 1$ d) 1, 2
440. If the vertices P, Q, R of a ΔPQR are rational points, which of the following points of the ΔPQR is (are) always rational points?
(A rational point is a point both of whose coordinates are rational numbers)
a) Centroid b) Incentre c) Circumcentre d) Orthocentre
441. Which of the following pieces of data does not uniquely determine acute angled ΔABC (R = circumradius)?
a) $a, \sin A, \sin B$ b) a, b, c c) $a, \sin B, R$ d) $a, \sin A, R$
442. If a ΔABC , $2ca \sin \frac{A-B+C}{2}$ is equal to
a) $a^2 + b^2 - c^2$ b) $c^2 + a^2 - b^2$ c) $b^2 - c^2 - a^2$ d) $c^2 - a^2 - b^2$
443. In any ΔABC under usual notation, $a(b \cos C - c \cos B)$ is equal to
a) $b^2 - c^2$ b) $c^2 - b^2$ c) $\frac{b^2 - c^2}{2}$ d) $\frac{c^2 - b^2}{2}$
444. In ΔABC , G is the centroid, D is the mid point of BC . If $A = (2, 3)$ and $G(7, 5)$, then the point D is
a) $(\frac{9}{2}, 4)$ b) $(\frac{19}{2}, 6)$ c) $(\frac{11}{2}, \frac{11}{2})$ d) $(8, \frac{13}{2})$
445. The transformed equation of $3x^2 + 3y^2 + 2xy = 2$, when the coordinate axes are rotated through an angle of 45° , is
a) $x^2 + 2y^2 = 1$ b) $2x^2 + y^2 = 1$ c) $x^2 + y^2 = 1$ d) $x^2 + 3y^2 = 1$
446. The equation of the locus of a point equidistant from the points (a_1, b_1) and (a_2, b_2) is $(a_1 - a_2)x + (b_1 - b_2)y + c = 0$, then the value of c is
a) $\sqrt{(a_1^2 + b_1^2 + c_1^2)}$ b) $a_1^2 - b_1^2 - c_1^2$
c) $\frac{1}{2}(a_2^2 + b_2^2 - a_1^2 - b_1^2)$ d) None of the above
447. ABC is a right angled triangle with $\angle B = 90^\circ$, $a = 6$ cm. If the radius of the circumcircle is 5 cm. Then the area of ΔABC is
a) 25 cm^2 b) 30 cm^2 c) 36 cm^2 d) 24 cm^2
448. The transformed equation of $x^2 + 6xy + 8y^2 = 10$ when the axes are rotated through an angle $\frac{\pi}{4}$ is
a) $15x^2 - 14xy + 3y^2 = 20$ b) $15x^2 + 14xy - 3y^2 = 20$
c) $15x^2 + 14xy + 3y^2 = 20$ d) $15x^2 - 14xy - 3y^2 = 20$
449. The angle of elevation of the top of a TV tower from three points A, B and C in a straight line through the foot of the tower are $\alpha, 2\alpha$ and 3α respectively. If $AB = a$, then height of the tower is
a) $a \tan \alpha$ b) $a \sin \alpha$ c) $a \sin 2\alpha$ d) $a \sin 3\alpha$
450. The ratio in which the x -axis divides the line segment joining $(3, 6)$ and $(4, -3)$ is
a) $2 : 1$ b) $1 : 2$ c) $3 : 4$ d) None of these
451. AB is vertical tower. The point A is on the ground and C is the middle point of AB . The part CB subtend an angle α at a point P on the ground. If $AP = n AB$, then the correct relation is
a) $n = (n^2 + 1) \tan \alpha$ b) $n = (2n^2 - 1) \tan \alpha$ c) $n^2 = (2n^2 + 1) \tan \alpha$ d) $n = (2n^2 + 1) \tan \alpha$
452. If the points $(-2, -5)$, $(2, -2)$, $(8, a)$ are collinear, then the value of a is
a) $-\frac{5}{2}$ b) $\frac{5}{2}$ c) $\frac{3}{2}$ d) $\frac{1}{2}$

453. If in a ΔABC , $\cos A + 2 \cos B + \cos C = 2$, then a, b, c are in
 a) AP b) GP c) HP d) None of these
454. In a ΔABC , $a = 5, b = 4$ and $\cos(A + B) = \frac{31}{32}$. In this triangle, c is equal to
 a) $\sqrt{6}$ b) 36 c) 6 d) None of these
455. In a triangle $r_1 > r_2 > r_3$, then
 a) $a > b > c$ b) $a < b < c$ c) $a > b$ and $b < c$ d) $a < b$ and $b > c$
456. In a ΔABC , if $a = 2x, b = 2y$ and $\angle C = 120^\circ$, then the area of the triangle is
 a) xy sq unit b) $xy\sqrt{3}$ sq unit c) $3xy$ sq unit d) $2xy$ sq unit
457. If the points $(a, b), (a', b')$ and $(a - a', b - b')$ are collinear, then
 a) $ab' = a'b$ b) $ab = a'b'$ c) $aa' = bb'$ d) $a^2 + b^2 = 1$
458. If $A(-a, 0)$ and $B(a, 0)$ are two fixed points, then the locus of the point at which AB subtends a right angle, is
 a) $x^2 + y^2 = 2a^2$ b) $x^2 - y^2 = a^2$ c) $x^2 + y^2 + a^2 = 0$ d) $x^2 + y^2 = a^2$
459. If the coordinates of two vertices of an equilateral triangle are $(2, 4)$ and $(2, 6)$, then the coordinates of its third vertex are
 a) $(\sqrt{3}, 5)$ b) $(2\sqrt{3}, 5)$ c) $(2 + \sqrt{3}, 5)$ d) $(2, 5)$
460. If point (x, y) is equidistant from $(a + b, b - a)$ and $(a - b, a + b)$, then
 a) $ax + by = 0$ b) $ax - by = 0$ c) $bx + ay = 0$ d) $bx - ay = 0$
461. If A, B, C, D are the angles of a quadrilateral, then $\frac{\sum \tan A}{\sum \cot A}$ is equal to
 a) $\prod \tan A$ b) $\prod \cot A$ c) $\sum \tan^2 A$ d) $\sum \cot^2 A$
462. If the angles of a triangle are in the ratio 3:4:5, then the sides are in the ratio
 a) $2:\sqrt{6}:\sqrt{3} + 1$ b) $\sqrt{2}:\sqrt{6}:\sqrt{3} + 1$ c) $2:\sqrt{3}:\sqrt{3} + 1$ d) 3:4:5
463. A line joining $A(2, 0)$ and $B(3, 1)$ is rotated about A in anti-clockwise direction through 15° . Find the equation of the line in the new position. If B goes to C in the new position, then coordinates of C are
 a) $\left(2 + \frac{1}{\sqrt{2}}, \frac{\sqrt{3}}{\sqrt{2}}\right)$ b) $\left(2 - \frac{1}{\sqrt{2}}, \frac{\sqrt{3}}{\sqrt{2}}\right)$ c) $\left(2 + \frac{1}{\sqrt{2}}, \frac{\sqrt{3}}{2}\right)$ d) None of these
464. ABC is a triangle with vertices $A(-1, 4), B(6, -2)$ and $C(-2, 4)$. D, E and F are the points which divide each AB, BC , and CA respectively in the ratio 3 : 1 internally. Then, the centroid of ΔDEF is
 a) (3, 6) b) (1, 2) c) (4, 8) d) (-3, 6)
465. A house subtends a right angle at the window of an opposite house and the angle of elevation of the window from the bottom of the first house is 60° . If the distance between the two houses be 6 m, then the height of the first house is
 a) $8\sqrt{3}$ m b) $6\sqrt{3}$ m c) $4\sqrt{3}$ m d) None of these
466. The orthocentre of the triangle whose vertices are $\{at_1 t_2, a(t_1 + t_2)\}, \{at_2 t_3, a(t_2 + t_3)\}, \{at_3 t_1, a(t_3 + t_1)\}$ is
 a) $\{-a, a(t_1 + t_2 + t_3 + t_1 t_2 t_3)\}$ b) $\{-a, a(t_1 + t_2 + t_3 + t_1 t_2 t_3)\}$
 c) $\{-a, a(t_1 - t_2 - t_3 - t_1 t_2 t_3)\}$ d) $\{-a, a(t_1 + t_2 - t_3 - t_1 t_2 t_3)\}$
467. A flag staff is upon the top of a building. If at a distance of 40 m from the base of building the angles of elevation of the top of the flag staff and building are 60° and 30° respectively, then the height of the flag staff is
 a) 46.19 m b) 50 m c) 25 m d) None of these
468. A person observes the angle of elevation of a building as 30° . The person proceeds towards the building with a speed of $25(\sqrt{3} - 1)$ m/h. After two hours, he observes the angle of elevation as 45° . The height of the building (in metres) is
 a) $50(\sqrt{3} - 1)$ b) $50(\sqrt{3} + 1)$ c) 50 d) 100
469. In a ΔABC , $\sum (b + c) \tan \frac{A}{2} \tan \left(\frac{B-C}{2}\right)$ is equal to
 a) a b) b c) c d) 0



470. In a ΔABC , angle A is greater than angle B . If the measures of angles A and B satisfy the equation $3 \sin x - 4 \sin^3 x - k = 0$, $0 < k < 1$, then the measure of angle C is
 a) $\frac{\pi}{3}$ b) $\frac{\pi}{2}$ c) $\frac{2\pi}{3}$ d) $\frac{5\pi}{6}$
471. In a ΔABC , a, c, A are given and b_1, b_2 are two values of third side b such that $b = 2b_1$. Then, $\sin A$ is equal to
 a) $\sqrt{\frac{9a^2 - c^2}{8a^2}}$ b) $\sqrt{\frac{9a^2 - c^2}{8c^2}}$ c) $\sqrt{\frac{9a^2 + c^2}{8a^2}}$ d) None of these
472. The incentre of a triangle with vertices $(7, 1)$, $(-1, 5)$ and $(3 + 2\sqrt{3}, 3 + 4\sqrt{3})$ is
 a) $(3 + \frac{2}{\sqrt{3}}, 3 + \frac{4}{\sqrt{3}})$ b) $(1 + \frac{2}{3\sqrt{3}}, 1 + \frac{4}{3\sqrt{3}})$ c) $(7, 1)$ d) None of the above
473. In any triangle ABC , $\frac{\tan \frac{A}{2} - \tan \frac{B}{2}}{\tan \frac{A}{2} + \tan \frac{B}{2}}$ is equal to
 a) $\frac{a-b}{a+b}$ b) $\frac{a-b}{2}$ c) $\frac{a-b}{a+b+c}$ d) $\frac{c}{a+b}$
474. If orthocentre and circumcentre of triangle are respectively $(1, 1)$ and $(3, 2)$, then the coordinates of its centroid are
 a) $(\frac{7}{3}, \frac{5}{3})$ b) $(\frac{5}{3}, \frac{7}{3})$ c) $(7, 5)$ d) None of these
475. In ΔABC , AD is median and $\angle A = 60^\circ$, then $4 AD^2$ is equal to



- a) $b^2 + c^2 - bc$ b) $2b^2 + c^2 - 2bc$ c) $b^2 + c^2 + 2bc$ d) None of these
476. The circumcentre of the triangle formed by the lines $y = x, y = 2x$ and $y = 3x + 4$ is
 a) $(6, 8)$ b) $(6, -8)$ c) $(3, 4)$ d) $(-3, -4)$
477. If in a ΔABC , $\cot \frac{A}{2} = \frac{b+c}{a}$, then the ΔABC is
 a) Isosceles b) Equilateral c) Right angled d) None of these
478. Area of the triangle formed by the lines $3x^2 - 4xy + y^2 = 0, 2x - y = 6$ is
 a) 16 sq units b) 25 sq units c) 36 sq units d) 49 sq units
479. The shadow of a tower is found to be 60 m shorter when the sun's altitude changes from 30° to 60° . The height of the tower from the ground is approximately equal to
 a) 62 m b) 301 m c) 101 m d) 52 m
480. Let $0 < \alpha < \frac{\pi}{2}$ be a fixed angle. If $P = (\cos \theta, \sin \theta)$ and $Q = \{\cos(\alpha - \theta), \sin(\alpha - \theta)\}$. Then Q is obtained from P by
 a) Clockwise rotation around the origin through angle α b) Anti-clockwise rotation around origin through angle α
 c) Reflection in the line through the origin with slope $\tan \alpha$ d) Reflection in the line through the origin with slope $\tan \frac{\alpha}{2}$
481. The equation of the base of an equilateral triangle is $x + y = 2$ and the vertex is $(2, -1)$. The area of triangle is
 a) $2\sqrt{3}$ b) $\frac{\sqrt{3}}{6}$ c) $\frac{1}{\sqrt{3}}$ d) $\frac{2}{\sqrt{3}}$
482. If $(0, 1)$ is the orthocentre and $(2, 3)$ is the centroid of a triangle. Then, its circumcentre is
 a) $(3, 2)$ b) $(1, 0)$ c) $(4, 3)$ d) $(3, 4)$



483. The angle of elevation of the top of the tower observed from each of the three points A, B, C on the ground forming a triangle is the same angle α . If R is the circumradius of the ΔABC , then the height of the tower is
 a) $R \sin \alpha$ b) $R \cos \alpha$ c) $R \cot \alpha$ d) $R \tan \alpha$
484. The circumcentre of a triangle formed by the lines $xy + 2x + 2y + 4 = 0$ and $x + y + 2 = 0$ is
 a) $(-1, -1)$ b) $(0, -1)$ c) $(1, 1)$ d) $(-1, 0)$
485. If the distance of any point P from the points $A(a + b, a - b)$ and $B(a - b, a + b)$ are equal, then the locus of P is
 a) $ax + by = 0$ b) $x - y = 0$ c) $x + y = 0$ d) $bx - ay = 0$
486. In an equilateral triangle of side $2\sqrt{3}$ cm, the circumcentre is
 a) 1 cm b) $\sqrt{3}$ cm c) 2 cm d) $2\sqrt{3}$ cm
487. Let ABC be a triangle, two of whose vertices are $(15, 0)$ and $(0, 10)$. If the orthocenter is $(6, 9)$, then the third vertex is
 a) $(15, 10)$ b) $(10, -15)$ c) $(0, 0)$ d) None of these
488. An aeroplane flying at a height of 300 metres above the ground passes vertically above another plane at an instant when the angles of elevation of the two planes from the same point on the ground are 60° and 45° respectively. The height of the lower plane from the ground (in metres) is
 a) $100\sqrt{3}$ b) $\frac{100}{\sqrt{3}}$ c) 50 d) $150(\sqrt{3} + 1)$
489. If the sides of a triangle are in ratio 3:7:8, then $R:r$ is equal to
 a) 2:7 b) 7:2 c) 3:7 d) 7:3
490. In a ΔABC , $(b + c - a) \tan \frac{A}{2}$ is equal to
 a) $\frac{2\Delta}{s}$ b) $\frac{\Delta}{s}$ c) $\frac{\Delta s}{bc}$ d) $\frac{s}{a} R$
491. The sum of the radii of inscribed and circumscribed circles for an n sides regular polygon of side a , is
 a) $a \cot\left(\frac{\pi}{n}\right)$ b) $\frac{a}{2} \cot\left(\frac{\pi}{2n}\right)$ c) $a \cot\left(\frac{\pi}{2n}\right)$ d) $\frac{a}{4} \cot\left(\frac{\pi}{2n}\right)$
492. If $\Delta = a^2 - (b - c)^2$, where Δ is the area of ΔABC , then $\tan A$ is equal to
 a) $\frac{15}{16}$ b) $\frac{8}{17}$ c) $\frac{8}{15}$ d) $\frac{1}{2}$
493. The median BE and AD of a triangle with vertices $A(0, b), B(0, 0), C(a, 0)$ are perpendicular to each other, if
 a) $a = \frac{b}{2}$ b) $b = \frac{a}{2}$ c) $ab = 1$ d) $a = \pm\sqrt{2}b$
494. If a, b, c be the sides of a ΔABC and if roots of the equation $a(b - c)x^2 + b(c - a)x + c(a - b) = 0$ are equal, then $\sin^2\left(\frac{A}{2}\right), \sin^2\left(\frac{B}{2}\right), \sin^2\left(\frac{C}{2}\right)$ are in
 a) AP b) GP c) HP d) AGP
495. In ΔABC , $\frac{1 + \cos(A - B) \cos C}{1 + \cos(A - C) \cos B}$ is equal to
 a) $\frac{a - b}{a - c}$ b) $\frac{a + b}{a + c}$ c) $\frac{a^2 - b^2}{a^2 - c^2}$ d) $\frac{a^2 + b^2}{a^2 + c^2}$
496. In a ΔABC , $a = 13$ cm, $b = 12$ cm and $c = 5$ cm. The distance of A from BC is
 a) $\frac{144}{13}$ b) $\frac{65}{12}$ c) $\frac{60}{13}$ d) $\frac{25}{13}$
497. If in a ΔABC , $r_1 < r_2 < r_3$, then
 a) $a < b < c$ b) $a > b > c$ c) $b < a < c$ d) $a < c < b$
498. A ladder rests against a wall making an angle α with the horizontal. The foot of the ladder is pulled away from the wall through a distance x , so that it slides a distance y down the wall making an angle β with the horizontal. The correct relation is
 a) $x = y \tan\left(\frac{\alpha + \beta}{2}\right)$ b) $y = x \tan\left(\frac{\alpha + \beta}{2}\right)$ c) $x = y \tan(\alpha + \beta)$ d) $y = x \tan(\alpha + \beta)$

499. Let $A(h, k)$, $B(1, 1)$ and $C(2, 1)$ be the vertices of a right angled triangle with AC as its hypotenuse. If the area of the triangle is 1, then the set of values which ' k ' can take is given by
 a) $\{1, 3\}$ b) $\{0, 2\}$ c) $\{-1, 3\}$ d) $\{-3, -2\}$
500. A variable line $\frac{x}{a} + \frac{y}{b} = 1$ is such that $a + b = 4$. The locus of the mid point of the portion of the line intercepted between the axes is
 a) $x + y = 4$ b) $x + y = 8$ c) $x + y = 1$ d) $x + y = 2$
501. If in a ΔABC , $\tan\left(\frac{A}{2}\right)$, $\tan\left(\frac{B}{2}\right)$, $\tan\left(\frac{C}{2}\right)$ are in HP, then the sides a, b, c are in
 a) AP b) GP c) HP d) None of these
502. The shadow of tower standing on a level ground is x metres long when the sun's altitude is 30° , while it is y metres long when the altitude is 60° . If the height of the tower is $45 \cdot \frac{\sqrt{3}}{2}$ m, then $x - y$ is
 a) 45 m b) $45\sqrt{3}$ m c) $\frac{45}{\sqrt{3}}$ m d) $45 \cdot \frac{\sqrt{3}}{2}$ m
503. All points lying inside the triangle formed by the points $(1, 3)$, $(5, 0)$ and $(-1, 2)$ satisfy
 a) $3x + 2y \geq 0$ b) $2x + y - 13 < 0$ c) $2x - 3y - 12 \leq 0$ d) All of these
504. If the area of the triangle with vertices $(x, 0)$, $(1, 1)$ and $(0, 2)$ is 4 sq unit, then the value of x is
 a) -2 b) -4 c) -6 d) 8
505. If the centroid of the triangle formed by the points $(0, 0)$, $(\cos \theta, \sin \theta)$ and $(\sin \theta, -\cos \theta)$ lies on the line $y = 2x$, then θ is equal to
 a) $\tan^{-1} 2$ b) $\tan^{-1} 3$ c) $\tan^{-1}(-3)$ d) $\tan^{-1}(-2)$
506. In order to remove first degree terms from the equation $2x^2 + 7y^2 + 8x - 14y + 4 = 0$, the origin is shifted at the point
 a) $(-2, 1)$ b) $(1, 2)$ c) $(2, 1)$ d) $(1, -2)$
507. If $t_1 + t_2 + t_3 = -t_1 t_2 t_3$, then orthocentre of the triangle formed by the points $[at_1 t_2, a(t_1 + t_2)]$, $[at_2 t_3, a(t_2 + t_3)]$ and $[at_3 t_1, a(t_3 + t_1)]$, lies on
 a) $(a, 0)$ b) $(-a, 0)$ c) $(0, a)$ d) $(0, -a)$
508. If $A(-5, 0)$ and $B(3, 0)$ are two vertices of a triangle ABC . Its area is 20 sq cm. The vertex C lies on the line $x - y = 2$. The coordinates of C are
 a) $(-7, -5)$ or $(3, 5)$ b) $(-3, -5)$ or $(-5, 7)$ c) $(7, 5)$ or $(3, 5)$ d) $(-3, -5)$ or $(7, 5)$
509. The locus of a point P which moves such that $2PA = 3PB$, where coordinates of points A and B are $(0, 0)$ and $(4, -3)$, is
 a) $5x^2 - 5y^2 - 72x + 54y + 225 = 0$ b) $5x^2 + 5y^2 - 72x + 54y + 225 = 0$
 c) $5x^2 + 5y^2 + 72x - 54y + 225 = 0$ d) $5x^2 + 5y^2 - 72x - 54y - 225 = 0$
510. The coordinates of the centroid of a triangle having its circumcentre and orthocentre at $(7/2, 5/2)$ and $(2, 1)$ respectively, are
 a) $(3, 2)$ b) $(13/6, 3/2)$ c) $(5/2, 3/2)$ d) $(3/2, 5/2)$
511. The base angle of triangle are $22\frac{1}{2}^\circ$ and $112\frac{1}{2}^\circ$. If b is the base and h is the height of the triangle, then
 a) $b = 2h$ b) $b = 3h$ c) $b = (1 + \sqrt{3})h$ d) $b = (2 + \sqrt{3})h$
512. Let ABC be a triangle such that $\angle ACB = \frac{\pi}{6}$ and let a, b and c denote the lengths of the sides opposite to A, B and C respectively. The value (s) of x for which $a = x^2 + x + 1$, $b = x^2 - 1$ and $c = 2x + 1$ is (are)
 a) $-(2 + \sqrt{3})$ b) $1 + \sqrt{3}$ c) $2 + \sqrt{3}$ d) $4\sqrt{3}$
513. If the distance between the points $P(a \cos 48^\circ, 0)$ and $Q(0, a \cos 12^\circ)$ is d , then $d^2 - a^2 =$
 a) $\frac{a^2}{4}(\sqrt{5} - 1)$ b) $\frac{a^2}{4}(\sqrt{5} + 1)$ c) $\frac{a}{8}(\sqrt{5} - 1)$ d) $\frac{a^2}{8}(\sqrt{5} + 1)$
514. In a ΔABC , $a(b \cos C - c \cos B)$ is equal to
 a) a^2 b) $b^2 - c^2$ c) 0 d) None of these
515. If the points (a_1, b_1) , (a_2, b_2) and (a_3, b_3) are collinear, then lines $a_i x + b_i y + 1 = 0$ for $i = 1, 2, 3$ are
 a) Concurrent b) Identical c) Parallel d) None of these
516. In ΔABC , $(b + c) \cos A + (c + a) \cos B + (a + b) \cos C$ is equal to

517. If in a ΔABC $a = 5, b = 4, A = \frac{\pi}{2} + B$, then C
- a) 0 b) 1 c) $a + b + c$ d) $2(a + b + c)$
- a) is $\tan^{-1}\left(\frac{1}{9}\right)$ b) is $\tan^{-1}\left(\frac{9}{40}\right)$ c) Cannot be evaluated d) is $2 \tan^{-1}\left(\frac{1}{9}\right)$
518. If p_1, p_2, p_3 are altitudes of a ΔABC drawn from the vertices A, B, C and Δ the area of the triangle, then $p_1^{-2} + p_2^{-2} + p_3^{-2}$ is equal to
- a) $\frac{a + b + c}{\Delta}$ b) $\frac{a^2 + b^2 + c^2}{4\Delta^2}$ c) $\frac{a^2 + b^2 + c^2}{\Delta^2}$ d) None of these

CO-ORDINATE GEOMETRY

: ANSWER KEY :

1)	a	2)	a	3)	d	4)	b	153)	d	154)	b	155)	a	156)	b
5)	a	6)	b	7)	d	8)	a	157)	c	158)	a	159)	c	160)	d
9)	d	10)	b	11)	b	12)	a	161)	d	162)	c	163)	d	164)	d
13)	c	14)	c	15)	b	16)	d	165)	c	166)	b	167)	a	168)	a
17)	c	18)	d	19)	c	20)	b	169)	b	170)	c	171)	c	172)	a
21)	c	22)	a	23)	c	24)	a	173)	c	174)	c	175)	a	176)	a
25)	c	26)	c	27)	b	28)	b	177)	d	178)	a	179)	c	180)	c
29)	a	30)	a	31)	c	32)	b	181)	b	182)	a	183)	d	184)	d
33)	b	34)	a	35)	b	36)	a	185)	b	186)	a	187)	a	188)	d
37)	b	38)	d	39)	d	40)	a	189)	a	190)	a	191)	b	192)	a
41)	a	42)	a	43)	c	44)	c	193)	d	194)	d	195)	b	196)	d
45)	c	46)	a	47)	d	48)	a	197)	c	198)	b	199)	c	200)	a
49)	d	50)	a	51)	d	52)	b	201)	b	202)	a	203)	a	204)	c
53)	a	54)	d	55)	b	56)	a	205)	a	206)	b	207)	a	208)	a
57)	c	58)	d	59)	a	60)	b	209)	c	210)	d	211)	c	212)	c
61)	b	62)	d	63)	a	64)	c	213)	b	214)	b	215)	c	216)	d
65)	c	66)	a	67)	b	68)	d	217)	a	218)	d	219)	b	220)	d
69)	c	70)	d	71)	c	72)	c	221)	a	222)	b	223)	b	224)	c
73)	a	74)	a	75)	d	76)	b	225)	c	226)	c	227)	a	228)	d
77)	c	78)	b	79)	d	80)	a	229)	a	230)	a	231)	a	232)	c
81)	b	82)	d	83)	a	84)	c	233)	c	234)	d	235)	d	236)	a
85)	a	86)	d	87)	c	88)	c	237)	a	238)	d	239)	a	240)	d
89)	c	90)	c	91)	a	92)	d	241)	c	242)	c	243)	a	244)	c
93)	c	94)	c	95)	a	96)	a	245)	a	246)	a	247)	d	248)	d
97)	b	98)	c	99)	c	100)	c	249)	d	250)	d	251)	d	252)	b
101)	a	102)	a	103)	a	104)	b	253)	c	254)	c	255)	c	256)	d
105)	c	106)	b	107)	c	108)	d	257)	a	258)	a	259)	c	260)	c
109)	b	110)	a	111)	c	112)	b	261)	a	262)	c	263)	d	264)	b
113)	a	114)	b	115)	d	116)	b	265)	b	266)	a	267)	c	268)	b
117)	a	118)	a	119)	b	120)	b	269)	d	270)	c	271)	a	272)	d
121)	a	122)	c	123)	d	124)	b	273)	a	274)	d	275)	d	276)	b
125)	c	126)	b	127)	d	128)	d	277)	a	278)	b	279)	b	280)	c
129)	c	130)	c	131)	c	132)	b	281)	c	282)	b	283)	c	284)	c
133)	b	134)	d	135)	c	136)	c	285)	a	286)	d	287)	b	288)	a
137)	b	138)	c	139)	d	140)	c	289)	a	290)	d	291)	b	292)	d
141)	c	142)	b	143)	d	144)	d	293)	a	294)	a	295)	d	296)	c
145)	c	146)	d	147)	b	148)	c	297)	c	298)	a	299)	a	300)	b
149)	b	150)	a	151)	c	152)	a	301)	d	302)	a	303)	b	304)	b



305) a	306) c	307) d	308) d	417) d	418) a	419) a	420) c
309) c	310) a	311) d	312) c	421) b	422) c	423) b	424) a
313) b	314) a	315) d	316) c	425) c	426) b	427) a	428) c
317) a	318) b	319) a	320) d	429) b	430) c	431) c	432) d
321) b	322) c	323) c	324) b	433) c	434) b	435) b	436) d
325) b	326) b	327) d	328) a	437) b	438) b	439) c	440) a
329) d	330) c	331) d	332) a	441) d	442) b	443) a	444) b
333) d	334) d	335) a	336) a	445) b	446) c	447) d	448) c
337) a	338) b	339) b	340) c	449) c	450) a	451) d	452) b
341) c	342) b	343) a	344) d	453) a	454) d	455) a	456) b
345) c	346) d	347) d	348) a	457) a	458) d	459) c	460) d
349) d	350) a	351) c	352) a	461) a	462) a	463) a	464) b
353) b	354) b	355) b	356) c	465) a	466) b	467) a	468) c
357) c	358) c	359) d	360) d	469) d	470) c	471) b	472) a
361) b	362) b	363) b	364) a	473) b	474) a	475) d	476) b
365) b	366) a	367) c	368) b	477) c	478) c	479) d	480) d
369) d	370) a	371) b	372) a	481) b	482) d	483) d	484) a
373) d	374) b	375) b	376) b	485) b	486) c	487) c	488) a
377) a	378) c	379) a	380) d	489) b	490) a	491) b	492) c
381) d	382) a	383) d	384) a	493) d	494) c	495) d	496) c
385) a	386) b	387) d	388) b	497) a	498) a	499) c	500) d
389) c	390) b	391) b	392) b	501) a	502) a	503) d	504) c
393) b	394) b	395) b	396) d	505) c	506) a	507) b	508) d
397) b	398) a	399) b	400) b	509) b	510) a	511) a	512) b
401) b	402) b	403) c	404) c	513) d	514) b	515) a	516) c
405) b	406) b	407) b	408) b	517) b	518) b		
409) c	410) c	411) a	412) c				
413) c	414) d	415) d	416) a				



CO-ORDINATE GEOMETRY

: HINTS AND SOLUTIONS :

1 (a)

$$\begin{aligned}
 & a^2(\cos^2 B - \cos^2 C) + b^2(\cos^2 C - \cos^2 A) \\
 & \quad + c^2(\cos^2 A - \cos^2 B) \\
 &= a^2(1 - \sin^2 B - 1 + \sin^2 C) \\
 & \quad + b^2(1 - \sin^2 C - 1 + \sin^2 A) \\
 &+ c^2(1 - \sin^2 A - 1 + \sin^2 B) \\
 &= a^2(\sin^2 C - \sin^2 B) + b^2(\sin^2 A - \sin^2 C) \\
 & \quad + c^2(\sin^2 B - \sin^2 A) \\
 &= k^2 a^2(c^2 - b^2) + k^2 b^2(a^2 - c^2) + k^2 c^2(b^2 - c^2) \\
 &= 0
 \end{aligned}$$

2 (a)

Let $\sin A = 3k, \sin B = 4k, \sin C = 5k$

$$\begin{aligned}
 \therefore \frac{\sin A}{a} &= \frac{\sin B}{b} = \frac{\sin C}{c} = p \quad [\text{say}] \\
 \Rightarrow \frac{3k}{a} &= \frac{4k}{b} = \frac{5k}{c} = p \\
 \Rightarrow a &= 3\left(\frac{k}{p}\right), b = 4\left(\frac{k}{p}\right), c = 5\left(\frac{k}{p}\right) \\
 \Rightarrow a &= 3l, b = 4l, c = 5l \quad \left[\text{let } l = \frac{k}{p}\right]
 \end{aligned}$$

$$\begin{aligned}
 \therefore \cos A &= \frac{b^2 + c^2 - a^2}{2bc} \\
 &= \frac{16 + 25 - 9}{2 \times 4 \times 5} = \frac{32}{40} = \frac{4}{5} \\
 \therefore \cos B &= \frac{c^2 + a^2 - b^2}{2ac} \\
 &= \frac{25 + 9 - 16}{2 \times 3 \times 5} = \frac{18}{30} = \frac{3}{5}
 \end{aligned}$$

Now, $\cos A : \cos B = \frac{4}{5} : \frac{3}{5} = 4 : 3$

4 (b)

Slope of perpendicular to the line joining the points $(a \cos \alpha, a \sin \alpha)$ and $(a \cos \beta, a \sin \beta) =$

$$\begin{aligned}
 & -\frac{\cos \alpha - \cos \beta}{\sin \alpha - \sin \beta} \\
 &= \tan \frac{\alpha + \beta}{2}
 \end{aligned}$$

Hence, equation of perpendicular is

$$y = \tan \left(\frac{\alpha + \beta}{2} \right) x \quad \dots(i)$$

Now, on solving the equation of line with Eq. (i), we get


$$\left[\frac{a}{2}(\cos \alpha + \cos \beta), \frac{a}{2}(\sin \alpha + \sin \beta) \right]$$

5 (a)

$$\begin{aligned}
 \text{Area of } \frac{\Delta PBC}{\Delta ABC} &= \frac{\{-3(-2-y) + 4(y-5) + x(5+2)\}}{\{6(5+2) - 3(-2-3) + 4(3-5)\}} \\
 &= \left| \frac{7x + 7y - 14}{49} \right| = \left| \frac{x + y - 2}{7} \right|
 \end{aligned}$$

6 (b)

Let PQ and RS be the poles of height 20 m and 80 m subtending angles α and β at R and P respectively. Let h be the height of the point T , the intersection of QR and PS



$$\begin{aligned}
 \text{Then, } PR &= h \cot \alpha + h \cot \beta \\
 &= 20 \cot \alpha + 80 \cot \beta \\
 \Rightarrow \cot \alpha &= 4 \cot \beta \\
 \Rightarrow \frac{\cot \alpha}{\cot \beta} &= 4
 \end{aligned}$$

$$\begin{aligned}
 \text{Again, } h \cot \alpha + h \cot \beta &= 20 \cot \alpha \\
 \Rightarrow (h - 20) \cot \alpha &= -h \cot \beta \\
 \Rightarrow \frac{\cot \alpha}{\cot \beta} &= \frac{h}{20 - h} = 4 \\
 \Rightarrow h &= 80 - 4h \\
 \Rightarrow h &= 16 \text{ m}
 \end{aligned}$$

8 (a)

Since, α, β, γ are the roots of the equation $x^3 - 3px^2 + 3qx - 1 = 0$

$$\begin{aligned}
 \therefore \alpha + \beta + \gamma &= 3p \\
 \alpha\beta + \beta\gamma + \gamma\alpha &= 3q \\
 \text{and } \alpha\beta\gamma &= 1
 \end{aligned}$$

Let $G(x, y)$ be the centroid of the given triangle

$$\therefore x = \frac{\alpha + \beta + \gamma}{3} = p$$

$$\text{and } y = \frac{\frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma}}{3}$$

$$= \frac{\beta\gamma + \gamma\alpha + \alpha\beta}{3\alpha\beta\gamma} = q$$

Hence, coordinates of the centroid of triangle are (p, q)

- 9 **(d)**
Let $O(0, 0)$ be the orthocenter, $A(h, k)$ be the third vertex and $B(-2, 3)$ and $C(5, -1)$ the other two vertices. Then, the slope of the line through A and O is $\frac{k}{h}$, while the line through B and C has the slope $\frac{(-1-3)}{(5+2)} = -\frac{4}{7}$. By the property of the

orthocenter, these two lines must be perpendicular, so we have

$$\left(\frac{k}{h}\right)\left(-\frac{4}{7}\right) = -1 \Rightarrow \frac{k}{h} = \frac{7}{4} \dots(i)$$

$$\text{Also, } \frac{5-2+h}{3} + \frac{-1+3+k}{3} = 7$$

$$\Rightarrow h + k = 16 \dots(ii)$$

Which is not satisfied by the points given in the options (a), (b) or (c)

- 10 **(b)**
Let (h, k) be the point
According to question,
 $4\sqrt{(h-h)^2 + k} = h^2 + k^2$
 $\Rightarrow 4|k| = h^2 + k^2$
Locus of the point is
 $4|y| = x^2 + y^2 \Rightarrow x^2 + y^2 - 4|y| = 0$

- 12 **(a)**
Given points are $P(4, -2)$, $A(2, -4)$ and $B(7, 1)$
Suppose P divides AB in the ratio $\lambda : 1$. Then,
 $\frac{7\lambda + 2}{\lambda + 1} = 4 \Rightarrow \lambda = \frac{2}{3}$

Thus, P divides AB internally in the ratio $2 : 3$
The coordinates of the point dividing AB

$$\text{externally in the ratio } 2 : 3 \text{ are } \left(\frac{2 \times 7 - 3 \times 2}{2 - 3}, \frac{2 \times 1 - 3 \times -4}{2 - 3}\right) = (-8, -14)$$

Hence, the harmonic conjugate of R with respect to A and B is $(-8, -14)$

- 13 **(c)**
If O is the origin and $P(x_1, y_1)$, $Q(x_2, y_2)$ are two points, then
 $OP \times OQ \cos \angle POQ = x_1x_2 + y_1y_2$
 $\therefore OP \times OQ \times \sin \angle POQ$
 $= \sqrt{OP^2 \times OQ^2 - OP^2 \times OQ^2 \times \cos^2 \angle POQ}$
 $= \sqrt{(x_1^2 + y_1^2)(x_2^2 + y_2^2) - (x_1x_2 + y_1y_2)^2}$
 $= \sqrt{(x_1y_2 - x_2y_1)^2} = |x_1y_2 - x_2y_1|$

- 14 **(c)**

$$\cos B = \frac{(3)^2 + (5)^2 - (4)^2}{2 \times 3 \times 5} = \frac{3}{5}$$

$$\Rightarrow \sin B = \sqrt{1 - \frac{9}{25}} = \frac{4}{5}$$

$$\therefore \sin 2B = 2 \sin B \cos B$$

$$= 2 \times \frac{4}{5} \times \frac{3}{5} = \frac{24}{25}$$

- 16 **(d)**
Given that, $\angle A = 45^\circ$, $\angle B = 75^\circ$
 $\angle C = 180^\circ - 45^\circ - 75^\circ = 60^\circ$

$$\therefore a + c\sqrt{2} = k(\sin A + \sqrt{2} \sin C)$$

$$= k(\sin 45^\circ + \sqrt{2} \sin 60^\circ)$$

$$= k\left(\frac{1}{\sqrt{2}} + \sqrt{2} \frac{\sqrt{3}}{2}\right) = k\left(\frac{1+\sqrt{3}}{\sqrt{2}}\right) \dots(i)$$


$$\text{And } k = \frac{b}{\sin B}$$

$$= \frac{b}{\sin 75^\circ} = \frac{2\sqrt{2}b}{\sqrt{3} + 1}$$

On putting the value of k in Eq. (i), we get

$$a + c\sqrt{2} = 2b$$

- 18 **(d)**
From figure $ABCD$ is a square




Whose diagonals AC and BD are of length 2 unit

$$\text{Hence, required area} = \frac{1}{2} AC \times BD$$

$$= \frac{1}{2} \times 2 \times 2 = 2 \text{ sq units}$$

- 19 **(c)**
In $\triangle APD$,
 $\tan 45^\circ = \frac{a}{AP} \Rightarrow AP = a$



and in $\triangle BPC$,

$$\tan 45^\circ = \frac{b}{PB}$$

$$\Rightarrow PB = b$$

$$\therefore DE = a + b \text{ and } CE = b - a$$

In $\triangle DEC$,

$$DC^2 = DE^2 + EC^2$$

$$= (a + b)^2 + (b - a)^2$$

$$= 2(a^2 + b^2)$$

20 (b)

If the axes are rotated through 30° , we have

$$x = X \cos 30^\circ - Y \sin 30^\circ = \frac{\sqrt{3}X - Y}{2}$$

$$\text{and, } y = X \sin 30^\circ + Y \cos 30^\circ = \frac{X + \sqrt{3}Y}{2}$$


Substituting these values in $x^2 + 2\sqrt{3}xy - y^2 = 2a^2$, we get

$$(\sqrt{3}X - Y)^2 + 2\sqrt{3}(\sqrt{3}X - Y)(X + \sqrt{3}Y) - (X + \sqrt{3}Y)^2 = 8a^2$$

$$\Rightarrow X^2 - Y^2 = a^2$$

21 (c)

Since, F, E and D are the mid points of the sides AB, AC and BC of triangle ABC respectively, then the vertices of triangle are $A(0, 0), B(0, 2), C(2, 0)$



$$\text{Now, } AB = c = \sqrt{0^2 + 2^2} = 2$$

$$BC = a = \sqrt{2^2 + 2^2} = 2\sqrt{2}$$

$$\text{And } CA = b = \sqrt{2^2 + 0^2} = 2$$

\therefore x coordinates of incentre

$$= \frac{ax_1 + bx_2 + cx_3}{a + b + c}$$

$$= \frac{2\sqrt{2}(0) + 2(0) + 2(2)}{2\sqrt{2} + 2 + 2}$$

$$= \frac{2}{2 + \sqrt{2}} = 2 - \sqrt{2}$$

22 (a)

Let (x, y) be the coordinates of vertex C and (x_1, y_1) be the coordinates of centroid of the triangle.

$$\therefore x_1 = \frac{x+2-2}{3} \text{ and } y_1 = \frac{y-3+1}{3}$$

$$\Rightarrow x_1 = \frac{x}{3} \text{ and } y_1 = \frac{y-2}{3}$$

Since, the centroid lies on the line $2x + 3y = 1$

$$\therefore 2x_1 + 3y_1 = 1$$


$$\Rightarrow \frac{2x}{3} + \frac{3(y-2)}{3} = 1$$

$$\Rightarrow 2x + 3y = 9$$

This equation represents the locus of the vertex C

23 (c)

Let the height of the tower be h



$$\text{In } \Delta PAO, \tan 60^\circ = \frac{h}{OA}$$

$$\Rightarrow OA = h \cot 60^\circ = \frac{h}{\sqrt{3}} \quad \dots(i)$$

$$\text{In } \Delta PBO, \tan 30^\circ = \frac{h}{OB}$$

$$\Rightarrow OB = \frac{h}{\frac{1}{\sqrt{3}}}$$

$$\Rightarrow AB + AO = \sqrt{3}h$$

$$\Rightarrow 20 + \frac{h}{\sqrt{3}} = \sqrt{3}h \quad [\text{using Eq.(i)}]$$

$$\Rightarrow h = \frac{20}{\sqrt{3} - \frac{1}{\sqrt{3}}}$$

$$\Rightarrow h = \frac{20\sqrt{3}}{2}$$

$$\Rightarrow h = 10\sqrt{3} \text{ m}$$

24 (a)

$$\text{We have, } \sin \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{bc}}$$

$$\Rightarrow bc \sin^2 \frac{A}{2} = (s-b)(s-c)$$

$$\text{On comparing with } x \sin^2 \frac{A}{2} = (s-b)(s-c)$$

$$\text{We get, } x = bc$$

25 (c)

$$\therefore p_1^2 + p_2^2 = \frac{4a^2}{\sec^2 \alpha + \operatorname{cosec}^2 \alpha} + \frac{a^2 \cos^2 2\alpha}{\cos^2 \alpha + \sin^2 \alpha}$$

$$= a^2 \left(\frac{4 \cos^2 \alpha \sin^2 \alpha}{\cos^2 \alpha + \sin^2 \alpha} + \frac{\cos^2 2\alpha}{1} \right)$$

$$= a^2 (\sin^2 2\alpha + \cos^2 2\alpha) = a^2$$

$$\text{and } p_1^2 p_2^2 = a^4 \sin^2 2\alpha \cos^2 2\alpha = \left(\frac{1}{4}\right) a^4 \sin^2 4\alpha$$

$$\therefore \left(\frac{p_1}{p_2} + \frac{p_2}{p_1}\right)^2 = \frac{(p_1^2 + p_2^2)^2}{p_1^2 p_2^2}$$

$$= \frac{4}{\sin^2 4\alpha} = 4 \operatorname{cosec}^2 4\alpha$$

26 (c)

$$\text{Let } A(2, 1), B(-2, 4)$$

$$\therefore AB = 5$$

Hence, the locus is the line segment AB

27 (b)

$$\frac{1}{r^2} + \frac{1}{r_1^2} + \frac{1}{r_2^2} + \frac{1}{r_3^2}$$

$$= \frac{s^2 + (s-a)^2 + (s-b)^2 + (s-c)^2}{\Delta^2}$$

$$= \frac{4s^2 + a^2 + b^2 + c^2 - 2s(a + b + c)}{\Delta^2}$$

$$= \frac{a^2 + b^2 + c^2}{\Delta^2}$$

28 (b)

Let $a = 4$ cm, $b = 5$ cm and $c = 6$ cm

$$\therefore s = \frac{4 + 5 + 6}{2} = \frac{15}{2}$$

Hence, area of triangle = $\sqrt{s(s-a)(s-b)(s-c)}$

$$= \sqrt{\left(\frac{15}{2}\right)\left(\frac{15}{2} - 4\right)\left(\frac{15}{2} - 5\right)\left(\frac{15}{2} - 6\right)}$$

$$= \sqrt{\frac{15}{2} \times \frac{7}{2} \times \frac{5}{2} \times \frac{3}{2}} = \frac{15}{4} \sqrt{7} \text{ cm}^2$$

30 (a)


Let CD be the tower

In $\triangle ACM$, $\tan \beta = \frac{b}{AC}$

$$\Rightarrow AC = b \cot \beta$$

and in $\triangle ADC$, $\tan \alpha = \frac{CD}{AC}$

$$\Rightarrow CD = b \cot \beta \tan \alpha$$



31 (c)

$$\text{Since, } R = \frac{b}{2 \sin B} = \frac{2}{2 \sin 30^\circ} = \frac{2}{1}$$

Area of circumcircle = πR^2

$$= \pi \times (2)^2 = 4\pi \text{ sq unit}$$

33 (b)

$$\frac{b^2 - c^2}{2aR} = \frac{4R^2(\sin^2 B - \sin^2 C)}{4R^2 \sin A}$$


$$= \frac{\sin(B+C) \sin(B-C)}{\sin A}$$

$$= \sin(B-C)$$

34 (a)

Given curve is

$$y = 1 - |x|$$



\therefore Area of $\triangle ABC = 2$ area of $\triangle AOB$

$$= 2 \times \frac{1}{2} \times 1 \times 1 = 1 \text{ sq unit}$$

35 (b)

Given, angles A, B, C of $\triangle ABC$ are in AP with d (common difference) = 15°

$$\therefore B = A + 15^\circ \text{ and } C = A + 30^\circ$$

Also, $A + B + C = 180^\circ$

$$\Rightarrow A + A + 15^\circ + A + 30^\circ = 180^\circ$$

$$\Rightarrow \angle A = 45^\circ$$

$$\therefore \angle B = 45^\circ + 15^\circ = 60^\circ$$

36 (a)

$$\text{Since, } \left(1 - \frac{r_1}{r_2}\right) \left(1 - \frac{r_1}{r_3}\right) = 2$$

$$\therefore \left(1 - \frac{s-b}{s-a}\right) \left(1 - \frac{s-c}{s-a}\right) = 2$$

$$\Rightarrow \frac{(b-a)(c-a)}{(s-a)^2} = 2$$

$$\Rightarrow 2bc - 2ab - 2ac + 2a^2$$

$$= b^2 + c^2 + a^2 + 2bc - 2ab - 2ac$$

$$\Rightarrow a^2 = b^2 + c^2$$

So, triangle is right angled

37 (b)


Let the height of the tower be $BC = h$, then length of shadow of tower $AB = \sqrt{3}h$.

In $\triangle ABC$, $\tan \alpha = \frac{BC}{AB}$

$$\Rightarrow \tan \alpha = \frac{h}{\sqrt{3}h}$$

$$\Rightarrow \tan \alpha = \tan 30^\circ$$

$$\Rightarrow \alpha = 30^\circ$$



38 (d)

Let the coordinates of P be (h, k) . Then,

$$x = X + h, y = Y + k$$

Substituting these in $2x^2 + y^2 - 4x - 4y = 0$, we get

$$2X^2 + Y^2 + 4(h-1)X + 2(k-2)Y + 2h^2 + k^2 - 4h - 4k = 0$$

Comparing this equation with

$$2X^2 + Y^2 - 8X - 8Y + 18 = 0, \text{ we get}$$

$$h-1 = -2, (k-2) = -4 \text{ and } 2h^2 + k^2 - 4h - 4k = 18$$

$$\Rightarrow h = -1, k = -2$$

40 (a)

$$\text{We have, } \tan \frac{C}{2} = \sqrt{\frac{(s-a)(s-b)}{s(s-c)}}$$

$$\because (s-a)(s-b) = s(s-c) \text{ (given)}$$

$$\therefore \tan \frac{C}{2} = \sqrt{\frac{s(s-c)}{s(s-c)}}$$


$$\Rightarrow \tan \frac{C}{2} = \tan \frac{\pi}{4}$$

$$\Rightarrow \angle C = 90^\circ$$

41 (a)

$$\text{Given that, } \cot \alpha = \frac{3}{5} \text{ and } \cot \beta = \frac{2}{5}$$

$$\text{In } \triangle BCD, \tan \beta = \frac{h}{BC}$$



$$\Rightarrow BC = h \cot \beta \Rightarrow BC = \frac{2h}{5} \dots(i)$$

$$\text{and in } \triangle ACD, \tan \alpha = \frac{h}{32+BC}$$

$$\Rightarrow h = \left(32 + \frac{2h}{5}\right) \frac{5}{3} \text{ [using Eq.(i)]}$$

$$\Rightarrow 3h = 160 + 2h$$

$$\Rightarrow h = 160 \text{ m}$$

42 (a)

The vertices of quadrilateral ABCD is

$$A(2, 3), B(3, 4), C(4, 5)$$

$$\text{and } D(5, 6)$$

$$\therefore AB = \sqrt{(3-2)^2 + (4-3)^2}$$

$$= \sqrt{(1)^2 + (1)^2} = \sqrt{2}$$

$$\text{Similarly, } BC = \sqrt{2}, CD = \sqrt{2}, \text{ and } DA = 3\sqrt{2}$$

$$\therefore a = b = c = \sqrt{2} \text{ and } d = 3\sqrt{2}$$

$$\text{and } s = \frac{a+b+c+d}{2}$$

$$= \frac{\sqrt{2} + \sqrt{2} + \sqrt{2} + 3\sqrt{2}}{2}$$

$$= \frac{6\sqrt{2}}{2} = 3\sqrt{2}$$

$$\therefore \text{Area of quadrilateral}$$

$$= \sqrt{(s-a)(s-b)(s-c)(s-d)}$$

$$= \sqrt{(3\sqrt{2} - \sqrt{2})(3\sqrt{2} - \sqrt{2})}$$

$$(3\sqrt{2} - \sqrt{2})(3\sqrt{2} - \sqrt{2})$$

$$= 0$$

43 (c)

$$\text{We have, } \Delta = \frac{1}{2}bc \sin A$$

$$\Rightarrow \frac{1}{2}k^2 \sin B \sin C \sin A = \Delta \dots(i)$$

$$\therefore a^2 \sin 2B + b^2 \sin 2A$$

$$= 2(a^2 \sin B \cos B + b^2 \sin A \cos A)$$

$$= 2k^2(\sin^2 A \sin B \cos B + \sin^2 B \sin A \cos A)$$

$$= 2k^2(\sin A \sin B \sin C) = 4\Delta \text{ [from Eq. (i)]}$$

44 (c)

$$1. \frac{\sin A}{\sin C} = \frac{\sin(A-B)}{\sin(B-C)}$$

$$\Rightarrow \sin(B+C) \sin(B-C) = \sin(A+B) \sin(A-B)$$

$$\Rightarrow \sin^2 B - \sin^2 C = \sin^2 A - \sin^2 B$$

$$\Rightarrow b^2 - c^2 = a^2 - b^2$$

$$\Rightarrow 2b^2 = a^2 + c^2$$

$$\Rightarrow a^2, b^2, c^2 \text{ are in AP}$$

$$2. r_1, r_2, r_3 \text{ are in HP}$$

$$\Rightarrow \frac{1}{r_1}, \frac{1}{r_2}, \frac{1}{r_3} \text{ are in AP}$$

$$\Rightarrow \frac{2}{r_2} = \frac{1}{r_1} + \frac{1}{r_3}$$

$$\Rightarrow 2(s-b) = s-a + s-c$$

$$\Rightarrow 2b = (a+c)$$

$$\Rightarrow a, b, c \text{ are in AP}$$

Hence, both of these statements are correct

45 (c)

The largest side of triangle is $\sqrt{p^2 + q^2 + pq}$

Greatest angle will be opposite to largest side. Let

θ be greatest angle, then

$$\cos \theta = \frac{p^2 + q^2 - p^2 - q^2 - pq}{2pq} = -\frac{1}{2}$$

$$\Rightarrow \theta = \frac{2\pi}{3}$$

46 (a)


Let area of triangle be Δ , then according to question

$$\Delta = \frac{1}{2}ax = \frac{1}{2}by = \frac{1}{2}cz$$

$$\begin{aligned} \therefore \frac{bx}{c} + \frac{cy}{a} + \frac{az}{b} &= \frac{b}{c} \left(\frac{2\Delta}{a} \right) + \frac{c}{a} \left(\frac{2\Delta}{b} \right) + \frac{a}{b} \left(\frac{2\Delta}{c} \right) \\ &= \frac{2\Delta(b^2 + c^2 + a^2)}{abc} \\ &= \frac{2(a^2 + b^2 + c^2)}{abc} \cdot \frac{abc}{4R} \quad \left(\because \Delta = \frac{abc}{4R} \right) \\ &= \frac{a^2 + b^2 + c^2}{2R} \end{aligned}$$

49 (d)

In ΔABC the vertices are $A(-3, 0)$, $B(4, -1)$ and $C(5, 2)$



$$\begin{aligned} \therefore BC &= \sqrt{(5-4)^2 + (2+1)^2} \\ &= \sqrt{1+9} = \sqrt{10} \end{aligned}$$

Area of ΔABC


$$\begin{aligned} &= \frac{1}{2} [x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2)] \\ &= \frac{1}{2} [-3(-1 - 2) + 4(2 - 0) + 5(0 + 1)] \\ &= \frac{1}{2} [9 + 8 + 5] = 11 \end{aligned}$$

As we know that, area of $\Delta = \frac{1}{2} \times BC \times AL$

$$\begin{aligned} \Rightarrow 11 &= \frac{1}{2} \times \sqrt{10} \times AL \\ \Rightarrow AL &= \frac{2 \times 11}{\sqrt{10}} = \frac{22}{\sqrt{10}} \end{aligned}$$

50 (a)

As the line divides the ΔABC in equal to area. Mid point of $AB(51, 30)$ which lies on $y = kx$



$$\therefore 30 = 51k \Rightarrow k = \frac{30}{51}$$

52 (b)

Let (h, k) be the point

According to question, $4\sqrt{(h-h)^2 + k^2} = h^2 + k^2$


$$\Rightarrow 4|k| = h^2 + k^2$$

Locus of the point is $4|y| = x^2 + y^2$

$$\Rightarrow x^2 + y^2 - 4|y| = 0$$

53 (a)

In ΔCBD , $\tan 60^\circ = \frac{h}{x}$



$$\Rightarrow h = x\sqrt{3} \dots (i)$$

and in ΔCAD , $\tan 30^\circ = \frac{h}{40+x}$

$$\Rightarrow h\sqrt{3} = 40 + x$$

$$\Rightarrow 3x = 40 + x \text{ [from Eq. (i)]}$$

$$\Rightarrow x = 20 \text{ m}$$

54 (d)

(a) We know, $\tan A + \tan B + \tan C = \tan A \tan B \tan C$

Since, $\tan A + \tan B + \tan C = 0$

\Rightarrow Let either of $\tan A$, $\tan B$ or $\tan C$ is zero *ie*, one angle is 0

So, it cannot be a triangle

$$(b) \frac{\sin A}{2} = \frac{\sin B}{3} = \frac{\sin C}{1}$$

$$\Rightarrow a : b : c = 2 : 3 : 1$$

Let $a = 2k$, $b = 3k$, $c = k$, $a + c = b$ (so triangle not possible)

$$(c) \sin A \sin B = \frac{\sqrt{3}}{4} = \cos A \cos B$$

Either $\sin A$, $\sin B$ are both positive or both negative but, if both are positive $\sin A + \sin B > 0$ but $\sin A + \sin B$ is negative so both negative but, if both are negative, then $\angle A$ and $\angle B$ are more than 90° , so it cannot be a triangle

$$(d) (a + b)^2 = c^2 + ab$$

$$\Rightarrow a^2 + b^2 + 2ab = c^2 + ab$$

$$\Rightarrow a^2 + b^2 - c^2 = -ab$$

$$\Rightarrow \frac{a^2 + b^2 - c^2}{2ab} = -\frac{1}{2} = \cos C$$

$$\Rightarrow \angle C = 120^\circ$$

$$\sin A + \cos A = \frac{\sqrt{3}}{\sqrt{2}}$$


$$\Rightarrow 1 + \sin 2A = \frac{3}{2} \Rightarrow \sin 2A = \frac{1}{2}$$

$$\Rightarrow 2A = 30^\circ \Rightarrow \angle A = 15^\circ$$

So, it can form a triangle

55 (b)

$$\text{In } \Delta PMC, \tan \alpha = \frac{h-a}{PM}$$



$$\Rightarrow PM = (h-a) \cot \alpha \dots (i)$$

$$\text{In } \Delta PMC', \tan \beta = \frac{h+a}{PM}$$

$$\Rightarrow h+a = PM \tan \beta$$

$$\Rightarrow h = (h-a) \cot \alpha \tan \beta - a$$


$$\Rightarrow h(1 - \cot \alpha \tan \beta) = -a(1 + \cot \alpha \tan \beta)$$

$$\Rightarrow h = \frac{a(\sin \alpha \cos \beta + \cos \alpha \sin \beta)}{\sin \beta \cos \alpha - \sin \alpha \cos \beta}$$

$$= \frac{a \sin(\alpha + \beta)}{\sin(\beta - \alpha)} m$$

56 (a)

Line perpendicular to OA passing through B is $x = 4$




$$\text{Slope of } AB = -\frac{3}{2}$$

Line perpendicular to AB through origin is $y = \frac{2}{3}x$

\therefore The point of intersection of a line $x = 4$ and $y = \frac{2}{3}x$ is $(4, \frac{8}{3})$

57 (c)

Since, $(0, 1)$, $(1, 1)$ and $(1, 0)$ are mid points of sides AB , BC and CA respectively



\therefore Coordinates of A, B and C are $(0, 0)$, $(0, 2)$ and $(2, 0)$ respectively

Now, $AB = 2, BC = 2\sqrt{2}, CA = 2$

\therefore x -coordinate of incentre

$$= \frac{0 + 0 + 2 \cdot 2}{2 + 2\sqrt{2} + 2} \left(\because x = \frac{ax_1 + bx_2 + cx_3}{a + b + c} \right)$$

$$= \frac{2}{2 + \sqrt{2}} = 2 - \sqrt{2}$$

58 (d)

Let $P(x, y)$ is equidistant from the mid points $A(a+b, b-a)$ and $(a-b, a+b)$


$$\therefore PA^2 = PB^2$$

$$\Rightarrow (a+b-x)^2 + (b-a-y)^2 = (a-b-x)^2 + (a+b-y)^2$$

$$\Rightarrow bx - ay = 0$$

59 (a)

Let the locus of a point in a plane be $P(h, k)$



According to the question,

$$|PA| + |PB| = 1 \Rightarrow |h| + |k| = 1$$


Hence, locus of a point is

$$|x| + |y| = 1$$

Which represents the equation of square

60 (b)

Let BC be the height of tower and CD be height of the flagstaff



$$\text{In } \Delta BAC, \tan \theta = \frac{x}{y} \dots (i)$$

$$\text{In } \Delta DAB, \tan 2\theta = \frac{x+h}{y}$$

$$\Rightarrow \frac{2 \tan \theta}{1 - \tan^2 \theta} = \frac{x+h}{y} \Rightarrow \frac{2(\frac{x}{y})}{1 - \frac{x^2}{y^2}} = \frac{x+h}{y} \text{ [from Eq. (i)]}$$

$$\Rightarrow 2xy^2 - xy^2 + x^3 = (y^2 - x^2)h$$

$$\Rightarrow h = \frac{x(x^2 + y^2)}{(y^2 - x^2)}$$

61 (b)

$$2ac \sin\left(\frac{A-B+C}{2}\right)$$

$$= 2ac \sin\left(\frac{180^\circ - 2B}{2}\right)$$

$$= 2ac \sin(90^\circ - B) = 2ac \cos B = a^2 + c^2 - b^2$$

62 (d)

$$(x+1)^2 + y^2 + (x-2)^2 + y^2 = 2[(x-1)^2 + y^2]$$

On simplification, we get $2x + 3 = 0$

63 (a)

We know that centroid divides the line segment joining orthocenter and circumcentre in the ratio 2:1. Since, the coordinates of orthocenter and circumcentre are (1, 1) and (3, 2) respectively

∴ The coordinates of centroid are

$$\left(\frac{2.3 + 1.1}{2 + 1}, \frac{2.2 + 1.1}{2 + 1}\right) = \left(\frac{7}{3}, \frac{5}{3}\right)$$

64 (c)

Given equation are

$$x \cot \theta + y \operatorname{cosec} \theta = 2 \quad \dots(i)$$

$$\text{And } x \operatorname{cosec} \theta + y \cot \theta = 6 \quad \dots(ii)$$

On squaring and subtracting Eq. (i) from Eq. (ii), we get

$$x^2(\operatorname{cosec}^2 \theta - \cot^2 \theta) + y^2(\cot^2 \theta - \operatorname{cosec}^2 \theta) =$$

$$(6)^2 - (2)^2$$

$$\Rightarrow x^2 - y^2 = 32$$

It represents an equation of hyperbola

65 (c)

Since, $\cot A + \cot C = 2 \cot B$

$$\Rightarrow \frac{\cos A}{\sin A} + \frac{\cos C}{\sin C} = \frac{2 \cos B}{\sin B}$$

$$\Rightarrow \frac{b^2 + c^2 - a^2}{2bc(ka)} + \frac{a^2 + b^2 - c^2}{2ab(kc)} = 2 \frac{a^2 + c^2 - b^2}{2ac(kb)}$$


$$\Rightarrow a^2 + c^2 = 2b^2$$

Hence, a^2, b^2, c^2 are in AP

66 (a)

In $\triangle ABC$, $AC = (H - h) \cot 15^\circ \dots (i)$

and in $\triangle ACD$, $AC = (H + h) \cot 45^\circ \dots (ii)$



From Eqs. (i) and (ii).

$$(H - h) \cot 15^\circ = (H + h) \cot 45^\circ$$

$$\Rightarrow H = \frac{h(\cot 15^\circ + 1)}{\cot 15^\circ - 1}$$

$$\therefore H = \frac{2500(2 + \sqrt{3} + 1)}{(2 + \sqrt{3} - 1)}$$

$$= \frac{2500(3 + \sqrt{3})}{(\sqrt{3} + 1)} \times \frac{(\sqrt{3} - 1)}{(\sqrt{3} - 1)}$$

$$= 2500\sqrt{3} \text{ m}$$


67 (b)

Let OP be the clock tower standing at the mid point O of side BC of $\triangle ABC$. Let $\alpha = \angle PAO = \cot^{-1} 3.2$ and $\beta = \angle PBO = \operatorname{cosec}^{-1} 2.6$

Then, $\cot \alpha = 3.2$ and $\operatorname{cosec} \beta = 2.6$

$$\therefore \cot \beta = \sqrt{\operatorname{cosec}^2 \beta - 1} = \sqrt{(2.6)^2 - 1} = 2.4$$

In $\triangle PAO$ and $\triangle PBO$, we have



$$AO = h \cot \alpha = 3.2h$$

$$\text{and } BO = h \cot B = 2.4h$$

$$\text{In } \triangle ABO, AB^2 = OA^2 + OB^2$$

$$\Rightarrow 100^2 = (3.2h)^2 + (2.4h)^2$$

$$\Rightarrow 100^2 = 16h^2$$

$$\Rightarrow h^2 = 625 \Rightarrow h = 25 \text{ m}$$

68 (d)

$$\because \cos 30^\circ = \frac{3 + 1 - a^2}{2\sqrt{3}} \left[\because \cos A = \frac{b^2 + c^2 - a^2}{2bc} \right]$$

$$\Rightarrow \frac{\sqrt{3}}{2} = \frac{4 - a^2}{2\sqrt{3}} \Rightarrow a^2 = 1$$

$$\Rightarrow a = 1$$

Here, we see side b is largest, so

$\angle B$ must be greatest

$$\therefore \text{By sine rule, } \frac{b}{\sin B} = \frac{a}{\sin A}$$

$$\Rightarrow \frac{\sqrt{3}}{\sin B} = \frac{1}{\sin 30^\circ}$$

$$\Rightarrow \sin B = \frac{\sqrt{3}}{2}$$

$$\Rightarrow \angle B = 120^\circ$$

69 (c)

Let each side of equilateral triangle = a

$$\therefore \Delta = \frac{\sqrt{3}}{4} a^2, \quad S = \frac{3a}{2}$$

$$\text{Now, } r = \frac{\Delta}{S} = \frac{\frac{\sqrt{3}}{4} a^2}{\frac{3a}{2}} = \frac{a}{2\sqrt{3}}$$

$$R = \frac{abc}{4\Delta} = \frac{a^3}{\sqrt{3}a^2} = \frac{a}{\sqrt{3}}$$

$$r_1 = \frac{\Delta}{s - a} = \frac{\frac{\sqrt{3}}{4} a^2}{\frac{2}{a}} = \frac{\sqrt{3}}{2} a$$

$$\therefore R:r_1:r_2 = \frac{a}{\sqrt{3}} : \frac{a}{2\sqrt{3}} : \frac{\sqrt{3}}{2}a$$

$$= 2:1:3$$

70 (d)

Given, $r_1 = 2r_2 = 3r_3$

$$\therefore \frac{\Delta}{s-a} = \frac{2\Delta}{s-b} = \frac{3\Delta}{s-c} = \frac{\Delta}{k} \text{ [say]}$$

Then, $s-a = k, s-b = 2k, s-c = 3k$

$$\Rightarrow 3s - (a+b+c) = 6k \Rightarrow s = 6k$$

$$\therefore \frac{a}{5} = \frac{b}{4} = \frac{c}{3} = k$$

$$\therefore \frac{a}{b} + \frac{b}{c} + \frac{c}{a} = \frac{5}{4} + \frac{4}{3} + \frac{3}{5} = \frac{191}{60}$$

71 (c)

Let sides of the triangle are $4x, 5x, 6x$

$$s = \frac{4x + 5x + 6x}{2} = \frac{15}{2}x$$

$$\Delta = \sqrt{\frac{15}{2}x \left(\frac{15}{2}x - 4x\right) \left(\frac{15}{2}x - 5x\right) \left(\frac{15}{2}x - 6x\right)}$$

$$= \sqrt{\frac{15}{2}x \times \frac{7}{2}x \times \frac{5}{2}x \times \frac{3}{2}x}$$

$$= \frac{15\sqrt{7}x^2}{4}$$

Circumradius, $R = \frac{4x \times 5x \times 6x}{4 \times \frac{15\sqrt{7}x^2}{4}}$

$$= \frac{8}{\sqrt{7}}x$$


Inradius, $r = \frac{\frac{15\sqrt{7}x^2}{4}}{\frac{15}{2}x}$

$$= \frac{\sqrt{7}}{2}x$$

$$\frac{R}{r} = \frac{\frac{8x}{\sqrt{7}}}{\frac{\sqrt{7}x}{2}} = \frac{16}{7}$$

72 (c)

In $\triangle DAP$, $\tan 60^\circ = \frac{1}{AP}$ [$\because EQ = DP = 1$]



$$\Rightarrow AP = \frac{1}{\sqrt{3}}$$

In $\triangle EAQ$, $\tan 30^\circ = \frac{EQ}{AP+PQ}$

$$\Rightarrow \frac{1}{\sqrt{3}} + PQ = \sqrt{3}$$

$$\Rightarrow PQ = \frac{2}{\sqrt{3}} \text{ km}$$

$$\therefore \text{Speed of plane} = \frac{\text{Distance}}{\text{Time}}$$

$$= \frac{\frac{2}{\sqrt{3}}}{\frac{10}{60 \times 60}} = 240\sqrt{3} \text{ km/h}$$

74 (a)

$$\text{Area of triangle} = \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix}$$

= A rational number if vertices have integral coordinates only

If triangle is equilateral, then area

$$= \frac{\sqrt{3}}{4} [(x_1 - x_2)^2 + (y_1 - y_2)^2]$$

= irrational quantity

So, triangle cannot be equilateral

75 (d)

$$\frac{b-c}{a} = \frac{k(\sin B + \sin C)}{k \sin A}$$


$$= \frac{2 \sin \left(\frac{B+C}{2}\right) \cos \left(\frac{B-C}{2}\right)}{2 \sin \frac{A}{2} \cos \frac{A}{2}}$$

$$\Rightarrow \frac{b+c}{a} = \frac{\cos \left(\frac{B-C}{2}\right)}{\sin \frac{A}{2}}$$

Similarly, $\frac{b-c}{a} = \frac{\sin \left(\frac{B-C}{2}\right)}{\cos \frac{A}{2}}$

76 (b)

Given, $\theta = \tan^{-1} \frac{3}{5} \Rightarrow \tan \theta_2 = \frac{3}{5}$... (i)



In $\triangle AOC$, $\tan \theta_1 = \frac{AC}{AO} = \frac{h}{160}$... (ii)

and in $\triangle AOB$, $\tan(\theta_1 + \theta_2) = \frac{h}{40}$

$$\Rightarrow \frac{\tan \theta_1 + \tan \theta_2}{1 - \tan \theta_1 \tan \theta_2} = \frac{h}{40}$$

$$\Rightarrow \frac{\frac{h}{160} + \frac{3}{5}}{1 - \frac{h}{160} \times \frac{3}{5}} = \frac{h}{40} \text{ [from Eqs. (i) and (ii)]}$$

$$\Rightarrow \frac{5[h + 96]}{800 - 3h} = \frac{h}{40}$$

$$\Rightarrow h^2 - 200h + 6400 = 0$$

$$\Rightarrow (h - 160)(h - 40) = 0$$

$$\Rightarrow h = 160 \text{ or } h = 40$$

Hence, height of the vertical pole is 40 m

77 (c)

Let $A(x_1, y_1)$ and $B(x_2, y_2)$ be two points. Let C and D be the points of internal and external division of AB in the ratio $\lambda : 1$. Then, the coordinates of C and D are

$$\left(\frac{\lambda x_2 + x_1}{\lambda + 1}, \frac{\lambda y_2 + y_1}{\lambda + 1}\right) \text{ and } \left(\frac{\lambda x_2 - x_1}{\lambda - 1}, \frac{\lambda y_2 - y_1}{\lambda - 1}\right) \text{ respectively}$$

$$\therefore AC = \frac{\lambda}{\lambda + 1} AB \text{ and } AD = \frac{\lambda}{\lambda - 1} AB$$

Clearly, $\frac{1}{AC} + \frac{1}{AD} = \frac{2}{AB} \Rightarrow AC, AB, AD$ are in H.P.

79 (d)

We have,

$$A + B + C = 180^\circ$$

$$\Rightarrow 3B = 180 \quad [\because A, B, C \text{ are in AP}]$$

$$\Rightarrow B = 60^\circ$$

$$\Rightarrow \cos B = \frac{1}{2}$$

$$\Rightarrow \frac{AB^2 + BC^2 - AC^2}{2AB \cdot BC} = \frac{1}{2}$$

$$\Rightarrow 36 + 49 - AC^2 = 6 \times 7 \Rightarrow AC^2 = 43 \Rightarrow AC = \sqrt{43}$$


80 (a)

Let the point be $P(h, k)$

It is given that difference of the distance from points $A(3, 0)$

and $B(-3, 0)$ is 4 i.e., $PA - PB = 4$

$$\Rightarrow \sqrt{(h-3)^2 + k^2} - \sqrt{(h+3)^2 + k^2} = 4$$



$$\Rightarrow \sqrt{(h-3)^2 + k^2} = 4 + \sqrt{(h+3)^2 + k^2}$$

On squaring both sides, we get

$$(h-3)^2 + k^2 = 16 + (h+3)^2 + k^2 + 8\sqrt{(h+3)^2 + k^2}$$

$$\Rightarrow h^2 + 9 - 6h + k^2 = 16 + h^2 + 9 + 6h + k^2 + 8\sqrt{(h+3)^2 + k^2}$$

$$\Rightarrow -6h = 16 + 6h + 8\sqrt{(h+3)^2 + k^2}$$

$$\Rightarrow -8\sqrt{(h+3)^2 + k^2} = 12h + 16$$

Again, squaring both sides, we get

$$64(h+3)^2 + k^2 = (12h+16)^2$$

$$\Rightarrow 64(h^2 + 9 + 6h + k^2) = 144h^2 + 256 + 2 \cdot 16 \cdot 12h$$

$$\Rightarrow 64(h^2 + 9 + 6h + k^2) = 16(9h^2 + 16 + 24h)$$

$$\Rightarrow 4(h^2 + 9 + 6h + k^2) = 9h^2 + 16 + 24h$$

$$\Rightarrow 4h^2 + 36 + 24h + 4k^2 = 9h^2 + 16 + 24h$$

$$\Rightarrow 5h^2 - 4k^2 = 20$$

$$\Rightarrow \frac{h^2}{4} - \frac{k^2}{5} = 1$$

Hence, the locus of points P is $\frac{x^2}{4} - \frac{y^2}{5} = 1$

81 (b)

$$\because \cos C = \frac{a^2 + b^2 - c^2}{2ab}$$

$$\Rightarrow \cos^2 C = \left(\frac{a^2 + b^2 - c^2}{2ab}\right)^2$$

$$\Rightarrow \cos^2 C$$

$$= \frac{[a^4 + b^4 + c^4 + 2a^2b^2 - 2c^2(a^2 + b^2)]}{4a^2b^2}$$

$$\Rightarrow \cos^2 C = \frac{1}{2} [\because a^4 + b^4 + c^4$$

$$= 2c^2(a^2 + b^2) \text{ given}]$$

$$\Rightarrow \cos C = \pm \frac{1}{\sqrt{2}}$$

$$\Rightarrow \angle C = 45^\circ \text{ or } 135^\circ$$

82 (d)

Locus, of P is $|\sqrt{x^2 + y^2 - 8y + 16} -$

$$\sqrt{x^2 + y^2 + 8y + 16}| = 6$$

On squaring, we get

$$x^2 + y^2 - 2$$

$$= \sqrt{x^2 + y^2 + 8y + 16} \sqrt{x^2 + y^2 - 8y + 16}$$

$$\Rightarrow (x^2 + y^2 - 2)^2 = (x^2 + y^2 + 16)^2 - (8y)^2$$

On simplification, we get

$$\frac{y^2}{9} - \frac{x^2}{7} = 1$$

83 (a)

Given, $\sin \frac{A}{2} \sin \frac{C}{2} = \sin \frac{B}{2}$

$$\Rightarrow \sqrt{\frac{(s-b)(s-c)}{bc}} \sqrt{\frac{(s-a)(s-b)}{ab}}$$

$$= \sqrt{\frac{(s-a)(s-c)}{ac}}$$

$$\Rightarrow \frac{s-b}{b} = 1$$


$$\Rightarrow s = 2b$$

84 (c)

In ΔPRQ


$$\tan 60^\circ = \frac{30}{x}$$

$$\Rightarrow x = 10\sqrt{3} \text{ m}$$



85 (a)

$$\text{In } \triangle ABD, \tan \alpha = \frac{h}{2h}$$



$$\Rightarrow \tan \alpha = \frac{1}{2} \quad \dots(i)$$

$$\text{In } \triangle ABC, \tan 2\alpha = \frac{h+p}{2h}$$

$$\Rightarrow \frac{2 \tan \alpha}{1 - \tan^2 \alpha} = \frac{h+p}{2h}$$

$$\Rightarrow \frac{2 \left(\frac{1}{2}\right)}{1 - \left(\frac{1}{2}\right)^2} = \frac{h+p}{2h}$$


$$\Rightarrow \frac{4}{3} = \frac{h+p}{2h}$$

$$\Rightarrow 8h = 3h + 3p$$

$$\Rightarrow 5h = 3p \Rightarrow p = \frac{5h}{3} \text{ m}$$

86 (d)

$$\text{In } \triangle ABD, \tan \beta = \frac{60}{d}$$



$$\Rightarrow d = 60 \cot \beta \quad \dots(i)$$

$$\text{In } \triangle DEC, \tan \alpha = \frac{DC}{EC}$$

$$\Rightarrow DC = d \tan \alpha$$

$$\Rightarrow 60 - h = d \tan \alpha \quad (\because BC = EA = h)$$

$$\Rightarrow 60 - h = 60 \cot \beta \tan \alpha \quad [\text{from Eq.(i)}]$$

$$\Rightarrow h = 60 \left(1 - \frac{\cos \beta}{\sin \beta} \cdot \frac{\sin \alpha}{\cos \alpha}\right)$$

$$\Rightarrow h = \frac{60 \sin(\beta - \alpha)}{\cos \alpha \sin \beta}$$

$$\Rightarrow \frac{60 \sin(\beta - \alpha)}{x} = \frac{60 \sin(\beta - \alpha)}{\cos \alpha \sin \beta} \quad (\text{given})$$

$$\Rightarrow x = \cos \alpha \sin \beta$$

87 (c)

We know that, in triangle larger side has an larger angle opposite to it. Since, angles $\angle A, \angle B$ and $\angle C$ are in AP

$$\Rightarrow 2B = A + C$$

$$\because A + B + C = \pi$$

$$\Rightarrow B = 60^\circ$$

$$\therefore \cos B = \frac{a^2 + c^2 - b^2}{2ac}$$

$$\Rightarrow \cos 60^\circ = \frac{1}{2} = \frac{100 + a^2 - 81}{20a}$$

$$\Rightarrow a^2 + 19 = 10a$$

$$\Rightarrow a^2 - 10a + 19 = 0$$

$$\therefore a = \frac{10 \pm \sqrt{100 - 76}}{2} = 5 \pm \sqrt{6}$$

88 (c)

Here, $\frac{1}{\sin^2 \frac{A}{2}}, \frac{1}{\sin^2 \frac{B}{2}}, \frac{1}{\sin^2 \frac{C}{2}}$ are in AP

$$\Rightarrow \frac{1}{\sin^2 \frac{C}{2}} - \frac{1}{\sin^2 \frac{B}{2}} = \frac{1}{\sin^2 \frac{B}{2}} - \frac{1}{\sin^2 \frac{A}{2}}$$

$$\Rightarrow \frac{ab}{(s-a)(s-b)} - \frac{ac}{(s-a)(s-c)}$$

$$= \frac{ac}{(s-a)(s-c)} - \frac{bc}{(s-b)(s-c)}$$

$$\Rightarrow \left(\frac{a}{s-a}\right) \left(\frac{b(s-c) - c(s-b)}{(s-b)(s-c)}\right)$$

$$= \left(\frac{c}{s-c}\right) \left(\frac{a(s-b) - b(s-a)}{(s-a)(s-b)}\right)$$

$$\Rightarrow ab + bc = 2ac \Rightarrow \frac{1}{c} + \frac{1}{a} = \frac{2}{b}$$

Hence, a, b, c are in HP

90 (c)

Let the vertices of triangle be $P(1, 1), Q(-1, -1)$ and $R(-\sqrt{3}, \sqrt{3})$

$$\therefore PQ = \sqrt{(1+1)^2 + (1+1)^2} = 2\sqrt{2}$$

$$QR = \sqrt{(-\sqrt{3}+1)^2 + (\sqrt{3}+1)^2}$$

$$= \sqrt{3+1-2\sqrt{3}+3+1+2\sqrt{3}} = 2\sqrt{2}$$

$$\text{and } RP = \sqrt{(-\sqrt{3}-1)^2 + (\sqrt{3}-1)^2}$$

$$= \sqrt{3+1+2\sqrt{3}+3+1-2\sqrt{3}}$$

$$= 2\sqrt{2}$$

$$\Rightarrow PQ = QR = RP$$

\therefore Triangle is an equilateral triangle

91 (a)

Let the third vertex be (a, b)

$$\therefore \text{Area of } \Delta = \frac{1}{2} \begin{vmatrix} 0 & 0 & 1 \\ a & b & 1 \\ 6 & 8 & 1 \end{vmatrix} = \frac{1}{2} |[8a - 6b]|$$

As (a, b) are integers, so we take

$(0, 0), (1, 1), (1, 2)$

At $(0, 0), \Delta = 0$, it is not possible

At $(1, 1) \Delta = 1$

At $(1, 2), \Delta = 2$


Here, we see that minimum area is 1

92 (d)

$$\begin{aligned} & \left(\cot \frac{A}{2} + \cot \frac{B}{2} \right) \left(a \sin^2 \frac{B}{2} + b \sin^2 \frac{A}{2} \right) \\ &= \left(\frac{\cos \frac{A}{2} + \sin \frac{B}{2} + \cos \frac{B}{2} + \sin \frac{A}{2}}{\sin \frac{A}{2} \sin \frac{B}{2}} \right) \left(a \sin^2 \frac{B}{2} \right. \\ & \quad \left. + b \sin^2 \frac{A}{2} \right) \\ &= \frac{\sin \left(\frac{A+B}{2} \right) \left(a \sin^2 \frac{B}{2} + b \sin^2 \frac{A}{2} \right)}{\sin \frac{A}{2} \sin \frac{B}{2}} \\ &= \left(\cos \frac{C}{2} \right) \left(a \frac{\sin \frac{B}{2}}{\sin \frac{A}{2}} + b \frac{\sin \frac{A}{2}}{\sin \frac{B}{2}} \right) \\ &= \sqrt{\frac{s(s-c)}{ab}} \left(a \sqrt{\frac{(s-a)(s-c)}{ac}} + b \sqrt{\frac{(s-b)(s-c)}{bc}} \right) \\ &= \sqrt{\frac{s(s-c)}{ab}} \left(\sqrt{\frac{(s-a)}{(s-b)}} ab + \sqrt{\frac{(s-b)}{(s-a)}} ab \right) \\ &= \sqrt{s(s-c)} \left(\frac{s-a+s-b}{\sqrt{(s-a)(s-b)}} \right) \\ &= \sqrt{s(s-c)} \left(\frac{2s-a-b}{\sqrt{(s-a)(s-b)}} \right) \\ &= c \sqrt{\frac{s(s-c)}{(s-a)(s-b)}} = c \cot \frac{C}{2} \end{aligned}$$

93 (c)

Given lines $y = mx, y = 2, y = 6$



Coordinates of points A and B are $\left(\frac{2}{m}, 2\right), \left(\frac{6}{m}, 6\right)$ respectively

$$\therefore AB = \sqrt{\left(\frac{2}{m} - \frac{6}{m}\right)^2 + (2-6)^2} < 5 \quad [\text{given}]$$

$$\Rightarrow \left(\frac{2}{m} - \frac{6}{m}\right)^2 + (4)^2 < 25$$

$$\Rightarrow \left(\frac{2}{m} - \frac{6}{m}\right)^2 < 9 \Rightarrow -3 < \frac{2}{m} - \frac{6}{m} < 3$$

$$\Rightarrow -\frac{4}{3} > m > \frac{4}{3}$$

$$\therefore m \in \left] -\infty, -\frac{4}{3} \cup \frac{4}{3}, \infty \right[$$

94 (c)

By sine rule

$$\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c} \quad (\text{say})$$

$$\therefore (b-c) \sin A + (c-a) \sin B + (a-b) \sin C$$

$$= (b-c)ak + (c-a)bk + (a-b)kc$$

$$= k[ab - ac + bc - ab + ac - bc]$$

$$= 0$$

95 (a)

Given that, $a = 3, b = 5, c = 6$

$$\text{Now, } s = \frac{a+b+c}{2} = 7$$

$$\therefore \Delta = \sqrt{s(s-a)(s-b)(s-c)}$$

$$= \sqrt{7(7-3)(7-5)(7-6)}$$

$$= \sqrt{7 \cdot 4 \cdot 2 \cdot 1} = 2\sqrt{14}$$

$$\therefore r = \frac{\Delta}{s} = \frac{2\sqrt{14}}{7} = \sqrt{\frac{8}{7}}$$

96 (a)

$$\cos B = \frac{3^2 + 4^2 - 5^2}{2(3)(4)} = \frac{9 + 16 - 25}{2(3)(4)} = 0$$


$$\Rightarrow \angle B = 90^\circ$$

$$\therefore \sin \frac{B}{2} + \cos \frac{B}{2} = \sin 45^\circ + \cos 45^\circ$$

$$= \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} = \sqrt{2}$$

97 (b)

$$\text{In } \Delta CAD, \tan 30^\circ = \frac{CD}{AC}$$



$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{h}{120 + x}$$

$$\Rightarrow \sqrt{3}h = 120 + x \quad \dots(i)$$

$$\text{and in } \Delta CBD, \tan 60^\circ = \frac{CD}{BC}$$

$$\Rightarrow \sqrt{3} = \frac{h}{x} \Rightarrow h = \sqrt{3}x \dots(ii)$$

From Eqs. (i) and (ii), we get, $x = 60$ m

On putting $x = 60$ in Eq.(i), we get

$$h = 60\sqrt{3} \text{ m}$$

98 (c)

$$\text{Area of triangle} = \frac{1}{2}[x(1-2) + 1(2-0) + 0(0-1)]$$

$$= \frac{1}{2}[-x + 2 + 0] = 4 \quad [\text{given}]$$

$$\Rightarrow 2 - x = 8 \Rightarrow x = -6$$


102 (a)

The sum of the distance of a point P from two perpendicular lines in a plane is 1, then the locus of P is a rhombus

103 (a)

$$\text{In } \triangle DCE, \tan 30^\circ = \frac{150}{CD}$$

$$\Rightarrow CD = \sqrt{3} \times 150$$



Now, In $\triangle DCF$,

$$\tan \theta = \frac{DF}{CD} = \frac{200}{\sqrt{3} \cdot 150} = \frac{4}{3\sqrt{3}}$$

104 (b)

$$2 \left(a \sin^2 \frac{C}{2} + c \sin^2 \frac{A}{2} \right)$$

$$= 2 \left(a \frac{(s-a)(s-b)}{ab} + c \frac{(s-b)(s-c)}{bc} \right)$$


$$= 2 \left(\frac{(s-b)}{b} (s-a + s-c) \right)$$

$$= \frac{2}{b} (s-b)b$$

$$= 2(s-b) = a - b + c$$

105 (c)

Let H be the orthocenter of $\triangle OAB$



$$\therefore (\text{slope of } OP) \cdot (\text{slope of } BA) = -1$$

$$\Rightarrow \left(\frac{y-0}{3-0} \right) \cdot \left(\frac{4-0}{3-4} \right) = -1$$


$$\Rightarrow -\frac{4}{3}y = -1$$

$$\Rightarrow y = \frac{3}{4}$$

$$\therefore \text{Required orthocentre} = (3, y) = \left(3, \frac{3}{4} \right)$$

106 (b)

Let the fourth vertex be $D(x, y)$



We know that two diagonals of a parallelogram are bisect each other

$$\therefore \frac{-1+7}{2} = \frac{2+x}{2} \Rightarrow x = 4$$

$$\text{and } \frac{-6+2}{2} = \frac{-5+y}{2} \Rightarrow y = 1$$

\therefore Fourth vertex of D is $(4, 1)$

107 (c)

We have, $\cos A \cos B + \sin A \sin B \sin C = 1$

$$\Rightarrow 2 \cos A \cos B + 2 \sin A \sin B \sin C = 2$$

$$\Rightarrow 2 \cos A \cos B + 2 \sin A \sin B \sin C$$

$$= \cos^2 A + \sin^2 A + \cos^2 B + \sin^2 B$$

$$\Rightarrow (\cos A - \cos B)^2 + (\sin A - \sin B)^2$$

$$+ 2 \sin A \sin B (1 - \sin C) = 0$$

$$\Rightarrow \cos A - \cos B = 0, \quad \sin A - \sin B = 0$$

$$\text{and } 1 - \sin C = 0$$

$$\Rightarrow A = B \text{ and } C = 90^\circ$$

$$\Rightarrow a = b \text{ and } C = 90^\circ$$

108 (d)

We have, $A + B + C = 180^\circ$

$$\Rightarrow A = 180^\circ - (B + C)$$

$$\Rightarrow \tan A = \tan(180^\circ - B - C)$$

$$\Rightarrow \tan 90^\circ = -\tan(B + C)$$

$$\Rightarrow \infty = -\frac{\tan B + \tan C}{1 - \tan B \tan C}$$

$$\Rightarrow 1 - \tan B \tan C = 0$$

$$\Rightarrow \tan B \tan C = 1$$

109 (b)

Since, the given points lies on a line, then

$$\begin{vmatrix} 1 & 1 & 1 \\ -5 & 5 & 1 \\ 13 & \lambda & 1 \end{vmatrix} = 0$$

$$\Rightarrow 1(5 - \lambda) - 1(-5 - 13) + 1(-5\lambda - 65) = 0$$

$$\Rightarrow -6\lambda = 42 \Rightarrow \lambda = -7$$

110 (a)


The vertices of triangle are $(0, 0)$, $(3, 0)$ and $(0, 4)$.

It is a right angled triangle, therefore

circumcentre is $\left(\frac{3}{2}, 2 \right)$

111 (c)

$$BC = 5, BA = 10$$



Let D divides AC in the ratio 2:1


\therefore Coordinate of D is $(\frac{1}{3}, \frac{1}{3})$

The bisector is the line joining B and D is

$$\frac{y-1}{x-5} = \frac{1}{7} \text{ or } x-7y+2=0$$

112 (b)

In $\triangle ECD$, $\tan 3\alpha = \frac{h}{CD}$



$$\Rightarrow CD = h \cot 3\alpha \dots (i)$$

In $\triangle EBD$, $\tan 2\alpha = \frac{h}{BD}$

$$\Rightarrow BD = h \cot 2\alpha \dots (ii)$$

In $\triangle EAD$, $\tan \alpha = \frac{h}{AD}$

$$\Rightarrow AD = h \cot \alpha \dots (iii)$$

From Eqs. (ii) and (iii),

$$AD - BD = h \cot \alpha - h \cot 2\alpha$$

$$AB = h(\cot \alpha - \cot 2\alpha) \dots (iv)$$

From Eqs. (i) and (ii),

$$BD - CD = h \cot 2\alpha - h \cot 3\alpha$$

$$\Rightarrow BC = h(\cot 2\alpha - \cot 3\alpha) \dots (v)$$

From Eqs. (iv) and (v),

$$\frac{AB}{BC} = \frac{h(\cot \alpha - \cot 2\alpha)}{h(\cot 2\alpha - \cot 3\alpha)}$$

$$\frac{\cos \alpha}{\sin \alpha} - \frac{\cos 2\alpha}{\sin 2\alpha}$$

$$= \frac{\sin 2\alpha - \cos 2\alpha \sin \alpha}{\sin 2\alpha \sin \alpha}$$

$$= \frac{\sin(2\alpha - \alpha)}{\sin 2\alpha \sin \alpha}$$

$$= \frac{\sin \alpha \sin 2\alpha}{\sin(3\alpha - 2\alpha)}$$


$$= \frac{\sin 2\alpha \sin 3\alpha}{\sin 3\alpha}$$

$$= \frac{\sin 2\alpha}{\sin \alpha} = 3 - 4 \sin^2 \alpha$$

$$= 3 - 2(1 - \cos 2\alpha)$$

$$= 1 + 2 \cos 2\alpha$$

114 (b)



In $\triangle AEF$, $\tan \phi = \frac{AE}{EF} = \frac{150}{EF}$

$$\frac{5}{2} = \frac{150}{EF}$$

$$\Rightarrow EF = 60 \text{ m}$$

and in $\triangle ACD$

$$\tan \theta = \frac{AC}{CD}$$

$$\Rightarrow \frac{4}{3} = \frac{150 - h}{60} \quad [\because CD = EF]$$

$$\Rightarrow 80 = 150 - h$$

$$\Rightarrow h = 70 \text{ m}$$

$$\therefore AC = 80 \text{ m and } CD = 60 \text{ m}$$


$$\Rightarrow AD = \sqrt{AC^2 + CD^2}$$

$$= \sqrt{6400 + 3600}$$

$$= \sqrt{10000} = 100 \text{ m}$$

115 (d)

In $\triangle APC$, $\sin(\angle PAC) = \frac{CP}{AC}$



$$\Rightarrow AC = \frac{r}{\sin \frac{\alpha}{2}} = r \operatorname{cosec} \frac{\alpha}{2} \dots (i)$$

Again, in $\triangle ABC$, $\sin \beta = \frac{BC}{AC}$

$$\Rightarrow BC = AC \sin \beta$$

$$\Rightarrow H = r \operatorname{cosec} \left(\frac{\alpha}{2} \right) \sin \beta \text{ [from Eq.(i)]}$$

116 (b)

Since, $\cos A = \frac{b^2 + c^2 - a^2}{2bc}$

$$\Rightarrow b^2 - 2bc \cos A + (c^2 - a^2) = 0$$

It is given that b_1 and b_2 are the roots of this equation

Therefore, $b_1 + b_2 = 2c \cos A$ and $b_1 b_2 = c^2 - a^2$

$$\Rightarrow 3b_1 = 2c \cos A \text{ and } 2b_1^2 = c^2 - a^2$$

$$[\because b_2 = 2b_1]$$

$$\Rightarrow 2 \left(\frac{2c}{3} \cos A \right)^2 = c^2 - a^2$$

$$\Rightarrow 8c^2 (1 - \sin^2 A) = 9c^2 - 9a^2$$

$$\Rightarrow \sin A = \sqrt{\frac{9a^2 - c^2}{8c^2}}$$

117 (a)

$$\because (\sqrt{a} + \sqrt{b} + \sqrt{c})(\sqrt{a} + \sqrt{b} - \sqrt{c})$$

$$= (\sqrt{a} + \sqrt{b})^2 - c$$

$$= a + b - c + 2\sqrt{ab} > 0$$

$$\therefore \sqrt{a} + \sqrt{b} > \sqrt{c}$$

118 (a)

In ΔABC , $\angle A = 30^\circ$, $BC = 10$ cm

O is the centre of circle

$$\therefore \angle BOC = 60^\circ$$

and OB and OC are the radius

$$\therefore \angle OBC = \angle OCB = 60^\circ$$


$\Rightarrow \Delta OBC$ is an equilateral triangle

\therefore radius of circle is

$$OB = OC = BC = 10 \text{ cm}$$

Now, area of the circumcircle is πr^2

$$= \pi(10)^2 = 100\pi \text{ sq cm}$$



119 (b)

We have, $R = \frac{abc}{4\Delta}$ and $r = \frac{\Delta}{s}$

$$\therefore \frac{R}{r} = \frac{abc}{4\Delta} \cdot \frac{s}{\Delta}$$

$$= \frac{abc \cdot s}{4\Delta^2}$$

Since, $a : b : c = 4 : 5 : 6$

$$\Rightarrow \frac{a}{4} = \frac{b}{5} = \frac{c}{6} = k \text{ (say)}$$

$$\text{Thus, } \frac{R}{r} = \frac{(4k)(5k)(6k)}{4\left(\frac{15k}{2} - 4k\right)\left(\frac{15k}{2} - 5k\right)\left(\frac{15k}{2} - 6k\right)}$$


$$= \frac{120k^3 \cdot 2}{k^3 \cdot 7 \cdot 5 \cdot 3} = \frac{16}{7}$$

120 (b)

Given equation of lines are

$$x = 0, y = 0 \text{ and } 3x + 4y = 12$$

Incentre is on the line $y = x$ (Angled bisector of OA and OB)



Angle bisector of $y = 0$ and $3x + 4y = 12$ is

$$\pm 5y = 3x + 4y - 12$$

$$\Rightarrow 3x + 9y = 12$$

$$\text{and } 3x - y = 12$$

Here, $3x + 9y = 12$ internal bisector

So, intersection point of $y = x$ and $3x + 9y = 12$ is $(1, 1)$

\therefore The required point of the incentre of triangle is $(1, 1)$

121 (a)

$$BP - AP = \pm 6 \text{ or } BP = AP \pm 6$$

$$\Rightarrow \sqrt{x^2 + (y + 4)^2} = \sqrt{x^2 + (y - 4)^2} \pm 6$$

Squaring and simplification, we get

$$4y - 9 = \pm 3\sqrt{x^2 + (y - 4)^2}$$

Again squaring, we get

$$9x^2 - 7y^2 + 63 = 0$$


122 (c)

Let OP be the tower whose height is h metres

$$\text{In } \Delta OAP, \tan \alpha = \frac{OP}{OA}$$

$$\Rightarrow OA = h \cot \alpha \dots(i)$$

$$\text{In } \Delta OBP, \tan \beta = \frac{OP}{OB}$$



$$\Rightarrow OB = h \cot \beta \dots(ii)$$

Now, in ΔOAB , $AB^2 = OA^2 + OB^2$

$$\Rightarrow d^2 = h^2 (\cot^2 \alpha + \cot^2 \beta) \text{ [from Eq. (i) and (ii)]}$$

$$\Rightarrow h = \frac{d}{\sqrt{\cot^2 \alpha + \cot^2 \beta}}$$

123 (d)

Since, the coordinates of P are $(1, 0)$

Let any point Q on $y^2 = 8x$ is $(2t^2, 4t)$

Again, let mid point of PQ is (h, k) , so

$$h = \frac{2t^2 + 1}{2} \Rightarrow 2h = 2t^2 + 1 \dots(i)$$

$$\text{and } k = \frac{4t + 0}{2} \Rightarrow t = \frac{k}{2} \dots(ii)$$

on putting the value of t from Eq. (ii) in Eq. (i), we get


$$2h = \frac{2k^2}{4} + 1 \Rightarrow 4h = k^2 + 2$$

Hence, locus of (h, k) is $y^2 - 4x + 2 = 0$

124 (b)

Let $P(t, t)$ divides AB in the ratio $k : 1$, then

$$\frac{3k+k}{k+1} = t \text{ and } \frac{5k+2}{k+1} = t$$



$$\Rightarrow \frac{3k+k}{k+1} = \frac{5k+2}{k+1}$$

$$\Rightarrow 4k - 5k = 2$$

$$\Rightarrow k = -2$$

125 (c)

Since, a, b and c , the sides of a triangle are in AP

$$\therefore 2b = a + c \dots(i)$$

$$\text{We know that, } \cos A = \frac{b^2+c^2-a^2}{2bc}$$

$$\Rightarrow \cos A = \frac{b^2+c^2-(2b-c)^2}{2bc} \quad [\text{from Eq. (i)}]$$

$$\Rightarrow \cos A = \frac{b^2+c^2-4b^2-c^2+4bc}{2bc}$$

$$\Rightarrow \cos A = \frac{4c-3b}{2c}$$

126 (b)

The intersection points of given lines are

$$(0, 0), \left(\frac{5}{2}, 5\right), \left(\frac{5}{3}, 5\right)$$

$$\therefore \text{Area of } \Delta = \frac{1}{2} \begin{vmatrix} 0 & 0 & 1 \\ \frac{5}{2} & 5 & 1 \\ \frac{5}{3} & 5 & 1 \end{vmatrix}$$


$$= \frac{1}{2} \left[1 \left(\frac{25}{2} - \frac{25}{3} \right) \right]$$

$$= \frac{1}{2} \times \frac{25}{6} = \frac{25}{12} \text{ sq units}$$

127 (d)

Given that,

$$\tan \theta = \frac{4}{3} \text{ and } \tan \phi = \frac{5}{2} \dots(i)$$



$$\text{In } \Delta ABE, \tan \phi = \frac{150}{d}$$

$$\Rightarrow d = 150 \cot \phi$$

$$= 150 \times \frac{2}{5} = 60 \text{ m} \dots(ii)$$

$$\text{In } \Delta DCE, \tan \theta = \frac{h}{d}$$

$$\Rightarrow \frac{4}{3} = \frac{h}{d} \quad [\text{from Eq. (i)}]$$

$$\Rightarrow h = \frac{4}{3} (60) \quad [\text{from Eq. (ii)}]$$

$$\Rightarrow h = 80 \text{ m}$$


$$\text{Now, in } \Delta DCE, DE^2 = DC^2 + CE^2$$

$$\Rightarrow x^2 = 60^2 + 80^2 = 10000$$

$$\Rightarrow x = 100 \text{ m}$$

128 (d)

$$\text{Given, } A - B = 60^\circ$$



By sine rule,

$$\frac{2a}{\sin A} = \frac{a}{\sin B}$$

$$\Rightarrow \sin A - 2 \sin B = 0$$

$$\Rightarrow \sin(60^\circ + B) - 2 \sin B = 0$$

$$\Rightarrow \frac{\sqrt{3}}{2} \cos B + \frac{1}{2} \sin B - 2 \sin B = 0$$

$$\Rightarrow \frac{\sqrt{3}}{2} \cos B - \frac{3}{2} \sin B = 0$$

$$\Rightarrow \sqrt{3} \left(\frac{1}{2} \cos B - \frac{\sqrt{3}}{2} \sin B \right) = 0$$

$$\Rightarrow \sqrt{3} [\cos(60^\circ + B)] = 0$$

$$\Rightarrow 60^\circ + B = 90^\circ$$

$$\Rightarrow B = 30^\circ$$

$$\Rightarrow A = 90^\circ$$

Hence, it is right angled triangle

129 (c)

$$\begin{vmatrix} 3q & 0 & 1 \\ 0 & 3p & 1 \\ 1 & 1 & 1 \end{vmatrix} = 0$$

$$\Rightarrow 3q(3p-1) + 1(0-3p) = 0$$

$$\Rightarrow 9pq = 3p + 3q$$

$$\Rightarrow \frac{1}{p} + \frac{1}{q} = 3$$

130 (c)

$$\text{Here, } s = \frac{15+36+39}{2} = 45$$

$$\therefore \sin \frac{C}{2} = \sqrt{\frac{(s-a)(s-b)}{ab}}$$

$$\Rightarrow \sin \frac{C}{2} = \sqrt{\frac{(45-15)(45-36)}{15 \times 36}}$$

$$= \sqrt{\frac{30 \times 9}{15 \times 36}} = \frac{1}{\sqrt{2}}$$

131 (c)

$$\text{Since, } \frac{c}{\sin C} = 2R \Rightarrow c = 2R \quad [\because C = 90^\circ] \dots(i)$$

$$\text{And } \tan \frac{C}{2} = \frac{r}{s-c}$$

$$\Rightarrow \tan \frac{\pi}{4} = \frac{r}{s-c}$$

$$\therefore r = s - c$$


$$\Rightarrow a + b - c = 2r \dots(ii)$$

From Eqs. (i) and (ii), we get

$$2(r+R) = a+b$$

132 (b)

Let BC be the light house



$$\text{In } \triangle ABC, \tan 15^\circ = \frac{60}{d}$$

$$\Rightarrow d = 60 \cot 15^\circ = 60 \left(\frac{\sqrt{3} + 1}{\sqrt{3} - 1} \right) \text{ m}$$

133 (b)

$$\text{Given that, } \cos^2 A + \cos^2 C = \sin^2 B$$

Obviously it is not an equilateral triangle because $A = B = C = 60^\circ$ does not satisfy the given condition. But $B = 90^\circ$, then $\sin^2 B = 1$ and

$$\cos^2 A + \cos^2 C = \cos^2 A + \cos^2 \left(\frac{\pi}{2} - A \right)$$

$$= \cos^2 A + \sin^2 A = 1$$

Hence, this satisfies the condition, so it is a right angled triangle but not necessary isosceles triangle

134 (d)

$$\text{Given, } a : b : c = 1 : \sqrt{3} : 2$$

$$\text{Here, } c^2 = a^2 + b^2$$

\therefore Triangle is right angled at C

$$\therefore \angle C = 90^\circ$$

$$\text{and } \frac{a}{b} = \frac{1}{\sqrt{3}}$$

$$\Rightarrow \tan A = \frac{1}{\sqrt{3}}$$

$$\Rightarrow \angle A = 30^\circ \text{ and } \angle B = 60^\circ \text{ [as } \angle A + \angle B = 90^\circ]$$


$$\therefore \text{Ratio of angles} = \angle A : \angle B : \angle C = 90^\circ$$

$$= 30^\circ : 60^\circ : 90^\circ = 1 : 2 : 3$$

135 (c)

$$\text{Given, } \tan \theta = \frac{1}{9}$$

$$\text{In } \triangle ABC, \tan \phi = \frac{3h}{3} = h \dots (i)$$



$$\text{In } \triangle ABD, \tan(\theta + \phi) = \frac{4h}{3}$$

$$\Rightarrow \frac{\tan \theta + \tan \phi}{1 - \tan \theta \tan \phi} = \frac{4h}{3}$$

$$\Rightarrow \frac{\frac{1}{9} + h}{1 - \frac{h}{9}} = \frac{4h}{3}$$

$$\Rightarrow \frac{1 + 9h}{9 - h} = \frac{4h}{3}$$

$$\Rightarrow 3 + 27h = 36h - 4h^2$$

$$\Rightarrow 4h^2 - 9h + 3 = 0$$

$$\Rightarrow h = \frac{9 \pm \sqrt{81 - 48}}{2 \times 4} = \frac{9 \pm \sqrt{33}}{8}$$

136 (c)

$$\text{Let } a = 60^\circ - d, B = 60^\circ, C = 60^\circ + d$$

$$\therefore \frac{b}{c} = \frac{\sin B}{\sin C} = \sqrt{\frac{3}{2}}$$

$$\Rightarrow \frac{\sin 60^\circ}{\sin(60^\circ + d)} = \sqrt{\frac{3}{2}}$$

$$\Rightarrow \frac{\sqrt{3}}{2 \sin(60^\circ + d)} = \sqrt{\frac{3}{2}}$$

$$\Rightarrow \sin(60^\circ + d) = \frac{1}{\sqrt{2}}$$


$$\Rightarrow 60^\circ + d = 45^\circ \Rightarrow d = -15^\circ$$

$$\text{So, } \angle A = 75^\circ$$

137 (b)

$$\text{In } \triangle ABC, \tan 30^\circ = \frac{87}{x}$$

$$\Rightarrow x = 87 \times \sqrt{3}$$



$$\therefore \text{Speed} = \frac{\text{Distance}}{\text{Time}}$$

$$\Rightarrow \text{Time} = \frac{87 \times \sqrt{3} \times 60}{5.8 \times 1000} = \frac{9\sqrt{3}}{10} \text{ min}$$

138 (c)

It is given that the centroid of the triangle formed by the points (a, b) , (b, c) and (c, a) is at the origin

$$\therefore \left(\frac{a + b + c}{3}, \frac{a + b + c}{3} \right) = (0, 0)$$

$$\Rightarrow a + b + c = 0 \Rightarrow a^3 + b^3 + c^3 = 3abc$$

139 (d)

We have,


$$\Delta = \text{Area of } \triangle ABC$$

$$\Rightarrow \Delta = \frac{1}{2} \times AB \times \dots = \frac{1}{2} \times 3 \times 4 = 6 \text{ sq. units}$$

$$s = \text{Semi-perimeter} = \frac{1}{2}(3 + 4 + 5) = 6 \text{ units}$$

$$\therefore r = \text{In-radius} = \frac{\Delta}{s} = 1$$

Hence, the coordinates of the incentre are $(1, 1)$



140 (c)

Given, $a^4 + b^4 + c^4 = 2c^2(a^2 + b^2)$... (i)

$$\therefore \cos C = \frac{a^2 + b^2 - c^2}{2ab}$$

$$\Rightarrow \cos^2 C$$

$$= \left[\frac{a^4 + b^4 + c^4 + 2a^2b^2 - 2c^2(a^2 + b^2)}{4a^2b^2} \right]$$

$$= \left[\frac{2c^2(a^2 + b^2) + 2a^2b^2 - 2c^2(a^2 + b^2)}{4a^2b^2} \right]$$

[from Eq. (i)]

$$\Rightarrow \cos^2 C = \frac{1}{2}$$


$$\Rightarrow \cos C = \pm \frac{1}{\sqrt{2}}$$

$$\Rightarrow \angle C = 45^\circ \text{ or } 135^\circ$$

141 (c)

Let $AB = AC$ and $\angle A = 120^\circ$

$$\therefore \text{Area of triangle} = \frac{1}{2} a^2 \sin 120^\circ$$



Where, $a = AD + BD$

$$= \sqrt{3} \tan 30^\circ + \sqrt{3} \cot 15^\circ$$

$$= 1 + \sqrt{3} \left(\frac{1 + \tan 45^\circ \tan 30^\circ}{\tan 45^\circ - \tan 30^\circ} \right)$$

$$= 1 + \sqrt{3} \left(\frac{\sqrt{3} + 1}{\sqrt{3} - 1} \right)$$

$$\therefore a = 4 + 2\sqrt{3}$$

$$\Rightarrow \text{Area of triangle} = \frac{1}{2} (4 + 2\sqrt{3})^2 \left(\frac{\sqrt{3}}{2} \right) = 12 + 7\sqrt{3}$$

142 (b)


$$\text{Area} = \frac{1}{2} \times \text{base} \times \text{altitude}$$

$$= \frac{1}{2} \times (2x \cos \theta) \times (x \sin \theta)$$

$$= \frac{1}{2} x^2 \sin 2\theta$$

(Since, maximum value of $\sin 2\theta$ is 1)

$$\therefore \text{Maximum area} = \frac{1}{2} x^2$$



143 (d)

$$\text{Here, } \tan \frac{A}{2} \tan \frac{C}{2} = \frac{1}{3}$$

$$\Rightarrow \sqrt{\frac{(s-b)(s-c)}{s(s-a)}} \cdot \sqrt{\frac{(s-b)(s-a)}{s(s-c)}} = \frac{1}{3}$$

$$\Rightarrow \frac{s-b}{s} = \frac{1}{3} \Rightarrow 2s = 3b$$

$$\Rightarrow 2b = a + c$$

$$\Rightarrow a, b, c \text{ are in AP}$$

144 (d)

Let the vertices of a triangle are $P(2, 1)$, $Q(5, 2)$ and $R(3, 4)$ and $A(x, y)$ be the circumcentre of ΔPQR

$$\therefore AP^2 = AQ^2$$

$$\Rightarrow (2-x)^2 + (1-y)^2 = (5-x)^2 + (2-y)^2$$

$$\Rightarrow 4 + x^2 - 4x + 1 + y^2 - 2y$$

$$= 25 + x^2 - 10x + 4 + y^2 - 4y$$

$$\Rightarrow 6x + 2y = 24$$

$$\Rightarrow 3x + y = 12 \quad \dots (i)$$

and $AP^2 = AR^2$

$$\Rightarrow (2-x)^2 + (1-y)^2 = (3-x)^2 + (4-y)^2$$

$$\Rightarrow 4 + x^2 - 4x + 1 + y^2 - 2y$$

$$= 9 + x^2 - 6x + 16 + y^2 - 8y$$

$$\Rightarrow 2x + 6y = 20$$

$$\Rightarrow x + 3y = 10 \quad \dots (ii)$$

On solving Eqs. (i) and (ii), we get

$$x = \frac{13}{4} \text{ and } y = \frac{9}{4}$$

$$\therefore \text{Circumcentre is } \left(\frac{13}{4}, \frac{9}{4} \right)$$

145 (c)

$$(a + b + c)(b + c - a) = kbc$$

$$\Rightarrow 2s(2s - 2a) = kbc$$

$$\Rightarrow \frac{s(s-a)}{bc} = \frac{k}{4}$$

$$\Rightarrow \cos^2 \left(\frac{A}{2} \right) = \frac{k}{4}$$

$$\therefore 0 < \cos^2 \left(\frac{A}{2} \right) < 1$$

$$\therefore 0 < \frac{k}{4} < 1$$

$$\Rightarrow 0 < k < 4$$

146 (d)

$$\begin{aligned} \therefore \text{Area of } \Delta PBC &= \frac{1}{2} \begin{vmatrix} \alpha & \beta & 1 \\ -3 & 5 & 1 \\ 4 & -2 & 1 \end{vmatrix} \\ &= \frac{1}{2} |7\alpha + 7\beta - 14| \end{aligned}$$

$$\begin{aligned} \text{Also, Area of } \Delta ABC &= \frac{1}{2} \begin{vmatrix} 6 & -3 & 1 \\ -3 & 5 & 1 \\ 4 & -2 & 1 \end{vmatrix} \\ &= \frac{1}{2} |42 - 21 - 14| = \frac{7}{2} \end{aligned}$$


$$\therefore \frac{\text{Area of } \Delta PBC}{\text{Area of } \Delta ABC} = \frac{\frac{7}{2} |\alpha + \beta - 2|}{\frac{7}{2}}$$

$$= |\alpha + \beta - 2|$$

147 (b)

Let DP is clock tower standing at the middle point D of BC

$$\text{Let } \angle PAD = \alpha = \cot^{-1} 3.2 \Rightarrow \cot \alpha = 3.2$$



$$\text{and } \angle PBD = \beta = \operatorname{cosec}^{-1} 2.6$$

$$\Rightarrow \operatorname{cosec} \beta = 2.6$$

$$\therefore \cot \beta = \sqrt{(\operatorname{cosec}^2 \beta - 1)}$$

$$= \sqrt{5.76} = 2.4$$

In ΔPAD and PBD ,

$$AD = h \cot \alpha = 3.2h$$

$$\text{and } BD = h \cot \beta = 2.4h$$

$$\text{In } \Delta ABD, AB^2 = AD^2 + BD^2$$

$$\Rightarrow 100^2 = [(3.2)^2 + (2.4)^2]h^2 = 16h^2$$

$$\Rightarrow h = \frac{100}{4} \Rightarrow h = 25 \text{ m}$$

148 (c)

$$a \cot A + b \cot B + c \cot C$$

$$= \frac{a}{\sin A} \cos A + \frac{b}{\sin B} \cos B + \frac{c}{\sin C} \cos C$$

$$= 2R (\cos A + \cos B + \cos C)$$

$$= 2R \left(1 + \frac{r}{R}\right) = 2(r + R)$$

150 (a)

Given pair of lines are rotated about the origin by $\pi/6$ in the anti-clockwise sense.

$$\therefore x = x' \cos \frac{\pi}{6} - y' \sin \frac{\pi}{6} = \frac{\sqrt{3}x' - y'}{2}$$

$$\text{and } y = x' \sin \frac{\pi}{6} + y' \cos \frac{\pi}{6} = \frac{x' + \sqrt{3}y'}{2}$$

on putting the values of x and y in given pair of lines, we get

$$\begin{aligned} \sqrt{3} \left(\frac{\sqrt{3}x' - y'}{2} \right)^2 - 4 \left(\frac{\sqrt{3}x' - y'}{2} \right) \left(\frac{x' + \sqrt{3}y'}{2} \right) \\ + \sqrt{3} \left(\frac{x' + \sqrt{3}y'}{2} \right)^2 = 0 \\ \Rightarrow \sqrt{3}(3x'^2 + y'^2 - 2\sqrt{3}x'y') \\ - 4(\sqrt{3}x'^2 3x'y' - x'y' - \sqrt{3}y'^2) \\ + \sqrt{3}(x'^2 + 3y'^2 + 2\sqrt{3}x'y') = 0 \\ \Rightarrow 3\sqrt{3}x'^2 + \sqrt{3}y'^2 - 6x'y' - 4\sqrt{3}x'^2 - 8x'y' \\ + 4\sqrt{3}y'^2 + \sqrt{3}x'^2 + 3\sqrt{3}y'^2 \\ + 6x'y' = 0 \end{aligned}$$

$$\Rightarrow 8\sqrt{3}y'^2 - 8x'y' = 0$$

$$\Rightarrow \sqrt{3}y'^2 - x'y' = 0$$

$$\therefore \text{Required equation is } \sqrt{3}y^2 - xy = 0$$

151 (c)

Using sine rule,

$$\frac{\sin A}{a} = \frac{\sin B}{b}$$

$$\Rightarrow \frac{\frac{2}{3}}{2} = \frac{\sin B}{3}$$

$$\Rightarrow \sin B = 1$$

$$\Rightarrow B = 90^\circ$$

152 (a)

since, b, c and a are in AP

$$\text{By sine rule, } \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$\Rightarrow a = \frac{b}{\sin B} = \frac{c}{\sin C} \quad [\because \angle A = 90^\circ]$$

$$\Rightarrow \sin B = \frac{b}{a}, \sin C = \frac{c}{a}$$

153 (d)

$$2a^2 + 4b^2 + c^2 = 4ab + 2ac$$

$$\Rightarrow a^2 + (2b)^2 - 4ab + a^2 + c^2 - 2ac = 0$$

$$\Rightarrow (a - 2b)^2 + (a - c)^2 = 0$$

$$\Rightarrow a = 2c = c$$

$$\cos B = \frac{a^2 + c^2 - b^2}{2ac}$$

$$= \frac{c^2 + c^2 - \left(\frac{c}{2}\right)^2}{2 \times c \times c} = \frac{2c^2 - \frac{c^2}{4}}{2c^2}$$

$$\Rightarrow \cos B = \frac{7}{8}$$

154 (b)

Given, M divides AB in the ratio $b : a$ (externally)

$$\therefore x = \frac{ba \cos \beta - ba \cos \alpha}{b - a}$$

$$\text{and } y = \frac{ab \sin \beta - ab \sin \alpha}{b - a}$$


$$\Rightarrow \frac{x}{y} = \frac{\cos \beta - \cos \alpha}{\sin \beta - \sin \alpha}$$

$$\Rightarrow \frac{x}{y} = \frac{2 \sin\left(\frac{\alpha+\beta}{2}\right) \sin\left(\frac{\alpha-\beta}{2}\right)}{2 \cos\left(\frac{\alpha+\beta}{2}\right) \sin\left(\frac{\beta-\alpha}{2}\right)}$$

$$\Rightarrow x \cos\left(\frac{\alpha+\beta}{2}\right) + y \sin\left(\frac{\alpha+\beta}{2}\right) = 0$$

155 (a)

$$\text{In } \triangle DAB, \tan \theta = \frac{64}{d}$$



$$\Rightarrow d = 64 \cot \theta \dots (i)$$

$$\text{In } \triangle CDE, \tan(90^\circ - \theta) = \frac{(100-64)}{d}$$

$$\Rightarrow d = 36 \tan \theta \dots (ii)$$

On multiplying Eqs. (i) and (ii), we get

$$d^2 = 36 \times 64 \Rightarrow d = 48$$

156 (b)


Let BC be the declivity and BA be the tower

\therefore In $\triangle ABC$, on applying sine rule

$$\frac{BC}{\sin 75^\circ} = \frac{AB}{\sin 30^\circ}$$


$$\Rightarrow AB = \frac{80 \sin 30^\circ}{\sin 75^\circ}$$

$$= \frac{40 \times 2\sqrt{2}}{\sqrt{3} + 1} = 40(\sqrt{6} - \sqrt{2}) \text{ ft}$$



157 (c)

Let the points be $B(x_1, y_1)$ and $D(x_2, y_2)$ and coordinates of mid point of BD are $\left(\frac{x_1+x_2}{2}, \frac{y_1+y_2}{2}\right)$



And coordinates of mid point of AC are $(0, 1)$

We know that mid point of both the diagonals lie on the same point E .

$$\therefore \frac{x_1+x_2}{2} = 0 \text{ and } \frac{y_1+y_2}{2} = 1$$

$$\Rightarrow x_1 + x_2 = 0 \dots (i)$$

$$\text{and } y_1 + y_2 = 2 \dots (ii)$$

also, slope of $BD \times$ slope of $AC = -1$

$$\frac{(y_1 - y_2)}{(x_1 - x_2)} \times \frac{(3 + 1)}{(0 - 0)} = -1$$

$$\Rightarrow y_1 - y_2 = 0 \dots (iii)$$

On solving Eqs. (ii) and (iii), we get

$$y_1 = 1, y_2 = 1$$

Now, slope of $AB \times$ slope of $BC = -1$

$$\Rightarrow \frac{(y_1 + 1)}{(x_1 - 0)} \times \frac{(y_1 - 3)}{(x_1 - 0)} = -1$$

$$\Rightarrow (y_1 + 1)(y_1 - 3) = -x_1^2$$


$$\Rightarrow 2(-2) = -x_1^2 \quad [\because y_1 = 1]$$

$$\Rightarrow x_1 = \pm 2$$

\therefore The required points are $(2, 1)$ and $(-2, 1)$

158 (a)

The diagonals meet at the mid point of AC , i.e. at $(3, 2)$ which lies on $y = 2x + c$



$$\therefore c = -4$$

$$\text{Let } B = (\alpha, 2\alpha - 4)$$

$$\because AB \perp BC$$

$$\Rightarrow \left(\frac{2\alpha - 7}{\alpha - 1}\right) \left(\frac{2\alpha - 5}{\alpha - 5}\right) = -1$$

$$\therefore \alpha^2 - 6\alpha + 8 = 0$$

$$\Rightarrow \alpha = 2, 4$$

The other two vertices are $(2, 0)$ and $(4, 4)$

159 (c)

$$\text{Given, } r_3 - r = r_1 + r_2$$

$$\Rightarrow 4R \sin \frac{C}{2} \left(\cos \frac{A}{2} \cos \frac{B}{2} - \sin \frac{B}{2} \sin \frac{A}{2} \right)$$

$$= 4R \cos \frac{C}{2} \left[\sin \frac{A}{2} \cos \frac{B}{2} + \cos \frac{A}{2} \sin \frac{B}{2} \right]$$

$$\Rightarrow \sin \frac{C}{2} \left[\cos \left(\frac{A+B}{2} \right) \right] = \cos \frac{C}{2} \left[\sin \left(\frac{A+B}{2} \right) \right]$$

$$\Rightarrow \sin \frac{C}{2} \left[\cos \left(\frac{\pi - C}{2} \right) \right] = \cos \frac{C}{2} \left[\sin \left(\frac{\pi - C}{2} \right) \right]$$

$$\left[\because A + B + C = \pi \Rightarrow \frac{A}{2} + \frac{B}{2} = \frac{\pi - C}{2} \right]$$

$$\Rightarrow \sin^2 \frac{C}{2} = \cos^2 \frac{C}{2} \Rightarrow \tan \frac{C}{2} = 1$$


$$\Rightarrow \angle C = \frac{\pi}{2}$$

$$\text{We know, } A + B + C = \pi \Rightarrow A + B = \frac{\pi}{2}$$

160 (d)

$$\text{Given } a = 3, b = 4, c = 5$$

$$\Rightarrow c^2 = a^2 + b^2$$



Therefore, it is a right angled triangle at C

$$\therefore R = \frac{1}{2}c = \frac{5}{2}$$

$$\text{and } r = \frac{\Delta}{s} = \frac{\frac{1}{2} \times 3 \times 4}{\frac{12}{2}} = 1$$

\therefore Distance between incentre and circumcentre


$$= \sqrt{R^2 - 2Rr}$$

$$= \sqrt{\left(\frac{5}{2}\right)^2 - 2 \cdot \frac{5}{2} \cdot 1}$$

$$= \sqrt{\frac{5}{2} \sqrt{\frac{5}{2}} - 2} = \frac{\sqrt{5}}{2}$$

161 (d)

Area of $\Delta ABC = a^2$



$$\Rightarrow \frac{1}{2} (a-x)3a = a^2$$

$$\Rightarrow a-x = \frac{2}{3} a$$

$$\Rightarrow x = \frac{a}{3}$$

Hence, one of the line on which third vertex lies is

$$x = \frac{a}{3}$$

162 (c)

Draw BE perpendicular to CA produced, then

$$BD = DC = \frac{a}{2} \text{ and } EA = AC = b$$

In ΔAEB ,

$$\cos(\pi - A) = \frac{b}{c}$$


$$\Rightarrow \cos A = -\frac{b}{c}$$

$$\Rightarrow \frac{b^2 + c^2 - a^2}{2bc} = -\frac{b}{c}$$

$$\Rightarrow a^2 = 3b^2 + c^2$$

$$\therefore \cos B = \frac{c^2 + a^2 - b^2}{2ac}$$

$$= \frac{c^2 + 3b^2 + c^2 - b^2}{2ca} = \frac{b^2 + c^2}{ca}$$



164 (d)

Let the sides of ΔABC be $a = n, b = n + 1, c = n + 2$, where n is a natural number. Then, C is the greatest and A is the least angle

As given $C = 2A$

$$\therefore \sin C = \sin 2A = 2 \sin A \cos A$$

$$\therefore kc = 2ka \frac{b^2 + c^2 - a^2}{2bc}$$

$$\Rightarrow bc^2 = a(b^2 + c^2 - a^2)$$

On substituting the values of a, b, c , we get

$$(n+1)(n+2)^2 = n[(n+1)^2 + (n+2)^2 - n^2]$$

$$= n(n^2 + 6n + 5)$$

$$= n(n+1)(n+5)$$

$$\Rightarrow (n+1)[(n+2)^2 - n(n+5)] = 0$$

Since, $n \neq 1$

$$\text{Thus, } (n+2)^2 = n(n+5)$$

$$\Rightarrow n^2 + 4n + 4 = n^2 + 5n$$

$$\Rightarrow n = 4$$

Hence, the sides of the triangle are 4, 5 and 6

165 (c)

$$3. \quad \frac{b^2 - c^2}{a \sin(B-C)} = \frac{2R^2(\sin^2 B - \sin^2 C)}{2R \sin A \sin(B-C)}$$

$$= \frac{2R \sin(B+C) \sin(B-C)}{\sin(B+C) \sin(B-C)} = 2R$$

$$4. \quad a \sin(B-C) + b \sin(C-A) + c \sin(A-B) = 0$$

$$= 2R[\sin A \sin(B-C) + \sin B \sin(C-A) + \sin C \sin(A-B)]$$

$$= 2R[\sin(B+C) \sin(B-C)$$

$$+ \sin(C+A) \sin(C-A)$$

$$+ \sin(A+B) \sin(A-B)]$$

$$= 2R[\sin^2 B - \sin^2 C + \sin^2 C - \sin^2 A + \sin^2 A - \sin^2 B]$$

$$= 2R(0) = 0$$

Hence, both of statements are correct

166 (b)

Let the general equation of the circle be

$$x^2 + y^2 + 2gx + 2fy + c = 0$$

The equation of circle passing through (0, 0), (2, 0) and (0, -2)

$$c = 0 \quad \dots(i)$$

$$4 + 4g + c = 0 \quad \dots(ii)$$

$$\text{and } 4 - 4f + c = 0 \quad \dots(iii)$$

On solving Eqs. (i), (ii) and (iii), we get

$$c = 0, \quad g = -1, \quad f = 1$$

\therefore The equation of circle becomes $x^2 + y^2 - 2x + 2y = 0$

Since, it passes through (k, -2), we get

$$k^2 + 4 - 2k - 4 = 0 \Rightarrow k = 0, 2$$

We have already take a point (0, -2), so we take only $k = 2$

167 (a)

$$\text{Let } X = x - h, \quad Y = y - k$$

$$\Rightarrow 0 = 7 - h, \quad 0 = -4 - k$$

$$\Rightarrow h = 7, \quad k = -4$$

Hence, $X = x - 7$ and $Y = y + 4$, then the point (4, 5) shifted to (-3, 9)

168 (a)

$$(a + b + c) \left(\tan \frac{A}{2} + \tan \frac{B}{2} \right) \\ = 2 \left(s \tan \frac{A}{2} + s \tan \frac{B}{2} \right)$$

$$= 2 \left(\frac{\Delta}{s-a} + \frac{\Delta}{s-b} \right)$$

$$= 2\Delta \frac{2s - (a+b)}{(s-a)(s-b)}$$

$$= 2\Delta \left(\frac{c}{(s-a)(s-b)} \right)$$

$$= 2c \cot \frac{C}{2}$$

169 (b)

$$\therefore \frac{2}{1!9!} + \frac{2}{3!7!} + \frac{1}{5!5!} = \frac{8^a}{(2b)!}$$

$$\Rightarrow \frac{1}{1!9!} + \frac{1}{3!7!} + \frac{1}{5!5!} + \frac{1}{3!7!} + \frac{1}{9!1!} = \frac{8^a}{(2b)!}$$

$$\Rightarrow \frac{1}{10!} \left(\frac{10!}{1!9!} + \frac{10!}{3!7!} + \frac{10!}{5!5!} + \frac{10!}{7!3!} + \frac{10!}{9!1!} \right) \\ = \frac{8^a}{(2b)!}$$

$$\Rightarrow \frac{1}{10!} ({}^{10}C_1 + {}^{10}C_3 + {}^{10}C_5 + {}^{10}C_7 + {}^{10}C_9) \\ = \frac{8^a}{(2b)!}$$

$$\Rightarrow \frac{2^9}{10!} = \frac{8^a}{(2b)!} = \frac{2^{3a}}{(2b)!}$$

$$\Rightarrow a = 3, \quad b = 5$$

$$\text{Also, } 2b = a + c \Rightarrow 10 = 3 + c \Rightarrow c = 7$$

$$\therefore a = 3, \quad b = 5, \quad c = 7$$

$$\therefore \frac{\tan A + \tan B}{2} \geq \sqrt{\tan A \tan B} \quad \dots(i)$$

$$\text{Also, } \cos C = \frac{a^2 + b^2 - c^2}{2ab}$$

$$= \frac{9 + 25 - 49}{30} = -\frac{1}{2}$$

$$\Rightarrow \angle C = 120^\circ \text{ and } A, B < 60^\circ$$

$$\tan A + \tan B + \tan C = \tan A \tan B \tan C$$

$$\Rightarrow \tan A + \tan B - \sqrt{3} = -\sqrt{3} \tan A \tan B$$

$$\therefore \tan A + \tan B = \sqrt{3}(1 - \tan A \tan B) \quad \dots(ii)$$

$$\text{Also, } \tan A + \tan B > 0$$

$$\Rightarrow \sqrt{3}(1 - \tan A \tan B) > 0$$

$$\Rightarrow \tan A \tan B < 1 \quad \dots(iii)$$

From Eq. (i) and (ii),

$$\frac{\sqrt{3}(1 - \tan A \tan B)}{2} \geq \sqrt{(\tan A \tan B)}$$

$$\text{Let } \tan A \tan B = \lambda$$

$$\therefore \sqrt{3}(1 - \lambda) \geq 2\sqrt{\lambda}$$

$$\Rightarrow 3\lambda^2 - 10\lambda + 3 \geq 0$$

$$\Rightarrow (3\lambda - 1)(\lambda - 3) \geq 0$$

$$\therefore \lambda - 3 < 0 \quad [\text{from Eq. (iii)}]$$


$$\therefore 3\lambda - 1 \leq 0$$

$$\Rightarrow \lambda \leq \frac{1}{3}$$

$$\Rightarrow \tan A \tan B \leq \frac{1}{3}$$

170 (c)

$$\text{In } \triangle BCD, \tan 60^\circ = \frac{H_1}{d}$$



$$\Rightarrow H_1 = d \tan 60^\circ$$

$$\text{and in } \triangle ABE, \tan 30^\circ = \frac{H_2}{d}$$

$$\Rightarrow H_2 = d \tan 30^\circ$$

$$\therefore \frac{H_1}{H_2} = \frac{\tan 60^\circ}{\tan 30^\circ} = \frac{\sqrt{3}}{1/\sqrt{3}} = \frac{3}{1}$$

171 (c)

$$\text{We have, } \cos C = \frac{a^2 + b^2 - c^2}{2ab}$$

$$\Rightarrow \cos 60^\circ = \frac{a^2 + b^2 - c^2}{2ab}$$

$$\Rightarrow a^2 + b^2 - c^2 = ab$$

$$\Rightarrow b^2 + bc + a^2 + ac = ab + ac + bc + c^2$$

On dividing by $(a+c)(b+c)$ and add 2 on both sides, we get

$$1 + \frac{b}{a+c} + 1 + \frac{a}{b+c} = 3$$

$$\Rightarrow \frac{1}{a+c} + \frac{1}{b+c} = \frac{3}{a+b+c}$$

172 (a)

$$(a+c)^2 - b^2 = 3ac \Rightarrow a^2 + c^2 - b^2 = ac$$

$$\text{But } \cos B = \frac{a^2 + c^2 - b^2}{2ac} = \frac{1}{2} \Rightarrow \angle B = \frac{\pi}{3} = 60^\circ$$

173 (c)

Given, a^2, b^2, c^2 are in AP

$$\Rightarrow \sin^2 B - \sin^2 A = \sin^2 C - \sin^2 B$$

$$\Rightarrow \sin(B+A) \sin(B-A) = \sin(C+B) \sin(C-B)$$

$$\Rightarrow \sin C (\sin B \cos A - \cos B \sin A)$$

$$= \sin A (\sin C \cos B - \cos C \sin B)$$

On dividing by $\sin A \sin B \sin C$, we get

$$2 \cot B = \cot A + \cot C$$

$$\Rightarrow \cot A, \cot B, \cot C \text{ are in AP}$$

174 (c)

$$\text{Given, } \sin A \sin B = \frac{ab}{c^2}$$

$$\Rightarrow c^2 = \frac{ab}{\sin A \sin B} = \left(\frac{a}{\sin A}\right) \left(\frac{b}{\sin B}\right)$$

$$\Rightarrow c^2 = \left(\frac{c}{\sin C}\right)^2$$

$$\therefore \left(\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}\right)$$

$$\Rightarrow \sin^2 C = 1$$

$$\Rightarrow c = 90^\circ$$

Hence, ΔABC is a right angled triangle

175 (a)

Let a, b, c be the sides of triangle, then

$$a + b + c = \frac{6}{3} (\sin A + \sin B + \sin C)$$

$$\Rightarrow a + b + c = 2(\sin A + \sin B + \sin C)$$

$$\Rightarrow \frac{a}{2} = \sin A$$

$$\text{But } a = 1$$


$$\therefore \sin A = \frac{1}{2} \Rightarrow \angle A = \frac{\pi}{6}$$

176 (a)

$$\text{The centroid of } \Delta ABC = \left(\frac{2+8+5}{3}, \frac{3+10+5}{3}\right) = (5, 6)$$

177 (d)

Since, the tower OP makes equal angle at the vertices of the triangle, therefore foot of the tower is the circumcentre



$$\text{In } \Delta OAP, \tan \alpha = \frac{OP}{OA}$$

$$\Rightarrow OP = OA \tan \alpha$$

$$\Rightarrow OP = R \tan \alpha$$

178 (a)

In ΔABE ,


$$\sin \beta = \frac{BE}{b}$$

$$\Rightarrow BE = h_1 = b \sin \beta$$

Using sine rule in ΔAED ,

$$\frac{\sin(\alpha - \beta)}{ED} = \frac{\sin(\gamma - \alpha)}{b}$$

$$\Rightarrow ED = \frac{b \sin(\alpha - \beta)}{\sin(\gamma - \alpha)}$$



Now, in ΔFED ,

$$\sin \gamma = \frac{h_2}{ED}$$

$$\Rightarrow h_2 = \frac{b \sin(\alpha - \beta) \sin \gamma}{\sin(\gamma - \alpha)}$$

\therefore Total height, CD

$$= h_1 + h_2 = b \sin \beta + \frac{b \sin(\alpha - \beta) \sin \gamma}{\sin(\gamma - \alpha)}$$

$$= \frac{b[\sin \beta \sin(\gamma - \alpha) + \sin(\alpha - \beta) \sin \gamma]}{\sin(\gamma - \alpha)}$$

$$= \frac{b[\sin \beta \{\sin \gamma \cos \alpha - \cos \gamma \sin \alpha\} + \sin \gamma \{\sin \alpha \cos \beta \cos \alpha\}]}{\sin(\gamma - \alpha)}$$

$$= \frac{b[\sin \beta \sin \gamma \cos \alpha - \sin \beta \cos \gamma \sin \alpha + \{\sin \gamma \sin \alpha \cos \beta - \sin \gamma \sin \beta \cos \alpha\}]}{\sin(\gamma - \alpha)}$$

$$= \frac{b \sin \alpha \sin(\gamma - \beta)}{\sin(\gamma - \alpha)}$$

179 (c)

Given, $a = 1, b = 2, \angle C = 60^\circ$

$$\therefore \text{Area of triangle} = \frac{1}{2} ab \sin C$$

$$= \frac{1}{2} \times 1 \times 2 \times \sin 60^\circ$$

$$= \frac{\sqrt{3}}{2} \text{ sq unit}$$

180 (c)

The given points are collinear

$$\text{If } \begin{vmatrix} t_1 & 2at_1 + at_1^3 & 1 \\ t_2 & 2at_2 + at_2^3 & 1 \\ t_3 & 2at_3 + at_3^3 & 1 \end{vmatrix} = 0$$

$$\Rightarrow a \begin{vmatrix} t_1 & 2t_1 + t_1^3 & 1 \\ t_2 & 2t_2 + t_2^3 & 1 \\ t_3 & 2t_3 + t_3^3 & 1 \end{vmatrix} = 0$$

Applying $R_2 \rightarrow R_2 - R_1, R_3 \rightarrow R_3 - R_1$, we get

$$\begin{vmatrix} t_1 & 2t_1 + t_1^3 & 1 \\ t_2 - t_1 & 2(t_2 - t_1) + (t_2^3 - t_1^3) & 0 \\ t_3 - t_1 & 2(t_3 - t_1) + (t_3^3 - t_1^3) & 0 \end{vmatrix} = 0$$

$$\Rightarrow (t_2 - t_1)(t_3 - t_1) \begin{vmatrix} t_1 & 2t_1 + t_1^3 & 1 \\ 1 & 2 + t_2^2 + t_1^2 + t_1 t_2 & 0 \\ 1 & 2 + t_3^2 + t_1^2 + t_3 t_1 & 0 \end{vmatrix} = 0$$

$$\Rightarrow (t_2 - t_1)(t_3 - t_1)(t_3 - t_2)(t_3 + t_2 + t_1) = 0$$

$$\Rightarrow t_1 + t_2 + t_3 = 0$$

$$[\because t_1 \neq t_2 \neq t_3]$$

181 (b)

Let P is a point on the perpendicular bisector of AB , its equation is

$$(y - 1) = \frac{1}{3}(x - 4) \Rightarrow x - 3y - 1 = 0$$

So, general point is $P(3h + 1, h)$

$$\text{Area} = \frac{1}{2} \begin{vmatrix} 3h + 1 & h & 1 \\ 3 & 4 & 1 \\ 5 & -2 & 1 \end{vmatrix} = \pm 10$$

$$\Rightarrow h = 2, 0$$

Or position of points is $(7, 2)$ and $(1, 0)$

182 (a)

$$\text{Given, } \angle A = \frac{\pi}{2}, \text{ then } a^2 = b^2 + c^2 = 4^2 + 3^2 = 25$$

$$\text{and } \frac{a}{\sin A} = 2R \Rightarrow a = 2R \text{ and } a = 5$$

$$\text{also, } r = \frac{\Delta}{s} = \frac{bc}{a+b+c} \quad (\because \Delta = \frac{bc}{2})$$

$$\therefore \frac{R}{r} = \frac{a(a+b+c)}{2bc} = \frac{5 \times 12}{2 \times 4 \times 3} = \frac{5}{2}$$

183 (d)

Since A, B, C are in AP

$$\Rightarrow 2B = A + C \Rightarrow \angle B = 60^\circ$$

$$\therefore \frac{a}{2} (2 \sin C \cos C) + \frac{c}{a} (2 \sin A \cos A)$$

$$= 2k (a \cos C + c \cos A)$$

$$= 2k(b)$$

$$= 2 \sin B \text{ [using, } b = a \cos C + c \cos A]$$

$$= \sqrt{3}$$

184 (d)

Given equation is

$$x^2 - 5x + 6 = 0$$

$$\Rightarrow (x - 3)(x - 2) = 0$$

$$\Rightarrow x = 3, 2$$

These are the sides of a triangle

$$\text{Let } a = 3, b = 2, \angle C = \frac{\pi}{3}$$

$$\Rightarrow \cos\left(\frac{\pi}{3}\right) = \frac{3^2 + 2^2 - c^2}{2 \cdot 3 \cdot 2} \quad \left[\because \cos C = \frac{a^2 + b^2 - c^2}{2ab} \right]$$

$$\Rightarrow \frac{1}{2} = \frac{13 - c^2}{12} \Rightarrow c^2 = 7$$

$$\Rightarrow c = \sqrt{7} \text{ [sides cannot be negative]}$$

$$\therefore \text{Perimeter of a triangle} = a + b + c$$

$$= 3 + 2 + \sqrt{7} = 5 + \sqrt{7}$$

185 (b)

Given, $\angle A = 60^\circ, a = 5, b = 4$

$$\therefore \cos A = \frac{b^2 + c^2 - a^2}{2bc}$$


$$\Rightarrow \cos 60^\circ = \frac{1}{2} = \frac{16 + c^2 - 25}{8c}$$

$$\Rightarrow 4c = c^2 - 9$$

$$\Rightarrow c^2 - 4c - 9 = 0$$

186 (a)

Let PQ be the height h of the tower and A, B are the points of observations



$$\text{We have, } \angle QAP = \frac{\pi}{4}, \angle BAP = \frac{\pi}{6},$$

$$AB = 10 \text{ m, } BQ = 10 \text{ m}$$

$$\therefore \angle QAB = \frac{\pi}{12} = \angle AQB$$

$$\Rightarrow \angle ABQ = \pi - \frac{\pi}{6} = \frac{5\pi}{6}$$

On applying cosine rule in ΔABQ , we get

$$AQ^2 = AB^2 + BQ^2 - 2AB \cdot BQ \cos \frac{5\pi}{6}$$

$$= 100 + 100 + 200 \cdot \frac{\sqrt{3}}{2}$$

$$= 100(2 + \sqrt{2})$$

$$\Rightarrow AQ = 10\sqrt{2 + \sqrt{3}}$$

$$\text{In } \Delta APQ, AP = AQ \cos \frac{\pi}{4} = \frac{10\sqrt{2 + \sqrt{3}}}{\sqrt{2}}$$

$$= 5\sqrt{4 + 2\sqrt{3}} = 5\sqrt{(\sqrt{3} + 1)^2}$$

$$\Rightarrow AP = 5(1 + \sqrt{3}) \text{ m}$$

187 (a)

Let the points be $A = (a \cos \theta, a \sin \theta)$ and

$$B = (a \cos \phi, a \sin \phi)$$

$$\therefore AB$$

$$= \sqrt{(a \cos \theta - a \cos \phi)^2 + (a \sin \theta - a \sin \phi)^2}$$

$$= \sqrt{a^2 \cos^2 \theta + a^2 \cos^2 \phi - 2a^2 \cos \theta \cos \phi + a^2 \sin^2 \theta + a^2 \sin^2 \phi - 2a^2 \sin \theta \sin \phi}$$

$$= \sqrt{2a^2 - 2a^2(\cos \theta \cos \phi + \sin \theta \sin \phi)}$$

$$= \sqrt{2}a(\sqrt{1 - \cos(\theta - \phi)})$$

$$\Rightarrow 2a = \sqrt{2}a \sqrt{2} \sin\left(\frac{\theta - \phi}{2}\right)$$

$$\Rightarrow \sin\left(\frac{\theta - \phi}{2}\right) = 1$$

$$\Rightarrow \frac{\theta - \phi}{2} = n\pi \pm \frac{\pi}{2}$$

$$\Rightarrow \theta - \phi = 2n\pi \pm \pi$$

$$\Rightarrow \theta = 2n\pi \pm \pi - \phi$$

Where $n \in Z$

188 (d)

Area of pentagon

$$= \frac{1}{2} \left[\begin{array}{l} x_1y_2 + x_2y_3 + x_3y_4 + x_4y_5 \\ + x_5y_1 - (y_1x_2 + y_2x_3 + y_3x_4 \\ + y_4x_5 + y_5x_1) \end{array} \right]$$

$$= \frac{1}{2} [0(0) + 12(2) + 12(7) + 6(5) + 0(0) - \{0(0) + 0 + 2(6) + 7(0) + 5(0)\}]$$

$$= \frac{1}{2} [(24 + 84 + 30 - 12)]$$

$$= 63 \text{ sq unit}$$

189 (a)

Let the height of the flag be h

191 (b)

$$\frac{a \cos A + b \cos B + c \cos C}{a + b + c}$$

$$= \frac{(2R \sin A) \cos A + (2R \sin B) \cos B + 2R \sin C (\cos C)}{2R \sin A + 2R \sin B + 2R \sin C}$$

$$\left(\because R = \frac{a}{2 \sin A} = \frac{b}{2 \sin B} = \frac{c}{2 \sin C} = \frac{abc}{4\Delta} \right)$$

$$= \frac{R[2 \sin A \cos A + 2 \sin B \cos B + 2 \sin C \cos C]}{2R[\sin A + \sin B + \sin C]}$$


$$= \frac{1}{2} \frac{(\sin 2A + \sin 2B + \sin 2C)}{(\sin A + \sin B + \sin C)}$$

$$= \frac{2[4 \cos(A/2) \cos(B/2) \cos(C/2)]}{4 \sin A \sin B \sin C}$$

$$\left(\because \sin 2\alpha + \sin 2\beta + \sin 2\gamma = 4 \sin \alpha \sin \beta \sin \gamma \right)$$

$$\left(\text{and } \sin \alpha + \sin \beta + \sin \gamma = 4 \cos \frac{\alpha}{2} \cos \frac{\beta}{2} \cos \frac{\gamma}{2} \right)$$

$$= \frac{4 \left[2 \sin \frac{A}{2} \cos \frac{A}{2} \times 2 \sin \frac{B}{2} \cos \frac{B}{2} \times 2 \sin \frac{C}{2} \cos \frac{C}{2} \right]}{2 \times 4 \left[\cos \frac{A}{2} \cos \frac{B}{2} \cos \frac{C}{2} \right]}$$



$$\text{In } \Delta ARQ, \tan \alpha = \frac{b}{a} \dots (i)$$

$$\text{and in } \Delta PRQ, \tan 2\alpha = \frac{h+b}{a} \dots (ii)$$


$$\Rightarrow \frac{2 \tan \alpha}{1 - \tan^2 \alpha} = \frac{h+b}{a}$$

$$\Rightarrow \frac{2 \times \frac{b}{a}}{1 - \frac{b^2}{a^2}} = \frac{h+b}{a} \quad [\text{from Eq. (i)}]$$

$$\Rightarrow \frac{2ab}{a^2 - b^2} = \frac{h+b}{a} \Rightarrow h = \frac{b(a^2 + b^2)}{(a^2 - b^2)}$$

190 (a)

Let BC be the height of kite



$$\text{In } \Delta ABC, \sin 60^\circ = \frac{h}{120}$$

$$\Rightarrow \frac{\sqrt{3}}{2} = \frac{h}{120}$$

$$\Rightarrow h = 60\sqrt{3} \text{ m}$$

The height of the kite is $60\sqrt{3} \text{ m}$


$$= 4 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}$$

$$= \frac{r}{R} \left[\because r = 4R \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2} \right]$$

192 (a)

Let $\angle APC = \alpha$. Then,

$$\tan \alpha = \frac{AC}{AP} = \frac{AC}{n AB} = \frac{AB}{2n AB} = \frac{1}{2n}$$



In $\triangle APB$, we have

$$\tan(\alpha + \beta) = \frac{AB}{AP} = \frac{AB}{n AB} = \frac{1}{n}$$

Now, $\beta = \alpha + \beta - \alpha$

$$\Rightarrow \tan \beta = \frac{\tan(\alpha + \beta) - \tan \alpha}{1 + \tan(\alpha + \beta) \tan \alpha}$$

$$\Rightarrow \tan \beta = \frac{1/n - 1/2n}{1 + 1/n \cdot 1/2n} = \frac{n}{2n^2 + 1}$$

193 (d)

Suppose $P(3,7)$ divides the segment joining $A(1,1)$ and $B(6,16)$ in the ratio $\lambda : 1$. Then,

$$\frac{6\lambda + 1}{\lambda + 1} = 3 \text{ and } \frac{16\lambda + 1}{\lambda + 1} = 7$$

$$\Rightarrow \lambda = \frac{2}{3}$$

$\Rightarrow P$ divides AB internally in the ratio $2 : 3$

Thus, Q divides AB externally in the ratio $2 : 3$ and hence its coordinates are

$$\left(\frac{2 \times 6 - 3 \times 1}{2 - 3}, \frac{2 \times 16 - 3 \times 1}{2 - 3} \right) \equiv (-9, -29)$$

194 (d)

Let the angles of a triangle are $3x, 5x$ and $10x$

$$\therefore 3x + 5x + 10x = 180^\circ \Rightarrow x = 10^\circ$$

\therefore Smallest angle of a triangle = 30°

And the greatest angle = 100°

Required ratio = $\sin 30^\circ : \sin 100^\circ$

$$= \frac{1}{2} : \cos 10^\circ = 1 : 2 \cos 10^\circ$$

195 (b)

$$\text{Given, } \begin{vmatrix} 1 & a & b \\ 1 & c & a \\ 1 & b & c \end{vmatrix} = 0$$

$$\Rightarrow c^2 - ab - a(c - a) + b(b - c) = 0$$

$$\Rightarrow a^2 + b^2 + c^2 - ab - bc - ca = 0$$

$$\Rightarrow \frac{1}{2} [2a^2 + 2b^2 + 2c^2 - 2ab - 2bc - 2ca] = 0$$

$$\Rightarrow \frac{1}{2} [(a - b)^2 + (b - c)^2 + (c - a)^2] = 0$$

$$\Rightarrow a = b = c$$

$$\Rightarrow \angle A = 60^\circ, \angle B = 60^\circ, \angle C = 60^\circ$$


$$\therefore \sin^2 A + \sin^2 B + \sin^2 C$$

$$= \sin^2 60^\circ + \sin^2 60^\circ + \sin^2 60^\circ$$

$$= \left(\frac{\sqrt{3}}{2}\right)^2 + \left(\frac{\sqrt{3}}{2}\right)^2 + \left(\frac{\sqrt{3}}{2}\right)^2 = \frac{9}{4}$$

197 (c)

Let $\angle OAB = \theta$



Then, $OA + AB = 1 + 4 \cot \theta + 4 + \tan \theta$

$$= 5 + 4 \cot \theta + \tan \theta \geq 5 + 4 = 9$$

(using $AM \geq GM$)

199 (c)

Given $\angle A = 20^\circ$

$$\therefore \angle B = \angle C = 80^\circ$$

Then, $b = c$

$$\therefore \frac{a}{\sin 20^\circ} = \frac{b}{\sin 80^\circ} = \frac{c}{\sin 80^\circ}$$

$$\Rightarrow \frac{a}{\sin 20^\circ} = \frac{b}{\cos 10^\circ}$$

$$\Rightarrow a = 2b \sin 10^\circ \dots (i)$$

$$\therefore a^3 + b^3 = 8b^3 \sin^3 10^\circ + b^3$$

$$= b^3 \{2(4 \sin^3 10^\circ) + 1\}$$

$$= b^3 \{2(3 \sin 10^\circ - \sin 30^\circ) + 1\}$$

$$= b^3 \{6 \sin 10^\circ\}$$

$$= 3b^2 \{2b \sin 10^\circ\}$$

$$= 3b^2 a \text{ [from Eq. (i)]}$$

$$= 3ac^2 \text{ } (\because b = c)$$

200 (a)

$$a(\cos^2 B + \cos^2 C) + \cos A (c \cos C + b \cos B)$$

$$= \cos B (a \cos B + b \cos A)$$


$$+ \cos C (a \cos C + c \cos A)$$

$$= (\cos B)c + (\cos C)b$$

$$= a$$

201 (b)

Let $(x_1, y_1), (x_2, y_2)$ and (x_3, y_3) are the coordinates of the points D, E and F



$$\therefore x_1 = \frac{3 \times 6 - 1 \times 1}{4} = \frac{17}{4}$$

$$\text{and } y_1 = \frac{-2 \times 3 + 4}{4} = -\frac{2}{4} = -\frac{1}{2}$$

$$\text{similarly, } x_2 = 0, y_2 = \frac{5}{2}$$

$$\text{and } x_3 = -\frac{5}{4}, y_3 = 4$$

let (x, y) be the coordinates of centroid of $\triangle DEF$

$$\therefore x = \frac{1}{3} \left(\frac{17}{4} + 0 - \frac{5}{4} \right) = 1$$

$$\text{and } y = \left(-\frac{1}{2} + \frac{5}{2} + 4 \right) \frac{1}{3} = 2$$

\therefore Coordinates of centroid are $(1, 2)$

202 (a)

$$\text{We have, } \frac{1}{a+c} = \frac{1}{b+c} = \frac{3}{a+b+c}$$

$$\Rightarrow \frac{a+b+2c}{(a+c)(b+c)} = \frac{3}{a+b+c}$$

$$\Rightarrow a^2 + ab + ac + ab + b^2 + bc + 2ca + 2bc + 2c^2$$

$$= 3(ab + ac + bc + c^2)$$


$$\Rightarrow a^2 + b^2 - c^2 = ab$$

$$\Rightarrow \frac{a^2 + b^2 - c^2}{2ab} = \frac{1}{2} = \cos C$$

$$\Rightarrow \angle C = 60^\circ$$

203 (a)

Let CD be the tower of height H metre



$$\text{From } \triangle BCD, \frac{H}{d} = \tan 30^\circ$$

and from $\triangle ABD$,

$$\frac{h}{d} = \tan 60^\circ$$

$$\therefore \frac{H/d}{h/d} = \frac{\tan 30^\circ}{\tan 60^\circ} \Rightarrow H = \frac{h}{3}$$

204 (c)


Using sine rule in $\triangle ADC$,

$$\frac{\sin(y+z)}{DC} = \frac{\sin C}{AD}$$

$$\text{In } \triangle ABD, \frac{\sin x}{BD} = \frac{\sin B}{AD}$$

$$\text{In } \triangle AEC, \frac{\sin z}{EC} = \frac{\sin C}{AE}$$

$$\text{In } \triangle ABE, \frac{\sin(x+y)}{BE} = \frac{\sin B}{AE}$$



$$\therefore \frac{\sin(x+y) \sin(y+z)}{\sin x \sin z} = \frac{BE}{AE} \times \frac{DC}{AD} \times \frac{AD}{BD} \times \frac{AE}{EC}$$

$$= \frac{2BD \times 2EC}{BD \times EC} = 4$$

205 (a)

$$\therefore \frac{\sin A}{\sin C} = \frac{\sin(A-B)}{\sin(B-C)}$$

$$\Rightarrow \frac{\sin A}{\sin C} = \frac{\sin A \cos B - \cos A \sin B}{\sin B \cos C - \cos B \sin C}$$

$$\Rightarrow \frac{a}{c} = \frac{a \cos B - b \cos A}{b \cos C - c \cos B}$$

$$\Rightarrow ab \cos C + bc \cos A = 2ac \cos B$$

$$\Rightarrow \frac{a^2 2 + b^2 - c^2}{2} + \frac{b^2 + c^2 - a^2}{2} = 2 \frac{c^2 + a^2 - b^2}{2}$$

$$\Rightarrow a^2 + c^2 = 2b^2$$

$$\Rightarrow a^2, b^2, c^2 \text{ are in AP}$$

206 (b)

Let point (x_1, y_1) be on the line $3x + 4y = 5$

$$\therefore 3x_1 + 4y_1 = 5 \dots (i)$$

$$\text{Also, } (x_1 - 1)^2 + (y_1 - 2)^2 = (x_1 - 3)^2 + (y_1 - 4)^2$$

$$\Rightarrow x_1^2 + y_1^2 - 2x_1 - 4y_1 + 5$$

$$= x_1^2 + y_1^2 - 6x_1 - 8y_1 + 25$$

$$\Rightarrow 4x_1 + 4y_1 = 20 \dots (ii)$$

On solving Eqs. (i) and (ii), we get

$$x_1 = 15, y_1 = -10$$

207 (a)

Let $A(x_1, y_1), B(x_2, y_2)$ be two fixed points and

let $P(h, k)$ be a variable point such that

$$\angle APB = \frac{\pi}{2}$$

Then, slope of $AP \times$ slope of $BP = -1$

$$\Rightarrow \frac{k - y_1}{h - x_1} \cdot \frac{k - y_2}{h - x_2} = -1$$

$$\Rightarrow (h - x_1)(h - x_2) + (k - y_1)(k - y_2) = 0$$

Hence, locus of (h, k) is

$$(x - x_1)(x - x_2) + (y - y_1)(y - y_2) = 0$$


Which is a circle having AB as diameter

208 (a)

$$\text{Given, } a \cos^2 \frac{C}{2} + c \cos^2 \frac{A}{2} = \frac{3b}{c}$$

$$\begin{aligned} \Rightarrow a \left[\frac{s(s-c)}{ab} \right] + c \left[\frac{s(s-a)}{bc} \right] &= \frac{3b}{2} \\ \Rightarrow \frac{s(s-c+s-a)}{b} &= \frac{3b}{2} \\ \Rightarrow 2s(2s-c-a) &= 3b^2 \\ \Rightarrow (a+b+c)b &= 3b^2 \\ \Rightarrow 2b &= a+c \end{aligned}$$

209 (c)



$$\begin{aligned} \therefore \frac{a}{2R} &= \sin \frac{\pi}{n} \\ \text{and } \frac{a}{2r} &= \tan \frac{\pi}{n} \quad \therefore \frac{r}{R} = \cos \frac{\pi}{n} \end{aligned}$$

For $n = 3$ gives $\frac{r}{R} = \frac{1}{2}$
 For $n = 4$ gives $\frac{r}{R} = \frac{1}{\sqrt{2}}$
 For $n = 6$ gives $\frac{r}{R} = \frac{\sqrt{3}}{2}$


210 (d)

Given, the vertex angles are A, B, C are side BC

$$\begin{aligned} \therefore \text{Area} &= \frac{1}{2} a(k \sin C) \sin B \\ &= \frac{1}{2} \cdot \frac{a^2 k \sin C \sin B}{a} \\ &= \frac{1}{2} \cdot \frac{a^2 k \sin C \sin B}{k \sin A} \\ &= \frac{1}{2} a^2 \cdot \frac{\sin B \sin C}{\sin A} \end{aligned}$$

211 (c)

Let OP be the flag staff of height h standing at the centre O of the field



In $\triangle OEP$, $OE = h \cot 15^\circ = h(2 + \sqrt{3})$
 and in $\triangle OFP$, $OF = h \cot 45^\circ = h$

$$\begin{aligned} \therefore EF &= h \sqrt{1 + (2 + \sqrt{3})^2} \\ &= 2h \sqrt{2 + \sqrt{3}} \end{aligned}$$

Since, $AC = 1200$
 $\Rightarrow 2EF = 4h \sqrt{2 + \sqrt{3}}$
 $\Rightarrow h = \frac{300}{\sqrt{2 + \sqrt{3}}}$

$$= 300 \sqrt{2 - \sqrt{3}} \text{ m}$$

214 (b)

$$\begin{aligned} A + B + C &= \pi \\ 2B &= A + C \\ \Rightarrow 3B &= \pi \\ \Rightarrow B &= \frac{\pi}{3} \\ \therefore \frac{a+c}{b} &= \frac{\sin A + \sin C}{\sin B} \\ &= \frac{2 \sin \frac{A+C}{2} \cos \frac{A-C}{2}}{\sin \frac{\pi}{3}} \\ &= \frac{2 \sin \frac{\pi}{3} \cos \frac{A-C}{2}}{\sin \frac{\pi}{3}} \\ &= 2 \cos \frac{A-C}{2} \end{aligned}$$

215 (c)

The altitudes from the vertices A, B and C are $\frac{2\Delta}{a}, \frac{2\Delta}{b}$ and $\frac{2\Delta}{c}$ respectively

Also, these are in HP

$$\Rightarrow \frac{1}{a}, \frac{1}{b}, \frac{1}{c} \text{ are in HP}$$

$$\Rightarrow a, b, c \text{ are in AP}$$


$$\Rightarrow \sin A, \sin B, \sin C \text{ are in AP}$$

216 (d)

Given, the equation of circle is

$$x^2 + y^2 - 4x - 6y - 12 = 0$$

$$\begin{aligned} \text{Radius of this circle} &= \sqrt{4 + 9 + 12} \\ &= 5 \text{ unit} \end{aligned}$$



In $\triangle BOD$,

$$\cos 30^\circ = \frac{BD}{OB}$$

$$\Rightarrow BD = \frac{\sqrt{3}}{2} \times 5 = \frac{5\sqrt{3}}{2} \text{ unit}$$

$$\therefore BC = 2BD = 5\sqrt{3}$$

$$\begin{aligned} \text{Hence, area of } \triangle ABC &= \frac{\sqrt{3}}{4} (5\sqrt{3})^2 = \frac{75}{4} \sqrt{3} \text{ sq} \\ &\text{units} \end{aligned}$$

218 (d)

Let $A(1, \sqrt{3}), B(0, 0), C(2, 0)$ be the given points

$$\therefore BC = \sqrt{(2-0)^2 + (0-0)^2} = 2$$

$$CA = \sqrt{(2-1)^2 + (0-\sqrt{3})^2} = 2$$

$$\text{and } AB = \sqrt{1+3} = 2$$

∴ Triangle is equilateral

We know that in equilateral triangle incentre is the same as Centroid of the triangle

$$\therefore \text{Incentre is } \left(\frac{1+0+2}{3}, \frac{\sqrt{3}+0+0}{3} \right) = \left(1, \frac{1}{\sqrt{3}} \right)$$


219 (b)

We know that area of the triangle with polar coordinates

$$\begin{aligned} &= \frac{1}{2} \left| \sum r_1 r_2 \sin(\theta_1 - \theta_2) \right| \\ &= \frac{1}{2} \left| 2.1 \sin\left(0 - \frac{\pi}{3}\right) + 2.3 \sin\left(\frac{\pi}{3} - \frac{2\pi}{3}\right) \right. \\ &\quad \left. + 3.1 \sin\left(\frac{2\pi}{3} - 0\right) \right| \\ &= \frac{1}{2} \left| -2 \cdot \frac{\sqrt{3}}{2} - \frac{6\sqrt{3}}{2} + \frac{3\sqrt{3}}{2} \right| = \frac{5\sqrt{3}}{4} \text{ sq units} \end{aligned}$$

220 (d)

For finding the distance AP , first we find out the perpendicular bisector of AB and AC



Perpendicular bisector of $A(1, 3)$ and $B(-3, 5)$ is

$$\begin{aligned} &2x(x_1 - x_2) + 2y(y_1 - y_2) \\ &= (x_1^2 + y_1^2) - (x_2^2 + y_2^2) \\ \Rightarrow &2x(1 + 3) + 2y(3 - 5) = (1 - 9) - (9 - 25) \\ \Rightarrow &2x - y + 6 \dots(i) \end{aligned}$$

Similarly, perpendicular bisector of $A(1, 3)$ and $C(-3, 5)$ is

$$\begin{aligned} &2x(1 - 5) + 2y(3 + 1) = (1 + 9) - (25 + 1) \\ \Rightarrow &-8x + 8y = -16 \\ \Rightarrow &x - y - 2 = 0 \dots(ii) \end{aligned}$$

Point of intersection of Eqs. (i) and (ii) is

$$P = (-8, -10)$$

Then, the distance between P and A is

$$\begin{aligned} PA &= \sqrt{(1 + 8)^2 + (3 + 10)^2} \\ &= \sqrt{81 + 169} \\ &= \sqrt{250} = \sqrt{25 \times 10} = 5\sqrt{10} \end{aligned}$$

222 (b)

Let sides are $a = 13, b = 12, c = 5$

$$\begin{aligned} \text{Now, } &a^2 = b^2 + c^2 \\ \Rightarrow &(13)^2 = (12)^2 + 5^2 \\ \Rightarrow &169 = 169 \\ \therefore &\angle A = 90^\circ \end{aligned}$$

$$\text{Since, } R = \frac{a}{2 \sin A} = \frac{13}{2 \sin 90^\circ} = \frac{13}{2}$$

223 (b)

Since, the vertices of the triangle are

$(a \cos t, a \sin t), (b \sin t, -\cos t)$ and $(1, 0)$

Let the coordinate of centroid be

$$x = \frac{a \cos t + b \sin t + 1}{3}$$

$$\Rightarrow 3x - 1 = a \cos t + b \sin t \dots(i)$$

$$\text{and } y = \frac{a \sin t - b \cos t + 0}{3}$$

$$3y = a \sin t - b \cos t \dots(ii)$$

On squaring and adding Eqs. (i) and (ii), we get

$$\begin{aligned} (3x - 1)^2 + (3y)^2 &= a^2(\cos^2 t + \sin^2 t) + b^2(\sin^2 t \\ &\quad + \cos^2 t) \end{aligned}$$

$$\Rightarrow (3x - 1)^2 + (3y)^2 = a^2 + b^2$$


224 (c)

In $\triangle ABC, BC = h \cot 60^\circ$

and in $\triangle ABD, BD = h \cot 45^\circ$

Since, $BD - BC = DC$

$$\Rightarrow h \cot 45^\circ - h \cot 60^\circ = 7$$




$$\Rightarrow h = \frac{7}{\cot 45^\circ - \cot 60^\circ} = \frac{7}{\left(1 - \frac{1}{\sqrt{3}}\right)}$$

$$= \frac{7\sqrt{3}(\sqrt{3} + 1)}{2} \text{ m}$$

225 (c)

Here, coordinates are $A(2, 4), B(-2, -4)$ and $C(2, -4)$



$$\begin{aligned} \text{Now, } |AB| &= \sqrt{(-2 - 2)^2 + (-4 - 4)^2} \\ &= \sqrt{16 + 64} = \sqrt{80} = 4\sqrt{5} \end{aligned}$$

226 (c)

Let the vertex $A(x, y)$ is equidistant from B and C

$$\therefore (x - 1)^2 + (y - 3)^2 = (x + 2)^2 + (y - 7)^2$$

$$\Rightarrow 6x - 8y + 43 = 0$$

Only the option (c) satisfy it

227 (a)

Let the ratio be $k: 1$

$$\therefore \frac{-7k + 3}{k + 1} = \frac{1}{2} \Rightarrow k = \frac{1}{3}$$

Hence, ratio is 1: 3 internally

228 (d)

We know that orthocenter of the right angled triangle ABC , right angled at A is A

Here, triangle is right angled at $O(0,0)$


\therefore Orthocentre $= (0,0)$

229 (a)

Here, $x^2 - y^2 + 2y = 1$

$\Rightarrow x^2 = (y-1)^2$

$\Rightarrow x = y-1$ and $x = -y+1$




Which could be graphically as shown in figure
Which gives angle bisector as $y = 1$ and $x = 0$

\therefore Required area $= \frac{1}{2} \times 2 \times 2 = 2$ sq units

230 (a)

In $\triangle ABE$, $\tan 60^\circ = \frac{12}{AB} \Rightarrow AB = 4\sqrt{3}$ m



and in $\triangle ACE$

$\tan 45^\circ = \frac{12}{AC} \Rightarrow AC = 12$ m

In $\triangle ABC$, $BC = \sqrt{AC^2 - AB^2} = \sqrt{144 - 48} = 4\sqrt{6}$ m

\therefore Area of rectangular field $= AB \times BC = 4\sqrt{3} \times 4\sqrt{6} = 48\sqrt{2}$ sqm


231 (a)

In $\triangle ABD$,

$\cos 60^\circ = \frac{2^2 + 5^2 - BD^2}{2(2)(5)}$

$\Rightarrow BD^2 = 19$

Now, in $\triangle BCD$



$\cos 120^\circ = \frac{CD^2 + 9 - 19}{(2)(3)(CD)}$

$\Rightarrow CD^2 + 3CD - 10 = 0$

$\Rightarrow CD = -5, 2$

$\Rightarrow CD = 2$ ($\because CD \neq -5$)

232 (c)


We have, $\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c} = \frac{1}{2R}$

$\Rightarrow \sin A = \frac{a}{2R}$

$\therefore 2R^2 \sin A \sin B \sin C = 2R^2 \frac{a}{2R} \cdot \frac{b}{2R} \cdot \frac{c}{2R}$

$= \frac{abc}{4R} = \Delta$

233 (c)



In $\triangle BAD$,

$\cos(90^\circ - B) = \frac{AD}{c}$

$\Rightarrow AD = c \sin B$

Similarly, $BE = a \sin C$ and $CF = b \sin A$

Since, AD, BE, CF are in HP


$\therefore c \sin B, a \sin C, b \sin A$ are in HP

$\Rightarrow \frac{1}{\sin C \sin B}, \frac{1}{\sin A \sin C}, \frac{1}{\sin B \sin A}$ are in AP

$\Rightarrow \sin A, \sin B, \sin C$ are in AP

234 (d)

Let P be the middle point of the line segment joining $A(3, -1)$ and $B(1, 1)$ is $(2, 0)$



Let P be shifted to Q by 2 unit and y -coordinate of Q is greater than that of P

Now, slope of $AB = \frac{1 - (-1)}{1 - 3} = 1$

then slope of $PQ = -1$

The coordinate of Q becomes $(2 \pm 2 \cos \theta, 0 \pm 2 \sin \theta)$, where $\tan \theta = 1$

ie, $(2 \pm \sqrt{2}, \pm \sqrt{2})$

As, y -coordinates of Q is greater than that of P

\therefore We take $Q = (2 + \sqrt{2}, \sqrt{2})$

235 (d)

Area of triangle $= \frac{1}{2} \times 6 \times |\alpha| = 15$

$\Rightarrow |\alpha| = 5 \Rightarrow \alpha = \pm 5$

and β can take any real value

236 (a)

Let points are $A(2, 3), B(3, 4), C(4, 5), D(5, 6)$

$$\therefore AB = a = \sqrt{(3-2)^2 + (4-3)^2} = \sqrt{2}$$

Similarly, $BC = b = \sqrt{2}, CD = c = \sqrt{2}$

And $DA = d = 3\sqrt{2}$

$$\text{Now, } s = \frac{a+b+c+d}{2}$$

$$= \frac{\sqrt{2} + \sqrt{2} + \sqrt{2} + 3\sqrt{2}}{2} = 3\sqrt{2}$$

$$\therefore \text{Area} = \sqrt{(s-a)(s-b)(s-c)(s-d)}$$

$$= \sqrt{(3\sqrt{2} - \sqrt{2})(3\sqrt{2} - \sqrt{2})(3\sqrt{2} - \sqrt{2})(3\sqrt{2} - 3\sqrt{2})}$$

$$= 0$$

237 (a)

$$\sqrt{(2+5)^2 + (3-2)^2} = \sqrt{(x-1)^2 + (2-3)^2}$$


$$\Rightarrow 49 + 1 = (x-1)^2 + 1$$

$$\Rightarrow x-1 = \pm 7 \Rightarrow x = -6, 8$$

238 (d)

In an equilateral triangle length of median $CD =$

$$\frac{\sqrt{3}}{2} a$$



Also, centroid of triangle lies on the centre of circle

$$\therefore OC = \frac{2}{3} \times \frac{\sqrt{3}}{2} a = \frac{a}{\sqrt{3}}$$


$$\therefore \text{Area of circle} = \pi(OC)^2 = \frac{\pi a^2}{3} \text{ sq units}$$

239 (a)

$$\text{In } \triangle ABC, \tan 30^\circ = \frac{CB}{CA}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{CB}{6}$$

$$\Rightarrow CB = \frac{6\sqrt{3}}{3} = 2\sqrt{3}$$



$$\therefore \text{Area of } \triangle ACB = \frac{1}{2} \times CA \times CB$$

$$= \frac{1}{2} \times 6 \times 2\sqrt{3} = 6\sqrt{3} \text{ sq units}$$

240 (d)

$$\text{We have, } \frac{\cos A}{a} = \frac{\cos B}{b} = \frac{\cos C}{c}$$

$$\Rightarrow \frac{\cos A}{k \sin A} = \frac{\cos B}{k \sin B} = \frac{\cos C}{k \sin C}$$

$$\Rightarrow \cot A = \cot B = \cot C$$

$$\Rightarrow A = B = C = 60^\circ$$

$\triangle ABC$ is an equilateral triangle

$$\therefore \Delta = \frac{\sqrt{3}}{4} a^2 = \frac{\sqrt{3}}{4} (2)^2 \quad [\because a = 2 \text{ (given)}]$$

$$= \sqrt{3}$$

241 (c)

$$\text{We have, } \cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

$$\text{or } c^2 - 2bc \cos A + b^2 - a^2 = 0$$

$$\therefore c_1 + c_2 = 2b \cos A = 2 \times 2 \times \frac{\sqrt{3}}{2} = 2\sqrt{3}$$

$$\text{and } c_1 c_2 = b^2 - a^2 = 4 - 5 = -1$$

$$\therefore |c_1 - c_2| = \sqrt{(c_1 + c_2)^2 - 4c_1 c_2}$$

$$= \sqrt{12 + 4} = \sqrt{16} = 4$$

242 (c)

$$(a-b)^2 \cos^2 \frac{C}{2} + (a+b)^2 \sin^2 \frac{C}{2}$$

$$= (a^2 + b^2 - 2ab) \cos^2 \frac{C}{2}$$

$$+ (a^2 + b^2 + 2ab) \sin^2 \frac{C}{2}$$

$$= a^2 + b^2 + 2ab \left(\sin^2 \frac{C}{2} - \cos^2 \frac{C}{2} \right)$$

$$= a^2 + b^2 - 2ab \cos C$$

$$= a^2 + b^2 - (a^2 + b^2 - c^2) = c^2$$

243 (a)

$$1. \quad rr_1 r_2 r_3 = \frac{\Delta}{s} \cdot \frac{\Delta}{(s-a)} \cdot \frac{\Delta}{(s-b)} \cdot \frac{\Delta}{(s-c)} = \frac{\Delta^4}{\Delta^2} = \Delta^2$$

$$2. \quad \text{Now, } r_1 r_2 + r_2 r_3 + r_3 r_1$$

$$= \frac{\Delta \Delta}{(s-a)(s-b)} + \frac{\Delta \Delta}{(s-b)(s-c)}$$

$$+ \frac{\Delta \Delta}{(s-c)(s-a)}$$

$$= \Delta^2 \left[\frac{(s-c) + (s-a) + (s-b)}{(s-a)(s-b)(s-c)} \right]$$

$$= \frac{\Delta^2 [3s - (a+b+c)]}{\Delta^2 / s} = s^2$$

244 (c)

Let angles of a triangle are $x, 2x$ and $7x$ respectively

$$\therefore x + 2x + 7x = 180^\circ \Rightarrow x = 18^\circ$$

Hence, the angles are $18^\circ, 36^\circ, 126^\circ$

Greatest side $\propto \sin 18^\circ$

$$\begin{aligned} \therefore \text{Required ratio} &= \frac{\sin 126^\circ}{\sin 18^\circ} \\ &= \frac{\sin(90^\circ + 36^\circ)}{\sin 18^\circ} \\ &= \frac{\cos 36^\circ}{\sin 18^\circ} = \frac{\sqrt{5} + 1}{\sqrt{5} - 1} \end{aligned}$$

246 (a)

$$\text{Area of triangle} = \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ rx_1 & ry_1 & 1 \\ r^2x_1 & r^2y_1 & 1 \end{vmatrix} = 0$$

\therefore Points are collinear

247 (d)

Given, new coordinates = $(4, -3)$ and $\theta = 135^\circ$

$$\therefore 4 = x \cos 135^\circ + y \sin 135^\circ$$

$$\Rightarrow 4 = -\frac{x}{\sqrt{2}} + \frac{y}{\sqrt{2}} \quad \dots(i)$$

and $-3 = -x \sin 135^\circ + y \cos 135^\circ$

$$\Rightarrow -3 = -\frac{x}{\sqrt{2}} - \frac{y}{\sqrt{2}} \quad \dots(ii)$$

On adding Eqs. (i) and (ii), we get

$$1 = -\frac{2x}{\sqrt{2}} \Rightarrow x = -\frac{1}{\sqrt{2}}$$

On subtracting Eqs. (i) and (ii), we get

$$7 = \frac{2y}{\sqrt{2}} \Rightarrow y = \frac{7}{\sqrt{2}}$$

$$\text{Thus } (x, y) = \left(-\frac{1}{\sqrt{2}}, \frac{7}{\sqrt{2}}\right)$$

248 (d)

We know, if coordinate axes are rotated, then

$$p = (x \cos \theta - y \sin \theta, x \sin \theta + y \cos \theta)$$

It is rotated at an angle 135° i.e., $\theta = 135^\circ$ and the new point be

$$p = [4 \cos(90^\circ + 45^\circ)$$


$$+ 3 \sin(90^\circ + 45^\circ), 4 \sin(90^\circ + 45^\circ) - 3 \cos(90^\circ + 45^\circ)]$$

$$= [-4 \sin 45^\circ + 3 \cos 45^\circ, 4 \cos 45^\circ + 3 \sin 45^\circ]$$

$$= \left[4 \cdot \left(\frac{-1}{\sqrt{2}}\right) + 3 \cdot \frac{1}{\sqrt{2}}, 4 \cdot \frac{1}{\sqrt{2}} + 3 \cdot \frac{1}{\sqrt{2}}\right] = \left(-\frac{1}{\sqrt{2}}, \frac{7}{\sqrt{2}}\right)$$

249 (d)

Let the height of the tower be h meters



$$\text{In } \triangle BCD, \tan 45^\circ = \frac{h}{BC}$$

$$\Rightarrow BC = h \quad \dots(i)$$

$$\text{In } \triangle ACD, \tan 30^\circ = \frac{h}{AC}$$

$$\Rightarrow AC = \frac{h}{\tan 30^\circ}$$

$$\Rightarrow AB + BC = \sqrt{3} h$$

$$\Rightarrow 60 + h = \sqrt{3} h \quad [\text{from Eq.(i)}]$$

$$\Rightarrow h = \frac{60}{\sqrt{3} - 1}$$

$$\Rightarrow h = \frac{60(\sqrt{3} + 1)}{2}$$

$$\Rightarrow h = 30(\sqrt{3} + 1)\text{m}$$

251 (d)

$$\text{Given, } r_1 + r_3 = k \cos^2 \frac{B}{2}$$

$$\text{i.e., } s \tan \frac{A}{2} + s \tan \frac{C}{2} = k \cos^2 \frac{B}{2}$$

$$\Rightarrow s \left[\sqrt{\frac{(s-b)(s-c)}{s(s-a)}} + \sqrt{\frac{(s-a)(s-b)}{s(s-c)}} \right]$$

$$= k \frac{s(s-b)}{ac}$$

$$\Rightarrow k = \frac{ac}{s-b} \left[\sqrt{\frac{(s-b)(s-c)}{s(s-a)}} \right.$$

$$\left. + \sqrt{\frac{(s-a)(s-b)}{s(s-c)}} \right]$$

$$= \frac{ac}{s-b} \times \frac{\sqrt{s-b}}{\sqrt{s}} \left[\frac{\sqrt{(s-c)}}{\sqrt{(s-a)}} + \frac{\sqrt{(s-a)}}{\sqrt{(s-c)}} \right]$$

$$= \frac{ac(2s-a-c)}{\sqrt{s(s-a)(s-b)(s-c)}}$$

$$= \frac{ac(b)}{\Delta} = 4R$$

254 (c)

Area of $\triangle CAB$ = area of $\triangle CAD$ + area of $\triangle CDB$

$$\Rightarrow \frac{1}{2} ba \sin C = \frac{1}{2} b \cdot CD \sin \frac{C}{2} + \frac{1}{2} a \cdot CD \sin \frac{C}{2}$$

$$\Rightarrow CD + \frac{2ab}{a+b} \cos \frac{C}{2}$$


255 (c)

Let h be the height of a tower.

Since, $\angle AOB = 60^\circ$

$\therefore \triangle OAB$ is an equilateral triangle.

$$\therefore OA = OB = AB = a$$



$$\text{In } \triangle OAC, \tan 30^\circ = \frac{h}{a} \Rightarrow h = \frac{a}{\sqrt{3}}$$

256 (d)

$$\Delta = \begin{vmatrix} -\sin(\beta - \alpha) & -\cos \beta & 1 \\ \cos(\beta - \alpha) & \sin \beta & 1 \\ \cos(\beta - \alpha + \theta) & \sin(\beta - \theta) & 1 \end{vmatrix}$$

Clearly, $\Delta \neq 0$ for any value of α, β, θ , Hence, points are non-collinear

257 (a)

$$\begin{aligned} \text{Given, } 8R^2 &= a^2 + b^2 + c^2 \\ &= 4R^2(\sin^2 A + \sin^2 B + \sin^2 C) \\ \Rightarrow \sin^2 A + \sin^2 B + \sin^2 C &= 2 \\ \Rightarrow (\cos^2 A - \sin^2 C) + \cos^2 B &= 0 \\ \Rightarrow \cos(A - C) \cos(A + C) + \cos^2 B &= 0 \\ \Rightarrow 2 \cos B \cos A \cos C &= 0 \\ \Rightarrow \cos A = 0 \text{ or } \cos B = 0 \text{ or } \cos C &= 0 \\ \Rightarrow A = \frac{\pi}{2} \text{ or } B = \frac{\pi}{2} \text{ or } C = \frac{\pi}{2} \end{aligned}$$

258 (a)

$$\begin{aligned} \because AB + \sqrt{(4+1)^2 + (0+1)^2} &= \sqrt{26} \\ BC &= \sqrt{(3+1)^2 + (5+1)^2} = \sqrt{52} \\ CA &= \sqrt{(4-3)^2 + (0+5)^2} = \sqrt{26} \\ \text{So, in isosceles triangle side } AB &= CA \\ \text{Also, } (\sqrt{26})^2 + (\sqrt{26})^2 &= 52 \\ \text{So, triangle is right angled and also is isosceles} & \text{ triangle} \end{aligned}$$

259 (c)

By Standard results,


$$3. \quad r = 4R \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}$$

$$4. \quad r_1 = s \tan \frac{A}{2}$$

$$5. \quad r_3 = \frac{\Delta}{s-c}$$

260 (c)

On solving given equation of lines, we get the points $A(4, 0)$, $(0, 3)B(0, 3)$ and $O(0, 0)$



\therefore Area of ΔOAB

$$= \frac{1}{2} \times OA \times OB$$

$$= \frac{1}{2} \times 4 \times 3$$

$$= 6 \text{ sq units}$$

261 (a)

Let (x, y) denotes the coordinates in A, B and C plane

$$\text{Then, } \frac{(x-1)^2 + y^2}{(x+1)^2 + y^2} = \frac{1}{9}$$

$$\Rightarrow 9x^2 + 9y^2 - 18x + 9 = x^2 + y^2 + 2x + 1$$

$$\Rightarrow 8x^2 + 8y^2 - 20x + 8 = 0$$

$$\Rightarrow x^2 + y^2 - \frac{5}{2}x + 1 = 0$$

$$\therefore A, B, C \text{ lie on a circle with } C\left(\frac{5}{4}, 0\right)$$

262 (c)

Since, points (a, b) , (b, a) and $(a^2, -b^2)$ are collinear

$$\therefore \begin{vmatrix} a & b & 1 \\ b & a & 1 \\ a^2 & -b^2 & 1 \end{vmatrix} = 0$$

Applying, $R_2 \rightarrow R_2 - R_1$ and $R_3 \rightarrow R_3 - R_1$, we get

$$\begin{vmatrix} a & b & 1 \\ b-a & a-b & 0 \\ a^2-a & -b^2-b & 0 \end{vmatrix} = 0$$

$$\Rightarrow (a-b)(b^2 + b - a^2 + a) = 0$$

$$\Rightarrow (a-b)(a+b)(b+1-a) = 0$$

$$\Rightarrow \text{Either } a-b=0 \text{ or } (a+b)=0 \text{ or } (b+1-a) = 0$$

$$\Rightarrow a = b + 1$$

263 (d)


Let AA_1 , BB_1 and CC_1 be the towers and O be the circumcentre of ΔABC

$$\angle A_1OA = \theta_A, \angle B_1OB = \theta_B, \angle C_1OC = \theta_C$$

$$\text{Now, } OA = AA_1 \cot \theta_A$$

$$OB = BB_1 \cot \theta_B$$

$$\text{and } OC = CC_1 \cot \theta_C$$



Since, O is the circumcentre of a triangle

$$\therefore OA = OB = OC$$

$$\Rightarrow AA_1 \cot \theta_A = BB_1 \cot \theta_B = CC_1 \cot \theta_C$$

$$\Rightarrow \frac{\tan \theta_A}{AA_1} = \frac{\tan \theta_B}{BB_1} = \frac{\tan \theta_C}{CC_1}$$

Any other relationship between $\tan \theta_A$, $\tan \theta_B$ and $\tan \theta_C$ cannot be establish

264 (b)

$$\text{Let } a = 6, b = 5, c = \sqrt{13}$$

$$\therefore \cos C = \frac{6^2 + 5^2 - 13}{2 \times 6 \times 5} = \frac{4}{5}$$

$$\text{Now, } \sin C = \sqrt{1 - \frac{16}{25}} = \frac{3}{5}$$

$$\therefore \text{Area of } \Delta ABC = \frac{1}{2} ab \sin C$$

$$= \frac{1}{2} \times 6 \times 5 \times \frac{3}{5} = 9 \text{ sq unit}$$


265 (b)

Since, y_1, y_2, y_3 and x_1, x_2, x_3 are in AP
 $\therefore y_2 - y_1 = y_3 - y_2$ and $x_2 - x_1 = x_3 - x_2$
 So, $\frac{y_3 - y_2}{x_3 - x_2} = \frac{y_2 - y_1}{x_2 - x_1}$

So, points are collinear

266 (a)

Let $\angle RPQ = \theta$ and $\angle RQP = \phi$



$$\therefore \theta - \phi = 2\alpha$$

Let $RM \perp PQ$, so that $RM = k$,
 $MP = a - h$ and $MQ = a + h$

$$\text{Then, } \tan \theta = \frac{RM}{MP} = \frac{k}{a-h}$$

$$\text{and } \tan \phi = \frac{RM}{MQ} = \frac{k}{a+h}$$

Again now, $2\alpha = \theta - \phi$

$$\therefore \tan 2\alpha = \tan(\theta - \phi)$$

$$= \frac{\tan \theta - \tan \phi}{1 + \tan \theta \tan \phi}$$


$$= \frac{k(a+h) - k(a-h)}{a^2 - h^2 + k^2}$$

$$\Rightarrow a^2 - h^2 + k^2 = 2hk \cot 2\alpha$$

Hence, the locus is $x^2 - y^2 + 2xy \cot 2\alpha - a^2 = 0$

267 (c)

Since, the centroid G divide the line AD in the ratio 2:1



$$\therefore AG = \frac{8}{3} \text{ and } DG = \frac{4}{3}$$

$$\text{In } \triangle ABG, \tan \frac{\pi}{3} = \frac{AG}{BG}$$

$$\Rightarrow BG = AG \cot \frac{\pi}{3}$$

$$\Rightarrow BG = \frac{8}{3} \times \frac{1}{\sqrt{3}} = \frac{8}{3\sqrt{3}}$$

$$\text{Area of } \triangle ADB = \frac{1}{2} \times AD \times BG$$

$$= \frac{1}{2} \times 4 \times \frac{8}{3\sqrt{3}} = \frac{16}{3\sqrt{3}}$$


Since, median divides a triangle into two triangle of equal area. Therefore,

$$\text{Area of } \triangle ABC = 2 \times \text{area of } \triangle ADB$$

$$= 2 \times \frac{16}{3\sqrt{3}} = \frac{32}{3\sqrt{3}} \text{ sq units}$$

268 (b)

Since, Q is the image of P



$$\therefore PQ = 2PP''$$

$$= \frac{2|4 - (-1)|}{\sqrt{1^2 + 1^2}} = 5\sqrt{2}$$

269 (d)

$$\text{Area} = \frac{1}{2} \begin{vmatrix} a & b+c & 1 \\ b & c+a & 1 \\ c & a+b & 1 \end{vmatrix} = \frac{1}{2} \begin{vmatrix} a & a+b+c & 1 \\ b & a+b+c & 1 \\ c & a+b+c & 1 \end{vmatrix}$$

$$= \frac{(a+b+c)}{2} \begin{vmatrix} a & 1 & 1 \\ b & 1 & 1 \\ c & 1 & 1 \end{vmatrix} = 0$$

270 (c)

We have,

$$x^2 + 4xy + y^2 = aX^2 + bY^2$$

$$\Rightarrow (X \cos \theta + Y \sin \theta)^2$$

$$+ 14(X \cos \theta + Y \sin \theta)(X \sin \theta$$

$$- Y \cos \theta) + (X \sin \theta - Y \cos \theta)^2$$

$$= aX^2 + bY^2$$

$$\Rightarrow a = 1 + 4 \sin \theta \cos \theta, b = 1 - 4 \sin \theta \cos \theta \text{ and}$$

$$\sin^2 \theta - \cos^2 \theta = 0$$

$$\Rightarrow a = 1 + 4 \sin \theta \cos \theta, b = 1 - 4 \sin \theta \cos \theta \text{ and}$$

$$\theta = \frac{\pi}{4}$$

$$\Rightarrow a = 3, b = -1$$

271 (a)

Given, distance $r = \sqrt{2}$ and $\theta = 45^\circ$

$$\therefore x = 4 + \sqrt{2} \cos 45^\circ = 4 + \sqrt{2} \cdot \frac{1}{\sqrt{2}} = 5$$

$$\text{and } y = 3 + \sqrt{2} \sin 45^\circ = 3 + \sqrt{2} \cdot \frac{1}{\sqrt{2}} = 4$$


272 (d)

Let the height of the cliff be $BD = 50$ m and height of the tower be $AE = h$ metre.

In $\triangle DEC$,

$$\tan 30^\circ = \frac{50 - h}{x}$$


$$\Rightarrow x = \frac{50-h}{\frac{1}{\sqrt{3}}} = \sqrt{3}(50-h) \dots(i)$$



and in $\triangle BAD$,
 $\tan 45^\circ = \frac{50}{x} \Rightarrow x = 50 \text{ m}$
 From Eq. (i),
 $50 = \sqrt{3}(50 - h)$
 $\Rightarrow h = \frac{50(\sqrt{3} - 1)}{\sqrt{3}} \text{ m}$
 $\Rightarrow h = 50 \left(1 - \frac{\sqrt{3}}{3}\right) \text{ m}$

273 (a)

Let the vertex of triangle be $A(2, -1)$ and equation of BC is $x + 2y = 1$



Since, the triangle is equilateral triangle
 $\therefore \angle ABC = 60^\circ$

and $AD = \left| \frac{2(1) + 2(-1) - 1}{\sqrt{1+2^2}} \right| = \frac{1}{\sqrt{5}}$

In $\triangle ABD$, $\frac{AD}{AB} = \sin 60^\circ$

$\Rightarrow AB = \frac{1}{\frac{\sqrt{3}}{2}} = \frac{2}{\sqrt{3}}$

$\Rightarrow AB = BC = AC = \frac{2}{\sqrt{15}}$

275 (d)


Area of $\Delta = \frac{1}{2} \begin{vmatrix} 2 & 2 & 1 \\ 5 & 5 & 1 \\ 6 & 7 & 1 \end{vmatrix}$

$= \frac{1}{2} [2(5 - 7) - 2(5 - 6) + 1(35 - 30)]$

$= \frac{1}{2} (-4 + 2 + 5) = \frac{3}{2} \text{ sq units}$

276 (b)

Line perpendicular to OA passing through B is $x = 3$



Slope of $AB = \frac{3\sqrt{3}-0}{3-6} = -\sqrt{3}$

Line perpendicular to AB through origin is $y = \frac{1}{\sqrt{3}}x$

\therefore The point of intersection of a line $x = 3$ and $y = \frac{1}{\sqrt{3}}x$ is $(3, \sqrt{3})$, which is the required orthocenter

277 (a)

We know that, in any $\triangle ABC$

$\cos A + \cos B + \cos C = 1 + 4 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}$

and $r = 4R \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}$


$\therefore \cos A + \cos B + \cos C = 1 + \frac{r}{R}$

278 (b)

Given, $\sin P, \sin Q, \sin R$ are in AP

$\Rightarrow a, b, c$ are in AP

$\therefore \frac{\sin P}{a} = \frac{\sin Q}{b} = \frac{\sin R}{c} = \lambda$ (say)



Let p_1, p_2, p_3 be altitudes from P, Q, R

$\therefore p_1 = c \sin Q = \lambda bc$,

$p_2 = a \sin R = \lambda ac$,

$p_3 = b \sin P = \lambda ab$

Since, a, b, c are in AP,

$\therefore \frac{1}{a}, \frac{1}{b}, \frac{1}{c}$ are in HP

$\Rightarrow \frac{abc}{a}, \frac{abc}{b}, \frac{abc}{c}$ are in HP

$\Rightarrow bc, ac, ab$ are in HP


$\Rightarrow \lambda bc, \lambda ac, \lambda ab$ are in HP

$\Rightarrow p_1, p_2, p_3$ are in HP

280 (c)

In $\triangle ABD$, $\frac{BD}{\sin \theta} = \frac{AD}{\sin B}$

$\Rightarrow BD = AD = \frac{\sin \theta}{\sin B} \dots(i)$



In $\triangle ACD$, $\frac{CD}{\sin(A-\theta)} = \frac{AD}{\sin C}$

$$\Rightarrow CD = AD \frac{\sin(A - \theta)}{\sin C}$$

$$\therefore BD = CD$$

$$\Rightarrow AD \frac{\sin \theta}{\sin B} = AD \frac{\sin(A - \theta)}{\sin C}$$

$$\Rightarrow \sin(A - \theta) = \frac{\sin C}{\sin B} \sin \theta = \frac{c}{b} \sin \theta$$

282 (b)

Since, circle is inscribed in the quadrilateral $ABCD$, then

$$a + c = b + d \dots(i)$$

And quadrilateral is cyclic, then

$$A + C = \pi$$

$$\text{or } C = \pi - A \dots(ii)$$

now, in ΔABD

$$\cos A = \frac{a^2 + d^2 - (BD)^2}{2ad}$$

$$\Rightarrow BD^2 = a^2 + d^2 - 2ad \cos A \dots(iii)$$

And in ΔBCD ,

$$\cos C = \frac{b^2 + c^2 - (BD)^2}{2bc}$$

$$\Rightarrow (BD)^2 = b^2 + c^2 - 2bc \cos C$$

$$= b^2 + c^2 - 2bc \cos(\pi - A) \text{ [from Eq.(ii)]}$$

$$\Rightarrow (BD)^2 = b^2 + c^2 + 2bc \cos A \dots(iv)$$

From Eqs. (iii) and (iv), we get

$$\cos A = \frac{a^2 + d^2 - b^2 - c^2}{2(bc + ad)} \dots(v)$$

Now, from Eq. (i),

$$(a - d)^2 = (b - c)^2$$

$$\Rightarrow a^2 + d^2 - b^2 - c^2 = 2(ad - bc) \dots(vi)$$

\therefore From Eqs. (v) and (vi),

$$\cos A = \frac{(ad - bc)}{(ad + bc)}$$

283 (c)

We have, $a + b + c = \lambda = 2s$

$$\therefore s = \frac{\lambda}{2}$$

$$b \cos^2 \frac{C}{2} + c \cos^2 \frac{B}{2} = b \frac{s(s-c)}{ab} + c \frac{s(s-b)}{ac}$$

$$= \frac{s}{a} \{s - c + s - b\}$$

$$= \frac{s}{a} \cdot a = s = \frac{\lambda}{2}$$

284 (c)

$$\cos B = \frac{1}{\sqrt{2}} = \frac{c^2 + a^2 - b^2}{2ac}$$

$$\Rightarrow \sqrt{2}ac = c^2 + a^2 - b^2$$

$$\Rightarrow 2a^2c^2 = c^4 + a^4 + b^4 + 2c^2a^2$$

$$- 2(c^2 + a^2)b^2$$

$$\Rightarrow c^4 + a^4 + b^4 = 2(c^2 + a^2)b^2$$

285 (a)

Let the point of the line divides the line in the ratio $m:1$

$$\therefore \text{Coordinates of point are } \left(\frac{5m+1}{m+1}, \frac{7m-1}{m+1} \right)$$

Which lies on $y + x = 4$

$$\Rightarrow \frac{7m-1}{m+1} + \frac{5m+1}{m+1} = 4$$

$$\Rightarrow \frac{12m}{m+1} = 4 \Rightarrow 12m = 4m + 4$$

$$\Rightarrow 8m = 4 \Rightarrow m = \frac{1}{2}$$

\therefore Required ratio is 1:2


286 (d)

Since, line perpendicular to OA passing through B is $x = 4$

$$\text{Slope of } AB = -\frac{3}{2}$$

Line perpendicular to AB through origin is, $y =$

$$\frac{2}{3}x$$



\therefore The point of intersection of lines $x = 4$

And $y = \frac{2}{3}x$ is $(4, \frac{8}{3})$. Which is the required

orthocenter

288 (a)

$$\text{We have, } \frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$$

$$\therefore \frac{b+c}{a} = \frac{\sin B + \sin C}{\sin A}$$

$$= \frac{\sin\left(\frac{B+C}{2}\right) \cos\left(\frac{B-C}{2}\right)}{\sin\frac{A}{2} \cos\frac{A}{2}} = \frac{\cos\left(\frac{B-C}{2}\right)}{\sin\frac{A}{2}}$$

$$[A + B + C = \pi \Rightarrow A = \pi - (B + C)]$$

289 (a)

Let h be the mid point of BC . Since, $\angle TBH = 90^\circ$, then $TH^2 = BT^2 + BH^2 = 5^2 + 5^2 = 50$

Also since,


$$\angle THG = 90^\circ, TG^2 = TH^2 + GH^2 = 50 + 25 = 75$$

Let θ be the required angle of elevation of G at T

$$\text{Then, } \sin \theta = \frac{GH}{TG}$$

$$= \frac{5}{5\sqrt{3}} = \frac{1}{\sqrt{3}}$$

$$\Rightarrow \theta = \sin^{-1}(1/\sqrt{3})$$



290 (d)


$$\text{Given, } \Delta = \frac{1}{2}ap_1 = \frac{1}{2}bp_2 = \frac{1}{2}cp_3$$

$$\therefore p_1 = \frac{2\Delta}{a}, p_2 = \frac{2\Delta}{b}, p_3 = \frac{2\Delta}{c}$$

$$\therefore p_1 p_2 p_3 = \frac{8\Delta^3}{abc} = \frac{8}{abc} \left(\frac{abc}{4R}\right)^3 = \frac{a^2 b^2 c^2}{8R^2}$$

291 (b)

$$\text{In } \Delta BCD, \cot 60^\circ = \frac{BC}{300}$$



$$\Rightarrow BC = 300 \times \frac{1}{\sqrt{3}} \dots(i)$$

$$\text{In } \Delta ACD, \cot 30^\circ = \frac{AC}{300}$$

$$\Rightarrow AC = 300\sqrt{3} \dots(ii)$$

$$\therefore \text{Distance between two boats} = AB \\ = AC - BC = 300\sqrt{3} - \frac{300}{\sqrt{3}} \quad [\text{using Eqs.(i) and (ii)}]$$

$$= 300 \frac{(3-1)}{\sqrt{3}} = \frac{600 \times \sqrt{3}}{3} = 346.4 \text{ m}$$

292 (d)

$$\therefore 3^2 + 4^2 = 5^2$$

\Rightarrow Given triangle is a right angled triangle whose length of hypotenuse is 5 unit

$$\therefore R = \frac{5}{2} = 2.5$$

293 (a)

$$\text{Given, } \left(1 - \frac{r_1}{r_2}\right) \left(1 - \frac{r_1}{r_3}\right) = 2$$

$$\therefore \left(1 - \frac{s-b}{s-a}\right) \left(1 - \frac{s-c}{s-a}\right) = 2$$

$$\Rightarrow \frac{(b-a)(c-a)}{(s-a)^2} = 2$$

$$\Rightarrow a^2 = b^2 + c^2$$

Hence, triangle is a right angled triangle

294 (a)

From given, we get

$$a \left(\tan A - \tan \frac{A+B}{2} \right) = b \left(\tan \frac{A+B}{2} - \tan B \right)$$

$$\Rightarrow a \cdot \frac{\sin \left(A - \frac{A+B}{2} \right)}{\cos A \cdot \cos \frac{A+B}{2}} = b \cdot \frac{\sin \left(\frac{A+B}{2} - B \right)}{\cos \frac{A+B}{2} \cos B}$$

$$\Rightarrow \frac{a \sin \left(\frac{A-B}{2} \right)}{\cos A} = \frac{b \sin \left(\frac{A-B}{2} \right)}{\cos B}$$

$$\Rightarrow \sin \left(\frac{A-B}{2} \right) \{ a \cos B - b \cos A \} = 0$$

$$\Rightarrow \text{Either } \sin \frac{A-B}{2} = 0 \text{ or } a \cos B = b \cos A$$

$$\text{When, } \sin \frac{A-B}{2} = 0 \Rightarrow A = B$$

$$\text{or when } a \cos B = b \cos A$$


$$\Rightarrow \sin A \cos B - \cos A \sin B = 0$$

$$\Rightarrow \sin(A-B) = 0$$

$$\Rightarrow A = B$$

295 (d)

Triangle is right angled triangle. In right angled triangle mid point of hypotenuse is circumcentre
So, coordinates of the circumcentre are (5, 2)




296 (c)

$$\text{From } \Delta ABC, \frac{\sin(y+z)}{DC} = \frac{\sin C}{AD}$$

$$\text{From } \Delta ABD, \frac{\sin x}{BD} = \frac{\sin B}{AD}$$

$$\text{From } \Delta AEC, \frac{\sin z}{EC} = \frac{\sin C}{AE}$$

$$\text{From } \Delta ABE, \frac{\sin(x+y)}{BE} = \frac{\sin B}{AE}$$



$$\therefore \frac{\sin(x+y) \sin(y+z)}{\sin x \sin z} = \frac{\frac{BE \sin B}{AE} \times \frac{DC \sin C}{AD}}{\frac{BD \sin B}{AD} \times \frac{EC \sin C}{AE}}$$


$$= \frac{BE}{AE} \times \frac{DC}{AD} \times \frac{AD}{BD} \times \frac{AE}{EC} \\ = \frac{2BD \times 2EC}{BD \times EC} = 4$$

297 (c)

$$\text{In } \Delta ABC, \tan 60^\circ = \frac{h}{BC}$$

$$\Rightarrow BC = h \cot 60^\circ$$

$$\text{In } \Delta ABD, \tan 30^\circ = \frac{h}{BD}$$



$$\begin{aligned} \Rightarrow BD &= h \cot 30^\circ \\ \Rightarrow BC + CD &= h \cot 30^\circ \\ \Rightarrow CD &= h \cot 30^\circ - BC \\ \Rightarrow d &= h \cot 30^\circ - h \cot 60^\circ \quad [\text{from Eq. (i)}] \end{aligned}$$

$$\begin{aligned} \therefore \text{Speed of car} &= \frac{\text{distance from D to C}}{\text{time taken}} \\ &= \frac{d}{3} = \frac{h \cot 30^\circ - h \cot 60^\circ}{3} \end{aligned}$$

$$\therefore \text{Time taken from C to B} = \frac{\text{distance}}{\text{speed}}$$


$$= \frac{h \cot 60^\circ}{h(\cot 30^\circ - \cot 60^\circ)}$$

$$= \frac{\frac{1}{\sqrt{3}} \times 3}{\left(\sqrt{3} - \frac{1}{\sqrt{3}}\right)}$$

$$= \frac{3}{2} = 1.5 \text{ min}$$

298 (a)

Let h be the height of the tower, then $h = AQ \tan \alpha = BQ \tan \beta = CQ \tan \gamma$



$$\Rightarrow BC = BQ - CQ = h(\cot \beta - \cot \gamma),$$

$$CA = h(\cot \alpha - \cot \gamma)$$


$$\text{and } AB = h(\cot \alpha - \cot \beta)$$

$$\begin{aligned} \text{Now, } BC \cot \alpha - CA \cot \beta + AB \cot \gamma &= h[\cot \alpha (\cot \beta - \cot \gamma) - \cot \beta (\cot \alpha - \cot \gamma) \\ &\quad + \cot \gamma (\cot \alpha - \cot \beta)] \end{aligned}$$

$$= 0$$

299 (a)

On solving the given equations of sides, we get the coordinates of the vertices of the triangle, $A(2, -1)$, $B(6, -1)$ and $C(2, 1)$




The circumcentre of ΔABC is the mid point of BC

$$\text{ie, } M \equiv \left(\frac{8}{2}, \frac{0}{2}\right) = (4, 0)$$

300 (b)

Let h metres be the height of tree CD and x meters be the width of river



$$\text{In } \Delta BCD, \tan 60^\circ = \frac{h}{x} \Rightarrow h = \sqrt{3}x \dots\dots (i)$$

$$\text{and in } \Delta ACD, \tan 30^\circ = \frac{h}{x+20}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{h}{x+20}$$

$$\Rightarrow 3x = x + 20 \quad [\text{from Eq. (i)}]$$

$$x = 10 \text{ m}$$

302 (a)

We have,

$$2s = a + b + c, A^2 = s(s-a)(s-b)(s-c)$$

$$\therefore AM \geq GM$$

$$\Rightarrow \frac{s-a+s-b+s-c}{3}$$

$$\geq \sqrt[3]{(s-a)(s-b)(s-c)}$$


$$\Rightarrow \frac{3s-2s}{3} \geq \frac{(A^2)^{1/3}}{s^{1/3}}$$

$$\Rightarrow \frac{s^3}{27} \geq \frac{A^2}{s} \Rightarrow A \leq \frac{s^2}{3\sqrt{3}}$$

303 (b)

$$\text{In } \Delta DBC, \tan \alpha = \frac{5}{x} \dots(i)$$

$$\text{and in } \Delta DAC, \tan 2\alpha = \frac{30}{x}$$




$$\Rightarrow \frac{2 \tan \alpha}{1 - \tan^2 \alpha} = \frac{30}{x}$$

$$\begin{aligned} \Rightarrow \frac{2\left(\frac{5}{x}\right)}{1 - \frac{25}{x^2}} &= \frac{30}{x} \\ \Rightarrow \frac{10x}{x^2 - 25} &= \frac{30}{x} \\ \Rightarrow 10x^2 &= 30x^2 - 750 \\ \Rightarrow 20x^2 &= 750 \\ \Rightarrow x^2 &= \frac{75}{2} \\ \Rightarrow x &= 5\sqrt{\frac{3}{2}} \text{ m} \end{aligned}$$

305 (a)

Here, ratio of angles are 4:1:1
 $\Rightarrow 4x + x + x = 180^\circ$
 $\Rightarrow x = 30^\circ$
 $\therefore \angle A = 120^\circ, \quad \angle B = \angle C = 30^\circ$



Thus, the ratio of longest side to the perimeter =

$$\begin{aligned} \frac{a}{a+b+c} \\ \text{Let } b = c = x \\ \therefore a^2 = b^2 + c^2 - 2bc \cos A \\ \Rightarrow a^2 = 2x^2 - 2x^2 \cos A = 2x^2 (1 - \cos A) \\ \Rightarrow a^2 = 4x^2 \sin^2 \frac{A}{2} \\ \Rightarrow a = 2x \sin \frac{A}{2} \\ \Rightarrow a = 2x \sin 60^\circ = \sqrt{3}x \end{aligned}$$

Thus, required ratio is

$$\frac{a}{a+b+c} = \frac{\sqrt{3}x}{x+x+\sqrt{3}x} = \frac{\sqrt{3}}{2+\sqrt{3}}$$

306 (c)

Let the third angle is θ

In a triangle, $\frac{\pi}{4} + \tan^{-1} 2 + \theta = \pi$

$$\Rightarrow \theta = \pi - \frac{\pi}{4} - \tan^{-1} 2$$

$$\Rightarrow \tan \theta = \tan \left[\pi - \left(\frac{\pi}{4} + \tan^{-1} 2 \right) \right]$$


$$= -\tan \left(\frac{\pi}{4} + \tan^{-1} 2 \right)$$

$$\Rightarrow \tan \theta = -\frac{1+2}{1-2} = 3$$

$$\Rightarrow \theta = \tan^{-1} 3$$

307 (d)

$$\text{In } \triangle BCD, \tan 60^\circ = \frac{CD}{BC} = \frac{h}{x} \Rightarrow h = \sqrt{3}x \dots(i)$$



and in $\triangle ACD$,

$$\tan 30^\circ = \frac{CD}{AC} = \frac{h}{x+40} \Rightarrow \frac{1}{\sqrt{3}} = \frac{h}{40+x}$$

$$40+x = h\sqrt{3}$$


$$\Rightarrow 40+x = 3x \quad [\text{using Eq.(i)}]$$

$$\Rightarrow x = 20$$

On putting this value in Eq.(i), we get $h = 20\sqrt{3}$ m

308 (d)

Let $H(3, \alpha)$ is the orthocenter



\therefore Slope of $BH \times$ Slope of $AC = -1$

$$\Rightarrow -\alpha \cdot \frac{4}{3} = -1$$

$$\Rightarrow \alpha = \frac{3}{4}$$

Hence, orthocenter of a triangle is $\left(3, \frac{3}{4}\right)$

309 (c)


Let the axes be rotated through an angle θ . Then,

$$\tan 2\theta = \frac{2h}{a-b} = \frac{4\sqrt{3}}{5-9} = -\sqrt{3} \Rightarrow 2\theta = \frac{2\pi}{3} \Rightarrow \theta = \frac{\pi}{3}$$

310 (a)

Since, $AC = 2BC$


\therefore Coordinates of C are



$$\left(\frac{4-3}{2+1}, \frac{2+4}{2+1} \right) \text{ ie, } \left(\frac{1}{3}, 2 \right)$$

311 (d)

Let P be the image of the origin about the line $x + y = 1$. Since, $OA = OB$, therefore Q is mid point of AB



∴ Coordinates of Q are $(\frac{1}{2}, \frac{1}{2})$

Let the coordinates of P are (x_1, y_1)

Also, Q is the mid point of OP

$$\therefore \frac{0+x_1}{2} = \frac{1}{2} \text{ and } \frac{0+y_1}{2} = \frac{1}{2}$$

$$\Rightarrow x_1 = 1, \quad y_1 = 1$$

∴ The coordinates of P are $(1, 1)$

312 (c)

We have, $\operatorname{cosec} A(\sin B \cos C + \cos B \sin C)$

$$= \left(\frac{\sin B}{\sin A} \cos C + \frac{\sin C}{\sin A} \cos B \right)$$

$$= \left(\frac{b}{a} \cos C + \frac{c}{a} \cos B \right) = 1$$

(∵ $a = b \cos C + c \cos B$)

313 (b)

Let coordinates of fourth vertex are (x, y)

$$\therefore \frac{x-5}{2} = \frac{7+3}{2}$$

$$\Rightarrow x = 15$$


$$\text{and } \frac{y-4}{2} = \frac{10+5}{2}$$

$$\Rightarrow y = 19$$

∴ Coordinates of fourth vertex are $(15, 19)$

314 (a)

Let AB and DE be two towers of equal height ' h '



$$\therefore \text{In } \triangle ABC, \tan 60^\circ = \frac{h}{x} \Rightarrow x = \frac{h}{\sqrt{3}} \dots (i)$$

$$\text{Again, in } \triangle CDE, \tan 30^\circ = \frac{h}{60-x}$$

$$\Rightarrow 60 - x = h\sqrt{3}$$

$$\Rightarrow 60 = h\sqrt{3} + \frac{h}{\sqrt{3}} \text{ [from Eq. (i)]}$$


$$\Rightarrow h = 15\sqrt{3} \text{ m}$$

315 (d)

$$\text{In } \triangle DEC, \tan \theta = \frac{DE}{CE}$$

$$\Rightarrow \tan \theta = \frac{h-a}{CE}$$

$$\Rightarrow CE = \frac{h-a}{\tan \theta}$$



In $\triangle ABC$,

$$\tan 2\theta = \frac{a}{AB} = \frac{a}{CE} \quad (\because CE = AB)$$

$$\Rightarrow \frac{2 \tan \theta}{1 - \tan^2 \theta} = \frac{a}{h-a} \text{ [from Eq.(i)]}$$

$$\Rightarrow a \tan \theta = (h-a) \times \frac{2 \tan \theta}{1 - \tan^2 \theta}$$

$$\Rightarrow a \tan \theta (1 - \tan^2 \theta) - 2(h-a) \tan \theta = 0$$

$$\Rightarrow \tan \theta (a - a \tan^2 \theta - 2h + 2a) = 0$$

∵ $\tan \theta \neq 0$

$$\therefore \tan^2 \theta = \frac{-2h + 3a}{a}$$

$$\Rightarrow \theta = \tan^{-1} \sqrt{\frac{-2h + 3a}{a}}$$

$$= \tan^{-1} \sqrt{3 - \frac{2h}{a}}$$

316 (c)

Given vertices are $A(2a, 4a), B(2a, 6a)$ and $C(2a + \sqrt{3}a, 5a), a > 0$

$$\text{Now, } AB = \sqrt{(2a + 2a)^2 + (4a - 6a)^2} = 2a$$

$$BC = \sqrt{(\sqrt{3}a)^2 + a^2} = 2a$$


$$\text{and } CA = \sqrt{(\sqrt{3}a)^2 + (-a)^2} = 2a$$

$$\therefore AB = BC = CA$$

Hence, triangle is an equilateral triangle, therefore it is an acute angled triangle

317 (a)

Let D and E are the mid points of AB and AC . So, coordinates of B and C are $(-3, 3)$ and $(5, 3)$ respectively




$$\text{Centroid of triangle} = \left(\frac{1-3+5}{3}, \frac{1+3+3}{3} \right)$$

$$= \left(1, \frac{7}{3} \right)$$


318 (b)

R is mid point of PT



Point $(\frac{5a+3b}{8}, \frac{5x+3y}{8})$ divides PT in ratio 3 : 5 and that is mid point of QR

319 (a)



On solving the given equations of lines, we get the coordinates of the vertices of a ΔOPQ which are $O(0,0)$, $P(1,0)$ and $Q(0,1)$. Since, the triangle is right angled at $O(0,0)$, therefore $O(0,0)$ is its orthocenter

320 (d)

Let the circumcentre of triangle be $P(x, y)$ and let the vertices of a ΔABC be $A(0, 30)$, $B(4, 0)$ and $C(30, 0)$

$$\therefore PA^2 = PB^2 = PC^2$$

$$\Rightarrow (x-0)^2 + (y-30)^2 = (x-4)^2 + (y-0)^2 = (x-30)^2 + (y-0)^2$$

From Ist and IInd terms,

$$x^2 + y^2 - 60y + 900 = x^2 + y^2 - 8x + 16 \Rightarrow 8x - 60y + 884 = 0 \dots(i)$$

From IInd and IIIrd terms,

$$x^2 - 8x + 16 + y^2 = x^2 - 60x + 900 + y^2 \Rightarrow 52x = 884 \Rightarrow x = 17$$

On putting $x = 17$ in Eq. (i), we get


$$y = 17$$

Hence, required point is $(17, 17)$

321 (b)

$$\text{Since, } \tan \theta = \frac{1}{2}$$

$$\text{In } \Delta ABC, \tan \alpha = \frac{\frac{h}{3}}{AB} = \frac{h}{120} \dots(i)$$



$$\text{In } \Delta ADB, \tan \beta = \frac{3h}{120} \dots(ii)$$

$$\therefore \tan \theta = \tan(\beta - \alpha)$$

$$\Rightarrow \tan \theta = \frac{\tan \beta - \tan \alpha}{1 - \tan \beta \tan \alpha}$$

$$\Rightarrow \frac{1}{2} = \frac{\frac{3h}{120} - \frac{h}{120}}{1 + \frac{3h^2}{14400}} \quad (\because \tan \theta = \frac{1}{2}, \text{ given})$$

$$\Rightarrow \frac{1}{2} = \frac{\frac{2h}{120}}{\frac{14400+3h^2}{14400}}$$

$$\Rightarrow \frac{1}{2} = \frac{240h}{14400+3h^2}$$

$$\Rightarrow 14400 + 3h^2 = 480h$$

$$\Rightarrow 4800 + h^2 - 160h = 0$$

$$\Rightarrow (h-40)(h-120) = 0$$

Since, the height of the pole is more than 100 m

$$\therefore h = 120 \text{ ft}$$

322 (c)

Slope of line $OP = \frac{3}{4}$, let new position is $Q(x, y)$

$$\text{slope of } OQ = \frac{y}{x} \text{ also } x^2 + y^2 = OQ^2 = 25 = (OP^2)$$

$$\tan 45^\circ = \left| \frac{\frac{y}{x} - \frac{3}{4}}{1 + \frac{3y}{4x}} \right| \Rightarrow \pm 1 = \frac{4y-3x}{4x+3y}$$

Straight lines and pair of straight lines

$$4x + 3y = 4y - 3x$$

$$\text{or } -4x - 3y = 4y - 3x$$

$$x = \frac{1}{7}y \dots(i)$$

$$-x = 7y \dots(ii)$$

Correct relation is $x = \frac{1}{7}y$ as new point must lie in


Ist quadrant

$$x^2 + 49x^2 = 25$$

$$\Rightarrow x = -\frac{1}{\sqrt{2}}, y = \frac{7}{\sqrt{2}}$$

323 (c)

In ΔABC , $\tan 30^\circ = \frac{BC}{AC}$



$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{h-150}{60}$$

$$\Rightarrow h = (150 + 20\sqrt{3}) \text{ m}$$

324 (b)

Since, $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = k$ (say)

$$\therefore (a+b+c)(a+b-c) = 3ab$$


$$\Rightarrow a^2 + b^2 + 2ab - c^2 = 3ab$$

$$\Rightarrow \frac{a^2 + b^2 - c^2}{2ab} = \frac{1}{2} \Rightarrow \cos C = \cos \frac{\pi}{3}$$

$$\Rightarrow \angle C = \frac{\pi}{3}$$

325 (b)

$$\text{In } \triangle ABD, \tan \alpha = \frac{H}{d}$$



$$\Rightarrow d = H \cot \alpha \dots(i)$$

$$\text{In } \triangle ECD, \tan(\alpha - \beta) = \frac{H-h}{d}$$

$$\Rightarrow \tan(\alpha - \beta) = \frac{H-h}{H \cot \alpha} \quad [\text{from Eq.(i)}]$$

$$\Rightarrow H[1 - \cot \alpha \tan(\alpha - \beta)] = h$$

$$\Rightarrow H = \frac{h \cot(\alpha - \beta)}{\cot(\alpha - \beta) - \cot \alpha}$$

326 (b)

Let the equation of line be

$$\frac{x}{a} + \frac{y}{b} = 1$$

Since, it passes through $(\frac{1}{5}, \frac{1}{5})$

$$\therefore \frac{1}{5a} + \frac{1}{5b} = 1$$

$$\Rightarrow a + b = 5ab \dots(i)$$

Since, the point $P(x, y)$ divides AB joining $A(a, 0)$ and $B(0, b)$ internally in ratio 3:1

$$\therefore x = \frac{a}{4}, y = \frac{3b}{4} \Rightarrow a = 4x \text{ and } b = \frac{4y}{3}$$

On putting the value of a and b in Eq. (i), we get

$$4x + \frac{4y}{3} = 5(4x) \left(\frac{4y}{3}\right)$$

$$\Rightarrow 3x + y = 20xy$$

327 (d)


$$(3 - 10 + a)(9 - 20 + a) > 0$$

$$\text{or } (a - 7)(a - 11) > 0$$

$$\therefore a \in (-\infty, 7) \cup (11, \infty)$$

328 (a)

$$y = |x - 2| \Rightarrow y = x - 2 \text{ and } y = 2 - x$$



$$\text{Area of shaded region} = 2 \triangle ABC = 2 \cdot \frac{1}{2} \cdot 1 \cdot 1 = 1$$

329 (d)

$$\frac{\cot \frac{A}{2} \cot \frac{B}{2} - 1}{\cot \frac{a}{2} \cot \frac{b}{2}} = 1 - \tan \frac{A}{2} \tan \frac{B}{2}$$

$$= 1 - \sqrt{\frac{(s-b)(s-c)}{s(s-a)}} \sqrt{\frac{(s-a)(s-c)}{s(s-b)}}$$

$$= 1 - \frac{s-c}{s} = \frac{c}{s}$$

$$= \frac{2c}{a+b+c}$$


330 (c)

We have, $AG = 1$

$$\therefore GD = \frac{1}{2} AG = \frac{1}{2}$$

Hence, the coordinates of D are $(1, 1/2)$

Clearly, AG is parallel to y -axis and the triangle ABC is equilateral. Therefore, BC is parallel to x -axis at a distance of $\frac{1}{2}$ unit from it



Let a be the length of each side of $\triangle ABC$. Then,

$$AD = \frac{\sqrt{3}}{2} a \Rightarrow \frac{\sqrt{3}}{2} a = \frac{3}{2} \Rightarrow a = \sqrt{3}$$

Since D is the mid-point of BC and $BC = \sqrt{3}$

$$\therefore BD = CD = \frac{\sqrt{3}}{2}$$

Hence, the coordinates of B and C are $(1 - \frac{\sqrt{3}}{2}, \frac{1}{2})$

and $(1 + \frac{\sqrt{3}}{2}, \frac{1}{2})$ respectively

331 (d)

$$c^2 \sin 2B + b^2 \sin 2C$$

$$= c^2 (2 \sin B \cos B) + b^2 (2 \sin C \cos C)$$


$$= 2c^2 \left(\frac{2\Delta}{ac} \cos B\right) + 2b^2 \left(\frac{2\Delta}{ab} \cos C\right)$$

$$= 4\Delta \left(\frac{c \cos B + b \cos C}{a}\right)$$

$$= 4\Delta \left(\frac{a}{a}\right) = 4\Delta$$

332 (a)

$$\text{In } \triangle DAB, \tan \theta = \frac{64}{d}$$



$$\Rightarrow d = 64 \cot \theta \quad \dots(i)$$

$$\text{In } \triangle CDE, \tan(90^\circ - \theta) = \frac{(100-64)}{d}$$

$$\Rightarrow d = 36 \tan \theta \quad \dots(ii)$$

From Eqs. (i) and (ii), we get

$$d^2 = 36 \times 64 \Rightarrow d = 48 \text{ m}$$

334 (d)

We know that area of circle is πr^2

If radius, $r = a$, then $A = \pi a^2$

And the area of the segment of angle $2\pi = \pi a^2$

$$\therefore \text{Area of 1 angle} = \frac{\pi a^2}{2\pi}$$

$$\therefore \text{Area of } 2\alpha \text{ angle} = \frac{2\alpha\pi a^2}{2\pi} = \alpha a^2$$

336 (a)

Let $a = 5k, b = 6k$ and $c = 5k$

$$s = \frac{5k + 6k + 5k}{2} = 8k$$

$$\therefore r = \frac{\Delta}{s}$$

$$= \sqrt{\frac{8k(8k-5k)(8k-6k)(8k-5k)}{8k}}$$


$$= \sqrt{\frac{8k \cdot 3k \cdot 2k \cdot 3k}{8k}} = \frac{3k}{2}$$

$$\Rightarrow k = \frac{2r}{3} = \frac{2 \times 6}{3} = 4$$

338 (b)

Let the height of the vertical tower $PQ = h, C$ is the middle point of line segment AB . Since, PQ is perpendicular to the plane QAB ,

$\therefore \angle PQA = \angle PQC = \angle PQB = 90^\circ$, we get



$$\frac{PQ}{QA} = \tan \theta \Rightarrow QA = h \cot \theta$$

Similarly, $QB = h \cot \theta$ and $QC = h \cot \phi$

Since, $QA = QB$, the $\triangle QAB$ is an isosceles triangle

Here, QCA is a right angled triangle in which

$$\angle QCA = 90^\circ$$

$$\therefore OC^2 + AC^2 = QA^2$$

$$\Rightarrow h^2 \cot^2 \phi + a^2 = h^2 \cot^2 \theta$$

$$\text{or } h^2 = \frac{a^2}{\cot^2 \theta - \cot^2 \phi}$$

$$\Rightarrow h^2 = \frac{a^2}{(\operatorname{cosec}^2 \theta - 1) - (\operatorname{cosec}^2 \phi - 1)}$$

$$= \frac{a^2}{\operatorname{cosec}^2 \theta - \operatorname{cosec}^2 \phi}$$

$$= \frac{a^2}{\frac{1}{\sin^2 \theta} - \frac{1}{\sin^2 \phi}}$$

$$\Rightarrow h^2 = \frac{a^2 \sin^2 \theta \sin^2 \phi}{\sin^2 \phi - \sin^2 \theta} = \frac{a^2 \sin^2 \theta \sin^2 \phi}{\sin(\phi - \theta) \sin(\phi + \theta)}$$

Hence, the required height h of the peak

$$= \frac{a \sin \theta \sin \phi}{\sqrt{\sin(\phi + \theta) \sin(\phi - \theta)}}$$

339 (b)

$$\text{Given, } \frac{\cos A}{K \sin A} = \frac{\cos B}{K \sin B} = \frac{\cos C}{K \sin C}$$

$$\Rightarrow \cot A = \cot B = \cot C$$

$$\Rightarrow A = B = C = 60^\circ$$


$\Rightarrow \triangle ABC$ is an equilateral triangle.

$$\therefore \Delta \frac{\sqrt{3}}{4} a^2 = \frac{\sqrt{3}}{4} \times \frac{1}{6} = \frac{\sqrt{3}}{24} \text{ sq unit}$$

340 (c)

$$\therefore \frac{a_1}{b_1} = \tan\left(\frac{\alpha + \beta}{2}\right) \quad \dots(i)$$

$$\text{and } \frac{a_2}{b_2} = \tan\left(\frac{\beta + \gamma}{2}\right) \quad \dots(ii)$$



Since, $a_1 a_2 = b_1 b_2$

$$\Rightarrow \frac{a_1}{b_1} = \frac{b_2}{a_2}$$

$$\Rightarrow \tan\left(\frac{\alpha + \beta}{2}\right) = \frac{1}{\tan\left(\frac{\beta + \gamma}{2}\right)}$$

$$\Rightarrow \tan\left(\frac{\alpha + \beta}{2}\right) \tan\left(\frac{\beta + \gamma}{2}\right) = 1$$

$$\therefore \frac{\alpha + \beta}{2} + \frac{\beta + \gamma}{2} = \frac{\pi}{2}$$

$$\Rightarrow \alpha + \beta + \gamma = \pi - \beta < \pi$$

341 (c)

We know that, $2s = a + b + c$

$$\begin{aligned} \therefore \frac{(a+b+c)(b+c-a)(c+a-b)(a+b-c)}{4b^2c^2} \\ = \frac{2s(2s-2a)(2s-2b)(2s-2c)}{4b^2c^2} \\ = 4 \frac{s(s-a)}{bc} \times \frac{(s-b)(s-c)}{bc} \\ = 4 \cos^2 \frac{A}{2} \times \sin^2 \frac{A}{2} = \sin^2 A \end{aligned}$$

342 (b)

Let $P(h, k)$ be the required point, then $2PA = 3PB$

$$\begin{aligned} \Rightarrow 4PA^2 &= 9PB^2 \\ \Rightarrow 4[(h-0)^2 + (k-0)^2] &= 9[(h-4)^2 + (k+3)^2] \\ \Rightarrow 4(h^2 + k^2) &= 9[h^2 + 16 - 8h + k^2 + 9 + 6k] \\ \Rightarrow 5h^2 + 5k^2 - 72h + 54k + 225 &= 0 \\ \therefore \text{Required locus of } P(h, k) \text{ is} \\ 5x^2 + 5y^2 - 72x + 54y + 225 &= 0 \end{aligned}$$

343 (a)

$$\text{Here, } a = \sqrt{(16-5)^2 + (12-12)^2} = 11$$

$$b = \sqrt{(16-0)^2 + (12-0)^2} = 20$$

$$\text{And } c = \sqrt{(5-0)^2 + (12-0)^2} = 13$$

\therefore Incentre =

$$\left(\frac{11 \times 0 + 20 \times 5 + 13 \times 16}{11+20+13}, \frac{11 \times 0 + 20 \times 12 + 13 \times 12}{11+20+13} \right) = (7, 9)$$

344 (d)

$$\text{We have, } \tan \frac{A}{2} \tan \frac{C}{2} = \frac{1}{2}$$

$$\Rightarrow \sqrt{\frac{(s-b)(s-c)}{s(s-a)}} \sqrt{\frac{(s-a)(s-b)}{s(s-c)}} = \frac{1}{2}$$

$$\Rightarrow \frac{s-b}{s} = \frac{1}{2} \Rightarrow 2s - 2b - s = 0$$

$$\Rightarrow a + c - 3b = 0$$

345 (c)

Since, $\sin^2 \frac{A}{2}, \sin^2 \frac{B}{2}, \sin^2 \frac{C}{2}$ be in HP

$$\Rightarrow \frac{1}{\sin^2 \frac{A}{2}}, \frac{1}{\sin^2 \frac{B}{2}}, \frac{1}{\sin^2 \frac{C}{2}} \text{ are in AP}$$

$$\Rightarrow \frac{1}{\sin^2 \frac{C}{2}} - \frac{1}{\sin^2 \frac{B}{2}} = \frac{1}{\sin^2 \frac{B}{2}} - \frac{1}{\sin^2 \frac{A}{2}}$$

$$\Rightarrow \frac{ab}{(s-a)(s-b)} - \frac{ac}{(s-a)(s-c)} = \frac{ac}{(s-a)(s-c)} - \frac{bc}{(s-b)(s-c)}$$

$$\begin{aligned} \Rightarrow \left(\frac{a}{s-a} \right) \left(\frac{b(s-c) - c(s-b)}{(s-b)(s-c)} \right) \\ = \left(\frac{c}{s-c} \right) \left(\frac{a(s-b) - b(s-a)}{(s-a)(s-b)} \right) \end{aligned}$$

$$\begin{aligned} \Rightarrow abs - abc - acs + abc \\ = acs - abc - bcs + abc \end{aligned}$$

$$\Rightarrow ab - ac = ac - bc \Rightarrow ab + bc = 2ac$$

$$\Rightarrow \frac{1}{c} + \frac{1}{a} = \frac{2}{b}$$

$\Rightarrow a, b, c$ are in HP

346 (d)

It is given that O is the origin and $P(2,3)$ and

$Q(4,5)$ are two points

$$\therefore \cos \angle POQ = \frac{OP^2 + OQ^2 - PQ^2}{2(OP)(OQ)}$$

$$\Rightarrow OP \times OQ \cos \angle POQ = \frac{1}{2} \{OP^2 + OQ^2 - PQ^2\}$$

$$\Rightarrow OP \times OQ \cos \angle POQ = \frac{1}{2} \{13 + 41 - 8\} = 23$$

ALITER If O is the origin and $P(x_1, y_1), Q(x_2, y_2)$ are two points, then

$$OP \times OQ \times \cos \angle POQ = x_1 x_2 + y_1 y_2$$

Here, $x_1 = 2, y_1 = 3, x_2 = 4$ and $y_2 = 5$

$$\therefore OP \times OQ \times \cos \angle POQ = 8 + 15 = 23$$

347 (d)

$$\frac{\sin 3B}{\sin B} = \frac{3 \sin B - 4 \sin^3 B}{\sin B} = 3 - 4 \sin^2 B$$

$$= 3 - 4(1 - \cos^2 B)$$

$$= -1 + \frac{4(a^2 + c^2 - b^2)^2}{4(ac)^2}$$


$$= -1 + \frac{\left(\frac{a^2+c^2}{2}\right)^2}{(ac)^2}$$

$$= -1 + \frac{(a^2 + c^2)^2}{4(ac)^2} \quad [\because 2b^2 = a^2 + c^2 \text{ (given)}]$$

$$= \frac{(a^2 + c^2)^2 - 4a^2c^2}{4(ac)^2} = \left(\frac{c^2 - a^2}{2ac}\right)^2$$

348 (a)

Since, $\angle QPC = \alpha$



$$\therefore \angle QPB = \angle BPC = \frac{\alpha}{2}$$

$$\text{In } \Delta PQB, \sin \frac{\alpha}{2} = \frac{r}{l}$$

$$\Rightarrow l = r \sec \frac{\alpha}{2} \dots (i)$$


$$\text{and in } \Delta POB, \sin \beta = \frac{h}{l}$$

$$\Rightarrow h = l \sin \beta$$

$$\Rightarrow h = r \operatorname{cosec} \frac{\alpha}{2} \sin \beta \quad [\text{from Eq.(i)}]$$

349 (d)

$$\begin{aligned}\cos OPQ &= \frac{OP^2 + PQ^2 - OQ^2}{2OP \cdot PQ} \\ &= \frac{(4^2 + 2^2) + 2^2 - (4^2 + 4^2)}{2\sqrt{4^2 + 2^2} \cdot \sqrt{2^2}} \\ &= \frac{24 - 32}{42\sqrt{5}}\end{aligned}$$




$$\Rightarrow \angle OPQ = \cos^{-1}\left(-\frac{1}{\sqrt{5}}\right)$$

350 (a)

Let AB be a hill whose height is h metres and CD be a pillar of height h' metres.

$$\text{In } \triangle EDB, \tan \alpha = \frac{h-h'}{ED} \dots (i)$$

$$\text{and in } \triangle ACB, \tan \beta = \frac{h}{AC} = \frac{h}{ED} \dots (ii)$$



\therefore From Eqs. (i) and (ii),

$$\frac{\tan \alpha}{\tan \beta} = \frac{h-h'}{h}$$

$$\Rightarrow h \cdot \frac{\tan \alpha}{\tan \beta} = h - h' \Rightarrow h' = \frac{h(\tan \beta - \tan \alpha)}{\tan \beta}$$

351 (c)

We have,

$$\text{Area of } \triangle ABC = 3 \text{ Area of } \triangle GAB$$

Now,

$$\text{Area of } \triangle GAB = \frac{1}{2}$$

$$\times \text{Absolute value of } \begin{vmatrix} 1 & 4 & 1 \\ 4 & -3 & 1 \\ -9 & 7 & 1 \end{vmatrix}$$

$$\Rightarrow \text{Area of } \triangle GAB = \frac{1}{2} |-10 - 52 + 1| = \frac{61}{2} \text{ sq. units}$$

$$\text{Hence, Area of } \triangle ABC = \frac{183}{2} \text{ sq. units}$$

352 (a)

Using sine rule,

$$\frac{a}{\sin A} = \frac{b}{\sin B}$$

$$\Rightarrow \frac{2\sqrt{2}}{\sin 45^\circ} = \frac{6}{\sin B}$$

$$\begin{aligned}\Rightarrow \sin B &= \frac{6}{2\sqrt{2}} \times \frac{1}{\sqrt{2}} \\ &= \frac{6}{4} = \frac{3}{2} > 1\end{aligned}$$

Which is not possible

Hence, no triangle is possible


353 (b)

Let the height of pole $AD = BC = h$

$$\text{In } \triangle OBC, \tan 60^\circ = \frac{h}{x}$$

$$\Rightarrow x = \frac{h}{\sqrt{3}}$$

$$\text{In } \triangle AOD, \tan 30^\circ = \frac{h}{100-x}$$



$$\Rightarrow h = (100 - x) \frac{1}{\sqrt{3}}$$

$$= \left(100 - \frac{h}{\sqrt{3}}\right) \frac{1}{\sqrt{3}}$$

$$\Rightarrow 3h = 100\sqrt{3} - h \text{ [from Eq.(i)]}$$

$$\Rightarrow h = \frac{100\sqrt{3}}{4}$$

$$\Rightarrow h = 25\sqrt{3} \text{ m}$$

354 (b)

The vertices of a triangle are $(0, 2), \left(\frac{1}{2}, 1\right), (1, 1)$


$$\therefore \text{Area of triangle} = \frac{1}{2} \begin{vmatrix} 0 & 2 & 1 \\ \frac{1}{2} & 1 & 1 \\ 1 & 1 & 1 \end{vmatrix}$$

$$= \frac{1}{2} \left[-2 \left(\frac{1}{2} - 1 \right) + 1 \left(\frac{1}{2} - 1 \right) \right]$$

$$= \frac{1}{4} \text{ sq unit}$$

356 (c)

Since, triangle is isosceles, hence centroid is the desired point




$$\therefore \text{Coordinates of } R \left(3, \frac{4}{3} \right)$$

357 (c)

Let $AQ(=PQ)$ be the broken part of the tree OP .
It is given that $OA = 10$ m and $\angle OAQ = 45^\circ$

In $\triangle OAQ$, we have

$$\tan 45^\circ = \frac{OQ}{OA} \Rightarrow OQ = 10$$



$$\text{Also, } AQ^2 = OA^2 + OQ^2$$

$$\Rightarrow AQ = \sqrt{100 + 100} = 10\sqrt{2}$$

$$\therefore OP = OQ + PQ = OQ + AQ = 10 + 10\sqrt{2} \\ = 10(\sqrt{2} + 1) \text{ mts}$$

358 (c)

Let the new coordinates be $P(x', y')$ after shifting origin to $P(x', y')$ i.e., $x = x' + h$ and $y = y' + k$

$$\therefore (x' + h)^2 + (y' + k)^2 - 4(x' + h) + 6(y' + k) - 7 = 0$$

$$\Rightarrow (x')^2 + (y')^2 + 2(h - 2)x' + 2(k + 3)y' + (h^2 + k^2 - 4h + 6k - 7) = 0$$

According to the question,

$$h - 2 = 0 \text{ and } k + 3 = 0$$


$$\Rightarrow (h, k) = (2, -3)$$

359 (d)

$$\text{In } \triangle ABC, \tan 30^\circ = \frac{BC}{AC}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{h - 150}{60}$$

$$\Rightarrow h - 150 = \frac{60}{\sqrt{3}}$$



$$\Rightarrow h = (150 + 20\sqrt{3}) \text{ m}$$

360 (d)

The sum of the distance

$$= \frac{\frac{a}{a} + \frac{b}{a} - 1}{\sqrt{\left(\frac{1}{a}\right)^2 + \left(\frac{1}{b}\right)^2}} + \frac{\frac{a}{b} + \frac{b}{a} - 1}{\sqrt{\left(\frac{1}{b}\right)^2 + \left(\frac{1}{a}\right)^2}}$$

$$= \left(\frac{a}{b} + \frac{b}{a}\right) \frac{1}{\sqrt{\frac{1}{a^2} + \frac{1}{b^2}}} \\ = \sqrt{a^2 + b^2}$$

361 (b)

$$\text{Given, } \frac{1}{b+c} + \frac{1}{c+a} = \frac{3}{a+b+c}$$

$$\Rightarrow 1 + \frac{b}{a+c} + 1 + \frac{a}{b+c} = 3$$

$$\Rightarrow b(b+c) + a(a+c) = (a+c)(b+c)$$

$$\Rightarrow a^2 + b^2 - c^2 = ab$$

$$\therefore \cos C = \frac{a^2 + b^2 - c^2}{2ab} = \frac{ab}{2ab} \Rightarrow \angle C = 60^\circ$$

362 (b)

$$\text{Given, } \frac{\sin A}{1/4} = \frac{\sin B}{1/4} = \frac{\sin C}{1/3}$$

$$\therefore \frac{\sin A}{3} = \frac{\sin B}{3} = \frac{\sin C}{4}$$

Here, $a = 3k$, $b = 3k$ and $c = 4k$, where k is a proportionality constant

$$\therefore \cos C = \frac{9k^2 + 9k^2 - 16k^2}{2 \times 3k \times 3k} = \frac{1}{9}$$

363 (b)

$$(b+c)(bc) \cos A + (a+c)(ac) \cos B \\ + (a+b)(ab) \cos C$$


$$= (b+c)(bc) \left(\frac{b^2 + c^2 - a^2}{2bc}\right) + (a+c)(ac) \\ \times \left(\frac{a^2 + c^2 - b^2}{2ca}\right) \\ + (a+b)(ab) \left(\frac{a^2 + b^2 - c^2}{2ab}\right)$$

$$= \frac{1}{2} \{ (b+c)(b^2 + c^2 - a^2) + (a+c) \\ \times (a^2 + c^2 - b^2) + (a+b)(a^2 \\ + b^2 - c^2) \}$$

$$= \frac{1}{2} \{ 2a^3 + 2b^3 + 2c^3 \} = a^3 + b^3 + c^3$$

364 (a)

$$\text{In } \triangle DCB, \tan \theta = \frac{h}{x} \dots (i)$$



$$\text{and in } \triangle ECA, \tan \theta = \frac{h+6}{2\sqrt{3}+x}$$

$$\Rightarrow \frac{h}{x} = \frac{h+6}{2\sqrt{3}+x} \text{ [from Eq.(i)]}$$

$$\Rightarrow 2\sqrt{3}h + hx = hx + 6x$$

$$\Rightarrow 2\sqrt{3}h = 6x$$

$$\Rightarrow h = \frac{6x}{2\sqrt{3}}$$

From Eq. (i), we get

$$\tan \theta = \frac{6x}{2\sqrt{3}x} = \sqrt{3}$$

$$\Rightarrow \theta = 60^\circ$$

365 (b)

Since, angles A, B, C are in AP

$$\therefore 2B = A + C$$

$$\because A + B + C = 180^\circ$$

$$\Rightarrow B = 60^\circ$$

$$\text{Now, } \cos B = \frac{a^2 + c^2 - b^2}{2ac}$$

$$\Rightarrow \frac{1}{2} = \frac{a^2 + c^2 - b^2}{2ac}$$

$$\Rightarrow a^2 + c^2 - b^2 = ac$$

$$\Rightarrow b^2 = a^2 + c^2 - ac$$

366 (a)

Let $a = 7$ cm, $b = 4\sqrt{3}$ cm and $c = \sqrt{13}$ cm

Here, we see that the smallest side is c

Therefore, the smallest angle will be C

$$\therefore \cos C = \frac{(7)^2 + (4\sqrt{3})^2 - (\sqrt{13})^2}{2 \times 7 \times 4\sqrt{3}} = \frac{\sqrt{3}}{2}$$

$$\Rightarrow \angle C = \frac{\pi}{6}$$

367 (c)

On solving the given straight lines, we get vertices of $\triangle ABC$

Which are $A(2, -2), B(1, 1), C(-2, 2)$

$$\therefore AB = \sqrt{(1-2)^2 + (1+2)^2} = \sqrt{10}$$

$$BC = \sqrt{(-2-1)^2 + (2-1)^2} = \sqrt{10}$$

$$\text{And } CA = \sqrt{(2+2)^2 + (-2-2)^2} = \sqrt{32}$$

Here, $AB = BC$

\Rightarrow Triangle is isosceles triangle

369 (d)

$$\text{If } \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix} = 0, \text{ then the points are collinear}$$

370 (a)

Let $\angle A = 45^\circ$ and $\angle B = 60^\circ$

$$\therefore \angle C = 75^\circ$$

Let smallest and greatest sides are a and c

$$\therefore a : c = \sin 45^\circ : \sin 75^\circ$$


$$= \frac{1}{\sqrt{2}} : \frac{\sqrt{3}+1}{2\sqrt{2}} = 2 : \sqrt{3} + 1$$

$$= 2(\sqrt{3}-1) : (\sqrt{3}+1)(\sqrt{3}-1)$$

$$= (\sqrt{3}-1) : 1$$

371 (b)

$$\text{In } \triangle ADF, \tan 60^\circ = \frac{1}{AF}$$



$$\Rightarrow AF = \cot 60^\circ = \frac{1}{\sqrt{3}} \quad \dots(i)$$

$$\text{In } \triangle ABC, \tan 30^\circ = \frac{1}{AB}$$

$$\Rightarrow AB = \cot 30^\circ$$

$$\Rightarrow AF + FB = \sqrt{3}$$


$$\Rightarrow d = \sqrt{3} - \frac{1}{\sqrt{3}} = \frac{2}{\sqrt{3}} \quad [\text{using Eq.(i)}]$$

$$\therefore \text{Speed of aeroplane} = \frac{\text{distance from } D \text{ to } C}{\text{time taken}}$$

$$= \frac{\frac{2}{\sqrt{3}}}{\frac{1}{10}} \times 60 \times 60 = 240\sqrt{3} \text{ km/h}$$

372 (a)

Let CD is a tower of height h . AB is building of height a



$$\text{In } \triangle BLD, \tan 30^\circ = \frac{h-a}{LB}$$

$$\therefore LB = \frac{(h-a)}{\tan 30^\circ} = \sqrt{3}(h-a)$$

$$\text{In } \triangle ACD, \tan 45^\circ = \frac{h}{LB}$$

$$\text{or } h(\sqrt{3}-1) = \sqrt{3}a$$

$$\therefore h = \frac{\sqrt{3}a}{\sqrt{3}-1} = \frac{\sqrt{3}(\sqrt{3}+1)a}{2}$$

$$\therefore h = \left(\frac{3+\sqrt{3}}{2}\right)a$$

373 (d)

$$\frac{\sin 3B}{\sin B} = \frac{3 \sin B - 4 \sin^3 B}{\sin B}$$


$$= 3 - 4(1 - \cos^2 B)$$

$$= -1 + 4 \left(\frac{a^2 + c^2 - b^2}{2ac} \right)^2 = -1 + \frac{\left(\frac{a^2 + c^2}{2} \right)^2}{(ac)^2}$$

$$= \frac{(a^2 + c^2)^2 - 4a^2c^2}{4(ac)^2} = \left(\frac{c^2 - a^2}{2ac} \right)^2$$

375 (b)

$$\text{Since, } \frac{BD}{DC} = \frac{1}{3}$$



In $\triangle ABD$, by sine rule

$$AD = \frac{\sin \frac{\pi}{3}}{\sin \angle BAD} \cdot BD \quad \dots \text{(ii)}$$

In $\triangle ADC$, by sine rule

$$AD = \frac{\sin \frac{\pi}{4}}{\sin \angle DAC} \cdot DC \quad \dots \text{(iii)}$$

From Eqs. (ii) and (iii),

$$\frac{\sin \angle BAD}{\sin \angle DAC} = \frac{\sin \frac{\pi}{3}}{\sin \frac{\pi}{4}} \cdot \frac{BD}{DC} = \frac{\sqrt{3}}{2} \cdot \sqrt{2} \cdot \frac{1}{3} = \frac{1}{\sqrt{6}}$$

376 (b)

Since, reflection of the orthocenter of $\triangle ABC$ in base BC will always lie on the circumcircle of the triangle ABC , therefore coordinate of a point lying on the circumcircle is $(1 - \frac{1 \times 4}{2}, 1 - \frac{1 \times 4}{2})$ i.e., $(-1, -1)$ and coordinates of the circumcentre is $(2, 0)$

$$\therefore \text{Radius of the circumcentre of } \triangle ABC = \sqrt{(2+1)^2 + (1)^2} = \sqrt{10}$$

377 (a)

Given, $b + c = 2a, \angle A = 60^\circ$

Since, $\angle A = 60^\circ \angle B + \angle C = 120^\circ$

Also, $b + c = 2a$ [given]

$$\sin B + \sin C = 2 \sin 60^\circ$$

$$\Rightarrow \left[\because \frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c} = k \right]$$

$$\Rightarrow 2 \sin \left(\frac{B+C}{2} \right) \cos \left(\frac{B-C}{2} \right) = \frac{2\sqrt{3}}{2}$$

$$\Rightarrow 2 \sin 60^\circ \cos \left(\frac{B-C}{2} \right) = \sqrt{3} \left[\because \sin 60^\circ = \frac{\sqrt{3}}{2} \right]$$

$$\Rightarrow \cos \left(\frac{B-C}{2} \right) = 1 \Rightarrow \frac{B-C}{2} = 0$$

$$\Rightarrow \angle B = \angle C$$

Hence, triangle is an equilateral triangle

378 (c)

Let the vertices of the triangle are

$A(-2, -6), B(-2, 4)$ and $C(1, 3)$

$$\text{Now, } AB = \sqrt{(-2+2)^2 + (-6-4)^2} = 10$$

$$BC = \sqrt{(-2-1)^2 + (4-3)^2} = \sqrt{10}$$

$$\text{and } CA = \sqrt{(1+2)^2 + (3+6)^2} = \sqrt{90}$$

$$\text{Here, } AB^2 = BC^2 + CA^2$$

\therefore Triangle is right angled triangle at C

$$\therefore \angle C = 90^\circ$$

So, orthocenter is $(1, 3)$

379 (a)

$$\text{Area of } \Delta = \frac{1}{2} \begin{vmatrix} 0 & 0 & 1 \\ a \cos \theta & b \sin \theta & 1 \\ a \cos \theta & -b \sin \theta & 1 \end{vmatrix}$$

$$\Rightarrow \Delta = \frac{1}{2} [1(-ab \sin \theta \cos \theta - ab \sin \theta \cos \theta)] = \frac{ab \sin 2\theta}{2}$$

Since, maximum value of $\sin 2\theta$ is 1, when $\theta = \frac{\pi}{4}$

$$\therefore \Delta_{\max} = \frac{ab}{2}$$

380 (d)

Let $O(x, y)$ be the circumcentre. Then, G divides $O'O$ in the ratio $2 : 1$

$$\therefore \frac{2x+0}{2+1} = 2 \text{ and } \frac{2y+1}{2+1} = 3 \Rightarrow x = 3 \text{ and } y = 4$$

Hence, the coordinates of O are $(3, 4)$

381 (d)

Perpendicular bisector of $A(1, 3)$ and $B(-3, 5)$ is

$$2x - y + 6 = 0 \quad \dots \text{(i)}$$

And perpendicular bisector of $A(1, 3)$ and $C(5, -1)$ is

$$x - y - 2 = 0 \quad \dots \text{(ii)}$$

On solving Eqs. (i) and (ii), we get

$$x = -8, y = -10$$

\therefore Coordinates of P are $(-8, -10)$

$$\text{Thus, } PA = \sqrt{(1+8)^2 + (3+10)^2} = \sqrt{81 + 169} = 5\sqrt{10}$$

382 (a)

Given sides are $3x - 4y = 0, 5x + 12y = 0$ and $x - 15 = 0$. The vertices of a triangle are

$A(0, 0), B(20, 15), C(-36, 15)$

$$\text{Now, } AB = c = \sqrt{(20-0)^2 + (15-0)^2} = \sqrt{400 + 225}$$

$$= 25$$

$$BC = a = \sqrt{(20+36)^2 + (15-15)^2} = 56$$

$$CA = b = \sqrt{(-36-0)^2 + (15-0)^2} = \sqrt{1296 + 225} = 39$$

\therefore Incentre

$$= \left(\frac{ax_1 + bx_2 + cx_3}{a+b+c}, \frac{ay_1 + by_2 + cy_3}{a+b+c} \right)$$

$$= \left(\frac{56 \times 0 + 39 \times 20 - 36 \times 25}{56 + 39 + 25}, \frac{56 \times 0 + 39 \times 15 + 25 \times 15}{56 + 39 + 25} \right)$$

$$= \left(\frac{-120}{120}, \frac{960}{120} \right)$$

$$= (-1, 8)$$


383 (d)

The mid point of the line joining the points $(-10, 8)$ and $(-6, -12)$ is $\left(\frac{-10-6}{2}, \frac{8+12}{2}\right)$ i.e., $(-8, 10)$. Let $(-8, 10)$ divides the line joining the points $(4, -2)$ and $(-2, 4)$ in the ratio $m:n$

$$\begin{aligned} \text{Then, } \frac{m(-2)+n(4)}{m+n} &= -8 \\ \Rightarrow -2m + 4n &= -8m - 8n \\ \Rightarrow 6m &= -12n \Rightarrow \frac{m}{n} = \frac{-2}{1} \\ \therefore \text{Required ratio } &2:1 \text{ externally.} \end{aligned}$$

384 (a)

Given the circle is inscribed in square formed by the lines $x^2 - 8x + 12 = 0$ and $y^2 - 14y + 45 = 0$
 $\Rightarrow x = 6$ and $x = 2$, $y = 5$ and $y = 9$



$ABCD$ clearly forms a square
 \therefore Centre of inscribed circle
 = point of intersection of diagonals
 = mid point of AC or BD
 $= \left(\frac{2+6}{2}, \frac{5+9}{2}\right)$
 \Rightarrow Centre of inscribed circle = $(4, 7)$

385 (a)

$$\begin{aligned} \text{Area of a circle} &= \pi \times (\text{radius})^2 \\ \therefore A &= \pi r^3, A_1 = \pi r_1^2, A_2 = \pi r_2^2, A_3 = \pi r_3^2 \\ \therefore \frac{1}{\sqrt{A_1}} + \frac{1}{\sqrt{A_2}} + \frac{1}{\sqrt{A_3}} \\ &= \frac{1}{r_1\sqrt{\pi}} + \frac{1}{r_2\sqrt{\pi}} + \frac{1}{r_3\sqrt{\pi}} \\ &= \frac{1}{\sqrt{\pi}} \left(\frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{r_3}\right) \\ &= \frac{1}{\sqrt{\pi}} \left(\frac{s-a}{\Delta} + \frac{s-b}{\Delta} + \frac{s-c}{\Delta}\right) \\ &= \frac{1}{\sqrt{\pi}} \left(\frac{3s - (a+b+c)}{\Delta}\right) \\ &= \frac{1}{\sqrt{\pi}} \cdot \frac{3s - 2s}{\Delta} = \frac{1}{\sqrt{\pi}} \cdot \frac{s}{\Delta} \\ &= \frac{1}{r\sqrt{\pi}} = \frac{1}{\sqrt{A}} \end{aligned}$$

386 (b)

We have,

$$\begin{aligned} x^2 + 4xy + y^2 &= aX^2 + bY^2 \\ \Rightarrow (X \cos \theta + Y \sin \theta)^2 \\ &+ 14(X \cos \theta + Y \sin \theta)(X \sin \theta \\ &- Y \cos \theta) + (X \sin \theta - Y \cos \theta)^2 \\ &= aX^2 + bY^2 \end{aligned}$$

$$\begin{aligned} \Rightarrow a &= 1 + 4 \sin \theta \cos \theta, b = 1 - 4 \sin \theta \cos \theta \text{ and} \\ \sin^2 \theta - \cos^2 \theta &= 0 \\ \Rightarrow a &= 1 + 4 \sin \theta \cos \theta, b = 1 - 4 \sin \theta \cos \theta \text{ and} \\ \theta &= \frac{\pi}{4} \\ \Rightarrow a &= 3, b = -1 \end{aligned}$$


387 (d)

$$\begin{aligned} \text{Let } S(x, y), \text{ then} \\ (x+1)^2 + y^2 + (x-2)^2 + y^2 &= 2[(x-1)^2 + y^2] \\ \Rightarrow 2x + 1 + 4 - 4x &= -4x + 2 \\ \Rightarrow x &= \frac{-3}{2} \end{aligned}$$

Hence, it is a straight line parallel to y -axis

388 (b)

Given lines are



$$\begin{aligned} x^2 - 3y^2 &= 0 \dots (i) \\ \text{And } x &= 4 \dots (ii) \\ \text{Put } x &= 4 \text{ from Eq. (ii) in Eq. (i), we get} \\ 16 - 3y^2 &= 0 \\ \Rightarrow y^2 &= \frac{16}{3} \Rightarrow y = \pm \frac{4}{\sqrt{3}} \end{aligned}$$

The vertices of the triangle are

$$O(0, 0), P\left(4, \frac{4}{\sqrt{3}}\right), Q\left(4, -\frac{4}{\sqrt{3}}\right)$$

$$OP = \sqrt{4^2 + \left(\frac{4}{\sqrt{3}}\right)^2} = \frac{8}{\sqrt{3}}$$

$$OQ = \sqrt{4^2 + \left(\frac{4}{\sqrt{3}}\right)^2} = \frac{8}{\sqrt{3}}$$

$$PQ = \sqrt{\left(\frac{4}{\sqrt{3}} + \frac{4}{\sqrt{3}}\right)^2} = \frac{8}{\sqrt{3}}$$


$$\Rightarrow OP = OQ = PQ = \frac{8}{\sqrt{3}}$$

$\therefore \Delta OPQ$ is an equilateral triangle

389 (c)


Let BC be the incomplete and BD be the complete pillar. In ΔABC and ABD , we have

$$\begin{aligned} \tan 45^\circ &= \frac{BC}{AB} \text{ and } \tan 60^\circ = \frac{BD}{AB} \\ \Rightarrow BC &= 100 \text{ m and } BD = 100\sqrt{3} \\ \Rightarrow BC + CD &= 100\sqrt{3} \\ \Rightarrow 100 + x &= 100\sqrt{3} \Rightarrow x = 100(\sqrt{3} - 1) \text{ m} \end{aligned}$$



390 (b)

The coordinates of the vertices of ΔOAB are $O(0,0)$, $A(4,0)$ and $B(0,3)$
 $\therefore OA = 4$, $OB = 3$ and $AB = 5$



Hence, the coordinates of the excentre opposite to the vertex O are


$$\left(\frac{-5 \times 0 + 3 \times 4 + 4 \times 0}{-5 + 3 + 4}, \frac{-5 \times 0 + 3 \times 0 + 4 \times 3}{-5 + 3 + 4} \right) = (6, 6)$$

391 (b)

$$\text{In } \Delta AOT, \tan \alpha = \frac{h}{OA}$$

$$\Rightarrow OA = h \cot \alpha$$

$$\text{In } \Delta AOP, \tan \beta = \frac{l}{AO}$$



$$\Rightarrow \tan \beta = \frac{l}{h \cot \alpha} \text{ [from Eq.(i)]}$$

$$\Rightarrow h = l \tan \alpha \cot \beta$$

392 (b)

$$6. \quad b \cos^2 \frac{C}{2} + c \cos^2 \frac{B}{2} = b \cdot \frac{s(s-c)}{ab} + c \cdot \frac{s(s-b)}{ac}$$

$$= \frac{s}{a} [2s - (b+c)] = s$$

Hence, statement I is true

$$7. \quad \text{Let } \cot \frac{A}{2} = \frac{b+c}{2}$$

$$\Rightarrow \frac{\cos \frac{A}{2}}{\sin \frac{A}{2}} = \frac{\sin B + \sin C}{\sin A}$$

$$\Rightarrow \frac{\cos \frac{A}{2}}{\sin \frac{A}{2}} = \frac{2 \sin \left(\frac{B+C}{2} \right) \cos \left(\frac{B-C}{2} \right)}{2 \sin \frac{A}{2} \cos \frac{A}{2}}$$

$$\Rightarrow \cos \frac{A}{2} = \cos \left(\frac{B-C}{2} \right)$$

$$\Rightarrow \frac{A}{2} = \frac{B-C}{2} \Rightarrow A + C = B$$

$$\text{But } A + B + C = \pi, \text{ therefore } B = \frac{\pi}{2}$$


But given statement is

$$\cot \frac{A}{2} = \frac{b+c}{2} \Rightarrow \angle B = 90^\circ$$

Hence, statement II is not true

393 (b)

$\because \angle CDA = \beta$, $\angle DCB = \gamma$,
and $\angle ACB = \pi - (\alpha + \beta + \gamma)$



In ΔABC , on applying sine rule, we get

$$\frac{AB}{\sin[\pi - (\alpha + \beta + \gamma)]} = \frac{CA}{\sin \gamma}$$

$$\Rightarrow CA = \frac{a \sin \gamma}{\sin(\alpha + \beta + \gamma)}$$

Now, in ΔCAD , on applying sine rule

$$\frac{CA}{\sin \beta} = \frac{CD}{\sin \alpha}$$

$$\Rightarrow CD = \frac{CA \cdot \sin \alpha}{\sin \beta}$$

$$= \frac{a \sin \gamma \sin \alpha}{\sin \beta \sin(\alpha + \beta + \gamma)}$$

394 (b)


Let PD be a pole

$$\text{In } \Delta DAP, \tan 30^\circ = \frac{DP}{AD}$$

$$\Rightarrow DP = \frac{a}{\sqrt{3}}$$

$$\text{In } \triangle PDB, \tan \theta = \frac{DP}{BD}$$

$$\Rightarrow \tan \theta = \frac{a/\sqrt{3}}{\sqrt{2}a} = \frac{1}{\sqrt{6}}$$



395 (b)

$$\text{Since, } \frac{a}{b} = \frac{2}{\sqrt{3}-1} \text{ and } A = 3B$$

$$\text{By sine rule, } \frac{\sin A}{\sin B} = \frac{a}{b}$$

$$\Rightarrow \frac{\sin 3B}{\sin B} = \frac{2}{\sqrt{3}-1}$$

$$\Rightarrow \frac{3 \sin B - 4 \sin^3 B}{\sin B} = \frac{2}{\sqrt{3}-1}$$

$$\Rightarrow 3 - 4 \sin^2 B = \frac{2(\sqrt{3}+1)}{2}$$

$$\Rightarrow \frac{2-\sqrt{3}}{4} = \sin^2 B$$


$$\Rightarrow \sin B = \frac{\sqrt{3}-1}{2\sqrt{2}}$$

$$\Rightarrow \angle B = 15^\circ, \angle A = 45^\circ \text{ and } \angle C = 120^\circ$$

396 (d)

$$\text{In } \triangle ABC, \tan \alpha = \frac{h}{x+d}$$

$$\Rightarrow x+d = h \cot \alpha$$



and in $\triangle ABD$,

$$\tan \beta = \frac{h}{x} \Rightarrow x = h \cot \beta$$

On putting this value in Eq. (i), we get

$$h \cot \beta + d = h \cot \alpha$$


$$\Rightarrow h = \frac{d}{\cot \alpha - \cot \beta}$$

397 (b)

$$\tan(\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta}$$

$$= \frac{\frac{y}{z} + \frac{x}{z}}{1 - \frac{y}{z} \times \frac{x}{z}}$$

$$= \frac{\frac{3}{6} + \frac{2}{6}}{1 - \frac{3}{6} \times \frac{2}{6}} = 1$$



$$\Rightarrow \alpha + \beta = \angle QPR = \frac{\pi}{4}$$

398 (a)

Let (x, y) be the original coordinates of P . Then,

$$x = 1 + \cos \theta \text{ and } y = 1 + \cos \phi$$

$$\Rightarrow x = 2 \cos^2 \frac{\theta}{2}, y = 2 \cos^2 \frac{\phi}{2}$$

399 (b)

$$\text{Given, } r_1 = 2r_2 = 3r_3$$

$$\Rightarrow \frac{\Delta}{(s-a)} = \frac{2\Delta}{(s-b)} = \frac{3\Delta}{(s-c)}$$

$$\Rightarrow (s-b) = 2(s-a) \text{ and } (s-c) = 3(s-a)$$

$$\Rightarrow \frac{a+b+c}{2} - b = 2\left(\frac{a+b+c}{2} - a\right)$$

$$\text{and } \frac{a+b+c}{2} - c = 3\left(\frac{a+b+c}{2} - a\right)$$

$$\Rightarrow a+c-b = 2(-a+b+c)$$

$$\text{and } a+b-c = 3(-a+b+c)$$

$$\Rightarrow 3a = 3b+c \text{ and } 2a = b+2c$$

$$\Rightarrow 4a = 5b \Rightarrow \frac{a}{b} = \frac{5}{4}$$

400 (b)

Since a, b and x are in AP

$$\Rightarrow 2b = a+c$$

$$\Rightarrow 3b = 2s \Rightarrow s = \frac{3b}{2}$$

$$\text{Now, } \frac{\sin \frac{A}{2} \sin \frac{C}{2}}{\sin \frac{B}{2}} = \sqrt{\frac{ac(s-b)(s-c)(s-a)}{(s-a)(s-c) \times bc \times ab}}$$

$$= \frac{s-b}{b} = \frac{\frac{3b}{2} - b}{b} = \frac{1}{2}$$


401 (b)

Let AD be the building of height h and BP be the hill. Then,

$$\tan q = \frac{h+x}{y} \dots(i)$$

$$\text{and } \tan p = \frac{x}{y}$$

$$\Rightarrow y = x \cot p \dots(ii)$$



From Eq. (i) and (ii), we get

$$\tan q = \frac{h+x}{x \cot p}$$

$$\Rightarrow x \cot p = (h+x) \cot q$$

$$\Rightarrow x = \frac{h \cot q}{\cot p - \cot q}$$

$$\Rightarrow h+x = \frac{h \cot q}{\cot p - \cot q} + h$$

$$\therefore \text{Height of hill} = \frac{h \cot p}{\cot p - \cot q}$$

402 (b)

$$\cos^2 A = 1 - \sin^2 A = 1 - \left(\frac{3}{5}\right)^2 = \frac{16}{25}$$

$$\Rightarrow \cos A = \frac{4}{5}$$

$$\Rightarrow \frac{(20)^2 + (21)^2 - a^2}{2 \cdot 20 \cdot 21} = \frac{4}{5}$$

$$\Rightarrow a^2 = 169 \Rightarrow a = 13$$

403 (c)

Given, $\angle C = 60^\circ$, $a = 2$, $b = 4$

$$\therefore \cos C = \frac{a^2 + b^2 - c^2}{2ab}$$

$$\Rightarrow 2ab \cos 60^\circ = a^2 + b^2 - c^2$$

$$\Rightarrow ab = a^2 + b^2 - c^2$$

$$\Rightarrow 8 = 4 + 16 - c^2 \Rightarrow c^2 = 12$$

$$\Rightarrow c = \sqrt{12} = 2\sqrt{3}$$

$$\text{We have, } \sin A = \frac{a \sin C}{c} = \frac{2 \cdot \frac{\sqrt{3}}{2}}{2\sqrt{3}} = \frac{1}{2}$$

$$\Rightarrow A = 30^\circ$$

$$\text{and } \sin B = \frac{b \sin C}{c} = \frac{4 \cdot \frac{\sqrt{3}}{2}}{2\sqrt{3}} = 1 \Rightarrow B = 90^\circ$$

405 (b)

$$\frac{b - c \cos A}{c - b \cos A} = \frac{b - \frac{b^2 + c^2 - a^2}{2b}}{c - \frac{b^2 + c^2 - a^2}{2c}}$$

$$= \frac{\left(\frac{b^2 + a^2 - c^2}{c^2 + a^2 - b^2}\right)}{\frac{c}{b}}$$

$$= \frac{b^2 + a^2 - c^2}{2ab} \cdot \frac{2ac}{c^2 + a^2 - b^2} = \frac{\cos C}{\cos B}$$

406 (b)

$$\text{We have, } \frac{s(s-a)}{bc} - \frac{(s-b)(s-c)}{bc}$$

$$= \cos^2 \frac{A}{2} - \sin^2 \frac{A}{2}$$

$$= \cos \frac{2A}{2} = \cos A$$

407 (b)

Let the third vertex of the triangle be (x, y) , then

$$\frac{x+4+(-2)}{3} = 2 \Rightarrow x = 4$$

$$\text{and } \frac{y+8+6}{3} = 7 \Rightarrow y = 7$$

\therefore The coordinates of the third vertex are $(4, 7)$

408 (b)


Let $ABCD$ be a square of each side of length a . It is given that $\angle BPC = 60^\circ$. Let M be the midpoint of BC . Then $\angle BPM = \angle CPM = 30^\circ$

In $\triangle BPM$, we have

$$\tan \angle BPM = \frac{BM}{PM}$$

$$\Rightarrow PM = \sqrt{3} BM = \frac{\sqrt{3}}{2} a$$

In $\triangle OPM$, we have



$$PM^2 = OM^2 + OP^2 \Rightarrow \frac{3a^2}{4} = \frac{a^2}{4} + h^2 \Rightarrow a^2 = 2h^2$$

409 (c)

$$(a-b)^2 \cos^2 \frac{C}{2} + (a+b)^2 \sin^2 \frac{C}{2}$$

$$= (a^2 + b^2 - 2ab) \cos^2 \frac{C}{2}$$

$$+ (a^2 + b^2 + 2ab) \sin^2 \frac{C}{2}$$

$$= a^2 + b^2 + 2ab \left(\sin^2 \frac{C}{2} - \cos^2 \frac{C}{2} \right)$$

$$= a^2 + b^2 - 2ab \cos C$$

$$= a^2 + b^2 - (a^2 + b^2 - c^2)$$

$$= c^2 \left(\because \cos C = \frac{a^2 + b^2 - c^2}{2ab} \right)$$

410 (c)

$$\text{Given, } \Delta = a^2 - (b-c)^2$$

$$\begin{aligned}
 &= (a + b - c)(a - b + c) \\
 &= 2(s - c) \cdot 2(s - b) \\
 \sqrt{s(s - a)(s - b)(s - c)} &= 4(s - b)(s - c) \\
 \Rightarrow \frac{1}{4} &= \sqrt{\frac{(s - b)(s - c)}{s(s - a)}} = \tan \frac{A}{2} \\
 \therefore \tan \frac{A}{2} &= \frac{1}{4}
 \end{aligned}$$

411 (a)


If O' is the orthocentre of a ΔABC , then the points O', A, B, C are such that each point is the orthocentre of the triangle formed by the remaining three points. So, the coordinates of the orthocentre of $\Delta O'AB$ are $(0, 0)$

412 (c)

$$\begin{aligned}
 &\cot B + \cot C - \cot A \\
 &= \frac{\cos B}{\sin B} + \frac{\cos C}{\sin C} - \cot A \\
 &= \frac{\sin C \cos B + \cos C \sin B}{\sin B \sin C} - \cot A \\
 &= \frac{\sin(B + C) \cos A}{\sin B \sin C} - \cot A \\
 &= \frac{\sin^2 A - \sin B \sin C \cos A}{\sin A \sin B \sin C} \quad (\because A + B + C = \pi) \\
 &= \frac{a^2 - bc \cos A}{k(abc)} \\
 &= \frac{a^2 - bc \frac{(b^2 + c^2 - a^2)}{2bc}}{(abc)k} \\
 &= \frac{2a^2 - (3a^2 - a^2)}{2(abc)k} \quad (\because b^2 + c^2 = 3a^2 \text{ given}) \\
 &= \frac{(a^2 - a^2)}{abck} = 0
 \end{aligned}$$

413 (c)

Let h be the height of the tower.



$$\begin{aligned}
 &= \frac{h}{1} \\
 \Rightarrow h &= \frac{\sqrt{3}(\sqrt{3} - 1)}{3 - 1} \\
 &= \frac{3 - \sqrt{3}}{2} \text{ m}
 \end{aligned}$$

414 (d)

$$\therefore \cos A = \frac{b^2 + c^2 - a^2}{2bc}$$


$$\begin{aligned}
 &= \frac{(8)^2 + (10)^2 - (6)^2}{2 \times 8 \times 10} = \frac{4}{5} \\
 \Rightarrow \sin A &= \frac{3}{5} \\
 \therefore \sin 2A &= 2 \sin A \cos A \\
 &= 2 \times \frac{3}{5} \times \frac{4}{5} = \frac{24}{25}
 \end{aligned}$$

415 (d)

$$\begin{aligned}
 &\frac{1 + \cos(A + B) \cos C}{1 + \cos(A - C) \cos B} \\
 &= \frac{1 + \cos[\pi - (A + B)] \cos(A - B)}{1 + \cos[\pi - (A + C)] \cos(A - C)} \\
 &= \frac{1 - \cos(A + B) \cos(A - B)}{1 - \cos(A - C) \cos(A + C)} \\
 &= \frac{1 - \cos^2 A + \sin^2 B}{1 - \cos^2 A + \sin^2 C} = \frac{\sin^2 A + \sin^2 B}{\sin^2 A + \sin^2 C} \\
 \therefore \frac{1 + \cos(A - B) \cos C}{1 + \cos(A - C) \cos B} &= \frac{a^2 + b^2}{a^2 + c^2} \quad [\text{by sine rule}]
 \end{aligned}$$

416 (a)

$$\text{In } \Delta AEC, \tan 60^\circ = \frac{500}{d_1} \Rightarrow d_1 = \frac{500}{\sqrt{3}} \text{ m} \dots (i)$$



$$\text{and in } \Delta BEC, \tan 30^\circ = \frac{500}{d_2}$$


$$\Rightarrow d_2 = 500\sqrt{3} \text{ m} \dots (ii)$$

\therefore Required diameter,

$$AB = d_1 + d_2 = \frac{500}{\sqrt{3}} + 500\sqrt{3} = \frac{2000}{\sqrt{3}} \text{ m}$$

417 (d)

Using sine rule in ΔACD ,



$$\frac{\sin(90^\circ - C)}{b} = \frac{\sin 90^\circ}{\frac{a}{c}} \Rightarrow \cos C = \frac{2b}{a}$$

And in ΔABD ,

$$\frac{\sin(A - 90^\circ)}{\frac{a}{2}} = \frac{\sin(90^\circ + C)}{c}$$


$$\Rightarrow -\frac{\cos A}{\frac{a}{2}} = \frac{\cos C}{c} \Rightarrow \cos A = -\frac{b}{c}$$

$$\Rightarrow \frac{b^2 + c^2 - a^2}{2bc} = -\frac{b}{c} \Rightarrow c^2 - a^2 = 3b^2$$

$$\therefore \cos A \cos C = -\frac{2b^2}{ac} = \frac{2}{3ac}(c^2 - a^2)$$

418 (a)

Let the vertices of triangle of A, B, C are $(x_1, y_1), (x_2, y_2), (x_3, y_3)$ respectively
Given mid points of the sides AB, BC and CA of ΔABC are $D(6, 1), E(3, 5)$ and $F(-1, -2)$ respectively



$$\therefore \frac{x_1 + x_2}{2} = 6, \frac{y_1 + y_2}{2} = 1$$

$$\Rightarrow \left. \begin{aligned} x_1 + x_2 &= 12, \\ y_1 + y_2 &= 2 \end{aligned} \right\} \dots(i)$$

$$\text{Similarly, } \left. \begin{aligned} x_2 + x_3 &= 6, \\ y_2 + y_3 &= 10 \end{aligned} \right\} \dots(ii)$$

$$\text{And } \left. \begin{aligned} x_1 + x_3 &= -2, \\ y_1 + y_3 &= -4 \end{aligned} \right\} \dots(iii)$$

On solving Eqs. (i), (ii) and (iii), we get

$$x_1 = 2, \quad x_2 = 10, \quad x_3 = -4$$

$$\text{And } y_1 = -6, \quad y_2 = 8, \quad y_3 = 2$$

Now, the vertex opposite to D is C i.e., $(-4, 2)$

419 (a)

Let $A(x_1, y_1), B(x_2, y_2)$ and $C(x_3, y_3)$ be the vertices of the triangle ABC and let $P(h, k)$ be any point on the locus,

Then, $PA^2 + PB^2 + PC^2 = c$ (constant)

$$\Rightarrow \sum_{i=1}^3 [(h - x_i)^2 + (k - y_i)^2] = c$$

$$\Rightarrow h^2 + k^2 - \frac{2h}{3}(x_1 + x_2 + x_3) - \frac{2k}{3}(y_1 + y_2 + y_3) + \sum_{i=1}^3 (x_i^2 + y_i^2) - c = 0$$

So, locus of (h, k) is

$$x^2 + y^2 - \frac{2x}{3}(x_1 + x_2 + x_3) - \frac{2y}{3}(y_1 + y_2 + y_3) + \lambda = 0$$

Where $\lambda = \sum_{i=1}^3 (x_i^2 + y_i^2) - c = 0$ (constant)

Clearly, the locus of a point is a circle with, centre at

$$\left(\frac{x_1 + x_2 + x_3}{3}, \frac{y_1 + y_2 + y_3}{3} \right)$$

420 (c)

$$\text{Since, } B = \frac{A+C}{2}$$

$$\Rightarrow B = 90^\circ - \frac{B}{2} \Rightarrow B = 60^\circ [\therefore A + B + C = 180^\circ]$$

$$\therefore \frac{\sin 60^\circ}{\sqrt{3}} = \frac{\sin C}{\sqrt{2}} \quad [\text{by sine rule}]$$

$$\Rightarrow \sin C = \frac{1}{\sqrt{2}} \Rightarrow C = 45^\circ$$

$$\therefore \angle A = 180^\circ - (60^\circ + 45^\circ) = 75^\circ$$

421 (b)

Since, $A(0, 1), B(0, -1)$ and $C(x, 0)$ are the vertices of an equilateral ΔABC .

$$\therefore AB = BC$$

$$\Rightarrow \sqrt{0^2 + 4} = \sqrt{x^2 + 1}$$

$$\Rightarrow x^2 = 3$$

$$\Rightarrow x = \pm\sqrt{3}$$

422 (c)

Let $a = \sin \alpha, b = \cos \alpha, c = \sqrt{1 + \sin \alpha \cos \alpha}$

Here, we see that the greatest side is c

$$\therefore \cos C = \frac{a^2 + b^2 - c^2}{2ab}$$


$$\Rightarrow \cos C = \frac{\sin^2 \alpha \cos^2 \alpha - 1 - \sin \alpha \cos \alpha}{2 \sin \alpha \cos \alpha}$$

$$\Rightarrow \cos C = -\frac{\sin \alpha \cos \alpha}{2 \sin \alpha \cos \alpha}$$

$$\Rightarrow \cos C = -\frac{1}{2} \Rightarrow \angle C = 120^\circ$$

423 (b)

Let the distance of two consecutive stones are $x, x + 1$



In ΔBCD ,

$$\tan 60^\circ = \frac{h}{x} \Rightarrow x = \frac{h}{\sqrt{3}} \dots(i)$$

$$\text{In } \Delta ABC, \tan 30^\circ = \frac{h}{x+1}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{h}{x+1} \Rightarrow \frac{h}{\sqrt{3}} + 1 = \sqrt{3}h \quad [\text{from Eq. (i)}]$$


$$\Rightarrow h = \frac{\sqrt{3}}{2} \text{ km}$$

424 (a)

$$\text{Given, } \tan \phi = 0.5 = \frac{1}{2}$$

$$\text{In } \Delta ABC, \tan \theta = \frac{10}{AB}$$

$$\Rightarrow AB = \frac{10}{\tan \theta}$$



In $\triangle ABD$,

$$\begin{aligned} \tan(\theta + \phi) &= \frac{30}{AB} \\ \Rightarrow \frac{\tan \theta + \tan \phi}{1 - \tan \theta \tan \phi} &= \frac{30 \tan \theta}{10} \\ \Rightarrow \tan \theta + \frac{1}{2} &= 3 \tan \theta - \frac{3}{2} \tan^2 \theta \\ \Rightarrow 3 \tan^2 \theta - 4 \tan \theta + 1 &= 0 \\ \Rightarrow (3 \tan \theta - 1)(\tan \theta - 1) &= 0 \\ \Rightarrow \tan \theta &= \frac{1}{3}, 1 \\ \therefore \tan \theta &= 1 \end{aligned}$$

426 (b)

Since, the sides of a triangle are in AP

$$\therefore 2b = a + c \text{ and } c = 7 \text{ cm}$$

$$\therefore \cos C = \frac{a^2 + b^2 - c^2}{2ab}$$

$$\Rightarrow \cos 120^\circ = -\frac{1}{2} = \frac{a^2 + \frac{a^2 + c^2 + 2ac}{4} - c^2}{2a \frac{(a+c)}{2}}$$

$$\Rightarrow -2a(a+c) = 4a^2 + a^2 + c^2 + 2ac - 4c^2$$

$$\Rightarrow -2a^2 - 2ac = 5a^2 - 3c^2 + 2ac$$

$$\Rightarrow 7a^2 + 4ac - 3c^2 = 0$$

$$\Rightarrow 7a^2 + 28a - 147 = 0 \quad (\because c = 7)$$

$$\Rightarrow a^2 + 4a - 21 = 0$$

$$\Rightarrow (a+7)(a-3) = 0$$

$$\Rightarrow a = 3 \text{ and } a \neq -7$$

$$\therefore b = 5$$

$$\text{Now, } s = \frac{a+b+c}{2} = \frac{3+5+7}{2} = \frac{15}{2}$$

$$\therefore \text{Area of triangle} = \sqrt{s(s-a)(s-b)(s-c)}$$

$$= \sqrt{\frac{15}{2} \left(\frac{15}{2} - 3\right) \left(\frac{15}{2} - 5\right) \left(\frac{15}{2} - 7\right)}$$

$$= \sqrt{\frac{15}{2} \cdot \frac{9}{2} \cdot \frac{5}{2} \cdot \frac{1}{2}} = \frac{15}{4} \sqrt{3} \text{ cm}^2$$

427 (a)

$$\text{Given, } \frac{b+c}{11} = \frac{a+c}{12} = \frac{a+b}{13} = k \text{ [say]}$$

$$\Rightarrow b+c = 11k, c+a = 12k, a+b = 13k$$

$$\Rightarrow 2(a+b+c) = 36k$$


$$\Rightarrow a+b+c = 18k$$

$$\Rightarrow a = 7k, b = 6k, c = 5k$$

$$\therefore \cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

$$= \frac{36 + 25 - 49}{2 \times 6 \times 5} = \frac{1}{5}$$

428 (c)




Angle AOM is 30° . Hence, required point of B is $(-3, 3\sqrt{3})$

429 (b)

Let $AB = BC = 2x$,

Then $BD = DC = x$ and $AD = (\sqrt{5})x$




Applying sine rule, $\frac{x}{\sin \beta} = \frac{AD}{\sin 45^\circ}$

$$\Rightarrow \sin \beta = \frac{\frac{x}{\sqrt{2}}}{(\sqrt{5})x} = \frac{1}{\sqrt{10}}$$

$$\text{and } \sin \alpha = \frac{x}{\sqrt{5}x} = \frac{1}{\sqrt{5}} \Rightarrow \frac{\sin \alpha}{\sin \beta} = \frac{\sqrt{2}}{1}$$

430 (c)

We know that, in a parallelogram diagonals cut each other at middle point



$$\therefore \frac{a+4}{2} = \frac{1+5}{2} \Rightarrow a = 2$$

$$\text{and } \frac{b+6}{2} = \frac{2+7}{2} \Rightarrow b = 3$$

431 (c)

Assume that angles are $4x, x$ and x

As $4x + x + x = 180^\circ$ [$\because \angle A + \angle B + \angle C = 180^\circ$]

$$\Rightarrow x = 30^\circ$$

\therefore Angles are $120^\circ, 30^\circ$ and 30°

Ratio of sides = $\sin A : \sin B : \sin C$

$$= \sin 120^\circ : \sin 30^\circ : \sin 30^\circ$$

$$= \sqrt{3} : 1 : 1$$

$$\therefore \text{Required ratio} = \frac{\sqrt{3}}{1+1+\sqrt{3}} = \frac{\sqrt{3}}{2+\sqrt{3}}$$

433 (c)

$$\text{Here, } s = \frac{18+24+30}{2} = 36$$

$$\text{Now, } \Delta = \sqrt{36(36-18)(36-24)(36-30)}$$

$$\Delta = \sqrt{36 \times 18 \times 12 \times 6} = 216$$

So, radius of the incircle

$$r = \frac{\Delta}{s} = \frac{216}{36} = 6 \text{ cm}$$

434 (b)

We know that the mid point of diagonals lies on line $y = 2x + c$, here mid point is $(3, 2)$, hence $c = -4$

435 (b)

Clearly, ΔOAB is an isosceles right angled triangle. So, its orthocentre is at $O(0,0)$ and the circumcentre is the mid-point of AB having coordinates $(a/2, a/2)$

$$\text{Hence, required distance} = \sqrt{\frac{a^2}{4} + \frac{a^2}{4}} = \frac{a}{\sqrt{2}}$$

436 (d)

The lines of a triangle are $x = y$, $x - 2y = 3$ and $x + 2y = -3$. Intersection points at sides are $A(-3, -3)$, $B(-1, -1)$ and $C(0, -\frac{3}{2})$

$$\therefore AB = \sqrt{4 + 4} = 2\sqrt{2}$$

$$AC = \sqrt{9 + \frac{9}{4}} = \frac{3\sqrt{5}}{2}$$

$$\text{and } BC = \sqrt{1 + \frac{1}{4}} = \frac{\sqrt{5}}{2}$$

437 (b)

Let the vertices of a triangle be $A(6, 0)$, $B(0, 6)$ and $C(6, 6)$

$$\text{Now, } AB = \sqrt{6^2 + 6^2} = 6\sqrt{2}$$

$$BC = \sqrt{6^2 + 0} = 6$$

$$\text{And } CA = \sqrt{0 + 6^2} = 6$$

$$\text{Also, } AB^2 = BC^2 + CA^2$$

Therefore, ΔABC is right angled at C . So, mid point of AB is the circumcentre of ΔABC

\therefore Coordinate of circumcentre are $(3, 3)$

Coordinates of centroid are,

$$G\left(\frac{6+0+6}{3}, \frac{0+6+6}{3}\right), \text{ i.e., } (4, 4)$$

$$\therefore \text{Required distance} = \sqrt{(4-3)^2 + (4-3)^2} = \sqrt{2}$$

438 (b)

Given that, $a = 5$, $b = 7$ and $\sin A = \frac{3}{4}$

$$\text{As we know, } \frac{\sin A}{a} = \frac{\sin B}{b}$$

$$\Rightarrow \frac{3}{4 \times 5} = \frac{\sin B}{7} \Rightarrow \sin B = \frac{21}{20}$$

Which is not possible because its value is greater than one

440 (a)

Since, coordinates of the centroid are

$\left(\frac{x_1+x_2+x_3}{3}, \frac{y_1+y_2+y_3}{3}\right)$, the centroid is always a rational point

441 (d)

\therefore In a ΔABC ,

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin\{\pi - (A+B)\}} = 2R$$

$$\Rightarrow \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin(A+B)} = 2R$$

8. If we know $a, \sin A, \sin B$, then we can find b, c, A, B and C

9. We can find A, B, C by using cosine rule

10. $\therefore a, \sin B, R$ are given, then we can find $\sin A, b$

11. $a, \sin A, R$ are given, then we know only the ratio $\frac{b}{\sin B}$

or $\frac{c}{\sin(A+B)}$; we cannot determine the values of $b, c, \sin B, \sin C$ separately

$\therefore \Delta ABC$ cannot be determined in this case

442 (b)

$$\text{Since, } A + C = \pi - B \Rightarrow \frac{A-B+C}{2} = \frac{\pi}{2} - B$$

$$\therefore 2ac \sin\left(\frac{A-B+C}{2}\right) = 2ac \cos B$$

$$= 2ca \cdot \frac{c^2 + a^2 - b^2}{2ca}$$

$$= a^2 + c^2 - b^2$$

443 (a)

$$a(b \cos C - c \cos B)$$

$$= ab \left(\frac{a^2 + b^2 - c^2}{2ab}\right) - ac \left(\frac{a^2 + c^2 - b^2}{2ac}\right)$$

$$= \frac{a^2 + b^2 - c^2}{2} - \frac{a^2 + c^2 - b^2}{2}$$


$$= b^2 - c^2$$

444 (b)

Since, D is the midpoint of BC . So, coordinate of D are $\left(\frac{x_2+x_3}{2}, \frac{y_2+y_3}{2}\right)$

Given, $G(7, 5)$ is the centroid of ΔABC

$$\therefore 7 = \frac{2+x_2+x_3}{3} \text{ and } 5 = \frac{3+y_2+y_3}{3}$$



$$\Rightarrow x_2 + x_3 = 21 - 2 \text{ and } y_2 + y_3 = 15 - 3$$

$$\Rightarrow \frac{x_2 + x_3}{2} = \frac{19}{2} \text{ and } \frac{y_2 + y_3}{2} = 6$$

$$\therefore \text{Coordinates of } D \left(\frac{19}{2}, 6 \right)$$

445 (b)

Since, the axes are rotated through an angle 45° , then we replace (x, y) by $(x \cos 45^\circ - y \sin 45^\circ, x \sin 45^\circ + y \cos 45^\circ)$ i.e., $\left(\frac{x}{\sqrt{2}} - \frac{y}{\sqrt{2}}, \frac{x}{\sqrt{2}} + \frac{y}{\sqrt{2}} \right)$ in the given equation

$$3x^2 + 3y^2 + 2xy = 2$$

$$\therefore 3 \left(\frac{x}{\sqrt{2}} - \frac{y}{\sqrt{2}} \right)^2 + 3 \left(\frac{x+y}{\sqrt{2}} \right)^2 + 2 \left(\frac{x-y}{\sqrt{2}} \right) \left(\frac{x+y}{\sqrt{2}} \right) = 2$$

$$\Rightarrow \frac{3}{2} (x^2 - y^2 - 2xy) + \frac{3}{2} (x^2 + y^2 + 2xy) + \frac{2}{2} (x^2 - y^2) = 2$$

$$\Rightarrow 4x^2 + 2y^2 = 2$$

$$\Rightarrow 2x^2 + y^2 = 1$$

446 (c)

Let $P(x, y)$ be any point on the line

Also, $(a_1 - a_2)x + (b_1 - b_2)y + c = 0 \dots(i)$

Since, $(x - a_1)^2 + (y - b_1)^2 = (x - a_2)^2 + (y - b_2)^2$

$$\Rightarrow x^2 + a_1^2 - 2a_1x + y^2 + b_1^2 - 2b_1y = x^2 + a_2^2 - 2a_2x + y^2 + b_2^2 - 2b_2y$$

$$\Rightarrow 2(a_2 - a_1)x + 2(b_2 - b_1)y = a_2^2 + b_2^2 - a_1^2 - b_1^2$$

$$\Rightarrow (a_1 - a_2)x + (b_1 - b_2)y + \frac{(a_2^2 + b_2^2 - a_1^2 - b_1^2)}{2} = 0 \dots(ii)$$

Since, Eqs. (i) and (ii) represents the same equation of line

$$\therefore c = \frac{a_2^2 + b_2^2 - a_1^2 - b_1^2}{2}$$

447 (d)

Let D be the centre of circumcircle


$$\therefore BD = 5 \text{ cm}$$

In $\triangle ABC$,

$$AC^2 + AB^2 + BC^2 = 100 = AB^2 + 36$$

$$\Rightarrow AB^2 = 64 \Rightarrow AB = 8$$

$$\therefore \text{Area of } \triangle ABC = \frac{1}{2} \times AB \times BC = \frac{1}{2} \times 8 \times 6 = 24 \text{ cm}^2$$



448 (c)

The given equation is $x^2 + 6xy + 8y^2 = 10 \dots(i)$

Since, axes are rotated through an angle $\frac{\pi}{4}$

$$\therefore x = x_1 \cos \frac{\pi}{4} - y_1 \sin \frac{\pi}{4} = \frac{x_1 - y_1}{\sqrt{2}}$$

$$\text{and } y = x_1 \sin \frac{\pi}{4} + y_1 \cos \frac{\pi}{4} = \frac{x_1 + y_1}{\sqrt{2}}$$

on putting the value of x and y in Eq. (i)

$$\left(\frac{x_1 - y_1}{\sqrt{2}} \right)^2 + 6 \left(\frac{x_1 - y_1}{\sqrt{2}} \right) \left(\frac{x_1 + y_1}{\sqrt{2}} \right) + 8 \left(\frac{x_1 + y_1}{\sqrt{2}} \right)^2 = 10$$

$$\Rightarrow x_1^2 + y_1^2 - 2x_1y_1 + 6x_1^2 - 6y_1^2 + 8x_1^2 + 8y_1^2 + 16x_1y_1 = 20$$

$$\Rightarrow 15x_1^2 + 3y_1^2 + 14x_1y_1 = 20$$


\therefore Required equation is

$$15x^2 + 14xy + 3y^2 = 20$$

449 (c)

In $\triangle ABE$, $\angle BAE = \angle AEB$

$\therefore AB = BE$



In $\triangle BCE$, using sine rule,

$$\frac{BE}{\sin(180^\circ - 3\alpha)} = \frac{CE}{\sin 2\alpha}$$

$$\Rightarrow CE = \frac{a \sin 2\alpha}{\sin 3\alpha} \dots (i)$$

Now, In $\triangle DCE$, $\sin 3\alpha = \frac{h}{CE}$

$$\Rightarrow \sin 3\alpha = \frac{h}{a \sin 2\alpha / \sin 3\alpha} \text{ [from Eq. (i)]}$$

$$\Rightarrow h = a \sin 2\alpha$$

450 (a)


We know that the x -axis divides the segment

joining $P(x_1, y_1)$ and (x_2, y_2) in the ratio $-y_1 : y_2$.

So, the required ratio is $-6 : -3$ i.e. $2 : 1$

451 (d)

$$\text{In } \Delta PAB, \tan \beta = \frac{AB}{AP}$$



$$\text{In } \Delta PAC, \tan \theta = \frac{AC}{AP}$$

$$\therefore \tan \alpha = \tan(\beta - \theta)$$

$$= \frac{\tan \beta - \tan \theta}{1 + \tan \beta \tan \theta}$$

$$= \frac{\frac{AB}{AP} - \frac{AC}{AP}}{1 + \frac{AB}{AP} \cdot \frac{AC}{AP}} \quad \dots(i)$$

$\therefore AP = n(AB) = n(2AC)$ ($\because C$ is the mid point of BA)

From Eq. (i), we get

$$\tan \alpha = \frac{\frac{1}{n} - \frac{1}{2n}}{1 + \frac{1}{2n^2}} = \frac{n}{2n^2 + 1}$$

$$\Rightarrow n = (2n^2 + 1) \tan \alpha$$

452 (b)

Let the three points be $A(-2, -5)$, $B(2, -2)$ and $C(8, a)$

If three points are collinear, then

slope of AB = slope of BC

$$\Rightarrow \frac{-2 + 5}{2 + 2} = \frac{a + 2}{8 - 2}$$

$$\Rightarrow \frac{3}{4} = \frac{a + 2}{6} \Rightarrow 9 = 2a + 4 \Rightarrow a = \frac{5}{2}$$

453 (a)

From given relation, we can write

$$\cos A + \cos C = 2(1 - \cos B)$$

$$\Rightarrow 2 \cos \frac{A+C}{2} \cos \frac{A-C}{2} = 2 \cdot 2 \sin^2 \frac{B}{2}$$

$$\Rightarrow \cos \frac{A-C}{2} = 2 \sin \frac{B}{2}$$

$$\Rightarrow 2 \sin \frac{A+C}{2} \cdot \cos \frac{A-C}{2} = 4 \sin \frac{B}{2} \cos \frac{B}{2}$$

$$\Rightarrow \sin A + \sin C = 2 \sin B$$

$$\Rightarrow a, b, c \text{ are in AP}$$

454 (d)

$$\text{Given, } \cos(A+B) = \frac{31}{32} \Rightarrow \cos(\pi - C) = \frac{31}{32}$$

$$\Rightarrow -\cos C = -\frac{a^2 + b^2 - c^2}{2ab} = \frac{31}{32}$$

$$\Rightarrow \frac{(5)^2 + (4)^2 - c^2}{2 \times 5 \times 4} = \frac{31}{32}$$

$$\Rightarrow 41 - c^2 = -\frac{155}{4}$$

$$\Rightarrow c^2 = \frac{319}{4}$$

$$\Rightarrow c = \frac{\sqrt{319}}{2}$$

455 (a)

$$\text{We know, } r_1 = \frac{\Delta}{s-a}, r_2 = \frac{\Delta}{s-b}, r_3 = \frac{\Delta}{s-c}$$

Given that, $r_1 > r_2 > r_3$

$$\Rightarrow \frac{\Delta}{s-a} > \frac{\Delta}{s-b} > \frac{\Delta}{s-c}$$

$$\Rightarrow \frac{1}{s-a} > \frac{1}{s-b} > \frac{1}{s-c}$$

$$\Rightarrow s-a < s-b < s-c$$

($s-a, s-b, s-c$ are positive)

$$\Rightarrow -a < -b < -c$$

$$\Rightarrow a > b > c$$

456 (b)

We have, $a = 2x$, $b = 2y$ and $\angle C = 120^\circ$

$$\text{Area of triangle, } \Delta = \frac{1}{2} ab \sin C$$

$$= \frac{1}{2} \times 2x \times 2y \sin 120^\circ$$

$$= xy\sqrt{3} \text{ sq unit}$$

457 (a)

$$\frac{a - a' - a'}{a' - a} = \frac{b - b' - b'}{b' - b} \quad \left(\because \frac{x_3 - x_2}{x_2 - x_1} = \frac{y_3 - y_2}{y_2 - y_1} \right)$$

$$\Rightarrow \frac{a - 2a'}{a' - a} = \frac{b - 2b'}{b' - b}$$

$$\Rightarrow \frac{a}{a'} = \frac{b}{b'}$$

$$\Rightarrow ab' = a'b$$

459 (c)

If two vertices of an equilateral triangle have the coordinates (x_1, y_1) and (x_2, y_2) , then the coordinates of its third vertex are

$$\left(\frac{x_1 + x_2 \pm \sqrt{3}(y_1 - y_2)}{2}, \frac{y_1 + y_2 \pm \sqrt{3}(x_1 - x_2)}{2} \right)$$

Here, we have

$$x_1 = 2, y_1 = 4, x_2 = 2 \text{ and } y_2 = 6$$

Hence, the coordinates of the third vertex are

$$\left(\frac{4 \pm \sqrt{3}(-2)}{2}, \frac{10 \pm \sqrt{3} \times 0}{2} \right) = (2 \mp \sqrt{3}, 5)$$

461 (a)

$$\because A + B + C + D = 2\pi$$

$$\Rightarrow \tan(A + B + C + D) = 0$$

$$\Rightarrow \frac{\sum \tan A - \sum \tan A \tan B \tan C}{1 - \sum \tan A \tan B + \tan A \tan B \tan C \tan D} = 0$$

$$\Rightarrow \sum \tan A - \sum \tan A \tan B \tan C = 0$$

$$\Rightarrow \sum \tan A = \tan A \tan B \tan C \tan D \sum \cot A$$

$$\Rightarrow \frac{\sum \tan A}{\sum \cot A} = \prod \tan A$$

462 (a)

Let the angles of a triangle are $30^\circ, 40^\circ, 50^\circ$

$$\therefore 30^\circ + 40^\circ + 50^\circ = 180^\circ \Rightarrow \theta = 15^\circ$$

\therefore Angles of a triangle are $45^\circ, 60^\circ, 75^\circ$

$$\text{Now, } \sin A = \sin 45^\circ = \frac{1}{\sqrt{2}}$$

$$\sin B = \sin 60^\circ = \frac{\sqrt{3}}{2}$$

$$\text{and } \sin C = \sin 75^\circ = \frac{\sqrt{3}+1}{2\sqrt{2}}$$


$$\therefore a : b : c = \sin A : \sin B : \sin C$$

$$= \frac{1}{\sqrt{2}} : \frac{\sqrt{3}}{2} : \frac{\sqrt{3}+1}{2\sqrt{2}}$$

$$= 2 : \sqrt{6} : \sqrt{3} + 1$$

463 (a)

Since, $AB = AC = \sqrt{2}$



The slope AB is 1. Hence, AB is inclined at 45° with the x -axis and AC is inclined at 60° with the x -axis. Equation of AC is

$$y = \sqrt{3}(x - 2)$$

The coordinates of C is $(2 + \sqrt{2} \cos 60^\circ, 0 + \sqrt{2} \sin 60^\circ)$


$$\text{or } \left(2 + \frac{1}{\sqrt{2}}, \sqrt{\frac{3}{2}} \right)$$

464 (b)

Let $(x_1, y_1), (x_2, y_2)$ and (x_3, y_3) are coordinates of the points D, E and F , which divide each AB, BC and CA respectively in the ratio 3:1 (internally)

$$\therefore x_1 = \frac{3 \times 6 - 1 \times 1}{4} = \frac{17}{4}$$

$$y_1 = \frac{-2 \times 3 + 4 \times 1}{4} = -\frac{2}{4} = -\frac{1}{2}$$



$$(6, -2)B \quad E(x_2, y_2) \quad C(-2, 4)$$

$$\text{Similarly, } x_2 = 0, y_2 = \frac{5}{2}$$

$$\text{and } x_3 = -\frac{5}{4}, y_3 = 4$$

Let (x, y) be the coordinates of centroid of ΔDEF

$$\therefore x = \frac{1}{3} \left(\frac{17}{4} + 0 - \frac{5}{4} \right) = 1$$

$$\text{and } y = \frac{1}{3} \left(-\frac{1}{2} + \frac{5}{2} + 4 \right) = 2$$

\therefore Coordinates of Centroid are $(1, 2)$

465 (a)

Let D be the position of window of second house and BC be the position of the first house.


$$\text{In } \Delta ADB, \tan 60^\circ = \frac{AD}{AB}$$

$$\Rightarrow AD = 6\sqrt{3}$$

$$\text{and } DB^2 = (6\sqrt{3})^2 + (6)^2$$

$$\Rightarrow DB = 12m$$

$$\text{In } \Delta DCB, \sin 60^\circ = \frac{12}{h}$$



$$h = \frac{12}{\sin 60^\circ} = 8\sqrt{3}m$$

466 (b)

Let the vertices be C, A, B respectively. The altitude from A is

$$\frac{y - a(t_2 + t_3)}{x - at_2t_3} = -t_1$$

$$\Rightarrow xt_1 + y = at_1t_2t_3 + a(t_2 + t_3) \dots(i)$$

The altitude from B is

$$xt_2 + y = at_1t_2t_3 + a(t_3 + t_1) \dots(ii)$$

Subtracting Eq. (ii) from Eq. (i), $x = -a$


$$\text{Hence, } y = a(t_1 + t_2 + t_3 + t_1t_2t_3)$$

\therefore The orthocenter is $\{-a, a(t_1 + t_2 + t_3 + t_1t_2t_3)\}$

467 (a)

$$\text{In } \Delta ABC, \tan 30^\circ = \frac{BC}{40}$$

$$\Rightarrow BC = \frac{40}{\sqrt{3}} \dots(i)$$



$$\text{In } \Delta DAB, \tan 60^\circ = \frac{BD}{40}$$

$$\Rightarrow BD = 40\sqrt{3}$$

$$\Rightarrow DC + BC = 40\sqrt{3}$$


$$\Rightarrow DC + \frac{40}{\sqrt{3}} = 40\sqrt{3} \text{ [from Eq.(i)]}$$

$$\Rightarrow DC = 40 \left(\sqrt{3} - \frac{1}{\sqrt{3}} \right) = \frac{80}{\sqrt{3}}$$

$$\Rightarrow DC = 46.19 \text{ m}$$

468 (c)

Let DC be the height of building



$$\therefore AB = \text{Speed} \times \text{Time}$$

$$= 25(\sqrt{3} - 1) \cdot 2$$

$$= 50(\sqrt{3} - 1)$$

$$\therefore \text{In } \triangle DBC, \tan 45^\circ = \frac{DC}{BC} \Rightarrow BC = h$$

$$\text{In } \triangle DAC, \tan 30^\circ = \frac{h}{50(\sqrt{3}-1)+h}$$

$$\Rightarrow 50(\sqrt{3} - 1) + h = \sqrt{3}h$$

$$\Rightarrow h = 50 \text{ m}$$

469 (d)

$$(b+c) \tan \frac{A}{2} \tan \left(\frac{B-C}{2} \right)$$

$$= (b+c) \tan \frac{A}{2} \cdot \frac{(b-c)}{(b+c)} \cot \frac{A}{2}$$

$$= b-c$$

$$\therefore \sum (b+c) \tan \left(\frac{B-C}{2} \right) \tan \frac{A}{2}$$

$$= b-c + c-a + a-b = 0$$

470 (c)

As, $A > B$ and $3 \sin x - 4 \sin^3 x - k = 0, 0 < k < 1$

$$1$$

$$\Rightarrow \sin 3x = k$$

As A and B satisfy given equation

$$\therefore \sin 3A = k, \quad \sin 3B = k$$

$$\Rightarrow \sin 3A - \sin 3B = k$$

$$\Rightarrow 2 \cos \left(\frac{3A+3B}{2} \right) \sin \left(\frac{3A-3B}{2} \right) = 0$$

$$\Rightarrow \cos \left(\frac{3A+3B}{2} \right) = 0 \quad \text{or} \quad \sin \left(\frac{3A-3B}{2} \right) = 0$$

$$\Rightarrow \frac{3A+3B}{2} = 90^\circ \quad \text{or} \quad \frac{3A-3B}{2} = 0$$

$$\Rightarrow A+B = 60^\circ \quad \text{or} \quad A=B$$

But given, $A > B$ (\therefore neglecting $A=B$)

Thus, $A+B = 60^\circ$

and $A+B+C = 180^\circ \Rightarrow \angle C = 120^\circ$

471 (b)

$$\text{We have, } \cos A = \frac{b^2+c^2-a^2}{2bc}$$

$$\Rightarrow b^2 - 2bc \cos A + (c^2 - a^2) = 0$$

It is given that b_1 and b_2 are roots of this equation

$$\therefore b_1 + b_2 = 2c \cos A \quad \text{and} \quad b_1 b_2 = c^2 - a^2$$

$$\Rightarrow 3b_1 = 2c \cos A \quad \text{and} \quad 2b_1^2 = c^2 - a^2 \quad [\because b_2 = 2b_1 \text{ (given)}]$$

$$\Rightarrow 2 \left(\frac{2c}{3} \cos A \right)^2 = c^2 - a^2$$

$$\Rightarrow 8c^2(1 - \sin^2 A) = 9c^2 - 9a^2$$

$$\Rightarrow \sin^2 A = \left(1 - \frac{9c^2 - 9a^2}{8c^2} \right)$$

$$\Rightarrow \sin A = \sqrt{\frac{9a^2 - c^2}{8c^2}}$$

472 (a)

Let the vertices of the $\triangle ABC$ are $A(7, 1), B(-1, 5)$

and $C(3 + 2\sqrt{3}, 3 + 4\sqrt{3})$

$$\text{Now, } AB = \sqrt{(7+1)^2 + (1-5)^2}$$

$$= \sqrt{64 + 16} = \sqrt{80}$$

$$BC = \sqrt{(-1-3-2\sqrt{3})^2 + (5-3-4\sqrt{3})^2}$$

$$= \sqrt{16 + 12 + 16\sqrt{3} + 4 + 48 - 16\sqrt{3}}$$

$$= \sqrt{80}$$

$$\text{and } CA = \sqrt{(3+2\sqrt{3}-7)^2 + (3+4\sqrt{3}-1)^2}$$

$$= \sqrt{16 + 12 - 16\sqrt{3} + 4 + 48 + 16\sqrt{3}}$$

$$= \sqrt{80}$$

Here, $AB = BC = CA = \sqrt{80}$

\therefore Hence, it is an equilateral triangle, so incentre and centroid coincides

$$\text{So, incentre} = \left(\frac{7-1+3+2\sqrt{3}}{3}, \frac{1+5+3+4\sqrt{3}}{3} \right)$$

$$= \left(\frac{9+2\sqrt{3}}{3}, \frac{9+4\sqrt{3}}{3} \right)$$

$$= \left(3 + \frac{2}{\sqrt{3}}, 3 + \frac{4}{\sqrt{3}} \right)$$

473 (b)

$$\tan \frac{A}{2} - \tan \frac{B}{2} = \frac{\sqrt{\frac{(s-b)(s-c)}{s(s-a)}} - \sqrt{\frac{(s-a)(s-c)}{s(s-b)}}}{\tan \frac{A}{2} + \tan \frac{B}{2} = \frac{\sqrt{\frac{(s-b)(s-c)}{s(s-a)}} + \sqrt{\frac{(s-a)(s-c)}{s(s-b)}}}$$

$$= \frac{(s-b)\sqrt{s(s-c)} - (s-a)\sqrt{s(s-c)}}{(s-b)\sqrt{s(s-c)} + (s-a)\sqrt{s(s-c)}}$$

$$= \frac{\sqrt{s(s-c)}(s-b-s+a)}{\sqrt{s(s-c)}(s-b-s+a)} = \frac{a-b}{c}$$

474 (a)

As we know that orthocenter, centroid and circumcentre are collinear and the centroid

divides the line segment joining orthocenter and circumcentre in the ratio 2:1. If the coordinates of

orthocentre and circumcentre are $(1, 1)$ and $(3, 2)$

respectively, the coordinate of centroid is

$$\left(\frac{2.3 + 1.1}{2 + 1}, \frac{2.2 + 1.1}{2 + 1}\right) = \left(\frac{7}{3}, \frac{5}{3}\right)$$

475 (d)

$$\because \angle A = 60^\circ, \cos 60^\circ = \frac{b^2 + c^2 - a^2}{2bc}$$

$$\Rightarrow b^2 + c^2 - a^2 = bc$$

$$\text{Now, } AB^2 + AC^2 = 2(AD^2 + BD^2)$$

$$\Rightarrow c^2 + b^2 = 2AD^2 + 2\left(\frac{a}{2}\right)^2$$

$$\Rightarrow 2b^2 + 2c^2 - a^2 = 4AD^2$$

$$\Rightarrow b^2 + c^2 + bc = 4AD^2 \quad (\because b^2 + c^2 - a^2 = bc)$$

476 (b)

The intersection of lines are $(0, 0)$, $(-4, -8)$ and $(-2, -2)$

Let circumcentre is (x_1, y_1)

$$\therefore x_1^2 + y_1^2 = (x_1 + 4)^2 + (y_1 + 8)^2$$

$$\Rightarrow 8x_1 + 16y_1 + 80 = 0 \dots(i)$$

$$\text{And } x_1^2 + y_1^2 = (x_1 + 2)^2 + (y_1 + 2)^2$$

$$\Rightarrow 4x_1 + 4y_1 + 8 = 0 \dots(ii)$$

On solving Eqs. (i) and (ii), we get

$$x_1 = 6 \text{ and } y_1 = -8$$

477 (c)

$$\text{Given, } \cot \frac{A}{2} = \frac{b+c}{a}$$

$$\Rightarrow \frac{\cos \frac{A}{2}}{\sin \frac{A}{2}} = \frac{\sin B + \sin C}{\sin A}$$

$$= \frac{2 \sin \frac{B+C}{2} \cos \frac{B-C}{2}}{2 \sin \frac{A}{2} \cos \frac{A}{2}}$$

$$\Rightarrow \cos \frac{A}{2} = \cos \frac{B-C}{2}$$

$$\therefore \frac{A}{2} = \frac{B-C}{2}$$

$$\Rightarrow A + C = B \Rightarrow \angle B = \frac{\pi}{2} \quad (\because A + B + C = \pi)$$

478 (c)

Given lines are

$$3x^2 - 4xy + y^2 = 0$$

$$\Rightarrow (3x - y)(x - y) = 0$$

$$\Rightarrow 3x - y = 0, x - y = 0 \text{ and } 2x - y = 6$$

The points of intersection of these lines are

$$(0, 0), (-6, -18) \text{ and } (6, 6)$$


$$\therefore \text{Area of triangle} = \frac{1}{2} \begin{vmatrix} 0 & 0 & 1 \\ -6 & -18 & 1 \\ 6 & 6 & 1 \end{vmatrix}$$

$$= \frac{1}{2}(-36 + 108) = \frac{1}{2}(72)$$

$$= 36 \text{ sq units}$$

479 (d)

$$\text{In } \triangle ABC, \tan 60^\circ = \frac{h}{x} \Rightarrow h\sqrt{3}x \dots(i)$$



and in $\triangle ABD$,

$$\tan 30^\circ = \frac{h}{x + 60} \Rightarrow x + 60 = \sqrt{3}h \dots(ii)$$


From Eqs. (i) and (ii),

$$h = \sqrt{3} \times 30 = 51.96 \text{ m}$$

$$= 52 \text{ m (approx)}$$

480 (d)

OP is inclined at angle θ with x -axis and OQ is inclined at angle $\alpha - \theta$ with x -axis




The bisector of angle POQ is inclined at angle

$$\frac{\theta + \alpha - \theta}{2} = \frac{\alpha}{2} \text{ with } x\text{-axis}$$

481 (b)

$$\text{The altitude } h \text{ is } \frac{|2-1-2|}{\sqrt{2}} = \frac{1}{\sqrt{2}}$$




$$\text{In } \triangle ABC, BC = 2h \cot 60^\circ = \sqrt{\frac{2}{3}}$$

$$\therefore \text{Area of } \triangle = \frac{h}{2} BC$$

$$= \frac{1}{2\sqrt{2}} \sqrt{\frac{2}{3}} = \frac{\sqrt{3}}{6}$$

483 (d)

Let OP be the tower. Since, the tower make equal angles at the vertices of the triangle, therefore foot of tower is at the circumcentre



$$\text{In } \triangle OAP, \tan \alpha = \frac{OP}{OA}$$

$$\Rightarrow OP = OA \tan \alpha$$

$$\Rightarrow OP = R \tan \alpha \quad (\because OA = R, \text{ given})$$

484 (a)


$$\text{Given, } xy + 2x + 2y + 4 = 0 \quad \dots(i)$$

$$\text{and } x + y + 2 = 0 \quad \dots(ii)$$

From Eqs. (i) and (ii),

$$xy = 0$$

$$\Rightarrow x = y = 0$$



\therefore vertices of triangle are $(-2, 0), (0, 0), (0, -2)$

Since, this triangle is right angled triangle and in a right angled triangle circumcentre is mid point of hypotenuse.

$\therefore (-1, -1)$ is the circumcentre

485 (b)

Let P be (x, y) and we have

$$A = (a + b, a - b), \quad B = (a - b, a + b)$$

$$\text{Here, } PA = PB \Rightarrow PA^2 = PB^2$$

$$\Rightarrow [x - (a + b)]^2 + [y - (a - b)]^2$$

$$= [x - (a - b)]^2 + [y - (a + b)]^2$$

$$\Rightarrow [x - (a + b)]^2 - [x - (a - b)]^2$$

$$= [y - (a + b)]^2 - [y - (a - b)]^2$$

$$\Rightarrow [x - (a + b) + x - (a - b)][x - (a + b) - x + a - b]$$

$$= [y - (a + b) + y - (a - b)][y - (a + b) - y + a - b]$$

$$\Rightarrow (2x - 2a)(-2b) = (2y - 2a)(-2b)$$

$$\Rightarrow x - y = 0$$

486 (c)

$$\text{Since, } R = \frac{a}{2 \sin A}$$

$$\Rightarrow R = \frac{2\sqrt{3}}{2 \sin 60^\circ} = 2 \text{ cm}$$

487 (c)

Let two of vertices of a triangle are $A(15, 0)$ and $B(0, 10)$ and third vertex is $C(h, k)$

We know that line passing through A and B should be perpendicular to line through C and orthocenter O

$$\therefore \left(\frac{-2}{3}\right)\left(\frac{9-k}{6-h}\right) = -1$$

$$\Rightarrow 2k = 3h$$

Which is satisfied by $(0, 0)$. Hence, coordinates of third vertex are $(0, 0)$


488 (a)

Let P and Q be the positions of two planes. It is given that $OP = 300$ m. From triangle OAQ , we have $OA = OQ$.

From ΔOAP , we have

$$\tan 60^\circ = \frac{OP}{OA} \Rightarrow \sqrt{3} = \frac{300}{OQ} \Rightarrow OQ = \frac{300}{\sqrt{3}}$$

$$= 100\sqrt{3} \text{ mts}$$



489 (b)

Let the sides be $a = 3x, b = 7x, c = -8x$. Then,

$$2s = a + b + c \Rightarrow s = 9x$$

$$\therefore \Delta = \sqrt{s(s-a)(s-b)(s-c)}$$

$$= \sqrt{9x \times 6x \times 2x \times x} = 6\sqrt{3}x^2$$

$$\text{Now, } R = \frac{abc}{4\Delta} \text{ and } r = \frac{\Delta}{s}$$

$$\therefore \frac{R}{r} = \frac{sabc}{4\Delta^2} = \frac{9x \times 3x \times 7x \times 8x}{4 \times 108 \times x^2} = \frac{7}{2}$$

490 (a)

$$(b + c - a) \tan \frac{A}{2} = (2s - 2a) \tan \frac{A}{2}$$

$$= 2(s - a) \sqrt{\frac{(s-b)(s-c)}{s(s-a)}}$$


$$= 2 \frac{\sqrt{s(s-a)(s-b)(s-c)}}{s} = \frac{2\Delta}{s}$$

491 (b)

Let $AB = a$

$ON \perp AB$

and $AN = BN$



$$\text{In } \Delta AON, \tan \frac{\pi}{n} = \frac{AN}{ON}$$

$$\Rightarrow ON = AN \cot \frac{\pi}{n} = \frac{a}{2} \cot \frac{\pi}{n}$$

$$\text{and } \sin \frac{\pi}{n} = \frac{AN}{OA}$$

$$\Rightarrow OA = \frac{a}{2} \operatorname{cosec} \frac{\pi}{n} \quad \dots(ii)$$

Now, sum of the radii = $ON + OA$

$$= \frac{a}{2} \cot \frac{\pi}{n} + \frac{a}{2} \operatorname{cosec} \frac{\pi}{n}$$

[from Eqs. (i) and (ii)]

$$= \frac{a}{2} \left[\frac{1 + \cos \frac{\pi}{n}}{\sin \frac{\pi}{n}} \right]$$

$$= \frac{a}{n} \left[\frac{2 \cos^2 \frac{\pi}{2n}}{2 \sin \frac{\pi}{2n} \cos \frac{\pi}{2n}} \right]$$

$$= \frac{a}{2} \cot \frac{\pi}{2n}$$

492 (c)

$$\Delta = 2bc - (b^2 + c^2 - a^2) = 2bc(1 - \cos A)$$

$$= 2bc \cdot 2 \sin^2 \frac{A}{2} \dots (i)$$

$$\text{But } \Delta = \frac{1}{2} bc \sin A = \frac{1}{2} bc \cdot 2 \sin \frac{A}{2} \cos \frac{A}{2}$$

$$\Rightarrow \Delta = bc \sin \frac{A}{2} \cos \frac{A}{2} \dots (ii)$$


On dividing Eq. (ii) by Eq. (i), we get

$$\tan \frac{A}{2} = \frac{1}{4}$$

$$\therefore \tan A = \frac{2 \tan \frac{A}{2}}{1 - \tan^2 \frac{A}{2}} = \frac{\frac{1}{2}}{1 - \frac{1}{16}} = \frac{8}{15}$$

493 (d)

The vertices of a ΔABC are $A(0, b)$, $B(0, 0)$ and $C(a, 0)$



Mid point of E and D are $(\frac{a}{2}, \frac{b}{2})$ and $(\frac{a}{2}, 0)$

The slope of median BE , $m_1 = \frac{b}{a}$

and slope of AD , $m_2 = -\frac{2b}{a}$

Since, the medians are perpendicular to each other

$$\therefore m_1 m_2 = -1$$

$$\Rightarrow \frac{b}{a} \times \left(-\frac{2b}{a} \right) = -1$$

$$\Rightarrow -2b^2 = -a^2 \Rightarrow a = \pm \sqrt{2}b$$

494 (c)

$$\therefore a(b - c) + b(c - a) + c(a - b) = 0$$

$\therefore x = 1$ is a root of the equation

$$a(b - c)x^2 + b(c - a)x + c(a - b) = 0$$

Then, other root = 1 (\because roots are equal)

$$\therefore \text{Product of roots} = 1 = \frac{c(a - b)}{a(b - c)}$$

$$\Rightarrow ab - ac = ca - bc$$

$$\Rightarrow b = \frac{2ac}{a + c}$$

$\therefore a, b, c$ are in HP

Then, $\frac{1}{a}, \frac{1}{b}, \frac{1}{c}$ are in AP

$\Rightarrow \frac{s}{a}, \frac{s}{b}, \frac{s}{c}$ are in AP

$\Rightarrow \frac{s}{a} - 1, \frac{s}{b} - 1, \frac{s}{c} - 1$ are in AP

$\Rightarrow \frac{(s-a)}{a}, \frac{(s-b)}{b}, \frac{(s-c)}{c}$ are in AP

Multiplying each by $\frac{abc}{(s-a)(s-b)(s-c)}$

Then $\frac{bc}{(s-b)(s-c)}, \frac{ca}{(s-c)(s-a)}, \frac{ab}{(s-a)(s-b)}$ are in AP

$\Rightarrow \frac{(s-b)(s-c)}{bc}, \frac{(s-c)(s-a)}{ca}, \frac{(s-a)(s-b)}{ab}$ are in HP

$\Rightarrow \sin^2 \left(\frac{A}{2} \right), \sin^2 \left(\frac{B}{2} \right), \sin^2 \left(\frac{C}{2} \right)$ are in HP

495 (d)

$$\frac{1 + \cos C \cos(A - B)}{1 + \cos(A - C) \cos B}$$


$$= \frac{1 - \cos(A + B) \cos(A - B)}{1 - \cos(A - C) \cos(A + C)} \quad (\because A + B + C = \pi)$$

$$\Rightarrow \frac{1 - \cos^2 A + \sin^2 B}{1 - \cos^2 A + \sin^2 C} = \frac{\sin^2 A + \sin^2 B}{\sin^2 A + \sin^2 C}$$

$$= \frac{a^2 + b^2}{a^2 + c^2}$$

496 (c)

$$\therefore b^2 + c^2 = a^2$$



$\therefore \Delta ABC$ is right angled triangle and right angled at A

Clearly, from figure,

$$\frac{1}{2}(AB)(AC) = \frac{1}{2}(AD)(BC)$$

$$\Rightarrow AD = \frac{(AB)(AC)}{BC}$$

$$= \frac{5 \cdot 12}{13} = \frac{60}{13}$$

497 (a)

Since, $r_1 < r_2 < r_3$

$$\Rightarrow \frac{1}{r_1} > \frac{1}{r_2} > \frac{1}{r_3}$$

$$\Rightarrow \frac{s-a}{\Delta} > \frac{s-b}{\Delta} > \frac{s-c}{\Delta}$$

$$\Rightarrow (s-a) > (s-b) > (s-c)$$

$$\Rightarrow -a > -b > -c$$

$$\Rightarrow a < b < c$$


498 (a)

Let l be the length of the ladder

ie, $BP = CQ = 1$

In ΔPAB , $\cos \alpha = \frac{PA}{PB}$ and $\sin \alpha = \frac{AB}{PB}$
 $\Rightarrow PA = l \cos \alpha$ and $AB = l \sin \alpha$... (i)

In ΔQAC , $\cos \beta = \frac{AQ}{QC}$
 $\Rightarrow AQ = l \cos \beta$ and $AC = l \sin \beta$... (ii)



Now, $CB = AB - CA$
 $= l \sin \alpha - l \sin \beta$ [from Eqs. (i) and (ii)]
 $= l(\sin \alpha - \sin \beta)$

and $QP = AQ - PA$
 $= l \cos \beta - l \cos \alpha$

$= l(\cos \beta - \cos \alpha)$
 $\therefore \frac{CB}{QP} = \frac{l(\sin \alpha - \sin \beta)}{l(\cos \beta - \cos \alpha)}$


$\Rightarrow \frac{y}{x} = \frac{2 \sin \left(\frac{\alpha - \beta}{2}\right) \cos \left(\frac{\alpha + \beta}{2}\right)}{2 \sin \left(\frac{\alpha + \beta}{2}\right) \cos \left(\frac{\alpha - \beta}{2}\right)}$

$\Rightarrow \frac{y}{x} = \cot \left(\frac{\alpha + \beta}{2}\right)$

$\Rightarrow x = y \tan \left(\frac{\alpha + \beta}{2}\right)$

499 (c)

$\because A(h, k), B(1, 1)$ and $C(2, 1)$ are the vertices of a right angled triangle ABC



Now, $AB = \sqrt{(1-h)^2 + (1-k)^2}$

Or $BC = \sqrt{(2-1)^2 + (1-1)^2} = 1$

Or $CA = \sqrt{(h-2)^2 + (k-1)^2}$

Now, Pythagoras theorem

$AC^2 = AB^2 + BC^2$

$\Rightarrow 4 + h^2 - 4h + k^2 + 1 - 2k$
 $= h^2 + 1 - 2h + k^2 + 1 - 2k + 1$

$\Rightarrow 5 - 4h = 3 - 2h$

$\Rightarrow h = 1$... (i)

Now, given that area of the triangle is 1

Then, area (ΔABC) = $\frac{1}{2} \times AB \times BC$

$\Rightarrow 1 = \frac{1}{2} \times \sqrt{(1-h)^2 + (1-k)^2} \times 1$

$\Rightarrow 2 = \sqrt{(1-h)^2 + (1-k)^2}$... (ii)

On putting $h = 1$ from Eq. (i), we get

$2 = \sqrt{(k-1)^2}$

On squaring both the sides, we get


$4 = k^2 + 1 - 2k$

$\Rightarrow k^2 - 2k - 3 = 0 \Rightarrow k = -1, 3$

Thus, the set of values of k is $\{-1, 3\}$

500 (d)

Let the coordinate of mid point of AB is (x_1, y_1)



$\therefore x_1 = \frac{a+0}{2}, y_1 = \frac{0+b}{2}$

$\Rightarrow a = 2x_1, b = 2y_1$

Given, $a + b = 4 \Rightarrow x_1 + y_1 = 2$

Hence, the locus of the mid point is $x + y = 2$

501 (a)

$\because \tan \frac{A}{2}, \tan \frac{B}{2}, \tan \frac{C}{2}$ are in HP

$\therefore \cot \frac{A}{2}, \cot \frac{B}{2}, \cot \frac{C}{2}$ are in AP

$\Rightarrow \cot \frac{B}{2} - \cot \frac{A}{2} = \cot \frac{C}{2} - \cot \frac{B}{2}$

$\Rightarrow \frac{s(s-b)}{\Delta} - \frac{s(s-a)}{\Delta} = \frac{s(s-c)}{\Delta} - \frac{s(s-b)}{\Delta}$


$\Rightarrow s - b - s + a = s - c - s + b$

$\Rightarrow 2b = a + c$

$\Rightarrow a, b, c$ are in AP

502 (a)

In ΔACD , $\tan 30^\circ = \frac{45\sqrt{3}/2}{x}$



$\Rightarrow x = \frac{45\sqrt{3}}{2 \times \frac{1}{\sqrt{3}}} = \frac{135}{2} \text{ m}$

and in ΔABD ,

$\tan 60^\circ = \frac{45\sqrt{3}/2}{y} \Rightarrow y = \frac{45}{2}$

$\therefore x - y = \frac{135}{2} - \frac{45}{2} = 45 \text{ m}$

503 (d)

For point $(1, 3)$, $3x + 2y = 3 + 6 > 0$

For point (5, 0), $3 \times 5 + 0 > 0$
 and for point (-1, 2), $-3 + 4 > 0$
 Similarly, other inequalities also hold
 Hence, option (d) is correct

504 (c)

Given that, $x_1 = x$, $x_2 = 1$, $x_3 = 0$
 and $y_1 = 0$, $y_2 = 1$, $y_3 = 2$

$$\begin{aligned} \therefore \text{Area of triangle} &= \frac{1}{2} [x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2)] \\ &= \frac{1}{2} [x(1 - 2) + 1(2 - 0) + 0(0 - 1)] \\ &= \frac{1}{2} [-x + 2 + 0] = \frac{1}{2} (2 - x) \end{aligned}$$

But area of triangle is 4 sq unit

$$\therefore \frac{1}{2} (2 - x) = 4$$

$$\Rightarrow 2 - x = 8 \Rightarrow x = -6$$

505 (c)

Given vertices of triangle are
 $O(0, 0)$, $A(\cos \theta, \sin \theta)$ and $B(\sin \theta, -\cos \theta)$, then
 coordinates of Centroid are

$$\left(\frac{\cos \theta + \sin \theta}{3}, \frac{\sin \theta - \cos \theta}{3} \right)$$

Since, Centroid lies on the line $y = 2x$

$$\therefore \frac{\sin \theta - \cos \theta}{3} = \frac{2 \cos \theta + 2 \sin \theta}{3}$$

$$\Rightarrow \sin \theta = -3 \cos \theta$$

$$\Rightarrow \theta = \tan^{-1}(-3)$$

506 (a)

In order to remove first degree terms from the
 equation $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$
 the origin is shifted at $(-g/a, -f/a)$

In the equation $2x^2 + 7y^2 + 8x - 14y + 4 = 0$,
 we have

$$a = 2, b = 7, g = 4 \text{ and } f = -7$$

Hence, the coordinates of the required point are
 $(-4/2, -7/7) = (-2, 1)$

507 (b)

Let $A[at_1t_2, a(t_1 + t_2)]$, $B[at_2t_3, a(t_2 + t_3)]$,
 $C[at_3t_1, a(t_3 + t_1)]$

$$\text{Slope of } AB (m_{AB}) = \frac{a(t_3 - t_1)}{at_2(t_3 - t_1)} = \frac{1}{t_2}$$

Equation of line through C perpendicular to AB is

$$y - a(t_3 + t_1) = -t_2(x - at_3t_1)$$

$$\Rightarrow y - a(t_3 + t_1) = -t_2x + at_1t_2t_3 \quad \dots(i)$$

Similarly, equation of line through B

perpendicular to CA is

$$y - a(t_2 + t_3) = -t_1(x - at_2t_3)$$

$$\Rightarrow y - a(t_2 + t_3) = -t_1x + at_1t_2t_3 \quad \dots(ii)$$

Using $t_1t_2t_3 = -(t_1 + t_2 + t_3)$ in Eqs. (i) and (ii),
 we get

$$y = -t_2x - at_2$$

$$\text{and } y = -t_1x - at_1$$

$$\Rightarrow t_2(x + a) = t_1(x + a)$$

$$\Rightarrow x = -a, y = 0$$

508 (d)

Let the coordinates of the third vertex C be (h, k) .

Then,

$$\frac{1}{2} \begin{vmatrix} h & k & 1 \\ -5 & 0 & 1 \\ 3 & 0 & 1 \end{vmatrix} = \pm 20 \Rightarrow k = \pm 5$$

Since, (h, k) lies on $x - y = 2$. Therefore, $h - k = 2$

For $k = 5$, $h = 7$. For $k = -5$, $h = -3$

Hence, the coordinates of the third vertex C are
 $(-3, -5)$ or $(7, 5)$

509 (b)

Let $P(h, k)$ be the required point, then

$$4PA^2 = 9PB^2$$

$$\Rightarrow 4(h^2 + k^2) = 9(h - 4)^2 + 9(k + 3)^2$$

$$\Rightarrow 4h^2 + 4k^2 = 9(h^2 + 16 - 8h)$$

$$+ 9(k^2 + 9 + 6k)$$

$$\Rightarrow 5h^2 + 5k^2 - 72h + 54k + 225 = 0$$

\therefore Required locus of $P(h, k)$ is

$$5x^2 + 5y^2 - 72x + 54y + 225 = 0$$

510 (a)

The coordinates of the orthocentre O' and

circumcentre O are $(2, 1)$ and $(\frac{7}{2}, \frac{5}{2})$ respectively


We know that the centroid G divides OO' in the
 ratio $1 : 2$

So, the coordinates of G are

$$\left(\frac{1 \times 2 + 2 \times \frac{7}{2}}{1 + 2}, \frac{1 \times 1 + 2 \times \frac{5}{2}}{1 + 2} \right) \equiv (3, 2)$$

511 (a)

$$\text{In } \Delta POR, \frac{PR}{\sin 90^\circ} = \frac{h}{\sin 67\frac{1}{2}^\circ} \quad \dots(i)$$



And in ΔPQR

$$\frac{PR}{\sin 22\frac{1}{2}^\circ} = \frac{b}{\sin 45^\circ} \quad \dots(ii)$$


From Eqs. (i) and (ii)

$$\frac{\sin 45^\circ h}{\sin 67\frac{1}{2}^\circ b} = \frac{\sin 22\frac{1}{2}^\circ}{\sin 90^\circ} \Rightarrow \frac{h}{b} = \frac{\frac{1}{2} \sin 45^\circ}{\sin 45^\circ}$$

$$\Rightarrow 2h = b$$

512 (b)

Using, $\cos C = \frac{a^2 + b^2 - c^2}{2ab}$



$$\Rightarrow \frac{\sqrt{3}}{2} = \frac{(x^2 + x + 1)^2 + (x^2 - 1)^2 - (2x + 1)^2}{2(x^2 + x + 1)(x^2 - 1)}$$

$$\Rightarrow (x + 2)(x + 1)(x - 1)x + (x^2 - 1)^2 = \sqrt{3}(x^2 + x + 1)(x^2 - 1)$$

$$\Rightarrow x^2 + 2x + (x^2 - 1) = \sqrt{3}(x^2 + x + 1)$$

$$\Rightarrow 2(2 - \sqrt{3})x^2 + (2 - \sqrt{3})x - (\sqrt{3} + 1) = 0$$

$$\Rightarrow x = -(2 + \sqrt{3}) \text{ and } x = 1 + \sqrt{3}$$

But, $x = -(2 + \sqrt{3}) \Rightarrow c$ is negative
 $\therefore x = 1 + \sqrt{3}$ is the only solution

513 (d)

It is given that the distance between the points $P(a \cos 48^\circ, 0)$ and $Q(0, a \cos 12^\circ)$ is d
 i.e., $PQ = d$
 $\Rightarrow PQ^2 = d^2$
 $\Rightarrow a^2 \cos^2 48^\circ + a^2 \cos^2 12^\circ = d^2$
 $\Rightarrow a^2(1 + \cos 96^\circ) + a^2(1 + \cos 24^\circ) = 2d^2$
 $\Rightarrow 2a^2 + a^2(\cos 96^\circ + \cos 24^\circ) = 2d^2$
 $\Rightarrow 2a^2 + 2a^2 \cos 60^\circ \cos 36^\circ = 2d^2$
 $\Rightarrow 2a^2 + a^2 \left(\frac{\sqrt{5} + 1}{4}\right) = 2d^2 \Rightarrow d^2 - a^2 = \frac{a^2}{8}(\sqrt{5} + 1)$

514 (b)

$$\frac{ab \cos C - ac \cos B}{\frac{a^2 + b^2 - c^2}{2} - \frac{c^2 + a^2 - b^2}{2}}$$

$$= \frac{a^2 + b^2 - c^2 - c^2 - a^2 + b^2}{2}$$

$$= \frac{2b^2 - 2c^2}{2} = b^2 - c^2$$

515 (a)

For collinearity, $\begin{vmatrix} a_1 & b_1 & 1 \\ a_2 & b_2 & 1 \\ a_3 & b_3 & 1 \end{vmatrix} = 0$

For concurrency to lines $a_i x + b_i y + 1 = 0, i = 1, 2, 3$ we have

$$\begin{vmatrix} a_1 & b_1 & 1 \\ a_2 & b_2 & 1 \\ a_3 & b_3 & 1 \end{vmatrix} = 0, \text{ so lines are concurrent}$$

516 (c)

$$(b + c) \cos A + (c + a) \cos B + (a + b) \cos C$$

$$= b \cos A + c \cos A + c \cos B + a \cos B + a \cos C + b \cos C$$

$$= (b \cos A + a \cos B) + (c \cos A + a \cos C) + (c \cos B + b \cos C)$$

$$= c + b + a$$

517 (b)

By sine rule, $\frac{\sin A}{a} = \frac{\sin B}{b}$

Here, $\frac{\sin(\frac{\pi}{2} + B)}{5} = \frac{\sin B}{4}$ [by sine rule]

$$\Rightarrow \tan B = \frac{4}{5}$$

Also, $\angle A + \angle B + \angle C = \pi$

$$\Rightarrow \frac{\pi}{2} + 2\angle B + \angle C = \pi$$

$$\Rightarrow 2 \tan^{-1}\left(\frac{4}{5}\right) + \angle C = \frac{\pi}{2}$$

$$\Rightarrow \angle C = \frac{\pi}{2} - 2 \tan^{-1}\left(\frac{4}{5}\right)$$

$$\Rightarrow \angle C = \frac{\pi}{2} - \tan^{-1}\left(\frac{\frac{8}{5}}{1 - \frac{16}{25}}\right)$$

$$\Rightarrow = \frac{\pi}{2} - \tan^{-1}\left(\frac{40}{9}\right) = \cot^{-1}\left(\frac{40}{9}\right)$$

$$\Rightarrow \angle C = \tan^{-1}\left(\frac{9}{40}\right)$$

518 (b)

We have, $\frac{1}{2} a p_1 = \Delta, \frac{1}{2} b p_2 = \Delta, \frac{1}{2} c p_3 = \Delta$

Where a, b, c are the sides of a triangle

$$\Rightarrow p_1 = \frac{2\Delta}{a}, \quad p_2 = \frac{2\Delta}{b}, \quad p_3 = \frac{2\Delta}{c}$$

$$\therefore \frac{1}{p_1^2} + \frac{1}{p_2^2} + \frac{1}{p_3^2}$$

$$= \frac{a^2}{4\Delta^2} + \frac{b^2}{4\Delta^2} + \frac{c^2}{4\Delta^2} + \frac{a^2 + b^2 + c^2}{4\Delta^2}$$