

Chemical Bonding and Molecular Structure

1. A diatomic molecule has a dipole moment of 1.2 D, if the bond distance is 1 Å, what percentage of electronic charge exists on each atom?
2. The dipole moments of a diatomic molecule AB is 0.41D and bond distance is 2.82, calculate the % ionic character of AB.
3. The bond angle between two hybrid orbitals is 105°. Calculate the percentage of *s*-character of hybrid orbital.
4. In O_2^- , O_2 and O_2^{2-} molecular species. Find the total number of antibonding electrons.
5. There are two groups of compounds *A* and *B*. Group *A* contains three compounds Px_4 , Qy_3 , Rz_2 . Group *B* also contains three compounds Sx_4 , Ty_3 , Uz_2 . Hybridization of each central atom of group *A* compounds is same as that of iodine in $I\text{BrCl}^-$ while in group *B* compounds it is same as that of iodine in $I\text{BrCl}^+$. Substituents *X*, *Y* and *Z* exhibit covalency of one in ground state. Then find the value of x/y .

Where, *x* and *y* are total number of lone pairs present at central atoms of compounds of group *A* and *B* respectively.

6. Consider the following three compounds (i) AX_{2n}^{n-} , (ii) AX_{3n} and (iii) AX_{4n}^{n+} , where central atom *A* is 15th group element and their maximum covalency is 3*n*. If total number of proton in surrounding atom *X* is *n* and value of *n* is one, then calculate value of " $x^3 + y^2 + z$ ". (Where *x*, *y* and *z* are total number of lone pair at central atom in compound (i), (ii) and (iii) respectively.
7. Calculate the value of " $x + y - z$ " here *x*, *y* and *z* are total number of non-bonded electron pair(s), pie(π) bond(s) and sigma (σ) bonds in hydrogen phosphite ion respectively.
8. Consider the following compounds:


(i) IF_5	(ii) ClF_4^-
(iii) XeO_2F_2	(iv) NH_2^-
(v) BCl_3	(vi) BeCl_2
(vii) AsCl_4^+	(viii) $\text{B}(\text{OH})_3$
(ix) NO_2^-	(x) ClO_2^+

The value of " $x + y - z$ " is

where *x*, *y* and *z* are total number of compounds in given compounds in which central atom used their all three *p*-orbitals, only two *p*-orbitals and only one *p*-orbital in hybridisation respectively.

9. Total number of species which used all three *p*-orbitals in hybridisation of central atoms and should be non-polar

also are



10. Consider the following orbitals $3s, 2p_x, 4d_{xy}, 4d_z^2, 3d_{x^2-y^2}, 3p_y, 4s, 4p_z$ and find total number of orbital(s) having even number of nodal plane.
11. For the following molecules :
 $\text{PCl}_5, \text{BrF}_3, \text{ICl}_2^-, \text{XeF}_5^-, \text{NO}_3^-, \text{XeO}_2\text{F}_2, \text{PCl}_4^+, \text{CH}_3^+$

Calculate the value of $\frac{a+b}{c}$

a = Number of species having $sp^3 d$ -hybridisation

b = Number of species which are planar

c = Number of species which are non-planar

12. Find total number of orbital which can overlap colaterally, (if inter nuclear axis is *z*) $s, p_x, p_y, p_z, d_{xy}, d_{yz}, d_{xz}, d_z^2, d_{x^2-y^2}$
13. The total number of lone-pairs of electrons in melamine is
14. Among the triatomic molecules/ions, $\text{BeCl}_2, \text{N}_3^-, \text{N}_2\text{O}, \text{NO}_2^+, \text{O}_3, \text{SCl}_2, \text{ICl}_2^-, \text{I}_3^-$ and XeF_2 , the total number of linear molecule(s)/ion(s) where the hybridization of the central atom does not have contribution from the *d*-orbital(s) is

[Atomic number : S = 16, Cl = 17, I = 53 and Xe = 54]

15. The sum of the number of lone pairs of electrons on each central atom in the following species is
 $[\text{TeBr}_6]^{2-}, [\text{BrF}_2]^+, \text{SNF}_3$ and $[\text{XeF}_3]^-$
 (Atomic numbers: N = 7, F = 9, S = 16, Br = 35, Te = 52, Xe = 54)

SOLUTIONS

1. (25) $\mu = e \times d$

$$\therefore e = \frac{\mu}{d} = \frac{1.2 \text{ D}}{1.0 \text{ \AA}} = \frac{1.2 \times 10^{-18} \text{ esu cm}}{1.0 \times 10^{-8} \text{ cm}}$$

$$= 1.2 \times 10^{-10} \text{ esu}$$

Percentage of electronic charge

$$= \frac{1.2 \times 10^{-10} \text{ esu}}{4.8 \times 10^{-10} \text{ esu}} \times 100 = 25\%$$

2. (76.94) Dipole moment = electric charge \times bond length

$$= 4.8 \times 10^{-10} \times 2.82 \times 10^{-8} = 13.53 \text{ D}$$

Now % ionic character

$$= \frac{\text{Actual dipole moment of the bond}}{\text{Dipole moment of pure ionic compound}} \times 100$$

$$\text{then \% ionic character in AB} = \frac{10.41}{13.53} \times 100 = 76.94\%$$

3. (21.44) s -character \propto bond angle

For 25% s character (as in sp^3 hybrid orbital), bond angle is 109.5° , for 33.3% s character (as in sp^2 hybrid orbital), bond angle is 120° and for 50% s character (as in sp hybrid orbital), bond angle is 180° .

Similarly, when the bond angle decreases below 109.5° , the s -character will decrease accordingly.

Decreasing in angle = $120^\circ - 109.5^\circ = 10.5^\circ$

Decrease in s -character = $33.3 - 25 = 8.3$

Actual decrease in bond angle







$$= 109.5^\circ - 105^\circ = 4.5^\circ$$

Expected decrease in s -character

$$= \frac{8.3}{10.5} \times 4.5 = 3.56\%$$

Thus, the s -character should decrease by about 3.56% i.e., s -character = $25 - 3.56 = 21.44\%$

4. (21) Molecular orbital electronic configuration of these species are :



Hence number of antibonding electrons are 7, 6 and 8 respectively.

5. (2) $\text{I}_3^- \Rightarrow sp^3 d$

Group A]

Hybridisation $sp^3 d$

Structure : Linear

$R_2 +$

$x = \text{Total l.p.} = 6$

Group B]

Hybridisation sp

Structure : Linear

$x = \text{Total l.p.} = 0$

Only BeCl_2 , N_3^- , N_2^+ and NO_2 are linear with

sp hybridisation.

$y = \text{Total l.p.} = 3$

$x/y = 6/3 = 2$

Group A]

Hybridisation $sp^3 d$

Structure : Linear

$x =$

$y =$

$x/y =$

Group B]

Hybridisation sp

Structure : Linear

$x =$

$y =$


$x/y =$

6. (9) $n = 1$, then $X = \text{H}$; $A = \text{N}$



lone pair $x = 2, y = 1, z = 0$


$$x^3 + y^2 + z = (2)^3 + (1)^2 + (0) = 9$$



$$x = 7, y = 1, z = 5$$

$$7 + 1 - 5 = 3$$

8. (8) (i) $\text{IF}_5 (sp^3 d)$ (ii) $\text{ClH}_4^- (sp^3 d^2)$




$$x = (sp^3)2 + sp^3 d(1) + sp^3 d^2(2) = 5$$

$$y = 4, z = 1;$$


$$\therefore x + y - z = 5 + 4 - 1 = 8$$

9. (2) $\text{XeO}_4, \text{XeF}_7^+$




$sp^3 d$

Polar




sp^2

Polar



$sp^3 d^2$

Polar




sp^3

Polar




sp^3

Non-polar




sp^3

Polar



$sp^3 d^3$

Non-polar



$sp^3 d$


Polar

10. (2) $3s : 0$ Nodal plane
 $4d_z : 0$ Nodal plane
 $4s : 0$ Nodal plane
 $2p_x : 1$ Nodal plane
 $3d_{x^2-y^2} : 2$ Nodal plane
 $4p_z : 1$ Nodal plane
 $4d_{xy} : 2$ Nodal plane
 $3p_y : 1$ Nodal plane
 $4d_{xy}, 4d_{z^2}, 3d_{x^2-y^2}$, (Two)

11. (3) $\text{PCl}_5 \longrightarrow sp^3d$, non-planar
 $\text{BrF}_3 \longrightarrow sp^3d$, bent, T-shape, planar
 $\text{ICl}_2 \longrightarrow sp^3d$, linear, planar
 $\text{XeF}_5^- \longrightarrow sp^3d^3$, pentagonal planar
 $\text{NO}_3^- \longrightarrow sp^2$, planar
 $\text{XeO}_2\text{F}_2 \longrightarrow sp^3d$, see-saw, non-planar
 $\text{PCl}_4^+ \longrightarrow sp^3$, tetrahedral, non-planar
 $\text{CH}_3^+ \longrightarrow sp^2$, Trigonal planar
 $a = 4, b = 5, c = 3$
 so, $\frac{a+b}{c} = 3$




12. (6) $(p_x, p_y, d_{xy}, d_{yz}, d_{xz}, d_{x^2-y^2})$

13. (6) Structure of melamine is as follows,



Total no. of lone pairs of electron is '6'.

14. (4) $\text{Cl}-\text{Be}-\text{Cl}$
 Hybridization sp
 Structure : linear
- $\text{O}=\text{N}^+ \rightarrow \text{O}$
 Hybridisation sp
 Structure : Linear
- $\text{Cl}-\overset{\cdot\cdot}{\text{S}}-\text{Cl}$
 Hybridisation sp^3
 Structure : Angular
- $\text{N} \equiv \text{N} - \overset{\cdot\cdot}{\text{N}}^-$
 Hybridisation sp
 Structure : linear
- $\text{O}=\overset{\cdot\cdot}{\text{O}}-\overset{\cdot\cdot}{\text{O}}$
 Hybridisation sp^2
 Structure : Trigonal planar
- $[\text{Cl}-\overset{\cdot\cdot}{\text{I}}-\text{Cl}]$
 Hybridisation sp^3d
 Structure : linear



Only BeCl_2 , N_3^- , N_2O and NO_2 are linear with sp -hybridisation.

15. (6) Number of electron pairs around the central atom

$$= \frac{V + M \pm C}{2}$$

Compounds	No. of lone pairs on central atom
$[\text{TeBr}_6]^{2-}$	$\frac{(6+6+2)}{2} - 6 = 1$
$[\text{BrF}_2]^+$	$\frac{(7+2-1)}{2} - 2 = 2$
SNF_3	$\frac{6-1+3}{2} - 4 = 0$
$[\text{XeF}_3]^-$	$\frac{(8+3+1)}{2} - 3 = 3$

\therefore Sum of number of lone pairs = $1 + 2 + 0 + 3 = 6$