

## DAY EIGHTEEN

# Differential Equations

### Learning & Revision for the Day

- Solution of Differential Equation
- Solutions of Differential Equations of First Order and First Degree
- Linear Differential Equation
- Inspection Method

An equation involving independent variables, a dependent variable and the derivatives of dependent variable w.r.t. independent variables, is called **differential equation**.

A differential equation which contains only one independent variable, is called an ordinary differential equation.

### Order and Degree of a Differential Equation

- The **order** of a differential equation is the order of the highest derivative occurring in the differential equation.
- The **degree** of differential equation is the degree of the highest order derivative, when differential coefficients are made free from radicals and fractions.

### Solution of Differential Equation

A solution of the differential equation is a relation between the dependent and independent variables of the equation not containing the derivatives, but satisfying the given differential equation.

### General Solution

A general solution of a differential equation is a relation between the variables (not involving the derivatives) which contains the same number of the arbitrary constants as the order of the differential equation.

### Particular Solution

Particular solution of the differential equation is obtained from the general solution by assigning particular values to the arbitrary constant in the general solution.

## Solutions of Differential Equations of the First Order and First Degree

A differential equation of first order and first degree may be of the following types:

### Differential Equation with Variable Separable

'Variable separable method' is used to solve such an equation in which variable can be separated completely, e.g. the term  $f(x)$  with  $dx$  and the term  $g(y)$  with  $dy$

i.e.  $g(y)dy = f(x)dx$

### Reducible to Variable Separable Form

Differential equations of the form  $\frac{dy}{dx} = f(ax + by + c)$  can be reduced to variable separable form by the substitution  $ax + by + c = z$ .

$$\Rightarrow a + b \frac{dy}{dx} = \frac{dz}{dx} \Rightarrow \left(\frac{dz}{dx} - a\right) \frac{1}{b} = f(z) \Rightarrow \frac{dz}{dx} = a + bf(z)$$

### Homogeneous Differential Equation

A function  $f(x, y)$  is said to be a homogeneous function of degree  $n$  if  $f(\lambda x, \lambda y) = \lambda^n f(x, y)$  for some non-zero constant  $\lambda$ .

- Let  $f(x, y)$  and  $g(x, y)$  be two homogeneous functions of same degree, then a differential equation expressible in the form

$$\frac{dy}{dx} = \frac{f(x, y)}{g(x, y)}$$

is called a **homogeneous differential equation**.

- To solve a homogeneous differential equation of the type

$$\frac{dy}{dx} = \frac{f(x, y)}{g(x, y)} = h\left(\frac{y}{x}\right), \text{ we put } y = vx \text{ and to solve a}$$

homogeneous differential equation of type

$$\frac{dx}{dy} = \frac{f(x, y)}{g(x, y)} = h\left(\frac{x}{y}\right), \text{ we put } x = vy.$$

### Linear Differential Equation

A differential equation of the form  $\frac{dy}{dx} + Py = Q$ ,

where,  $P$  and  $Q$  are the functions of  $x$  (or constants), is called a linear differential equation.

To solve such an equation, first find integrating Factor,

$IF = e^{\int P dx}$ . Then, the solution of the differential equation is given by  $y(IF) = \int Q(IF) dx + C$ .

#### NOTE

- Sometimes given equation becomes a linear differential equation of the form  $\frac{dx}{dy} + Rx = S$ , where  $R$  and  $S$  are function of  $y$  (or constants).
- The integrating factor in this case is  $IF = e^{\int R dy}$  and the solution is given by  $x \cdot (IF) = \int (S \times IF) dy + C$ .

### Bernoulli's Differential Equation (Reducible to Linear Form)

Let the differential equation be of the form

$$\frac{dy}{dx} + Py = Qy^n \Rightarrow y^{-n} \frac{dy}{dx} + Py^{-n+1} = Q$$

Here, put  $y^{-n+1} = z \Rightarrow (-n+1)y^{-n} \frac{dy}{dx} = \frac{dz}{dx}$

$$\Rightarrow \frac{dz}{dx} + (1-n)Pz = (1-n)Q$$

which is a linear differential equation in  $z$ .

The solution of this equation is given by

$$ze^{\int (1-n)P dx} = \int [(1-n) \cdot Q \cdot e^{\int (1-n)P dx}] dx + C$$

### Inspection Method

If we can write the differential equation in the form  $f\{f_1(x, y)\} d\{f_1(x, y)\} + \phi\{f_2(x, y)\} d\{f_2(x, y)\} + \dots = 0$ , then each term can be easily integrated separately.

For this use the following results.

- (i)  $d(x \pm y) = dx \pm dy$       (ii)  $d(xy) = x dy + y dx$
- (iii)  $d\left(\frac{x}{y}\right) = \frac{y dx - x dy}{y^2}$
- (iv)  $d\left(\tan^{-1} \frac{x}{y}\right) = \frac{y dx - x dy}{x^2 + y^2}$
- (v)  $d[\log(xy)] = \frac{x dy + y dx}{xy}$
- (vi)  $d\left[\log\left(\frac{x}{y}\right)\right] = \frac{y dx - x dy}{xy}$
- (vii)  $d\left[\frac{1}{2} \log(x^2 + y^2)\right] = \frac{x dx + y dy}{x^2 + y^2}$
- (viii)  $d\left(-\frac{1}{xy}\right) = \frac{x dy + y dx}{x^2 y^2}$
- (ix)  $d\left(\frac{e^x}{y}\right) = \frac{y e^x dx - e^x dy}{y^2}$
- (x)  $d(\sqrt{x^2 + y^2}) = \frac{x dx + y dy}{\sqrt{x^2 + y^2}}$
- (xi)  $d\left(\frac{1}{2} \log \frac{x+y}{x-y}\right) = \frac{x dy - y dx}{x^2 - y^2}$

DAY PRACTICE SESSION 1

## FOUNDATION QUESTIONS EXERCISE

- 1** The order of the differential equation whose general solution is given by

$$y = c_1 e^{2x+c_2} + c_3 e^x + c_4 \sin(x+c_5), \text{ is}$$

- (a) 5      (b) 4      (c) 3      (d) 2

- 2** The degree of the differential equation

$$\frac{d^2 y}{dx^2} + 3 \left[ \frac{dy}{dx} \right]^2 = x^2 \log \left[ \frac{d^2 y}{dx^2} \right], \text{ is}$$

- (a) 1      (b) 2  
(c) 3      (d) None of these

- 3** Order and degree of the differential equation, representing the family of curves  $y^2 = 2c(x + \sqrt{c})$ , where  $c$  is a positive parameter, are respectively equal to

- (a) 1, 3      (b) 2, 3  
(c) 2, 4      (d) 1, 2

- 4** The differential equation of all circles in the first quadrant which touch the coordinate axes is of order

- (a) 1      (b) 2  
(c) 3      (d) None of these

- 5** The differential equation of the family of curves  $v = \frac{A}{r} + B$ ,

where  $A$  and  $B$  are arbitrary constants, is

- (a)  $\frac{d^2 v}{dr^2} + \frac{1}{r} \frac{dv}{dr} = 0$       (b)  $\frac{d^2 v}{dr^2} - \frac{2}{r} \frac{dv}{dr} = 0$   
(c)  $\frac{d^2 v}{dr^2} + \frac{2}{r} \frac{dv}{dr} = 0$       (d) None of these

- 6** The solution of the differential equation

$$x dx + y dy + \frac{x dy - y dx}{x^2 + y^2} = 0, \text{ is}$$

- (a)  $y = x \tan \left( \frac{x^2 + y^2 + C}{2} \right)$   
(b)  $x = y \tan \left( \frac{x^2 + y^2 + C}{2} \right)$   
(c)  $y = x \tan \left( \frac{C - x^2 - y^2}{2} \right)$   
(d) None of the above

- 7** The differential equation of the family of parabolas with focus at the origin and the  $X$ -axis as axis, is

- (a)  $y \left( \frac{dy}{dx} \right)^2 + 4x \frac{dy}{dx} = 4y$       (b)  $y \left( \frac{dy}{dx} \right)^2 = 2x \frac{dy}{dx} - y$   
(c)  $y \left( \frac{dy}{dx} \right)^2 + y = 2xy \frac{dy}{dx}$       (d)  $y \left( \frac{dy}{dx} \right)^2 + 2xy \frac{dy}{dx} + y = 0$

- 8** The differential equation whose solution is  $x^2 + y^2 + 2ax + 2by + c = 0$ , where  $a, b, c$  are arbitrary constants, is

- (a)  $3y_1 y_2 - (1 + y_1^2) y_3 = 0$       (b)  $3y_1^2 y_2 - (1 + y_1^2) y_3 = 0$   
(c)  $3y_1 y_2^2 + (1 + y_1^2) y_3 = 0$       (d)  $3y_1 y_2^2 - (1 + y_1^2) y_3 = 0$

- 9** If  $x dy = y(dx + y dy)$ ,  $y(1) = 1$  and  $y(x) > 0$ .

Then,  $y(-3)$  is equal to

- (a) 3      (b) 2  
(c) 1      (d) 0

- 10** If  $\frac{dy}{dx} = y + 3 > 0$  and  $y(0) = 2$ , then  $y(\log 2)$  is equal to

- (a) 5      (b) 13      (c) -2      (d) 7

- 11** The differential equation  $\frac{dy}{dx} = \frac{\sqrt{1-y^2}}{y}$  determines family

of circles with,

- (a) variable radii and a fixed centre at  $(0, 1)$   
(b) variable radii and a fixed centre at  $(0, -1)$   
(c) fixed radius 1 and variable centre along the  $X$ -axis  
(d) fixed radius 1 and variable centre along the  $Y$ -axis

- 12** If the solution of the differential equation  $\frac{dy}{dx} = \frac{ax+3}{2y+f}$

represents a circle, then the value of  $a$  is

- (a) 2      (b) -2      (c) 3      (d) -4

- 13** Solution of the differential equation  $y - x \frac{dy}{dx} = y^2 + \frac{dy}{dx}$  is

- (a)  $Cy = (1-x)(1-y)$       (b)  $Cx = (1+x)(1-y)$   
(c)  $Cy = (1+x)(1-y)$       (d)  $Cx = (1-x)(1+y)$

- 14** Solution of the equation  $\ln \left( \frac{dy}{dx} \right) = ax + by$ , is

- (a)  $ae^{-by} + be^{ax} + C = 0$   
(b)  $ae^{by} + be^{ax} + C = 0$   
(c)  $ae^{by} + be^{-ax} + C = 0$   
(d) None of the above

- 15** The solution of the differential equation

$$\frac{dy}{dx} = e^{x-y} + x^2 e^{-y} \text{ is}$$

- (a)  $e^y = e^x + \frac{x^3}{3} + C$       (b)  $e^y = e^x + 2x + C$   
(c)  $e^y = e^x + x^3 + C$       (d)  $y = e^x + C$

- 16** If  $(2 + \sin x) \frac{dy}{dx} + (y+1) \cos x = 0$  and  $y(0) = 1$ , then  $y\left(\frac{\pi}{2}\right)$

is equal to

→ JEE Mains 2017

- (a)  $-\frac{1}{3}$       (b)  $\frac{4}{3}$       (c)  $\frac{1}{3}$       (d)  $-2/3$

- 17** If a curve passes through the point  $\left(2, \frac{7}{2}\right)$  and has slope

$\left(1 - \frac{1}{x^2}\right)$  at any point  $(x, y)$  on it, then the ordinate of the

point on the curve, whose abscissa is  $-2$ , is

→ JEE Mains 2013

- (a)  $-\frac{3}{2}$       (b)  $\frac{3}{2}$       (c)  $\frac{5}{2}$       (d)  $-\frac{5}{2}$

18 Solution of the equation  $(x+y)^2 \frac{dy}{dx} = 4$ ,  $y(0) = 0$  is

- (a)  $y = 2 \tan^{-1}\left(\frac{x+y}{2}\right)$       (b)  $y = 4 \tan^{-1}\left(\frac{x+y}{4}\right)$   
 (c)  $y = 4 \tan^{-1}\left(\frac{x+y}{2}\right)$       (d) None of these

19 The solution of  $\frac{dy}{dx} + 1 = e^{x+y}$  is

- (a)  $e^{-(x+y)} + x + C = 0$       (b)  $e^{-(x+y)} - x + C = 0$   
 (c)  $e^{x+y} + x + C = 0$       (d)  $e^{x+y} - x + C = 0$

20 Solution of the equation  $xdy = (y + xf(y/x))/f'(y/x) dx$  is

- (a)  $|f(y/x)| = C|x|$ ,  $C \in \mathbb{R}$       (b)  $|f(y/x)| = |x| + C$ ,  $C > 0$   
 (c)  $|f(y/x)| = C|x|$ ,  $C > 0$       (d) None of these

21 If  $x \frac{dy}{dx} = y(\log y - \log x + 1)$ , then the solution of the equation is

- (a)  $\log\left(\frac{x}{y}\right) = Cy$       (b)  $\log\left(\frac{y}{x}\right) = Cx$   
 (c)  $x \log\left(\frac{y}{x}\right) = Cy$       (d)  $y \log\left(\frac{x}{y}\right) = Cx$

22 The general solution of  $y^2 dx + (x^2 - xy + y^2) dy = 0$ , is

- (a)  $\tan^{-1}\left(\frac{x}{y}\right) + \log y + C = 0$   
 (b)  $2 \tan^{-1}\left(\frac{x}{y}\right) + \log x + C = 0$   
 (c)  $\log(y + \sqrt{x^2 + y^2}) + \log y + C = 0$   
 (d)  $\sin^{-1}\left(\frac{x}{y}\right) + \log y + C = 0$

23 Solution of the differential equation  $x^2 dy + y(x+y) dx = 0$  is

- (a)  $y + 2x = c^2 x^2 y$       (b)  $y - 2x = c^2 x^2 / y$   
 (c)  $y + 2x = c^2 x^2 / y$       (d) None of these

24 If  $(x^2 + y^2) dy = xy dx$  and  $y(x_0) = e$ ,  $y(1) = 1$ , then  $x_0$  is

- (a)  $e\sqrt{3}$       (b)  $\sqrt{2e^2 - 1}/\sqrt{2}$   
 (c)  $\sqrt{e^2 - 1}/\sqrt{2}$       (d)  $\sqrt{e^2 + 1}/\sqrt{2}$

25 If a curve  $y = f(x)$  passes through the point  $(1, -1)$  and satisfies the differential equation  $y(1+xy) dx = x dy$ , then  $f\left(-\frac{1}{2}\right)$  is equal to

→ JEE Mains 2016

- (a)  $-\frac{2}{5}$       (b)  $-\frac{4}{5}$       (c)  $\frac{2}{5}$       (d)  $\frac{4}{5}$

26 The equation of the curve passing through the origin and satisfying the differential equation

$$(1+x^2) \frac{dy}{dx} + 2xy = 4x^2 \text{ is}$$

→ JEE Mains 2013

- (a)  $(1+x^2)y = x^3$       (b)  $3(1+x^2)y = 2x^3$   
 (c)  $(1+x^2)y = 3x^3$       (d)  $3(1+x^2)y = 4x^3$

27 The solution of the differential equation

$$\frac{dy}{dx} + \frac{3x^2}{1+x^3} y = \frac{\sin^2 x}{1+x^3}, \text{ is}$$

- (a)  $y(1+x^3) = x + \frac{1}{2} \sin 2x + C$   
 (b)  $y(1+x^3) = Cx + \frac{1}{2} \sin 2x$   
 (c)  $y(1+x^3) = Cx - \frac{1}{2} \sin 2x$   
 (d)  $y(1+x^3) = \frac{x}{2} - \frac{1}{4} \sin 2x + C$

28 If  $y(x)$  satisfies the differential equation  $y' - y \tan x = 2x \sec x$  and  $y(0) = 0$ , then

→ IIT 2012

- (a)  $y\left(\frac{\pi}{4}\right) = \frac{\pi^2}{8\sqrt{2}}$       (b)  $y\left(\frac{\pi}{4}\right) = \frac{\pi^2}{18}$   
 (c)  $y\left(\frac{\pi}{3}\right) = \frac{\pi^2}{9}$       (d)  $y\left(\frac{\pi}{3}\right) = \frac{4\pi}{3} + \frac{\pi^2}{3\sqrt{3}}$

29 The solution of the equation

$$\frac{dy}{dx} + y \tan x = x^m \cos x, \text{ is}$$

- (a)  $(m+1)y = x^{m+1} \cos x + C(m+1) \cos x$   
 (b)  $my = (x^m + C) \cos x$   
 (c)  $y = (x^{m+1} + C) \cos x$   
 (d) None of the above

30 Let the population of rabbits surviving at a time  $t$  be governed by the differential equation  $\frac{dP(t)}{dt} = \frac{1}{2} P(t) - 200$ .

If  $P(0) = 100$ , then  $P(t)$  is equal to

→ JEE Mains 2014

- (a)  $400 - 300e^{\frac{t}{2}}$       (b)  $300 - 200e^{-\frac{t}{2}}$   
 (c)  $600 - 500e^{\frac{t}{2}}$       (d)  $400 - 300e^{-\frac{t}{2}}$

31 The solution of differential equation

$$(xy^5 + 2y) dx - x dy = 0, \text{ is}$$

- (a)  $9x^8 + 4x^9 y^4 = 9y^4 C$       (b)  $9x^8 - 4x^9 y^4 - 9y^4 C = 0$   
 (c)  $x^8(9 + 4y^4) = 10y^4 C$       (d) None of these

32 Let  $Y = y(x)$  be the solution of the differential equation

$$\sin x \frac{dy}{dx} + y \cos x = 4x, x \in (0, \pi). \text{ If } y\left(\frac{\pi}{2}\right) = 0, \text{ then}$$

$y(\pi/6)$  is equal to

→ JEE Mains 2018

- (a)  $-\frac{8}{9\sqrt{3}} \pi^2$       (b)  $-\frac{8}{9} \pi^2$       (c)  $-\frac{4}{9} \pi^2$       (d)  $\frac{4}{9\sqrt{3}} \pi^2$

33 **Statement I** The slope of the tangent at any point  $P$  on a parabola, whose axis is the axis of  $x$  and vertex is at the origin, is inversely proportional to the ordinate of the point  $P$ .

**Statement II** The system of parabolas  $y^2 = 4ax$  satisfies a differential equation of degree 1 and order 1.

→ JEE Mains 2013

- (a) Statement I is true, Statement II is true; Statement II is a correct explanation for Statement I

- (b) Statement I is true; Statement II is true; Statement II is not a correct explanation for Statement I  
 (c) Statement I is true; Statement II is false  
 (d) Statement I is false; Statement II is true

**34 Statement I** The elimination of four arbitrary constants in  $y = (c_1 + c_2 + c_3 e^{c_4})x$  results into a differential equation of the first order  $x \frac{dy}{dx} = y$ .

**Statement II** Elimination of  $n$  independent arbitrary constants results in a differential equation of the  $n$ th order.

- (a) Statement I is true; Statement II is true; Statement II is a correct explanation for Statement I  
 (b) Statement I is true; Statement II is true; Statement II is not a correct explanation for Statement I  
 (c) Statement I is true; Statement II is false  
 (d) Statement I is false; Statement II is true

## DAY PRACTICE SESSION 2

# PROGRESSIVE QUESTIONS EXERCISE

**1** The differential equation which represents the family of curves  $y = c_1 e^{c_2 x}$  where  $c_1$  and  $c_2$  are arbitrary constant, is

- (a)  $y'' = y'y$  (b)  $yy'' = y'$   
 (c)  $yy'' = (y')^2$  (d)  $y' = y^2$

**2** The population  $p(t)$  at time  $t$  of a certain mouse species satisfies the differential equation  $\frac{dp(t)}{dt} = 0.5(t) - 450$ . If

$p(0) = 850$ , then the time at which the population becomes zero is

- (a)  $2 \log 18$  (b)  $\log 9$   
 (c)  $\frac{1}{2} \log 18$  (d)  $\log 18$

**3** A curve passes through the point  $\left(1, \frac{\pi}{6}\right)$ . Let the slope of

the curve at each point  $(x, y)$  be  $\frac{y}{x} + \sec\left(\frac{y}{x}\right)$ ,  $x > 0$ .

Then the equation of the curve is → JEE Advanced 2013

- (a)  $\sin\left(\frac{y}{x}\right) = \log x + \frac{1}{2}$   
 (b)  $\operatorname{cosec}\left(\frac{y}{x}\right) = \log x + 2$   
 (c)  $\sec\left(\frac{2y}{x}\right) = \log x + 2$   
 (d)  $\cos\left(\frac{2y}{x}\right) = \log x + \frac{1}{2}$

**4** Let  $y'(x) + y(x)g'(x) = g(x)g'(x)$ ,  $y(0) = 0$ ,  $x \in R$ , where  $f'(x)$  denotes  $\frac{df(x)}{dx}$  and  $g(x)$  is a given non-constant

differentiable function on  $R$  with  $g(0) = g(2) = 0$ . Then the value of  $y(2)$  is

- (a) 2 (b) 4  
 (c) 5 (d) 0

**5** Consider the differential equation

$y^2 dx + \left(x - \frac{1}{y}\right) dy = 0$ . If  $y(1) = 1$ , then  $x$  is given by

- (a)  $1 - \frac{1}{y} + \frac{e^y}{e}$  (b)  $4 - \frac{2}{y} - \frac{1}{e}$   
 (c)  $3 - \frac{1}{y} + \frac{1}{e}$  (d)  $1 + \frac{1}{y} - \frac{1}{e}$

**6** The equation of the curve  $y = f(x)$  passing through the origin which satisfies the differential equation

$\frac{dy}{dx} = \sin(10x + 6y)$  is

- (a)  $y = \frac{1}{3} \left\{ \tan^{-1} \frac{5 \tan 4x}{4 - 3 \tan 4x} - 5x \right\}$   
 (b)  $y = \frac{1}{3} \left\{ \tan^{-1} \frac{5 \tan 4x}{4 - 3 \tan 4x} + 5x \right\}$   
 (c)  $y = \frac{1}{3} \left\{ \tan^{-1} \frac{5 \tan 4x}{4 + 3 \tan 4x} - 5x \right\}$   
 (d) None of the above

**7** Let  $I$  be the purchase value of an equipment and  $V(t)$  be the value after it has been used for  $t$  yr. The value  $V(t)$  depreciates at a rate given by differential equation

$\frac{dV(t)}{dt} = -k(T - t)$ , where  $k > 0$  is a constant and  $T$  is the

total life in years of the equipment. Then, the scrap value  $V(T)$  of the equipment is

- (a)  $I - \frac{kT^2}{2}$  (b)  $I - \frac{k(T - t)^2}{2}$   
 (c)  $e^{-kT}$  (d)  $T^2 - \frac{1}{k}$

**8** The degree of the differential equation satisfying the

relation  $\sqrt{1+x^2} + \sqrt{1+y^2} = \lambda (x\sqrt{1+y^2} - y\sqrt{1+x^2})$ , is

- (a) 1 (b) 2  
 (c) 3 (d) None of these

**9** If length of tangent at any point on the curve  $y = f(x)$  intercepted between the point of contact and  $X$ -axis is of length 1, the equation of the curve is

- (a)  $\sqrt{1-y^2} + \ln \left| \frac{1-\sqrt{1-y^2}}{y} \right| = \pm x + C$   
 (b)  $\sqrt{1-y^2} - \ln \left| \frac{1-\sqrt{1-y^2}}{y} \right| = \pm x + C$   
 (c)  $\sqrt{1-y^2} + \ln \left| \frac{1+\sqrt{1-y^2}}{y} \right| = \pm x + C$   
 (d) None of the above

**10** The solution of differential equation

$$\cos x \, dy = y(\sin x - y) \, dx, \text{ where, } 0 < x < \frac{\pi}{2}, \text{ is}$$

- (a)  $\sec x = (\tan x + C)y$       (b)  $y \sec x = \tan x + C$   
 (c)  $y \tan x = \sec x + C$       (d)  $\tan x = (\sec x + C)y$

**11** The curves satisfying the differential equation

$$(1-x^2)y' + xy = ax \text{ are}$$

- (a) ellipse and parabola      (b) ellipse and circles  
 (c) ellipse and hyperbola      (d) None of these

**12** A curve passes through (2, 0) and the slope of tangent at a point  $P(x, y)$  is equal to  $\frac{(x+1)^2 + y - 3}{(x+1)}$ . Then equation of the curve is → IIT 2004

- (a)  $y = x^2 + 2x$       (b)  $y = x^2 - 2x$   
 (c)  $y = 2x^2 - x$       (d) None of these

**13** The solution of the differential equation

$$\frac{x + \frac{x^3}{3!} + \frac{x^5}{5!} + \dots}{1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \dots} = \frac{dx - dy}{dx + dy} \text{ is}$$

- (a)  $2ye^{2x} = Ce^{2x} + 1$       (b)  $2ye^{2x} = Ce^{2x} - 1$   
 (c)  $ye^{2x} = Ce^{2x} + 2$       (d) None of these

**14** Let  $f$  be a real-valued differentiable function on  $R$  (the set of all real numbers) such that  $f(1) = 1$ . If the y-intercept of the tangent at any point  $P(x, y)$  on the curve  $y = f(x)$  is equal to the cube of the abscissa of  $P$ , then the value of  $f(-3)$  is equal to

- (a) 3      (b) 6      (c) 9      (d) 1

**15** Let  $f(x)$  be differentiable in the interval  $(0, \pi)$  such that  $f(1) = 1$  and  $\lim_{t \rightarrow x} \frac{t^2 f(x) - x^2 f(t)}{t - x} = 1$  for each  $x > 0$ . Then  $f(x)$

is

- (a)  $\frac{1}{3x} + \frac{2x^2}{3}$       (b)  $-\frac{1}{3x} + \frac{4x^2}{3}$   
 (c)  $-\frac{1}{2x} + \frac{2}{x^2}$       (d)  $\frac{1}{x}$

**16** Let a solution  $y = y(x)$  of the differential equation

$$x\sqrt{x^2 - 1} \, dy - y\sqrt{y^2 - 1} \, dx = 0 \text{ satisfy } y(2) = \frac{2}{\sqrt{3}}.$$

**Statement I**  $y(x) = \sec \left( \sec^{-1} x - \frac{\pi}{6} \right)$ .

**Statement II**  $y(x)$  is given by  $\frac{1}{y} = \frac{2\sqrt{3}}{x} - \sqrt{1 - \frac{1}{x^2}}$ .

- (a) Statement I is true, Statement II is true; Statement II is a correct explanation for Statement I  
 (b) Statement I is true, Statement II is true; Statement II is not a correct explanation for Statement I  
 (c) Statement I is true; Statement II is false  
 (d) Statement I is false; Statement II is true

## ANSWERS

### SESSION 1

- |         |         |         |         |         |         |         |         |         |         |
|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| 1. (b)  | 2. (d)  | 3. (a)  | 4. (a)  | 5. (c)  | 6. (c)  | 7. (b)  | 8. (d)  | 9. (a)  | 10. (d) |
| 11. (c) | 12. (b) | 13. (c) | 14. (a) | 15. (a) | 16. (c) | 17. (a) | 18. (a) | 19. (a) | 20. (c) |
| 21. (b) | 22. (a) | 23. (a) | 24. (a) | 25. (d) | 26. (d) | 27. (d) | 28. (a) | 29. (a) | 30. (a) |
| 31. (a) | 32. (b) | 33. (b) | 34. (a) |         |         |         |         |         |         |

### SESSION 2

- |         |         |         |         |         |         |        |        |        |         |
|---------|---------|---------|---------|---------|---------|--------|--------|--------|---------|
| 1. (c)  | 2. (a)  | 3. (a)  | 4. (d)  | 5. (d)  | 6. (a)  | 7. (a) | 8. (a) | 9. (a) | 10. (a) |
| 11. (c) | 12. (b) | 13. (b) | 14. (c) | 15. (a) | 16. (c) |        |        |        |         |

# Hints and Explanations

## SESSION 1

1 Given,

$$\begin{aligned} y &= c_1 \cdot e^{2x+c_2} + c_3 e^x + c_4 \sin(x+c_5) \\ &= c_1 \cdot e^{c_2} e^{2x} + c_3 e^x \\ &\quad + c_4 (\sin x \cos c_5 + \cos x \sin c_5) \\ &= Ae^{2x} + c_3 e^x + B \sin x + D \cos x \end{aligned}$$

Here,  $A = c_1 e^{c_2}$ ,  $B = c_4 \cos c_5$  and

$D = c_4 \sin c_5$

Since, equation consists four arbitrary constants.

So, the order of differential equation is 4.

2 Since, the equation is not a polynomial in all differential coefficients, so its degree is not defined.

3 Given,  $y^2 = 2c(x + \sqrt{c})$  ... (i)

On differentiating both side w.r.t.  $x$ , we get

$$2y \frac{dy}{dx} = 2c \Rightarrow c = y \frac{dy}{dx}$$

On putting in Eq. (i),

$$\begin{aligned} y^2 &= 2xy \frac{dy}{dx} + 2y^{3/2} \left( \frac{dy}{dx} \right)^{3/2} \\ \Rightarrow 8y^3 \left( \frac{dy}{dx} \right)^3 &= \left( y^2 - 2xy \frac{dy}{dx} \right)^2 \end{aligned}$$

Which is the differential equation of order one and degree 3.

4 Clearly, the equation of family of circle which touch both the axes is  $(x-a)^2 + (y-a)^2 = a^2$ , where  $a$  is a parameter.

Since, there is only one parameter, therefore order of differential equation representing this family is 1.

5 We have,  $v = \frac{A}{r} + B$ , where  $A$  and  $B$  are parameters

On differentiating twice w.r.t.  $r$ , we get

$$\Rightarrow \frac{dv}{dr} = \frac{-A}{r^2} \quad \dots (i)$$

$$\text{and } \frac{d^2v}{dr^2} = \frac{2A}{r^3} \quad \dots (ii)$$

Now, on substituting the value of  $A$  from Eq. (i) in Eq. (ii), we get

$$\frac{d^2v}{dr^2} = \frac{2}{r^3} \left( -r^2 \frac{dv}{dr} \right) = \frac{-2}{r} \frac{dv}{dr}$$

$\Rightarrow \frac{d^2v}{dr^2} + \frac{2}{r} \frac{dv}{dr} = 0$ , which is the required differential equation.

6 We have,  $x dx + y dy + \frac{xdy - ydx}{x^2 + y^2} = 0$

$$\Rightarrow \frac{1}{2} d(x^2 + y^2) + d \left( \tan^{-1} \frac{y}{x} \right) = 0$$

On integrating, we get

$$\begin{aligned} \frac{1}{2} (x^2 + y^2) + \tan^{-1} \left( \frac{y}{x} \right) &= \frac{C}{2} \\ \Rightarrow \frac{C - x^2 - y^2}{2} &= \tan^{-1} \left( \frac{y}{x} \right) \\ \therefore y &= x \tan \left( \frac{C - x^2 - y^2}{2} \right) \end{aligned}$$

7 Equation of family of parabolas with focus at  $(0, 0)$  and axis as  $X$ -axis is  $y^2 = 4a(x-a)$  ... (i)

On differentiating Eq. (i) w.r.t.  $x$ , we get

$$\begin{aligned} 2yy_1 &= 4a \\ \therefore y^2 &= 2yy_1 \left( x - \frac{yy_1}{2} \right) \end{aligned}$$

$$\begin{aligned} \Rightarrow y &= 2xy_1 - y y_1^2 \\ \Rightarrow y y_1^2 &= 2xy_1 - y \end{aligned}$$

8  $x^2 + y^2 + 2ax + 2by + c = 0$  ... (i)

On differentiating Eq. (i) three times, we get

$$\begin{aligned} 2x + 2yy_1 + 2a + 2by_1 &= 0 \\ \Rightarrow x + yy_1 + a + by_1 &= 0 \quad \dots (ii) \\ 1 + y_1^2 + yy_2 + by_2 &= 0 \quad \dots (iii) \\ 3y_1 y_2 + yy_3 + by_3 &= 0 \quad \dots (iv) \end{aligned}$$

On eliminating  $b$  from Eqs. (iii) and (iv), we get

$$3y_1 y_2^2 - y_3 - y_1^2 y_3 = 0$$

9 We have,  $x dy = y(dx + y dy)$ ,  $y > 0$

$$\therefore \frac{x dy - y dx}{y^2} = dy$$

$$\Rightarrow \frac{x}{y} = -y + C \quad [\text{integrating}]$$

Now,  $y(1) = 1 \Rightarrow C = 2$

$$\therefore \frac{x}{y} + y = 2$$

For  $x = -3$ ,  $-3 + y^2 = 2y$

$$\Rightarrow y^2 - 2y - 3 = 0$$

$$\Rightarrow (y+1)(y-3) = 0$$

$$\therefore y = 3 \quad [ \because y > 0 ]$$

10 Here,  $\frac{dy}{dx} = y + 3 > 0$  and  $y(0) = 2$

$$\Rightarrow \int \frac{dy}{y+3} = \int dx$$

$$\Rightarrow \log_e |y+3| = x + C$$

But  $y(0) = 2$

$$\therefore \log_e |2+3| = 0 + C$$

$$\Rightarrow C = \log_e 5$$

$$\Rightarrow \log_e |y+3| = x + \log_e 5$$

When  $x = \log_e 2$ , then

$$\begin{aligned} \Rightarrow \log_e |y+3| &= \log_e 2 + \log_e 5 \\ &= \log_e 10 \end{aligned}$$

$$\therefore y + 3 = 10 \Rightarrow y = 7$$

11 Clearly,  $\int \frac{y}{\sqrt{1-y^2}} dy = \int dx$

$$\Rightarrow -\sqrt{1-y^2} = x + C$$

$$\Rightarrow (x+C)^2 + y^2 = 1$$

Hence, the centre is  $(-C, 0)$  and radius is 1.

12 We have,  $\frac{dy}{dx} = \frac{ax+3}{2y+f}$

$$\Rightarrow (ax+3)dx = (2y+f)dy$$

On integrating, we obtain

$$a \cdot \frac{x^2}{2} + 3x = y^2 + fy + C$$

$$\Rightarrow -\frac{a}{2}x^2 + y^2 - 3x + fy + C = 0$$

This will represent a circle, if  $-\frac{a}{2} = 1$

[ $\because$  coefficient of  $x^2 =$  coefficient of  $y^2$ ]

$$\Rightarrow a = -2$$

13  $(x+1)\frac{dy}{dx} = y - y^2$

$$\Rightarrow \frac{dy}{y(1-y)} = \frac{dx}{x+1}$$

$$\Rightarrow \left( \frac{1}{y} + \frac{1}{1-y} \right) dy = \frac{dx}{x+1}$$

$$\begin{aligned} \Rightarrow \log y - \log(1-y) &= \log(x+1) + \log C \\ &= \log(x+1) + \log C \end{aligned}$$

$$\Rightarrow \frac{y}{1-y} = C(x+1)$$

$$\Rightarrow (x+1)(1-y) = Cy.$$

14 We have,  $\frac{dy}{dx} = e^{ax} \cdot e^{by}$

$$\Rightarrow e^{-by} dy = e^{ax} dx$$

$$\Rightarrow \frac{-1}{b} e^{-by} = \frac{1}{a} e^{ax} + C$$

$$\Rightarrow be^{ax} + ae^{-by} + C' = 0 \quad [C' = abC]$$

15 We have,

$$\frac{dy}{dx} = e^{x-y} + x^2 e^{-y} = e^{-y}(x^2 + e^x)$$

On separating the variables, we get

$$e^y dy = (x^2 + e^x) dx$$

On integrating both sides, we get

$$e^y = \frac{x^3}{3} + e^x + C$$

16 We have,

$$(2 + \sin x) \frac{dy}{dx} + (y+1) \cos x = 0$$

$$\Rightarrow \frac{dy}{y+1} = -\frac{\cos x}{2 + \sin x} dx$$

$$\Rightarrow \int \frac{dy}{y+1} = \int -\frac{\cos x}{2 + \sin x} dx$$

$$\Rightarrow \ln(y+1) = -\ln(2 + \sin x) + \ln C$$

$$\Rightarrow (y+1)(2 + \sin x) = C$$

Now,  $y(0) = 1$



$$\begin{aligned} \Rightarrow (2)(2+0) &= C \\ \Rightarrow C &= 4 \\ \text{Thus, } (y+1)(2+\sin x) &= 4 \\ \text{Now, at } x = \frac{\pi}{2}, (y+1)\left(2+\sin\frac{\pi}{2}\right) &= 4 \\ (y+1)(2+1) &= 4 \\ y = \frac{4}{3} - 1 &= \frac{1}{3} \end{aligned}$$

**17** Given,  $\frac{dy}{dx} = 1 - \frac{1}{x^2}$

$$\Rightarrow dy = \left(1 - \frac{1}{x^2}\right) dx$$

$$\Rightarrow y = x + \frac{1}{x} + C \quad [\text{integrating}]$$

Since, the curve passing through the point  $\left(2, \frac{7}{2}\right)$ .

i.e.  $\frac{7}{2} = 2 + \frac{1}{2} + C \Rightarrow C = 1$

$$\therefore y = x + \frac{1}{x} + 1 \quad \dots(i)$$

Now, at  $x = -2$

Ordinate,  $y = -2 - \frac{1}{2} + 1 = -3/2$

**18** We have,  $(x+y)^2 \frac{dy}{dx} = 4$

On putting  $x+y = v$  and  $1 + \frac{dy}{dx} = \frac{dv}{dx}$ ,

we get

$$\Rightarrow v^2 \left(\frac{dv}{dx} - 1\right) = 4$$

$$\Rightarrow v^2 \frac{dv}{dx} = v^2 + 4.$$

$$\therefore \frac{v^2 + 4 - 4}{v^2 + 4} dv = dx$$

$$\Rightarrow v - 2 \tan^{-1}(v/2) = x + C$$

$$\Rightarrow x + y - 2 \tan^{-1}\left(\frac{x+y}{2}\right) = x + C$$

Now,  $y(0) = 0 \Rightarrow C = 0$ .

$$\therefore y = 2 \tan^{-1}\left(\frac{x+y}{2}\right)$$

**19** We have,  $\frac{dy}{dx} + 1 = e^{x+y}$

On putting  $x+y = z$  and  $1 + \frac{dy}{dx} = \frac{dz}{dx}$ ,

we get

$$\therefore \frac{dz}{dx} = e^z$$

$$\Rightarrow e^{-z} dz = dx$$

On integrating both sides, we get

$$\frac{e^{-z} - 1}{-1} = x + C$$

$$\Rightarrow x + e^{-(x+y)} + C = 0$$

**20**  $\frac{dy}{dx} = \frac{y}{x} + \frac{f(y/x)}{f'(y/x)}$

On putting  $y/x = v$  and  $\frac{dy}{dx} = v + x \frac{dv}{dx}$ ,

we get

$$\frac{f'(v)}{f(v)} dv = \frac{dx}{x}$$

$$\Rightarrow \log |f(v)| = \log |x| + \log C, C > 0$$

$$\Rightarrow |f(y/x)| = C |x|, C > 0.$$

**21** Given,  $\frac{dy}{dx} = \left(\frac{y}{x}\right) \left[\log\left(\frac{y}{x}\right) + 1\right]$

Put  $y = tx$ ,

and  $\frac{dy}{dx} = t + x \frac{dt}{dx}$

Then, we get  $t + x \frac{dt}{dx} = t \log t + t$

$$\Rightarrow \frac{dt}{t \log t} = \frac{dx}{x}$$

$$\Rightarrow \log \log t = \log x + \log C$$

$$\therefore \log\left(\frac{y}{x}\right) = Cx$$

**22** Given,  $\frac{dx}{dy} + \frac{x^2 - xy + y^2}{y^2} = 0$

$$\Rightarrow \frac{dx}{dy} + \left(\frac{x}{y}\right)^2 - \left(\frac{x}{y}\right) + 1 = 0$$

Put  $v = \frac{x}{y}$  or  $x = vy$

and  $\frac{dx}{dy} = v + y \frac{dv}{dy}$

Then, we get  $v + y \frac{dv}{dy} + v^2 - v + 1 = 0$

$$\Rightarrow \frac{dv}{v^2 + 1} + \frac{dy}{y} = 0$$

$$\Rightarrow \tan^{-1}(v) + \log y + C = 0 \quad [\text{integrating}]$$

$$\therefore \tan^{-1}\left(\frac{x}{y}\right) + \log y + C = 0$$

**23** We have  $\frac{dy}{dx} + \frac{y(x+y)}{x^2} = 0$

On putting  $y = vx$ , we get

$$v + x \frac{dv}{dx} + v(1+v) = 0$$

or  $\frac{dx}{x} + \frac{dv}{v(v+2)} = 0$

or  $\frac{dx}{x} + \frac{1}{2} \left(\frac{1}{v} - \frac{1}{v+2}\right) dv = 0$

On integrating, we get

$$\log x + \frac{1}{2} [\log v - \log(v+2)] + \log C = 0$$

or  $\log(v+2) = \log x^2 v C^2$

or  $y + 2x = C^2 x^2 y$ .

**24** Given,  $\frac{dy}{dx} = \frac{xy}{x^2 + y^2}$

Put  $y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx}$ , we get

$$\Rightarrow v + x \frac{dv}{dx} = \frac{v}{1+v^2}$$

$$\Rightarrow x \frac{dv}{dx} = \frac{v}{1+v^2} - v = -\frac{v^3}{1+v^2}$$

$$\Rightarrow \frac{1+v^2}{v^3} dv = -\frac{dx}{x}$$

$$\Rightarrow -\frac{1}{2v^2} + \log v = -\log x + C$$

$$\Rightarrow \frac{x^2}{2y^2} + C = \log \frac{y}{x} + \log x = \log y.$$

Now,  $x = 1, y = 1$

$$\Rightarrow C = -1/2$$

and  $x = x_0, y = e$

$$\Rightarrow \frac{x_0^2}{2e^2} - \frac{1}{2} = 1$$

$$\Rightarrow x_0^2 = 3e^2.$$

**25** We have,

$$y(1+xy) dx = x dy$$

$$\Rightarrow \frac{dy}{dx} = \frac{y(1+xy)}{x} = \frac{y}{x} + y^2$$

On putting  $y = vx$  and  $\frac{dy}{dx} = v + x \frac{dv}{dx}$ ,

we get

$$v + x \frac{dv}{dx} = v + v^2 x^2$$

$$\Rightarrow \frac{dv}{v^2} = x dx$$

$$\Rightarrow \int \frac{dv}{v^2} = \int x dx$$

$$\Rightarrow -\frac{1}{v} = \frac{x^2}{2} + C$$

$$\Rightarrow -\frac{x}{y} = \frac{x^2}{2} + C$$

Put  $(1, -1)$ , then

$$C = \frac{1}{2}$$

$$\therefore -\frac{x}{y} = \frac{x^2}{2} + \frac{1}{2}$$

Now, put  $x = -\frac{1}{2}, y = \frac{4}{5}$

$$\therefore f\left(-\frac{1}{2}\right) = \frac{4}{5}$$

**26** Given,  $(1+x^2) \frac{dy}{dx} + 2xy = 4x^2$

$$\Rightarrow \frac{dy}{dx} + \frac{2x}{1+x^2} y = \frac{4x^2}{1+x^2}$$

$$\therefore \text{IF} = e^{\int \frac{2x}{1+x^2} dx} = e^{\log(1+x^2)} = 1+x^2$$

and the solution is

$$y \cdot (1+x^2) = \int 4x^2 dx + C$$

$$y(1+x^2) = \frac{4x^3}{3} + C$$

Now,  $x = 0, y = 0$

$$\Rightarrow C = 0$$

$$\therefore y(1+x^2) = \frac{4x^3}{3}$$

$$\Rightarrow 3y(1+x^2) = 4x^3$$

**27** We have,  $\frac{dy}{dx} + \frac{3x^2}{1+x^3} y = \frac{\sin^2 x}{1+x^3}$

Since, it is a linear equation with

$$P = \frac{3x^2}{1+x^3}$$

$$\therefore \text{IF} = e^{\int P dx} = e^{\log(1+x^3)} = 1+x^3$$



and the solution is

$$y(1+x^3) = \int \frac{\sin^2 x}{1+x^3} (1+x^3) dx$$

$$= \int \frac{1-\cos 2x}{2} dx$$

$$\therefore y(1+x^3) = \frac{1}{2}x - \frac{\sin 2x}{4} + C$$

**28** Given,  $\frac{dy}{dx} - y \tan x = 2x \sec x, y(0) = 0$

$$\text{IF} = e^{-\int \tan x dx} = e^{-\log \sec x}$$

$$\text{IF} = \cos x$$

$$\therefore \cos x \cdot y = \int 2x \sec x \cdot \cos x dx$$

$$\Rightarrow \cos x \cdot y = x^2 + C$$

$$\Rightarrow y(0) = 0 \Rightarrow C = 0$$

$$\therefore y = x^2 \sec x$$

and  $y' = 2x \cdot \sec x + x^2 \sec x \cdot \tan x$

$$y\left(\frac{\pi}{4}\right) = \frac{\pi^2}{16} \cdot \sqrt{2} = \frac{\pi^2}{8\sqrt{2}}$$

$$y'\left(\frac{\pi}{4}\right) = \frac{\pi}{2} \cdot \sqrt{2} + \frac{\pi^2}{16} \cdot \sqrt{2}$$

$$y\left(\frac{\pi}{3}\right) = \frac{\pi^2}{9} \cdot 2 = \frac{2\pi^2}{9}$$

$$y'\left(\frac{\pi}{3}\right) = 2 \cdot \frac{\pi}{3} \cdot 2 + \frac{\pi^2}{9} \cdot 2 \cdot \sqrt{3}$$

$$= \frac{4\pi}{3} + \frac{2\pi^2\sqrt{3}}{9}$$

**29** This is the linear equation of the form  $\frac{dy}{dx} + Py = Q$ .

where,  $P = \tan x$  and  $Q = x^m \cos x$   
Now, integrating factor (IF)

$$= e^{\int P dx} = e^{\int \tan x dx}$$

$$= e^{\log \sec x} = \sec x$$

and the solution is given by,

$$y \cdot e^{\int P dx} = \int Q \cdot e^{\int P dx} dx + C$$

$$\Rightarrow y \cdot \sec x = \int x^m \cdot \cos x \cdot \sec x dx + C$$

$$\Rightarrow y \sec x = \frac{x^{m+1}}{m+1} + C$$

$$\therefore (m+1)y = x^{m+1} \cos x + C(m+1) \cos x$$

**30** Given differential equation is

$\frac{dP}{dt} - \frac{1}{2}P(t) = -200$ , which is a linear differential equation.

Here,  $P(t) = \frac{-1}{2}$  and  $Q(t) = -200$

$$\text{Now, IF} = e^{\int -\frac{1}{2} dt} = e^{-\frac{t}{2}}$$

and the solution is

$$P(t) \cdot \text{IF} = \int Q(t) \text{IF} dt + K$$

$$P(t) \cdot e^{-\frac{t}{2}} = -\int 200 e^{-\frac{t}{2}} dt + K$$

$$P(t) \cdot e^{-\frac{t}{2}} = 400 e^{-\frac{t}{2}} + K$$

$$\Rightarrow P(t) = 400 + Ke^{\frac{t}{2}}$$

If  $P(0) = 100$ , then  $K = -300$

$$\Rightarrow P(t) = 400 - 300e^{\frac{t}{2}}$$

**31** We have,  $(xy^5 + 2y)dx = xdy$

$$\Rightarrow x \frac{dy}{dx} - 2y = xy^5$$

$$\Rightarrow \frac{dy}{dx} - \frac{2y}{x} = y^5$$

$$\Rightarrow y^{-5} \frac{dy}{dx} - \frac{2y^{-4}}{x} = 1 \quad \dots(i)$$

$$\text{Put, } y^{-4} = t$$

$$\Rightarrow -4y^{-5} \frac{dy}{dx} = \frac{dt}{dx}$$

$$\Rightarrow y^{-5} \frac{dy}{dx} = \frac{-1}{4} \frac{dt}{dx} \quad \dots(ii)$$

From Eqs. (i) and (ii), we get

$$-\frac{1}{4} \frac{dt}{dx} - \frac{2t}{x} = 1$$

$$\Rightarrow \frac{dt}{dx} + \frac{8t}{x} = -4$$

$$\text{Now, IF} = e^{\int \frac{8}{x} dx} = e^{8 \log x} = x^8$$

and the solution is

$$t \cdot x^8 = \int (-4)x^8 dx + C$$

$$\Rightarrow \frac{x^8}{y^4} = -\frac{4 \cdot x^9}{9} + C$$

$$\Rightarrow 9x^8 + 4x^9 \cdot y^4 = 9y^4 C$$

**32** We have,  $\sin x \frac{dy}{dx} + y \cos x = 4x$

$$\Rightarrow \sin x dy + y \cos x dx = 4x dx$$

$$\Rightarrow d(y \sin x) = 4x dx$$

On integrating both sides, we get

$$y \sin x = 2x^2 + C$$

Since, it passes through  $\left(\frac{\pi}{2}, 0\right)$

$$\therefore 0 = \frac{\pi^2}{2} + C \Rightarrow C = -\frac{\pi^2}{2}$$

$$\Rightarrow y \sin x = 2x^2 - \frac{\pi^2}{2}$$

$$\Rightarrow y = 2x^2 \operatorname{cosec} x - \frac{\pi^2}{2} \operatorname{cosec} x$$

$$\Rightarrow y(\pi/6) = 2 \left(\frac{\pi^2}{36}\right) \operatorname{cosec} \frac{\pi}{6} - \frac{\pi^2}{2} \operatorname{cosec} \frac{\pi}{6}$$

$$= 2 \left(\frac{\pi^2}{36}\right) 2 - \frac{\pi^2}{2} \cdot 2$$

$$= -\frac{8\pi^2}{9}$$

**33 Statement I** Let the equation of parabola whose axis is the axis of  $x$  and vertex at the origin is

$$y^2 = 4ax$$

$$\Rightarrow 2y \frac{dy}{dx} = 4a$$

$$\Rightarrow \frac{dy}{dx} = \frac{2a}{y}$$

$$\Rightarrow \frac{dy}{dx} \propto \frac{1}{y}$$

[where,  $a \rightarrow$  parameter]

**Statement II**  $y^2 = 4ax \quad \dots(i)$

$$\Rightarrow 2y \frac{dy}{dx} = 4a$$

$$\Rightarrow a = \frac{y}{2} \cdot \frac{dy}{dx} \quad [\text{from Eq. (i)}]$$

$$\Rightarrow y^2 = 4x \cdot \frac{y}{2} \cdot \frac{dy}{dx}$$

$$\Rightarrow y^2 = 2xy \frac{dy}{dx}$$

$$\Rightarrow y = 2x \cdot \frac{dy}{dx}$$

which has order = 1 and degree = 1

**34** Let  $c_1 + c_2 + c_3 e^{c_4} = A$  [constant]

Then,  $y = Ax$

$$\Rightarrow \frac{dy}{dx} = A$$

$$\Rightarrow y = x \frac{dy}{dx}$$

$$\therefore x \frac{dy}{dx} = y$$

## SESSION 2

**1** Given,  $y = c_1 e^{c_2 x}$

$$\Rightarrow y' = c_1 c_2 e^{c_2 x} \Rightarrow c_2 = \frac{y'}{y}$$

$$\text{and } y'' = c_1 c_2^2 e^{c_2 x} \Rightarrow y'' = y \cdot \left(\frac{y'}{y}\right)^2$$

$$\Rightarrow yy'' = (y')^2$$

**2** Given,

$$p'(t) = \frac{dp(t)}{dt} = 0.5 p(t) - 450$$

$$\Rightarrow \frac{2dp(t)}{p(t) - 900} = dt$$

$$\Rightarrow \int \frac{2dp(t)}{p(t) - 900} = \int dt$$

$$\Rightarrow 2 \log |p(t) - 900| = t + C$$

To find the value of  $C$ , let's substitute  $t = 0$  and  $p(0) = 850$

$$\Rightarrow 2 \log |p(0) - 900| = 0 + C$$

$$\Rightarrow C = 2 \log |850 - 900|$$

$$\Rightarrow C = 2 \log 50$$

Now,  $2 \log |p(t) - 900| = t + 2 \log 50$

Now, put  $p(t) = 0$ , then

$$2 \log |0 - 900| = t + 2 \log 50$$

$$\Rightarrow t = 2 \log \left| \frac{900}{50} \right| = 2 \log 18$$

**3** Given slope at  $(x, y)$  is

$$\frac{dy}{dx} = \frac{y}{x} + \sec(y/x)$$

$$\text{Let } \frac{y}{x} = t \Rightarrow y = xt$$

$$\text{and } \frac{dy}{dx} = t + x \frac{dt}{dx}$$

$$\text{Now, } t + x \frac{dt}{dx} = t + \sec(t)$$

$$\int \cos t \, dt = \int \frac{1}{x} dx$$

$$\sin t = \ln x + C$$

$$\sin(y/x) = \ln x + C$$

$\therefore$  This curve passes through  $(1, \pi/6)$

$$\sin(\pi/6) = \ln(1) + C \Rightarrow C = \frac{1}{2}$$

Thus,  $\sin\left(\frac{y}{x}\right) = \ln x + \frac{1}{2}$

4 We have,  $y'(x) + y(x)g'(x) = g(x)g'(x)$

Linear differential equation with integrating factor  $e^{g(x)}$

$$\Rightarrow y(x) \cdot e^{g(x)} = \int g(x) \cdot g'(x) \cdot e^{g(x)} dx$$

$$\Rightarrow y(x) \cdot e^{g(x)} = e^{g(x)}(g(x) - 1) + C$$

Since,  $y(0) = 0$  and  $g(0) = 0$ , therefore  $C = 1$

$$\Rightarrow y(x) = (g(x) - 1) + e^{-g(x)}$$

$$\Rightarrow y(2) = (g(2) - 1) + e^{-g(2)} = 0,$$

as  $g(2) = 0$ .

5 Here,  $\frac{dx}{dy} + \frac{1}{y^2} \cdot x = \frac{1}{y^3}$

[linear differential equation in  $x$ ]

Clearly, IF =  $e^{\int \frac{1}{y^2} dy} = e^{-\frac{1}{y}}$

Now, complete solution is,

$$x \cdot e^{-\frac{1}{y}} = \int \frac{1}{y^3} \cdot e^{-\frac{1}{y}} dy$$

$$\Rightarrow x \cdot e^{-\frac{1}{y}} = \int \frac{1}{y} \cdot \frac{1}{y^2} \cdot e^{-\frac{1}{y}} dy$$

Put  $-\frac{1}{y} = t \Rightarrow \frac{1}{y^2} dy = dt$

$$\Rightarrow x e^{-\frac{1}{y}} = \int -t \cdot e^{-t} dt$$

$$\Rightarrow x e^{-\frac{1}{y}} = -\{t \cdot e^{-t} - \int 1 \cdot e^{-t} dt\} + C$$

$$\Rightarrow x e^{-\frac{1}{y}} = -te^{-t} + e^{-t} + C$$

$$\Rightarrow x e^{-\frac{1}{y}} = \frac{1}{y} \cdot e^{-\frac{1}{y}} + e^{-\frac{1}{y}} + C$$

$$\Rightarrow e^{-1} = e^{-1} + e^{-1} + C \quad [\because y(1) = 1]$$

$$\Rightarrow C = -\frac{1}{e}$$

$$\Rightarrow x e^{-\frac{1}{y}} = \frac{1}{y} \cdot e^{-\frac{1}{y}} + e^{-\frac{1}{y}} - \frac{1}{e}$$

$\therefore x = \frac{1}{y} + 1 - \frac{1}{e} \cdot e^{\frac{1}{y}}$

6 On putting  $10x + 6y = t$  and

$10 + 6 \frac{dy}{dx} = \frac{dt}{dx}$ , we get

$$\frac{dt}{dx} = 6 \sin t + 10 = 6 \left( \frac{2 \tan t/2}{1 + \tan^2 t/2} \right) + 10$$

$$\Rightarrow \frac{\sec^2 t/2}{10 \tan^2 t/2 + 12 \tan t/2 + 10} dt = dx$$

$$\Rightarrow \frac{dz}{5z^2 + 6z + 5} = dx \quad [z = \tan(t/2)]$$

$$\Rightarrow \frac{1}{5} \int \frac{dz}{\left(z + \frac{3}{5}\right)^2 + (4/5)^2} = \int dx$$

$$\Rightarrow \frac{1}{4} \tan^{-1} \frac{5z + 3}{4} = x + C$$

$$\Rightarrow 5 \tan t/2 + 3 = 4 \tan 4(x + C)$$

$$\Rightarrow 5 \tan(5x + 3y) + 3 = 4 \tan 4(x + C)$$

Now,  $x = 0, y = 0 \Rightarrow C = \frac{1}{4} \tan^{-1}(3/4)$

Hence, equation of the curve is

$$5 \tan(5x + 3y) = \frac{25 \tan 4x}{4 - 3 \tan 4x}$$

$$\Rightarrow y = \frac{1}{3} \left\{ \tan^{-1} \frac{5 \tan 4x}{4 - 3 \tan 4x} - 5x \right\}$$

7 Given,  $\frac{d\{V(t)\}}{dt} = -k(T - t)$

$$\therefore d\{V(t)\} = -k(T - t) dt \quad \dots(i)$$

$$\Rightarrow \int_0^T d\{V(t)\} = \int_0^T -k(T - t) dt$$

$$\Rightarrow V(T) - V(0) = k \left[ \frac{(t - T)^2}{2} \right]_0^T$$

$$\Rightarrow V(T) - I = \frac{k}{2} [(T - T)^2 - (0 - T)^2]$$

$$[\because \text{when } t = 0, \text{ then } V(t) = I]$$

$$\therefore V(T) = I - \frac{k}{2} T^2$$

8 On putting  $x = \tan A$ , and  $y = \tan B$  in

the given relation, we get

$$\cos A + \cos B = \lambda (\sin A - \sin B)$$

$$\Rightarrow \tan \left( \frac{A - B}{2} \right) = \frac{1}{\lambda}$$

$$\Rightarrow \tan^{-1} x - \tan^{-1} y = 2 \tan^{-1} \left( \frac{1}{\lambda} \right)$$

On differentiating w.r.t.  $x$ , we get

$$\frac{1}{1 + x^2} - \frac{1}{1 + y^2} \cdot \frac{dy}{dx} = 0$$

$$\Rightarrow \frac{dy}{dx} = \frac{1 + y^2}{1 + x^2}$$

Clearly, it is a differential equation of degree 1.

9  $|y| \sqrt{1 + (dx/dy)^2} = 1$

$$\Rightarrow y^2 \left( 1 + \left( \frac{dx}{dy} \right)^2 \right) = 1$$

$$\therefore \frac{dy}{dx} = \pm \frac{y}{\sqrt{1 - y^2}}$$

$$\Rightarrow \int \frac{\sqrt{1 - y^2}}{y} dy = \pm x + C.$$

On putting  $y = \sin \theta$  and  $dy = \cos \theta d\theta$ , we get

$$C \pm x = \int \frac{\cos \theta \cdot \cos \theta d\theta}{\sin \theta}$$

$$= \int (\operatorname{cosec} \theta - \sin \theta) d\theta$$

$$\Rightarrow C \pm x = \log |\operatorname{cosec} \theta - \cot \theta| + \cos \theta$$

$$\Rightarrow C \pm x = \log \left| \frac{1 - \sqrt{1 - y^2}}{y} \right| + \sqrt{1 - y^2}.$$

10 Given,  $\cos x \, dy = y \sin x \, dx - y^2 \, dx$

$$\Rightarrow \frac{1}{y^2} \frac{dy}{dx} - \frac{1}{y} \tan x = -\sec x$$

Put  $-\frac{1}{y} = z \Rightarrow \frac{1}{y^2} \frac{dy}{dx} = \frac{dz}{dx}$

$$\therefore \frac{dz}{dx} + (\tan x)z = -\sec x$$

This is a linear differential equation.

Now, IF =  $e^{\int \tan x \, dx} = e^{\log \sec x} = \sec x$

Hence, the solution is

$$z \cdot (\sec x) = \int -\sec x \cdot \sec x \, dx + C_1$$

$$\Rightarrow -\frac{1}{y} \sec x = -\tan x + C_1$$

$$\therefore \sec x = y(\tan x + C)$$

where  $C = -C_1$ .

11 We have,  $(1 - x^2) \frac{dy}{dx} + xy = ax$

$$\Rightarrow \frac{dy}{dx} + \frac{x}{1 - x^2} y = \frac{ax}{1 - x^2} \quad [\text{L. D. E.}]$$

$$\text{IF} = e^{\int \frac{x}{1 - x^2} dx} = e^{-\frac{1}{2} \log |1 - x^2|}$$

$$= \begin{cases} \frac{1}{\sqrt{1 - x^2}}, & \text{if } -1 < x < 1 \\ \frac{1}{\sqrt{x^2 - 1}}, & \text{if } x < -1 \text{ or } x > 1. \end{cases}$$

If  $-1 < x < 1$ , then solution is

$$y \cdot \frac{1}{\sqrt{1 - x^2}} = a \int \frac{x}{(1 - x^2)^{3/2}} dx + C$$

$$y \cdot \frac{1}{\sqrt{1 - x^2}} = \frac{a}{\sqrt{1 - x^2}} + C$$

$$\Rightarrow y = a + C \sqrt{1 - x^2}$$

$$\Rightarrow (y - a)^2 = C^2(1 - x^2)$$

$$\Rightarrow C^2 x^2 + (y - a)^2 = C^2$$

which is an ellipse.

If  $x < -1$  or  $x > 1$ , solution is

$$y \cdot \frac{1}{\sqrt{x^2 - 1}} = C + a \int \frac{x}{(1 - x^2)\sqrt{x^2 - 1}} dx$$

$$= C - a \int \frac{x}{(x^2 - 1)^{3/2}} dx = C + \frac{a}{\sqrt{x^2 - 1}}$$

$$\Rightarrow y = a + C \sqrt{x^2 - 1}$$

$$\Rightarrow (y - a)^2 = C^2 x^2 - c^2$$

$$\Rightarrow C^2 x^2 - (y - a)^2 = C^2$$

which represents a hyperbola.

12 Given,

$$\frac{dy}{dx} = \frac{(x + 1)^2 + y - 3}{x + 1} = (x + 1) + \frac{y - 3}{x + 1}$$

Putting  $x + 1 = X, y - 3 = Y, \frac{dy}{dx} = \frac{dY}{dX}$ ,

the equation becomes

$$\frac{dY}{dX} = X + \frac{Y}{X}$$

or  $\frac{dY}{dX} - \frac{1}{X} \cdot Y = X$  [L. D. E.]

IF =  $e^{\int (-1/X) dX} = e^{-\log X} = X^{-1} = \frac{1}{X}$

∴ The solution is

$$Y \cdot \left(\frac{1}{X}\right) = C + \int X \cdot \left(\frac{1}{X}\right) dx = C + X$$

or  $\frac{(y-3)}{(x+1)} = C + x + 1$

Now,  $x = 2, y = 0$

$$\Rightarrow \frac{0-3}{2+1} = C + 2 + 1$$

$$\Rightarrow C = -4$$

∴ The equation of the curve is

$$\frac{y-3}{x+1} = x-3 \text{ or } y = x^2 - 2x.$$

**13** We have,  $\frac{x + \frac{x^3}{3!} + \frac{x^5}{5!} + \dots}{1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \dots} = \frac{dx-dy}{dx+dy}$

On applying componendo and dividendo, we get

$$\frac{\left(x + \frac{x^3}{3!} + \frac{x^5}{5!} + \dots\right) + \left(1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \dots\right)}{\left(x + \frac{x^3}{3!} + \frac{x^4}{4!} + \dots\right) - \left(1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \dots\right)}$$

$$= \frac{(dx-dy) + (dx+dy)}{(dx-dy) - (dx-dy)}$$

$$\Rightarrow \frac{\left(1+x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots\right)}{-\left(1-x + \frac{x^2}{2!} - \frac{x^3}{3!} + \dots\right)} = \frac{2dx}{-2dy} = -\frac{dx}{dy}$$

$$\Rightarrow \frac{e^x}{-e^{-x}} = -\frac{dx}{dy}$$

$$\Rightarrow dy = e^{-2x} dx$$

On integrating both sides, we get

$$y = \frac{e^{-2x}}{-2} + C_1$$

$$\Rightarrow 2ye^{2x} = -1 + 2C_1 e^{2x}$$

$$\Rightarrow 2ye^{2x} = Ce^{2x} - 1, \text{ where } C = 2C_1$$

**14** Equation of tangent is  $Y - y = \frac{dy}{dx}(X - x)$

Y-intercept of tangent is  $y - x \frac{dy}{dx}$

From given condition,  $y - x \frac{dy}{dx} = x^3$ ,

$$\Rightarrow \frac{dy}{dx} - \frac{1}{x} \cdot y = -x^2 \quad \dots(i)$$

Now, IF =  $e^{\int \frac{1}{x} dx} = e^{-\log_e x} = \frac{1}{x}$

and solution is

$$\frac{1}{x} \cdot y = -\int x dx = -\frac{x^2}{2} + C$$

$$\Rightarrow f(x) = \frac{-x^3}{2} + Cx$$

Now,  $f(1) = 1 \Rightarrow C = 1 + \frac{1}{2} = \frac{3}{2}$

$$\therefore f(x) = \frac{-x^3}{2} + \frac{3}{2}x$$

$$\Rightarrow f(-3) = \frac{27}{2} - \frac{9}{2} = 9$$

**15**  $\lim_{t \rightarrow x} \frac{t^2 f(x) - x^2 f(x) + x^2 f(x) - x^2 f(t)}{t - x} = 1$

$$\lim_{t \rightarrow x} \left[ (t+x)f(x) + x^2 \left( \frac{f(x) - f(t)}{t-x} \right) \right] = 1$$

$$2xf(x) + x^2(-f'(x)) = 1$$

$$\Rightarrow \frac{x^2 \frac{dy}{dx} - 2xy}{x^4} = -\frac{1}{x^4},$$

where  $y = f(x)$

$$\frac{d}{dx} \left( \frac{y}{x^2} \right) = -\frac{1}{x^4}$$

$$\Rightarrow x^{-2} y = \frac{1}{3x^3} + C$$

$$\Rightarrow x^{-2} y = \frac{1}{3x^3} + \frac{2}{3} \quad [\because f(1) = 1]$$

$$\Rightarrow y = \frac{1}{3x} + \frac{2x^2}{3}$$

**16** Given,  $\frac{dy}{dx} = \frac{y\sqrt{y^2-1}}{x\sqrt{x^2-1}}$

$$\Rightarrow \int \frac{dy}{y\sqrt{y^2-1}} = \int \frac{dx}{x\sqrt{x^2-1}}$$

$$\Rightarrow \sec^{-1} y = \sec^{-1} x + C$$

Now,  $x = 2, y = \frac{2}{\sqrt{3}}$

$$\Rightarrow \frac{\pi}{6} = \frac{\pi}{3} + C \Rightarrow C = -\frac{\pi}{6}$$

$$\therefore y = \sec \left( \sec^{-1} x - \frac{\pi}{6} \right)$$

$$\Rightarrow \frac{1}{y} = \cos \left( \cos^{-1} \frac{1}{x} - \cos^{-1} \frac{\sqrt{3}}{2} \right)$$

$$\Rightarrow \frac{1}{y} = \cos \left[ \cos^{-1} \left( \frac{\sqrt{3}}{2x} + \sqrt{1 - \frac{1}{x^2}} \right) \right]$$

$$\left[ \sqrt{1 - \frac{3}{4}} \right]$$

$$\therefore \frac{1}{y} = \frac{\sqrt{3}}{2x} + \frac{1}{2} \sqrt{1 - \frac{1}{x^2}}$$