

- c) they must be mutually exclusive and the sum of their probabilities must be equal to 1
d) the sum of their probabilities must be equal to 1
both are correct
6. The area bounded by the curves $y^2 = 4x$ and $y = x$ is equal to [1]
a) $\frac{8}{3}$ b) $\frac{35}{6}$
c) none of these d) $\frac{1}{3}$
7. A bag contains 5 red and 3 blue balls. If 3 balls are drawn at random without replacement the probability of getting exactly one red ball is [1]
a) $\frac{45}{196}$ b) $\frac{15}{56}$
c) $\frac{15}{29}$ d) $\frac{135}{392}$
8. If $\vec{a} = (\hat{i} - 2\hat{j} + 3\hat{k})$ and $\vec{b} = (\hat{i} - 3\hat{k})$ and then $|\vec{b} \times 2\vec{a}| = ?$ [1]
a) $2\sqrt{23}$ b) $5\sqrt{17}$
c) $10\sqrt{3}$ d) $4\sqrt{19}$
9. The equation of the line passing through the points $a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$ and $b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$ is [1]
a) $\vec{r} = (a_1\hat{i} + a_2\hat{j} + a_3\hat{k}) - t(b_1\hat{i} + b_2\hat{j} + b_3\hat{k})$ b) $\vec{r} = (a_1\hat{i} + a_2\hat{j} + a_3\hat{k}) + \lambda(b_1\hat{i} + b_2\hat{j} + b_3\hat{k})$
c) None Of These d) $\vec{r} = a_1(1-t)\hat{i} + a_2(1-t)\hat{j} + a_3(1-t)\hat{k} + t(b_1\hat{i} + b_2\hat{j} + b_3\hat{k})$
10. The general solution of the DE $x \frac{dy}{dx} = y + x \tan \frac{y}{x}$ is [1]
a) $\sin\left(\frac{y}{x}\right) = C$ b) $\sin\left(\frac{y}{x}\right) = Cy$
c) none of these d) $\sin\left(\frac{y}{x}\right) = Cx$
11. $\int_0^{\frac{\pi}{4}} \log(1 + \tan x) dx = ?$ [1]
a) $\frac{\pi}{8} \log 2$ b) $\frac{\pi}{4} \log 2$
c) $\frac{\pi}{4}$ d) 0
12. If the area cut off from a parabola by any double ordinate is k times the corresponding rectangle contained by that double ordinate and its distance from [1]

the vertex, then k is equal to

a) $\frac{2}{3}$

b) 3

c) $\frac{1}{3}$

d) $\frac{3}{2}$

13. The function $f(x) = \tan x - x$ [1]

a) always increases

b) never increases

c) always decreases

d) sometimes increases and sometimes decreases.

14. If $A = \frac{1}{3} \begin{bmatrix} 1 & 1 & 2 \\ 2 & 1 & -2 \\ x & 2 & y \end{bmatrix}$ satisfies $A^T A = I$, then $x + y =$ [1]

a) -3

b) none of these

c) 0

d) 3

15. If $A = \begin{bmatrix} 1 & 2 & x \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & -2 & y \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ and $AB = I_3$, then $x + y$ equals [1]

a) -1

b) 0

c) none of these

d) 2

16. The principal value of $\sin^{-1}(\sin \frac{3\pi}{4}) = \dots\dots$ [1]

a) $\frac{\pi}{4}$

b) $\frac{3\pi}{4}$

c) $\frac{5\pi}{4}$

d) $-\frac{\pi}{4}$

17. For the differential equation $\left(\frac{dy}{dx}\right)^2 - x\left(\frac{dy}{dx}\right) + y = 0$, which one of the following is not its solution? [1]

a) $4y = x^2$

b) $y = -x - 1$

c) $y = x - 1$

d) $y = x$

18. **Assertion (A):** $f(x) = 2x^3 - 9x^2 + 12x - 3$ is increasing outside the interval (1, 2). [1]
Reason (R): $f'(x) < 0$ for $x \in (1, 2)$.

a) Both A and R are true and R is the correct explanation of A.

b) Both A and R are true but R is not the correct explanation of A.

c) A is true but R is false.

d) A is false but R is true.



19. If $A = \begin{bmatrix} 2 & \lambda & -3 \\ 0 & 2 & 5 \\ 1 & 1 & 3 \end{bmatrix}$, then A^{-1} exists if. [1]

- a) $\lambda = 2$
c) None of these
b) $\lambda \neq -2$
d) $\lambda \neq 2$

20. **Assertion (A):** If $\begin{vmatrix} x & 2 \\ 18 & x \end{vmatrix} = \begin{vmatrix} 6 & 2 \\ 18 & 6 \end{vmatrix}$ then $x = \pm 6$. [1]

Reason (R): If A is a skew-symmetric matrix of odd order, then $|A| = 0$.

- a) Both A and R are true and R is the correct explanation of A.
- b) Both A and R are true but R is not the correct explanation of A.
- c) A is true but R is false.
- d) A is false but R is true.

Section B

21. Using the principal values, write the value of $\cos^{-1}\left(\frac{1}{2}\right) + 2\sin^{-1}\left(\frac{1}{2}\right)$. [2]

22. Find the general solution of the differential equation $\frac{dy}{dx} - 2y = \cos 3x$. [2]

23. Evaluate: $\begin{vmatrix} \cos 65^\circ & \sin 65^\circ \\ \sin 25^\circ & \cos 25^\circ \end{vmatrix}$. [2]

OR

Evaluate $\Delta = \begin{bmatrix} 0 & \sin \alpha & -\cos \alpha \\ -\sin \alpha & 0 & \sin \beta \\ \cos \alpha & -\sin \beta & 0 \end{bmatrix}$

24. Show that $(\vec{a} \times \vec{b})^2 = |\vec{a}|^2 |\vec{b}|^2 - (\vec{a} \cdot \vec{b})^2 = \begin{vmatrix} \vec{a} \cdot \vec{a} & \vec{a} \cdot \vec{b} \\ \vec{a} \cdot \vec{b} & \vec{b} \cdot \vec{b} \end{vmatrix}$ [2]

25. An electronic assembly consists of two sub-systems say A and B. From previous testing procedures, the following probabilities are assumed to be known: [2]

$$P(A \text{ fails}) = 0.2$$

$$P(\text{B fails alone}) = 0.15$$

$$P(A \text{ and } B \text{ fail}) = 0.15$$

Evaluate the following probabilities.

$$(1) P(\overline{A}|\overline{B})$$

(2) $P(A \text{ fails alone})$.

Section C

26. Find the general solution of the differential equation: [3]

$$(1 - x^2) \frac{dy}{dx} + xy = x\sqrt{1 - x^2}$$

OR

Verify that $y^2 = 4ax$ is a solution of the differential equation $y = x \frac{dy}{dx} + a \frac{dx}{dy}$

27. Evaluate: $\int \frac{x}{\sqrt{x^2+x+1}} dx$ [3]

28. Evaluate the integral: $\int (2x+3)\sqrt{x^2+4x+3} dx$ [3]

OR

By using the properties of definite integrals, evaluate the integral $\int_{-5}^5 |x+2| dx$

29. If the function $f(x)$ given by $f(x) = \begin{cases} 3ax+b, & \text{if } x > 1 \\ 11, & \text{if } x = 1 \\ 5ax-2b, & \text{if } x < 1 \end{cases}$ is continuous at $x = 1$, [3]

then find the values of a and b .

30. Sketch the graph $y = |x-5|$. Evaluate $\int_0^1 |x-5| dx$. What does this value of the [3]
integral represent on the graph.

31. If $\vec{a}, \vec{b}, \vec{c}$ are three vectors such that $|\vec{a}| = 5, |\vec{b}| = 12$ and $|\vec{c}| = 13$, and [3]
 $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ find the value of $\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}$

OR

The vectors $\vec{a} = 3\hat{i} + x\hat{j} - \hat{k}$ and $\vec{b} = 2\hat{i} - \hat{j} + y\hat{k}$ are mutually \perp . Given $|\vec{a}| = |\vec{b}|$
find x and y .

Section D

32. Solve the Linear Programming Problem graphically: [5]
Maximize $Z = 3x + 4y$ subject to the constraints: $x + y \leq 4, x \geq 0, y \geq 0$

33. Let n be a fixed positive integer. Define a relation R on Z as follows: [5]
 $(a, b) \in R \Leftrightarrow a - b$ is divisible by n . Show that R is an equivalence relation on Z .

OR

Let R be relation defined on the set of natural number N as follows:

$R = \{(x, y) : x \in N, y \in N, 2x + y = 41\}$. Find the domain and range of the relation R .
Also verify whether R is reflexive, symmetric and transitive.

34. Differentiate $\sin^{-1}(4x\sqrt{1-4x^2})$ with respect to $\sqrt{1-4x^2}$, if $x \in \left(-\frac{1}{2}, -\frac{1}{2\sqrt{2}}\right)$ [5]

35. Show that the straight lines whose direction cosines are given by the equations al [5]
 $+bm+cn=0$ and $ul^2+vm^2+wn^2=0$ are perpendicular, if
 $a^2(v+w)+b^2(u+w)+c^2(u+v)=0$ and, parallel, if $\frac{a^2}{u} + \frac{b^2}{v} + \frac{c^2}{w} = 0$

OR




$\vec{AB} = 3\hat{i} - \hat{j} + \hat{k}$ and $\vec{CD} = -3\hat{i} + 2\hat{j} + 4\hat{k}$ are two vectors. The position vectors of the points A and C are $6\hat{i} + 7\hat{j} + 4\hat{k}$ and $-9\hat{j} + 2\hat{k}$, respectively. Find the position vector of a point P on the line AB and a point Q on the line CD such that \vec{PQ} is perpendicular to \vec{AB} and \vec{CD} both.

Section E

36. Read the text carefully and answer the questions:

[4]

A gardener wants to construct a rectangular bed of garden in a circular patch of land. He takes the maximum perimeter of the rectangular region as possible.
(Refer to the images given below for calculations)



- Find the perimeter of rectangle in terms of any one side and radius of circle.
- Find critical points to maximize the perimeter of rectangle?
- Check for maximum or minimum value of perimeter at critical point.

OR

If a rectangle of the maximum perimeter which can be inscribed in a circle of radius 10 cm is square, then the perimeter of region.

37. Read the text carefully and answer the questions:

[4]

Three schools A, B and C organized a mela for collecting funds for helping the rehabilitation of flood victims. They sold handmade fans, mats, and plates from recycled material at a cost of ₹ 25, ₹ 100 and ₹ 50 each. The number of articles sold by school A, B, C are given below.



Article	School	A	B	C
Fans		40	25	35
Mats		50	40	50
Plates		20	30	40

- Represent the sale of handmade fans, mats and plates by three schools A, B and C and the sale prices (in ₹) of given products per unit, in matrix form.

- (ii) Find the funds collected by school A, B and C by selling the given articles.
- (iii) If they increase the cost price of each unit by 20%, then write the matrix representing new price.

OR

Find the total funds collected for the required purpose after 20% hike in price.

38. **Read the text carefully and answer the questions:**

[4]

To teach the application of probability a maths teacher arranged a surprise game for 5 of his students namely Govind, Girish, Vinod, Abhishek and Ankit. He took a bowl containing tickets numbered 1 to 50 and told the students go one by one and draw two tickets simultaneously from the bowl and replace it after noting the numbers.



- (i) Teacher ask Govind, what is the probability that tickets are drawn by Abhishek, shows a prime number on one ticket and a multiple of 4 on other ticket?
- (ii) Teacher ask Girish, what is the probability that tickets drawn by Ankit, shows an even number on first ticket and an odd number on second ticket?

SOLUTION

Section A

1. (a) $\frac{\pi}{2}$

Explanation: Let's consider the first parallel vector to be $\vec{a} = a\hat{i} + b\hat{j} + c\hat{k}$ and second parallel vector be $\vec{b} = (b-c)\hat{i} + (c-a)\hat{j} + (a-b)\hat{k}$

For the angle, we can use the formula $\cos\alpha = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| \times |\vec{b}|}$

For that, we need to find the magnitude of these vectors

$$|\vec{a}| = \sqrt{a^2 + b^2 + c^2}$$

$$= \sqrt{a^2 + b^2 + c^2}$$

$$|\vec{b}| = \sqrt{(b-c)^2 + (c-a)^2 + (a-b)^2}$$

$$= \sqrt{2(a^2 + b^2 + c^2 - ab - bc - ca)}$$

$$\Rightarrow \cos\alpha = \frac{(a\hat{i} + b\hat{j} + c\hat{k}) \cdot ((b-c)\hat{i} + (c-a)\hat{j} + (a-b)\hat{k})}{\sqrt{2(a^2 + b^2 + c^2 - ab - bc - ca)} \times \sqrt{a^2 + b^2 + c^2}}$$

$$\Rightarrow \cos\alpha = \frac{ab - ac + bc - ba + ca - cb}{\sqrt{2(a^2 + b^2 + c^2 - ab - bc - ca)} \times \sqrt{a^2 + b^2 + c^2}}$$

$$\Rightarrow \cos\alpha = \frac{0}{\sqrt{2(a^2 + b^2 + c^2 - ab - bc - ca)} \times \sqrt{a^2 + b^2 + c^2}}$$

$$\Rightarrow \alpha = \cos^{-1}(0)$$

$$\therefore \alpha = \frac{\pi}{2}$$

2. (a) $\cos^{-1}\left(\frac{31}{50}\right)$

Explanation: Given vectors $\vec{a} = \hat{i} + 2\hat{j} - 3\hat{k}$ and $\vec{b} = 3\hat{i} - \hat{j} + 2\hat{k}$

$(2\vec{a} + \vec{b}) = (5\hat{i} + 3\hat{j} - 4\hat{k})$ and $(\vec{a} + 2\vec{b}) = (7\hat{i} + 0\hat{j} + \hat{k})$

$$\cos\theta = \frac{(5 \times 7 + 3 \times 0 - 4 \times 1)}{\sqrt{50} \times \sqrt{50}} = \frac{31}{50} \Rightarrow \theta = \cos^{-1}\left(\frac{31}{50}\right)$$

3. (c) $\frac{e^x}{x+4} + C$

Explanation: $I = \int \frac{x+3}{(x+4)^2} e^x dx$

$$I = \int \left(\frac{x+4-1}{(x+4)^2} \right) e^x dx$$

$$I = \int \left(\frac{1}{x+4} - \frac{1}{(x+4)^2} \right) e^x dx$$

$$f(x) = \frac{1}{x+4} \Rightarrow f'(x) = -\frac{1}{(x+4)^2}$$

$$\Rightarrow I = \frac{e^x}{x+4} + c$$

4. (d) $-\frac{1}{2} x \cos 2x + \frac{1}{4} \sin 2x + C$

Explanation: $I = \int x \sin 2x dx$

By using IBP Formula we get,

$$I = x \int \sin 2x dx - \int \left(\frac{d}{dx} x \right) \int \sin 2x dx$$

$$= -\frac{x}{2} \cos 2x + \frac{1}{4} \sin 2x + c$$

5. (a) None of these

Explanation: If two events A and B are independent, then we know that

$$P(A \cap B) = P(A) \cdot P(B), P(A) \neq 0, P(B) \neq 0$$

Since, A and B have a common outcome.

Further, mutually exclusive events never have a common outcome.

In other words, two independent events having non-zero probabilities of occurrence cannot be mutually exclusive and conversely, i.e., two mutually exclusive events having non-zero probabilities of outcome cannot be independent.

6. (a) $\frac{8}{3}$

Explanation: The two curves $y^2 = 4x$ and $y = x$ meet where $x^2 = 4x$ i.e. where $x = 0$ or $x = 4$. Moreover, the parabola lies above the line $y = x$ between $x = 0$ and $x = 4$. Hence, the required area is:

$$\begin{aligned} \int_0^4 (\sqrt{4x} - x) dx &= \int_0^4 \left(2x^{\frac{1}{2}} - x \right) dx \\ &= \left[\frac{2x^{\frac{3}{2}}}{\frac{3}{2}} - \frac{x^2}{2} \right]_0^4 \\ &= \frac{4}{3} \left(4^{\frac{3}{2}} \right) - \frac{16}{2} = \frac{32}{3} - 8 = \frac{8}{3} \end{aligned}$$

7. (b) $\frac{15}{56}$

Explanation: Probability of getting exactly one red (R) ball

$$= P_R \cdot P_B \cdot P_B + P_B \cdot P_R \cdot P_R + P_B \cdot P_B \cdot P_R$$

$$= \frac{5}{8} \cdot \frac{3}{7} \cdot \frac{2}{6} + \frac{3}{8} \cdot \frac{5}{7} \cdot \frac{2}{6} + \frac{3}{8} \cdot \frac{2}{7} \cdot \frac{5}{6}$$

$$= \frac{15}{4 \cdot 7 \cdot 6} + \frac{15}{4 \cdot 7 \cdot 6} + \frac{15}{4 \cdot 7 \cdot 6}$$

$$= \frac{5}{56} + \frac{5}{56} + \frac{5}{56} = \frac{15}{56}$$

Which is the required solution

8. (d) $4\sqrt{19}$

Explanation: $2\vec{a} = (2\hat{i} - 4\hat{j} + 6\hat{k})$ and $\vec{b} = (\hat{i} - 3\hat{k})$

$$\begin{aligned} \text{Now, } |\vec{b} \times 2\vec{a}| &= \left| \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & 0 & 3 \\ 2 & -4 & 6 \end{vmatrix} \right| = | -12\vec{i} - 12\vec{j} - 4\vec{k} | \\ &= \sqrt{(144) + (144) + 16} = \sqrt{304} \\ &= 4\sqrt{19} \end{aligned}$$

9. (d) $\vec{r} = a_1(1-t)\hat{i} + a_2(1-t)\hat{j} + a_3(1-t)\hat{k} + t(b_1\hat{i} + b_2\hat{j} + b_3\hat{k})$

Explanation: $\vec{r} = a_1(1-t)\hat{i} + a_2(1-t)\hat{j} + a_3(1-t)\hat{k} + t(b_1\hat{i} + b_2\hat{j} + b_3\hat{k})$

Equation of the line passing through the points having position vectors



$a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$ and $b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$ is

$\vec{r} = (a_1\hat{i} + a_2\hat{j} + a_3\hat{k}) + t \{ (b_1\hat{i} + b_2\hat{j} + b_3\hat{k}) - (a_1\hat{i} + a_2\hat{j} + a_3\hat{k}) \}$, where t is a parameter

$$= (a_1\hat{i} + a_2\hat{j} + a_3\hat{k}) - t(a_1\hat{i} + a_2\hat{j} + a_3\hat{k}) + t(b_1\hat{i} + b_2\hat{j} + b_3\hat{k})$$

$$= a_1(1-t)\hat{i} + a_2(1-t)\hat{j} + a_3(1-t)\hat{k} + t(b_1\hat{i} + b_2\hat{j} + b_3\hat{k})$$

10. (d) $\sin\left(\frac{y}{x}\right) = Cx$

Explanation: Given DE: $x \frac{dy}{dx} = y + x \tan \frac{y}{x}$

Now, Dividing both sides by x , we obtain $\frac{dy}{dx} = \frac{y}{x} + \tan \frac{y}{x}$

Let $y = v x$ Differentiating both sides, we get

$$\frac{dy}{dx} = v + x \frac{dv}{dx}$$

Now, our differential equation becomes,

$$v + x \frac{dv}{dx} = v + \tan v$$

On separating the variables, we obtain $\frac{dv}{\tan v} = \frac{dx}{x}$

Integrating both sides, we get, $\sin v = Cx$

Substituting the value of v we get, $\sin\left(\frac{y}{x}\right) = Cx$

11. (a) $\frac{\pi}{8} \log 2$

Explanation: let $I = \int_0^{\pi} \log(1 + \tan x) dx$

We know that,

$$\therefore \int_0^a f(x) dx = \int_0^a f(a-x) dx = 1$$

$$\therefore f(a-x) = \log\left(1 + \tan\left(\frac{\pi}{4} - x\right)\right)$$

$$= \log\left(1 + \frac{\left(\tan \frac{\pi}{4} - \tan x\right)}{1 + \tan \frac{\pi}{4} \tan x}\right)$$

$$= \log \left(1 + 1(1 - \tan x) \frac{1}{1 + \tan x} \right)$$

$$= \log \frac{2}{1 + \tan x}$$

$$\therefore \int_0^{\pi/4} f(a-x) dx = 1$$

$$= \int_0^{\pi/4} \log \frac{2}{1 + \tan x} dx$$

$$= \int_0^{\pi/4} \log 2 dx - \int_0^{\pi/4} (1 + \tan x) dx$$

$$\therefore I = \int_0^{\pi/4} \log 2 dx - 1$$

$$\therefore 2I = \frac{\pi}{4} \log 2$$

$$\therefore I = \frac{\pi}{8} \log 2$$

12. (a) $\frac{2}{3}$

Explanation: Required area :

$$= 2 \int_0^a \sqrt{4ax} dx$$

$$= k a (2 \sqrt{4aa})$$

$$= \frac{8\sqrt{a}}{3} a \frac{3}{2}$$

$$= 4\sqrt{aka} \frac{3}{2} \Rightarrow k = \frac{2}{3}$$

13. (a) always increases

Explanation: We have, $f(x) = \tan x - x$

$$\therefore f'(x) = \sec^2 x - 1$$

$$\Rightarrow f'(x) \geq 0, \forall x \in R$$

So, $f(x)$ always increases

14. (b) none of these

Explanation: We have, $A = \frac{1}{3} \begin{bmatrix} 1 & 1 & 2 \\ 2 & 1 & -2 \\ x & 2 & y \end{bmatrix}$

$$\Rightarrow A^T = \frac{1}{3} \begin{bmatrix} 1 & 2 & x \\ 1 & 1 & 2 \\ 2 & -2 & y \end{bmatrix}$$

Now, $A^T A = I$

$$\Rightarrow \begin{bmatrix} x^2 + 5 & 2x + 3 & xy - 2 \\ 3 + 2x & 6 & 2y \\ xy - 6 & 2y & y^2 + 8 \end{bmatrix} = \begin{bmatrix} 9 & 0 & 0 \\ 0 & 9 & 0 \\ 0 & 0 & 9 \end{bmatrix}$$

The corresponding elements of two equal matrices are not equal.

Thus, the matrix A is not orthogonal.

15. (b) 0

Explanation: $AB = \begin{bmatrix} 1 & 2 & x \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \times \begin{bmatrix} 1 & -2 & y \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

$$= \begin{bmatrix} 1+0+0 & -2+2+0 & y+0+x \\ 0+0+0 & 0+1+0 & 0+0+0 \\ 0+0+0 & 0+0+0 & 0+0+1 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 0 & x+y \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$I_3 = \begin{bmatrix} 1 & 0 & x+y \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Hence $x + y = 0$

16. (a) $\frac{\pi}{4}$

Explanation: $\sin^{-1}(\sin \frac{3\pi}{4}) = \sin^{-1}(\sin(\pi - \frac{\pi}{4}))$

$$= \sin^{-1}(\sin \frac{\pi}{4}) = \frac{\pi}{4}.$$

17. (d) $y = x$

Explanation: The given differential equation is

$$\left(\frac{dy}{dx}\right)^2 - x\left(\frac{dy}{dx}\right) + y = 0 \dots(i)$$

$$y = x \Rightarrow \frac{dy}{dx} = 1$$

From Eq. (i), $(1)^2 + x(1) + x = 1 \neq 0$

So, $y = x$ is not a solution of Eq. (i).

18. (b) Both A and R are true but R is not the correct explanation of A.

Explanation: Assertion: We have, $f(x) = 2x^3 - 9x^2 + 12x - 3$

$$\Rightarrow f'(x) = 6x^2 - 18x + 12$$

For increasing function, $f'(x) \geq 0$

$$\therefore 6(x^2 - 3x + 2) \geq 0$$

$$\Rightarrow 6(x - 2)(x - 1) \geq 0$$

$$\Rightarrow x \leq 1 \text{ and } x \geq 2$$

$\therefore f(x)$ is increasing outside the interval $(1, 2)$, therefore it is a true statement.

Reason: Now, $f'(x) < 0$

$$\Rightarrow 6(x - 2)(x - 1) < 0$$

$$\Rightarrow 1 < x < 2$$

\therefore Assertion and Reason are both true but Reason is not the correct explanation of Assertion.

19. (c) None of these

Explanation: We have,

$$A = \begin{bmatrix} 2 & \lambda & -3 \\ 0 & 2 & 5 \\ 1 & 1 & 3 \end{bmatrix}$$

A^{-1} exists if $|A| \neq 0$

$$\text{Now } |A| = 2(6 - 5) - \lambda(-5) - 3(-2) = 8 + 5\lambda \neq 0$$

$$\Rightarrow 5\lambda \neq -8$$

$$\Rightarrow \lambda \neq \frac{-8}{5}$$

So, A^{-1} exists if and only if $\lambda \neq \frac{-8}{5}$

20. (b) Both A and R are true but R is not the correct explanation of A.

Explanation: Both A and R are true but R is not the correct explanation of A.

Section B

$$21. \text{ We have, } \cos^{-1}\left(\frac{1}{2}\right) = \cos^{-1}\left(\cos\frac{\pi}{3}\right)$$

$$= \frac{\pi}{3} \left[\because \frac{\pi}{3} \in [0, \pi] \right]$$

$$\text{Also } \sin^{-1}\left(-\frac{1}{2}\right) = \sin^{-1}\left(-\sin \frac{\pi}{6}\right)$$

$$= \sin^{-1}\left(\sin\left(-\frac{\pi}{6}\right)\right)$$

$$= -\frac{\pi}{6} \left[\because -\frac{\pi}{6} \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \right]$$

$$\therefore \cos^{-1}\left(\frac{1}{2}\right) - 2\sin^{-1}\left(-\frac{1}{2}\right) = \frac{\pi}{3} - 2\left(-\frac{\pi}{6}\right)$$

$$= \frac{\pi}{3} + \frac{\pi}{3} = \frac{2\pi}{3}$$

22. The given differential equation is of the form $\frac{dy}{dx} + Py = Q$, where $P = -2$ and $Q = \cos 3x$.

Thus, the given differential equation is linear.

$$\text{IF} = e^{\int P dx} = e^{\int -2 dx} = e^{-2x}$$

So, the required solution is

$$y \times \text{IF} = \int \{Q \times F\} dx + C,$$

$$\text{i.e., } y \times e^{-2x} = \int e^{-2x} \cos 3x dx + C$$

$$= e^{-2x} \left[\frac{-2 \cos 3x + 3 \sin 3x}{\{(-2)^2 + 3^2\}} \right] + C$$

$$\left[\because \int e^{ax} \cos bxdx = e^{ax} \left\{ \frac{a \cos bx + b \sin bx}{(a^2 + b^2)} \right\} \right]$$

$$\therefore y = \frac{(3 \sin 3x - 2 \cos 3x)}{13} + Ce^{2x}, \text{ which is the required solution.}$$

23. Given:

$$\begin{vmatrix} \cos 65^\circ & \sin 65^\circ \\ \sin 25^\circ & \cos 25^\circ \end{vmatrix}$$

By directly opening this determinant we get

$$\cos 65^\circ \times \cos 25^\circ - \sin 25^\circ \times \sin 65^\circ$$

$$= \cos (65^\circ + 25^\circ) \because \cos A \cos B - \sin A \sin B = \cos (A + B)$$

$$= \cos 90^\circ$$

$$= 0$$

OR

Expanding along R_1 , we get,

$$\Delta = 0 \begin{vmatrix} 0 & \sin\beta \\ -\sin\beta & 0 \end{vmatrix} - \sin\alpha \begin{vmatrix} -\sin\alpha & \sin\beta \\ \cos\alpha & 0 \end{vmatrix} - \cos\alpha \begin{vmatrix} -\sin\alpha & 0 \\ \cos\alpha & -\sin\beta \end{vmatrix}$$

$$= 0 - \sin\alpha(0 - \cos\alpha\sin\beta) - \cos\alpha(\sin\alpha\sin\beta - 0) = \sin\alpha\cos\alpha\sin\beta - \cos\alpha\sin\alpha\sin\beta = 0$$

24. We have,

$$(\vec{a} \times \vec{b})^2 = |\vec{a} \times \vec{b}|^2$$

$$\Rightarrow (\vec{a} \times \vec{b})^2 = \{|\vec{a}| |\vec{b}| \sin\theta\}^2$$

$$\Rightarrow (\vec{a} \times \vec{b})^2 = |\vec{a}|^2 |\vec{b}|^2 \sin^2\theta$$

$$\Rightarrow (\vec{a} \times \vec{b})^2 = \{|\vec{a}|^2 |\vec{b}|^2\} (1 - \cos^2\theta)$$

$$\Rightarrow (\vec{a} \times \vec{b})^2 = |\vec{a}|^2 |\vec{b}|^2 - |\vec{a}|^2 |\vec{b}|^2 \cos^2\theta$$

$$\Rightarrow (\vec{a} \times \vec{b})^2 = (\vec{a} \cdot \vec{a})(\vec{b} \cdot \vec{b}) - (\vec{a} \cdot \vec{b})(\vec{a} \cdot \vec{b}) [\because \vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos\theta]$$

$$\Rightarrow (\vec{a} \times \vec{b})^2 = \begin{vmatrix} \vec{a} \cdot \vec{a} & \vec{a} \cdot \vec{b} \\ \vec{a} \cdot \vec{b} & \vec{b} \cdot \vec{b} \end{vmatrix}$$

$$\text{Hence, } (\vec{a} \times \vec{b})^2 = |\vec{a}|^2 |\vec{b}|^2 - (\vec{a} \cdot \vec{b})^2 = \begin{vmatrix} \vec{a} \cdot \vec{a} & \vec{a} \cdot \vec{b} \\ \vec{a} \cdot \vec{b} & \vec{b} \cdot \vec{b} \end{vmatrix}$$

25. Event A fails and B fails denoted by \bar{A} and \bar{B} respectively.

$$\therefore P(\bar{A}) = 0.2 \text{ and } P(\bar{A} \text{ and } \bar{B}) = 0.15$$

$$\Rightarrow P(\bar{A} \cap \bar{B}) = 0.15$$

$$\therefore P(\bar{B} \text{ above}) = P(\bar{B}) - P(\bar{A} \cap \bar{B})$$

$$\Rightarrow 0.15 = P(\bar{B}) - 0.15$$

$$\Rightarrow P(\bar{B}) = 0.30$$

$$\text{i. } P(\bar{A}|\bar{B}) = \frac{P(\bar{A} \cap \bar{B})}{P(\bar{B})} = \frac{0.15}{0.30} = \frac{1}{2} = 0.5$$

$$\text{ii. } P(A \text{ fails alone}) = P(A \text{ alone}) = P(\bar{A}) - P(\bar{A} \cap \bar{B}) = 0.20 - 0.15 = 0.05$$

Section C

26. The given differential equation is,

$$\begin{aligned} (1-x^2) \cdot \frac{dy}{dx} + xy &= x\sqrt{1-x^2} \\ \Rightarrow \frac{dy}{dx} + \frac{x}{1-x^2} \cdot y &= \frac{x}{\sqrt{1-x^2}} \end{aligned}$$

This is of the form $\frac{dy}{dx} + Py = Q$, where,

$$P = \frac{x}{1-x^2} \text{ and } Q = \frac{x}{\sqrt{1-x^2}}$$

Thus, the given differential equation is linear

Now, $IF = e^{\int P dx}$

$$\Rightarrow IF = e^{\int \frac{x}{1-x^2} dx}$$

$$\Rightarrow I.F = e^{-\frac{1}{2} \int \frac{-2x}{1-x^2} dx} \Rightarrow I.F = e^{-\frac{1}{2} \log(1-x^2)}$$

$$\Rightarrow IF = (1-x^2)^{-\frac{1}{2}} \Rightarrow I.F = \frac{1}{\sqrt{1-x^2}}$$

Therefore the solution is given by

$$(I.F) \cdot y = \int (I.F)Q + C$$

$$\Rightarrow \frac{1}{\sqrt{1-x^2}} \cdot y = \int \frac{1}{\sqrt{1-x^2}} \cdot \frac{x}{\sqrt{1-x^2}} dx + C$$

$$\Rightarrow \frac{y}{\sqrt{1-x^2}} = \int \frac{x}{1-x^2} dx + C \Rightarrow \frac{y}{\sqrt{1-x^2}} = -\frac{1}{2} \int \frac{-2x}{1-x^2} dx + C$$

$$\Rightarrow \frac{y}{\sqrt{1-x^2}} = -\frac{1}{2} \log(1-x^2) + C$$

$$\Rightarrow y = -\frac{1}{2} \sqrt{1-x^2} \cdot \log(1-x^2) + C$$

OR

We have, $y^2 = 4ax \dots(i)$

Differentiating both sides of (i) with respect to x, we have,

$$2y \frac{dy}{dx} = 4a$$

$$\Rightarrow \frac{dy}{dx} = \frac{2a}{y} \dots(ii)$$

Differentiating both sides of (i) with respect to y, we have,

$$2y = 4a \frac{dx}{dy}$$

$$\Rightarrow \frac{dx}{dy} = \frac{y}{2a} \dots(iii)$$

$$\therefore x \frac{dy}{dx} + a \frac{dx}{dy} = x \left(\frac{2a}{y} \right) + a \left(\frac{y}{2a} \right) \dots[\text{Using (ii) and (iii)}]$$

$$\Rightarrow x \frac{dy}{dx} + a \frac{dx}{dy} = \frac{2ax}{y} + \frac{y}{2}$$

$$\Rightarrow x \frac{dy}{dx} + a \frac{dx}{dy} = \frac{y^2}{2y} + \frac{y}{2} \dots[\text{Using (i)}]$$

$$\Rightarrow x \frac{dy}{dx} + a \frac{dx}{dy} = \frac{y}{2} + \frac{y}{2}$$

$$\Rightarrow x \frac{dy}{dx} + a \frac{dx}{dy} = y$$

$$\Rightarrow y = x \frac{dy}{dx} + a \frac{dx}{dy}$$

Therefore, the given function is the solution to the given differential equation.

27. Let the given integral be, $I = \int \frac{x}{\sqrt{x^2+x+1}} dx$

$$\text{Let } x = \lambda \frac{d}{dx} (x^2 + x + 1) + \mu$$

$$= \lambda(2x + 1) + \mu$$

$$\Rightarrow x = (2\lambda)x + \lambda + \mu$$

Comparing the coefficients of like powers of x,

$$2\lambda = 1 \Rightarrow \lambda = \frac{1}{2}$$

$$\lambda + \mu = 0$$

$$\Rightarrow \left(\frac{1}{2}\right) + \mu = 0$$

$$\Rightarrow \mu = -\frac{1}{2}$$

$$\text{So, } I = \int \frac{\frac{1}{2}(2x+1) - \frac{1}{2}}{\sqrt{x^2+x+1}} dx$$

$$= \frac{1}{2} \int \frac{(2x+1)}{\sqrt{x^2+x+1}} dx - \frac{1}{2} \int \frac{1}{\sqrt{x^2+2x\left(\frac{1}{2}\right) + \left(\frac{1}{2}\right)^2 - \left(\frac{1}{2}\right)^2 + 1}} dx$$

$$= \frac{1}{2} \int \frac{2x+1}{\sqrt{x^2+x+1}} dx - \frac{1}{2} \int \frac{1}{\sqrt{\left(x+\frac{1}{2}\right)^2 - \left(\frac{\sqrt{3}}{2}\right)^2}} dx$$

$$= \frac{1}{2} \times 2\sqrt{x^2+x+1} - \frac{1}{2} \log \left| x + \frac{1}{2} + \sqrt{\left(x+\frac{1}{2}\right)^2 - \left(\frac{\sqrt{3}}{2}\right)^2} \right| + c \text{ [Since,}$$

$$\int \frac{1}{\sqrt{x}} dx = 2\sqrt{x} + c, \int \frac{1}{\sqrt{x^2-a^2}} dx = \log |x + \sqrt{x^2-a^2}| + c]$$

$$\Rightarrow I = \sqrt{x^2+x+1} - \frac{1}{2} \log \left| x + \frac{1}{2} + \sqrt{x^2+x+1} \right| + c$$

28. Let the given integral be,

$$I = \int (x+1)\sqrt{x^2+x+1} dx$$

$$\text{Also, } x+1 = \lambda \frac{d}{dx} (x^2+x+1) + \mu$$

$$\Rightarrow x+1 = \lambda(2x+1) + \mu$$

$$\Rightarrow x+1 = (2\lambda)x + \lambda + \mu$$

Equating coefficient of like terms

$$2\lambda = 1$$

$$\Rightarrow \lambda = \frac{1}{2}$$

And

$$\lambda + \mu = 1$$

$$\Rightarrow \frac{1}{2} + \mu = 1$$

$$\therefore \mu = \frac{1}{2}$$

$$\begin{aligned}\therefore I &= \frac{1}{2} \int (2x+1) \sqrt{x^2+x+1} dx + \frac{1}{2} \int \sqrt{x^2+x+1} dx \\ &= \frac{1}{2} \int (2x+1) \sqrt{x^2+x+1} dx + \frac{1}{2} \int \sqrt{x^2+x + \left(\frac{1}{2}\right)^2 - \left(\frac{1}{2}\right)^2 + 1} dx \\ &= \frac{1}{2} \int (2x+1) \sqrt{x^2+x+1} dx + \frac{1}{2} \int \sqrt{\left(x+\frac{1}{2}\right)^2 + \left(\frac{\sqrt{3}}{2}\right)^2} dx\end{aligned}$$

Now, let $x^2 + x + 1 = t$

$$\Rightarrow (2x+1) dx = dt$$

Then,

$$\begin{aligned}I &= \frac{1}{2} \int \sqrt{t} dt + \frac{1}{2} \left[\frac{x+\frac{1}{2}}{2} \sqrt{\left(x+\frac{1}{2}\right)^2 + \left(\frac{\sqrt{3}}{2}\right)^2} + \right. \\ &\quad \left. \frac{3}{8} \log \left| \left(x+\frac{1}{2}\right) + \sqrt{\left(x+\frac{1}{2}\right)^2 + \left(\frac{\sqrt{3}}{2}\right)^2} \right| \right] + C \\ &= \frac{1}{2} \times \frac{2}{3} t^{\frac{3}{2}} + \frac{1}{2} \left[\left(\frac{2x+1}{4}\right) \sqrt{x^2+x+1} + \frac{3}{8} \log \left| \left(x+\frac{1}{2}\right) + \sqrt{x^2+x+1} \right| \right] + C \\ &= \frac{1}{3} (x^2+x+1)^{\frac{3}{2}} + \frac{1}{2} \left[\left(\frac{2x+1}{4}\right) \sqrt{x^2+x+1} + \frac{3}{8} \log \left| \left(x+\frac{1}{2}\right) + \sqrt{x^2+x+1} \right| \right] + C\end{aligned}$$

OR

$$\text{Let } I = \int_{-5}^5 |x+2| dx \dots (i)$$

Putting $x+2=0$

$$\Rightarrow x = -2 \in (-5, 5)$$

\therefore From eq. (i),

$$I = \int_{-5}^{-2} |x+2| dx + \int_{-2}^5 |x+2| dx$$

$$\begin{aligned}
&= \int_{-5}^{-2} -(x+2)dx + \int_{-2}^5 (x+2)dx \\
&= -\left(\frac{x^2}{2} + 2x\right)_{-5}^{-2} + \left(\frac{x^2}{2} + 2x\right)_{-2}^5 \\
&= -\left[\left(\frac{4}{2} - 4\right) - \left(\frac{25}{2} - 10\right)\right] + \left[\left(\frac{25}{2} + 10\right) - \left(\frac{4}{2} - 4\right)\right] \\
&= -\left(-2 - \frac{5}{2}\right) + \left(\frac{45}{2} + 2\right) \\
&= 2 + \frac{5}{2} + \frac{45}{2} + 2 \\
&= 4 + 25 = 29
\end{aligned}$$

29. Given, $f(x) = \begin{cases} 3ax + b, & \text{if } x > 1 \\ 11, & \text{if } x = 1 \\ 5ax - 2b, & \text{if } x < 1 \end{cases}$

Since $f(x)$ is continuous at $x=1$, therefore,

$$\text{LHL} = \text{RHL} = f(1) \dots \dots \dots (i)$$

$$\text{Now, LHL} = \lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (5ax - 2b)$$

$$= \lim_{h \rightarrow 0} [5a(1-h) - 2b]$$

$$= \lim_{h \rightarrow 0} (5a - 5ah - 2b) = 5a - 2b$$

$$\text{and RHL} = \lim_{x \rightarrow 1^+} (3ax + b) = \lim_{h \rightarrow 0} [3a(1+h) + b]$$

$$= \lim_{h \rightarrow 0} (3a + 3ah + b) = 3a + b$$

Also, given that $f(1) = 11$

On substituting these values in Eq. (i), we get

$$5a - 2b = 3a + b = 11$$

$$\Rightarrow 3a + b = 11 \dots \dots \dots (ii)$$

$$\text{and } 5a - 2b = 11 \dots \dots (iii)$$

On subtracting $3 \times \text{Eq. (iii)}$ from $5 \times \text{Eq. (ii)}$, we get

$$15a + 5b - 15a + 6b = 55 - 33$$


$$\Rightarrow 11b = 22 \Rightarrow b = 2$$

On putting the value of b in Eq. (ii), we get,

$$3a + 2 = 11 \Rightarrow 3a = 9 \Rightarrow a = 3$$

Hence, $a = 3$ and $b = 2$

30.



We are given that

$y = |x - 5|$ intersect $x = 0$ and $x = 1$ at $(0, 5)$ and $(1, 4)$

Now, $y = |x - 5|$

$= -(x - 5)$ For all $x \in (0, 1)$

Integration represents the area enclosed by the graph from $x = 0$ to $x = 1$

Now area denoted by A , is given by

$$A = \int_0^1 |y| dx$$

$$= \int_0^1 |x - 5| dx$$

$$= \int_0^1 -(x - 5) dx$$

$$= - \int_0^1 (x - 5) dx$$

$$= - \left[\frac{x^2}{2} - 5x \right]_0^1$$

$$= - \left[\left(\frac{1}{2} - 5 \right) - (0 - 0) \right]$$

$$= - \left(-\frac{9}{2} \right)$$

$$= \frac{9}{2} \text{ sq. units}$$

31. If $|\vec{a}| = 5$, $|\vec{b}| = 12$, $|\vec{c}| = 13$

If $(\vec{a} + \vec{b} + \vec{c}) = 0$(i)

Find $\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a} = ?$

Squaring the given equation (i) We get ,

$$(\vec{a} + \vec{b} + \vec{c}) \cdot (\vec{a} + \vec{b} + \vec{c}) = 0$$

$$\vec{a} \cdot \vec{a} + \vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c} + \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{a} \cdot \vec{c} + \vec{c} \cdot \vec{b} + \vec{c} \cdot \vec{c}$$

$$|\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + 2(\vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}) = 0$$

$$25 + 144 + 169 + 2(x) = 0$$

$$338 + 2x = 0$$

$$2x = -338$$

$$x = -169$$

hence, the required term is equal to -169.

OR

$$\vec{a} \cdot \vec{b} = 0 \quad [\because \vec{a} \perp \vec{b}]$$

$$\Rightarrow (3\hat{i} + x\hat{j} - \hat{k}) \cdot (2\hat{i} - \hat{j} + y\hat{k})$$

$$\Rightarrow 6 - x - y = 0$$

$$\Rightarrow y + x = 6 \dots\dots (1)$$

$$|\vec{a}| = |\vec{b}| \quad [\text{Given}]$$

$$3^2 + x^2 + 1 = 2^2 + 1^2 + y^2$$

$$y^2 - x^2 = 5$$

$$(y - x)(y + x) = 5$$

$$6(y - x) = 5$$

$$y - x = \frac{5}{6} \dots\dots (2)$$


From (1) and (2), we get,

$$x = \frac{31}{12}, y = \frac{41}{12}$$

Section D

32. As $x \geq 0, y \geq 0$, therefore we shall shade the other inequalities in the first quadrant only.

Now $x + y \leq 4$



Let $x + y = 4$

$$\Rightarrow \frac{x}{4} + \frac{y}{4} = 1$$

Thus the line has 4 and 4 as intercepts along the axes. Now, $(0, 0)$ satisfies the inequation, i.e., $0 + 0 \leq 4$. Therefore, shaded region OAB is the feasible solution.

Its corners are O $(0, 0)$, A $(4, 0)$, B $(0, 4)$

At O $(0, 0)$ $Z = 0$

At A $(4, 0)$ $Z = 3 \times 4 = 12$

At B $(0, 4)$ $Z = 4 \times 4 = 16$

Hence, max $Z = 16$ at $x = 0, y = 4$.

33. $R = \{(a, b) : a - b \text{ is divisible by } n\}$ on Z .

Now,

Reflexivity: Let $a \in Z$

$$\Rightarrow a - a = 0 \times n$$

$$\Rightarrow a - a \text{ is divisible by } n$$

$$\Rightarrow (a, a) \in R$$

$\Rightarrow R$ is reflexive

Symmetric: Let $(a, b) \in R$

$$\Rightarrow a - b = np \text{ for some } p \in \mathbb{Z}$$

$$\Rightarrow b - a = n(-p)$$

$\Rightarrow b - a$ is divisible by n

$$\Rightarrow (b, a) \in R$$

$\Rightarrow R$ is symmetric

Transitive: Let $(a, b) \in R$ and $(b, c) \in R$

$$\Rightarrow a - b = np \text{ and } b - c = nq \text{ for some } p, q \in \mathbb{Z}$$

$$\Rightarrow a - c = n(p + q)$$

$\Rightarrow a - c$ is divisible by n

$$\Rightarrow (a, c) \in R$$

$\Rightarrow R$ is transitive

Thus, R being reflexive, symmetric and transitive on \mathbb{Z} .

Hence, R is an equivalence relation on \mathbb{Z}

OR

Given that,

$$R = \{(1, 39), (2, 37), (3, 35) \dots (19, 3), (20, 1)\}$$

$$\text{Domain} = \{1, 2, 3, \dots, 20\}$$

$$\text{Range} = \{1, 3, 5, 7, \dots, 39\}$$

R is not reflexive as $(2, 2) \notin R$ as

$$2 \times 2 + 2 \neq 41$$

R is not symmetric

as $(1, 39) \in R$ but $(39, 1) \notin R$

R is not transitive

as $(11, 19) \in R, (19, 3) \in R$

But $(11, 3) \notin R$

Hence, R is neither reflexive, nor symmetric and nor transitive.

34. Let $u = \sin^{-1}(4x\sqrt{1-4x^2})$

Put $2x = \cos\theta$

$$\Rightarrow u = \sin^{-1}(2 \times \cos\theta \sqrt{1 - \cos^2\theta})$$

$$\Rightarrow u = \sin^{-1}(2\cos\theta\sin\theta)$$

$$\Rightarrow u = \sin^{-1}(\sin 2\theta) \dots (i)$$

Let $v = \sqrt{1 - 4x^2} \dots (ii)$

Here,

$$x \in \left(-\frac{1}{2}, -\frac{1}{2\sqrt{2}}\right)$$

$$\Rightarrow 2x \in \left(-1, -\frac{1}{\sqrt{2}}\right)$$

$$\Rightarrow \theta \in \left(\frac{3\pi}{4}, \pi \right)$$

So, from equation (i),

$$u = \pi - 2\theta \left[\text{since, } \sin^{-1}(\sin\theta) = \pi - \theta, \text{ if } \theta \in \left(\frac{\pi}{2}, \frac{3\pi}{2} \right) \right]$$

$$\Rightarrow u = \pi - 2\cos^{-1}(2x) \text{ [since, } 2x = \cos\theta \text{]}$$

Differentiate it with respect to x,

$$\frac{du}{dx} = 0 - 2 \left(\frac{-1}{\sqrt{1-(2x)^2}} \right) \frac{d}{dx}(2x)$$

$$\Rightarrow \frac{du}{dx} = \frac{2}{\sqrt{1-4x^2}} (2)$$

$$\Rightarrow \frac{du}{dx} = \frac{4}{\sqrt{1-4x^2}} \dots \text{(iii)}$$

from equation (ii),

$$\frac{dv}{dx} = \frac{-4x}{\sqrt{1-4x^2}}$$

$$\text{but } x \in \left(-\frac{1}{2}, -\frac{1}{2\sqrt{2}} \right)$$

$$\therefore \frac{dv}{dx} = \frac{-4(-x)}{\sqrt{1-4(-x)^2}}$$

$$\Rightarrow \frac{dv}{dx} = \frac{4x}{\sqrt{1-4x^2}} \dots \text{(iv)}$$

Dividing equation (iii) by (iv)

$$\frac{\frac{du}{dx}}{\frac{dv}{dx}} = \frac{4}{\sqrt{1-4x^2}} \times \frac{\sqrt{1-4x^2}}{4x}$$

$$\therefore \frac{du}{dv} = \frac{1}{x}.$$

35. The given equations are

$$al + bm + cn = 0 \dots \text{(i)}$$

$$\text{and, } ul^2 + vm^2 + wn^2 = 0 \dots \text{(ii)}$$

From (i), we get

$$n = -\left(\frac{al+bm}{c}\right)$$

Substituting $n = -\left(\frac{al+bm}{c}\right)$ in (ii), we get

$$ul^2 + vm^2 + w \frac{(al+bm)^2}{c^2} = 0$$

$$\Rightarrow (c^2u + a^2w)l^2 + 2abwlm + (c^2v + b^2w)m^2 = 0$$

$$\Rightarrow (a^2w + c^2u)\left(\frac{l}{m}\right)^2 + 2abw\left(\frac{l}{m}\right) + (b^2w + c^2v) = 0 \dots(iii)$$

This is a quadratic equation in $\frac{l}{m}$. So, it gives two values of $\frac{l}{m}$. Suppose the two

values be $\frac{l_1}{m_1}$ and $\frac{l_2}{m_2}$.

$$\therefore \frac{l_1}{m_1}, \frac{l_2}{m_2} = \frac{b^2w + c^2v}{a^2w + c^2u} \Rightarrow \frac{l_1l_2}{b^2w + c^2v} = \frac{m_1m_2}{a^2w + c^2u} \dots(iv)$$

Similarly, by making a quadratic equation in $\frac{m}{n}$, we obtain

$$\frac{m_1m_2}{a^2w + c^2u} = \frac{n_1n_2}{a^2v + b^2u} \dots(v)$$

From (iv) and (v), we get

$$\frac{l_1l_2}{b^2w + c^2v} = \frac{m_1m_2}{a^2w + c^2u} = \frac{n_1n_2}{a^2v + b^2u} = \lambda \text{ (say)}$$

$$\Rightarrow l_1l_2 = \lambda(b^2w + c^2v), m_1m_2 = \lambda(a^2w + c^2u), n_1n_2 = \lambda(a^2v + b^2u)$$

For the given lines to be perpendicular, we must have

$$l_1l_2 + m_1m_2 + n_1n_2 = 0$$

$$\Rightarrow \lambda(b^2w + c^2v) + \lambda(a^2w + c^2u) + \lambda(a^2v + b^2u) = 0$$

$$\Rightarrow a^2(v+w) + b^2(u+w) + c^2(u+v) = 0$$

For the given lines to be parallel, the direction cosines must be equal and so the roots of the equation (iii) must be equal.

$$\therefore 4a^2b^2w^2 - 4(a^2w + c^2u)(b^2w + c^2v) = 0 \text{ [On equating discriminant to zero]}$$

$$\Rightarrow a^2c^2vw + b^2c^2uw + c^4uv = 0$$

$$\Rightarrow a^2vw + b^2c^2uw + c^2uv = 0$$

$\Rightarrow \frac{a^2}{u} + \frac{b^2}{v} + \frac{c^2}{w} = 0$ [Dividing throughout by uvw] Hence the required result is proved

OR

We have, $\vec{AB} = 3\hat{i} - \hat{j} + \hat{k}$ and $\vec{CD} = -3\hat{i} + 2\hat{j} + 4\hat{k}$

Also, the position vectors of A and C are $6\hat{i} + 7\hat{j} + 4\hat{k}$ and $-9\hat{j} + 2\hat{k}$, respectively.

Since, PQ is perpendicular to both \vec{AB} and \vec{CD} .

So, P and Q will be foot of perpendicular to both the lines through A and C.

Now, equation of the line through A and parallel to the vector \vec{AB} is,

$$\vec{r} = (6\hat{i} + 7\hat{j} + 4\hat{k}) + \lambda(3\hat{i} - \hat{j} + \hat{k})$$

And the line through C and parallel to the vector \vec{CD} is given by

$$\vec{r} = -9\hat{j} + 2\hat{k} + \mu(-3\hat{i} + 2\hat{j} + 4\hat{k}) \dots (i)$$

$$\text{Let } \vec{r} = (6\hat{i} + 7\hat{j} + 4\hat{k}) + \lambda(3\hat{i} - \hat{j} + \hat{k})$$

$$\text{and } \vec{r} = -9\hat{j} + 2\hat{k} + \mu(-3\hat{i} + 2\hat{j} + 4\hat{k}) \dots (ii)$$

Let $P(6 + 3\lambda, 7 - \lambda, 4 + \lambda)$ is any point on the first line and Q be any point on second line is given by $(-3\mu, -9 + 2\mu, 2 + 4\mu)$.

$$\begin{aligned} \therefore \vec{PQ} &= (-3\mu - 6 - 3\lambda)\hat{i} + (-9 + 2\mu - 7 + \lambda)\hat{j} + (2 + 4\mu - 4 - \lambda)\hat{k} \\ &= (-3\mu - 6 - 3\lambda)\hat{i} + (2\mu + \lambda - 16)\hat{j} + (4\mu - \lambda - 2)\hat{k} \end{aligned}$$

If PQ is perpendicular to the first line, then

$$\begin{aligned} 3(-3\mu - 6 - 3\lambda) - (2\mu + \lambda - 16) + (4\mu - \lambda - 2) &= 0 \\ \Rightarrow -9\mu - 18 - 9\lambda - 2\mu - \lambda + 16 + 4\mu - \lambda - 2 &= 0 \\ \Rightarrow -7\mu - 11\lambda - 4 &= 0 \dots (iii) \end{aligned}$$

If PQ is perpendicular to the second line, then

$$\begin{aligned} -3(-3\mu - 6 - 3\lambda) + (2\mu + \lambda - 16) + (4\mu - \lambda + 2) &= 0 \\ \Rightarrow 9\mu + 18 + 9\lambda + 4\mu + 2\lambda - 32 + 16\mu - 4\lambda - 8 &= 0 \\ \Rightarrow 29\mu + 7\lambda - 22 &= 0 \dots (iv) \end{aligned}$$

On solving Eqs. (iii) and (iv), we get

$$\begin{aligned} -49\mu - 77\lambda - 28 &= 0 \\ \Rightarrow 319\mu + 77\lambda - 242 &= 0 \\ \Rightarrow 270\mu - 270 &= 0 \\ \Rightarrow \mu &= 1 \end{aligned}$$

Using μ in Eq. (iii), we get


$$\begin{aligned} -7(1) - 11\lambda - 4 &= 0 \\ \Rightarrow -7 - 11\lambda - 4 &= 0 \\ \Rightarrow -11 - 11\lambda &= 0 \\ \Rightarrow \lambda &= -1 \end{aligned}$$

$$\begin{aligned} \therefore \vec{PQ} &= [-3(1) - 6 - 3(-1)]\hat{i} + [2(1) + (-1) - 16]\hat{j} + [4(1) - (-1) - 2]\hat{k} \\ &= -6\hat{i} - 15\hat{j} + 3\hat{k} \end{aligned}$$

Section E

36. Read the text carefully and answer the questions:

A gardener wants to construct a rectangular bed of garden in a circular patch of land. He takes the maximum perimeter of the rectangular region as possible. (Refer to the images given below for calculations)



- (i) Let 'y' be the breadth and 'x' be the length of rectangle and 'a' is radius of given circle.

$$\text{From fig } 4a^2 = x^2 + y^2$$

$$\Rightarrow y^2 = 4a^2 - x^2$$

$$\Rightarrow y = \sqrt{4a^2 - x^2}$$

$$\text{Perimeter (P)} = 2x + 2y = 2\left(x + \sqrt{4a^2 - x^2}\right)$$

- (ii) We know that $P = 2\left(x + \sqrt{4a^2 - x^2}\right)$

Critical points to maximize perimeter $\frac{dP}{dx} = 0$

$$\Rightarrow \frac{dp}{dx} = 2\left(1 + \frac{1}{2\sqrt{4a^2 - x^2}}(-2x)\right) = 0$$

$$2\left(\frac{\sqrt{4a^2 - x^2} - x}{\sqrt{4a^2 - x^2}}\right) = 0$$

$$\Rightarrow \sqrt{4a^2 - x^2} = x$$

$$\Rightarrow 4a^2 - x^2 = x^2$$

$$\Rightarrow 2a^2 = x^2$$

$$\Rightarrow x = \pm\sqrt{2a}$$

$$\text{when } x = \sqrt{2a}, y = \sqrt{2a}$$

when $x = -\sqrt{2a}$ not possible as 'x' is length critical point is $(\sqrt{2a}, \sqrt{2a})$

(iii)

$$\frac{dp}{dx} = 2 \left(1 + \frac{1}{2\sqrt{4a^2 - x^2}}(-2x) \right)$$

$$\frac{d^2P}{dx^2} = -2 \left(\frac{\sqrt{4a^2 - x^2} - (x) \left(\frac{-2x}{2\sqrt{4a^2 - x^2}} \right)}{(4a^2 - x^2)} \right)$$

$$= -2 \left(\frac{(4a^2 - x^2) + x^2}{(4a^2 - x^2)^{3/2}} \right)$$

$$\Rightarrow \left. \frac{d^2P}{dx^2} \right]_{x=a\sqrt{2}} = -2 \left(\frac{4a^2}{(4a^2 - 2a^2)^{3/2}} \right) = \frac{-2}{(2\sqrt{2})a} < 0$$

Perimeter is maximum at a critical point.

OR

From the above results know that $x = y = \sqrt{2}a$

a = radius

Here, $x = y = 10\sqrt{2}$

Perimeter = $P = 4 \times \text{side} = 40\sqrt{2}$ cm

37. Read the text carefully and answer the questions:

Three schools A, B and C organized a mela for collecting funds for helping the rehabilitation of flood victims. They sold handmade fans, mats, and plates from recycled material at a cost of ₹ 25, ₹ 100 and ₹ 50 each. The number of articles sold by school A, B, C are given below.



Article	School	A	B	C
Fans		40	25	35
Mats		50	40	50
Plates		20	30	40

(i) *Fans Mats Plates*

$$P = \begin{matrix} A \\ B \\ C \end{matrix} \begin{bmatrix} 40 & 50 & 20 \\ 25 & 40 & 30 \\ 35 & 50 & 40 \end{bmatrix}$$

$$Q = \begin{bmatrix} 25 \\ 100 \\ 50 \end{bmatrix} \begin{matrix} \textit{Fans} \\ \textit{Mats} \\ \textit{Plates} \end{matrix}$$

(ii) Clearly, total funds collected by each school is given by the matrix

$$PQ = \begin{bmatrix} 40 & 50 & 20 \\ 25 & 40 & 30 \\ 35 & 50 & 40 \end{bmatrix} \begin{bmatrix} 25 \\ 100 \\ 50 \end{bmatrix}$$

$$= \begin{bmatrix} 1000 + 5000 + 1000 \\ 625 + 4000 + 1500 \\ 875 + 5000 + 2000 \end{bmatrix} = \begin{bmatrix} 7000 \\ 6125 \\ 7875 \end{bmatrix}$$

\therefore Funds collected by school A is ₹7000.

Funds collected by school B is ₹6125.

Funds collected by school C is ₹7875.

(iii)

$$\text{New price matrix } Q = 20\% \times \begin{bmatrix} 25 \\ 100 \\ 50 \end{bmatrix} \begin{matrix} \textit{Fans} \\ \textit{Mats} \\ \textit{Plates} \end{matrix}$$

$$\Rightarrow Q = \begin{bmatrix} 25 + 25 \times 0.20 \\ 100 + 100 \times 0.20 \\ 50 + 50 \times 0.20 \end{bmatrix} \begin{matrix} \textit{Fans} \\ \textit{Mats} \\ \textit{Plates} \end{matrix}$$

$$Q = \begin{bmatrix} 30 \\ 120 \\ 60 \end{bmatrix} \begin{matrix} \textit{Fans} \\ \textit{Mats} \\ \textit{Plates} \end{matrix}$$

$$\text{New price matrix } Q = 20\% \times \begin{bmatrix} 25 \\ 100 \\ 50 \end{bmatrix} \begin{matrix} \textit{Fans} \\ \textit{Mats} \\ \textit{Plates} \end{matrix}$$

$$\Rightarrow Q = \begin{bmatrix} 25 + 25 \times 0.20 \\ 100 + 100 \times 0.20 \\ 50 + 50 \times 0.20 \end{bmatrix} \begin{matrix} \text{Fans} \\ \text{Mats} \\ \text{Plates} \end{matrix}$$

$$Q = \begin{bmatrix} 30 \\ 120 \\ 60 \end{bmatrix} \begin{matrix} \text{Fans} \\ \text{Mats} \\ \text{Plates} \end{matrix}$$

OR

$$PQ = \begin{bmatrix} 40 & 50 & 20 \\ 25 & 40 & 30 \\ 35 & 50 & 40 \end{bmatrix} \begin{bmatrix} 30 \\ 120 \\ 60 \end{bmatrix}$$

$$PQ = \begin{bmatrix} 1200 + 6000 + 1200 \\ 750 + 4800 + 1800 \\ 1050 + 6000 + 2400 \end{bmatrix} = \begin{bmatrix} 8400 \\ 7350 \\ 9450 \end{bmatrix}$$

Total fund collected = 8400 + 7350 + 9450 = ₹25,200

38. Read the text carefully and answer the questions:

To teach the application of probability a maths teacher arranged a surprise game for 5 of his students namely Govind, Girish, Vinod, Abhishek and Ankit. He took a bowl containing tickets numbered 1 to 50 and told the students go one by one and draw two tickets simultaneously from the bowl and replace it after noting the numbers.



- (i) Required probability = P(one ticket with prime number and other ticket with a multiple of 4)

$$= 2 \left(\frac{15}{50} \times \frac{12}{49} \right) = \frac{36}{245}$$

- (ii) P(First ticket shows an even number and second ticket shows an odd number)

$$= \frac{25}{50} \times \frac{25}{49} = \frac{25}{98}$$