

Sample Question Paper - 10
Mathematics (041)
Class- XII, Session: 2021-22
TERM II

Time Allowed: 2 hours

Maximum Marks: 40

General Instructions:

1. This question paper contains three sections – A, B and C. Each part is compulsory.
2. Section - A has 6 short answer type (SA1) questions of 2 marks each.
3. Section – B has 4 short answer type (SA2) questions of 3 marks each.
4. Section - C has 4 long answer-type questions (LA) of 4 marks each.
5. There is an internal choice in some of the questions.
6. Q 14 is a case-based problem having 2 sub-parts of 2 marks each.

Section A

1. Prove that: $\int_0^{\pi/2} \frac{\cos^{1/4} x}{(\sin^{1/4} x + \cos^{1/4} x)} dx = \frac{\pi}{4}$ [2]

OR

Evaluate: $\int x^3 e^x dx$

2. Write the order and degree of the differential equation $y = x \frac{dy}{dx} + a \sqrt{1 + \left(\frac{dy}{dx}\right)^2}$. [2]
3. For what value of 'a' the vectors $2\hat{i} - 3\hat{j} + 4\hat{k}$ and $a\hat{i} + 6\hat{j} - 8\hat{k}$ are collinear? [2]
4. Find the cartesian and vector equations of the planes through the line of intersection of the planes $\vec{r} \cdot (\hat{i} - \hat{j}) + 6 = 0$ and $\vec{r} \cdot (3\hat{i} + 3\hat{j} - 4\hat{k}) = 0$ which are at a unit distance from the origin. [2]
5. A can solve 90% of the problems given in a book and B can solve 70%. What is the probability that at least one of them will solve the problem, selected at random from the book? [2]
6. An electronic assembly consists of two sub-systems say A and B. From previous testing procedures, the following probabilities are assumed to be known: [2]

$P(A \text{ fails}) = 0.2$

$P(B \text{ fails alone}) = 0.15$

$P(A \text{ and } B \text{ fail}) = 0.15$

Evaluate the following probabilities.

(1) $P(\overline{A}|\overline{B})$

(2) $P(A \text{ fails alone})$.

Section B

7. Evaluate: $\int \frac{(x^2+1)}{(x-1)^2(x+3)} dx$. [3]
8. Solve the following differential equation $\frac{dy}{dx} = 1 + x^2 + y^2 + x^2 y^2$, given that $y = 1$, when $x =$ [3]



0.

OR

Show that the family of curves for which the slope of the tangent at any point (x, y) on it is $\frac{dy}{dx} = \frac{x^2+y^2}{2xy}$, is given by $x^2 - y^2 = cx$.

9. Find the value of λ so that the four points A, B, C and D with position vectors $4\hat{i} + 5\hat{j} + \hat{k}$, $-\hat{j} - \hat{k}$, $3\hat{i} + \lambda\hat{j} + 4\hat{k}$ and $-4\hat{i} + 4\hat{j} + 4\hat{k}$, respectively are coplanar. [3]
10. Find the foot of perpendicular from the point (2, 3, -8) to the line $\frac{4-x}{2} = \frac{y}{6} = \frac{1-z}{3}$. Also, find the perpendicular distance from the given point to the line. [3]

OR

Find the distance of the point $(-1, -5, -10)$ from the point of intersection of the line $\vec{r} = 2\hat{i} - \hat{j} + 2\hat{k} + \lambda(3\hat{i} + 4\hat{j} + 2\hat{k})$ and the plane $\vec{r} \cdot (\hat{i} - \hat{j} + \hat{k}) = 5$

Section C

11. Evaluate: $\int \frac{(3 \sin x - 2) \cos x}{5 - \cos^2 x - 4 \sin x} dx$ [4]
12. Find the area common to the circle $x^2 + y^2 = 16$ and the parabola $y^2 = 6ax$. [4]

OR

Find the area enclosed by the parabola $4y = 3x^2$ and the line $2y = 3x + 12$

13. By computing the shortest distance determine whether the pairs of lines intersect or not: $\vec{r} = (\hat{i} - \hat{j}) + \lambda(2\hat{i} + \hat{k})$ and $\vec{r} = (2\hat{i} - \hat{j}) + \mu(\hat{i} + \hat{j} - \hat{k})$ [4]

CASE-BASED/DATA-BASED

14. The probability that a certain person will buy a shirt is 0.2, the probability that he will buy a trouser is 0.3, and the probability that he will buy a shirt given that he buys a trouser is 0.4. [4]



- i. Find the probability that he will buy both a shirt and a trouser.
- ii. Find also the probability that he will buy a trouser given that he buys a shirt.

Solution
MATHEMATICS 041
Class 12 - Mathematics

Section A

1. Let $y = \int_0^{\pi/2} \frac{\cos^{\frac{1}{4}} x}{\sin^{\frac{1}{4}} x + \cos^{\frac{1}{4}} x} dx \dots (i)$

Use King theorem of definite integral

$$\int_a^b f(x) dx = \int_a^b f(a+b-x) dx$$

$$y = \int_0^{\pi/2} \frac{\cos^{\frac{1}{4}} \left(\frac{\pi}{2} - x\right)}{\sin^{\frac{1}{4}} \left(\frac{\pi}{2} - x\right) + \cos^{\frac{1}{4}} \left(\frac{\pi}{2} - x\right)} dx$$

$$y = \int_0^{\pi/2} \frac{\sin^{\frac{1}{4}} x}{\sin^{\frac{1}{4}} x + \cos^{\frac{1}{4}} x} dx \dots (ii)$$

Adding eq.(i) and eq.(ii), we get

$$2y = \int_0^{\pi/2} \frac{\cos^{\frac{1}{4}} x}{\sin^{\frac{1}{4}} x + \cos^{\frac{1}{4}} x} dx + \int_0^{\pi/2} \frac{\sin^{\frac{1}{4}} x}{\sin^{\frac{1}{4}} x + \cos^{\frac{1}{4}} x} dx$$

$$2y = \int_0^{\pi/2} \frac{\cos^{\frac{1}{4}} x + \sin^{\frac{1}{4}} x}{\sin^{\frac{1}{4}} x + \cos^{\frac{1}{4}} x} dx$$

$$2y = \int_0^{\pi/2} 1 dx$$

$$2y = (x)_0^{\pi/2}$$

$$y = \frac{\pi}{4}$$

OR

Let $I = \int_I x^3 e^x dx$, then we have

$$I = x^3 e^x - \int 3x^2 e^x dx = x^3 e^x - 3 \int_I x^2 e^x dx$$

$$\Rightarrow I = x^3 e^x - 3 \{x^2 e^x - \int 2x e^x dx\} = x^3 e^x - 3 \{x^2 e^x - 2 \int_I x e^x dx\}$$

$$\Rightarrow I = x^3 e^x - 3 [x^2 e^x - 2 \{x e^x - \int 1 \cdot e^x dx\}]$$

$$\Rightarrow I = x^3 e^x - 3 x^2 e^x + 6x e^x - 6e^x + C$$

$$\Rightarrow I = (x^3 - 3x^2 + 6x - 6) e^x + C$$

2. We have

$$y = x \frac{dy}{dx} + a \sqrt{1 + \left(\frac{dy}{dx}\right)^2}$$

$$\left(y - x \frac{dy}{dx}\right) = a \sqrt{1 + \left(\frac{dy}{dx}\right)^2}$$

Squaring both the sides,

$$y^2 - x^2 \left(\frac{dy}{dx}\right)^2 - 2xy \left(\frac{dy}{dx}\right) = a^2 \left(1 + \left(\frac{dy}{dx}\right)^2\right)$$

$$x^2 \left(\frac{dy}{dx}\right)^2 - a^2 \left(\frac{dy}{dx}\right)^2 - 2xy \left(\frac{dy}{dx}\right) + y^2 - a^2 = 0$$

$$\left(\frac{dy}{dx}\right)^2 (x^2 - a^2) - 2xy \left(\frac{dy}{dx}\right) + y^2 - a^2 = 0$$

The highest order differential coefficient is 2 so,

Order of differential equation is 1

Degree of differential equation is 2

3. Here, it is given that two vectors, let $\vec{p} = 2\hat{i} - 3\hat{j} + 4\hat{k}$ and $\vec{q} = a\hat{i} + 6\hat{j} - 8\hat{k}$ are collinear

Since the given vectors are collinear, we have, $\vec{p} = \lambda \vec{q}$

$$\Rightarrow 2\hat{i} - 3\hat{j} + 4\hat{k} = \lambda(a\hat{i} + 6\hat{j} - 8\hat{k})$$

$$\Rightarrow 2\hat{i} - 3\hat{j} + 4\hat{k} = a\lambda\hat{i} + 6\lambda\hat{j} - 8\lambda\hat{k}$$

$$\Rightarrow \lambda a = 2, 6\lambda = -3 \text{ and } -8\lambda = 4$$

$$\Rightarrow \lambda = -\frac{1}{2} \text{ and } a = -4$$

4. Given, equations of plane are $(x\hat{i} + y\hat{j} + z\hat{k}) \cdot (\hat{i} - \hat{j}) + 6 = 0$ and $(x\hat{i} + y\hat{j} + z\hat{k}) \cdot (3\hat{i} + 3\hat{j} - 4\hat{k}) = 0$

$$\Rightarrow x - y + 6 = 0 \text{ and } 3x + 3y - 4z = 0$$

Any plane through their intersection is $(x - y + 6) + \lambda(3x + 3y - 4z) = 0$

$$\Rightarrow (1 + 3\lambda)x + (3\lambda - 1)y + 4\lambda z + 6 = 0$$

$$\therefore \frac{6}{\sqrt{(1+3\lambda)^2 + (3\lambda-1)^2 + (-4\lambda)^2}} = 1$$

$$\Rightarrow 34\lambda^2 + 2 = 36$$

$$\Rightarrow \lambda^2 = 1 \Rightarrow \lambda = \pm 1$$

Therefore, the required planes are $2x + y - 2z + 3 = 0$ and $x + 2y - 2z - 3 = 0$ In vector form they are

$$\vec{r} \cdot (2\hat{i} + \hat{j} - 2\hat{k}) + 3 = 0 \text{ and } \vec{r} \cdot (\hat{i} + 2\hat{j} - 2\hat{k}) - 3 = 0.$$

5. Let E and F be the events defined as follows:

E = A solves the problem, F = B solves the problem.

Clearly, E and F are independent events such that

$$P(E) = \frac{90}{100} = \frac{9}{10} \text{ and } P(F) = \frac{70}{100} = \frac{7}{10}$$

Problem will be solved if atleast one of them solves it

Therefore, required probability = $P(E \cup F)$

$$= P(E) + P(F) - P(E \cap F)$$

$$= P(E) + P(F) - P(E)P(F) \text{ [as E and F are independent]}$$

$$= P(E) + P(F) [1 - P(E)]$$

$$= \frac{9}{10} + \frac{7}{10} [1 - \frac{9}{10}]$$

$$= \frac{9}{10} + \frac{7}{10} \times \frac{1}{10} = \frac{97}{100} = 0.97$$

Which is the required solution.

6. Event A fails and B fails denoted by \bar{A} and \bar{B} respectively.

$$\therefore P(\bar{A}) = 0.2 \text{ and } P(A \text{ and } B \text{ fails}) = 0.15$$

$$\Rightarrow P(A \cap B) = 0.15$$

$$\therefore P(\bar{B} \text{ above}) = P(\bar{B}) - P(A \cap B)$$

$$\Rightarrow 0.15 = P(\bar{B}) - 0.15$$

$$\Rightarrow P(\bar{B}) = 0.30$$

$$\text{i. } P(\bar{A}|\bar{B}) = \frac{P(\bar{A} \cap \bar{B})}{P(\bar{B})} = \frac{0.15}{0.30} = \frac{1}{2} = 0.5$$

$$\text{ii. } P(A \text{ fails alone}) = P(\bar{A} \text{ alone}) = P(\bar{A}) - P(\bar{A} \cap \bar{B}) = 0.20 - 0.15 = 0.05$$

Section B

7. Let the given integral be, $I = \int \frac{x^2+1}{(x+3)(x-1)^2} dx$

Now using partial fractions by putting, $\frac{x^2+1}{(x+3)(x-1)^2} = \frac{A}{(x+3)} + \frac{B}{(x-1)} + \frac{C}{(x-1)^2} \dots (1)$

$$A(x-1)^2 + B(x+3)(x-1) + C(x+3) = x^2 + 1$$

Now put $x - 1 = 0$

Therefore, $x = 1$

$$A(0) + B(0) + C(4) = 2$$

$$c = \frac{1}{2}$$

Now put $x + 3 = 0$

Therefore, $x = -3$

$$A(-3-1)^2 + B(0) + C(0) = 9 + 1 = 10$$

$$A = \frac{5}{8}$$

By equating the coefficient of x^2 , we get, $A + B = 1$

$$\frac{5}{8} + B = 1$$

$$B = 1 - \frac{5}{8} = \frac{3}{8}$$

From equation (1), we get,

$$\begin{aligned}\frac{x^2+1}{(x+3)(x-2)^2} &= \frac{5}{8} \times \frac{1}{(x+3)} + \frac{3}{8} \times \frac{1}{(x-2)} + \frac{1}{(x-2)^2} \\ \int \frac{x^2+1}{(x+3)(x-2)^2} dx &= \frac{5}{8} \int \frac{1}{(x+3)} dx + \frac{3}{8} \int \frac{1}{(x-2)} dx + \int \frac{1}{(x-2)^2} dx \\ &= \frac{5}{8} \log|x+3| + \frac{3}{8} \log|x-2| - \frac{1}{2(x-2)} + C\end{aligned}$$

8. According to the question ,

Given differential equation is ,

$$\frac{dy}{dx} = 1 + x^2 + y^2 + x^2 y^2$$

$$\Rightarrow \frac{dy}{dx} = 1(1+x^2) + y^2(1+x^2)$$

$$\Rightarrow \frac{dy}{dx} = (1+x^2)(1+y^2)$$

$$\Rightarrow \frac{dy}{1+y^2} = (1+x^2) dx$$

On integrating both sides, we get

$$\int \frac{dy}{1+y^2} = \int (1+x^2) dx$$

$$\Rightarrow \tan^{-1} y = x + \frac{x^3}{3} + C \dots (i)$$

Given that $y = 1$, when $x = 0$.

On putting $x = 0$ and $y = 1$ in Eq. (i), we get

$$\tan^{-1} 1 = C$$

$$\Rightarrow \tan^{-1}(\tan \pi/4) = C \quad [\because \tan \frac{\pi}{4} = 1]$$

$$\Rightarrow C = \frac{\pi}{4}$$

On putting the value of C in Eq. (i), we get

$$\tan^{-1} y = x + \frac{x^3}{3} + \frac{\pi}{4}$$

$$\therefore y = \tan\left(x + \frac{x^3}{3} + \frac{\pi}{4}\right)$$

which is the required solution of differential equation.

OR

$$\text{We have, } \frac{dy}{dx} = \frac{x^2+y^2}{2xy}$$

Clearly, each of the function $x^2 + y^2$ and $2xy$ is a homogeneous function of degree 2, so the given equation is homogeneous.

$$\text{Put } y = vx \text{ and } \frac{dy}{dx} = v + x \frac{dv}{dx}$$

The given equation becomes

$$v + x \frac{dv}{dx} = \frac{x^2+v^2x^2}{2vx^2}$$

$$\Rightarrow v + x \frac{dv}{dx} = \frac{v^2+1}{2v}$$

$$\Rightarrow x \frac{dv}{dx} = \left(\frac{v^2+1}{2v} - v\right)$$

$$\Rightarrow x \frac{dv}{dx} = \frac{v^2+1-2v^2}{2v} = \frac{1-v^2}{2v}$$

$$\Rightarrow x \frac{dv}{dx} = \frac{-(v^2-1)}{2v} \Rightarrow -\frac{2v}{v^2-1} dv = \frac{dx}{x} \quad [\text{using variable separable form}]$$

On integrating both sides, we get

$$-\log|v^2-1| = \log x - \log C_1$$

$$\Rightarrow -\log|v^2-1| - \log x = -\log C_1$$

$$\Rightarrow \log|x(v^2-1)| = \log C_1 \Rightarrow x(v^2-1) = C_1$$

$$\Rightarrow x\left(\frac{y^2}{x^2}-1\right) = C_1 \Rightarrow x\left(\frac{y^2-x^2}{x^2}\right) = C_1$$

$$\Rightarrow \frac{y^2 - x^2}{x} = C_1 \Rightarrow x^2 - y^2 = -C_1 x$$

$$\Rightarrow x^2 - y^2 = Cx \quad [\because C = -C_1]$$

9. According to the question,

Given, $\vec{OA} = 4\hat{i} + 5\hat{j} + \hat{k}$,

$\vec{OB} = -\hat{j} - \hat{k}$,

$\vec{OC} = 3\hat{i} + \lambda\hat{j} + 4\hat{k}$ and

$\vec{OD} = -4\hat{i} + 4\hat{j} + 4\hat{k}$.

Now, $\vec{AB} = \vec{OB} - \vec{OA} = -\hat{j} - \hat{k} - (4\hat{i} + 5\hat{j} + \hat{k})$

$= -4\hat{i} - 6\hat{j} - 2\hat{k}$

$\vec{AC} = \vec{OC} - \vec{OA} = 3\hat{i} + \lambda\hat{j} + 4\hat{k} - (4\hat{i} + 5\hat{j} + \hat{k})$

$= -\hat{i} + (\lambda - 5)\hat{j} + 3\hat{k}$

and $\vec{AD} = \vec{OD} - \vec{OA}$

$= -4\hat{i} + 4\hat{j} + 4\hat{k} - (4\hat{i} + 5\hat{j} + \hat{k})$

$= -8\hat{i} - \hat{j} + 3\hat{k}$

Since, vectors $\vec{OA}, \vec{OB}, \vec{OC}$ and \vec{OD} are coplanar.

$\therefore [\vec{AB} \vec{AC} \vec{AD}] = 0$

$\therefore \begin{vmatrix} -4 & -6 & -2 \\ -1 & (\lambda - 5) & 3 \\ -8 & -1 & 3 \end{vmatrix} = 0$

$-4(3\lambda - 15 + 3) + 6(-3 + 24) - 2(1 + 8\lambda - 40) = 0$

$\Rightarrow -4(3\lambda - 12) + 6(21) - 2(8\lambda - 39) = 0$

$\Rightarrow -12\lambda + 48 + 126 - 16\lambda + 78 = 0$

$\Rightarrow -28\lambda + 252 = 0$

$\Rightarrow \lambda = 9$

10. We have equation of line is $\frac{4-x}{2} = \frac{y}{6} = \frac{1-z}{3}$

$\Rightarrow \frac{x-4}{-2} = \frac{y}{6} = \frac{z-1}{-3} = \lambda$

$\Rightarrow x = -2\lambda + 4, y = 6\lambda$ and $z = -3\lambda + 1$

Let the coordinates of L be $(4 - 2\lambda, 6\lambda, 1 - 3\lambda)$, then, direction ratios of PL are proportional to $(4 - 2\lambda - 2, 6\lambda - 3, 1 - 3\lambda + 8)$ i.e., $(2 - 2\lambda, 6\lambda - 3, 9 - 3\lambda)$.


Also, direction ratios are proportional to -2, 6, -3. Since, PL is perpendicular to give line.

$\therefore -2(2 - 2\lambda) + 6(6\lambda - 3) - 3(9 - 3\lambda) = 0$

$\Rightarrow -4 + 4\lambda + 36\lambda - 18 - 27 + 9\lambda = 0$

$\Rightarrow 49\lambda = 49 \Rightarrow \lambda = 1$

So, the coordinates of L are $(4 - 2\lambda, 6\lambda, 1 - 3\lambda)$ i.e., $(2, 6, -2)$.



Also, length of PL $= \sqrt{(2-2)^2 + (6-3)^2 + (-2+8)^2}$
 $= \sqrt{0 + 9 + 36} = 3\sqrt{5} \text{ units}$

OR

Given: A point P (say) $(-1, -5, -10)$

and equation of the line $\vec{r} = 2\hat{i} - \hat{j} + 2\hat{k} + \lambda(3\hat{i} + 4\hat{j} + 2\hat{k}) \dots(i)$

equation of the plane is $\vec{r} \cdot (\hat{i} - \hat{j} + \hat{k}) = 5$

Putting the value of \vec{r} from eq. (i) in eq. (ii),

$$\left[(2\hat{i} - \hat{j} + 2\hat{k}) + \lambda(3\hat{i} + 4\hat{j} + 2\hat{k}) \right] \cdot (\hat{i} - \hat{j} + \hat{k}) = 5$$

$$\Rightarrow (2\hat{i} - \hat{j} + 2\hat{k}) \cdot (\hat{i} - \hat{j} + \hat{k}) + \lambda(3\hat{i} + 4\hat{j} + 2\hat{k}) \cdot (\hat{i} - \hat{j} + \hat{k}) = 5$$

$$\Rightarrow 2 + 1 + 2 + \lambda(3 - 4 + 2) = 5$$

$$\Rightarrow 5 + \lambda = 5$$

$$\Rightarrow \lambda = 0$$

Putting $\lambda = 0$ in eq. (i), $\vec{r} = 2\hat{i} - \hat{j} + 2\hat{k} + 0(3\hat{i} + 4\hat{j} + 2\hat{k})$

$$\Rightarrow \vec{r} = 2\hat{i} - \hat{j} + 2\hat{k}$$

Therefore, Point of intersection is $(-2, 1, 2)$

\therefore Distance of the given point $P(-1, -5, -10)$ from the point of intersection is

$$\sqrt{(2+1)^2 + (-1+5)^2 + (2+10)^2}$$

$$= \sqrt{9 + 16 + 144}$$

$$= \sqrt{169} = 13 \text{ units}$$

Section C

11. Let the given integral be

$$I = \int \frac{(3 \sin x - 2) \cos x}{5 - \cos^2 x - 4 \sin x} dx$$

$$\therefore I = \int \frac{(3 \sin x - 2) \cos x}{5 - (1 - \sin^2 x) - 4 \sin x} dx$$

$$\Rightarrow 1 = \int \frac{(3 \sin x - 2) \cos x}{5 - 1 + \sin^2 x - 4 \sin x} dx$$

Substitute $\sin x = t$

$$\Rightarrow \cos x dx = dt$$

Thus,

$$I = \int \frac{(3t-2)}{4+t^2-4t} dt$$

$$\Rightarrow I = \int \frac{(3t-2)}{t^2-4t+4} dt$$

$$\Rightarrow I = \int \frac{(3t-2)}{(t-2)^2} dt$$

Now let us separate the integrand into the simplest form using partial fractions.

$$\frac{(3t-2)}{(t-2)^2} = \frac{A}{(t-2)} + \frac{B}{(t-2)^2}$$

$$= \frac{A(t-2)+B}{(t-2)^2}$$

$$= \frac{At-2A+B}{(t-2)^2}$$

$$= 3t - 2 = At - 2A + B$$

Comparing the coefficients, we have,

$$A = 3 \text{ and } -2A + B = -2$$

Substituting the value of $A = 3$ in the above equation, we have,

$$\Rightarrow -2 \times 3 + B = -2$$

$$\Rightarrow -6 + B = -2$$

$$\Rightarrow B = 6 - 2$$

$$\Rightarrow B = 4$$

Therefore we have, $I = \int \frac{(3t-2)}{(t-2)^2} dt$ becomes

$$I = \int \frac{3}{(t-2)} dt + \int \frac{4}{(t-2)^2} dt$$

$$= 3 \log |t-2| - 4 \left(\frac{1}{t-2} \right) + C$$

$$= 3 \log |2-t| + 4 \left(\frac{1}{2-t} \right) + C$$

Now substituting $t = \sin x$, we have,

$$I = 3 \log |2 - \sin x| + 4 \left(\frac{1}{2 - \sin x} \right) + C$$

12. To find: Area enclosed by


$$x^2 + y^2 = 16 \dots (i)$$

$$\text{and } y^2 = 6ax \dots (ii)$$

Equation (i) represents a circle with centre (0, 0) and meets X-axis $(\pm 4a, 0)$.

Equation (ii) represents a parabola with vertex (0, 0) and axis as x-axis, Points of intersection of circle and parabola are $(2a, 2\sqrt{3}a), (2a, -2\sqrt{3}a)$.

A rough sketch of curves is given as:-



Required ODCO is sliced into rectangles of area $y_1 \Delta x$ and it slides from $x = 0$ to $x = 2a$.

Region BCDB is sliced into rectangle of area $y_2 \Delta x$ it slides from $x = 2a$ to $x = 4a$. So,

Required area = 2 [Region ODCO + Region BCDB]

$$= 2 \left[\int_0^{2a} y_1 dx + \int_{2a}^{4a} y_2 dx \right]$$

$$= 2 \left[\int_0^{2a} \sqrt{6ax} dx + \int_{2a}^{4a} \sqrt{16a^2 - x^2} dx \right]$$

$$= 2 \left[\sqrt{6a} \left(\frac{2}{3} x \sqrt{x} \right) \right]_0^{2a} + \left[\frac{x}{2} \sqrt{16a^2 - x^2} + \frac{16a^2}{2} \sin^{-1} \left(\frac{x}{4a} \right) \right]_{2a}^{4a}$$

$$= \frac{2\sqrt{2}}{\sqrt{3}} a \cdot 2a \cdot \sqrt{2a} + aa^2 \cdot \sin^{-1}(1) - \frac{2a}{2} \cdot \sqrt{12a^2} - 8a^2 \cdot \sin^{-1} \left(\frac{1}{2} \right)$$

$$= 2 \left[\left(\sqrt{6a} \cdot \frac{2}{3} 2a \sqrt{2a} \right) + \left[\left(0 + 8a^2 \cdot \frac{\pi}{2} \right) - \left(a\sqrt{12a^2} + 8a^2 \cdot \frac{\pi}{6} \right) \right] \right]$$

$$= 2 \left[\frac{8\sqrt{3}a^2}{3} + 4a^2\pi - 2\sqrt{3}a^2 - \frac{4}{3}a^2\pi \right]$$


$$= 2 \left[\frac{2\sqrt{3}a^2}{3} + \frac{8a^2\pi}{3} \right]$$

$$A = \frac{4a^2}{3} (4\pi + \sqrt{3}) \text{ sq. units.}$$

OR

Equation of the parabola is

$$4y = 3x^2 \dots (i)$$



$$\Rightarrow x^2 = \frac{4}{3}y$$

Equation of the line is $2y = 3x + 12 \dots(ii)$

$$\Rightarrow y = \frac{3x+12}{2} = \frac{3x}{2} + 6$$

In the graph, points of intersection are B (4, 12) and C (-2, 3).

$$\text{Now, Area ABCD} = \left| \int_{-2}^4 \left(\frac{3}{2}x + 6 \right) dx \right|$$

$$= \left[\frac{3}{4}x^2 + 6x \right]_{-2}^4$$

$$= (12 + 24) - (3 - 12)$$

$$= 45 \text{ sq units}$$

$$\text{Again, Area CDO} + \text{Area OAB} = \left| \int_{-2}^4 \left(\frac{3}{4}x^2 \right) dx \right|$$

$$= \left[\frac{3}{4} \cdot \frac{x^3}{3} \right]_{-2}^4$$

$$= \frac{1}{4}[64 - (-8)] = 18 \text{ sq. units}$$

\therefore Required area = Area ABCD - (Area CDO + Area OAB)

$$= 45 - 18 = 27 \text{ sq. units}$$

13. Equation of line in vector form

$$\text{Line I: } \vec{r} = (\hat{i} - \hat{j} + 0\hat{k}) + \lambda(2\hat{i} + 0\hat{j} + \hat{k})$$

$$\text{Line II: } \vec{r} = (2\hat{i} - \hat{j}) + \mu(\hat{i} + \hat{j} - \hat{k})$$

Here,

$$\vec{a}_1 = \hat{i} - \hat{j} + 0\hat{k}$$

$$\vec{a}_2 = 2\hat{i} - \hat{j}$$

$$\vec{b}_1 = 2\hat{i} + 0\hat{j} + \hat{k}$$

$$\vec{b}_2 = \hat{i} + \hat{j} - \hat{k}$$

We know that the shortest distance between lines is

$$d = \frac{|(\vec{a}_2 - \vec{a}_1)(\vec{b}_1 \times \vec{b}_2)|}{|\vec{b}_1 \times \vec{b}_2|}$$

$$(\vec{a}_2 - \vec{a}_1) = (2\hat{i} - \hat{j}) - (\hat{i} - \hat{j} + 0\hat{k})$$

$$(\vec{a}_2 - \vec{a}_1) = \hat{i} + 0\hat{j} + 0\hat{k}$$

$$\vec{b}_1 \times \vec{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 0 & 1 \\ 1 & 1 & -1 \end{vmatrix}$$

$$\vec{b}_1 \times \vec{b}_2 = (0 - 1)\hat{i} - (-2 - 1)\hat{j} + (2 - 0)\hat{k}$$

$$\Rightarrow \vec{b}_1 \times \vec{b}_2 = -\hat{i} + 3\hat{j} + 2\hat{k}$$

$$|\vec{b}_1 \times \vec{b}_2| = \sqrt{(-1)^2 + 3^2 + 2^2}$$

$$\Rightarrow |\vec{b}_1 \times \vec{b}_2| = \sqrt{14}$$

$$|(\vec{a}_2 - \vec{a}_1)(\vec{b}_1 \times \vec{b}_2)| = |(\hat{i} + 0\hat{j} + 0\hat{k})(-\hat{i} + 3\hat{j} + 2\hat{k})|$$

$$\Rightarrow |(\vec{a}_2 - \vec{a}_1)(\vec{b}_1 \times \vec{b}_2)| = 1$$

Substituting these values in the expression,

$$d = \frac{|(\vec{a}_2 - \vec{a}_1)(\vec{b}_1 \times \vec{b}_2)|}{|\vec{b}_1 \times \vec{b}_2|}$$

$$d = \frac{1}{\sqrt{14}}$$

$$d = \frac{1}{\sqrt{14}} \text{ units}$$

Shortest distance d between the lines is not 0. Hence the given lines are not intersecting.

CASE-BASED/DATA-BASED

14. Let S = Shirt, T = Trouser

$$P(S) = 0.2, P(T) = 0.3 \text{ and } P\left(\frac{S}{T}\right) = 0.4$$

We need to find $P(S \cap T)$ and $P\left(\frac{T}{S}\right)$

$$\text{We know, } P\left(\frac{S}{T}\right) = \frac{P(S \cap T)}{P(T)}$$

From given data, $0.4 = P(S \cap T) / 0.3$

$$P(S \cap T) = 0.4 \times 0.3 = 0.12$$

Also, we have,

$$P\left(\frac{T}{S}\right) = \frac{P(T \cap S)}{P(S)} = \frac{0.12}{0.2} = \frac{12}{20} = \frac{6}{10} = \frac{3}{5} = 0.6$$

