

Sample Question Paper - 5
Mathematics (041)
Class- XII, Session: 2021-22
TERM II

Time Allowed: 2 hours

Maximum Marks: 40

General Instructions:

1. This question paper contains three sections – A, B and C. Each part is compulsory.
2. Section - A has 6 short answer type (SA1) questions of 2 marks each.
3. Section – B has 4 short answer type (SA2) questions of 3 marks each.
4. Section - C has 4 long answer-type questions (LA) of 4 marks each.
5. There is an internal choice in some of the questions.
6. Q 14 is a case-based problem having 2 sub-parts of 2 marks each.

Section A

1. Evaluate: $\int \cos^3 x \sin 2x \, dx$. [2]

OR

Evaluate: $\int (x + 1) \log x \, dx$

2. Solve differential equation: $\frac{dy}{dx} - y \cot x = \operatorname{cosec} x$ [2]
3. Find the value of a for which the vector $\vec{r} = (a^2 - 4)\hat{i} + 2\hat{j} - (a^2 - 9)\hat{k}$ make acute angles with the coordinate axes. [2]
4. Find the equation of a plane passing through the point $P(6, 5, 9)$ and parallel to the plane determined by the points $A(3, -1, 2)$, $B(5, 2, 4)$ and $(-1, -1, 6)$. Also, find the distance of this plane from the point A . [2]
5. If A and B are two independent events such that $P(\bar{A} \cap B) = \frac{2}{15}$ and $P(A \cap \bar{B}) = \frac{1}{6}$, then find $P(B)$. [2]
6. If two events A and B are such that $P(\bar{A}) = 0.3$, $P(B) = 0.4$ and $P(A \cap \bar{B}) = 0.5$, find $P(\frac{B}{A \cap B})$ [2]

Section B

7. Evaluate the integral: $\int \frac{1}{x\sqrt{1+x^n}} dx$ [3]
8. Verify that $y^2 = 4a(x + a)$ is a solution of the differential equation $y \left\{ 1 - \left(\frac{dy}{dx} \right)^2 \right\} = 2x \frac{dy}{dx}$. [3]

OR

Find one-parameter families of solution curves of the differential equation: $x \frac{dy}{dx} - y = (x + 1) e^{-x}$

9. If $\vec{a} = \hat{i} + \hat{j} + 2\hat{k}$ and $\vec{b} = 2\hat{i} + \hat{j} - 2\hat{k}$, find the unit vector in the direction of $6\vec{b}$. [3]
10. Find the vector and Cartesian equations of the plane passing through the point $(3, -1, 2)$ and parallel to the lines $\vec{r} = (-\hat{j} + 3\hat{k}) + \lambda(2\hat{i} - 5\hat{j} - \hat{k})$ and $\vec{r} = (\hat{i} - 3\hat{j} + \hat{k}) + \mu(-5\hat{i} + 4\hat{j})$. [3]

OR



Prove that the line through A (0, -1, -1) and B (4, 5, 1) intersects the line through C (3, 9, 4) and D (-4, 4, 4).

Section C

11. Prove that: $\int_0^{\pi/2} x \cot x dx = \frac{\pi}{4}(\log 2)$. [4]

12. Find the area of circle $4x^2 + 4y^2 = 9$ which is interior to the parabola $x^2 = 4y$. [4]

OR

Find the area between the curves $y = x$ and $y = x^2$

13. Find the vector equation of the line passing through the point (1, 2, 3) and parallel to the planes $\vec{r} \cdot (\hat{i} - \hat{j} + 2\hat{k}) = 5$ and $\vec{r} \cdot (3\hat{i} + \hat{j} + \hat{k}) = 6$. [4]

CASE-BASED/DATA-BASED

14. In pre-board examination of class XII, commerce stream with Economics and Mathematics of a particular school, 50% of the students failed in Economics, 35% failed in Mathematics and 25% failed in both Economics and Mathematics. A student is selected at random from the class. [4]



Based on the above information, answer the following questions.

- The probability that the selected student has failed in Economics, if it is known that he has failed in Mathematics?
- The probability that the selected student has failed in Mathematics, if it is known that he has failed in Economics?



Solution
MATHEMATICS BASIC 041
Class 12 - Mathematics

Section A

1. Here,

$$I = \int \sin 2x \cos^3 x \, dx$$

$$\Rightarrow \int 2 \sin x \cos x \cos^2 x \, dx$$

$$\Rightarrow \int 2 \sin x \cos^4 x \, dx$$

Now put $\cos x = t$

$$\Rightarrow -\sin x \, dx = dt$$

$$\Rightarrow -2 \int t^4 \, dt$$

$$\Rightarrow -2 \times \frac{t^5}{5} + c$$

Re-substituting the value of $t = \cos x$ we get,

$$\Rightarrow \frac{-2 \cos^5 x}{5} + c$$

OR

Let $I = \int (x+1) \log x \, dx$, then we have

$$I = \log x \int (x+1) \, dx - \int \left(\frac{1}{x}\right) (x+1) \, dx$$

$$= \left(\frac{x^2}{2} + x\right) \log x - \int \frac{1}{x} \left(\frac{x^2}{2} + x\right) \, dx$$

$$= \left(\frac{x^2}{2} + x\right) \log x - \frac{1}{2} \int x \, dx - \int dx$$

$$= \left(\frac{x^2}{2} + x\right) \log x - \frac{1}{2} \times \frac{x^2}{2} - x + C$$

$$I = \left(\frac{x^2}{2} + x\right) \log x - \left(\frac{x^2}{4} + x\right) + C$$

2. Given that $\frac{dy}{dx} - y \cot x = \operatorname{cosec} x$

It is linear differential equation.

Comparing it with $\frac{dy}{dx} + py = Q$

$$P = -\cot x, Q = \operatorname{cosec} x$$

$$\text{I.F.} = e^{\int P \, dx}$$

$$= e^{-\int \cot x \, dx}$$

$$= e^{-|\log|\sin x|}$$

$$= \operatorname{cosec} x$$

Solution of the given equation is given by,

$$y (\text{I.F.}) = \int Q \times (\text{I.F.}) \, dx + c$$

$$y \operatorname{cosec} x = \int \operatorname{cosec} x \times \operatorname{cosec} x \, dx + c$$

$$y \operatorname{cosec} x = \int \csc^2 x \, dx + c$$

$$y \operatorname{cosec} x = -\cot x + c$$

3. We know that, For vector \vec{r} to be inclined with acute angles with the coordinate axes, we must have,

$$\vec{r} \cdot \hat{i} > 0, \vec{r} \cdot \hat{j} > 0 \text{ and } \vec{r} \cdot \hat{k} > 0$$

$$\Rightarrow \vec{r} \cdot \hat{i} > 0 \text{ and } \vec{r} \cdot \hat{k} > 0 \left[\because \vec{r} \cdot \hat{j} = 2 > 0 \right]$$

$$\Rightarrow (a^2 - 4) > 0 \text{ and } -(a^2 - 9) > 0 \left[\because \vec{r} \cdot \hat{i} = a^2 - 4 \text{ and } \vec{r} \cdot \hat{k} = -(a^2 - 9) \right]$$

$$\Rightarrow (a - 2)(a + 2) > 0 \text{ and } (a + 3)(a - 3) < 0$$

$$\Rightarrow a < -2 \text{ or } a > 2 \text{ and } -3 < a < 3$$

$$\Rightarrow a \in (-3, -2) \cup (2, 3)$$

4. We have a vector \vec{n} normal to the plane determined by the points A (3, -1, 2), B(5, 2, 4) and C(-1, -1, 6) is given

$$\text{by } \vec{n} = \vec{AB} \times \vec{AC}$$



$$\therefore \vec{n} = \vec{AB} \times \vec{AC} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 3 & 2 \\ -4 & 0 & 4 \end{vmatrix} = 12\hat{i} - 16\hat{j} + 12\hat{k}$$

Clearly, $\vec{n} = 12\hat{i} - 16\hat{j} + 12\hat{k}$ is also normal to the plane passing through P(6,5, 9) and parallel to the plane determined by point A, B and C. So, its equation is $\vec{r} \cdot \vec{n} = \vec{a} \cdot \vec{n}$ and $\vec{a} = 6\hat{i} + 5\hat{j} + 9\hat{k}$

$$\text{or, } \vec{r} \cdot (12\hat{i} - 16\hat{j} + 12\hat{k}) = (12\hat{i} - 16\hat{j} + 12\hat{k}) \cdot (6\hat{i} + 5\hat{j} + 9\hat{k})$$

$$\text{or, } \vec{r} \cdot (12\hat{i} - 16\hat{j} + 12\hat{k}) = 72 - 80 + 108$$

$$\text{or, } \vec{r} \cdot (3\hat{i} - 4\hat{j} + 3\hat{k}) = 25$$

The cartesian equation of this plane is $3x - 4y + 3z = 25$

Hence The required distance d of this plane from point A(3, -1, 2) is given by

$$d = \left| \frac{3 \times 3 - 4 \times -1 + 3 \times 2 - 25}{\sqrt{9+16+9}} \right| = \frac{6}{\sqrt{34}}$$

5. Let: $P(A) = x$, $P(B) = y$

$$P(\bar{A} \cap B) = \frac{2}{15}$$

$$\Rightarrow P(\bar{A}) \times P(B) = \frac{2}{15}$$

$$\Rightarrow (1 - x)y = \frac{2}{15} \dots(i)$$

$$P(A \cap \bar{B}) = \frac{1}{6}$$

$$\Rightarrow P(A) \times P(B) = \frac{1}{6}$$

$$\Rightarrow (1 - y)x = \frac{1}{6} \dots(ii)$$

subtracting (i) from (ii), we get,

$$x - y = \frac{1}{30}$$

$$x = y + \frac{1}{30}$$

putting the value of x in (ii), we have,

$$\left(y + \frac{1}{30}\right)(1 - y) = \frac{1}{6}$$

$$\Rightarrow 30y^2 - 29y + 4 = 0$$

$$\Rightarrow y = \frac{1}{6}, \frac{4}{5}$$

6. According to Baye's Theorem

$$P\left(\frac{B}{\bar{A} \cap \bar{B}}\right) = \frac{P(B \cap (\bar{A} \cap \bar{B}))}{P(\bar{A} \cap \bar{B})}$$

$$= \frac{P(B \cap (\overline{A \cup B}))}{P(\bar{A} \cap \bar{B})}$$

$$= \frac{P(\bar{B} \cap (A \cup B))}{P(\bar{A} \cap \bar{B})}$$

$$= \frac{P(\bar{A} \cap B)}{P(B \cup (A \cup B))}$$

$$= \frac{P(\bar{A} \cap B)}{P(\bar{A} \cap B)}$$

$$\text{Now } \bar{B} \cup B = \bar{U} = \phi$$

$$\text{So } P(\bar{B} \cup (A \cup B)) = \phi$$

$$\text{Therefore } P\left(\frac{B}{\bar{A} \cap \bar{B}}\right) = 0$$

Section B

7. Let the given integral be,

$$I = \int \frac{dx}{x\sqrt{1+x^n}}$$

$$= \int \frac{x^{n-1}dx}{x^{n-1}x^1\sqrt{1+x^n}}$$

$$= \int \frac{x^{n-1}dx}{x^n\sqrt{1+x^n}}$$

Putting $x^n = t$

$$\Rightarrow n x^{n-1} dx = dt$$

$$\Rightarrow x^{n-1} dx = \frac{dt}{n}$$

$$\therefore I = \frac{1}{n} \int \frac{dt}{t\sqrt{1+t}}$$

$$\text{let } 1 + t = p^2$$

$$\begin{aligned}
&\Rightarrow dp = 2p \, dp \\
&\therefore I = \frac{1}{n} \int \frac{2p \, dp}{(p^2-1)p} \\
&= \frac{2}{n} \int \frac{dp}{p^2-1^2} \\
&= \frac{2}{n} \times \frac{1}{2} \log \left| \frac{p-1}{p+1} \right| + C \\
&= \frac{1}{n} \log \left| \frac{\sqrt{1+t}-1}{\sqrt{1+t}+1} \right| + C \\
&= \frac{1}{n} \log \left| \frac{\sqrt{1+x^n}-1}{\sqrt{1+x^n}+1} \right| + C
\end{aligned}$$

8. The given functional relation is,

$$y^2 = 4a(x + a)$$

Differentiating above equation with respect to x

$$2y \frac{dy}{dx} = 4a$$

Substituting above results in

$$y \left(1 - \left(\frac{dy}{dx} \right)^2 \right) = 2x \frac{dy}{dx}, \text{ we get,}$$

$$y \left(1 - \left(\frac{dy}{dx} \right)^2 \right) = 4 \frac{ax}{y}$$

$$\Rightarrow y - \frac{4a^2}{y} = 4 \frac{ax}{y}$$

$$\Rightarrow \frac{y^2 - 4a(a+x)}{y} = 0$$

$$\Rightarrow \frac{4a(a+x) - 4a(a+x)}{y} = 0$$

$$\Rightarrow 0 = 0, \text{ which is true.}$$

$$\therefore y^2 = 4a(x + a) \text{ is the solution of } y \left(1 - \left(\frac{dy}{dx} \right)^2 \right) = 2x \frac{dy}{dx}$$

OR

The given differential equation is,

$$x \frac{dy}{dx} - y = (x+1) e^{-x}$$

$$\frac{dy}{dx} - \frac{y}{x} = \left(\frac{x+1}{x} \right) e^{-x}$$

It is a linear differential equation. Comparing it with,

$$\frac{dy}{dx} + Py = Q$$

$$P = -\frac{1}{x}, Q = \left(\frac{x+1}{x} \right) e^{-x}$$

$$\text{I.F.} = e^{\int P dx}$$

$$= e^{-\int \frac{1}{x} dx}$$

$$= e^{-\log |x|}$$

$$= e^{\log \left(\frac{1}{x} \right)}$$

$$= \frac{1}{x}, x > 0$$

Solution of the equation is given by,

$$y \times (\text{I.F.}) = \int Q \times (\text{I.F.}) dx + c$$

$$y \times \left(\frac{1}{x} \right) = \int \left(\frac{x+1}{x} \right) e^{-x} \times \left(\frac{1}{x} \right) dx + c$$

$$\frac{y}{x} = \int \left(\frac{1}{x} + \frac{1}{x^2} \right) e^{-x} dx + c$$

$$\text{Let } -x = t$$

$$-dx = dt$$

$$y \left(-\frac{1}{x} \right) = \int \left(-\frac{1}{t} + \frac{1}{t^2} \right) e^t dt + c$$

$$y \left(-\frac{1}{x} \right) = -\frac{1}{t} e^t + c$$

$$[\text{Since } \int \{f(x) + f'(x)\} e^x dx = f(x) e^x + c]$$

$$-\frac{y}{x} = \frac{1}{x} e^{-x} + c$$

$$y = -(e^{-x} + cx)$$

$$y = -e^x + c_1x, \text{ where } c_1 = -c$$

9. We have,

$$\vec{a} = \hat{i} + \hat{j} + 2\hat{k}$$

$$\vec{b} = 2\hat{i} + \hat{j} - 2\hat{k}$$

We need to find the unit vector in the direction of $6\vec{b}$.

First, let us calculate $6\vec{b}$.

As we have,

$$\vec{b} = 2\hat{i} + \hat{j} - 2\hat{k}$$

Multiply it by 6 on both sides.

$$\Rightarrow 6\vec{b} = 6(2\hat{i} + \hat{j} - 2\hat{k})$$

For finding unit vector, we have the formula:

$$\hat{6b} = \frac{6\vec{b}}{|6\vec{b}|}$$

Now we know the value of $6\vec{b}$, so just substitute the value in the above equation.

$$\Rightarrow \hat{6b} = \frac{12\hat{i} + 6\hat{j} - 12\hat{k}}{|12\hat{i} + 6\hat{j} - 12\hat{k}|}$$

$$\text{Here, } |12\hat{i} + 6\hat{j} - 12\hat{k}| = \sqrt{12^2 + 6^2 + (-12)^2}$$

$$\Rightarrow \hat{6b} = \frac{12\hat{i} + 6\hat{j} - 12\hat{k}}{\sqrt{144 + 36 + 144}}$$

$$\Rightarrow \hat{6b} = \frac{12\hat{i} + 6\hat{j} - 12\hat{k}}{\sqrt{324}}$$

$$\Rightarrow \hat{6b} = \frac{12\hat{i} + 6\hat{j} - 12\hat{k}}{18}$$

Let us simplify.

$$\Rightarrow \hat{6b} = \frac{6(2\hat{i} + \hat{j} - 2\hat{k})}{18}$$

$$\Rightarrow \hat{6b} = \frac{2\hat{i} + \hat{j} - 2\hat{k}}{3}$$

Thus, unit vector in the direction of $6\vec{b}$ is $\frac{2\hat{i} + \hat{j} - 2\hat{k}}{3}$.

10. We know that $(\vec{r} - \vec{a})(\vec{b} \times \vec{c}) = 0$

$$\text{Here } \vec{a} = 3\hat{i} - \hat{j} + 2\hat{k}$$

$$\vec{b} = 2\hat{i} - 5\hat{j} - \hat{k} \text{ and } \vec{c} = -5\hat{i} + 4\hat{j}$$

$$\text{Now, } \vec{b} \times \vec{c} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & -5 & -1 \\ -5 & 4 & 0 \end{vmatrix} = \begin{vmatrix} -5 & -1 \\ 4 & 0 \end{vmatrix} \hat{i} - \begin{vmatrix} 2 & -1 \\ -5 & 0 \end{vmatrix} \hat{j} + \begin{vmatrix} -2 & -5 \\ -5 & 4 \end{vmatrix} \hat{k} = 4\hat{i} + 5\hat{j} - 17\hat{k}$$

Therefore the required equation is $(\vec{r} - \vec{a}) \cdot (\vec{b} \times \vec{c}) = 0$

$$\Rightarrow [(x - 3)\hat{i} + (y + 1)\hat{j} + (z - 2)\hat{k}] \cdot (4\hat{i} + 5\hat{j} - 17\hat{k}) = 0$$

$$\Rightarrow 4(x - 3) + 5(y + 1) - 17(z - 2) = 0$$

$$\Rightarrow 4x - 12 + 5y + 5 - 17z + 34 = 0$$

$$\Rightarrow 4x + 5y - 17z + 27 = 0$$

This is the Cartesian of plane

The required vector equation of the plane is $\vec{r} \cdot (4\hat{i} + 5\hat{j} - 17\hat{k}) + 27 = 0$

OR

The equation of line through $A(0, -1, -1)$ and $B(4, 5, 1)$ is

$$\frac{x-0}{4-0} = \frac{y+1}{5+1} = \frac{z+1}{1+1}$$

$$\text{i.e. } \frac{x}{4} = \frac{y+1}{6} = \frac{z+1}{2} \dots (i)$$

Equation of line through $C(3, 9, 4)$ and $D(-4, 4, 4)$ is

$$\frac{x-3}{-4-3} = \frac{y-9}{4-9} = \frac{z-4}{0}$$

$$\text{i.e., } \frac{x-3}{-7} = \frac{y-9}{-5} = \frac{z-4}{0} \dots \text{(ii)}$$

We know that, the lines

$$\frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1} \text{ and}$$

$$\frac{x-x_2}{a_2} = \frac{y-y_2}{b_2} = \frac{z-z_2}{c_2} \text{ will intersect, if}$$

$$\text{if } \begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

∴ The given lines will intersect, if

$$\begin{vmatrix} 3-0 & 9-(-1) & 4-(-1) \\ 4 & 6 & 2 \\ -7 & -5 & 0 \end{vmatrix} = 0$$

Now,

$$\begin{vmatrix} 3-0 & 9-(-1) & 4-(-1) \\ 4 & 6 & 2 \\ -7 & -5 & 0 \end{vmatrix} = \begin{vmatrix} 3 & 10 & 5 \\ 4 & 6 & 2 \\ -7 & -5 & 0 \end{vmatrix}$$

$$= 3(0 + 10 - 10(0 + 14)) + 5(-20 + 42)$$

$$= 30 - 140 + 110 = 0$$

Hence, the given lines intersect.

Section C

11. To solve this we Use integration by parts that is,

$$\int I \times II dx = I \int II dx - \int \frac{d}{dx} I (\int II dx) dx$$

$$y = x \int \cot x dx - \int \frac{d}{dx} x (\int \cot x dx) dx$$

$$y = (x \log \sin x)_0^{\frac{\pi}{2}} - \int_0^{\frac{\pi}{2}} \log \sin x dx$$

$$\text{Let, } I = \int_0^{\frac{\pi}{2}} \log \sin x dx \dots \text{(i)}$$

Use King theorem of definite integral

$$\int_a^b f(x) dx = \int_a^b f(a+b-x) dx$$

$$I = \int_0^{\frac{\pi}{2}} \log \sin\left(\frac{\pi}{2} - x\right) dx$$

$$I = \int_0^{\frac{\pi}{2}} \log \cos x dx \dots \text{(ii)}$$

Adding eq. (i) and (ii) we get,

$$2I = \int_0^{\frac{\pi}{2}} \log \sin x dx + \int_0^{\frac{\pi}{2}} \log \cos x dx$$

$$2I = \int_0^{\frac{\pi}{2}} \log \frac{2 \sin x \cos x}{2} dx$$

$$2I = \int_0^{\frac{\pi}{2}} \log \sin 2x - \log 2 dx$$

$$\text{Let, } 2x = t$$

$$\Rightarrow 2dx = dt$$

$$\text{At } x = 0, t = 0$$

$$\text{At } x = \frac{\pi}{2}, t = \pi$$

$$2I = \frac{1}{2} \int_0^{\pi} \log \sin t dt - \frac{\pi}{2} \log 2$$

$$2I = \frac{2}{2} \int_0^{\frac{\pi}{2}} \log \sin x dx - \frac{\pi}{2} \log 2$$

$$2I = I - \frac{\pi}{2} \log 2$$


$$I = \int_0^{\frac{\pi}{2}} \log \sin x dx = -\frac{\pi}{2} \log 2$$

$$y = (x \log \sin x)_0^{\frac{\pi}{2}} - \int_0^{\frac{\pi}{2}} \log \sin x dx$$

$$y = \frac{\pi}{2} \log \sin \frac{\pi}{2} - \left(-\frac{\pi}{2} \log 2\right)$$

$$y = \frac{\pi}{2} \log 2$$

Hence proved..



Solving the given equation of circle, $4x^2 + 4y^2 = 9$, and parabola, $x^2 = 4y$, we obtain the point of intersection as $B(\sqrt{2}, \frac{1}{2})$ and $D(-\sqrt{2}, \frac{1}{2})$

It can be observed that the required area is symmetrical about y-axis.

\therefore Area OBCDO = $2 \times$ Area OBCO

We draw BM perpendicular to OA.


Therefore, the coordinates of M are $(\frac{\sqrt{2}}{2}, 0)$

Therefore, Area OBCO = Area OMBCO - Area OMBO

$$\begin{aligned} &= \int_0^{\sqrt{2}} \sqrt{\frac{(9-4x^2)}{4}} dx - \int_0^{\sqrt{2}} \frac{x^2}{4} dx \\ &= \frac{1}{2} \int_0^{\sqrt{2}} \sqrt{9-4x^2} dx - \frac{1}{4} \int_0^{\sqrt{2}} x^2 dx \\ &= \frac{1}{4} \left[x\sqrt{9-4x^2} + \frac{9}{2} \sin^{-1} \frac{2x}{3} \right]_0^{\sqrt{2}} - \frac{1}{4} \left[\frac{x^3}{3} \right]_0^{\sqrt{2}} \\ &= \frac{1}{4} \left[\sqrt{2}\sqrt{9-8} + \frac{9}{2} \sin^{-1} \frac{2\sqrt{2}}{3} \right] - \frac{1}{12} (\sqrt{2})^3 \\ &= \frac{\sqrt{2}}{4} + \frac{9}{8} \sin^{-1} \frac{2\sqrt{2}}{3} - \frac{\sqrt{2}}{6} \\ &= \frac{\sqrt{2}}{12} + \frac{9}{8} \sin^{-1} \frac{2\sqrt{2}}{3} \\ &= \frac{1}{2} \left(\frac{\sqrt{2}}{6} + \frac{9}{4} \sin^{-1} \frac{2\sqrt{2}}{3} \right) \end{aligned}$$

Therefore, the required area OBCDO = $2 \times \frac{1}{2} \left[\frac{\sqrt{2}}{6} + \frac{9}{4} \sin^{-1} \frac{2\sqrt{2}}{3} \right] = \left[\frac{\sqrt{2}}{6} + \frac{9}{4} \sin^{-1} \frac{2\sqrt{2}}{3} \right]$ sq. units.
OR

Equation of one curve (straight line) is $y = x$ (i)



Equation of second curve (parabola) is $y = x^2$... (ii)

Solving eq. (i) and (ii), we get $x = 0$ or $x = 1$ and $y = 0$ or $y = 1$

\therefore Points of intersection of line (i) and parabola (ii) are O (0, 0) and A (1, 1).

Now Area of triangle OAM

= Area bounded by line (i) and x - axis

$$\begin{aligned} &= \left| \int_0^1 y dx \right| = \left| \int_0^1 x dx \right| = \left(\frac{x^2}{2} \right)_0^1 \\ &= \frac{1}{2} - 0 = \frac{1}{2} \text{ sq units} \end{aligned}$$

Also Area OBAM = Area bounded by parabola (ii) and x - axis

$$= \left| \int_0^1 y dx \right| = \left| \int_0^1 x^2 dx \right| = \left(\frac{x^3}{3} \right)_0^1$$

$$= \frac{1}{3} - 0 = \frac{1}{3} \text{ sq. units}$$

∴ Required area OBA between line (i) and parabola (ii)

= Area of triangle OAM – Area of OBAM

$$= \frac{1}{2} - \frac{1}{3} = \frac{3-2}{6} = \frac{1}{6} \text{ sq. units}$$

13. Suppose the required line is parallel to vector \vec{b}

Which is given by $\vec{b} = b_1 \hat{i} + b_2 \hat{j} + b_3 \hat{k}$

We know that the position vector of the point (1, 2, 3) is given by

$$\vec{a} = \hat{i} + 2\hat{j} + 3\hat{k}$$

The equation of line passing through (1, 2, 3) and parallel to \vec{b} is given by

$$\vec{r} = \vec{a} + \lambda \vec{b}$$

$$\Rightarrow \vec{r} = (\hat{i} + 2\hat{j} + 3\hat{k}) + \lambda (b_1 \hat{i} + b_2 \hat{j} + b_3 \hat{k}) \dots (i)$$

The equation of the given planes are

$$\vec{r} \cdot (\hat{i} - \hat{j} + 2\hat{k}) = 5 \dots (ii)$$

$$\text{and } \vec{r} \cdot (3\hat{i} + \hat{j} + \hat{k}) = 6 \dots (iii)$$

The line in Equation (i) and plane in Eq. (ii) are parallel.

Therefore, the normal to the plane of Eq. (ii) is perpendicular to the given line

$$\therefore (\hat{i} - \hat{j} + 2\hat{k}) \cdot (b_1 \hat{i} + b_2 \hat{j} + b_3 \hat{k}) = 0$$

$$\Rightarrow (b_1 - b_2 + 2b_3) = 0 \dots (iv)$$

Similarly, from Eqs. (i) and (iii), we get

$$(3\hat{i} + \hat{j} + \hat{k}) \cdot (b_1 \hat{i} + b_2 \hat{j} + b_3 \hat{k}) = 0$$

$$\Rightarrow (3b_1 + b_2 + b_3) = 0 \dots (v)$$

On solving Equations (iv) and (v) by cross-multiplication, we get

$$\frac{b_1}{(-1) \times 1 - 1 \times 2} = \frac{b_2}{2 \times 3 - 1 \times 1} = \frac{b_3}{1 \times 1 - 3(-1)}$$

$$\Rightarrow \frac{b_1}{-3} = \frac{b_2}{5} = \frac{b_3}{4}$$

Therefore, the direction ratios of \vec{b} are (-3, 5, 4).

$$\vec{b} = -3\hat{i} + 5\hat{j} + 4\hat{k} \left[\because \vec{b} = b_1 \hat{i} + b_2 \hat{j} + b_3 \hat{k} \right]$$

On substituting the value of b in Equation (i), we get

$$\vec{r} = (\hat{i} + 2\hat{j} + 3\hat{k}) + \lambda (-3\hat{i} + 5\hat{j} + 4\hat{k})$$

which is the equation of the required line.

CASE-BASED/DATA-BASED

14. Let E denote the event that student has failed in Economics and M denote the event that student has failed in Mathematics.

$$\therefore P(E) = \frac{50}{100} = \frac{1}{2}, P(M) = \frac{35}{100} = \frac{7}{20} \text{ and } P(E \cap M) = \frac{25}{100} = \frac{1}{4}$$

i. Required probability = P(E|M)

$$= \frac{P(E \cap M)}{P(M)} = \frac{\frac{1}{4}}{\frac{7}{20}} = \frac{1}{4} \times \frac{20}{7} = \frac{5}{7}$$

ii. Required probability = P(M|E)

$$= \frac{P(M \cap E)}{P(E)} = \frac{\frac{1}{4}}{\frac{1}{2}} = \frac{1}{2}$$