

SAMPLE QUESTION PAPER

BLUE PRINT

Time Allowed : 3 hours

Maximum Marks : 80

S. No.	Chapter	VSA / Case based (1 mark)	SA-I (2 marks)	SA-II (3 marks)	LA (5 marks)	Total
1.	Relations and Functions	2(2)	–	1(3)	–	3(5)
2.	Inverse Trigonometric Functions	1(1)	1(2)	–	–	2(3)
3.	Matrices	2(2)	–	–	1(5)*	3(7)
4.	Determinants	1(1)	1(2)*	–	–	2(3)
5.	Continuity and Differentiability	1(1)*	1(2)	2(6)	–	4(9)
6.	Application of Derivatives	1(4)	1(2)	1(3)*	–	3(9)
7.	Integrals	1(1)*	1(2)*	1(3)	–	3(6)
8.	Application of Integrals	–	1(2)	1(3)	–	2(5)
9.	Differential Equations	1(1)*	1(2)	1(3)*	–	3(6)
10.	Vector Algebra	2(2)#	1(2)*	–	–	3(4)
11.	Three Dimensional Geometry	5(5)#	–	–	1(5)*	6(10)
12.	Linear Programming	–	–	–	1(5)*	1(5)
13.	Probability	1(4)	2(4)	–	–	3(8)
	Total	18(24)	10(20)	7(21)	3(15)	38(80)

*It is a choice based question.

#Out of the two or more questions, one/two question(s) is/are choice based.

PDF Watermark Remover DEMO : Purchase from www.PDFWatermarkRemover.com to remove the watermark

MATHEMATICS

*Time allowed : 3 hours**Maximum marks : 80***General Instructions :**

1. This question paper contains two parts A and B. Each part is compulsory. Part-A carries 24 marks and Part-B carries 56 marks.
2. Part-A has Objective Type Questions and Part-B has Descriptive Type Questions.
3. Both Part-A and Part-B have internal choices.

Part - A :

1. It consists of two Sections-I and II.
2. Section-I comprises of 16 very short answer type questions.
3. Section-II contains 2 case study-based questions.

Part - B :

1. It consists of three Sections-III, IV and V.
2. Section-III comprises of 10 questions of 2 marks each.
3. Section-IV comprises of 7 questions of 3 marks each.
4. Section-V comprises of 3 questions of 5 marks each.
5. Internal choice is provided in 3 questions of Section-III, 2 questions of Section-IV and 3 questions of Section-V. You have to attempt only one of the alternatives in all such questions.

PART - A

Section - I

1. If the function $f(x) = \begin{cases} kx^2, & \text{if } x \leq 2 \\ 3, & \text{if } x > 2 \end{cases}$ is continuous at $x = 2$, then find the value of k .

OR

If $y = \log_7 (\log x)$, then find $\frac{dy}{dx}$.

2. If $\tan^{-1}(\cot \theta) = 2\theta$, then find the value of θ .
3. Find the value of $(\hat{i} + \hat{j}) \times (\hat{j} + \hat{k}) \cdot (\hat{k} + \hat{i})$.

OR

If lines $\frac{x-1}{-3} = \frac{y-2}{2k} = \frac{z-3}{2}$ and $\frac{x-1}{3k} = \frac{y-5}{1} = \frac{z-6}{-5}$ are mutually perpendicular, then find the value of k .

4. If a line makes angles 90° , 135° , 45° with the X, Y, Z axes respectively, then find its direction cosines.

5. Evaluate : $\int \frac{dx}{5-8x-x^2}$

OR

Evaluate : $\int_{-\pi/4}^{\pi/4} |\sin x| dx$

6. For matrix $A = \begin{bmatrix} 3 & 4 & -2 \\ -4 & 5 & -3 \\ 2 & 7 & 9 \end{bmatrix}$, find $\frac{1}{2}(A - A')$. (where A' is the transpose of the matrix A)

7. Find the direction cosines of the side AC of a ΔABC whose vertices are given by $A(3, 5, 4)$, $B(-2, -2, -2)$ and $C(3, -5, 4)$.

OR

Show that three points $A(-2, 3, 5)$, $B(1, 2, 3)$ and $C(7, 0, -1)$ are collinear.

8. If $A = \{1, 5, 6\}$, $B = \{7, 9\}$ and $R = \{(a, b) \in A \times B : |a - b| \text{ is even}\}$. Then write the relation R .

9. Find the degree and order of the differential equation : $5x \left(\frac{dy}{dx} \right)^2 - \frac{d^2y}{dx^2} - 6y = \log x$.

OR

Solve the differential equation $(1 + x^2) \frac{dy}{dx} = e^y$.

10. If A and B are the points $(-3, 4, -8)$ and $(5, -6, 4)$ respectively, then find the ratio in which yz -plane divides the line joining the points A and B .

11. If A is a square matrix such that $A^2 = A$, then find $(I + A)^3 - 7A$.

12. A line makes an angle of $\pi/4$ with each of X -axis and Y -axis. What angle does it make with Z -axis?

13. If $P = \begin{bmatrix} 10 & -2 \\ -5 & 1 \end{bmatrix}$, then check whether P^{-1} exists or not.

14. Write the projection of $\vec{b} + \vec{c}$ on \vec{a} , where $\vec{a} = 2\hat{i} - 2\hat{j} + \hat{k}$, $\vec{b} = \hat{i} + 2\hat{j} - 2\hat{k}$ and $\vec{c} = 2\hat{i} - \hat{j} + 4\hat{k}$.


15. Let $n(A) = 4$ and $n(B) = 6$, then find the number of one-one functions from A to B .

16. A line makes 45° with OX , and equal angles with OY and OZ . Find the sum of these three angles.

Section - II

Case study-based questions are compulsory. Attempt any 4 sub parts from each question. Each sub-part carries 1 mark.

17. A card is lost from a pack of 52 cards. From the remaining cards of pack two cards are drawn and are found to be both spades.




Based on the above information, answer the following questions :

- (i) The probability of drawing two spades, given that a card of spade is missing, is
 (a) $\frac{21}{425}$ (b) $\frac{22}{425}$ (c) $\frac{23}{425}$ (d) $\frac{1}{425}$
- (ii) The probability of drawing two spades, given that a card of club is missing, is
 (a) $\frac{26}{425}$ (b) $\frac{22}{425}$ (c) $\frac{19}{425}$ (d) $\frac{23}{425}$
- (iii) Let A be the event of drawing two spades from remaining 51 cards and E_1, E_2, E_3 and E_4 be the events that lost card is of spade, club, diamond and heart respectively, then the value of $\sum_{i=1}^4 P(A / E_i)$ is
 (a) 0.17 (b) 0.24 (c) 0.25 (d) 0.18
- (iv) All of a sudden, missing card is found and, then two cards are drawn simultaneously without replacement. Probability that both drawn cards are aces is
 (a) $\frac{1}{52}$ (b) $\frac{1}{221}$ (c) $\frac{1}{121}$ (d) $\frac{2}{221}$
- (v) If two card are drawn from a well shuffled pack of 52 cards, with replacement, then probability of getting not a king in 1st and 2nd draw is
 (a) $\frac{144}{169}$ (b) $\frac{12}{169}$ (c) $\frac{64}{169}$ (d) none of these

18. Arun got a rectangular parallelopiped shaped box and spherical ball inside it as his birthday present. Sides of the box are $x, 2x$, and $x/3$, while radius of the ball is r cm.

Based on the above information, answer the following questions :



- (i) If S represents the sum of volume of parallelopiped and sphere, then S can be written as
 (a) $\frac{4x^3}{3} + \frac{2}{2}\pi r^2$ (b) $\frac{2x^2}{3} + \frac{4}{3}\pi r^2$
 (c) $\frac{2x^3}{3} + \frac{4}{3}\pi r^3$ (d) $\frac{2}{3}x + \frac{4}{3}\pi r$
- (ii) If sum of the surface areas of box and ball are given to be constant, then x is equal to
 (a) $\sqrt{\frac{k^2 - 4\pi r^2}{6}}$ (b) $\sqrt{\frac{k^2 - 4\pi r}{6}}$ (c) $\sqrt{\frac{k^2 - 4\pi}{6}}$ (d) none of these
- (iii) The radius of the ball, when S is minimum, is
 (a) $\sqrt{\frac{k^2}{54 + \pi}}$ (b) $\sqrt{\frac{k^2}{54 + 4\pi}}$ (c) $\sqrt{\frac{k^2}{64 + 3\pi}}$ (d) $\sqrt{\frac{k^2}{4\pi + 3}}$
- (iv) Relation between length of the box and radius of the ball can be represented as
 (a) $x = 2r$ (b) $x = \frac{r}{2}$ (c) $x = \frac{r}{2}$ (d) $x = 3r$
- (v) Minimum volume of the ball and box together is
 (a) $\frac{k^2}{2(3\pi + 54)^{2/3}}$ (b) $\frac{k}{(3\pi + 54)^{3/2}}$ (c) $\frac{k^3}{3(4\pi + 54)^{1/2}}$ (d) none of these

PART - B

Section - III

19. Find the intervals on which the function $f(x) = 2x^3 + 9x^2 + 12x + 20$ is increasing.



20. A vector \vec{r} is inclined at equal angles to OX , OY and OZ . If the magnitude of \vec{r} is 6 units, then find \vec{r} .

OR

Find the value of λ such that the vectors $\vec{a} = 2\hat{i} + \lambda\hat{j} + \hat{k}$ and $\vec{b} = \hat{i} - 2\hat{j} + 3\hat{k}$ are perpendicular to each other.

21. If A and B are two independent events, such that $P(A) = \frac{1}{2}$ and $P(B) = \frac{1}{5}$, then find the value of $P(A|A \cup B)$.

22. If $x \in [0, 1]$, then find the value of $\frac{1}{2} \cos^{-1} \left(\frac{1-x}{1+x} \right)$.

23. Evaluate : $\int \frac{\sqrt{16 + (\log x)^2}}{x} dx$

OR

Evaluate : $\int_0^{\pi/2} \frac{\sin x}{1 + \cos^2 x} dx$

24. Solve the differential equation : $\frac{dy}{dx} = \frac{3e^{2x} + 3e^{4x}}{e^x + e^{-x}}$

25. A and B are two events such that $P(A) \neq 0$. Find $P(B/A)$ if

- (i) A is a subset of B (ii) $A \cap B = \phi$

26. Find the derivative of $[\sqrt{1-x^2} \sin^{-1} x - x]$ w.r.t. x .

27. Find the area bounded by the curve $x^2 + y^2 = 1$ in the first quadrant.

28. Compute the adjoint of the matrix $\begin{bmatrix} 2 & 0 & -1 \\ 5 & 1 & 0 \\ 1 & 1 & 3 \end{bmatrix}$.

OR

If the matrix $\begin{bmatrix} 1 & a & 2 \\ 1 & 2 & 5 \\ 2 & 1 & 1 \end{bmatrix}$ is not invertible, then find the value of a .

Section - IV

29. Let $A = R - \{2\}$ and $B = R - \{1\}$. If $f: A \rightarrow B$ is a mapping defined by $f(x) = \frac{x-1}{x-2}$, then show that f is bijective.

30. Consider $f(x) = \begin{cases} \frac{\cos^2 x - \sin^2 x - 1}{\sqrt{x^2 + 1} - 1}, & \text{for } x \neq 0 \\ k, & \text{for } x = 0 \end{cases}$. If $f(x)$ is continuous at $x = 0$, then find the value of k .

31. Find the values of x for which $f(x) = (x(x-2))^2$ is an increasing function. Also, find the points on the curve, where the tangent is parallel to x -axis.

OR

An open box with a square base is to be made out of a given quantity of cardboard of area c^2 square units.

Show that the maximum volume of the box is $\frac{c^3}{6\sqrt{3}}$ cubic units.

32. Evaluate : $\int_0^1 \{\tan^{-1} x + \tan^{-1}(1-x)\} dx$

Mathematics

163

33. If $y = x \log \left(\frac{x}{a+bx} \right)$, then prove that $x^3 \frac{d^2 y}{dx^2} = \left(x \frac{dy}{dx} - y \right)^2$.

34. Solve the differential equation $\frac{dy}{dx} = \frac{e^x (\sin^2 x + \sin 2x)}{y(2 \log y + 1)}$.

OR

Find the solution of the equation $\frac{dy}{dx} = \frac{y^2 - y - 2}{x^2 + 2x - 3}$.

35. Find the area enclosed between the curve $y = \log_e (x + e)$ and the coordinates axes.

Section - V

36. Find the image of the point having position vector $\hat{i} + 3\hat{j} + 4\hat{k}$ in the plane $\vec{r} \cdot (2\hat{i} - \hat{j} + \hat{k}) + 3 = 0$.

OR

Find the points on the line $\frac{x+2}{1} = \frac{y+1}{2} = \frac{z-3}{2}$ at a distance of 2 units from the point $(-2, -1, 3)$.

37. Solve the following linear programming problem (LPP) graphically.

Maximize $Z = 4x + 6y$

Subject to constraints:

$$x + 2y \leq 80, \quad 3x + y \leq 75; \quad x, y \geq 0$$

OR

Solve the following linear programming problem (LPP) graphically.

Minimize $Z = 30x + 20y$

Subject to constraints : $x + y \leq 8, x + 4y \geq 12, 5x + 8y \geq 20; x, y \geq 0$

38. If $A = \begin{bmatrix} 0 & 6 & 7 \\ -6 & 0 & 8 \\ 7 & -8 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 2 \\ 1 & 2 & 0 \end{bmatrix}$ and $C = \begin{bmatrix} 2 \\ -2 \\ 3 \end{bmatrix}$, then calculate AC , BC and $(A + B)C$. Also verify that $(A + B)C = AC + BC$.

OR

Find the matrix A satisfying the matrix equation $\begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix} A \begin{bmatrix} 4 & 7 \\ 3 & 5 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$.

1. Since, $f(x)$ is continuous at $x = 2$.

$$\therefore \lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^+} f(x) \Rightarrow k(2)^2 = 3 \Rightarrow k = \frac{3}{4}$$

OR

$$y = \log_7(\log x) = \frac{\log(\log x)}{\log 7}$$

$$\therefore \frac{dy}{dx} = \frac{1}{\log 7} \cdot \frac{1}{\log x} \cdot \frac{1}{x} \Rightarrow \frac{dy}{dx} = \frac{1}{x \log 7 \log x}$$

2. We have, $\tan^{-1}(\cot \theta) = 2\theta \Rightarrow \cot \theta = \tan 2\theta$

$$\Rightarrow \cot \theta = \cot\left(\frac{\pi}{2} - 2\theta\right)$$

$$\Rightarrow \theta = \frac{\pi}{2} - 2\theta \Rightarrow 3\theta = \frac{\pi}{2} \Rightarrow \theta = \frac{\pi}{6}$$

3. We have,

$$\begin{aligned} (\hat{i} + \hat{j}) \times (\hat{j} + \hat{k}) \cdot (\hat{k} + \hat{i}) &= (\hat{i} \times \hat{j} + \hat{i} \times \hat{k} + \hat{j} \times \hat{k}) \cdot (\hat{k} + \hat{i}) \\ &= (\hat{k} - \hat{j} + \hat{i}) \cdot (\hat{k} + \hat{i}) = \hat{k} \cdot \hat{k} + \hat{i} \cdot \hat{i} \quad (\because \hat{i} \cdot \hat{j} = \hat{j} \cdot \hat{k} = \hat{k} \cdot \hat{i} = 0) \\ &= |\hat{k}|^2 + |\hat{i}|^2 = 1 + 1 = 2 \end{aligned}$$

OR

$$\text{Lines } \frac{x-1}{-3} = \frac{y-2}{2k} = \frac{z-3}{2} \text{ and}$$

$$\frac{x-1}{3k} = \frac{y-5}{1} = \frac{z-6}{-5} \text{ are perpendicular if}$$

$$a_1a_2 + b_1b_2 + c_1c_2 = 0.$$

$$\Rightarrow -3(3k) + 2k + 2(-5) = 0 \Rightarrow k = -\frac{10}{7}$$

4. Here $\alpha = 90^\circ$, $\beta = 135^\circ$, $\gamma = 45^\circ$

Direction cosines are $l = \cos \alpha = \cos 90^\circ = 0$,

$$m = \cos \beta = \cos 135^\circ = \frac{-1}{\sqrt{2}}, n = \cos \gamma = \cos 45^\circ = \frac{1}{\sqrt{2}}$$

$$\begin{aligned} 5. \text{ Let } I &= \int \frac{dx}{5-8x-x^2} = \int \frac{dx}{21-(x+4)^2} \\ &= \int \frac{dx}{(\sqrt{21})^2 - (x+4)^2} = \frac{1}{2\sqrt{21}} \log \left| \frac{\sqrt{21} + x + 4}{\sqrt{21} - x - 4} \right| + C \end{aligned}$$

OR

$$\text{Let } I = \int_{-\pi/4}^{\pi/4} |\sin x| dx = 2 \int_0^{\pi/4} \sin x dx$$

$$= 2[-\cos x]_0^{\pi/4} = -2\left[\frac{1}{\sqrt{2}} - 1\right] = 2 - \sqrt{2}$$

Mathematics

$$6. \text{ We have, } A = \begin{bmatrix} 3 & 4 & -2 \\ -4 & 5 & -3 \\ 2 & 7 & 9 \end{bmatrix} \Rightarrow A' = \begin{bmatrix} 3 & -4 & 2 \\ 4 & 5 & 7 \\ -2 & -3 & 9 \end{bmatrix}$$

$$\therefore \frac{1}{2}(A - A') = \frac{1}{2} \begin{bmatrix} 0 & 8 & -4 \\ -8 & 0 & -10 \\ 4 & 10 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 4 & -2 \\ -4 & 0 & -5 \\ 2 & 5 & 0 \end{bmatrix}$$

7. The direction cosines of the line AC are

$$\begin{aligned} &\frac{3-3}{\sqrt{0^2 + (-10)^2 + (0)^2}}, \frac{-5-5}{\sqrt{0^2 + (-10)^2 + 0^2}}, \\ &\frac{4-4}{\sqrt{0^2 + (-10)^2 - 0^2}} \\ &= 0, -1, 0 \end{aligned}$$

OR

Direction ratios of the line AB = 3, -1, -2,

Direction ratios of the line BC = 6, -2, -4

$$\text{Now, } \frac{3}{6} = \frac{-1}{-2} = \frac{-2}{-4}$$

Since the direction cosines of the line AB and BC are proportional and B is the common point. Hence, the points are collinear.

8. We have, $A \times B = \{(1, 7), (1, 9), (5, 7), (5, 9), (6, 7), (6, 9)\}$

$$\therefore R = \{(1, 7), (1, 9), (5, 7), (5, 9)\}$$

9. Here, highest order derivative is $\frac{d^2y}{dx^2}$, so its order

is 2 and power of $\frac{d^2y}{dx^2}$ is one, so its degree is 1.

OR

$$\text{We have, } (1+x^2) \frac{dy}{dx} = e^y$$

$$\Rightarrow \frac{dy}{e^y} = \frac{dx}{1+x^2} \Rightarrow \int \frac{dy}{e^y} = \int \frac{dx}{1+x^2}$$

$$\Rightarrow -e^{-y} = \tan^{-1}x + C \Rightarrow e^{-y} + \tan^{-1}x + C_1 = 0.$$

10. Let λ be the ratio in which yz-plane divides the line joining the points $(-3, 4, -8)$ and $(5, -6, 4)$. The co-ordinates of any point on the line joining the two

points are $\left(\frac{5\lambda-3}{\lambda+1}, \frac{-6\lambda+4}{\lambda+1}, \frac{4\lambda-8}{\lambda+1}\right)$. If the point is in yz-plane, then its x-coordinate should be zero.

$$\therefore \frac{5\lambda-3}{\lambda+1} = 0 \Rightarrow 5\lambda-3=0 \Rightarrow \lambda = \frac{3}{5}$$

So, the required ratio is 3 : 5.

11. We have, $A^2 = A$... (i)
 Now, $(I + A)^3 - 7A = I^3 + A^3 + 3A^2I + 3AI^2 - 7A$
 $= I + A^2A + 3A^2I + 3AI - 7A$
 $= I + AA + 3A + 3A - 7A$ [Using (i)]
 $= I + A^2 - A = I + A - A$ [Using (i)]
 $= I$

12. Let γ be the required angle. Then
 $\cos^2\alpha + \cos^2\beta + \cos^2\gamma = 1$
 $\Rightarrow \cos^2\gamma = 1 - \frac{1}{2} - \frac{1}{2} = 0 \Rightarrow \cos\gamma = 0$
 $\Rightarrow \gamma = \frac{\pi}{2}$

13. Since $|P| = \begin{vmatrix} 10 & -2 \\ -5 & 1 \end{vmatrix} = 10 - 10 = 0$
 $\therefore P^{-1}$ does not exist.

14. Here, $\vec{a} = 2\hat{i} - 2\hat{j} + \hat{k}$
 $\vec{b} = \hat{i} + 2\hat{j} - 2\hat{k}$ and $\vec{c} = 2\hat{i} - \hat{j} + 4\hat{k}$
 $\Rightarrow \vec{b} + \vec{c} = 3\hat{i} + \hat{j} + 2\hat{k}$
 \therefore Projection of $\vec{b} + \vec{c}$ on \vec{a}
 $= \frac{(\vec{b} + \vec{c}) \cdot \vec{a}}{|\vec{a}|} = \frac{(3\hat{i} + \hat{j} + 2\hat{k}) \cdot (2\hat{i} - 2\hat{j} + \hat{k})}{|2\hat{i} - 2\hat{j} + \hat{k}|}$
 $= \frac{3 \times 2 + 1 \times (-2) + 2 \times 1}{\sqrt{2^2 + (-2)^2 + 1^2}} = \frac{6}{3} = 2$

15. Number of one-one functions from A to B
 $= {}^6P_4 = 6 \cdot 5 \cdot 4 \cdot 3 = 360$

16. Here $\alpha = 45^\circ$ and $\beta = \gamma$
 $\therefore \cos\alpha = \frac{1}{\sqrt{2}}$ and $\cos\beta = \cos\gamma$
 Since, $\cos^2\alpha + \cos^2\beta + \cos^2\gamma = 1$
 $\Rightarrow 1/2 + \cos^2\beta + \cos^2\beta = 1$
 $\Rightarrow 2\cos^2\beta = 1/2 \Rightarrow \cos\beta = \frac{1}{2} \Rightarrow \beta = \gamma = 60^\circ$
 $\therefore \alpha + \beta + \gamma = 45^\circ + 60^\circ + 60^\circ = 165^\circ$

17. (i) (b) : Required probability = $\frac{{}^{12}C_2}{{}^{51}C_2}$
 $= \frac{12 \times 11}{51 \times 50} = \frac{22}{425}$

(ii) (a) : Required probability = $\frac{{}^{13}C_2}{{}^{51}C_2} = \frac{13 \times 12}{51 \times 50} = \frac{26}{425}$

(iii) (b) : We have, $P(E_1) = P(E_2) = P(E_3) = P(E_4)$
 $= \frac{13}{52} = \frac{1}{4}$

$$P(A/E_1) = \frac{{}^{12}C_2}{{}^{51}C_2} = \frac{22}{425}$$

$$P(A/E_2) = \frac{{}^{13}C_2}{{}^{51}C_2} = \frac{26}{425}$$

$$P(A/E_3) = P(A/E_4) = \frac{26}{425}$$

$$\therefore \sum_{i=1}^4 P(A/E_i) = \frac{22}{425} + \frac{26}{425} + \frac{26}{425} + \frac{26}{425} = \frac{100}{425} = 0.24$$

(iv) (b) : $P(\text{getting both aces}) = \frac{{}^4C_2}{{}^{52}C_2} = \frac{4 \times 3}{52 \times 51} = \frac{1}{221}$

(v) (a) : $P(\text{drawing a king}) = \frac{4}{52} = \frac{1}{13}$

$$\therefore P(\text{not drawing a king}) = 1 - \frac{1}{13} = \frac{12}{13}$$

$$\therefore \text{Required probability} = \frac{12}{13} \times \frac{12}{13} = \frac{144}{169}$$

18. (i) (c) : Let S be the sum of volume of parallelepiped and sphere, then

$$S = x(2x) \left(\frac{x}{3} \right) + \frac{4}{3}\pi r^3 = \frac{2x^3}{3} + \frac{4}{3}\pi r^3 \quad \dots (1)$$

(ii) (a) : Since, sum of surface area of box and sphere is given to be constant.

$$\therefore 2 \left(x \times 2x + 2x \times \frac{x}{3} + \frac{x}{3} \times x \right) + 4\pi r^2 = k^2 \text{ (say)}$$

$$\Rightarrow 6x^2 + 4\pi r^2 = k^2$$

$$\Rightarrow x^2 = \frac{k^2 - 4\pi r^2}{6} \Rightarrow x = \sqrt{\frac{k^2 - 4\pi r^2}{6}} \quad \dots (2)$$

(iii) (b) : From (1) and (2), we get

$$\begin{aligned} S &= \frac{2}{3} \left(\frac{k^2 - 4\pi r^2}{6} \right)^{3/2} + \frac{4}{3}\pi r^3 \\ &= \frac{2}{3 \times 6\sqrt{6}} (k^2 - 4\pi r^2)^{3/2} + \frac{4}{3}\pi r^3 \\ \Rightarrow \frac{dS}{dr} &= \frac{1}{9\sqrt{6}} \cdot \frac{3}{2} (k^2 - 4\pi r^2)^{1/2} (-8\pi r) + 4\pi r^2 \\ &= 4\pi r \left[r - \frac{1}{3\sqrt{6}} \sqrt{k^2 - 4\pi r^2} \right] \end{aligned}$$

For maximum/minimum, $\frac{dS}{dr} = 0$

$$\Rightarrow \frac{-4\pi r}{3\sqrt{6}} \sqrt{k^2 - 4\pi r^2} = -4\pi r^2$$

$$\Rightarrow k^2 - 4\pi r^2 = 54r^2$$

$$\Rightarrow r^2 = \frac{k^2}{54+4\pi} \Rightarrow r = \sqrt{\frac{k^2}{54+4\pi}} \quad \dots (3)$$

$$(iv) (d): \text{Since, } x^2 = \frac{k^2 - 4\pi r^2}{6} = \frac{1}{6} \left[k^2 - 4\pi \left(\frac{k^2}{54+4\pi} \right) \right]$$

[From (2) and (3)]

$$= \frac{9k^2}{54+4\pi} = 9 \left(\frac{k^2}{54+4\pi} \right) = 9r^2 = (3r)^2$$

$$\Rightarrow x = 3r$$

(v) (c): Minimum volume is given by

$$V = \frac{2}{3}x^3 + \frac{4}{3}\pi r^3 = \frac{2}{3}(3r)^3 + \frac{4}{3}\pi r^3$$

$$= 18r^3 + \frac{4}{3}\pi r^3 = \left(18 + \frac{4}{3}\pi \right) r^3$$

$$= \left(18 + \frac{4}{3}\pi \right) \left(\frac{k^2}{54+4\pi} \right)^{3/2} \quad [\text{Using (3)}]$$

$$= \frac{1}{3} \frac{k^3}{(54+4\pi)^{1/2}}$$

19. Given, $f(x) = 2x^3 + 9x^2 + 12x + 20$

$$\Rightarrow f'(x) = 6x^2 + 18x + 12$$

$$= 6(x^2 + 3x + 2) = 6(x+1)(x+2)$$

For $f(x)$ to be increasing, $f'(x) > 0$

$$\Rightarrow 6(x+1)(x+2) > 0$$

$$\Rightarrow (x+1)(x+2) > 0$$

$$\Rightarrow x+1 > 0, x+2 > 0 \text{ or } x+1 < 0, x+2 < 0$$

$$\Rightarrow x > -1 \text{ or } x < -2$$

$$\Rightarrow x \in (-1, \infty) \text{ or } x \in (-\infty, -2)$$

$\therefore f$ is increasing in $(-\infty, -2) \cup (-1, \infty)$.

20. Suppose \vec{r} makes an angle α with each of the axes OX, OY and OZ. Then, its direction cosines are

$$l = \cos \alpha, m = \cos \alpha, n = \cos \alpha \Rightarrow l = m = n$$

$$\text{Now, } l^2 + m^2 + n^2 = 1 \Rightarrow 3l^2 = 1 \Rightarrow l = \pm \frac{1}{\sqrt{3}}$$

$$\therefore \vec{r} = |\vec{r}| (l\hat{i} + m\hat{j} + n\hat{k})$$

$$\Rightarrow \vec{r} = 6 \left(\pm \frac{1}{\sqrt{3}}\hat{i} \pm \frac{1}{\sqrt{3}}\hat{j} \pm \frac{1}{\sqrt{3}}\hat{k} \right) = 2\sqrt{3}(\pm\hat{i} \pm \hat{j} \pm \hat{k}).$$

OR

If the vectors \vec{a} and \vec{b} are perpendicular to each other, then $\vec{a} \cdot \vec{b} = 0$.

$$\Rightarrow (2\hat{i} + \lambda\hat{j} + \hat{k}) \cdot (\hat{i} - 2\hat{j} + 3\hat{k}) = 0$$

$$\Rightarrow (2)(1) + \lambda(-2) + (1)(3) = 0$$

$$\Rightarrow -2\lambda + 5 = 0 \Rightarrow \lambda = \frac{5}{2}$$

Mathematics

$$21. \text{ We have, } P(A) = \frac{1}{2}, P(B) = \frac{1}{5}$$

$$\text{Now, } P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$= \frac{1}{2} + \frac{1}{5} - \left(\frac{1}{2} \right) \cdot \left(\frac{1}{5} \right) \quad (A \text{ and } B \text{ are independent events})$$

$$= \frac{3}{5}$$

$$\therefore P(A/A \cup B) = \frac{P(A \cap (A \cup B))}{P(A \cup B)}$$

$$= \frac{P(A)}{P(A \cup B)} = \frac{1/2}{3/5} = \frac{5}{6}$$

$$22. \text{ Let } x = \tan^2 \theta \Rightarrow \sqrt{x} = \tan \theta \Rightarrow \theta = \tan^{-1} \sqrt{x}$$

$$\text{Now, } \frac{1}{2} \cos^{-1} \left(\frac{1-x}{1+x} \right) = \frac{1}{2} \cos^{-1} \left(\frac{1-\tan^2 \theta}{1+\tan^2 \theta} \right)$$

$$= \frac{1}{2} \cos^{-1}(\cos 2\theta) = \frac{1}{2}(2\theta) = \theta = \tan^{-1} \sqrt{x}$$

$$23. \text{ Let } I = \int \frac{\sqrt{16+(\log x)^2}}{x} dx$$

$$\text{Put } \log x = t \Rightarrow \frac{1}{x} dx = dt$$

$$\therefore I = \int \sqrt{16+t^2} dt$$

$$= \frac{t}{2} \sqrt{16+t^2} + \frac{16}{2} \log |t + \sqrt{16+t^2}| + c$$

$$\therefore I = \frac{1}{2} \log x \sqrt{16+(\log x)^2}$$

$$+ 8 \log |\log x + \sqrt{16+(\log x)^2}| + c$$

OR

$$\text{Let } I = \int_0^{\pi/2} \frac{\sin x}{1+\cos^2 x} dx$$

$$\text{Put } \cos x = t \Rightarrow -\sin x dx = dt$$

$$\text{When } x = 0, t = 1 \text{ and when } x = \frac{\pi}{2}, t = 0$$

$$\therefore I = - \int_1^0 \frac{dt}{1+t^2} = - [\tan^{-1} t]_1^0$$

$$= -[\tan^{-1} 0 - \tan^{-1} 1] = \frac{\pi}{4}$$

$$24. \text{ We have, } \frac{dy}{dx} = \frac{3e^{2x} + 3e^{4x}}{e^x + e^{-x}}$$

$$\Rightarrow \int dy = \int \frac{3e^{2x}(1+e^{2x})}{e^{-x}(e^{2x}+1)} dx \quad [\text{Integrating both sides}]$$

$$\Rightarrow y = \int 3e^{3x} dx = \frac{3e^{3x}}{3} + c \Rightarrow y = e^{3x} + c$$



25. (i) Since, A is a subset of B . $\therefore A \subset B$

$$\Rightarrow A \cap B = A$$

$$\therefore P(A \cap B) = P(A) \quad \dots (i)$$

$$\text{Now, } P(B/A) = \frac{P(B \cap A)}{P(A)} = \frac{P(A)}{P(A)} \quad [\text{Using (i)}]$$

$$= 1$$

$$(ii) \text{ If } A \cap B = \phi \Rightarrow P(A \cap B) = 0$$

$$\therefore P(B/A) = \frac{P(B \cap A)}{P(A)} = \frac{0}{P(A)} = 0$$

26. We have, $\frac{d}{dx}[(\sqrt{1-x^2})\sin^{-1}x - x]$

$$= (\sqrt{1-x^2}) \cdot \frac{d}{dx}(\sin^{-1}x) + (\sin^{-1}x) \cdot \frac{d}{dx}(\sqrt{1-x^2}) - 1$$

$$= (\sqrt{1-x^2}) \cdot \frac{1}{(\sqrt{1-x^2})} + (\sin^{-1}x) \cdot \frac{1}{2}(1-x^2)^{-1/2} \cdot (-2x) - 1$$

$$= 1 - \frac{x \sin^{-1}x}{\sqrt{1-x^2}} - 1 = \frac{-x \sin^{-1}x}{\sqrt{1-x^2}}$$

27. We have, $x^2 + y^2 = 1$, a circle with centre $(0, 0)$ and radius = 1.


Required area

= area of shaded region

$$A = \int_0^1 \sqrt{1-x^2} dx$$

$$= \left[\frac{x}{2} \sqrt{1-x^2} + \frac{1}{2} \sin^{-1} \frac{x}{1} \right]_0^1$$

$$= \left[\frac{1}{2} \sin^{-1} 1 \right] = \left(\frac{1}{2} \times \frac{\pi}{2} \right) = \frac{\pi}{4} \text{ sq. unit}$$



28. Let $A = \begin{bmatrix} 2 & 0 & -1 \\ 5 & 1 & 0 \\ 1 & 1 & 3 \end{bmatrix}$

$$A_{11} = (-1)^{1+1} \begin{vmatrix} 1 & 0 \\ 1 & 3 \end{vmatrix} = 3; \quad A_{12} = (-1)^{1+2} \begin{vmatrix} 5 & 0 \\ 1 & 3 \end{vmatrix} = -15$$

$$A_{13} = (-1)^{1+3} \begin{vmatrix} 5 & 1 \\ 1 & 1 \end{vmatrix} = 4; \quad A_{21} = (-1)^{2+1} \begin{vmatrix} 0 & -1 \\ 1 & 3 \end{vmatrix} = -1$$

$$A_{22} = (-1)^{2+2} \begin{vmatrix} 2 & -1 \\ 1 & 3 \end{vmatrix} = 7; \quad A_{23} = (-1)^{2+3} \begin{vmatrix} 2 & 0 \\ 1 & 1 \end{vmatrix} = -2;$$

$$A_{31} = (-1)^{3+1} \begin{vmatrix} 0 & -1 \\ 1 & 0 \end{vmatrix} = 1; \quad A_{32} = (-1)^{3+2} \begin{vmatrix} 2 & -1 \\ 5 & 0 \end{vmatrix} = -5;$$

$$A_{33} = (-1)^{3+3} \begin{vmatrix} 2 & 0 \\ 5 & 1 \end{vmatrix} = 2$$

$$\therefore \text{adj } A = \begin{bmatrix} 3 & -15 & 4 \\ -1 & 7 & -2 \\ 1 & -5 & 2 \end{bmatrix} = \begin{bmatrix} 3 & -1 & 1 \\ -15 & 7 & -5 \\ 4 & -2 & 2 \end{bmatrix}$$

OR

$$\text{The matrix is not invertible if } \begin{vmatrix} 1 & a & 2 \\ 1 & 2 & 5 \\ 2 & 1 & 1 \end{vmatrix} = 0$$

$$\Rightarrow 1(2-5) - a(1-10) + 2(1-4) = 0$$

$$\Rightarrow -3 + 9a - 6 = 0 \Rightarrow a = 1$$

29. We have, $f(x) = \frac{x-1}{x-2}$

For one-one : Let $x, y \in A$ and consider $f(x) = f(y)$

$$\Rightarrow \frac{x-1}{x-2} = \frac{y-1}{y-2}$$

$$\Rightarrow (x-1)(y-2) = (x-2)(y-1)$$

$$\Rightarrow xy - y - 2x + 2 = xy - x - 2y + 2 \Rightarrow x = y$$

Thus, $f(x) = f(y) \Rightarrow x = y$ for all $x, y \in A$

So, f is one-one.

For onto : Let y be an arbitrary element of B . Then,

$$f(x) = y \Rightarrow \frac{x-1}{x-2} = y \Rightarrow (x-1) = y(x-2) \Rightarrow x = \frac{1-2y}{1-y}$$

Clearly, $x = \frac{1-2y}{1-y}$ is a real number for all $y \neq 1$.

Also, $\frac{1-2y}{1-y} \neq 2$ for any y , for, if we take $\frac{1-2y}{1-y} = 2$,

then we get $1 = 2$, which is wrong.

So, f is onto. Hence, f is a bijective.

30. $f(0) = k$ (Given) ...(i)

Since, $f(x)$ is continuous at $x = 0$.

$$\therefore f(0) = \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x)$$

$$\text{Now, } \lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \frac{\cos^2 x - \sin^2 x - 1}{\sqrt{x^2 + 1} - 1}$$

$$= \lim_{x \rightarrow 0} \frac{1 - \sin^2 x - \sin^2 x - 1}{\sqrt{x^2 + 1} - 1} \times \frac{\sqrt{x^2 + 1} + 1}{\sqrt{x^2 + 1} + 1}$$

$$= \lim_{x \rightarrow 0} \frac{-2 \sin^2 x (\sqrt{x^2 + 1} + 1)}{x^2 + 1 - 1}$$

$$= \lim_{x \rightarrow 0} -2 \frac{\sin^2 x}{x^2} \cdot (\sqrt{x^2 + 1} + 1)$$

$$= -2 \left(\lim_{x \rightarrow 0} \frac{\sin^2 x}{x^2} \right) \times \lim_{x \rightarrow 0} (\sqrt{x^2 + 1} + 1)$$

$$= -2(1)^2 (1 + 1) = -4$$

...(ii)

From (i) and (ii), we get $k = -4$.

31. Given, $f(x) = (x(x-2))^2 = x^2(x-2)^2$, $D_f = R$.


Differentiating w.r.t. x , we get

$$f'(x) = x^2 \cdot 2(x-2) + (x-2)^2 \cdot 2x$$

$$= 2x(x-2)(x+x-2) = 2x(x-2)(2x-2) = 4x(x-1)(x-2)$$

Now, the given function f is (strictly) increasing iff $f'(x) > 0$

$$\Rightarrow x \in (0, 1) \cup (2, \infty)$$



Further, the tangents will be parallel to x -axis iff $f'(x) = 0$

$$\Rightarrow x = 0, 1, 2$$

The given curve is $y = x^2(x - 2)^2$

When $x = 0$, $y = 0$;

When $x = 1$, $y = 1^2(1 - 2)^2 = 1 \times (-1)^2 = 1 \times 1 = 1$;

When $x = 2$, $y = 2^2(2 - 2)^2 = 4 \times 0 = 0$.

\therefore The points on the given curve, where the tangents are parallel to x -axis are $(0, 0)$, $(1, 1)$ and $(2, 0)$.

OR

Let h be height and x be the side of the square base of the open box.

Then its area $= x \times x + 4h \times x = c^2$

$$\Rightarrow h = \frac{c^2 - x^2}{4x}$$

Now $V =$ volume of the box

$$= x^2 h = x^2 \cdot \frac{c^2 - x^2}{4x} = \frac{1}{4}(c^2 x - x^3)$$

$$\Rightarrow \frac{dV}{dx} = \frac{1}{4}(c^2 - 3x^2) \text{ and } \frac{d^2V}{dx^2} = \frac{1}{4}(-6x) = -\frac{3}{2}x$$

For maxima or minima $\frac{dV}{dx} = 0 \Rightarrow x^2 = \frac{c^2}{3}$

$$\Rightarrow x = \frac{c}{\sqrt{3}} \quad (\because x \neq 0)$$

For this value of x , $\frac{d^2V}{dx^2} < 0$

$\Rightarrow V$ is maximum at $x = \frac{c}{\sqrt{3}}$ and its maximum volume is,

$$V = \frac{1}{4}x(c^2 - x^2) = \frac{1}{4} \cdot \frac{c}{\sqrt{3}} \left(c^2 - \frac{c^2}{3} \right) = \frac{c^3}{6\sqrt{3}} \text{ cubic units.}$$


32. Consider, $\int_0^1 \{\tan^{-1}x + \tan^{-1}(1-x)\} dx$

$$= \int_0^1 \tan^{-1}x dx + \int_0^1 \tan^{-1}(1-x) dx$$

$$= \int_0^1 \tan^{-1}x dx + \int_0^1 \tan^{-1}\{1-(1-x)\} dx$$

$$\left[\because \int_0^a f(x) dx = \int_0^a f(a-x) dx \right]$$

$$= \int_0^1 \tan^{-1}x dx + \int_0^1 \tan^{-1}x dx = 2 \int_0^1 \tan^{-1}x dx$$



$$\begin{aligned} &= 2[(\tan^{-1}x) \cdot x]_0^1 - 2 \int_0^1 \frac{x}{(1+x^2)} dx \\ &= 2[(\tan^{-1}1) \cdot 1 - 0] - [\log(1+x^2)]_0^1 \\ &= \left(2 \times \frac{\pi}{4}\right) - (\log 2 - \log 1) = \left(\frac{\pi}{2} - \log 2\right) \end{aligned}$$

33. Here, $y = x \log \left(\frac{x}{a+bx} \right)$... (i)

$$\Rightarrow y = x[\log x - \log(a+bx)] = x \log x - x \log(a+bx)$$

$$\Rightarrow \frac{dy}{dx} = x \cdot \frac{1}{x} + 1 \cdot \log x - \left[1 \cdot \log(a+bx) + x \cdot \frac{1}{a+bx} \cdot b \right]$$

$$= 1 - \frac{bx}{a+bx} + \log x - \log(a+bx)$$

$$\Rightarrow \frac{dy}{dx} = \frac{a}{a+bx} + \log \left(\frac{x}{a+bx} \right)$$

$$\Rightarrow \frac{dy}{dx} = \frac{a}{a+bx} + \frac{y}{x} \quad [\text{Using (i)}] \quad \dots (ii)$$

Again differentiating (ii) w.r.t. x , we get

$$\begin{aligned} \frac{d^2y}{dx^2} &= a \cdot (-1)(a+bx)^{-2} \cdot b + \frac{x \frac{dy}{dx} - y}{x^2} \\ &= \frac{-ab}{(a+bx)^2} + \frac{a}{x(a+bx)} = \frac{-abx + a(a+bx)}{x(a+bx)^2} = \frac{a^2}{x(a+bx)^2} \end{aligned}$$

$$\text{Now, R.H.S.} = \left(x \frac{dy}{dx} - y \right)^2$$

$$= \left\{ x \cdot \left[\frac{a}{a+bx} + \frac{y}{x} \right] - y \right\}^2 = \left(\frac{ax}{a+bx} \right)^2$$

$$\text{and L.H.S.} = x^3 \frac{d^2y}{dx^2} = \frac{a^2 x^2}{(a+bx)^2} = \left[\frac{ax}{a+bx} \right]^2 = \text{R.H.S.}$$

34. We have, $\frac{dy}{dx} = \frac{e^x (\sin^2 x + \sin 2x)}{y(2 \log y + 1)}$

$$\Rightarrow \int y(2 \log y + 1) dy = \int e^x (\sin^2 x + \sin 2x) dx$$

$$\Rightarrow 2 \int y \log y dy + \int y dy = \int e^x (\sin^2 x + \sin 2x) dx$$

$$\Rightarrow 2 \left[\log |y| \cdot \frac{y^2}{2} - \int \frac{1}{y} \times \frac{y^2}{2} dy \right] + \frac{y^2}{2} = e^x \sin^2 x + C$$

$$[\because \int e^x (f(x) + f'(x)) dx = e^x f(x) + C]$$

$$\Rightarrow y^2 \log |y| - \frac{y^2}{2} + \frac{y^2}{2} = e^x \sin^2 x + C$$

$$\Rightarrow y^2 \log |y| = e^x \sin^2 x + C, \text{ which is required solution.}$$

OR

$$\text{We have } \frac{dy}{y^2 - y - 2} = \frac{dx}{x^2 + 2x - 3}$$

Integrating both sides, we get

$$\begin{aligned} \int \frac{dy}{y^2 - y - 2} &= \int \frac{dx}{x^2 + 2x - 3} \\ \Rightarrow \int \frac{dy}{\left(y - \frac{1}{2}\right)^2 - \left(\frac{3}{2}\right)^2} &= \int \frac{dx}{(x+1)^2 - 2^2} + c \\ \Rightarrow \frac{1}{2 \cdot \frac{3}{2}} \log \left| \frac{y - \frac{1}{2} - \frac{3}{2}}{y - \frac{1}{2} + \frac{3}{2}} \right| &= \frac{1}{2 \cdot 2} \log \left| \frac{x+1-2}{x+1+2} \right| + c \\ \Rightarrow \frac{1}{3} \log \left| \frac{y-2}{y+1} \right| &= \frac{1}{4} \log \left| \frac{x-1}{x+3} \right| + c \end{aligned}$$

35. The bounded area is as shown in figure.

Curve is $y = \log_e(x+e)$

If $y = 0$, then $x = 1 - e$

$A \equiv (1 - e, 0)$

Required area is

$$A = \int_{1-e}^0 \log_e(x+e) dx$$

Put $x + e = t \Rightarrow dx = dt$ and $x = 1 - e \Rightarrow t = 1$ and $x = 0 \Rightarrow t = e$

$$\begin{aligned} A &= \int_1^e \log t dt = [t \log t - t]_1^e = e \log e - e - 0 + 1 \\ &= 1 \text{ sq. unit} \end{aligned}$$

36. Let Q be the image of the point $P(\hat{i} + 3\hat{j} + 4\hat{k})$ in the plane $\vec{r} \cdot (2\hat{i} - \hat{j} + \hat{k}) + 3 = 0$

Then, PQ is normal to the plane. Since PQ passes through P and is normal to the given plane, therefore equation of line PQ is

$$\vec{r} = (\hat{i} + 3\hat{j} + 4\hat{k}) + \lambda(2\hat{i} - \hat{j} + \hat{k})$$

Since Q lies on line PQ , so let the position vector of Q be $(\hat{i} + 3\hat{j} + 4\hat{k}) + \lambda(2\hat{i} - \hat{j} + \hat{k})$

$$= (1 + 2\lambda)\hat{i} + (3 - \lambda)\hat{j} + (4 + \lambda)\hat{k}$$

Since, R is the mid-point of PQ . Therefore, position vector of R is

$$\begin{aligned} &\frac{[(1 + 2\lambda)\hat{i} + (3 - \lambda)\hat{j} + (4 + \lambda)\hat{k}] + [\hat{i} + 3\hat{j} + 4\hat{k}]}{2} \\ &= (\lambda + 1)\hat{i} + \left(3 - \frac{\lambda}{2}\right)\hat{j} + \left(4 + \frac{\lambda}{2}\right)\hat{k} \end{aligned}$$

Since R lies on the plane $\vec{r} \cdot (2\hat{i} - \hat{j} + \hat{k}) + 3 = 0$

$$\Rightarrow \left\{ (\lambda + 1)\hat{i} + \left(3 - \frac{\lambda}{2}\right)\hat{j} + \left(4 + \frac{\lambda}{2}\right)\hat{k} \right\} \cdot (2\hat{i} - \hat{j} + \hat{k}) + 3 = 0$$

$$\Rightarrow 2\lambda + 2 - 3 + \frac{\lambda}{2} + 4 + \frac{\lambda}{2} + 3 = 0 \Rightarrow \lambda = -2$$

Thus, the position vector of Q is

$$(\hat{i} + 3\hat{j} + 4\hat{k}) - 2(2\hat{i} - \hat{j} + \hat{k}) = -3\hat{i} + 5\hat{j} + 2\hat{k}$$

OR

$$\text{The given line is } \frac{x+2}{1} = \frac{y+1}{2} = \frac{z-3}{2} \quad \dots(i)$$

Let $P(-2, -1, 3)$ lies on the line.

The direction ratios of line (i) are 1, 2, 2

\therefore The direction cosines of line are $\frac{1}{3}, \frac{2}{3}, \frac{2}{3}$

Equation (i) may be written as

$$\frac{x+2}{\frac{1}{3}} = \frac{y+1}{\frac{2}{3}} = \frac{z-3}{\frac{2}{3}} \quad \dots(ii)$$

Coordinates of any point on the line (ii) may be taken as

$$\left(\frac{1}{3}r - 2, \frac{2}{3}r - 1, \frac{2}{3}r + 3 \right)$$

$$\text{Let } Q \equiv \left(\frac{1}{3}r - 2, \frac{2}{3}r - 1, \frac{2}{3}r + 3 \right)$$

Given $|r| = 2$, $\therefore r = \pm 2$

Putting the values of r , we have

$$Q \equiv \left(-\frac{4}{3}, \frac{1}{3}, \frac{13}{3} \right) \text{ or } Q \equiv \left(\frac{-8}{3}, \frac{-7}{3}, \frac{5}{3} \right)$$


37. We have maximize $Z = 4x + 6y$.

Subject to constraints :

$$x + 2y \leq 80, 3x + y \leq 75 \text{ and } x \geq 0, y \geq 0$$

Now we draw the graphs of the lines

$$l_1 : x + 2y = 80, l_2 : 3x + y = 75 \text{ and } x = 0, y = 0.$$



We obtain shaded region as the feasible region.

The lines l_1 and l_2 intersect at $Q(14, 33)$.

Thus, the vertices of the feasible region are $P(0, 40)$, $Q(14, 33)$, $R(25, 0)$ and $O(0, 0)$.

Corner Points	Value of $Z = 4x + 6y$
$P(0, 40)$	240
$Q(14, 33)$	254 (Maximum)
$R(25, 0)$	100
$O(0, 0)$	0

Thus, Z has maximum value 254 at $Q(14, 33)$.

OR


We have minimize $Z = 30x + 20y$.

Subject to constraints :

$$x + y \leq 8, x + 4y \geq 12, 5x + 8y \geq 20, x, y \geq 0$$

Now, we draw the graphs of

$$l_1 : x + y = 8, l_2 : x + 4y = 12, l_3 : 5x + 8y = 20 \text{ and } x = 0, y = 0$$



Shaded region ABC is the required feasible region.

$B\left(\frac{20}{3}, \frac{4}{3}\right)$ is the point of intersection of the lines l_1 and l_2 .

Thus, the vertices of the feasible region are

$$A(0, 3), B\left(\frac{20}{3}, \frac{4}{3}\right) \text{ and } C(0, 8).$$

Corner Points	Value of $Z = 30x + 20y$
$A(0, 3)$	60 (Minimum)
$B(20/3, 4/3)$	226.6
$C(0, 8)$	160

$\therefore Z$ has minimum value 60 at $A(0, 3)$.

$$38. AC = \begin{bmatrix} 0 & 6 & 7 \\ -6 & 0 & 8 \\ 7 & -8 & 0 \end{bmatrix} \begin{bmatrix} 2 \\ -2 \\ 3 \end{bmatrix}$$

$$= \begin{bmatrix} 0 \cdot 2 + 6 \cdot (-2) + 7 \cdot 3 \\ (-6) \cdot 2 + 0 \cdot (-2) + 8 \cdot 3 \\ 7 \cdot 2 + (-8) \cdot (-2) + 0 \cdot 3 \end{bmatrix} = \begin{bmatrix} 9 \\ 12 \\ 30 \end{bmatrix}$$

$$BC = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 2 \\ 1 & 2 & 0 \end{bmatrix} \begin{bmatrix} 2 \\ -2 \\ 3 \end{bmatrix} = \begin{bmatrix} 0 \cdot 2 + 1 \cdot (-2) + 1 \cdot 3 \\ 1 \cdot 2 + 0 \cdot (-2) + 2 \cdot 3 \\ 1 \cdot 2 + 2 \cdot (-2) + 0 \cdot 3 \end{bmatrix} = \begin{bmatrix} 1 \\ 8 \\ -2 \end{bmatrix}$$



Mathematics

$$A + B = \begin{bmatrix} 0 & 6 & 7 \\ -6 & 0 & 8 \\ 7 & -8 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 2 \\ 1 & 2 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} 0+0 & 6+1 & 7+1 \\ -6+1 & 0+0 & 8+2 \\ 7+1 & -8+2 & 0+0 \end{bmatrix} = \begin{bmatrix} 0 & 7 & 8 \\ -5 & 0 & 10 \\ 8 & -6 & 0 \end{bmatrix}$$

$$\therefore (A + B)C = \begin{bmatrix} 0 & 7 & 8 \\ -5 & 0 & 10 \\ 8 & -6 & 0 \end{bmatrix} \begin{bmatrix} 2 \\ -2 \\ 3 \end{bmatrix}$$

$$= \begin{bmatrix} 0 \cdot 2 + 7 \cdot (-2) + 8 \cdot 3 \\ (-5) \cdot 2 + 0 \cdot (-2) + 10 \cdot 3 \\ 8 \cdot 2 + (-6) \cdot (-2) + 0 \cdot 3 \end{bmatrix} = \begin{bmatrix} 10 \\ 20 \\ 28 \end{bmatrix} \quad \dots(i)$$

$$\text{Now, } AC + BC = \begin{bmatrix} 9 \\ 12 \\ 30 \end{bmatrix} + \begin{bmatrix} 1 \\ 8 \\ -2 \end{bmatrix} = \begin{bmatrix} 9+1 \\ 12+8 \\ 30-2 \end{bmatrix} = \begin{bmatrix} 10 \\ 20 \\ 28 \end{bmatrix} \quad \dots(ii)$$

From (i) and (ii), we get

$$(A + B)C = AC + BC$$

OR

$$\text{Let } B = \begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix} \text{ and } C = \begin{bmatrix} 4 & 7 \\ 3 & 5 \end{bmatrix}$$

$$\text{Now, } |B| = 3 - 4 = -1 \neq 0$$

$$|C| = 20 - 21 = -1 \neq 0$$

Hence B^{-1} and C^{-1} exist.

\therefore The given matrix equation becomes $BAC = I$

$$\Rightarrow B^{-1}(BAC)C^{-1} = B^{-1}I C^{-1} \Rightarrow IAI = B^{-1}C^{-1}$$

$$\Rightarrow A = B^{-1}C^{-1} \quad \dots(i)$$

$$\text{Now, adj } B = \begin{bmatrix} 3 & -2 \\ -2 & 1 \end{bmatrix}$$

$$\therefore B^{-1} = \frac{1}{|B|} (\text{adj } B) = \frac{1}{-1} \begin{bmatrix} 3 & -2 \\ -2 & 1 \end{bmatrix} = \begin{bmatrix} -3 & 2 \\ 2 & -1 \end{bmatrix}$$

$$\text{Also, adj } C = \begin{bmatrix} 5 & -3 \\ -7 & 4 \end{bmatrix} = \begin{bmatrix} 5 & -7 \\ -3 & 4 \end{bmatrix}$$

$$\therefore C^{-1} = \frac{1}{|C|} (\text{adj } C) = \frac{1}{-1} \begin{bmatrix} 5 & -7 \\ -3 & 4 \end{bmatrix} = \begin{bmatrix} -5 & 7 \\ 3 & -4 \end{bmatrix}$$

$$\text{Now, from (i), } A = B^{-1}C^{-1} = \begin{bmatrix} -3 & 2 \\ 2 & -1 \end{bmatrix} \begin{bmatrix} -5 & 7 \\ 3 & -4 \end{bmatrix}$$

$$= \begin{bmatrix} 15+6 & -21-8 \\ -10-3 & 14+4 \end{bmatrix} = \begin{bmatrix} 21 & -29 \\ -13 & 18 \end{bmatrix}$$