

**Sample Question Paper - 19**  
**Physics (042)**  
**Class- XII, Session: 2021-22**  
**TERM II**

Time : 2 Hours

Max. Marks : 35

**General Instructions :**


- (i) There are 12 questions in all. All questions are compulsory.
- (ii) This question paper has three sections: Section A, Section B and Section C.
- (iii) Section A contains three questions of two marks each, Section B contains eight questions of three marks each, Section C contains one case study-based question of five marks.
- (iv) There is no overall choice. However, an internal choice has been provided in one question of two marks and two questions of three marks. You have to attempt only one of the choices in such questions.
- (v) You may use log tables if necessary but use of calculator is not allowed.

**SECTION - A**

1. How does a light emitting diode (LED) work? Give two advantages of LED's over the conventional incandescent lamps.
2. In an experiment on  $\alpha$ -particle scattering by a thin foil of gold, draw a graph showing, the number of particles scattered versus the scattering angle  $\theta$ . Why is it that a very small fraction of the particles are scattered at  $\theta > 90^\circ$ ? Write two important conclusions that can be drawn regarding the structure of the atom from the study of this experiment.

**OR**

The figure shows energy level diagram of hydrogen atom.



- (a) Find out the transition which results in the emission of a photon of wavelength 496 nm.
  - (b) Which transition corresponds to the emission of radiation of maximum wavelength? Justify your answer.
3. Distinguish between 'intrinsic' and 'extrinsic' semiconductors.


**SECTION - B**

4. Write the two processes that take place in the formation of a  $p$ - $n$  junction. Explain with the help of a diagram, the formation of depletion region and barrier potential in a  $p$ - $n$  junction.



5. The maximum kinetic energy of the photoelectrons emitted is doubled when the wavelength of light incident on the photosensitive surface changes from  $\lambda_1$  to  $\lambda_2$ . Deduce expressions for the threshold wavelength and work function for the metal surface in terms of  $\lambda_1$  and  $\lambda_2$ .

6. Calculate for how many years will the fusion of 2.0 kg deuterium keep 800 W electric lamp glowing. Take the fusion reaction as



7. The electric field of an electromagnetic wave in free space is given by  $\vec{E} = 10 \cos(10^7 t + kx) \hat{j}$  V/m, where  $t$  and  $x$  are in seconds and metres respectively. Find the wavelength  $\lambda$  and wave number  $k$ ?

OR

The magnetic component of a wave of light is  $B_x = (4.0 \times 10^{-6} \text{ T}) \sin [(1.57 \times 10^7 \text{ m}^{-1}) y + \omega t]$   
Find the intensity of light.

8. Dictate the phenomenon of total internal reflection applied in cable fibre optics.


9. Deduce, with the help of Young's arrangement to produce interference pattern and an expression for the fringe width.

OR

Using Huygens principle, obtain the law of refraction at a plane interface when light passes from a rarer to a denser medium.

10. The energy levels of a hypothetical atom are shown in the given figure. Which of the shown transitions will result in the emission of a photon of wavelength 275 nm?

Which of these transitions correspond to emission of radiation of (i) maximum and (ii) minimum wavelength?



11. Two similar thin equi-convex lenses, of focal length  $f$  each, are kept coaxially in contact with each other such that the focal length of the combination is  $F_1$ . When the space between the two lenses is filled with glycerin (which has the same refractive index ( $\mu = 1.5$ ) as that of glass) then the equivalent focal length is  $F_2$ . Find the ratio of  $F_1$  and  $F_2$ .

## SECTION - C

### 12. CASE STUDY : ELECTROMAGNETIC SPECTRUM

All the known radiations from a big family of electromagnetic waves which stretch over a large range of wavelengths. Electromagnetic wave include radio waves, microwaves, visible light waves, infrared rays, UV rays, X-rays and gamma rays. The orderly distribution of the electromagnetic waves in accordance with their wavelength or frequency into distinct groups having widely differing properties is electromagnetic spectrum.

(i) Which wavelength of the Sun is used finally as electric energy?

- |                   |                    |
|-------------------|--------------------|
| (a) Radio waves   | (b) Infrared waves |
| (c) Visible light | (d) Microwaves     |

- (ii) Which of the following electromagnetic radiations have the longest wavelength?
- (a) X-rays (b)  $\gamma$ -rays  
(c) Microwaves (d) Radiowaves
- (iii) Which one of the following is not electromagnetic in nature?
- (a) X-rays (b) Gamma rays  
(c) Cathode rays (d) Infrared rays
- (iv) Which of the following has minimum wavelength ?
- (a) X-rays (b) Ultraviolet rays  
(c)  $\gamma$ -rays (d) Cosmic rays
- (v) The decreasing order of wavelength of infrared, microwave, ultraviolet and gamma rays is
- (a) microwave, infrared, ultraviolet, gamma rays  
(b) gamma rays, ultraviolet, infrared, microwave  
(c) microwave, gamma rays, infrared, ultraviolet  
(d) infrared, microwave, ultraviolet, gamma rays.



## Solution


### PHYSICS - 042

#### Class 12 - Physics

1. A light emitting diode is simply a forward biased  $p$ - $n$  junction which emits spontaneous light radiation. At the junction, energy is released in the form of photons due to the recombination of the excess minority charge carrier with the majority charge carrier.

Advantages :

- (i) Low operational voltage and less power.
- (ii) Fast action and no warm up time required.



A very small fraction of  $\alpha$ -particles are scattered at  $\theta > 90^\circ$  because the size of nucleus is very small nearly  $1/8000$  times the size of atom. So, a few  $\alpha$ -particles experience a strong repulsive force and turn back.

Conclusions :

- (i) Entire positive charge and most of the mass of the atom is concentrated in the nucleus with the electrons some distance away.
- (ii) Size of the nucleus is about  $10^{-15}$  m to  $10^{-14}$  m, while size of the atom is  $10^{-10}$  m, so the electrons are at distance  $10^4$  m to  $10^5$  m from the nucleus, and being large empty space in the atom, most  $\alpha$  particles go through the empty space.

OR

(a)  $\lambda = 496 \text{ nm} = 496 \times 10^{-9} \text{ m}$

$$E = \frac{hc}{\lambda} = \frac{6.62 \times 10^{-34} \times 3 \times 10^8}{496 \times 10^{-9}} \text{ J}$$

$$= \frac{6.6 \times 10^{-34} \times 3 \times 10^8}{496 \times 10^{-9} \times 1.6 \times 10^{-19}} = 2.5 \text{ eV}$$

This energy corresponds to the transition  $A(n = 4 \text{ to } n = 3)$  for which the energy change = 2 eV

(b) Energy of emitted photon is given by,


$$E = \frac{hc}{\lambda} \therefore \lambda_{\max} \propto \frac{1}{E_{\min}}$$

Transition A, for which the energy emission is minimum, corresponds to the emission of radiation of maximum wavelength.

3.

	Intrinsic Semiconductors	Extrinsic Semiconductors
(i)	These are pure semiconducting tetravalent crystals.	These are semi-conducting tetravalent crystals doped with impurity atoms of group III or V.
(ii)	Their electrical conductivity is low.	Their electrical conductivity is high.
(iii)	There is no permitted energy state between valence and conduction bands.	There is permitted energy state of the impurity atom between valence and conduction bands.

4. Two processes that take place in the formation of a  $p$ - $n$  junction are diffusion and drift.



When  $p$ - $n$  junction is formed, then at the junction free electrons from  $n$ -type diffuse over to  $p$ -type, thereby filling in the holes in  $p$ -type. Due to this a layer of positive charge is built on  $n$ -side and a layer of negative charge is built on  $p$ -side of the  $p$ - $n$  junction. This layer sufficiently grows up within a very short time of the junction being formed, preventing any further movement of charge carriers (*i.e.*, electrons and holes) across the junction. Thus a potential difference  $V_0$  of the order of 0.1 to 0.3 V is set up across the  $p$ - $n$  junction called potential barrier or junction barrier. The thin region around the junction containing immobile positive and negative charges is known as depletion layer.

5. Since  $\frac{hc}{\lambda} = K_{\max} + \phi$

For case I:  $\frac{hc}{\lambda_1} = K + \frac{hc}{\lambda_T} \Rightarrow hc \left[ \frac{1}{\lambda_1} - \frac{1}{\lambda_T} \right] = K \quad \dots(i)$

For case II:  $\frac{hc}{\lambda_2} = 2K + \frac{hc}{\lambda_T} \Rightarrow hc \left[ \frac{1}{\lambda_2} - \frac{1}{\lambda_T} \right] = 2K \quad \dots(ii)$

Substituting (i) in (ii),

$$hc \left[ \frac{1}{\lambda_2} - \frac{1}{\lambda_T} \right] = 2hc \left[ \frac{1}{\lambda_1} - \frac{1}{\lambda_T} \right]$$

$$\Rightarrow \lambda_T = \frac{\lambda_1 \lambda_2}{2\lambda_2 - \lambda_1}$$

6. Given  $m = 2 \text{ kg}$ ,  $P = 800 \text{ W}$ .

Here two deuterium nuclei produce  $3.27 \text{ MeV}$  energy  
 $= 5.232 \times 10^{-13} \text{ J}$

$$\therefore \text{Energy per nuclei} = \frac{5.232 \times 10^{-13}}{2} = 2.616 \times 10^{-13} \text{ J}$$

No. of deuterium atom in  $2 \text{ kg}$

$$= \frac{6.023 \times 10^{23} \times 2000}{2} = 6.023 \times 10^{26} \text{ atom}$$

$$\therefore \text{Total energy} = 6.023 \times 10^{26} \times 2.616 \times 10^{-13}$$

$$= 15.75 \times 10^{13} \text{ J}$$

$$\text{Power} = \frac{\text{Energy}}{\text{Time}} \Rightarrow t = \frac{15.75 \times 10^{13}}{800} = 1.96 \times 10^{11} \text{ s}$$

$$= \frac{1.96 \times 10^{11}}{365 \times 24 \times 60 \times 60} = 6.2 \times 10^3 \text{ years}$$

7. As given

$$E = 10 \cos(10^7 t + kx)$$

Comparing it with standard equation of e.m. wave,

$$E = E_0 \cos(\omega t + kx)$$

Amplitude  $E_0 = 10 \text{ V/m}$  and  $\omega = 10^7 \text{ rad/s}$

$$\therefore c = v\lambda = \frac{\omega\lambda}{2\pi} \text{ or } \lambda = \frac{2\pi c}{\omega} = \frac{2\pi \times 3 \times 10^8}{10^7} = 188.4 \text{ m}$$

$$\text{Also, } c = \frac{\omega}{k} \text{ or } k = \frac{\omega}{c} = \frac{10^7}{3 \times 10^8} = 0.033 \text{ rad/m}$$

The wave is propagating along  $y$  direction.

OR

$$B_x = (4.0 \times 10^{-6} \text{ T}) \sin[(1.57 \times 10^7 \text{ m}^{-1})y + \omega t]$$

Comparing the given equation with

$$B_x = B_0 \sin(ky + \omega t)$$

We get,  $B_0 = 4.0 \times 10^{-6} \text{ T}$

Intensity of light is

$$I = \frac{1}{2\mu_0} B_0^2 c = \frac{(4.0 \times 10^{-6})^2 \times (3 \times 10^8)}{2 \times 4\pi \times 10^{-7}}$$


$$= 1.9 \times 10^3 \text{ W/m}^2 = 1.9 \text{ kW/m}^2$$

8. Optical fibre is made up of very fine quality glass or quartz of refractive index about 1.7.


A light beam incident on one end of an optical fibre at

appropriate angle refracts into the fibre and undergoes repeated total internal reflection.

This is because the angle of incidence is greater than critical angle. The beam of light is received at other end of fibre with nearly no loss in intensity. To send a complete image, the image of different portion is send through separate fibres and thus a complete image can be transmitted through an optical fibre.



9.



Consider a point  $P$  on the screen at distance  $x$  from the centre  $O$ . The nature of the interference at the point  $P$  depends on path difference,

$$p = S_2P - S_1P$$

From right-angled  $\Delta S_2BP$  and  $\Delta S_1AP$ ,

$$(S_2P)^2 - (S_1P)^2 = [S_2B^2 + PB^2] - [S_1A^2 + PA^2]$$

$$= \left[ D^2 + \left( x + \frac{d}{2} \right)^2 \right] - \left[ D^2 + \left( x - \frac{d}{2} \right)^2 \right]$$

$$\text{or } (S_2P - S_1P)(S_2P + S_1P) = 2xd$$

$$\text{or } S_2P - S_1P = \frac{2xd}{S_2P + S_1P}$$

In practice, the point  $P$  lies very close to  $O$ , therefore  $S_1P \approx S_2P \approx D$ .

$$\text{Hence, } p = S_2P - S_1P = \frac{2xd}{2D}$$

$$\text{or } p = \frac{xd}{D}$$

Positions of bright fringes : For constructive interference,

$$p = \frac{xd}{D} = n\lambda$$

$$\text{or } x = \frac{nD\lambda}{D} \text{ where } n = 0, 1, 2, 3, \dots$$

Positions of dark fringes : For destructive interference,

$$p = \frac{xd}{D} = (2n-1) \frac{\lambda}{2}$$

$$\text{or } x = (2n-1) \frac{D\lambda}{2d} \text{ where } n = 1, 2, 3$$

Width of a dark fringe = Separation between two consecutive bright fringes

$$= x_n - x_{n-1} = \frac{nD\lambda}{d} - \frac{(n-1)D\lambda}{d} = \frac{D\lambda}{d}$$

Width of bright fringe = Separation between two consecutive dark fringes

$$= x'_n - x'_{n-1} = (2n-1) \frac{D\lambda}{2d} - [2(n-1)-1] \frac{D\lambda}{2d} = \frac{D\lambda}{d}$$


Clearly, both the bright and dark fringes are of equal width.

Hence the expression for the fringe width in Young's double slit experiment can be written as

$$\beta = \frac{D\lambda}{d}$$

OR

Snell's law of refraction : Let  $PP'$  represents the surface separating medium 1 and medium 2 as shown in figure.



Let  $v_1$  and  $v_2$  represents the speed of light in medium 1 and medium 2 respectively. We assume a plane wavefront  $AB$  propagating in the direction  $A'A$  incident on the interface at an angle  $i$ . Let  $t$  be the time taken by the wavefront to travel the distance  $BC$ .

$$\therefore BC = v_1 t \quad [ \because \text{distance} = \text{speed} \times \text{time} ]$$

In order to determine the shape of the refracted wavefront, we draw a sphere of radius  $v_2 t$  from the point  $A$  in the second medium (the speed of the wave in second medium is  $v_2$ ).

Let  $CE$  represents a tangent plane drawn from the point  $C$ . Then

$$AE = v_2 t$$

$\therefore CE$  would represent the refracted wavefront.

In  $\Delta ABC$  and  $\Delta AEC$ , we have

$$\sin i = \frac{BC}{AC} = \frac{v_1 t}{AC} \quad \text{and} \quad \sin r = \frac{AE}{AC} = \frac{v_2 t}{AC}$$

Where  $i$  and  $r$  are the angles of incident and refraction respectively.

$$\therefore \frac{\sin i}{\sin r} = \frac{v_1 t}{AC} \cdot \frac{AC}{v_2 t}$$

$$\frac{\sin i}{\sin r} = \frac{v_1}{v_2}$$

If  $c$  represents the speed of light in vacuum, then


$$\mu_1 = \frac{c}{v_1} \quad \text{and} \quad \mu_2 = \frac{c}{v_2}$$

$$\Rightarrow v_1 = \frac{c}{\mu_1} \quad \text{and} \quad v_2 = \frac{c}{\mu_2}$$

Where  $\mu_1$  and  $\mu_2$  are the refractive indices of medium 1 and medium 2.

$$\therefore \frac{\sin i}{\sin r} = \frac{c/\mu_1}{c/\mu_2} \Rightarrow \frac{\sin i}{\sin r} = \frac{\mu_2}{\mu_1} \Rightarrow \mu_1 \sin i = \mu_2 \sin r$$

This is the Snell's law of refraction.



The wavelength of emitted radiation from state ( $n = 4$ ) to the state ( $n = 2$ ) is

$$\lambda = \frac{hc}{(E_4 - E_2)} = \frac{6.6 \times 10^{-34} \times 3 \times 10^8}{[0 - (-4.5)] \times 1.6 \times 10^{-19}}$$

$$= 2.75 \times 10^{-7} \text{ m} = 275 \times 10^{-9} \text{ m} = 275 \text{ nm}$$

Hence, transition shown by arrow  $B$  corresponds to emission of wavelength = 275 nm.

(i) The maximum wavelength of emitted radiation from state  $n = 4$  to  $n = 3$  is

$$\lambda = \frac{hc}{[0 - (-2)] \text{ eV}}$$

$$\Rightarrow \lambda = \frac{6.6 \times 10^{-34} \times 3 \times 10^8}{2 \times 1.6 \times 10^{-19}}$$

$$= 6.18 \times 10^{-7} \text{ m} = 618 \times 10^{-9} \text{ m} = 618 \text{ nm}$$

Hence transition  $A$  corresponds to maximum wavelength.

(ii) The minimum wavelength of emitted radiation from state  $n = 3$  to  $n = 1$  is

$$\lambda = \frac{6.6 \times 10^{-34} \times 3 \times 10^8}{[-2 \text{ eV} - (-10 \text{ eV})]} = \frac{6.6 \times 10^{-34} \times 3 \times 10^8}{8 \times 1.6 \times 10^{-19}}$$

$$\lambda = 1.55 \times 10^{-7} \text{ m} = 155 \text{ nm}$$

Hence transition  $D$  corresponds to minimum wavelength.

11. According to lens maker's formula

$$\frac{1}{f} = (\mu - 1) \left( \frac{1}{R_1} - \frac{1}{R_2} \right)$$

$$\frac{1}{f} = (\mu - 1) \left( \frac{1}{R} - \frac{1}{-R} \right) = (1.5 - 1) \left( \frac{2}{R} \right) = \frac{1}{R}$$



Two similar equi-convex lenses of focal length  $f$  each are held in contact with each other.

The focal length  $F_1$  of the combination is given by

$$\frac{1}{F_1} = \frac{1}{f} + \frac{1}{f} = \frac{2}{f}; \quad F_1 = \frac{f}{2} = \frac{R}{2} \quad \dots (i)$$

For glycerin in between lenses, there are three lenses, one concave and two convex.

Focal length of the concave lens is given by

$$\frac{1}{f'} = (1.5 - 1) \left( \frac{-2}{R} \right) = -\frac{1}{R}$$

Now, equivalent focal length of the combination is,

$$\frac{1}{F_2} = \frac{1}{f} + \frac{1}{f'} + \frac{1}{f}; \quad \frac{1}{F_2} = \frac{1}{R} - \frac{1}{R} + \frac{1}{R} = \frac{1}{R}$$

$$F_2 = R$$



... (ii)

Dividing equation (i) by (ii), we get  $\frac{F_1}{F_2} = \frac{1}{2}$

12. (i) (b) : Infrared rays can be converted into electric energy as in solar cell.

(ii) (d) : Radiowaves have longest wavelength.

(iii) (c) : Cathode rays are invisible fast moving streams of electrons emitted by the cathode of a discharge tube which is maintained at a pressure of about 0.01 mm of mercury.

(iv) (c) :  $\gamma$ -rays have minimum wavelength.

(v) (a) :  $\lambda_{\text{micro}} > \lambda_{\text{infra}} > \lambda_{\text{ultra}} > \lambda_{\text{gamma}}$