

# SAMPLE QUESTION PAPER

## BLUE PRINT

Time Allowed : 3 hours

Maximum Marks : 70

S. No.	Chapter	VSA/ AR/ Case Based (1 mark)	SA-I (2 marks)	SA-II (3 marks)	LA (5 marks)	Total
1.	Electrostatics	3(3)	2(4)	1(3)	–	8(16)
2.	Current Electricity	1(1)	–	–	1(5)	
3.	Magnetic Effects of Current and Magnetism	–	–	1(3)	–	7(17)
4.	Electromagnetic Induction and Alternating Current	4(7)	1(2)	–	1(5)	
5.	Electromagnetic Waves	2(2)	1(2)	1(3)	–	9(18)
6.	Optics	3(6)	1(2)	1(3)	–	
7.	Dual Nature of Radiation and Matter	1(1)	1(2)	–	–	6(12)
8.	Atoms and Nuclei	2(2)	1(2)	–	1(5)	
9.	Electronic Devices	–	2(4)	1(3)	–	3(7)
	<b>Total</b>	<b>16(22)</b>	<b>9(18)</b>	<b>5(15)</b>	<b>3(15)</b>	<b>33(70)</b>



# PHYSICS

**Time allowed : 3 hours****Maximum marks : 70**

- (i) All questions are compulsory. There are 33 questions in all.
- (ii) This question paper has five sections: Section A, Section B, Section C, Section D and Section E.
- (iii) Section A contains ten very short answer questions and four assertion reasoning MCQs of 1 mark each. Section B has two case based questions of 4 marks each, Section C contains nine short answer questions of 2 marks each, Section D contains five short answer questions of 3 marks each and Section E contains three long answer questions of 5 marks each.
- (iv) There is no overall choice. However internal choice is provided. You have to attempt only one of the choices in such questions.

## SECTION - A

**All questions are compulsory. In case of internal choices, attempt any one of them.**

1. What is the source of stellar energy?
2. Distinguish between polar and non-polar dielectric.


**OR**

A hollow insulated conduction sphere is given a positive charge of  $10 \mu\text{C}$ . What will be the electric field at the centre of the sphere if its radius is 2 metres ?

3. State Huygens' principle.
4. What is the value of time constant of a C-R circuit?
5. A particle is dropped from a height  $H$ . Find the dependence of the de Broglie wavelength of the particle on height  $H$ .

**OR**

Figure shows a plot of  $\frac{1}{\sqrt{V}}$ , where  $V$  is the accelerating potential, versus the de-Broglie wavelength ' $\lambda$ ' in the case of two particles having same charge ' $q$ ' but different masses  $m_1$  and  $m_2$ . Which line (A or B) represents a particle of large mass?



6. Define ionisation energy. What is its value for a hydrogen atom?



7. What is the electric flux linked with closed surface given in the figure?



8. Why is the penetrating power of gamma rays very large?

OR

What are the directions of electric and magnetic field vectors relative to each other and relative to the direction of propagation of electromagnetic waves?

9. Define 1 ampere current in terms of number of electrons flowing per second.
10. An ac voltage is applied to a resistance  $R$  and an inductor  $L$  in series. If  $R$  and the inductive reactance are both equal to  $3 \Omega$ , calculate the phase difference between the applied voltage and the current in the circuit.

OR

An series  $L$ - $C$ - $R$  circuit is connected to a source of A.C. current. At resonance, what will be the phase difference between the applied voltage and the current in the circuit?

**For question numbers 11, 12, 13 and 14, two statements are given-one labelled Assertion (A) and the other labelled Reason (R). Select the correct answer to these questions from the codes (a), (b), (c) and (d) as given below.**

- (a) Both A and R are true and R is the correct explanation of A  
(b) Both A and R are true but R is NOT the correct explanation of A  
(c) A is true but R is false  
(d) A is false and R is also false
11. **Assertion (A)** : Gauss's law shows diversion when inverse square law is not obeyed.  
**Reason (R)** : Gauss's law is a consequence of conservation of charges.
12. **Assertion (A)** : A ray of light is incident from outside on a glass sphere surrounded by air. This ray may suffer total internal reflection at second interface.  
**Reason (R)** : If a ray of light goes from denser to rarer medium, it bends away from the normal.
13. **Assertion (A)** : The possibility of an electric bulb fusing is lower at the time of switching on and off.  
**Reason (R)** : Inductive effects produce a decline in current at the time of switch-off and switch-on
14. **Assertion (A)** : Accelerated charge radiate electromagnetic waves.  
**Reason (R)** : As the wave propagate through the space, the oscillating electric and magnetic field regenerate each other.

## SECTION - B

**Questions 15 and 16 are Case Study based questions and are compulsory. Attempt any 4 sub parts from each question. Each question carries 1 mark.**

15. Optical power (also referred to as dioptric power, refractive power, focusing power, or convergence power) is the degree to which a lens, mirror, or other optical system converges or diverges light. The SI unit for optical power is the inverse meter ( $\text{m}^{-1}$ ), which is commonly called the dioptre. Converging lenses have positive optical power, while diverging lenses have negative power. When a lens is immersed in a refractive medium, its optical power and focal length change.



- (i) What is the relation between the object distance  $u$ , image distance  $v$  and power  $P$  in a lens?
- (a)  $\frac{uv}{u-v}$                       (b)  $\frac{u-v}{uv}$                       (c)  $\frac{u}{v}$                       (d)  $\frac{u+v}{u-v}$
- (ii) The radius of curvature of each surface of a convex lens of refractive index 1.5 is 40 cm. Its power is
- (a) 2.5 D                      (b) 2 D                      (c) 1.5 D                      (d) 1 D
- (iii) A convex lens of focal length 20 cm made of glass is immersed in a liquid of refractive index 2.3. Its power will be
- (a) positive  
(b) can be positive or Negative depending on the depth  
(c) zero  
(d) negative.
- (iv) Focal length of a convex lens of refractive index 1.5 is 2 cm. Focal length of lens when immersed in a liquid of refractive index of 1.25 will be
- (a) 5 cm                      (b) 4 cm                      (c) 2.5 cm                      (d) 1.2 cm
- (v) A convex lens of focal length 15 cm is placed on a plane mirror. An object is placed at 30 cm from the lens. The image is
- (a) real, at 30 cm in front of the mirror                      (b) real, at 30 cm behind the mirror  
(c) real, at 10 cm in front of the mirror                      (d) virtual, at 10 cm behind the mirror.
- 16.** A thermal power plant produced electric power of 600 kW at 4000 V, which is to be transported to a place 20 km away from the power plant for consumer's usage. It can be transported either directly with a cable of large current carrying capacity or by using a combination of step-up and step-down transformers at the two ends. The drawback of the direct transmission is the large energy dissipation. In the method using transformers, the dissipation is much smaller. In this method a step-up transformer is used at the plant side so that the current is reduced to a smaller value. At the consumers 'end, a step-down transformer is used to supply power to the consumers at the specified lower voltage. It is reasonable to assume that the power cable is purely resistive, and the transformers are ideal with power factor unity. All the currents and voltage mentioned are rms values.
- In the method using the transformers assume that the ratio of the number of turns in the primary to that in the secondary in the step-up transformer is 1 : 10 and the power to the consumers has to be supplied at 200 V.
- (i) The voltage induced in the primary coil is
- (a) 4000 V                      (b) 40000 V                      (c) 200 V                      (d) 2000 V
- (ii) If the direct transmission method with a cable of resistance  $0.4 \Omega \text{ km}^{-1}$  is used, the total current in the transmission line is
- (a) 150 A                      (b) 100 A                      (c) 40 A                      (d) 200 A
- (iii) The power dissipation (in %) during transmission is
- (a) 20                      (b) 30                      (c) 40                      (d) 50
- (iv) The ratio of the number of turns in the primary to that in the secondary in the step-down transformer is
- (a) 200 : 1                      (b) 150 : 1                      (c) 100 : 1                      (d) 50 : 1
- (v) The purpose of providing an iron core in a transformer is to
- (a) provide support to windings                      (b) reduce hysteresis loss  
(c) decrease the reluctance of the magnetic path                      (d) reduce eddy current losses




## SECTION - C

All questions are compulsory. In case of internal choices, attempt anyone.

17. A charged conical conductor loses its charge earlier than a similarly charged sphere. Why?
18. Name the electromagnetic waves used for studying crystal structure of solids. What is its frequency range?
19. Draw a ray diagram showing the image formation by an astronomical telescope when the final image is formed at infinity.
20. Consider the fusion reaction:  ${}^4\text{He} + {}^4\text{He} \rightarrow {}^8\text{Be}$   
For the reaction, find (i) mass defect (ii) Q-value (iii) Is such a fusion energetically favourable?  
Atomic mass of  ${}^8\text{Be}$  is 8.0053 u and that of  ${}^4\text{He}$  is 4.0026 u.


OR

In the following reaction, determine the energy released.



Given : Mass of  ${}^1_1\text{H} = 1.007825$  u, Mass of  ${}^4_2\text{He} = 4.002603$  u, Mass of  $e^+ = 0.000548$  u

21. If photons of frequency  $\nu$  are incident on the surfaces of metals A and B of threshold frequencies  $\nu/2$  and  $\nu/3$  respectively, find the ratio of the maximum kinetic energy of electrons emitted from A to that from B.
22. A battery of emf 2 V is applied across the block of a semiconductor. The length of the block is 0.1 m and the area of cross-section is  $1 \times 10^{-4} \text{ m}^2$ . If the block is of intrinsic silicon at 300 K, find the electron and hole currents. [Take  $n_e = n_h = 1.5 \times 10^{16} / \text{m}^3$ ;  $\mu_e = 0.135 \text{ m}^2\text{V}^{-1}\text{s}^{-1}$ ;  $\mu_h = 0.048 \text{ m}^2\text{V}^{-1}\text{s}^{-1}$ ]
23. What is the effective capacitance between points X and Y?




OR

An electric dipole of moment  $\vec{p}$  is lying along a uniform electric field  $\vec{E}$ . Find the work done in rotating the dipole by  $90^\circ$ .


24. What are eddy currents? Write any two applications of eddy currents.

OR

A bar magnet is moved in the direction indicated by the arrow between two coils PQ and CD. Predict the direction of the induced current in each coil.



25. In the given circuit, find the current through the battery.



## SECTION - D

All questions are compulsory. In case of internal choices, attempt any one.

26. How does an oscillating charge produce electromagnetic wave? Explain. Draw a sketch showing the propagation of plane *e.m.* wave along the *Z*-direction, clearly depicting the directions of oscillating electric and magnetic field vectors.
27. A solenoid has a core of a material with relative permeability 400. The windings of the solenoid are insulated from the core and carry a current of 2 A. If the number of turns is 1000 per metre, calculate (i) *H* (ii) *B* (iii) the magnetising current,  $I_m$ .

OR


Where on the earth's surface is the value of vertical component of the earth's magnetic field zero? The horizontal component of the earth's magnetic field at a given place is  $0.4 \times 10^{-4} \text{ Wb m}^{-2}$  and angle of dip is  $30^\circ$ . Calculate the value of (i) vertical component, (ii) the total intensity of the earth's magnetic field.

28. A convex lens made of a material of refractive index  $\mu_1$  is kept in a medium of refractive index  $\mu_2$ . Parallel rays of light are incident on the lens. Complete the path of rays of light emerging from the convex lens if
- (a)  $\mu_1 > \mu_2$  (b)  $\mu_1 = \mu_2$   
(c)  $\mu_1 < \mu_2$
29. Mobilities of electrons and holes in a sample of intrinsic germanium at room temperature are  $0.36 \text{ m}^2 \text{ V}^{-1} \text{ s}^{-1}$  and  $0.17 \text{ m}^2 \text{ V}^{-1} \text{ s}^{-1}$ . The electron and hole densities are each equal to  $2.5 \times 10^{19} \text{ m}^{-3}$ . Determine the electrical conductivity of germanium.


OR

Write any two distinguishing features between conductors, semiconductors and insulators on the basis of energy band diagrams.

30. An electric dipole is kept first to the left and then to the right of a negatively charged infinite plane sheet having a uniform surface charge density. The arrows  $p_1$  and  $p_2$  show the directions of its electric dipole moment in the two cases. (a) Identify for each case, whether the dipole is in stable or unstable equilibrium. Justify each answer.



- (b) Next, the dipole is kept in a similar way (as shown), near an infinitely long straight wire having uniform negative linear charge density. Will the dipole be in equilibrium at these two positions? Justify your answer.



## SECTION - E

All questions are compulsory. In case of internal choices, attempt any one.

31. A gas of hydrogen like atoms can absorb radiations of 68 eV. Consequently, the atoms emit radiations of only three different wavelength. All the wavelengths are equal or smaller than that of the absorbed photon.
- Determine the initial state of the gas atoms.
  - Identify the gas atoms.
  - Find the minimum wavelength of the emitted radiations.
  - Find the ionization energy and the respective wavelength for the gas atoms.

OR

Write shortcomings of Rutherford's atomic model. Explain how these were overcome by the postulates of Bohr's atomic model.

32. A  $2\ \mu\text{F}$  capacitor,  $100\ \Omega$  resistor and  $8\ \text{H}$  inductor are connected in series with an ac source
- What should be the frequency of the source such that current drawn in the circuit is maximum? What is this frequency called?
  - If the peak value of emf of the source is  $200\ \text{V}$ , find the maximum current.
  - Draw a graph showing variation of amplitude of circuit current with changing frequency of applied voltage in a series  $LCR$  circuit for two different values of resistance  $R_1$  and  $R_2$  ( $R_1 > R_2$ ).
  - Define the term 'Sharpness of Resonance'. Under what condition, does a circuit become more selective?

OR


A device  $X$  is connected across an ac source of voltage  $V = V_0 \sin \omega t$ . The current through  $X$  is given as

$$I = I_0 \sin \left( \omega t + \frac{\pi}{2} \right).$$

- Identify the device  $X$  and write the expression for its reactance.
  - Draw graphs showing variation of voltage and current with time over one cycle of ac, for  $X$ .
  - How does the reactance of the device  $X$  vary with frequency of the ac? Show this variation graphically.
  - Draw the phasor diagram for the device  $X$ .
33. State the two Kirchhoff's laws. Explain briefly how these rules are justified.

OR

- Space between two concentric spheres of radii  $r_1$  and  $r_2$ , such that  $r_1 < r_2$ , is filled with a material of resistivity  $\rho$ . Find the resistance between inner and outer surface of the material.



- The current density varies with radial distance  $r$  as  $J = ar^2$ , in a cylindrical wire of radius  $R$ . Determine the current passing through the wire between radial distance  $R/3$  and  $R/2$ .
- A wire whose cross-sectional area is increasing linearly from its one end to the other, is connected across a battery of  $V$  volts. Which of the following quantities remain constant in the wire?  
(a) drift speed      (b) current density      (c) electric current      (d) electric field

Justify your answer.



# SOLUTIONS

1. Nuclear fusion.

2. A dielectric whose molecules possess electric moment even when electric field is not applied is called polar dielectric. On the other hand, a dielectric whose molecules do not possess permanent dipole moment, is called non-polar dielectric.

OR

Field inside a conducting sphere = 0.

3. According to Huygens' principle, each point on a wavefront is a source of secondary waves, which add up to give a wavefront at any later time.

4. The time constant for resonance circuit,  $\tau = CR$

5. Velocity acquired by a particle while falling from a height  $H$  is

$$v = \sqrt{2gH} \quad \dots (i)$$

As  $\lambda = \frac{h}{mv} = \frac{h}{m\sqrt{2gH}}$  (Using (i))

OR

de-Broglie wavelength,  $\lambda = \frac{h}{\sqrt{2mqV}}$

or  $\lambda = \frac{h}{\sqrt{2mq}} \cdot \frac{1}{\sqrt{V}}$ . The graph of  $\lambda$  versus  $\frac{1}{\sqrt{V}}$  is

a straight line of slope  $\frac{h}{\sqrt{2mq}}$ . The slope of line  $B$  is smaller, so particle  $B$  has larger mass (charge is same).

6. Ionisation energy for an atom is defined as the energy required to remove an electron completely from the outermost shell of the atom.

The ionisation energy required to remove an electron from ground state of hydrogen atom is

$$E = E_{\infty} - E_1 = 0 - (-13.6) \text{ eV} = 13.6 \text{ eV}$$

7. According to Gauss's law

$$\text{Electric flux, } \phi = \frac{q}{\epsilon_0}$$

where,  $q$  = total charge enclosed by closed surface

$$\begin{aligned} \therefore \phi &= \frac{1.25 + 7 + 1 - 0.4}{\epsilon_0} \\ &= \frac{8.85 \text{ C}}{8.85 \times 10^{-12} \text{ C}^2 \text{ N}^{-1} \text{ m}^{-2}} = 10^{12} \text{ N m}^2 \text{ C}^{-1} \end{aligned}$$

8.  $\gamma$ -rays are em waves of very large energy, so they pass through the matter with less number of collisions with atoms or molecules of matter, thereby penetrating more into it.

OR

In an electromagnetic wave  $\vec{E}, \vec{B}$  and direction of propagation are mutually perpendicular.

9.  $Q = It$  Also  $Q = ne$  [ $e = 1.6 \times 10^{-19} \text{ C}$ ]

$$\begin{aligned} \therefore ne &= It \quad \text{or} \quad n = \frac{It}{e} = \frac{1 \text{ A} \times 1 \text{ s}}{1.6 \times 10^{-19}} \\ &= 6.25 \times 10^{18} \text{ electrons} \end{aligned}$$

10. (b) : Here,  $R = 3 \Omega$

Inductive reactance,  $X_L = 3 \Omega$

The phase difference between the applied voltage and the current in the circuit is

$$\tan \phi = \frac{X_L}{R} = \frac{3 \Omega}{3 \Omega} = 1$$

$$\phi = \tan^{-1}(1) \quad \text{or} \quad \phi = \frac{\pi}{4}$$

OR

For resonance condition, the impedance will be minimum and the current will be maximum. This is only possible when  $X_L = X_C$ .

Therefore,  $\tan \theta = \frac{X_L - X_C}{R} = 0$  or  $\theta = 0$ .

11. (b)

12. (b) : When a ray of light enters a spherical glass sphere, it is first refracted at first interface and then it strikes the inner surface of sphere (second interface) and gets totally internally reflected if the angle between the refracted ray and normal to the drop surface is greater than the critical angle ( $42^\circ$ , in this case).

13. (d)

14. (a) : Consider a charge oscillating with some frequency. This oscillating charge produces an oscillating electric field in space, which produces an oscillating magnetic field, which in turn, is a source of oscillating electric field and so on. The oscillating electric and magnetic field thus regenerate each other, as the wave propagates through the space.

15. (i) (b) : From lens maker formula,

$$\frac{1}{f} = \frac{1}{v} - \frac{1}{u} \Rightarrow f = \frac{uv}{u-v}$$

Since power,  $P = \frac{1}{f} \Rightarrow P = \frac{u-v}{uv}$



(ii) (a) : Power,  $P = \frac{1}{f} = (\mu - 1) \left[ \frac{1}{R_1} - \frac{1}{R_2} \right]$

Here,  $\mu = 1.5$ ,  $R_1 = 40 \text{ cm} = 0.4 \text{ m}$ ,  $R_2 = -0.4 \text{ m}$

$$P = (1.5 - 1) \left( \frac{1}{0.4} + \frac{1}{0.4} \right) = 2.5 \text{ D}$$

(iii) (d) : Given focal length,  $f_a = 20 \text{ cm}$

In air,

$$\frac{1}{f_a} = ({}^a\mu_g - 1) \left( \frac{1}{R_1} - \frac{1}{R_2} \right) \quad \dots(i)$$

In liquid

$$\frac{1}{f_l} = ({}^l\mu_g - 1) \left( \frac{1}{R_1} - \frac{1}{R_2} \right) \quad \dots(ii)$$

Dividing (i)/(ii),

$$\frac{1}{f_a} = \frac{({}^a\mu_g - 1)}{({}^l\mu_g - 1)} \Rightarrow \frac{f_l}{20} = \frac{(1.5 - 1)}{(2.3 - 1)}$$

$$\Rightarrow f_l = \frac{20(0.5)}{-0.348} = -28.75 \text{ cm}$$

$$\therefore P = \frac{100}{f_l} = \frac{-100}{28.75} = -3.4 \text{ D}$$

(iv) (a) : Given  $\mu_g = 1.5$ ,  $f_a = 2 \text{ cm}$ ,  $g_l = 1.25$

In air,

$$\frac{1}{f_a} = ({}^a\mu_g - 1) \left( \frac{1}{R_1} - \frac{1}{R_2} \right)$$

In liquid,

$$\frac{1}{f_l} = ({}^l\mu_g - 1) \left( \frac{1}{R_1} - \frac{1}{R_2} \right)$$

Dividing (i)/(ii),

$$\frac{f_l}{f_a} = \frac{{}^a\mu_g - 1}{{}^l\mu_g - 1} \Rightarrow f_l = \frac{2(1.5 - 1)}{\frac{1.5}{1.25} - 1} = 5 \text{ cm}$$

(v) (c) :  $\frac{1}{F} = \frac{1}{f_l} + \frac{1}{f_m} + \frac{1}{f_l} = \frac{2}{f_l} + \frac{1}{\infty} \Rightarrow F = \frac{15}{2} \text{ cm}$

$$\frac{2}{15} = \frac{1}{30} + \frac{1}{v}, v = 10 \text{ cm.}$$

$\therefore$  image is real, 10 cm in front of mirror.

16. (i) (b) : For step up transformer,

$$\frac{N_s}{N_p} = \frac{V_s}{V_p} \Rightarrow \frac{10}{1} = \frac{V_s}{4000}$$

$$\Rightarrow V_s = 4000 \times 10 = 40,000 \text{ V}$$

(ii) (a) : Total resistance =  $0.4 \times 20 = 8 \Omega$

Power,  $P = VI$

$$600 \times 10^3 = 4000 \times I = 150 \text{ A.}$$

Physics

(iii) (b) : Power,  $P = I^2 R$

$$= 150^2 \times 8 = 180 \text{ kW}$$

$$\% \text{ loss is power} = \frac{180}{600} \times 100 = 30\%$$

(iv) (a) : For step down transformer,


$$\frac{N_p}{N_s} = \frac{V_p}{V_s} = \frac{40,000}{200} = \frac{200}{1}$$

(v) (d)

17. In case of a conical conductor, the charge density is maximum at its pointed end. Due to this large density, the charge leaks away from this end. But in case of a similarly charged sphere, the charge density is uniform over its surface. Thus, the charged conical conductor loses its charge earlier than a similarly charged sphere.

18. X-rays,  $3 \times 10^{16} \text{ Hz}$  to  $3 \times 10^{20} \text{ Hz}$ .

19.



(i) 20.  ${}^4\text{He} + {}^4\text{He} \rightarrow {}^8\text{Be} + Q$

$$(i) \Delta m = 2 \times 4.0026 - 8.0053 = 8.0052 - 8.0053 = -0.0001 \text{ amu}$$


(ii)  $Q = (2m_{\text{He}} - m_{\text{Be}}) c^2 = (2 \times 4.0026 - 8.0053) \times 931 \text{ MeV}$

$$= -93.1 \text{ keV}$$

(iii) Since  $Q$  is negative, the fusion is not energetically favourable.

OR

The given nuclear reaction is



The energy released during the process is

$$Q = [4m({}^1_1\text{H}) - m({}^4_2\text{He}) - 2m(e^+)]c^2 = [4 \times 1.007825 - 4.002603 - 2 \times 0.000548]u \times c^2 = (0.027601)(931.5) \text{ MeV} = 25.7 \text{ MeV}$$

21.  $E = h\nu = h\nu_{0A} + (\text{KE})_{\text{max } A} = h\nu_{0B} + (\text{KE})_{\text{max } B}$

$$\therefore h\nu = \frac{h\nu}{2} + K_A = \frac{h\nu}{3} + K_B$$

$$\Rightarrow \frac{K_A}{K_B} = \frac{h\nu}{2} \times \frac{3}{2h\nu} = \frac{3}{4}$$

22. For silicon:

$$\text{Electron current, } I_e = \frac{n_e A e \mu_e V}{l}$$

$$= \frac{1.5 \times 10^{16} \times 10^{-4} \times 1.6 \times 10^{-19} \times 0.135 \times 2}{0.1}$$


$$= 6.48 \times 10^{-7} \text{ A}$$

$$\text{Hole current, } I_h = \frac{n_h A e \mu_h V}{l}$$

$$= \frac{1.5 \times 10^{16} \times 10^{-4} \times 1.6 \times 10^{-19} \times 0.048 \times 2}{0.1}$$

$$= 2.30 \times 10^{-7} \text{ A}$$

23. The given circuit can be simplified as



OR

Work done in deflecting a dipole through an angle  $\theta$  is given by

$$W = \int_0^\theta pE \sin \theta d\theta = pE(1 - \cos \theta)$$

Since  $\theta = 90^\circ$


$$\therefore W = pE(1 - \cos 90^\circ) \text{ or, } W = pE.$$

24. When magnetic flux linked with a metallic conductor changes, then induced currents are set up in the conductor in the form of closed loops known as 'Eddy currents'.

Two applications of eddy currents are


- (i) electromagnetic braking in trains
- (ii) induction furnace

OR



According to Lenz's law, direction of current in loop PQ is from P to Q and in loop CD is from C to D.

25. In the given circuit, diode  $D_1$  is reverse biased, so it will not conduct. Diodes  $D_2$  and  $D_3$  are forward biased, so they will conduct. The corresponding equivalent circuit is as shown in the figure. The equivalent resistance of the circuit is




$$R_{eq} = \frac{(5+5) \times 20}{(5+5) + 20} = \frac{10 \times 20}{10 + 20} = \frac{200}{30} = \frac{20}{3} \Omega$$

$$\text{Current through the battery, } I = \frac{10 \text{ V}}{\frac{20}{3} \Omega} = 1.5 \text{ A}$$

26. An oscillating or accelerated charge is supposed to be source of an electromagnetic wave. An oscillating charge produces an oscillating electric field in space which further produces an oscillating magnetic field which in turn is a source of electric field. These oscillating electric and magnetic field hence, keep on regenerating each other and an electromagnetic wave is produced.

The *e.m.* wave propagates along *z*-axis.



For an *e.m.* wave propagating in *Z*-direction, electric field is directed along *X*-axis and magnetic field is directed along *Y*-axis.

$$\hat{k} = \hat{i} \times \hat{j}$$

27. Here,  $\mu_r = 400$ ,  $I = 2 \text{ A}$ ,  $n = 1000$  per metre

(i)  $H = nI = 1000 \times 2 = 2 \times 10^3 \text{ A m}^{-1}$

(ii)  $B = \mu H = \mu_0 \mu_r H = 4\pi \times 10^{-7} \times 400 \times (2 \times 10^3)$   
 $= 1.0 \text{ T}$

(iii) The magnetising current  $I_m$  is the additional current that needs to be passed through the windings of the solenoid in the absence of the core, which would produce a  $B$  value as in the presence of the core.

Thus,  $B = \mu_0 n(I + I_m)$

$$1.0 = 4\pi \times 10^{-7} \times 1000 \times (2 + I_m)$$

$$I_m = \frac{1.0}{4\pi \times 10^{-4}} - 2 = 796 - 2 = 794 \text{ A}$$

OR

Vertical component of earth's magnetic field is zero at the equator.

$$B_H = 0.4 \times 10^{-4} \text{ Wb m}^{-2}$$

$$\delta = 30^\circ$$

(i)  $B_V = B \sin \delta$

$$B_H = B \cos \delta$$


$$B = \frac{B_H}{\cos \delta}$$

$$B_V = \frac{B_H \sin \delta}{\cos \delta} = B_H \tan \delta$$

$$B_V = 0.4 \times 10^{-4} \times \tan 30^\circ = \frac{0.4 \times 10^{-4}}{\sqrt{3}}$$

$$= 0.23 \times 10^{-4} \text{ Wb m}^{-2}$$


(ii)  $B = \frac{B_H}{\cos \delta} = \frac{0.4 \times 10^{-4}}{\cos 30^\circ} = 0.46 \times 10^{-4} \text{ Wb m}^{-2}$



28. In case (a)  $\mu_1 > \mu_2$ , the lens behaves as convergent lens.

In case (b)  $\mu_1 = \mu_2$ , the lens behaves as a plane plate.

In case (c)  $\mu_1 < \mu_2$ , the lens behaves as a divergent lens. The path of rays in all the three cases is shown in the figure.



29. The electrical conductivity of germanium is

$$\sigma = e[n_e\mu_e + n_h\mu_h]$$


Here,  $\mu_e = 0.36 \text{ m}^2 \text{ V}^{-1} \text{ s}^{-1}$ ,  $\mu_h = 0.17 \text{ m}^2 \text{ V}^{-1} \text{ s}^{-1}$ ,

$$n_e = n_h = 2.5 \times 10^{19} \text{ m}^{-3}$$

$$\begin{aligned} \therefore \sigma &= (1.6 \times 10^{-19} \text{ C})[(2.5 \times 10^{19} \text{ m}^{-3})(0.36 \text{ m}^2 \text{ V}^{-1} \text{ s}^{-1}) \\ &\quad + (2.5 \times 10^{19} \text{ m}^{-3})(0.17 \text{ m}^2 \text{ V}^{-1} \text{ s}^{-1})] \\ &= 2.12 \text{ S m}^{-1} \end{aligned}$$

OR

The band diagram for conductors, semiconductors and insulators are given as follows:



Two distinguishing features :

Physics

(i) In conductors, the valence band and conduction band tend to overlap (or nearly overlap) while in insulators they are separated by a large energy gap and in semiconductors they are separated by a small energy gap.

(ii) The conduction band of a conductor has a large number of electrons available for electrical conduction. However, the conduction band of insulators is almost empty while that of the semiconductor has only a (very) small number of such electrons available for electrical conduction.

30. (a)  $p_1$ : stable equilibrium  $p_2$ : unstable equilibrium  
The electric field, on either side, is directed towards the negatively charged sheet and its magnitude is independent of the distance of the field point from the sheet.

For position  $p_1$ , dipole moment and electric field are parallel, while for position  $p_2$ , they are antiparallel. Hence,  $p_1$  is in stable equilibrium and  $p_2$  is in unstable equilibrium.

(b) The dipole will not be in equilibrium in any of the two positions.

The electric field due to an infinite straight charged wire is non-uniform ( $E \propto 1/r$ ).

Hence there will be a net non-zero force on the dipole in each case.

31. (i)  $\frac{n(n-1)}{2} = 3 \quad \therefore n = 3$

i.e. after excitation, atom jumps to second excited state.

Hence,  $n_f = 3$ . So  $n_i$  can be 1 or 2.

If  $n_i = 1$  then energy emitted is either equal to, greater than or less than the energy absorbed. Hence, the emitted wavelength is either equal to, less than or greater than the absorbed wavelength.

Hence,  $n_i \neq 1$

If  $n_i = 2$ , then  $E_e \geq E_a$ . Hence  $\lambda_e \leq \lambda_0$

(ii)  $E_3 - E_2 = 68 \text{ eV}$

$$\therefore (13.6)(Z^2)\left(\frac{1}{4} - \frac{1}{9}\right) = 68$$

$$\therefore Z = 6$$

(iii)  $\lambda_{\min} = \frac{12400}{E_3 - E_1} = \frac{12400}{(13.6)(6)^2\left(1 - \frac{1}{9}\right)}$

$$= \frac{12400}{435.2} = 28.49 \text{ \AA}$$

(iv) Ionisation energy =  $(13.6)(6^2) = 489.6 \text{ eV}$


$$\lambda = \frac{12400}{489.6} = 25.33 \text{ \AA}$$

OR

Limitation of Rutherford's model :

Rutherford's atomic model is inconsistent with classical physics.

According to electromagnetic theory, an electron is a charged particle moving in the circular orbit around the nucleus and is accelerated, so it should emit radiation continuously and thereby lose energy. Due to this, radius of the electron would decrease continuously and also the atom should then produce continuous spectrum, and ultimately electron will fall into the nucleus and atom will collapse in  $10^{-8}$  s. But the atom is fairly stable and it emits line spectrum.



(ii) Rutherford's model is not able to explain the spectrum of even most simplest H-spectrum.

Bohr's postulates to resolve observed features of atomic spectrum :

(i) Quantum condition: Of all the possible circular orbits allowed by the classical theory, the electrons are permitted to circulate only in those orbits in which the angular momentum of an electron is an integral multiple of  $\frac{h}{2\pi}$ ,  $h$  being Planck's constant. Therefore, for any permitted orbit,

$$L = mvr = \frac{nh}{2\pi}, \quad n = 1, 2, 3, \dots,$$

where  $n$  is called the principal quantum number, and this equation is called Bohr's quantisation condition.

(ii) Stationary orbits: While revolving in the permissible orbits, an electron does not radiate energy. These non-radiating orbits are called stationary orbits.

(iii) Frequency condition: An atom can emit or absorb radiation in the form of discrete energy photons only when an electron jumps from a higher to a lower orbit or from a lower to a higher orbit, respectively.

$$h\nu = E_i - E_f$$

where  $\nu$  is frequency of radiation emitted,  $E_i$  and  $E_f$  are the energies associated with stationary orbits of principal quantum number  $n_i$  and  $n_f$  respectively (where  $n_i > n_f$ ).

32. (i) For maximum current, the impedance should be minimum, i.e.,  $X_L = X_C$ .

In such a condition the frequency is called as resonance frequency and is given by


$$\omega = \frac{1}{\sqrt{LC}}$$

Given:  $L = 8\text{H}$  and  $C = 2 \times 10^{-6}\text{F}$

$$f = \frac{1}{2\pi\sqrt{LC}} = \frac{1}{2 \times 3.14 \times \sqrt{8 \times 2 \times 10^{-6}}} = 39.88 \text{ s}^{-1}$$

(ii) Maximum current  $I = \frac{\mathcal{E}_0}{R} = \frac{200}{100} = 2\text{A}$

(iii)



(iv) Sharpness of resonance : It is defined as the ratio of the voltage developed across the inductance ( $L$ ) or capacitance ( $C$ ) at resonance to the voltage developed across the resistance ( $R$ ).


$$Q = \frac{1}{R} \sqrt{\frac{L}{C}}$$

It may also be defined as the ratio of resonant angular frequency to the bandwidth of the circuit.

$$Q = \frac{\omega_r}{2\Delta\omega}$$

Circuit become more selective if the resonance is more sharp, maximum current is more, the circuit is close to resonance for smaller range of ( $2\Delta\omega$ ) of frequencies. Thus, the tuning of the circuit will be good.

Figure shows the variation of  $i_m$  with  $\omega$  in a  $LCR$  series circuit for two values of resistance  $R_1$  and  $R_2$  ( $R_1 > R_2$ ),



The condition for resonance in the  $LCR$  circuit is,

$$X_L = X_C \Rightarrow \omega_0 L = \frac{1}{\omega_0 C} \Rightarrow \omega_0 = \frac{1}{\sqrt{LC}}$$

We see that the current amplitude is maximum at the resonant frequency. Since  $i_m = V_m / R$  at resonance, the current amplitude for case  $R_2$  is sharper to that for case  $R_1$ .

OR

(a) Here device  $X$  is a capacitor.

Capacitive reactance,  $X_C = \frac{1}{\omega C} = \frac{1}{2\pi\nu C}$

(b)  $V = V_0 \sin \omega t$

$C = \frac{q}{V}$

$q = CV_0 \sin \omega t$




$i = \frac{dq}{dt} = \frac{d}{dt}(CV_0 \sin \omega t)$

$= \omega CV_0 \cos \omega t = \frac{V_0}{\frac{1}{\omega C}} \cos \omega t$


$\therefore i = \frac{V_0}{X_C} \sin \left( \omega t + \frac{\pi}{2} \right)$

or  $i = i_0 \sin \left( \omega t + \frac{\pi}{2} \right)$

In pure capacitive circuit current leads voltage by  $\frac{\pi}{2}$ .



33. Kirchhoff's first rule : The algebraic sum of all the current passing through a junction of an electric circuit is zero.



Here,  $I_1, I_2, I_3, I_4$  and  $I_5$  are current in different branches of a circuit which meet at a junction.

$I_1 + I_2 - I_3 + I_4 - I_5 = 0$

This rule is based on the principle of conservation of charge.

Kirchhoff's second rule : The algebraic sum of the applied emf's of an electrical circuit is equal to the algebraic sum of potential drops across the resistors of the loop.


Mathematically,

$\Sigma \epsilon = \Sigma IR$

This is based on energy conservation principle

Using this rule,

$\epsilon_1 - \epsilon_2 = IR_1 + IR_2$



OR

(i) Since  $R = \rho \frac{l}{a}$   $\therefore R = \rho \frac{dl}{4\pi l^2}$ .

(where  $l$  is any radius and  $dl$  is small element).

Total resistance,

$R = \frac{\rho}{4\pi} \int_{r_1}^{r_2} \frac{dl}{l^2} = \frac{\rho}{4\pi} \left[ -\frac{1}{l} \right]_{r_1}^{r_2} = \frac{\rho}{4\pi} \left[ \frac{1}{r_1} - \frac{1}{r_2} \right]$ .

$R = \left[ \frac{r_2 - r_1}{r_1 r_2} \right] \frac{\rho}{4\pi}$ .


(ii) Current density,  $J = ar^2$

Current,  $I = \int J dA$

$I = \int_{R/3}^{R/2} ar^2 (2\pi r dr)$ , [where  $dA = 2\pi r dr$ ]

$= 2\pi a \left[ \frac{r^4}{4} \right]_{R/3}^{R/2} = \frac{65\pi a R^4}{2592}$

(iii) Electric current : Through the wire flow of charge through (a non-uniform conductor) a conductor is same, hence current remains constant.



As area of cross-section of the conductor is varying so current density through wire and drift velocity of electron will not be same.

