

CLASS – XI
CHEMISTRY
SAMPLE PAPER 3

Time: 3 Hours

Marks: 70

General Instructions

- All questions are compulsory.
 - Section A: Q.no. 1 to 5 are very short answer questions and carry 1 mark each.
 - Section B: Q.no. 6 to 12 are short answer questions and carry 2 marks each.
 - Section C: Q.no. 13 to 24 are also short answer questions and carry 3 marks each.
 - Section D: Q.no. 25 to 27 are long answer questions and carry 5 marks each.
 - There is no overall choice. However an internal choice has been provided in two questions of one mark, two questions of two marks, four questions of three marks and all the three questions of five marks weightage. You have to attempt only one of the choices in such questions.
 - Use of log tables if necessary, use of calculators is not allowed.
-

Section A

1. Arrange the following in the order of decreasing reactivity towards alkenes. [1]
HCl, HBr, HI, HF
2. Predict the shapes of the following molecules using VSEPR theory? [1]
BeCl₂, SiCl₄

OR

What is meant by bond pairs of electrons?

3. Give IUPAC name of allyl alcohol. [1]
4. What is biochemical oxygen demand? [1]

OR

What is the name of the compound formed when CO combines with blood?

5. Give the colours shown in the flame test by sodium and potassium. [1]





Section B



6. How does metallic and nonmetallic character vary on moving from left to right in the periodic table? [2]
7. Arrange the following in increasing order of size. Give reason for your answer. [2]
 Mg^{2+} O^{2-} Na^+ F^- Al^{3+} .
8. Calculate the wavelength of an electron moving with a velocity of $2.05 \times 10^7 \text{ m/s}$.
(Mass of electron = $9.1 \times 10^{-31} \text{ kg}$, $h = 6.63 \times 10^{-34} \text{ Js}$) [2]
9. At 273 K the density of a gaseous oxide at 2 bar is the same as that of nitrogen (Atomic mass = 14u) at 5 bar. Calculate the molar mass of the oxide [2]
10. Consider the reaction of water with F_2 and suggest, in terms of oxidation and reduction, which species are oxidized/ reduced. [2]

OR

Complete the following reactions:



11. How many neutrons and protons are there in following nuclei? [2]



12. If density of methanol is 0.793 kg/L, what is its volume needed for making 2.5 L of its 0.25 M solution? [2]

OR

The density of 3 M solution of NaCl is 1.25 g/ml. Calculate the molarity of the solution.



Section C

13. Calculate the energy associated with the first orbit of He^+ . What is the radius of this orbit? [3]

OR

An element with mass number 81 contains 3.7% more neutrons as compared to protons. Assign the symbol to the element.

14. Element X, Y and Z have 4, 5 and 7 valence electrons respectively. [3]

- (i) Write the molecular formula of the compounds formed by these elements individually with hydrogen.
- (ii) Which of these elements will have highest dipole moment?

OR

- (a) Which of the two is more stable and why? H_2^+ or H_2^-
- (b) All bonds in PCl_5 are not equal. Explain.
- (c) Which of the two is more ionic and why? NaCl or NaI

15. Give reasons: [3]

- (i) Evaporation causes cooling
- (ii) Falling liquid drops are spherical.
- (iii) Vapour pressure of acetone is less than that of ether at same temperature.

16. [3]

The combustion of one mole of methanol takes place at 298 K and 1 atm. After combustion CO_2 (g) and H_2O (l) are produced and 726 kJ of heat is liberated. Calculate the standard enthalpy of formation of one mole of CH_3OH (l). Standard enthalpies of formation of CO_2 (g) and H_2O (l) are -393 kJ mol^{-1} and -286 kJ mol^{-1} respectively.

17. How many grams of KBr be added to 1 L of 0.05 M solution of silver nitrate just to start the precipitation of AgBr ? K_{sp} of $\text{AgBr} = 5.0 \times 10^{-13}$ [3]

OR

Consider the following endothermic reaction:



- (i) Write expression for K_p for the above reaction.
- (ii) How will the equilibrium be affected by
 - Increasing the pressure
 - Using a catalyst?





18. Balance the given redox reaction in acidic medium. [3]
 $\text{MnO}_4^- + \text{SO}_2 \rightarrow \text{Mn}^{2+} + \text{HSO}_4^-$

OR


Balance $\text{P} + \text{HNO}_3 \longrightarrow \text{H}_3\text{PO}_4 + \text{NO}_2 + \text{H}_2\text{O}$ by oxidation number method.

19. [3]
(a) Name the class of hydrides to which H_2O and NaH belong.
(b) What is understood by hydride gap?
(c) What do you mean by 15 volume H_2O_2 solution?

20. Write the IUPAC names of the following: [3]
(a)



(c)




21. What are electrophiles? Explain electrophile substitution reaction with the help of example [3]
22. Comment on each of the following observations: [3]
(a) Lithium forms a nitride directly like magnesium. Give equation involved.
(b) BaO is soluble but BaSO_4 is insoluble in water.
23. [3]
(a) In which C-C bond of $\text{CH}_3\text{CH}_2\text{CH}_2\text{Br}$, the inductive effect is expected to be least?
(b) What type of isomerism is present in the following pairs?
(i) $\text{CH}_3\text{CH}_2\text{COCH}_2\text{CH}_3$ and $\text{CH}_3\text{COCH}_2\text{CH}_2\text{CH}_3$
(ii) $\text{CH}_3\text{CH}_2\text{OH}$ and CH_3OCH_3



24. Explain: [3]
- (a) What happens when:
- Quicklime is heated with silica
 - Calcium nitrate is heated
- (b) When is a cation highly polarising? Which alkali metal ion has the highest polarizing power?

Section D

25. Calculate enthalpy of change for the process: [5]



and calculate bond enthalpy of C-Cl in CCl_4

Given: $\Delta_{\text{vap}}H^\circ(\text{CCl}_4) = 30.5 \text{ kJ/mol}$

$\Delta_{\text{f}}H^\circ(\text{CCl}_4) = -135.5 \text{ kJ/mol}$

$\Delta_{\text{a}}H^\circ(\text{C}) = 715.0 \text{ kJ/mol}$

$\Delta_{\text{a}}H^\circ(\text{Cl}_2) = 242 \text{ kJ/mol}$

OR

- (a) What is bond energy? Why is it called enthalpy of atomisation?
- (b) Calculate bond energy of C-H bond, given that heat of formation of CH_4 , heat of sublimation of carbon and heat of dissociation of H_2 are -74.8 , $+719.6$, 435 kJ/mol respectively.
26. Explain why? [5]
- Boric acid is a monobasic lewis acid.
 - PbO_2 is a stronger oxidizing agent than SnO_2 .
 - CO_2 is a gas but SiO_2 is solid at room temperature.
 - SiF_6^{2-} is known but SiCl_6^{2-} is not known.
 - What is inorganic benzene and why it is so called?

OR

When metal X is treated with sodium hydroxide, a white precipitate (A) is obtained, which is soluble in excess of NaOH to give soluble complex (B). Compound (A) is soluble in dilute HCl to form compound (C). The compound (A) when heated strongly gives (D), which is used to extract metal. Identify (X), (A), (B), (C) and (D). Write suitable equations to support their identities.

27. [5]
- (a) Do the following conversions:
- Benzene to p-nitrobromobenzene
 - Ethyl chloride to ethane
- (b) Give mechanism of addition of HBr to propene.



(c) Write a note on Friedel-Crafts alkylation.

OR

(a) Out of n-hexane and ethyne which will be more acidic. Also give reason for this behaviour.

(b) Explain with example

(i) Wurtz reaction

(ii) Acidic dehydration

(c) Convert propyne to propanone

CBSE
Class XI Chemistry
Sample Paper – 3 Solution

Section A

1. $\text{HI} > \text{HBr} > \text{HCl} > \text{HF}$
2. BeCl_2 : Linear
 SiCl_4 : Tetrahedral

OR

The electron pairs involved in the bond formation are known as bond pairs or shared pairs.

3. IUPAC name of allyl alcohol: Prop-2-en-1-ol
4. The amount of oxygen required by bacteria to breakdown the organic matter present in a certain volume of a sample of water is called biochemical oxygen demand.

OR

Carboxyhaemoglobin is the compound formed when CO combines with blood.

5. Sodium (Na) – Yellow
Potassium (K) - Violet

Section B

6. Metallic character decreases and non metallic character increases in moving from left to right in a period. This is due to increase in ionization enthalpy and electron gain enthalpy.
7. Increasing order of size: $\text{Al}^{3+} < \text{Mg}^{2+} < \text{Na}^+ < \text{F}^- < \text{O}^{2-}$
This is an isoelectronic series i.e. the number electrons are the same in all the elements. Thus, as the effective nuclear charge decreases, electrons are held away from the nucleus and thus size increases.

8. Given:

Velocity of electron = 2.07×10^7 m/s

Mass of electron = 9.1×10^{-31} kg

We know,



$$\lambda = \frac{h}{mv}$$

$$= \frac{6.63 \times 10^{-34}}{9.1 \times 10^{-31} \times 2.05 \times 0.5 \times 10^7}$$

$$= 3.55 \times 10^{-11} \text{ m}$$

9. Given:

Pressure P = 5 bar

Molar mass of nitrogen M = 28 g/mol

Density of nitrogen,

$$\rho = \frac{PM}{RT}$$


Density of gaseous oxide is,

$$\frac{0.987 \times 5 \times 28}{273 \times 0.0821} = \frac{0.987 \times 5 \times (x)}{273 \times 0.0821}$$

$$x = \frac{0.987 \times 5 \times 28 \times 273 \times 0.0821}{273 \times 0.0821 \times 0.987 \times 5}$$



$$= 70$$

The molar mass of the oxide is 70 g/mol



In this reaction H_2O is getting oxidized to O_2 and F_2 is getting reduced to F^- ion. Therefore, F_2 is the oxidizing agent and H_2O is reducing agent.

OR



11. Given:

No.	Nuclei	No. of protons	No. of neutrons
1	${}^{56}_{26}\text{Fe}$	26	30
2	${}^{88}_{38}\text{Sr}$	38	50



12. Molar mass of methanol (CH_3OH) = 32 g/mol
= 0.032 kg/mol

Molarity of solution =

$$= \frac{0.793}{0.032}$$

$$= 24.78 \text{ mol/l}$$

We have,

$$M_1V_1 = M_2V_2$$

$$24.78 \times V_1 = 0.25 \times 2.5$$

$$V_1 = 25.22 \text{ ml}$$

The required volume is 25.22 ml

OR

Given:

$$M = 3 \text{ mol/lit}$$

$$\begin{aligned} \text{Mass of NaCl in 1 litre solution} &= 3 \times 58.5 \\ &= 175.5 \text{ g} \end{aligned}$$

$$\begin{aligned} \text{Mass of water in solution} &= 1000 \times 1.25 \\ &= 1250 \text{ g} \end{aligned}$$

$$\begin{aligned} \text{Mass of water in solution} &= 1250 - 175.5 \\ &= 1074.5 \text{ g} \end{aligned}$$

Molarity

$$= \frac{\text{Number of moles of solute}}{\text{Mass of solvent in kg}}$$

$$= \frac{3}{1.074}$$

$$= 2.79 \text{ m}$$



Section C

13.

We know,

$$E_n = \frac{-(2.18 \times 10^{-18})Z^2}{n^2}$$

For He^+ ,

$$n = 1, n = 2$$

$$\begin{aligned} E_1 &= \frac{-(2.18 \times 10^{-18})2^2}{1^2} \\ &= -8.72 \times 10^{-18} \text{ J} \end{aligned}$$

$$r_n = \frac{(0.0529)n^2}{Z}$$

Here, $n = 1, Z = 2$

$$\begin{aligned} r_n &= \frac{(0.0529)1^2}{2} \\ &= 0.02645 \text{ nm} \end{aligned}$$

Energy is $8.72 \times 10^{-18} \text{ J}$

Radius of the orbit is 0.02645 nm

OR

Let us find out the atomic number of element.

Let the number of protons = x

$$\text{Number of neutrons} = x + \frac{x \times 31.7}{100}$$

$$= (x + 0.317x)$$

Mass no. of element = No. of protons + No. of neutrons

$$81 = x + x + 0.317x$$

$$81 = 2.317x$$

$$x = 35$$

\therefore No. of protons = 35

No. of neutrons = $81 - 35$


$$= 46$$

Atomic number of element is 35

The element with atomic number 35 is bromine (Br).



14. The balanced chemical equation is



Volume of oxygen required to convert 2 x 22.4 L of CO at N.T.P. = 22.4 L

Volume of oxygen required to convert 5.2 L of CO at N.T.P. = $\frac{22.4}{2 \times 22.4} \times 5.2 = 2.6$ L

OR

- (i) H_2^+ is more stable than H_2^- as it contains no electron in antibonding MO while latter contains an electron in antibonding MO making it less st
- (ii) PCl_5 contains axial and equatorial bonds. Axial bonds are longer than equatorial bonds as they face more repulsion from equatorial bonds. Hence axial bonds are weaker than equatorial bonds.
- (iii) NaI is more covalent due to high polarizability of iodide ion due to its bigger size than chloride ion.


15.

- (i) This is due to the reason that the molecules which undergo evaporation are high energy molecules and therefore, the kinetic energy of the remaining molecules becomes less. Since the remaining molecules have lower average kinetic energy, their temperature becomes low.
- (ii) This is due to surface tension of liquids. Due to surface tension, the molecules of a liquid, try to make surface area to be minimum and for a given volume, sphere has the minimum surface area. Therefore the falling liquid drops are spherical.
- (iii) Intermolecular forces are stronger in acetone than in ether. Thus the vapour pressure of acetone is less than ether.



16.

(i) Combustion of methanol




(ii) Enthalpy of formation of CO_2



(iii) Enthalpy of formation of H_2O



Required reaction :




(Eq-2) + (2 x Eq-3) - (Eq-1) gives the required enthalpy

for formation of methanol

$$\begin{aligned} \Delta\text{H} &= (-572-393)+726 \\ &= -239 \text{ kJ mol}^{-1} \end{aligned}$$

17. For AgBr, $K_{sp} = 5.0 \times 10^{-13}$

Precipitation of AgBr will occur when ionic product $[\text{Ag}^+][\text{Br}^-]$ becomes larger than K_{sp} .



$$[\text{Ag}^+] = 0.05 \text{ M}$$

The concentration of Br^- required to start precipitation.

$$[\text{Br}^-] = \frac{K_{sp}}{[\text{Ag}^+]} = \frac{5.0 \times 10^{-13}}{0.05} = 1.0 \times 10^{-11}$$

$$\text{Now, } [\text{Br}^-] = [\text{KBr}] = 1.0 \times 10^{-11}$$

Molar mass of KBr = 120

$$\begin{aligned} \text{Therefore, the amount of KBr required} &= 1.0 \times 10^{-11} \times 120 \\ &= 1.20 \times 10^{-9} \text{ g} \end{aligned}$$

OR

For the reaction $\text{CH}_4(\text{g}) + \text{H}_2\text{O}(\text{g}) \rightleftharpoons \text{CO}(\text{g}) + 3\text{H}_2(\text{g})$

$$(i) K_p = \frac{(p_{\text{CO}})(p_{\text{H}_2})^3}{(p_{\text{CH}_4})(p_{\text{H}_2\text{O}})}$$


(ii) On increasing pressure, the reaction equilibrium will shift in the backward direction.

There is no effect of catalyst in equilibrium composition; however the equilibrium will be attained faster.




18. Reduction half reaction:

Step 1:





Oxidation half reaction:



Step 1:



Multiply by required coefficient and add the two equations



OR



$$\text{O} = 15 \quad \quad \quad \text{O} = 15$$

$$\text{H} = 5 \quad \quad \quad \text{H} = 5$$

Oxygen and Hydrogen atoms are balanced.

19.

- (a) H_2O is covalent hydride whereas NaH is ionic or saline hydride.
- (b) Group 7 to group 9 elements do not form hydrides. This region of periodic table from group 7 to 9 is called as hydride gap.
- (c) 1 L of H_2O_2 gives 15 L of O_2 at NTP.


20.

- (a) 2-Ethyl-3-methylpentan-1-ol
- (b) 1-Chloropropan-2-one
- (c) 2,4,6-Tribromophenol




21. A reagent which can accept an electron pair in a reaction is called an electrophile.

Examples: H^+ , Cl^+ , NO_2^+ , R_3C^+ .



Mechanism:




22.

(a) Lithium and magnesium follow diagonal relationship and so lithium like magnesium forms nitride while other alkali metals do not.

(b) Size of O^{2-} ion is smaller than SO_4^{2-} . Since a bigger anions stabilizes bigger cation more than a smaller cation stabilizes a bigger anion, lattice enthalpy of BaO is smaller than $BaSO_4$. BaO is soluble as hydration energy is more than lattice energy but $BaSO_4$ (as hydration energy is less than lattice energy) is insoluble in water.

23.

(a) The inductive effect is least in C_2-C_3 bond because the magnitude of inductive effect decreases as the number of intervening bonds increases.



(b)

(i) Metamerism


(ii) Functional group isomerism



24.

(a)

(i) When quicklime is heated with silica it gives calcium silicate.



(ii) When calcium nitrate is heated it forms CaO, NO₂ and O₂



(b) A cation is highly polarizing if its charge/radius ratio is high.

Li⁺ ion has the highest polarizing power among the alkali metal ions because it has the highest charge/radius ratio.

Section D

25.

Given:



By solving,

Equation(3) + 2(equation4) – equation (1) – equation(2)

Substituting the values,

$$\begin{aligned} \Delta H &= [715.0 + 2(242) - 30.5 - (-135.5)] \text{ kJ/mol} \\ &= 1304 \text{ kJ/mol} \end{aligned}$$

Bond enthalpy of C-Cl in CCl₄ (average value)

$$= \frac{1304}{4}$$


$$= 326 \text{ kJmol}^{-1}$$



Enthalpy of change for the process is 1304 kJ/mol

Bond enthalpy is 326 kJ/mol

OR

(a) Bond energy is the amount of energy required to dissociate one mole of bonds present between the atoms in the gaseous phase. As molecules dissociate completely into atoms in the gaseous phase therefore bond energy of a diatomic molecule is called enthalpy of atomization.





This gives the enthalpy of dissociation of four moles C-H bonds.

Hence bond energy for C-H bond

$$= \frac{1665.2}{4}$$

$$= 416.3 \text{ kJ/mol}$$





26.

- (a) This is because boric acid does not act as proton donor rather it accepts a lone pair of electrons from OH⁻ ions. There by acting as monobasic lewis acid.
- (b) PbO₂ and SnO₂ both are in +4 oxidation state. But due to stronger inert pair effect Pb²⁺ ion is more stable than Sn²⁺ ion.
- (c) In other way Pb²⁺ ions is more easily reduced to Pb⁰ ions. Thus PbO₂ acts as a stronger oxidizing agent than SnO₂.
- (d) The molecules of CO₂ are held together by weak Van der Waals forces of attraction which can be easily overcome by collisions of the molecules at room temperature. Consequently CO₂ is a gas.
While silicon atoms forms four single covalent bonds with O-atm which are tetrahedrally arranged and form a three- dimensional structure. Thus SiO₂ is a high melting solid.
- (e) SiF₆²⁻ is known but SiCl₆²⁻ is not known, because interaction between lone pair of chloride ion and Si⁴⁺ ion is not strong. Also six large chloride ions cannot be accommodated around Si⁴⁺ due to limitation of its size.
- (f) Borazine is called inorganic benzene, as its structure is similar to that of benzene. Its formula is B₃N₃H₆. The compound is isoelectric and isostructural with benzene.

OR

Metal X on treatment with sodium hydroxide gives white precipitate which dissolves in excess of NaOH to give soluble complex (B), therefore, the metal X is Al.








27.

(a)


(i) Benzene to p-Nitrobromobenzene




(ii) Ethyl chloride to ethene



(b) Mechanism of addition of HBr to propene




(c) Friedel-Crafts alkylation- It is the reaction of benzene with alkyl halide in presence of anhydrous aluminium chloride. The reaction results in the formation of alkyl benzene.




OR

(a)



n-hexane
 sp^3 hybridised
carbon
s-character 25%




Ethyne
 sp hybridized
carbon
s-character 50%

Since s-orbitals are closer to the nucleus, hence due to more s character in ethyne (sp hybridized) the hybridized orbital is nearest to this carbon atom in comparison to sp^2 hybridised carbon. This leads to the movement of C-H bond pair more towards sp hybridized carbon, leading to the development of partial positive charge on the hydrogen attached to sp hybridised carbon and eventually helps in release of proton (H^+). Thus, ethyne is more acidic than n-hexane.




(b)

(i) Wurtz reaction: Alkyl halides on treatment with sodium metal in dry ether medium give higher alkanes. This is called Wurtz reaction and is used for the preparation of alkanes with even number of carbon atoms.



(ii) Acidic dehydration: Alcohols on heating with conc. H_2SO_4 at 443 K form alkenes with elimination of one water molecule. Since a water molecule is lost in the presence of acid, the reaction is called acidic dehydration of alcohols.



(c)

