

**Class XI Session 2023-24**  
**Subject - Mathematics**  
**Sample Question Paper - 7**

**Time Allowed: 3 hours**

**Maximum Marks: 80**

**General Instructions:**

1. This Question paper contains - five sections A, B, C, D and E. Each section is compulsory. However, there are internal choices in some questions.
2. Section A has 18 MCQ's and 02 Assertion-Reason based questions of 1 mark each.
3. Section B has 5 Very Short Answer (VSA)-type questions of 2 marks each.
4. Section C has 6 Short Answer (SA)-type questions of 3 marks each.
5. Section D has 4 Long Answer (LA)-type questions of 5 marks each.
6. Section E has 3 source based/case based/passage based/integrated units of assessment (4 marks each) with sub parts.

**Section A**

1.  $\frac{\cos 8^\circ - \sin 8^\circ}{\cos 8^\circ + \sin 8^\circ} = ?$  [1]  
a)  $\tan 52^\circ$  b)  $\tan 37^\circ$   
c) None of these d)  $\tan 8^\circ$
2. The domain and range of real function  $f$  defined by  $f(x) = \sqrt{x-1}$  is given by [1]  
a) Domain =  $[\infty, \infty)$ , Range =  $[0, \infty)$  b) Domain =  $[1, \infty)$ , Range =  $(\infty, \infty)$   
c) Domain =  $[1, \infty)$ , Range =  $[0, \infty)$  d) Domain =  $(1, \infty)$ , Range =  $(0, \infty)$
3. The mean and the variance of 10 observations are given to be 4 and 2 respectively. If every observation is multiplied by 2, the mean and the variance of the new series will be respectively. [1]  
a) 8 and 4 b) 8 and 20  
c) 8 and 8 d) 80 and 40
4. If  $G(x) = \sqrt{25 - x^2}$  then  $\lim_{x \rightarrow 1} \frac{G(x) - G(1)}{x - 1}$  has the value [1]  
a)  $\frac{1}{24}$  b)  $-\sqrt{24}$   
c)  $\frac{-1}{\sqrt{24}}$  d)  $\frac{1}{5}$
5. The equations of the sides AB, BC and CA of  $\triangle ABC$  are  $y - x = 2$ ,  $x + 2y = 1$  and  $3x + y + 5 = 0$  respectively. [1]  
The equation of the altitude through B is  
a)  $3x - y + 2 = 0$  b)  $x - 3y + 4 = 0$   
c)  $x - 3y + 1 = 0$  d) None of these
6. Equation of y-axis is considered as [1]



- a)  $y = 0, z = 0$   
c)  $z = 0, x = 0$

b) none of these  
d)  $x = 0, y = 0$

7. If  $z = x + iy; x, y \in \mathbb{R}$  then : [1]  
a)  $z\bar{z} < |z|^2$   
c)  $z\bar{z} > |z|^2$   
b)  $z\bar{z} = |z|^2$   
d) none of these

8.  ${}^5C_1 + {}^5C_2 + {}^5C_3 + {}^5C_4 + {}^5C_5$  is equal to [1]  
a) 33  
c) 31  
b) 30  
d) 32

9.  $\lim_{x \rightarrow 0} \frac{\tan 2x - x}{3x - \sin x}$  is equal to [1]  
a)  $\frac{1}{2}$   
c)  $\frac{1}{4}$   
b) 2  
d)  $-\frac{1}{2}$

10. If the arcs of the same length in two circles subtend angles of  $60^\circ$  and  $75^\circ$  at their respective centres, the ratio of their radii is [1]  
a) 5 : 3  
c) 5 : 4  
b) 3 : 5  
d) 4 : 5

11. For any two sets A and B,  $A \cap (A \cup B) = \dots$  [1]  
a) none of these  
c)  $\phi$   
b) B  
d) A

12. The integral part of  $(\sqrt{2} + 1)^6$  is [1]  
a) 98  
c) 99  
b) 96  
d) 100

13. If  $x = 99^{50} + 100^{50}$  and  $y = (101)^{50}$  then [1]  
a)  $x < y$   
c)  $x = y$   
b)  $x > y$   
d)  $x \geq y$

14. If x and a are real numbers such that  $a > 0$  and  $|x| > a$ , then [1]  
a)  $x \in (-a, \infty)$   
c)  $x \in (-a, a)$   
b)  $x \in (-\infty, -a) \cup (a, \infty)$   
d)  $x \in [-\infty, a]$

15. Let R be set of points inside a rectangle of sides a and b ( $a, b > 1$ ) with two sides along the positive direction of x-axis and y-axis. Then [1]  
a)  $R = \{(x, y) : 0 \leq x \leq a, 0 \leq y \leq b\}$   
c)  $R = \{(x, y) : 0 < x < a, 0 < y < b\}$   
b)  $R = \{(x, y) : 0 \leq x < a, 0 \leq y \leq b\}$   
d)  $R = \{(x, y) : 0 \leq x \leq a, 0 < y < b\}$

16. If  $A + B + C = \pi$ , then  $\frac{\tan A + \tan B + \tan C}{\tan A \tan B \tan C}$  is equal to [1]  
a) 1  
c) None of these  
b)  $\tan A \tan B \tan C$   
d) 0

17.  $\lim_{x \rightarrow \frac{\pi}{4}} \frac{\tan x - 1}{x - \frac{\pi}{4}}$  is equal to
- a) 1                                  b)  $\frac{1}{2}$
- c) 0                                  d) 2
18. Find r if  $^{10}P_r = 2.^9P_r$  [1]
- a) 6                                  b) 4
- c) 3                                  d) 5
19. **Assertion (A):** If A = set of letters in Alloy B = set of letters in LOYAL, then set A & B are equal sets. [1]  
**Reason (R):** If two sets have exactly the same elements, they are called equal sets.
- a) Both A and R are true and R is the correct explanation of A.                      b) Both A and R are true but R is not the correct explanation of A.
- c) A is true but R is false.                      d) A is false but R is true.
20. **Assertion (A):** The sum of infinite terms of a geometric progression is given by  $S_\infty = \frac{a}{1-r}$ , provided  $|r| < 1$ . [1]  
**Reason (R):** The sum of n terms of Geometric progression is  $S_n = \frac{a(r^n-1)}{r-1}$ .
- a) Both A and R are true and R is the correct explanation of A.                      b) Both A and R are true but R is not the correct explanation of A.
- c) A is true but R is false.                      d) A is false but R is true.

## Section B

21. Let A and B be two non-empty sets such that  $n(A) = 5$ ,  $n(B) = 6$  and  $n(A \cap B) = 3$ . Find [2]
- i.  $n(A \times B)$
  - ii.  $n(B \times A)$
  - iii.  $n\{(A \times B) \cap (B \times A)\}$

OR

Find the values of a and b, if

- i.  $(2a - 5, 4) = (5, b + 6)$
- ii.  $(a - 3, b + 7) = (3, 7)$
22. Evaluate:  $\lim_{x \rightarrow 0} \frac{\sin^2 4x^2}{x^4}$ . [2]
23. If the odds against the occurrence of an event be 4 : 7, find the probability of the occurrence of the event. [2]

OR

A bag contains 8 red, 3 white and 9 blue balls. If three balls are drawn at random, determine the probability that all the three balls are blue balls.

24. For any two sets A and B prove by using properties of sets that:  $(A \cap B) \cup (A - B) = A$ . [2]
25. Find the equation of a line that has y-intercept 4 and is perpendicular to the line joining (2, -3) and (4, 2). [2]

## Section C

26. In how many ways can six persons be seated in a row? **[3]**
27. Verify that  $(-1, 2, 1)$ ,  $(1, -2, 5)$ ,  $(4, -7, 8)$  and  $(2, -3, 4)$  are the vertices of a parallelogram. **[3]**
28. Using binomial theorem, expand:  $\left(\frac{2x}{3} - \frac{3}{2x}\right)^6$  **[3]**

OR

Evaluate:  $(\sqrt{3} + 1)^5 - (\sqrt{3} - 1)^5$

29. Differentiate  $(x^2 + 1)(x - 5)$  from first principle. [3]

OR

Find the derivative of the following functions from first principle.  $\frac{x+1}{x-1}$

30. If A.M. and G.M. of roots of a quadratic equation are 8 and 5 respectively then obtain the quadratic equation. [3]

OR

If the sum of an infinite decreasing G.P. is 3 and the sum of the squares of its terms is  $\frac{9}{2}$ , then write its first term and common difference.

31. Out of 25 members in a family, 12 like to take tea, 15 like to take coffee and 7 like to take coffee and tea both. [3]

How many like

i. at least one of the two drinks

ii. only tea but not coffee

iii. only coffee but not tea

iv. neither tea nor coffee

#### Section D

32. Calculate the mean deviation about the mean of the set of first  $n$  natural numbers when  $n$  is an even number. [5]

33. Find the vertex, axis, focus, directrix, latus - rectum of the following parabolas. Also, draw their rough sketches: [5]

$$y = x^2 - 2x + 3.$$

OR

Find the eccentricity, centre, vertices, foci, minor axis, major axis, directrices and latus-rectum of the ellipse  $25x^2 + 9y^2 - 150x - 90y + 225 = 0$ .

34. Solve the following system of linear inequalities [5]

$$-2 - \frac{x}{4} \geq \frac{1+x}{3} \text{ and } 3 - x < 4(x-3)$$

35. If  $2 \tan \alpha = 3 \tan \beta$ , prove that  $\tan(\alpha - \beta) = \frac{\sin 2\beta}{5 - \cos 2\beta}$ . [5]

OR

If  $\sec(x + \alpha) + \sec(x - \alpha) = 2 \sec x$ , prove that  $\cos x = \pm \sqrt{2} \cos \frac{\alpha}{2}$ .

#### Section E

36. Read the text carefully and answer the questions: [4]

##### Method to Find the Sets When Cartesian Product is Given

For finding these two sets, we write first element of each ordered pair in first set say  $A$  and corresponding second element in second set  $B$  (say).

##### Number of Elements in Cartesian Product of Two Sets

If there are  $p$  elements in set  $A$  and  $q$  elements in set  $B$ , then there will be  $pq$  elements in  $A \times B$  i.e. if  $n(A) = p$  and  $n(B) = q$ , then  $n(A \times B) = pq$ .

(i) The Cartesian product  $A \times A$  has 9 elements among which are found  $(-1, 0)$  and  $(0, 1)$ . Find the set  $A$  and the remaining elements of  $A \times A$ .

(ii)  $A$  and  $B$  are two sets given in such a way that  $A \times B$  contains 6 elements. If three elements of  $A \times B$  are  $(1, 3)$ ,  $(2, 5)$  and  $(3, 3)$ , then find the remaining elements of  $A \times B$ .

(iii) If the set  $A$  has 3 elements and set  $B$  has 4 elements, then find the number of elements in  $A \times B$ .

OR

If  $A \times B = \{(a, 1), (b, 3), (a, 3), (b, 1), (a, 2), (b, 2)\}$ . Find  $A$  and  $B$ .

37. Read the text carefully and answer the questions: [4]



Four friends Dinesh, Yuvraj, Sonu, and Rajeev are playing cards. Dinesh, shuffling a cards and told to Rajeev choose any four cards.



- (i) What is the probability that Rajeev getting all face card.
- (ii) What is the probability that Rajeev getting two red cards and two black card.
- (iii) What is the probability that Rajeev getting one card from each suit.

**OR**

What is the probability that Rajeev getting two king and two Jack cards.

38. **Read the text carefully and answer the questions:**

[4]

We have,  $i = \sqrt{-1}$ . So, we can write the higher powers of  $i$  as follows

i.  $i^2 = -1$

ii.  $i^3 = i^2 \cdot i = (-1) \cdot i = -i$

iii.  $i^4 = (i^2)^2 = (-1)^2 = 1$

iv.  $i^5 = i^{4+1} = i^4 \cdot i = 1 \cdot i = i$

v.  $i^6 = i^{4+2} = i^4 \cdot i^2 = 1 \cdot i^2 = -1...$

In order to compute  $i^n$  for  $n > 4$ , write

$$i^n = i^{4q+r} \text{ for some } q, r \in \mathbb{N} \text{ and } 0 \leq r \leq 3. \text{ Then, } i^n = i^{4q} \cdot i^r$$

$$= (i^4)^q \cdot i^r = (1)^q \cdot i^r = i^r$$

In general for any integer  $k$

$$i^{4k} = 1, i^{4k+1} = i, i^{4k+2} = -1 \text{ and } i^{4k+3} = -i$$

- (i) Find the value of  $i^{30}$ .
- (ii) If  $z = i^{-39}$ , then find the simplest form of  $z$ .

## Solution

### Section A

1.  
(b)  $\tan 37^\circ$   
**Explanation:**  $\frac{\cos 8^\circ - \sin 8^\circ}{\cos 8^\circ + \sin 8^\circ} = \frac{1 - \tan 8^\circ}{1 + \tan 8^\circ} = \frac{\tan 45^\circ - \tan 8^\circ}{1 + \tan 45^\circ \tan 8^\circ} [\because 1 = \tan 45^\circ]$   
 $= \tan (45^\circ - 8^\circ) = \tan 37^\circ$
2.  
(c) Domain =  $[1, \infty)$ , Range =  $[0, \infty)$   
**Explanation:** We have,  $f(x) = \sqrt{x-1}$   
Clearly,  $f(x)$  is defined if  $x-1 \geq 0$   
 $\Rightarrow x \geq 1$   
 $\therefore$  Domain of  $f = [1, \infty)$   
Now for  $x \geq 1, x-1 \geq 0$   
 $\Rightarrow \sqrt{x-1} \geq 0$   
 $\Rightarrow$  Range of  $f = [0, \infty)$
3.  
(c) 8 and 8  
**Explanation:** Let the observations be  $x'_i$  s,  $i = 1, 2, \dots, 10$  and the mean and variance of  $y'_i$  s are  $\bar{x} = 4$  and  $\sigma^2 = 2$ .  
Now, let  $y_i = 2x'_i$  s and the mean and variance of  $y'_i$  s and  $\bar{y}$  and  $\sigma_1^2$  then  
 $\bar{y} = \frac{\sum 2x_i}{10} = 2 \frac{\sum x_i}{10} = 2\bar{x} = 8$  and  $\sigma_1^2 = \text{var}(y'_i \text{ s}) = \text{var}(2x'_i \text{ s})$   
 $= 4 \text{ var}(x'_i \text{ s}) = 4 \times 2 = 8$   
Thus, the mean and variance of new series are 8 and 8.
4.  
(c)  $\frac{-1}{\sqrt{24}}$   
**Explanation:** The equation is in the form of  $\frac{0}{0}$   
 $\frac{1}{2\sqrt{25-x^2}} \cdot (-2x)$   
Using L' Hospital rule we have  $\frac{1}{1}$   
Substituting  $x = 1$  we get  $\frac{-1}{\sqrt{24}}$
5.  
(b)  $x - 3y + 4 = 0$   
**Explanation:** The equation of the sides AB, AC and CA of  $\triangle ABC$  are  $y - x = 2$ ,  $x + 2y = 1$  and  $3x + y + 5 = 0$ , respectively.  
Solving the equations of AB and BC, i.e,  $y - x = 2$  and  $x + 2y = 1$ , we get  
 $x = -1, y = 1$   
So, the coordinates of B are  $(-1, 1)$   
 $\therefore$  Slope of AC = -3  
Thus, slope of the altitude through B is  $\frac{1}{3}$ .  
Equation of the required altitude is given below as per the general formula :  
 $y - 1 = \frac{1}{3}(x + 1)$   
 $\Rightarrow x - 3y + 4 = 0$ .
6.  
(c)  $z = 0, x = 0$   
**Explanation:** On y-axis consider as  $x = 0$  and  $z = 0$
7.  
(b)  $z\bar{z} = |z|^2$   
**Explanation:** If  $z = x + iy$  then  $\bar{z} = x - iy$   
Now  $z\bar{z} = (x + iy) \cdot (x - iy) = x^2 + y^2 = |z|^2$  [ $\because |z| = \sqrt{x^2 + y^2}$ ]

8.

(c) 31

**Explanation:**  ${}^5C_1 + {}^5C_2 + {}^5C_3 + {}^5C_4 + {}^5C_5$   
 $= {}^5C_1 + {}^5C_2 + {}^5C_2 + {}^5C_1 + {}^5C_5$   
 $= 2 \times {}^5C_1 + 2 \times {}^5C_2 + {}^5C_5$   
 $= 2 \times 5 + 2 \times \frac{5!}{2!3!} + 1$   
 $= 10 + 20 + 1$   
 $= 31.$

9. (a)  $\frac{1}{2}$

**Explanation:** Given,  $\lim_{x \rightarrow 0} \frac{\tan 2x - x}{3x - \sin x} = \lim_{x \rightarrow 0} \frac{x \left[ \frac{\tan 2x}{x} - 1 \right]}{x \left[ 3 - \frac{\sin x}{x} \right]}$   
 $\lim_{x \rightarrow 0} \frac{\frac{\tan 2x}{2x} \times 2 - 1}{3 - \frac{\sin x}{x}} = \frac{1 \cdot 2 - 1}{3 - 1} = \frac{2 - 1}{2} = \frac{1}{2}$

10.

(c) 5 : 4

**Explanation:**  $\theta_1 = 60^\circ = \left( 60 \times \frac{\pi}{180} \right)^c = \left( \frac{\pi}{3} \right)^c$  and  $\theta_2 = 75^\circ = \left( 75 \times \frac{\pi}{180} \right)^c = \left( \frac{5\pi}{12} \right)^c$   
 $\therefore l = r_1 \theta_1 = r_2 \theta_2$   
 $\Rightarrow r_1 \times \frac{\pi}{3} = r_2 \times \frac{5\pi}{12} \Rightarrow \frac{r_1}{r_2} = \left( \frac{5}{12} \times 3 \right) = \frac{5}{4} \Rightarrow r_1 : r_2 = 5 : 4$

11.

(d) A

**Explanation:** Common between set A and  $(A \cup B)$  is set A itself

12.

(c) 99

**Explanation:** We have  $(1 + x)^n = 1 + {}^nC_1(x) + {}^nC_2(x)^2 + \dots + (x)^n$   
Hence  $(\sqrt{2} + 1)^6 = 1 + {}^6C_1(\sqrt{2}) + {}^6C_2(\sqrt{2})^2 + {}^6C_3(\sqrt{2})^3 + {}^6C_4(\sqrt{2})^4 + {}^6C_5(\sqrt{2})^5 + (\sqrt{2})^6$   
 $\Rightarrow (\sqrt{2} + 1)^6 = 1 + 6(\sqrt{2}) + 15 \times 2 + 20 \times 2(\sqrt{2}) + 15 \times 4 + 6 \times 4(\sqrt{2}) + 8$   
 $= 99 + 70\sqrt{2}$   
Hence integral part of  $(\sqrt{2} + 1)^6 = 99$

13. (a)  $x < y$

**Explanation:** Given  $x = 99^{50} + 100^{50}$  and  $y = (101)^{50}$   
Now  $y = (101)^{50} = (100 + 1)^{50} = {}^{50}C_0(100)^{50} + {}^{50}C_1(100)^{49} + {}^{50}C_2(100)^{48} + \dots + {}^{50}C_{50} \dots (i)$   
Also  $(99)^{50} = (100 - 1)^{50} = {}^{50}C_0(100)^{50} - {}^{50}C_1(100)^{49} + {}^{50}C_2(100)^{48} - \dots + {}^{50}C_{50} \dots (ii)$   
Now subtract equation (ii) from equation (i), we get  
 $(101)^{50} - (99)^{50} = 2 \left[ {}^{50}C_1 (100)^{49} + {}^{50}C_3 (100)^{47} + \dots \right]$   
 $= 2 \left[ 50(100)^{49} + \frac{50 \times 49 \times 48}{3 \times 2 \times 1} (100)^{47} + \dots \right]$   
 $= (100)^{50} + 2 \left( \frac{50 \times 49 \times 48}{3 \times 2 \times 1} (100)^{47} \right)$   
 $\Rightarrow (101)^{50} - (99)^{50} > (100)^{50}$   
 $\Rightarrow (101)^{50} > (100)^{50} + (99)^{50}$   
 $\Rightarrow y > x$

14.

(b)  $x \in (-\infty, -a) \cup (a, \infty)$

**Explanation:**  $|x| > a$   
 $\Rightarrow x < -a$  or  $x > a$   
 $\Rightarrow x \in (-\infty, -a) \cup (a, \infty)$

15.

(c)  $R = \{(x, y) : 0 < x < a, 0 < y < b\}$

**Explanation:** We have, R be set of points inside a rectangle of sides a and b  
Since,  $a, b > 1$

a and b cannot be equal to 0

Thus,  $R = \{(x, y) : 0 < x < a, 0 < y < b\}$

16. (a) 1

**Explanation:**  $\pi = 180^\circ$

Using  $\tan(180 - A) = -\tan A$ , we get;

$$C = \pi - (A + B)$$

Now,

$$\begin{aligned} & \frac{\tan A + \tan B + \tan C}{\tan A \tan B \tan C} \\ &= \frac{\tan A + \tan B + \tan[\pi - (A + B)]}{\tan A \tan B \tan[\pi - (A + B)]} \\ &= \frac{\tan A + \tan B - \tan(A + B)}{-\tan A \tan B \tan(A + B)} \\ &= \frac{\tan A + \tan B - \frac{\tan A + \tan B}{1 - \tan A \tan B}}{-\tan A \tan B \times \frac{\tan A + \tan B}{1 - \tan A \tan B}} \\ &= \frac{\tan A + \tan B - \tan^2 A \tan B - \tan A \tan^2 B - \tan A - \tan B}{-\tan^2 A \tan B - \tan A \tan^2 B} \\ &= \frac{-\tan^2 A \tan B - \tan A \tan^2 B}{-\tan^2 A \tan B - \tan A \tan^2 B} \\ &= 1 \end{aligned}$$

17.

(d) 2

**Explanation:** Let  $x - \frac{\pi}{4} = t$

$$\begin{aligned} & \Rightarrow \lim_{t \rightarrow 0} \frac{\tan\left(\frac{\pi}{4} + t\right) - 1}{t} \\ & \Rightarrow \lim_{t \rightarrow 0} \frac{2 \tan t}{(1 - \tan t)(t)} \\ & = 2 \end{aligned}$$

18.

(d) 5

**Explanation:** Given  ${}^{10}P_r = 2 \cdot {}^9P_r$

$$\begin{aligned} & \Rightarrow \frac{10!}{(10-r)!} = 2 \cdot \frac{(9)!}{(9-r)!} \\ & \Rightarrow \frac{10 \times 9!}{(10-r) \times (9-r)!} = 2 \cdot \frac{(9)!}{(9-r)!} \\ & \Rightarrow \frac{10}{(10-r)} = 2 \\ & \Rightarrow 10 = 20 - 2r \\ & \Rightarrow 2r = 10 \\ & \Rightarrow r = 5 \end{aligned}$$

19. (a) Both A and R are true and R is the correct explanation of A.

**Explanation:** Both A and R are true and R is the correct explanation of A.

20.

(b) Both A and R are true but R is not the correct explanation of A.

**Explanation:** Both A and R are true but R is not the correct explanation of A.

### Section B

21. Here we are given that, A and B are two non-empty sets such that  $n(A) = 5$ ,  $n(B) = 6$  and  $n(A \cap B) = 3$

i.  $n(A \times B) = n(A) \times n(B) = (5 \times 6) = 30$

ii.  $n(B \times A) = n(B) \times n(A) = (6 \times 5) = 30$

iii. Given:  $n(A \cap B) = 3$

$\therefore$  A and B have 3 elements in common

So,  $(A \times B)$  and  $(B \times A)$  have  $3^2 = 9$  elements in common.

Hence,  $n\{(A \times B) \cap (B \times A)\} = 9$

OR

We know that two ordered pairs are equal if their corresponding elements are equal.

i.  $(2a - 5, 4) = (5, b + 6) \Rightarrow 2a - 5 = 5$  and  $4 = b + 6$  [equating corresponding elements]  
 $\Rightarrow 2a = 5 + 5$  and  $4 - 6 = b$



$$\Rightarrow 2a = 10 \text{ and } -2 = b \Rightarrow a = 5 \text{ and } b = -2$$

$$\text{ii. } (a - 3, b + 7) = (3, 7) \Rightarrow a - 3 = 3 \text{ and } b + 7 = 7 \text{ [equating corresponding elements]}$$

$$\Rightarrow a = 3 + 3 \text{ and } b = 7 - 7 \Rightarrow a = 6 \text{ and } b = 0$$

$$\begin{aligned} 22. \text{ We have: } & \lim_{x \rightarrow 0} \left[ \frac{\sin^2 4x^2}{x^4} \right] \\ &= \lim_{x \rightarrow 0} \left[ \frac{\sin(4x^2)}{x^2} \times \frac{\sin(4x^2)}{x^2} \right] \\ &= \lim_{x \rightarrow 0} \left[ \frac{\sin(4x^2)}{4x^2} \times 4 \times \frac{\sin(4x^2)}{4x^2} \times 4 \right] \\ &= 4 \times 4 \left[ \cdot \cdot \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \right] \\ &= 16 \end{aligned}$$

23. We know that,

If odds in favour of the occurrence an event are a: b, then the probability of an event to occur is  $\frac{a}{a+b}$ ,

similarly, if odds are not in the favor of the occurrence an event are a: b, then the probability of not occurrence of the event is  $\frac{a}{a+b}$

that is the probability of not occurring =  $\frac{a}{a+b}$

We also know that,

Probability of occurring = 1 - the probability of not occurring

$$\begin{aligned} &= 1 - \frac{a}{a+b} \\ &= \frac{b}{a+b} \end{aligned}$$

Given a = 4 and b = 7

$$\begin{aligned} \text{Probability of occurrence} &= \frac{7}{4+7} \\ &= \frac{7}{11} \end{aligned}$$

OR

We have to find the probability that all the three balls are blue balls

**Given:** bag which contains 8 red, 3 white, 9 blue balls

**Formula:**  $P(E) = \frac{\text{favourable outcomes}}{\text{total possible outcomes}}$

three balls are drawn at random therefore

Total possible outcomes of selecting two persons is  ${}^{20}C_3$

$$\text{Therefore } n(S) = {}^{20}C_3 = 1140$$

let E be the event that all the balls are blue

$$E = \{B, B, B\}$$

$$n(E) = {}^9C_3 = 84$$

$$\begin{aligned} P(E) &= \frac{n(E)}{n(S)} \\ P(E) &= \frac{84}{1140} = \frac{7}{95} \end{aligned}$$

24. We can write,  $(A \cap B) \cup (A - B) = (A \cap B) \cup (A \cap B')$

$$= X \cup (A \cap B'), \text{ where } X = A \cap B$$

$$= (X \cup A) \cap (X \cup B')$$

$$= A \cap (A \cup B') [\because X \cup A = (A \cap B) \cup A = A] [\because A \cap B \subset A]$$

$$= X \cup B' = (A \cap B) \cup B'$$

$$\Rightarrow (A \cup B') \cap (B \cup B')$$

$$\Rightarrow (A \cup B') \cap U = A \cup B'$$

$$= A [\because A \subset A \cup B']$$

25. Let m be the slope of the required line.

Since the required line is perpendicular to the line joining A (2, -3) and B (4, 2).

$$\therefore m \times \text{Slope of AB} = -1 \Rightarrow m \times \frac{2+3}{4-2} = -1 \Rightarrow m = -\frac{2}{5}$$

The required line cuts off an intercept of length 4 on y-axis. So, c = 4

Substituting these values in  $y = mx + c$ , we obtain that the equation of the required line is

$$y = -\frac{2}{5}x + 4$$

$$\text{or, } 2x + 5y - 20 = 0$$

which is the required equation of line.

Section C



26. Given: Six persons are to be arranged in a row.

Assume six seats, now in the first seat, any one of six members can be seated, so the total number of possibilities is  ${}^6C_1$

Similarly, in the second seat, any one of five members can be seated, so the total number of possibilities is  ${}^5C_1$

In the third seat, any one of four members can be seated, so the total number of possibilities is  ${}^4C_1$


In the fourth seat, any one of three members can be seated, so the total number of possibilities is  ${}^3C_1$

In the fifth seat, any one of two members can be seated, so the total number of possibilities is  ${}^2C_1$

In the sixth seat, only one remaining person can be seated, so the total number of possibilities is  ${}^1C_1$

Hence the total number of possible outcomes =  ${}^6C_1 \times {}^5C_1 \times {}^4C_1 \times {}^3C_1 \times {}^2C_1 \times {}^1C_1 = 6 \times 5 \times 4 \times 3 \times 2 \times 1 = 720$

27. Let A (-1, 2, 1), B(1, -2, 5), C(4, -7, 8) and D (2, -3, 4) are the vertices of a quadrilateral ABCD.



Then, mid-point of

$$AC = \left( \frac{-1+4}{2}, \frac{2-7}{2}, \frac{1+8}{2} \right) = \left( \frac{3}{2}, \frac{-5}{2}, \frac{9}{2} \right) \left[ \because \text{coordinates of mid-point } \left( \frac{x_1+x_2}{2}, \frac{y_1+y_2}{2}, \frac{z_1+z_2}{2} \right) \right]$$

$$\text{Similarly, mid-point of BD} = \left( \frac{3}{2}, -\frac{5}{2}, \frac{9}{2} \right)$$

Mid-points of both the diagonals are the same (i.e., they bisect each other).

Hence, ABCD is a parallelogram.

28. To find: Expansion of  $\left( \frac{2x}{3} - \frac{3}{2x} \right)^6$  by means of binomial theorem

$$\text{Formula used: } {}^nC_r = \frac{n!}{(n-r)!(r)!}$$

$$(a+b)^n = {}^nC_0 a^n + {}^nC_1 a^{n-1}b + {}^nC_2 a^{n-2}b^2 + \dots + {}^nC_{n-1} a b^{n-1} + {}^nC_n b^n$$

$$\text{Now here We have, } \left( \frac{2x}{3} - \frac{3}{2x} \right)^6$$

$$\begin{aligned} &= \left[ 6C_0 \left( \frac{2x}{3} \right)^{6-0} \right] + \left[ 6C_1 \left( \frac{2x}{3} \right)^{6-1} \left( -\frac{3}{2x} \right)^1 \right] + \left[ 6C_2 \left( \frac{2x}{3} \right)^{6-2} \left( -\frac{3}{2x} \right)^2 \right] \\ &+ \left[ 6C_3 \left( \frac{2x}{3} \right)^{6-3} \left( -\frac{3}{2x} \right)^3 \right] + \left[ 6C_4 \left( \frac{2x}{3} \right)^{6-4} \left( -\frac{3}{2x} \right)^4 \right] \\ &+ \left[ 6C_5 \left( \frac{2x}{3} \right)^{6-5} \left( -\frac{3}{2x} \right)^5 \right] + \left[ 6C_6 \left( -\frac{3}{2x} \right)^6 \right] \\ &= \left[ \frac{6!}{0!(6-0)!} \left( \frac{2x}{3} \right)^6 \right] - \left[ \frac{6!}{1!(6-1)!} \left( \frac{2x}{3} \right)^5 \left( \frac{3}{2x} \right) \right] + \left[ \frac{6!}{2!(6-2)!} \left( \frac{2x}{3} \right)^4 \left( \frac{9}{4x^2} \right) \right] - \left[ \frac{6!}{3!(6-3)!} \left( \frac{2x}{3} \right)^3 \left( \frac{27}{8x^3} \right) \right] \\ &+ \left[ \frac{6!}{4!(6-4)!} \left( \frac{2x}{3} \right)^2 \left( \frac{81}{16x^4} \right) \right] - \left[ \frac{6!}{5!(6-5)!} \left( \frac{2x}{3} \right)^1 \left( \frac{243}{32x^5} \right) \right] + \left[ \frac{6!}{6!(6-6)!} \left( \frac{729}{64x^6} \right) \right] \\ &= \left[ 1 \left( \frac{64x^6}{729} \right) \right] - \left[ 6 \left( \frac{32x^5}{243} \right) \left( \frac{3}{2x} \right) \right] + \left[ 15 \left( \frac{16x^4}{81} \right) \left( \frac{9}{4x^2} \right) \right] - \left[ 20 \left( \frac{8x^3}{27} \right) \right] \\ &+ \left[ 15 \left( \frac{4x^2}{9} \right) \left( \frac{81}{16x^4} \right) \right] - \left[ 6 \left( \frac{2x}{3} \right) \left( \frac{243}{32x^5} \right) \right] + \left[ 1 \left( \frac{729}{64x^6} \right) \right] \\ &= \frac{64}{729} x^6 - \frac{32}{27} x^4 + \frac{20}{3} x^2 - 20 + \frac{135}{4} \frac{1}{x^2} - \frac{243}{8} \frac{1}{x^4} + \frac{729}{64} \frac{1}{x^6} \end{aligned}$$

OR

$$\text{To find: Value of } (\sqrt{3} + 1)^5 - (\sqrt{3} - 1)^5$$

$$\text{Formula used: } {}^nC_r = \frac{n!}{(n-r)!(r)!}$$

$$(a+b)^n = {}^nC_0 a^n + {}^nC_1 a^{n-1}b + {}^nC_2 a^{n-2}b^2 + \dots + {}^nC_{n-1} a b^{n-1} + {}^nC_n b^n$$

$$(a+1)^5 = {}^5C_0 a^5 + {}^5C_1 a^{5-1}1 + {}^5C_2 a^{5-2}1^2 + {}^5C_3 a^{5-3}1^3 + {}^5C_4 a^{5-4}1^4 + {}^5C_5 1^5$$

$$= {}^5C_0 a^5 + {}^5C_1 a^4 + {}^5C_2 a^3 + {}^5C_3 a^2 + {}^5C_4 a + {}^5C_5 \dots (i)$$

$$(a-1)^5 = [{}^5C_0 a^5] + [{}^5C_1 a^{5-1}(-1)^1] + [{}^5C_2 a^{5-2}(-1)^2] + [{}^5C_3 a^{5-3}(-1)^3] + [{}^5C_4 a^{5-4}(-1)^4] + [{}^5C_5 (-1)^5]$$

$$= {}^5C_0 a^5 - {}^5C_1 a^4 + {}^5C_2 a^3 - {}^5C_3 a^2 + {}^5C_4 a - {}^5C_5 \dots (ii)$$

Subtracting (ii) from (i)

$$(a+1)^5 - (a-1)^5 = [{}^5C_0 a^5 + {}^5C_1 a^4 + {}^5C_2 a^3 + {}^5C_3 a^2 + {}^5C_4 a + {}^5C_5] - [{}^5C_0 a^5 - {}^5C_1 a^4 + {}^5C_2 a^3 - {}^5C_3 a^2 + {}^5C_4 a - {}^5C_5]$$



$$\begin{aligned}
&= 2[{}^5C_1a^4 + {}^5C_3a^2 + {}^5C_5] \\
&= 2 \left[ \left( \frac{5!}{1!(5-1)!} a^4 \right) + \left( \frac{5!}{3!(5-3)!} a^2 \right) + \left( \frac{5!}{5!(5-5)!} \right) \right] \\
&= 2[(5)a^4 + (10)a^2 + (1)] \\
&= 2[5a^4 + 10a^2 + 1] = (a+1)^5 - (a-1)^5 \\
&\text{Putting the value of } a, = \sqrt{3} \text{ in the above equation we get..} \\
&(\sqrt{3} + 1)^5 - (\sqrt{3} - 1)^5 = 2 [5(\sqrt{3})^4 + 10(\sqrt{3})^2 + 1] \\
&= 2[(5)(9) + (10)(3) + 1] \\
&= 2[45 + 30 + 1] \\
&= 152
\end{aligned}$$

29. We need to find the derivative of  $f(x) = (x^2 + 1)(x - 5)$

Derivative of a function  $f(x)$  from first principle is given by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \quad \{\text{where } h \text{ is a very small positive number}\}$$

$\therefore$  derivative of  $f(x) = (x^2 + 1)(x - 5)$  is given as

$$\begin{aligned}
f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
&\Rightarrow f'(x) = \lim_{h \rightarrow 0} \frac{\{(x+h)^2 + 1\}(x+h-5) - (x^2+1)(x-5)}{h} \\
&\Rightarrow f'(x) = \lim_{h \rightarrow 0} \frac{\{(x+h)^3 + x+h-5(x+h)^2 - 5\} - (x^3 - 5x^2 + x - 5)}{h}
\end{aligned}$$

Using  $(a+b)^2 = a^2 + 2ab + b^2$  and  $(a+b)^3 = a^3 + 3ab(a+b) + b^3$  we have:

$$\begin{aligned}
&\Rightarrow f'(x) = \lim_{h \rightarrow 0} \frac{\{x^3 + 3x^2h + 3h^2x + h^3 + x + h - 5x^2 - 10hx - 5h^2 - 5\} - (x^3 - 5x^2 + x - 5)}{h} \\
&\Rightarrow f'(x) = \lim_{h \rightarrow 0} \frac{\{3x^2h + 3h^2x + h^3 + h - 10hx - 5h^2\}}{h} \\
&\Rightarrow f'(x) = \lim_{h \rightarrow 0} \frac{h\{3x^2 + 3hx + h^2 + 1 - 10x - 5h\}}{h} \\
&\Rightarrow f'(x) = \lim_{h \rightarrow 0} \{3x^2 + 3hx + h^2 + 1 - 10x - 5h\} \\
&\Rightarrow f'(x) = 3x^2 + 3(0)x + 0^2 + 1 - 10x - 5(0) \\
&\Rightarrow f'(x) = 3x^2 - 10x + 1
\end{aligned}$$

OR

$$\text{Here } f(x) = \frac{x+1}{x-1}$$

$$\text{Then } f(x+h) = \frac{x+h+1}{x+h-1}$$

$$\text{We know that } f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$\begin{aligned}
&\Rightarrow f'(x) = \lim_{h \rightarrow 0} \frac{\frac{x+h+1}{x+h-1} - \frac{x+1}{x-1}}{h} \\
&= \lim_{h \rightarrow 0} \frac{(x+h+1)(x-1) - (x+1)(x+h-1)}{h(x+h-1)(x-1)} \\
&= \lim_{h \rightarrow 0} \frac{x^2 + x + xh - h + x - 1 - x^2 - xh + x - x - h + 1}{h(x+h-1)(x-1)} \\
&= \lim_{h \rightarrow 0} \frac{-2h}{h(x+h-1)(x-1)} = \frac{-2}{(x-1)^2}
\end{aligned}$$

30. Let  $a$  and  $b$  be the roots of required quadratic equation.

$$\text{Then A.M.} = \frac{a+b}{2} = 8$$

$\Rightarrow$

$$a + b = 16$$

$$\text{And G.M.} = \sqrt{ab} = 5$$

$$\Rightarrow ab = 25$$

Now, Quadratic equation  $x^2 - (\text{Sum of roots})x + (\text{Product of roots}) = 0$

$$\Rightarrow x^2 - (a+b)x + ab = 0$$

$$\Rightarrow x^2 - 16x + 25 = 0$$

Therefore, required equation is  $x^2 - 16x + 25 = 0$

OR

Let us take a G.P. whose first is  $a$  and common difference is  $r$ .

$$\therefore S_{\infty} = \frac{a}{1-r}$$

$$\Rightarrow \frac{a}{1-r} = 3 \dots (i)$$

And, sum of the terms of the G.P.  $a^2, (ar)^2, (ar^2)^2, \dots \infty$

$$S_{\infty} = \frac{a^2}{1-r^2}$$

$$\Rightarrow \frac{a^2}{1-r^2} = \frac{9}{2} \dots (ii)$$

$$\Rightarrow 2a^2 = 9(1-r^2)$$

$$\Rightarrow 2[3(1-r)]^2 = 9 - 9r^2 \text{ [From (i)]}$$

$$\Rightarrow 18(1+r^2-2r) = 9 - 9r^2$$

$$\Rightarrow 18 - 9 + 18r^2 + 9r^2 - 36r = 0$$

$$\Rightarrow 27r^2 - 36r + 9 = 0$$

$$\Rightarrow 3(9r^2 - 12r + 3) = 0$$

$$\Rightarrow 9r^2 - 12r + 3 = 0$$

$$\Rightarrow 9r^2 - 9r - 3r + 3 = 0$$

$$\Rightarrow 9r(r-1) - 3(r-1) = 0$$

$$\Rightarrow (9r-3)(r-1) = 0$$

$$\Rightarrow r = \frac{1}{3} \text{ and } r = 1$$

But,  $r = 1$  is not possible.

$$\therefore r = \frac{1}{3}$$

Now, substituting  $r = \frac{1}{3}$  in  $\frac{a}{1-r} = 3$

$$a = 3\left(1 - \frac{1}{3}\right)$$

$$\Rightarrow a = 3 \times \frac{2}{3} = 2$$

Therefore the first term is 2 and common difference is  $\frac{1}{3}$

31. Given that,  $n(T) = 12$

$$n(C) = 15$$

$$n(T \cap C) = 7$$

$$i. n(T \cup C) = n(T) + n(C) - n(T \cap C)$$

$$= 12 + 15 - 7$$

$$n(T \cup C) = 20$$

20 members like at least one of the two drinks.

ii. Only tea but not coffee

$$= n(T) - n(T \cap C)$$

$$= 12 - 7$$

$$= 5$$

iii. Only coffee but not tea

$$= n(C) - n(T \cap C)$$

$$= 15 - 7$$

$$= 8$$

iv. Neither tea nor coffee

$$= n(U) - n(T \cup C)$$

$$= 25 - 20$$

$$= 5$$

#### Section D

32. Given: set of first  $n$  natural numbers when  $n$  is an even number.

To find: the mean deviation about the mean

We know first  $n$  natural numbers are 1, 2, 3 ...,  $n$ . And given  $n$  is even number.

So mean is,

$$\bar{X} = \frac{1+2+3+\dots+n}{n} = \frac{\frac{n(n+1)}{2}}{n} = \frac{(n+1)}{2}$$



The deviations of numbers from the mean are as shown below,

$$1 - \frac{(n+1)}{2}, 2 - \frac{(n+1)}{2}, 3 - \frac{(n+1)}{2}, \dots, \frac{(n-2)}{2} - \frac{(n+1)}{2}, \frac{(n)}{2} - \frac{(n+1)}{2}, \frac{(n+2)}{2} - \frac{(n+1)}{2}, \dots, n - \frac{(n+1)}{2}$$

Or,

$$\frac{2-(n+1)}{2}, \frac{4-(n+1)}{2}, \frac{6-(n+1)}{2}, \dots, \frac{n-2-(n+1)}{2}, \frac{n-(n+1)}{2}, \frac{(n+2)-(n+1)}{2}, \dots, \frac{2n-(n+1)}{2}$$

$$\frac{2-(n+1)}{2}, \frac{4-(n+1)^2}{2}, \frac{6-(n+1)}{2}, \dots, \frac{-3}{2}, \frac{-1}{2}, \frac{1}{2}, \dots, \frac{2n-(n+1)}{2}$$

Or,

$$\frac{1-n}{2}, \frac{3-n}{2}, \frac{5-n}{2}, \dots, \frac{-3}{2}, \frac{-1}{2}, \frac{1}{2}, \dots, \frac{n-1}{2}$$

So the absolute values of deviation from the mean is

$$|x_i - \bar{x}| = \frac{(n-1)}{2}, \frac{(n-3)}{2}, \frac{(n-5)}{2}, \dots, \frac{3}{2}, \frac{1}{2}, \frac{1}{2}, \dots, \frac{n-1}{2}$$

The sum of absolute values of deviations from the mean, is

$$\sum |x_i - \bar{x}| = \frac{(n-1)}{2} + \frac{(n-3)}{2} + \frac{(n-5)}{2} + \dots + \frac{3}{2} + \frac{1}{2} + \frac{1}{2} + \dots + \frac{n-1}{2}$$

$$\sum |x_i - \bar{x}| = \left( \frac{1}{2} + \frac{3}{2} + \dots + \frac{(n-1)}{2} \right) \left( \frac{n}{2} \right)$$

Now we know sum of first n natural numbers =  $n^2$

Therefore, mean deviation about the mean is

$$M.D = \frac{\sum |x_i - \bar{x}|}{n} = \frac{\left( \frac{1}{2} + \frac{3}{2} + \dots + \frac{(n-1)}{2} \right) \left( \frac{n}{2} \right)}{n}$$

$$M.D = \frac{\sum |x_i - \bar{x}|}{n} = \frac{\left( \frac{n}{2} \right)^2}{n}$$

$$M.D = \frac{\sum |x_i - \bar{x}|}{n} = \frac{n^2}{4n} = \frac{n}{4}$$

33. Here the given equation are;

$$y = x^2 - 2x + 3$$

$$\Rightarrow x^2 - 2x = y - 3$$

$$\Rightarrow x^2 - 2x + 1 = y - 3 + 1$$


$$\Rightarrow (x - 1)^2 = y - 2 \quad \dots (i)$$

Now, shifting the origin to the point (1, 2) without rotating the axes and denoting new coordinates with respect to these axes by X and Y, we get,

$$x = X + 1, y = Y + 2 \quad \dots (ii)$$

Using these relations, equation (i) reduces to

$$X^2 = Y \quad \dots (iii)$$



This is of the form  $X^2 = 4aY$

Comparing, we get,

$$4a = 1 \text{ i.e., } a = 1/4$$

Vertex: Coordinates of the vertex with respect to the new axes are (X = 0, Y = 0).

So, the coordinates of the vertex with respect to the old axes are (1, 2) [Put X = 0, Y = 0 in (ii)]

Axis: The equation of the axis of the parabola with respect to the new axes is X = 0

So, the equation of the axis with respect to the old axes is x = 1 [Put X = 0 in (ii)]

Focus: The coordinates of the focus with respect to the new axes are (X = 0, Y = a) i.e.

$$(X = 0, Y = 1/4)$$

So, the coordinates of the focus S with respect to the old axes are

$$(1, 9/4) \quad [\text{Put } X = 0, Y = \frac{1}{4} \text{ in (ii)}]$$

Directrix: The equation of the directrix with respect to the new axes is Y = -a i.e. Y = -1/4

So, the equation of the directrix with respect to the old axes is

$$y = -\frac{1}{4} + 2 \text{ or } y = \frac{7}{4} \quad [\text{Put } Y = -\frac{1}{4} \text{ in (ii)}]$$

Latus-rectum: Length of the latus-rectum of given parabola is  $4a = 1$

OR

The equation of the ellipse is

$$25x^2 + 9y^2 - 150x - 90y + 225 = 0$$

$$\Rightarrow 25x^2 - 150x + 9y^2 - 90y = -225$$

$$\Rightarrow 25(x^2 - 6x) + 9(y^2 - 10y) = -225$$

$$\Rightarrow 25(x^2 - 6x + 9) + (y^2 - 10y + 25) = -225 + 225 + 225$$

$$\Rightarrow 25(x - 3)^2 + 9(y - 5)^2 = 225$$

$$\Rightarrow \frac{(x-3)^2}{9} + \frac{(y-5)^2}{25} = 1 \dots(i)$$

Shifting the origin at (3, 5) without rotating the coordinate axes and denoting the new coordinates with respect to the new axes by X and Y, we have .. (ii)

$$x = X + 3 \text{ and } y = Y + 5$$

Using these relations, equation (i) reduces to

$$\frac{X^2}{3^2} + \frac{Y^2}{5^2} = 1 \dots(iii)$$

Comparing equation (iii) with standard form  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ , we get

$$a^2 = 32 \text{ and } b^2 = 52.$$

$$\Rightarrow a = 4\sqrt{2} \text{ and } b = \sqrt{52}$$

Clearly,  $a < b$ . So, equation (iii) represents an ellipse whose major and minor axes along Y and X axes respectively.

Eccentricity:

$$e = \sqrt{1 - \frac{a^2}{b^2}} = \sqrt{1 - \frac{9}{25}} = \frac{4}{5}$$

Centre:

The coordinates of the centre with respect to new axes are (X = 0, Y = 0).

So, the coordinates of the centre with respect to old axes are (3, 5).

Vertices:

The vertices of the ellipse with respect to the new axes are (X = 0, Y =  $\pm b$ ) i.e. (X = 0, Y =  $\pm 5$ ). So, the vertices with respect to the old axes are

(3, 5  $\pm$  5) i.e. (3, 0) and (3, 10) [Putting X = 0, Y =  $\pm 5$  in (ii)]

Foci:

The coordinates of the foci with respect to the old axes are (X = 0, Y =  $\pm be$ ) i.e. (X = 0, Y =  $\pm 4$ ). So, the coordinates of the foci with respect to the old axes are

(3,  $\pm 4 + 5$ ) i.e. (3, 1) and (3, 9) [Putting X = 0, Y =  $\pm 4$  in (ii)]

Directrices:

The equations of the directrices with respect to the new axes are Y =  $\pm \frac{b}{e}$  i.e. Y =  $\pm \frac{25}{4}$

So, the equations of the directrices with respect to the old axes are

$$y = \pm \frac{25}{4} + 5 \text{ i.e. } y = -\frac{5}{4} \text{ and } y = \frac{45}{4} \text{ [Putting } Y = \pm \frac{25}{4} \text{ in (ii)]}$$

Axes:

Lengths of the major axis = 2b = 10,

Lengths of the Minor axis = 2a = 6.

Equation of the major axis with respect to the new axes is X = 0. So, the equation of the major axis with respect to the old axes is x = 3. [Putting X = 0 in (ii)]

The equation of the minor axis with respect to the new axes is Y = 0. So, the equation of the minor axis with respect to the old axes is y = 5. [Putting Y = 0 in (ii)]

$$\text{Latus-rectum: The length of the latus-rectum} = \frac{2a^2}{b} = \frac{2 \times 9}{5} = \frac{18}{5}$$

The equations of the latus-rectum with respect to the new axes are Y =  $\pm ae$  i.e. y =  $\pm Y \pm 4$ . So, the equations of the latus-rectum with respect to the old axes are

$$y = \pm 4 + 5 \text{ i.e. } y = 1 \text{ and } y = 9. \text{ [Putting } Y = \pm 4 \text{ in (ii)]}$$

34. The given system of linear inequalities is

$$-2 - \frac{x}{4} \geq \frac{1+x}{3} \dots (i)$$

$$\text{and } 3 - x < 4(x - 3) \dots (ii)$$


From inequality (i), we get

$$-2 - \frac{x}{4} \geq \frac{1+x}{3}$$

$$\begin{aligned}
&\Rightarrow -24 - 3x \geq 4 + 4x \text{ [multiplying both sides by 12]} \\
&\Rightarrow -24 - 3x - 4 \geq 4 + 4x - 4 \text{ [subtracting 4 from both sides]} \\
&\Rightarrow -28 - 3x \geq 4x \\
&\Rightarrow -28 - 3x + 3x \geq 4x + 3x \text{ [adding 3x on both sides]} \\
&\Rightarrow -28 \geq 7x \\
&\Rightarrow -\frac{28}{7} \geq \frac{7x}{7} \text{ [dividing both sides by 7]} \\
&\Rightarrow -4 \geq x \text{ or } x \leq -4 \dots \text{(iii)}
\end{aligned}$$

Thus, any value of  $x$  less than or equal to  $-4$  satisfied the inequality.

So, solution set is  $x \in (-\infty, -4]$




From inequality (ii), we get


$$\begin{aligned}
&3 - x < 4(x - 3) \\
&\Rightarrow 3 - x < 4x - 12 \\
&\Rightarrow 3 - x + 12 < 4x - 12 + 12 \text{ [adding 12 on both sides]} \\
&\Rightarrow 15 - x < 4x \\
&\Rightarrow 15 - x + x < 4x + x \text{ [adding x on both sides]} \\
&\Rightarrow 15 < 5x \\
&\Rightarrow 3 < x \text{ [dividing both sides by 5]} \\
&\text{or } x > 3 \dots \text{(iv)}
\end{aligned}$$

Thus, any value of  $x$  greater than 3 satisfies the inequality.

So, the solution set is  $x \in (3, \infty)$



The solution set of inequalities (i) and (ii) are represented graphically on number line as given below:



As no region is common, hence the given system has no solution.

$$\begin{aligned}
35. \text{ LHS} &= \tan(\alpha - \beta) = \frac{\sin 2\beta}{5 - \cos 2\beta} \\
&= \frac{\tan \alpha - \tan \beta}{1 + \tan \alpha \tan \beta} \\
&= \frac{\frac{3}{2} \tan \beta - \tan \beta}{1 + \frac{3}{2} \tan \beta \tan \beta} \dots [\because 2 \tan \alpha = 3 \tan \beta \Rightarrow \tan \alpha = \frac{3}{2} \tan \beta] \\
&= \frac{\tan \beta \left( \frac{3}{2} - 1 \right)}{1 + \frac{3}{2} \tan^2 \beta} \\
&= \frac{\frac{1}{2} \tan \beta}{1 + \frac{3}{2} \tan^2 \beta} \\
&= \frac{\frac{1}{2} \frac{\sin \beta}{\cos \beta}}{1 + \frac{3}{2} \left( \frac{\sin \beta}{\cos \beta} \right)^2} \dots [\because \tan \beta = \frac{\sin \beta}{\cos \beta}] \\
&= \frac{\frac{\sin \beta}{2 \cos \beta}}{1 + \frac{3 \sin^2 \beta}{2 \cos^2 \beta}} \\
&= \frac{\frac{\sin \beta}{2 \cos \beta}}{\frac{2 \cos^2 \beta + 3 \sin^2 \beta}{2 \cos^2 \beta}} \\
&= \frac{\sin \beta}{2 \cos \beta} \cdot \frac{2 \cos^2 \beta}{2 \cos^2 \beta + 3 \sin^2 \beta} \\
&= \frac{\sin \beta \cos \beta}{2 \cos^2 \beta + 3 \sin^2 \beta} \\
&= \frac{\sin 2\beta}{2(2 \cos^2 \beta + 3 \sin^2 \beta)} \dots \{ \because \sin 2x = 2(\sin x)(\cos x) \}
\end{aligned}$$

$$\begin{aligned}
&= \frac{\sin 2\beta}{2(1+\cos 2\beta)+3(1-\cos 2\beta)} \dots \{ \because 2 \cos^2 x = 1 + \cos 2x \text{ and } 2 \sin^2 x = 1 - \cos 2x \} \\
&= \frac{\sin 2\beta}{2+2 \cos 2\beta+3-3 \cos 2\beta} \\
&= \frac{\sin 2\beta}{5-\cos 2\beta}
\end{aligned}$$

LHS = RHS

Hence Proved

OR

We have to prove that  $\cos x = \pm \sqrt{2} \cos \frac{\alpha}{2}$ .

It is given that  $\sec(x + \alpha) + \sec(x - \alpha) = 2 \sec x$

$$\begin{aligned}
\Rightarrow \frac{1}{\cos(x+\alpha)} + \frac{1}{\cos(x-\alpha)} &= \frac{2}{\cos x} \dots \left[ \because \sec x = \frac{1}{\cos x} \right] \\
\Rightarrow \frac{\cos(x-\alpha) + \cos(x+\alpha)}{\cos(x+\alpha) \cos(x-\alpha)} &= \frac{2}{\cos x} \dots \left[ \because \cos A + \cos B = 2 \cos \frac{A+B}{2} \cos \frac{A-B}{2} \right] \\
\Rightarrow \frac{2 \cos \left( \frac{x+\alpha+x-\alpha}{2} \right) \cos \left( \frac{x+\alpha-x+\alpha}{2} \right)}{\cos(x+\alpha) \cos(x-\alpha)} &= \frac{2}{\cos x} \\
\Rightarrow \frac{2 \cos \left( \frac{2x}{2} \right) \cos \left( \frac{2\alpha}{2} \right)}{2 \cos(x+\alpha) \cos(x-\alpha)} &= \frac{1}{\cos x} \dots \{ \because 2 \cos A \cos B = \cos(A+B) + \cos(A-B) \} \\
\Rightarrow \frac{2 \cos x \cos \alpha}{\cos(x+\alpha+x-\alpha) + \cos(x+\alpha-x+\alpha)} &= \frac{1}{\cos x} \\
\Rightarrow \frac{2 \cos x \cos \alpha}{\cos 2x + \cos 2\alpha} &= \frac{1}{\cos x} \\
\Rightarrow 2 \cos^2 x \cos \alpha &= \cos 2x + \cos 2\alpha \\
\Rightarrow 2 \cos^2 x \cos \alpha &= 2 \cos^2 x - 1 + \cos 2\alpha \dots \{ \because \cos 2x = 2 \cos^2 x - 1 \} \\
\Rightarrow 2 \cos^2 x \cos \alpha - 2 \cos^2 x &= \cos 2\alpha - 1 \\
\Rightarrow 2 \cos^2 x (\cos \alpha - 1) &= 2 \cos 2\alpha - 1 - 1 \dots \{ \because \cos 2x = 2 \cos^2 x - 1 \} \\
\Rightarrow 2 \cos^2 x &= \frac{2 \cos^2 \alpha - 2}{\cos \alpha - 1} \\
\Rightarrow 2 \cos^2 x &= \frac{2(\cos^2 \alpha - 1)}{\cos \alpha - 1} \\
\Rightarrow 2 \cos^2 x &= \frac{(\cos \alpha - 1)(\cos \alpha + 1)}{\cos \alpha - 1} \\
\Rightarrow 2 \cos^2 x &= \cos \alpha + 1 \\
\Rightarrow 2 \cos^2 x &= 2 \cos^2 \frac{\alpha}{2} - 1 + 1 \dots \left[ \pm \sqrt{2} \cos \frac{\alpha}{2} \cos x = 2 \cos^2 \frac{x}{2} - 1 \right] \\
\Rightarrow 2 \cos^2 x &= 2 \cos^2 \frac{\alpha}{2} \\
\Rightarrow \cos x &= \pm \sqrt{2 \cos^2 \frac{\alpha}{2}} \\
\Rightarrow \cos x &= \pm \sqrt{2} \cos \frac{\alpha}{2}
\end{aligned}$$

Hence Proved.

### Section E

36. Read the text carefully and answer the questions:

#### Method to Find the Sets When Cartesian Product is Given

For finding these two sets, we write first element of each ordered pair in first set say A and corresponding second element in second set B (say).

#### Number of Elements in Cartesian Product of Two Sets

If there are p elements in set A and q elements in set B, then there will be pq elements in  $A \times B$  i.e. if  $n(A) = p$  and  $n(B) = q$ , then  $n(A \times B) = pq$ .

(i)  $n(A \times A) = 9$

$$\Rightarrow n(A) \times n(A) = 9 \Rightarrow n(A) = 3$$

$$(-1, 0) \in A \times A \Rightarrow -1 \in A, 0 \in A$$

$$(0, 1) \in A \times A \Rightarrow 0 \in A, 1 \in A$$

$$\Rightarrow -1, 0, 1 \in A$$

$$\text{Also, } n(A) = 3 \Rightarrow A = \{-1, 0, 1\}$$

$$\text{Hence, } A = \{-1, 0, 1\}$$

$$\text{Also, } A \times A = \{-1, 0, 1\} \times \{-1, 0, 1\}$$

$$= \{(-1, -1), (-1, 0), (-1, 1), (0, -1), (0, 0), (0, 1), (1, -1), (1, 0), (1, 1)\}$$

Hence, the remaining elements of  $A \times A$  are

$$(-1, -1), (-1, 1), (0, -1), (0, 0), (1, -1), (1, 0) \text{ and } (1, 1).$$



(ii) Given,  $(A \times B) = 6$  and  $(A \times B) = \{(1, 3), (2, 5), (3, 3)\}$

We know that Cartesian product of set  $A = \{a, b\}$  &  $B = \{c, d\}$  is  $A \times B = \{(a, c), (a, d), (b, c), (b, d)\}$

Therefore,  $A = \{1, 2, 3\}$  &  $B = \{3, 5\}$

$\Rightarrow A \times B = \{(1, 3), (1, 5), (2, 3), (2, 5), (3, 3), (3, 5)\}$

Thus, remaining elements are  $A \times B = \{(1, 5), (2, 3), (3, 5)\}$

(iii) If the set A has 3 elements and set B has 4 elements, then the number of elements in  $A \times B = 12$

OR

Clearly, A is the set of all first entries in ordered pairs in  $A \times B$  and B is the set of all second entries in ordered pairs in  $A \times B$

$\therefore A = \{a, b\}$  and  $B = \{1, 2, 3\}$

### 37. Read the text carefully and answer the questions:

Four friends Dinesh, Yuvraj, Sonu, and Rajeev are playing cards. Dinesh, shuffling a cards and told to Rajeev choose any four cards.



(i) Total number of possible outcomes  $= {}^{52}C_4$

We know that there are 12 face cards

$\therefore$  Number of favourable outcomes  $= {}^{12}C_4$

$\therefore$  Required probability  $= \frac{{}^{12}C_4}{{}^{52}C_4}$

(ii) Total number of possible outcomes  $= {}^{52}C_4$

We know that there are 26 red and 26 black cards.

$\therefore$  Number of favourable outcomes  $= {}^{26}C_2 \times {}^{26}C_2$

$\therefore$  Required probability  $= \frac{({}^{26}C_2)^2}{{}^{52}C_4}$

(iii) Total number of possible outcomes  $= {}^{52}C_4$

$\therefore$  Number of favourable outcomes  $= ({}^{13}C_1)^4$

$\therefore$  Required probability  $= \frac{(13)^4}{{}^{52}C_4}$

OR

Total number of possible outcomes  $= {}^{52}C_4$

In playing cards there are 4 king and 4 jack cards.

$\therefore$  Number of favourable outcomes  $= ({}^4C_2 \times {}^4C_2)$

$= 6 \times 6 = 36$

$\therefore$  Required probability  $= \frac{36}{{}^{52}C_4}$

### 38. Read the text carefully and answer the questions:

We have,  $i = \sqrt{-1}$ . So, we can write the higher powers of i as follows

i.  $i^2 = -1$

ii.  $i^3 = i^2 \cdot i = (-1) \cdot i = -i$

iii.  $i^4 = (i^2)^2 = (-1)^2 = 1$

iv.  $i^5 = i^{4+1} = i^4 \cdot i = 1 \cdot i = i$

v.  $i^6 = i^{4+2} = i^4 \cdot i^2 = 1 \cdot i^2 = -1...$

In order to compute  $i^n$  for  $n > 4$ , write

$i^n = i^{4q+r}$  for some  $q, r \in \mathbb{N}$  and  $0 \leq r \leq 3$ . Then,  $i^n = i^{4q} \cdot i^r$

$= (i^4)^q \cdot i^r = (1)^q \cdot i^r = i^r$

In general for any integer  $k$

$$i^{4k} = 1, i^{4k+1} = i, i^{4k+2} = -1 \text{ and } i^{4k+3} = -i$$

$$(i) \quad i^{30} = (i)^{4 \times 7 i^2} = -1$$

$$(ii) \quad i^{-39} = i(i^{-40})$$

$$= i((i^2)^{-20}) = i((-1)^{-20}) [\because i^2 = -1]$$

$$= i\left(\frac{1}{(-1)^{20}}\right) = i\left(\frac{1}{1}\right) = i = 0 + i(1)$$

Comparing with  $a + ib$ ,

$$a = 0, b = 1$$

$$0 + i$$

