

c) 150^0

d) 30^0

Section B**Attempt any 16 questions**

21. The distance between the graph of the equations $x = -3$ and $x = 2$ is [1]

a) 1

b) 3

c) 2

d) 5

22. Each side of an equilateral triangle measures 8 cm. The area of the triangle is [1]

a) $32\sqrt{3}\text{cm}^2$

b) 48cm^2

c) $16\sqrt{3}\text{cm}^2$

d) $8\sqrt{3}\text{cm}^2$

23. A linear equation in two variables is of the form $ax + by + c = 0$, where [1]


a) $a \neq 0$ and $b = 0$

b) $a = 0$ and $b = 0$

c) $a \neq 0$ and $b \neq 0$

d) $a = 0$ and $b \neq 0$

24. In the given figure, straight lines AB and CD intersect at O. If $\angle AOC + \angle BOD = 130^\circ$ then $\angle AOD = ?$ [1]



a) 110°

b) 65°

c) 115°

d) 125°

25. If $\sqrt{7} = 2.646$ then $\frac{1}{\sqrt{7}} = ?$ [1]

a) None of these

b) 0.375

c) 0.378

d) 0.441

26. The area of an isosceles triangle having base 2 cm and the length of one of the equal sides 4 cm, is [1]

a) $4\sqrt{15}\text{cm}^2$

b) $\sqrt{15}\text{cm}^2$

c) $2\sqrt{15}\text{cm}^2$

d) $\sqrt{\frac{15}{2}}\text{cm}^2$

27. The number of times a particular item occurs in a given data is called its. [1]

a) class-size

b) cumulative frequency

c) frequency

d) variation

28. $\frac{5^{n+2} - 6 \times 5^{n+1}}{13 \times 5^n - 2 \times 5^{n+1}}$ is equal to [1]

a) $\frac{3}{5}$

b) $-\frac{3}{5}$

c) $-\frac{5}{3}$

d) $\frac{5}{3}$

29. In $\triangle ABC$, $BD \perp AC$, $\angle CAE = 30^\circ$ and $\angle CBD = 40^\circ$. Then $\angle AEB = ?$ [1]



c) (ii) and (iii) are correct

d) (i) and (iii) are correct

36. The taxi fare in a city is as follows: For the first kilometer, the fare is ₹8 and for the subsequent distance it is ₹5 per kilometer. Taking the distance covered as x km and total fare as ₹ y , write a linear equation for this information. [1]


a) $y = 5x + 3$

b) $y = 5x - 3$

c) $x = 5y - 3$

d) $x = 5y + 3$

37. In figure, $x + y =$ [1]



a) 270°

b) 210°

c) 190°

d) 230°

38. The value of $\frac{x^{a(b-c)}}{x^{b(a-c)}} \div \left(\frac{x^b}{x^a}\right)^c$ is [1]

a) 1

b) 3

c) 4

d) 2

39. Let \bar{X} be the mean of x_1, x_2, \dots, x_n and \bar{Y} the mean of y_1, y_2, \dots, y_n . If \bar{Z} is the mean of $x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_n$, then \bar{Z} is equal to: [1]

a) $\frac{\bar{x} + \bar{y}}{n}$

b) $\frac{\bar{x} + \bar{y}}{2}$

c) $\bar{x} + \bar{y}$

d) $\frac{\bar{x} + \bar{y}}{2n}$

40. If the mean of x and $\frac{1}{x}$ is M , then the mean of x^2 and $\frac{1}{x^2}$ is [1]

a) $2M + 1$

b) $2M^2 + 1$


c) $2M - 1$

d) $2M^2 - 1$

Section C

Attempt any 8 questions

Question No. 41 to 45 are based on the given text. Read the text carefully and answer the questions:



Hareesh and Deep were trying to prove a theorem. For this they did the following;

- Draw a triangle ABC
- D and E are found as the mid points of AB and AC
- DE was joined and DE was extended to F so $DE = EF$

- a) x-axis
c) y-axis
- b) ordinate
d) origin
48. What the point **O (0,0)** is called? **[1]**
- a) x-axis
c) ordinate
- b) y-axis
d) origin
49. What is the ordinate of the ball of Arjun? **[1]**
- a) 2
c) 4
- b) 3
d) -3
50. What are the coordinates of the ball of Ashok? **[1]**
- a) (4, 3)
c) (3, 4)
- b) (4, 4)
d) (3, 3)

Solution

Section A

1. (d) 1

$$\begin{aligned} & \left(\frac{81}{16}\right)^{-\frac{3}{4}} \times \left\{ \left(\frac{25}{9}\right)^{-\frac{3}{2}} \div \left(\frac{5}{2}\right)^{-3} \right\} \\ & \Rightarrow \left(\frac{3}{2}\right)^{4 \times \frac{-3}{4}} \times \left\{ \left(\frac{5}{3}\right)^{2 \times \frac{-3}{2}} \div \left(\frac{5}{2}\right)^{-3} \right\} \\ & \Rightarrow \left(\frac{3}{2}\right)^{-3} \times \left\{ \left(\frac{5}{3}\right)^{-3} \div \left(\frac{5}{2}\right)^{-3} \right\} \end{aligned}$$

Explanation:

$$\begin{aligned} & \Rightarrow \left(\frac{3}{2}\right)^{-3} \times \left(\frac{5}{3} \times \frac{2}{5}\right)^{-3} \\ & \Rightarrow \left(\frac{3}{2}\right)^{-3} \times \left(\frac{2}{3}\right)^{-3} \\ & \Rightarrow \left(\frac{3}{2} \times \frac{2}{3}\right)^{-3} \\ & \Rightarrow (1)^{-3} = 1 \end{aligned}$$

2. (b) 4

Explanation: (2, 0) is a solution of the linear equation $2x + 3y = k$
 $\Rightarrow 4 = k$

3. (d) 80°

Explanation: We have :

$\angle AOC + \angle BOC = 180^\circ$ [Since AOB is a straight line]

$$\Rightarrow 4x + 5x = 180^\circ$$

$$\Rightarrow 9x = 180^\circ$$

$$\Rightarrow x = 20^\circ$$

$$\therefore \angle AOC = 4 \times 20^\circ = 80^\circ$$

4. (c) 500 cm^2

Explanation: Since diagonals of a rhombus divide it into 4 triangles of equal area. Therefore,
Area of rhombus = $4 \times$ Area of triangle
 $= 4 \times 125 = 500 \text{ sq. cm}$

5. (d) $\frac{\sqrt{7}+2}{3}$

Explanation: After rationalising:

$$\begin{aligned} \frac{1}{\sqrt{7}-2} &= \frac{1}{\sqrt{7}-2} \times \frac{\sqrt{7}+2}{\sqrt{7}+2} \\ &= \frac{\sqrt{7}+2}{(\sqrt{7})^2 - (2)^2} \\ &= \frac{\sqrt{7}+2}{7-4} \\ &= \frac{\sqrt{7}+2}{3} \end{aligned}$$

6. (a) Infinitely many

Explanation: There are many linear equations in 'x' and 'y' can be satisfied by $x = 1, y = 2$
for example

$$x + y = 3 \quad x - y = -1$$

$$2x + y = 4$$

and so on there are infinite number of examples

7. (b) 20°

Explanation: Let,

AB, CD and EF intersect at O

$$\angle COB = \angle AOD \text{ (Vertically opposite angle)}$$

$$\angle AOD = 3x + 10 \dots(i)$$

$$\angle AOE + \angle AOD + \angle DOF = 180^\circ \text{ (Linear pair)}$$

$$x + 3x + 10^\circ + 90^\circ = 180^\circ$$

$$4x + 100^\circ = 180^\circ$$

$$4x = 80^\circ$$

$$x = 20^\circ$$

8. (c) 25

Explanation: In the given figure $\angle CAD = \angle EAF$ (Vertically opposite angles)

$$\therefore \angle CAD = 30^\circ$$

In $\triangle ABD$,

$$\angle ABD + \angle BAD + \angle ADB = 180^\circ \text{ (Angle sum property)}$$

$$\Rightarrow (x + 10)^\circ + (x^\circ + 30^\circ) + 90^\circ = 180^\circ$$

$$\Rightarrow 2x + 130^\circ = 180^\circ$$

$$\Rightarrow 2x = 180^\circ - 130^\circ = 50^\circ$$

$$\Rightarrow x = 25$$

Thus, the value of x is 25.

Hence, the correct answer is 25.

9. (b) $\frac{1}{5}$

Explanation: $10^{2y} = 25$

$$10^{2y} = 5^2$$

$$(10^y)^2 = (5)^2$$

$$\Rightarrow 10^y = 5$$

Now 10^{-y}

$$= \frac{1}{10^y}$$

$$= \frac{1}{5}$$

10. (a) 93 marks

Explanation: Let, Vihaan obtains x marks in the fourth test.

So,

$$\frac{92+85+78+x}{4} = 87$$

$$\frac{255+x}{4} = 87$$

$$255 + x = 348$$

$$x = 348 - 255$$

$$x = 93 \text{ marks}$$

11. (c) 130°

Explanation: $\angle 2 = 180^\circ - \angle 1$

$$\angle 2 = 180^\circ - 50^\circ = 130^\circ$$

12. (c) $\frac{33}{2}$

Explanation: $g = t^{\frac{2}{3}} + 4t^{-\frac{1}{2}}$

$$= t^{\frac{2}{3}} + 4 \times \frac{1}{t^{\frac{1}{2}}}$$

$$= (64)^{\frac{2}{3}} + 4 \times \frac{1}{64^{\frac{1}{2}}}$$

$$= (4^3)^{\frac{2}{3}} + 4 \times \frac{1}{(8^2)^{\frac{1}{2}}}$$

$$= 4^{\frac{2}{3} \times 3} + 4 \times \frac{1}{8^{2 \times \frac{1}{2}}}$$

$$= 4^2 + \frac{4}{8}$$

$$= 16 + \frac{1}{2}$$

$$= \frac{33}{2}$$

13. (b) 4

Explanation: Let $\sqrt{a} + \frac{1}{\sqrt{a}} = x$

Then, squaring both side, we get

$$a + \frac{1}{a} + 2 = x^2$$

$$\Rightarrow \frac{a^2+1}{a} + 2 = x^2$$

Now, put the value of a,

$$\frac{(7-4\sqrt{3})^2+1}{7-4\sqrt{3}} + 2 = x^2$$

$$\Rightarrow \frac{49+48-56\sqrt{3}+1}{7-4\sqrt{3}} + 2 = x^2$$

$$\Rightarrow \frac{98-56\sqrt{3}}{7-4\sqrt{3}} + 2 = x^2$$

$$\Rightarrow 14 \left(\frac{7-4\sqrt{3}}{7-4\sqrt{3}} \right) + 2 = x^2$$

$$\Rightarrow 16 = x^2$$

$$\Rightarrow x = 4$$

$$\text{So, } x = \sqrt{a} + \frac{1}{\sqrt{a}} = 4$$

14. (c) 30°

Explanation: $40^\circ + x = 70^\circ$ (exterior angle)

$$\angle x = 70^\circ - 40^\circ$$

$$\angle x = 30^\circ$$

15. (c) 5 sq. units

Explanation: To find the area of the triangle formed by the line $2x + 5y = 10$ and co-ordinate axis

We put $x = 0$ in given equation at $x = 0$, we get $y = 2$

at $y = 0$ we get $x = 5$

So the line cut y-axis at 2 and x-axis at 5

So the height of the triangle is 2 unit and the base is 5 unit

Area of triangle = $\frac{1}{2}$ base \times height

$$= \frac{1}{2} \times 2 \times 5$$

$$= 5 \text{ sq. units}$$

16. (c) 38

Explanation: The mean of the six numbers is 23.

So the sum of six numbers is $23 \times 6 = 138$

After excluding one number, the mean of the remaining numbers is 20.


So the sum of five numbers is $20 \times 5 = 100$

The difference between them is

$$138 - 100 = 38$$

17. (a) $24\sqrt{5}$ cm

Explanation: Let ABC be a triangle in which sides $AB = 35$ cm, $BC = 54$ cm and $CA = 61$ cm



Now, semi-perimeter of a triangle,

$$s = \frac{a+b+c}{2} = \frac{35+54+61}{2} = \frac{150}{2} = 75 \text{cm}$$

$$[\because \text{semiperimeter, } s = \frac{a+b+c}{2}]$$

$$\therefore \text{Area of } \triangle ABC = \sqrt{s(s-a)(s-b)(s-c)} \text{ [by Heron's formula]}$$

$$\begin{aligned}
&= \sqrt{75(75 - 35)(75 - 54)(75 - 61)} \\
&= \sqrt{75 \times 40 \times 21 \times 14} \\
&= \sqrt{25 \times 3 \times 4 \times 2 \times 5 \times 7 \times 3 \times 7 \times 2} \\
&= 5 \times 2 \times 2 \times 3 \times 7\sqrt{5} \\
&= 420\sqrt{5}\text{cm}^2
\end{aligned}$$

Also, Area of $\triangle ABC = \frac{1}{2} \times AB \times \text{Altitude}$

$$\Rightarrow \frac{1}{2} \times 35 \times CD$$

$$\Rightarrow CD = \frac{420 \times 2\sqrt{5}}{35}$$

$$\therefore CD = 24\sqrt{5}$$

Hence, the length of altitude is $24\sqrt{5}\text{cm}$.

18. (b) $\left(a + \frac{1}{a}\right) \frac{\bar{x}}{2}$

Explanation: mean of ax_1, ax_2, \dots, ax_n , is $a\bar{x}$

mean of $\frac{x_1}{a}, \frac{x_2}{a}, \dots, \frac{x_n}{a}$ is $\frac{1}{a}\bar{x}$

so their mean is $\left(a + \frac{1}{a}\right) \frac{\bar{x}}{2}$

19. (b) $\frac{5}{3}$

Explanation: Let $x=1.666\dots$ ---(i)

multiply eq. (i) by 10, we get

$$10x = 16.666\dots$$
---(ii)

subtract eq(i) from (ii) we get

$$9x = 15$$

$$x = \frac{5}{3}$$

20. (d) 30^0

Explanation: Let one angle be x^0

Its supplementary angle will be $180^0 - x^0$

According to question

$$x = \frac{1}{5}(180^0 - x)$$

$$5x + x = 180^0$$

$$6x = 180^0$$

$$x = \frac{180}{6}$$

$$x = 30^0$$

Section B

21. (d) 5

Explanation: Distance between the graph of the equations $x = -3$ and $x = 2$ is $2 - (-3) = 5$ units

22. (c) $16\sqrt{3}\text{cm}^2$

Explanation: Area of equilateral triangle $= \frac{\sqrt{3}}{4} \times (\text{Side})^2$

$$= \frac{\sqrt{3}}{4} \times (8)^2$$

$$= \frac{\sqrt{3}}{4} \times 64$$

$$= 16\sqrt{3}\text{cm}^2$$

23. (c) $a \neq 0$ and $b \neq 0$

Explanation: A linear equation in two variables is of the form $ax + by + c = 0$ as a and b are coefficient of x and y

so if $a = 0$ and $b = 0$ or either of one is zero in that case the equation will be one variable or their will be no equation respectively.

therefore when $a \neq 0$ and $b \neq 0$ then only the equation will be in two variable

24. (c) 115^0

Explanation: We have:

$$\angle AOC = \angle BOD \text{ [Vertically-Opposite Angles]}$$

$$\begin{aligned} \therefore \angle AOC + \angle BOD &= 130^\circ \\ \Rightarrow \angle AOC + \angle AOC &= 130^\circ \quad [\because \angle AOC = \angle BOD] \\ \Rightarrow 2\angle AOC &= 130^\circ \\ \Rightarrow \angle AOC &= 65^\circ \end{aligned}$$

Now,

$$\begin{aligned} \angle AOC + \angle AOD &= 180^\circ \quad [\because \text{COD is a straight line}] \\ \Rightarrow 65^\circ + \angle AOD &= 180^\circ \\ \Rightarrow \angle AOC &= 115^\circ \end{aligned}$$

25. (c) 0.378

$$\begin{aligned} \text{Explanation: } \frac{1}{\sqrt{7}} &= \frac{1}{\sqrt{7}} \times \frac{\sqrt{7}}{\sqrt{7}} \\ &= \frac{\sqrt{7}}{7} \\ &= \frac{1}{7} \times \sqrt{7} \\ &= \frac{1}{7} \times 2.646 \\ &= 0.378 \end{aligned}$$

26. (b) $\sqrt{15} \text{ cm}^2$

$$\begin{aligned} \text{Explanation: } s &= \frac{4+4+2}{2} = 5 \text{ cm} \\ \text{Area of triangle} &= \sqrt{s(s-a)(s-b)(s-c)} \\ &= \sqrt{5(5-4)(5-4)(5-2)} \\ &= \sqrt{5 \times 1 \times 1 \times 3} \\ &= \sqrt{15} \text{ sq. cm} \end{aligned}$$

27. (c) frequency

Explanation: The number of times a particular item occurs in a given data is called its Frequency.

28. (c) $-\frac{5}{3}$

$$\begin{aligned} \text{Explanation: } &\frac{5^{n+2} - 6 \times 5^{n+1}}{13 \times 5^n - 2 \times 5^{n+1}} \\ &= \frac{5^n(5^2 - 6 \times 5^1)}{5^n(13 - 2 \times 5^1)} \\ &= \frac{5^2 - 6 \times 5}{13 - 2 \times 5} \\ &= \frac{25 - 30}{13 - 10} \\ &= \frac{-5}{3} \end{aligned}$$

29. (d) 80°

$$\begin{aligned} \text{Explanation: In BDC} \\ \angle BDC + \angle BCD + \angle DBC &= 180^\circ \\ BD \perp AC \\ \angle BCD &= 90^\circ, \angle DBC = 40^\circ \\ 90^\circ + \angle BCD + 40^\circ &= 180^\circ \\ \angle BCD + 130^\circ &= 180^\circ \\ \angle BCD &= 180^\circ - 130^\circ \\ \angle BCD &= 50^\circ \\ \angle AEB &= \angle CAE + \angle C \dots (\text{exterior angle}) \\ \angle CAE &= 30^\circ \\ \angle C &= 50^\circ \\ \angle AEB &= 30^\circ + 50^\circ \\ \angle AEB &= 80^\circ \end{aligned}$$

30. (b) $\frac{2x+3y}{5}$

Explanation: Average is equal to the sum of all the values in the data set divided by the number of values in the data set.

$$\begin{aligned} \text{Average} &= \frac{x+x+y+y+y}{5} \\ \text{Average} &= \frac{2x+3y}{5} \end{aligned}$$



31. (d) 20 cm

Explanation: Given: $s - a = 8$ cm, $s - b = 7$ cm and $s - c = 5$ cm

Adding all equations,

$$s - a + s - b + s - c = 8 + 7 + 5$$

$$\Rightarrow 3s - (a + b + c) = 20 \left[s = \frac{a+b+c}{2} \right]$$

$$\Rightarrow 3s - 2s = 20$$

$$\Rightarrow s = 20 \text{ cm}$$

32. (d) $5\sqrt{2}$

Explanation: $\sqrt{8} + 2\sqrt{32} - 5\sqrt{2}$

$$\Rightarrow 2\sqrt{2} + 2 \times 4\sqrt{2} - 5\sqrt{2}$$

$$\Rightarrow 10\sqrt{2} - 5\sqrt{2}$$

$$\Rightarrow 5\sqrt{2}$$

33. (a) $\frac{3}{2}x$

Explanation: From Figure, $\angle DOC = 180^\circ - \angle AOD$ (Both are Supplementary)

$$\Rightarrow \angle DOC = 180^\circ - 3y^\circ$$

Also, $\angle ACB = 180^\circ - \angle A - \angle B$

$$\Rightarrow \angle ACB = 180^\circ - x^\circ - 2x^\circ = 180^\circ - 3x^\circ$$

And $\angle ACD = 180^\circ - \angle ACB$

$$= 180^\circ - (180^\circ - 3x^\circ)$$

$$\Rightarrow \angle ACD = 3x^\circ$$

Now, in $\triangle OCD$

$$\angle DOC + \angle OCD + \angle D = 180^\circ$$

$$180^\circ - 3y^\circ + 3x^\circ + y^\circ = 180^\circ \quad [\angle OCD = \angle ACD]$$

$$\Rightarrow 2y^\circ = 3x^\circ$$

$$\Rightarrow y = \frac{3}{2}x$$

34. (a) $\bar{X} + k$

Explanation: Let us take n observations X_1, \dots, X_n

If \bar{X} be the mean of the n observations, then we have

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$$

$$\Rightarrow \sum_{i=1}^n X_i = n\bar{X}$$

Add a constant k to each of the observations. Then the observations becomes $X_i + k, \dots, X_n + k$

If \bar{Y} be the mean of the new observations, then we have

$$\bar{Y} = \frac{1}{n} \sum_{i=1}^n (X_i + k)$$

$$= \frac{1}{n} \sum_{i=1}^n X_i + \frac{1}{n} \sum_{i=1}^n k$$

$$= \bar{X} + \frac{1}{n} \cdot nk$$

$$= \bar{X} + k$$

35. (c) (ii) and (iii) are correct

Explanation: When two straight lines intersect them, Adjacent angles are supplementary and opposite angles are equal.

36. (a) $y = 5x + 3$

Explanation: Taxi fare for first kilometer = ₹8

Taxi fare for subsequent distance = ₹5

Total distance covered = x

Total fare = y

Since the fare for first kilometer = ₹8

According to problem, Fare for $(x - 1)$ kilometer = $5(x - 1)$

So, the total fare $y = 5(x - 1) + 8$

$$\Rightarrow y = 5(x - 1) + 8$$

$$\Rightarrow y = 5x - 5 + 8$$

$$\Rightarrow y = 5x + 3$$

Hence, $y = 5x + 3$ is the required linear equation.

37. (d) 230°

Explanation: In $\triangle ACO$

$$\angle ACO + \angle COA + \angle OAC = 180^\circ$$

Now, $\angle OAC = 180^\circ$

$$\Rightarrow 80^\circ + 40^\circ + 180^\circ - x^\circ = 180^\circ$$

$$\Rightarrow x^\circ = 120^\circ$$

$\angle BOD = \angle COA = 40^\circ$ (Opposite angles)

$$\angle BDO = 70^\circ$$

In $\triangle OBD$

$$\angle OBD = 180^\circ - 40^\circ - 70^\circ = 70^\circ$$

Also, $y^\circ = 180^\circ - \angle OBD = 180^\circ - 70^\circ = 110^\circ$

$$\Rightarrow x^\circ + y^\circ = 120^\circ + 110^\circ = 230^\circ$$

38. (a) 1

Explanation: $\frac{x^{a(b-c)}}{x^{b(a-c)}} \div \left(\frac{x^b}{x^a}\right)^c$

$$\Rightarrow \frac{x^{ab-ac}}{x^{ba-bc}} \div \left(\frac{x^{bc}}{x^{ac}}\right)$$

$$\Rightarrow x^{ab-ac-ab+bc} \div x^{bc-ac}$$

$$\Rightarrow x^{bc-ac} \div x^{bc-ac}$$

$$\Rightarrow 1$$

39. (b) $\frac{\bar{x} + \bar{y}}{2}$

Explanation: Since \bar{x} and \bar{y} are two numbers, though being means, their arithmetic mean is given by:

$$\bar{z} = \frac{\bar{x} \text{ and } \bar{y}}{2}$$

40. (d) $2M^2 - 1$

Explanation: Given, $\frac{x + \frac{1}{x}}{2} = M$

Taking square on both sides

$$\left(\frac{x + \frac{1}{x}}{2}\right)^2 = (M)^2$$

$$\left(x + \frac{1}{x}\right)^2 = (2M)^2$$

$$\left(x^2 + 2 + \frac{1}{x^2}\right) = (2M)^2$$

$$\left(x^2 + \frac{1}{x^2}\right) = 4M^2 - 2$$

Divide by 2 on both sides to get mean

$$\frac{\left(x^2 + \frac{1}{x^2}\right)}{2} = 2M^2 - 1$$

Section C

41. (a) SAS

Explanation: SAS

42. (d) $\angle ADE$

Explanation: $\angle ADE$

43. (c) $\angle DAE$

Explanation: $\angle DAE$

44. (a) BD

Explanation: BD

45. **(b)** BD
Explanation: BD
46. **(d)** (2, -3)
Explanation: (2, -3)
47. **(a)** x-axis
Explanation: x-axis
48. **(d)** origin
Explanation: origin
49. **(d)** -3
Explanation: -3
50. **(c)** (3, 4)
Explanation: (3, 4)

