

**Sample Question Paper - 23**  
**Mathematics-Basic (241)**  
**Class- X, Session: 2021-22**  
**TERM II**

**Time Allowed : 2 hours**


**Maximum Marks : 40**

**General Instructions :**

- The question paper consists of 14 questions divided into 3 sections A, B, C.
- Section A comprises of 6 questions of 2 marks each. Internal choice has been provided in two questions.
- Section B comprises of 4 questions of 3 marks each. Internal choice has been provided in one question.
- Section C comprises of 4 questions of 4 marks each. An internal choice has been provided in one question. It contains two case study based questions.


**SECTION - A**

- In the given figure, if  $\angle BAP = 150^\circ$ , then  $\angle AOB = k$ , then find the value of  $k$ .



**OR**

In the given figure,  $RS$  is the tangent to the circle at  $L$  and  $MN$  is a diameter. If  $\angle NML = 30^\circ$ , determine  $\angle RLM$ .



- Find the volume of the greatest sphere that can be cut off from a cylindrical log of wood of base radius 1 cm and height 5 cm.
- $(x + 2)$ ,  $x$  and  $(x - 1)$  are the frequencies of the numbers 12, 15 and 20 respectively. If the mean of the distribution is 14.5, then find the value of  $x$ .

**OR**

Mode for the following distribution is 17.5 and  $x$  is less than 6. Find  $x$ .


Class-interval	0 - 5	5 - 10	10 - 15	15 - 20	20 - 25
Frequency	5	2	3	6	$x$

- What is the common difference of an A.P. in which  $a_{21} - a_7 = 84$ ?

5. Solve :  $4x^2 - \sqrt{3}x - 5 = 0$
6. Find the value of mode, using an empirical relation, when it is given that mean and median are 10.5 and 9.6 respectively.


## SECTION - B

7. If  $p, q, r$  are in A.P., then find the value of  $p^3 + r^3 - 8q^3$  in terms of  $pqr$ .
8. In the given figure,  $O$  is the centre of a circle,  $AB$  is a chord and  $AT$  is the tangent at  $A$ . If  $\angle AOB = 110^\circ$ , then find  $\angle BAT$ .



OR

In the following figure,  $QS$  is the diameter and  $O$  is the centre of circle.  $APT$  is the tangent at  $P$ . Find  $\angle APQ$ .



9. A girl standing on the top of a 7 m high building observes that, the angle of elevation of the top of a tower is  $60^\circ$  and the angle of depression of the foot of the tower is  $30^\circ$ . Find the height of the tower.
10. Which term of the A.P. 4, 7, 10, 13, ....., is 49?

## SECTION - C

11. From the following data find the mode and median age of 150 residents of a colony who took part in swachh bharat abhiyan :

Age (in yrs.) more than or equal to	0	10	20	30	40	50
Number of residents	50	46	40	20	10	3

OR


Find the mean and mode of the following frequency distribution :

Class	0-10	10-20	20-30	30-40	40-50	50-60	60-70
Frequency	4	4	7	10	12	8	5


12. Draw a circle of radius 3 cm. From a point  $P$ , 7 cm away from its centre, draw two tangents to the circle. Measure the length of each tangent.

## Case Study - 1


13. Rohini went to a juice stall with her mother. While drinking juice she observed that shopkeeper has three types of glasses of inner diameter 5 cm to serve customers. The glass height is 10 cm and volume of type (A) glass is  $196.43 \text{ cm}^3$ .  $\left( \text{Use } \pi = \frac{22}{7} \right)$



(A)



(B)




(C)

- (i) Find the volume of type (B) glass.
- (ii) Which glass has minimum capacity?

### Case Study - 2

14. Two hoardings are put on two poles of equal heights standing on either side of the road. From a point between them on the road the angle of elevation of the top of poles are  $60^\circ$  and  $30^\circ$  respectively. Height of the each pole is 20 m.



Based on the above information, answer the following questions. (Take  $\sqrt{3} = 1.73$ ).

- (i) Find the length of  $PO$ .
- (ii) Find the width of the road.

## Solution

### MATHEMATICS BASIC 241

#### Class 10 - Mathematics

1.  $\angle OAP = 90^\circ$  [ $\because$  Tangent is perpendicular to the radius through the point of contact]  
 $\therefore \angle OAB = \angle BAP - \angle OAP = 150^\circ - 90^\circ = 60^\circ$   
 Now,  $OA = OB \Rightarrow \angle OAB = \angle OBA = 60^\circ$   
 $\therefore \angle AOB = 180^\circ - 2(60^\circ) = 60^\circ \Rightarrow k = 60$

**OR**

Join  $OL$ .


$OL \perp RS$ .

Also,  $OL = OM$  [Radii of the same circle]

$$\therefore \angle OML = \angle OLM$$

$$\Rightarrow \angle OLM = 30^\circ$$

$$\Rightarrow \angle RLM = 90^\circ - 30^\circ = 60^\circ$$



2. The radius of the greatest sphere that can be cut off from the cylinder = 1 cm

$$\therefore \text{Volume of the sphere} = \frac{4}{3}\pi r^3 = \frac{4}{3}\pi(1)^3 = \frac{4}{3}\pi \text{ cm}^3$$

$$3. \text{ We have, mean} = \frac{\sum f_i x_i}{\sum f_i}$$

$$\Rightarrow \frac{12(x+2)+15(x)+20(x-1)}{(x+2)+(x)+(x-1)} = 14.5$$

$$\Rightarrow 2(47x+4) = 29(3x+1)$$

$$\Rightarrow 94x+8 = 87x+29 \Rightarrow 7x = 21 \Rightarrow x = 3$$

**OR**

Given, mode = 17.5, which lies in the interval 15-20.

$\therefore$  15-20 is the modal class.

So,  $l = 15, f_0 = 3, f_1 = 6, f_2 = x$  and  $h = 5$

$$\therefore \text{Mode} = l + \left( \frac{f_1 - f_0}{2f_1 - f_0 - f_2} \right) \times h$$

$$\Rightarrow 17.5 = 15 + \left( \frac{6-3}{2 \times 6 - 3 - x} \right) \times 5$$

$$\Rightarrow 2.5 = \left( \frac{3}{9-x} \right) \times 5$$

$$\Rightarrow 22.5 - 2.5x = 15 \Rightarrow 2.5x = 7.5 \Rightarrow x = 3$$

Hence, the required value of  $x$  is 3.

4. Let  $a$  be 1<sup>st</sup> term and  $d$  be the common difference of the A.P.

According to the question,  $a_{21} - a_7 = 84$

$$\Rightarrow a + (21-1)d - (a + (7-1)d) = 84$$

$$\Rightarrow a + 20d - a - 6d = 84 \Rightarrow 14d = 84 \Rightarrow d = 6$$

$\therefore$  Common difference is 6.

5. We have,  $4x^2 - \sqrt{3}x - 5 = 0$

By quadratic formula, we have

$$x = \frac{-(-\sqrt{3}) \pm \sqrt{(-\sqrt{3})^2 - 4 \times (-5)(4)}}{2 \times 4} = \frac{\sqrt{3} \pm \sqrt{83}}{8}$$

6. We know that the empirical relationship is

$$\text{Mode} = 3 \text{ Median} - 2 \text{ Mean}$$

$$= 3(9.6) - 2(10.5) \quad [\because \text{Median} = 9.6 \text{ and Mean} = 10.5]$$

$$= 28.8 - 21.0 = 7.8$$

7. Since  $p, q, r$  are in A.P.

$$\therefore q - p = r - q \Rightarrow 2q = p + r \Rightarrow p + r - 2q = 0$$

$$\Rightarrow p^3 + r^3 + (-2q)^3 = 3 \times p \times r \times (-2q)$$

$$[\because \text{If } a + b + c = 0, \text{ then } a^3 + b^3 + c^3 = 3abc]$$

$$\Rightarrow p^3 + r^3 - 8q^3 = -6pqr$$

8. In  $\triangle OAB$ ,  $OA = OB$  [Radii of same circle]

$$\therefore \angle OAB = \angle OBA \quad [\because \text{Angles opposite to equal sides are equal}]$$

$$\text{Let } \angle OAB = \angle OBA = x$$

$$\text{In } \triangle AOB, \angle AOB + \angle OAB + \angle OBA = 180^\circ$$

$$\Rightarrow 110^\circ + x + x = 180^\circ \quad [\because \angle AOB = 120^\circ \text{ (Given)}]$$

$$\Rightarrow 2x = 180^\circ - 110^\circ \Rightarrow x = 35^\circ$$

$$\text{Now, } \angle OAT = \angle OAB + \angle BAT = 90^\circ$$

$[\because \text{Tangent is perpendicular to radius at point of contact}]$

$$\Rightarrow 35^\circ + \angle BAT = 90^\circ \Rightarrow \angle BAT = 55^\circ$$

**OR**

Join  $OP$

Now,  $OP \perp AT$

$$\Rightarrow \angle APO = 90^\circ$$

$[\because APT \text{ is tangent}]$


In  $\triangle QOP$

$$OQ = OP \text{ [Radii of same circle]}$$

$$\therefore \angle OPQ = \angle OQP = 30^\circ$$

$[\because \text{Angles opposite to equal sides of a triangle are equal}]$

$$\therefore \angle APQ = \angle APO - \angle OPQ = 90^\circ - 30^\circ = 60^\circ$$



9. Let  $AB = 7$  m be the height of building and  $CD$  be the height of tower. Now,  $AB = DE = 7$  m

Also,  $BD = AE$

In  $\triangle ABD$ ,

$$\frac{AB}{BD} = \tan 30^\circ$$

$$\Rightarrow \frac{7}{BD} = \frac{1}{\sqrt{3}}$$

$$\Rightarrow BD = 7\sqrt{3} \text{ m} = AE$$

$$\text{In } \triangle ACE, \frac{CE}{AE} = \tan 60^\circ$$

$$\Rightarrow \frac{CE}{7\sqrt{3}} = \sqrt{3} \Rightarrow CE = (\sqrt{3} \times 7\sqrt{3}) \text{ m} = 21 \text{ m}$$

$$\therefore CD = CE + ED = (21 + 7) \text{ m} = 28 \text{ m}$$

Thus, the height of the tower is 28 m.

10. The given A.P. is 4, 7, 10, 13, ...

Here,  $a = 4$ ,  $d = 7 - 4 = 3$

Let the  $n^{\text{th}}$  term of the A.P. be 49.

$$\text{Then, } a_n = a + (n - 1)d \Rightarrow 49 = 4 + (n - 1)(3)$$

$$\Rightarrow 45 = 3(n - 1) \Rightarrow n - 1 = 15 \Rightarrow n = 16$$

Hence, 16<sup>th</sup> term of the A.P. is 49.

11. The frequency distribution table for the given data can be drawn as :

Class	Cumulative frequency (c.f.)	Frequency ( $f_i$ )
0-10	50	4
10-20	46	6
20-30	40	20
30-40	20	10
40-50	10	7
50-60	3	3
Total		50

$$\text{Here } \frac{N}{2} = \frac{50}{2} = 25$$

$\therefore$  Median class is 20-30.

$$\therefore \text{Median} = 20 + \left( \frac{25 - 20}{20} \right) \times 10 = 20 + 2.5 = 22.5$$

Now, maximum frequency is 20.

$\therefore$  Modal class is 20-30

$$\therefore \text{Mode} = 20 + \left[ \frac{20 - 6}{2(20) - 6 - 10} \right] \times 10$$

$$= 20 + \left[ \frac{14}{24} \right] \times 10 = 25.83$$

OR

The frequency distribution table for the given data can be drawn as :

Class	( $x_i$ )	( $f_i$ )	$f_i x_i$
0-10	5	4	20
10-20	15	4	60
20-30	25	7	175
30-40	35	10	350
40-50	45	12	540
50-60	55	8	440
60-70	65	5	325
		$\Sigma f_i = 50$	$\Sigma f_i x_i = 1910$

$$\text{Mean} = \frac{\Sigma f_i x_i}{\Sigma f_i} = \frac{1910}{50} = 38.2$$

$$\text{Mode} = l + \left[ \frac{f_1 - f_0}{2f_1 - f_0 - f_2} \right] \times h$$


The maximum frequency is 12.

$\therefore$  Modal class is 40-50.

$$\begin{aligned} \therefore \text{Mode} &= 40 + \left[ \frac{12 - 10}{2(12) - 10 - 8} \right] \times 10 \\ &= 40 + \left[ \frac{2}{24 - 18} \right] \times 10 = 40 + \frac{10}{3} = \frac{130}{3} = 43.3 \end{aligned}$$

## 12. Steps of construction :

**Step-I :** Draw a circle of radius 3 cm, taking O as centre and OC be its radius.



**Step-II :** Produce OC to P such that OP = 7 cm.

**Step-III :** Draw perpendicular bisector of OP that meets OP at Q.

**Step-IV :** Taking Q as centre and radius QP draw a circle which intersect previous circle at points A and B.

**Step-V :** Join P to A and P to B.

Thus, PA and PB are the required tangents.

Now, join OA.

In  $\triangle AOP$ ,  $\angle OAP = 90^\circ$  [Angle in semicircle]

$\therefore AP^2 = OP^2 - OA^2$  [By Pythagoras theorem]

$$= 7^2 - 3^2 = 40$$

$$\Rightarrow AP = 6.32 \text{ cm}$$

Similarly,  $BP = 6.32$  cm

Thus, length of each tangent = 6.32 cm

13. Diameter of each glass = 5 cm

$\therefore$  Radius of each glass = 2.5 cm

Height of each glass = 10 cm

(i) Volume of type (B) glass

= Volume of type (A) glass - Volume of hemisphere

$$= 196.43 - \frac{2}{3} \pi r^3 = 196.43 - \frac{2}{3} \times \frac{22}{7} \times 2.5 \times 2.5 \times 2.5$$

$$= 196.43 - 32.74 = 163.7 \text{ cm}^3$$

(ii) Volume of type (C) glass = Volume of type (A) glass - Volume of cone

$$= 196.43 - \frac{1}{3} \pi r^2 h = 196.43 - \frac{1}{3} \times \frac{22}{7} \times 2.5 \times 2.5 \times 1.5$$

$$= 196.43 - 9.82 = 186.61 \text{ cm}^3$$

Glass (B) has minimum capacity.

14. (i) In  $\triangle OPQ$ , we have

$$\tan 60^\circ = \frac{PQ}{PO}$$

$$\Rightarrow \sqrt{3} = \frac{20}{PO} \Rightarrow PO = \frac{20}{\sqrt{3}} \text{ m}$$

(ii) In  $\triangle ORS$ , we have

$$\tan 30^\circ = \frac{RS}{OR} \Rightarrow \frac{1}{\sqrt{3}} = \frac{20}{OR} \Rightarrow OR = 20\sqrt{3} \text{ m}$$

Clearly, width of the road =  $PR$

$$= PO + OR = \left( \frac{20}{\sqrt{3}} + 20\sqrt{3} \right) \text{ m}$$

$$= 20 \left( \frac{4}{\sqrt{3}} \right) \text{ m} = \frac{80}{\sqrt{3}} \text{ m} = 46.24 \text{ m}$$

