

**Sample Question Paper - 3**  
**Class- X Session- 2021-22 TERM 1**  
**Subject- Mathematics (Basic)**

**Time Allowed: 1 hour and 30 minutes**

**Maximum Marks: 40**

**General Instructions:**

1. The question paper contains three parts A, B and C.
2. Section A consists of 20 questions of 1 mark each. Attempt any 16 questions.
3. Section B consists of 20 questions of 1 mark each. Attempt any 16 questions.
4. Section C consists of 10 questions based on two Case Studies. Attempt any 8 questions.
5. There is no negative marking.

**Section A**


**Attempt any 16 questions**

1. The decimal expansion of  $\frac{987}{10500}$  will terminate after: [1]
  - a) 2 decimal places
  - b) 3 decimal places
  - c) 1 decimal place
  - d) None of these
2. A fraction becomes  $\frac{9}{11}$ , if 2 is added to both the numerator and denominator. If 3 is added to both the numerator and denominator it becomes  $\frac{5}{6}$ , then the fraction is [1]
  - a)  $\frac{9}{7}$
  - b)  $\frac{-9}{7}$
  - c)  $\frac{7}{9}$
  - d)  $\frac{-7}{9}$
3. If  $\alpha$  and  $\beta$  are the zeroes of the polynomial  $3x^2 + 11x - 4$ , then the value of  $\frac{1}{\alpha} + \frac{1}{\beta}$  is [1]
  - a)  $\frac{13}{4}$
  - b)  $\frac{12}{4}$
  - c)  $\frac{11}{4}$
  - d)  $\frac{15}{4}$
4. A system of linear equations is said to be inconsistent if it has [1]
  - a) one solution
  - b) at least one solution
  - c) two solutions
  - d) no solution
5. If  $\tan \theta = \sqrt{3}$ , then  $\sec \theta =$  [1]
  - a)  $\sqrt{\frac{3}{2}}$
  - b) 2
  - c)  $\frac{2}{\sqrt{3}}$
  - d)  $\frac{1}{\sqrt{3}}$
6. If the LCM of a and 18 is 36 and the HCF of a and 18 is 2, then a = [1]
  - a) 1
  - b) 2
  - c) 4
  - d) 3





16. If  $\sin \theta - \cos \theta = 0$ , then the value of  $\theta$  is [1]  
 a)  $60^\circ$  b)  $30^\circ$   
 c)  $45^\circ$  d)  $90^\circ$
17. If a pair of linear equations in two variables is consistent, then the lines represented by two equations are [1]  
 a) parallel b) always coincident  
 c) intersecting d) intersecting or coincident
18. A card is selected from a deck of 52 cards. The probability of its being a red face card is [1]  
 a)  $\frac{3}{13}$  b)  $\frac{1}{2}$   
 c)  $\frac{2}{12}$  d)  $\frac{3}{26}$
19. The number  $\frac{\sqrt{5}+\sqrt{2}}{\sqrt{5}-\sqrt{2}}$  is [1]  
 a) an irrational number b) an integer  
 c) not a real number d) a rational number
20. In the figure, the area of  $\triangle ABC$  (in square units) is [1]



- a) 10 b) 2.5  
 c) 7.5 d) 15

### Section B

Attempt any 16 questions

21. The pair of equations  $5x - 15y = 8$  and  $3x - 9y = \frac{24}{5}$  has [1]  
 a) infinitely many solutions b) no solution  
 c) two solutions d) one solution
22. If  $f(x) = ax^2 + bx + c$  has no real zeros and  $a + b + c < 0$ , then [1]  
 a)  $c > 0$  b)  $c < 0$   
 c) None of these d)  $c = 0$
23. The least number that is divisible by all the numbers from 1 to 10 (both inclusive) is [1]  
 a) 100 b) 10  
 c) 504 d) 2520

24. If  $\sin\theta = \frac{a}{b}$  then  $\cos\theta = ?$  [1]  
 a)  $\frac{b}{\sqrt{b^2-a^2}}$  b)  $\frac{\sqrt{b^2-a^2}}{b}$   
 c)  $\frac{b}{a}$  d)  $\frac{a}{\sqrt{b^2-a^2}}$
25. The difference between two numbers is 26 and one number is three times the other. The numbers are [1]  
 a) 39 and 13 b) 30 and 10  
 c) 36 and 12 d) 36 and 10
26. If the zeroes of the quadratic polynomial  $x^2 + (a + 1)x + b$  are 2 and -3, then [1]  
 a)  $a = 0, b = -6$  b)  $a = 5, b = -1$   
 c)  $a = -7, b = -1$  d)  $a = 2, b = -6$
27. A semicircle is drawn on AC. Two chords AB and BC of length 8 cm and 6 cm respectively are drawn in the semicircle. What will be the measure of the diameter of the circle? [1]  
 a) 12 cm. b) 10 cm.  
 c) 14 cm. d) 11 cm.
28. If the distance between the points (2, -2) and (-1, x) is 5, one of the values of x is [1]  
 a) -2 b) -1  
 c) 1 d) 2
29. If  $2x = \sec A$  and  $\frac{2}{x} = \tan A$  then  $2\left(x^2 - \frac{1}{x^2}\right) = ?$  [1]  
 a)  $\frac{1}{2}$  b)  $\frac{1}{4}$   
 c)  $\frac{1}{16}$  d)  $\frac{1}{8}$
30. Half the perimeter of a rectangular garden, whose length is 4m more than its width is 36m. The area of the garden is [1]  
 a) 320 m<sup>2</sup> b) 300 m<sup>2</sup>  
 c) 400 m<sup>2</sup> d) 360 m<sup>2</sup>
31. If two positive integers m and n can be expressed as  $m = x^2y^5$  and  $n = x^3y^2$ , where x and y are prime numbers, then HCF(m, n) = [1]  
 a)  $x^2y^2$  b)  $x^2y^3$   
 c)  $x^3y^2$  d)  $x^3y^3$
32. If  $\triangle ABC$  and  $\triangle DEF$  are two triangles such that  $\frac{AB}{DE} = \frac{BC}{EF} = \frac{CA}{FD} = \frac{2}{5}$ , then Area ( $\triangle ABC$ ): Area ( $\triangle DEF$ ) = [1]  
 a) 2 : 5 b) 4 : 15  
 c) 8 : 125 d) 4 : 25
33. If  $7 \tan \theta = 4$  then  $\frac{(7 \sin \theta - 3 \cos \theta)}{(7 \sin \theta + 3 \cos \theta)} = ?$  [1]


- a)  $\frac{3}{7}$     b)  $\frac{1}{7}$   
 c)  $\frac{5}{14}$     d)  $\frac{5}{7}$
34. The mid-point of the line segment joining the points A (-2, 8) and B (- 6, - 4) is [1]  
 a) (- 4, -6)    b) (4, 2)  
 c) (2, 6)     d) (- 4, 2)
35. A piggy bank contains 100 fifty paise coins, 50 one rupee coins, 20 two rupee coins and 10 five rupee coins. One coin is drawn at random. The probability that the coin drawn will not be a five rupee coin is [1]  
 a)  $\frac{5}{9}$     b)  $\frac{7}{18}$   
 c)  $\frac{8}{9}$     d)  $\frac{17}{18}$
36. If the lines given by  $3x + 2ky = 2$  and  $2x + 5y + 1 = 0$  are parallel, then the value of k is [1]  
 a)  $-\frac{5}{4}$      b)  $\frac{3}{2}$   
 c)  $\frac{15}{4}$      d)  $\frac{2}{5}$
37. Which of the following is an irrational number? [1]  
 a) 3.141141114...                                     b)  $3.\overline{1416}$   
 c)  $\frac{22}{7}$      d) 3.1416
38. If  $x = a \cos \theta$  and  $y = b \sin \theta$ , then  $b^2x^2 + a^2y^2 =$  [1]  
 a)  $a^2 + b^2$     b)  $ab$   
 c)  $a^4b^4$      d)  $a^2b^2$
39. A die is thrown once. Find the probability of getting an even number: [1]  
 a)  $\frac{1}{2}$      b)  $\frac{1}{3}$   
 c)  $\frac{2}{3}$      d)  $\frac{1}{4}$
40. The point P which divides the line segment joining the points A(2, -5) and B(5, 2) in the ratio 2 : 3 lies in the quadrant. [1]  
 a) III    b) I  
 c) II     d) IV

**Section C**

**Attempt any 8 questions**

**Question No. 41 to 45 are based on the given text. Read the text carefully and answer the questions:**

Preeti visited to her uncles house. From point A, where Artina was standing, a bus and building come in a straight line as shown in the figure.



41. Which similarity criteria can be seen in this case, if bus and building are considered in a straight line? [1]
- a) SAS b) SSS  
c) ASA d) AA
42. If the distance between Preeti and the bus is twice as much as the height of the bus, then the height of the bus is [1]
- a) 15 m b) 12.5 m  
c) 40 m d) 25 m
43. If the distance of Preeti from the building is twelve times the height of the bus, then the ratio of the heights of bus and building is [1]
- a) 1 : 6 b) 3 : 1  
c) 1 : 4 d) 2 : 3
44. What is the ratio of the distance between Preeti and top of bus to the distance between the tops of bus and building? [1]
- a) 2 : 5 b) Can't be determined  
c) 1 : 6 d) 1 : 5
45. What is the height of the building? [1]
- a) 120 m b) 30 m  
c) 75 m d) 50 m

**Question No. 46 to 50 are based on the given text. Read the text carefully and answer the questions:**

While doing dusting a maid found a button whose upper face is of black colour, as shown in the figure. The diameter of each of the smaller identical circles is  $\frac{1}{4}$  of the diameter of the larger circle whose radius is 16 cm.



46. The area of each of the smaller circle is [1]
- a)  $52.3 \text{ cm}^2$  b)  $50.28 \text{ cm}^2$   
c)  $46.39 \text{ cm}^2$  d)  $40.28 \text{ cm}^2$
47. The area of the larger circle is [1]
- a)  $855.57 \text{ cm}^2$  b)  $804.57 \text{ cm}^2$   
c)  $704.57 \text{ cm}^2$  d)  $990.57 \text{ cm}^2$

48. The area of the black colour region is [1]
- a)  $623.45 \text{ cm}^2$                       b)  $603.45 \text{ cm}^2$   
c)  $610.45 \text{ cm}^2$                       d)  $600.45 \text{ cm}^2$
49. The area of quadrant of a smaller circle is [1]
- a)  $12 \text{ cm}^2$                               b)  $12.57 \text{ cm}^2$   
c)  $11.57 \text{ cm}^2$                       d)  $13.68 \text{ cm}^2$
50. If two concentric circles are of radii 2 cm and 5 cm, then the area between them is [1]
- a)  $66 \text{ cm}^2$                               b)  $60 \text{ cm}^2$   
c)  $68 \text{ cm}^2$                               d)  $63 \text{ cm}^2$

# Solution

## Section A

1. (b) 3 decimal places

**Explanation:**  $\frac{987}{10500} = \frac{47}{500} = \frac{47}{2^2 \times 5^3}$  Here, in the denominator of the given fraction the highest power of prime factor 5 is 3, therefore, the decimal expansion of the rational number  $\frac{47}{2^2 \times 5^3}$  will terminate after 3 decimal places.

2. (c)  $\frac{7}{9}$

**Explanation:** Let the fraction be  $\frac{x}{y}$ .

According to question

$$\frac{x+2}{y+2} = \frac{9}{11}$$

$$\Rightarrow 11x + 22 = 9y + 18$$

$$\Rightarrow 11x - 9y = -4 \dots (i)$$

And  $\frac{x+3}{y+3} = \frac{5}{6}$

$$\Rightarrow 6x + 18 = 5y + 15$$

$$\Rightarrow 6x - 5y = -3 \dots (ii)$$

On solving eq. (i) and eq. (ii), we get

$$x = 7, y = 9$$

Therefore, the fraction is  $\frac{7}{9}$

3. (c)  $\frac{11}{4}$

**Explanation:** Here  $a = 3, b = 11, c = -4$  Since  $\frac{1}{\alpha} + \frac{1}{\beta} = \frac{\alpha + \beta}{\alpha\beta}$

$$\alpha + \beta = \frac{-11}{3}, \alpha\beta = \frac{-4}{3}$$

So,  $\frac{\frac{-11}{3}}{\frac{-4}{3}} = \frac{11}{4}$

4. (d) no solution

**Explanation:** A system of linear equations is said to be inconsistent if it has no solution means two lines are running parallel and not cutting each other at any point.

5. (b) 2

**Explanation:** Since  $\sec \theta = \sqrt{1 + \tan^2 \theta}$

$$\therefore \sec \theta = \sqrt{1 + (\sqrt{3})^2}$$

$$= \sqrt{1 + 3} = \sqrt{4} = 2$$

6. (c) 4

**Explanation:** LCM (a, 18) = 36

$$\text{HCF (a, 18)} = 2$$

We know that the product of numbers is equal to the product of their HCF and LCM.

Therefore,

$$18a = 2(36)$$

$$a = \frac{2(36)}{18}$$

$$a = 4$$

7. (a)  $\frac{-9}{2}$

**Explanation:** For  $ax^2 + bx + c$ , we have  $\alpha\beta = \frac{c}{a}$

For  $2x^2 + 5x - 9$ , we have  $\alpha\beta = \frac{-9}{2}$





8. (a) 5

**Explanation:** Three vertices of a rectangle ABCD are B (4,0), C (4, 3) and D (0, 3) length of one of its diagonals

$$BD = \sqrt{(4-0)^2 + (0-3)^2} = \sqrt{4^2 + 3^2} \\ = \sqrt{16+9} = \sqrt{25} = 5$$

9. (b) cannot be both positive

**Explanation:**


Let  $p(x) = x^2 + kx + k, k \neq 0$

On comparing  $p(x)$  with  $ax^2 + bx + c$ , we get

$$a = 1, b = k \text{ and } c = k$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad \text{[by quadratic formula]}$$

$$= \frac{-k \pm \sqrt{k^2 - 4k}}{2 \times 1}$$

$$= \frac{-k \pm \sqrt{k(k-4)}}{2}, k \neq 0$$


Here, we see that  $k(k-4) > 0$

$$\Rightarrow k \in (-\infty, 0) \cup (4, \infty)$$

In quadratic polynomial  $ax^2 + bx + c$

If  $a > 0, b > 0, c > 0$  or  $a < 0, b < 0, c < 0$ , then the polynomial has always all negative zeroes.

and if  $a > 0, c < 0$  or  $a < 0, c > 0$ , then the polynomial has always zeroes of opposite sign.

Case I: If  $k \in (-\infty, 0)$  i.e.,  $k < 0$

$$\Rightarrow a = 1 > 0, \quad b, c = k < 0$$

So, both zeroes are of opposite sign.

Case II: If  $k \in (4, \infty)$  i.e.,  $k \geq 4$

$$\Rightarrow a = 1 > 0, b, c > 4$$

So, both zeroes are negative.

Hence, in any case zeroes of the given quadratic polynomial cannot both be positive.

10. (a)  $a > 0, b < 0$  and  $c > 0$

**Explanation:** Clearly,  $f(x) = ax^2 + bx + c$  represent a parabola opening upwards.

Therefore,  $a > 0$

The vertex of the parabola is in the fourth quadrant, therefore  $b < 0$

$y = ax^2 + bx + c$  cuts Y axis at P which lies on OY.

Putting  $x = 0$  in  $y = ax^2 + bx + c$ , we get  $y = c$ .

So the coordinates of P is (0, c).

Clearly, P lies on OY.  $\Rightarrow c > 0$

Hence,  $a > 0, b < 0$  and  $c > 0$

11. (d)  $\frac{1}{3}$

**Explanation:** Total outcomes of selecting a number from 30 numbers = 30

Favourable numbers (prime numbers) = 10,

i.e., (2, 3, 5, 7, 11, 13, 17, 19, 23, 29)

$$\therefore \text{Probability of selecting a prime number} = \frac{10}{30} = \frac{1}{3}$$

12. (c) 180

**Explanation:** It is given that:  $a = (2^2 \times 3^3 \times 5^4)$  and  $b = (2^3 \times 3^2 \times 5)$

$\therefore$  HCF (a, b) = Product of smallest power of each common prime factor in the numbers =  $2^2 \times 3^2 \times 5 = 180$

13. (a) 7

**Explanation:** Given that R is the mid- point of the line segment AB.

$$\text{Th y-coordinate of R} = \frac{5+y}{2}$$

$$\Rightarrow y = 7$$

14. (c) 6

**Explanation:** The given points are A(5,0), B(8, 0) and C(8, 4)

$$\therefore (x_1 = 5, y_1 = 0), (x_2 = 8, y_2 = 0) \text{ and } (x_3 = 8, y_3 = 4)$$

The area of the triangle



$$\begin{aligned}
&= \frac{1}{2} |x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2)| \\
&= \frac{1}{2} |(0 - 4) + 8(4 - 0) + 8(0)| \\
&= \frac{1}{2} |-20 + 32 + 9| \\
&= \frac{1}{2} \times 12 \\
&= 6 \text{ sq. units}
\end{aligned}$$

15. (d) 6 and 8

**Explanation:** Sum of the zeroes of the polynomial  $= \frac{-b}{a} = \frac{6}{1} = 6$   
 And Product of the zeroes of the polynomial  $= \frac{c}{a} = \frac{8}{1} = 8$

16. (c)  $45^\circ$

**Explanation:** Given:  $\sin \theta - \cos \theta = 0$   
 $\Rightarrow \sin \theta = \cos \theta$   
 $\Rightarrow \sin \theta = \sin(90^\circ - \theta)$   
 $\Rightarrow \theta = 90^\circ - \theta$   
 $\Rightarrow 2\theta = 90^\circ$   
 $\Rightarrow \theta = 45^\circ$

17. (d) intersecting or coincident

**Explanation:** If a pair of linear equations in two variables is consistent, then its solution exists.  
 $\therefore$  The lines represented by the equations are either intersecting or coincident.

18. (d)  $\frac{3}{26}$

**Explanation:** In a deck of 52 cards, there are 12 face cards i.e. 6 red (3 hearts and 3 diamonds) and 6 black cards (3 spade and 3 clubs)  
 So, probability of getting a red face card  $= \frac{6}{52} = \frac{3}{26}$

19. (a) an irrational number

**Explanation:**

$$\begin{aligned}
&\frac{\sqrt{5} + \sqrt{2}}{\sqrt{5} - \sqrt{2}} \\
&= \frac{\sqrt{5} + \sqrt{2}}{\sqrt{5} - \sqrt{2}} \times \frac{\sqrt{5} + \sqrt{2}}{\sqrt{5} + \sqrt{2}} \\
&= \frac{(\sqrt{5} + \sqrt{2})^2}{(\sqrt{5})^2 - (\sqrt{2})^2} \\
&= \frac{(\sqrt{5})^2 + (\sqrt{2})^2 + 2 \times \sqrt{5} \times \sqrt{2}}{5 - 2} \\
&= \frac{5 + 2 + 2\sqrt{10}}{3} \\
&= \frac{7 + 2\sqrt{10}}{3}
\end{aligned}$$

Here  $\sqrt{10} = \sqrt{2} \times \sqrt{5}$

Since  $\sqrt{2}$  and  $\sqrt{5}$  both are an irrational number

Therefore,  $\frac{\sqrt{5} + \sqrt{2}}{\sqrt{5} - \sqrt{2}}$  is an irrational number.

20. (c) 7.5

**Explanation:** Vertices of  $\triangle ABC$  are

A(1, 3), B(-1, 0), C(4, 0)

$$\begin{aligned}
\therefore \text{Area} &= \frac{1}{2} [(x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2))] \\
&= \frac{1}{2} [1(0 - 0) + (-1)(0 - 3) + 4(3 - 0)] \\
&= \frac{1}{2} [0 + 3 + 12] = \frac{15}{2} = 7.5
\end{aligned}$$

### Section B

21. (a) infinitely many solutions

**Explanation:** Given:  $a_1 = 5, a_2 = 3, b_1 = -15, b_2 = -9, c_1 = 8$  and  $c_2 = \frac{24}{5}$  Here

$$\frac{a_1}{a_2} = \frac{5}{3}, \frac{b_1}{b_2} = \frac{-15}{-9} = \frac{5}{3}, \frac{c_1}{c_2} = \frac{8}{\frac{24}{5}} = \frac{5}{3} \therefore \frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$$

Since all have the same answer  $\frac{5}{3}$ .

Therefore, the pair of given linear equations has infinitely many solutions.

22. (b)  $c < 0$

**Explanation:** We are given  $a+b+c < 0$

$$\Rightarrow f(1) < 0$$

So,  $f(x)$  must be negative for all  $x$ .

23. (d) 2520

**Explanation:** Factors of 1 to 10 numbers

$$1 = 1$$

$$2 = 1 \times 2$$

$$3 = 1 \times 3$$

$$4 = 1 \times 2 \times 2$$

$$5 = 1 \times 5$$

$$6 = 1 \times 2 \times 3$$

$$7 = 1 \times 7$$

$$8 = 2 \times 2 \times 2$$

$$9 = 1 \times 3 \times 3$$

$$\Rightarrow \text{LCM of number 1 to 10} = \text{LCM}(1, 2, 3, 4, 5, 6, 7, 8, 9, 10)$$

$$= 1 \times 2 \times 2 \times 2 \times 3 \times 3 \times 5 \times 7 = 2520$$

24. (b)  $\frac{\sqrt{b^2-a^2}}{b}$

**Explanation:**  $\cos^2 \theta = (1 - \sin^2 \theta) = \left(1 - \frac{a^2}{b^2}\right) = \frac{b^2-a^2}{b^2} \Rightarrow \cos \theta = \frac{\sqrt{b^2-a^2}}{b}$

25. (a) 39 and 13

**Explanation:** Let the two numbers be  $x$  and  $y$

According to question,  $x - y = 26$  and  $x = 3y$

Putting the value of  $x$  in  $x - y = 26$ , we get,

$$3y - y = 26$$

$$\Rightarrow y = 13 \text{ And } x = 3 \times 13 = 39$$

Therefore, the two numbers are 13 and 39.

26. (a)  $a = 0, b = -6$

**Explanation:** Zeroes of a polynomial are the values of  $x$  at which the polynomial is equal to zero.

2 and -3 are the zeroes of the polynomial  $p(x) = x^2 + (a + 1)x + b$

i.e.  $p(2) = 0$  and  $p(-3) = 0$

$$p(2) = (2)^2 + (a + 1)(2) + b = 0$$

$$\Rightarrow 4 + 2a + 2 + b = 0$$

$$\Rightarrow 6 + 2a + b = 0 \dots (1)$$

$$p(-3) = (-3)^2 + (a + 1)(-3) + b = 0$$

$$\Rightarrow 9 - 3a - 3 + b = 0$$

$$\Rightarrow 6 - 3a + b = 0 \dots (2)$$

Equating (1) & (2), as both the equations are equal to zero.

$$\therefore 6 + 2a + b = 6 - 3a + b$$

$$\Rightarrow 5a = 0$$

$$\Rightarrow a = 0$$


Putting the value of 'a' in (1)

$$6 + 2(0) + b = 0$$

$$\Rightarrow b = -6$$

27. (b) 10 cm.

**Explanation:** The diameter of circle is AC. Here  $\angle ABC$  is angle of semicircle.



$\therefore \angle ABC = 90^\circ \therefore \Delta ABC$  is a right angled triangle.

$\therefore$  By using Pythagoras theorem,

$$\therefore AC = \sqrt{AB^2 + BC^2} = \sqrt{(8)^2 + (6)^2} \Rightarrow AC = \sqrt{100} = 10 \text{ cm}$$

28. (d) 2

**Explanation:** 2

29. (a)  $\frac{1}{2}$

**Explanation:** We know that  $\sec^2 A - \tan^2 A = 1$ .

$$\therefore (2x)^2 - \left(\frac{2}{x}\right)^2 = 1 \Rightarrow 4x^2 - \frac{4}{x^2} = 1 \Rightarrow 4\left(x^2 - \frac{1}{x^2}\right) = 1$$

$$\Rightarrow \left(x^2 - \frac{1}{x^2}\right) = \frac{1}{4} \Rightarrow 2\left(x^2 - \frac{1}{x^2}\right) = 2 \times \frac{1}{4} = \frac{1}{2}$$

30. (a)  $320 \text{ m}^2$

**Explanation:** Let the width be  $x$   
then length be  $x + 4$

According to the question,

$$l + b = 36$$

$$x + (x + 4) = 36$$

$$2x + 4 = 36$$

$$2x = 36 - 4$$

$$2x = 32$$

$$x = 16.$$

Hence, The length of the garden will be 20 m and width will be 16 m.

$$\text{Area} = \text{length} \times \text{breadth} = 20 \times 16 = 320 \text{ m}^2$$

31. (a)  $x^2 y^2$

**Explanation:**  $x^2 y^5 = y^3 (x^2 y^2)$

$$x^3 y^3 = x(x^2 y^2)$$

Therefore HCF (m, n) is  $x^2 y^2$

32. (d) 4 : 25

**Explanation:** In  $\Delta ABC$  and  $\Delta DEF$

$$\frac{AB}{DE} = \frac{BC}{EF} = \frac{CA}{FD} = \frac{2}{5}$$

$\therefore$  The sides are proportional

$$\therefore \Delta ABC \sim \Delta DEF$$

$$\therefore \frac{\text{area of } \Delta ABC}{\text{area of } \Delta DEF} = \frac{AB^2}{DE^2}$$

$$= \left(\frac{2}{5}\right)^2 = \frac{4}{25}$$

$\therefore$  Ratio = 4 : 25

33. (b)  $\frac{1}{7}$

**Explanation:** Given,  $\tan \theta = \frac{4}{7}$

$$\therefore \frac{(7 \sin \theta - 3 \cos \theta)}{(7 \sin \theta + 3 \cos \theta)} = \frac{(7 \tan \theta - 3)}{(7 \tan \theta + 3)} \quad [\text{Dividing numerator and denom. by } \cos \theta]$$

$$= \frac{\left(7 \times \frac{4}{7} - 3\right)}{\left(7 \times \frac{4}{7} + 3\right)} = \frac{(4 - 3)}{(4 + 3)} = \frac{1}{7}$$

34. (d)  $(-4, 2)$

**Explanation:**  $(x, y) = \left\{ \frac{(-6+(-2))}{2}, \frac{(8+(-4))}{2} \right\}$

$$= \left( \frac{-8}{2}, \frac{4}{2} \right)$$

$$= (-4, 2)$$

35. (d)  $\frac{17}{18}$

**Explanation:** Number of total coins =  $100 + 50 + 20 + 10 = 180$

Number of coins except five rupee coins =  $180 - 10 = 170$

$$\therefore \text{Required Probability} = \frac{170}{180} = \frac{17}{18}$$

36. (c)  $\frac{15}{4}$

**Explanation:** Condition for parallel lines is

$$\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2} \dots (i)$$

Given lines,

$$3x + 2ky - 2 = 0 \text{ and}$$

$$2x + 5y - 1 = 0;$$

Comparing with standard form,

$$\text{Here, } a_1 = 3, b_1 = 2k, c_1 = -2$$

$$\text{and } a_2 = 2, b_2 = 5, c_2 = -1$$

From Eq. (i),

$$\frac{3}{2} = \frac{2k}{5}$$

$$k = \frac{15}{4}$$

37. (a) 3.141141114...

**Explanation:** 3.141141114 is an irrational number because it is a non-repeating and non-terminating decimal.

38. (d)  $a^2b^2$

**Explanation:**  $x = a \cos \theta, y = b \sin \theta$

$$bx = ab \cos \theta \dots (i)$$

$$ay = ab \sin \theta \dots (ii)$$

Squaring and adding (i) and (ii) we get,

$$b^2x^2 + a^2y^2 = a^2b^2 \cos^2 \theta + a^2b^2 \sin^2 \theta$$

$$= a^2b^2 (\cos^2 \theta + \sin^2 \theta)$$

$$= a^2b^2 \times 1$$

$$= a^2b^2$$

39. (a)  $\frac{1}{2}$

**Explanation:** Total number of outcomes =  $\{1, 2, 3, 4, 5, 6\}$

So, total outcomes = 6

Favourable outcomes in this case =  $\{2, 4, 6\}$

So, number of favourable outcomes = 3

$$\therefore P(\text{an even number}) = \frac{\text{Favourable outcomes}}{\text{Total outcomes}} = \frac{3}{6} = \frac{1}{2}$$

40. (d) IV

**Explanation:** The point p is given by  $P \left( \frac{2 \times 5 + 3 \times 2}{2+3}, \frac{2 \times 2 - 3 \times 5}{2+3} \right) = P \left( 3, \frac{-11}{5} \right)$


so, p lies in IV quadrant.

$(-, +)$	$(+\infty)$
	if
	if
$(-, -)$	$(+, -)$

### Section C

41. (d) AA

**Explanation:** Let BC represents the height of bus and DE represents the height of building.



In  $\triangle ABC$  and  $\triangle ADE$ ,


$\angle A = \angle A$  (Common)

$\angle B = \angle D$  (Corresponding angles)

$\therefore \triangle ABC \sim \triangle ADE$  (By AA similarity criteria)

42. (b) 12.5 m

**Explanation:** Let BC represents the height of bus and DE represents the height of building.




We have,  $AB = 2BC$

$$\Rightarrow BC = \frac{25}{2} = 12.5 \text{ m}$$

So, height of bus = 12.5 m

43. (a) 1 : 6

**Explanation:** Let BC represents the height of bus and DE represents the height of building.



We have,  $AD = 12 BC$

$$\Rightarrow AD = 12 \times 12.5 = 150 \text{ m}$$


$\therefore \triangle ABC \sim \triangle ADE$

$$\therefore \frac{AB}{AD} = \frac{BC}{DE} \Rightarrow \frac{BC}{DE} = \frac{25}{150} = \frac{1}{6}$$

So, ratio of heights of bus and building is 1 : 6.

44. (d) 1 : 5

**Explanation:** Let BC represents the height of bus and DE represents the height of building.



Since  $\triangle ABC \sim \triangle ADE$


$$\Rightarrow \frac{AB}{AD} = \frac{AC}{AE} \Rightarrow \frac{AC}{AE} = \frac{1}{6}$$

$$\Rightarrow \frac{AC}{AE-AC} = \frac{1}{6-1} \Rightarrow \frac{AC}{EC} = \frac{1}{5}$$

$\therefore$  Required ratio = 1 : 5

45. (c) 75 m

**Explanation:** Let BC represents the height of bus and DE represents the height of building.



Height of the building = DE

$$\text{Now, } \frac{BC}{DE} = \frac{1}{6}$$

$$\Rightarrow DE = 6BC = 6 \times 12.5 = 75 \text{ m}$$

46. (b) 50.28 cm<sup>2</sup>

**Explanation:** Let r and R be the radii of each smaller circle and larger circle respectively.

We have,  $d = \frac{1}{4}D$

$$\Rightarrow r = \frac{1}{4}R \Rightarrow r = \frac{1}{4} \times 16 \Rightarrow r = 4 \text{ cm}$$

Area of smaller circle =  $\pi r^2$

$$= \frac{22}{7} \times 4 \times 4 = 50.28 \text{ cm}^2$$

47. (b) 804.57 cm<sup>2</sup>

**Explanation:** Let r and R be the radii of each smaller circle and larger circle respectively.

We have,  $d = \frac{1}{4}D$

$$\Rightarrow r = \frac{1}{4}R \Rightarrow r = \frac{1}{4} \times 16 \Rightarrow r = 4 \text{ cm}$$

Area of larger circle =  $\pi R^2$

$$= \frac{22}{7} \times 16 \times 16 = \frac{5632}{7} = 804.57 \text{ cm}^2$$

48. (b) 603.45 cm<sup>2</sup>

**Explanation:** Let r and R be the radii of each smaller circle and larger circle respectively.

We have,  $d = \frac{1}{4}D$

$$\Rightarrow r = \frac{1}{4}R \Rightarrow r = \frac{1}{4} \times 16 \Rightarrow r = 4 \text{ cm}$$

Area of the black colour region = Area of larger circle - Area of 4 smaller circles

$$= 804.57 - 4 \times 50.28 = 603.45 \text{ cm}^2$$

49. (b) 12.57 cm<sup>2</sup>

**Explanation:** Let r and R be the radii of each smaller circle and larger circle respectively.

We have,  $d = \frac{1}{4}D$

$$\Rightarrow r = \frac{1}{4}R \Rightarrow r = \frac{1}{4} \times 16 \Rightarrow r = 4 \text{ cm}$$

Area of quadrant of a smaller circle

$$= \frac{1}{4} \times 50.28 = 12.57 \text{ cm}^2$$

50. (a) 66 cm<sup>2</sup>

**Explanation:** Let r and R be the radii of each smaller circle and larger circle respectively.

We have,  $d = \frac{1}{4}D$

$$\Rightarrow r = \frac{1}{4}R \Rightarrow r = \frac{1}{4} \times 16 \Rightarrow r = 4 \text{ cm}$$

Area between two concentric circles

$$= \pi(R^2 - r^2) = \frac{22}{7}(5^2 - 2^2)$$

$$= \frac{22}{7}(25 - 4) = \frac{22}{7} \times 21 = 66 \text{ cm}^2$$

