

Section C

Attempt any 8 questions

Question No. 41 to 45 are based on the given text. Read the text carefully and answer the questions:

While playing badminton Ronit seeing the barrier chains hung between two posts at the edge of the walk way of a street. It is hung in the shape of the parabola. Parabola is the graphical representation of a particular type of polynomial.



41. Which of the following polynomial is graphically represented by a parabola? [0.71]
a) Cubic polynomial b) Linear polynomial
c) None of these d) Quadratic polynomial
42. If a polynomial, represented by a parabola, intersects the x-axis at -3, 4 and y-axis at -2, [0.71]
then its zero(es) is/are
a) -3 and 3 b) -1, 2 and -2
c) 2 and -2 d) -1
43. If the barrier chains between two posts is represented by the polynomial $x^2 - x - 12$, then its [0.71]
zeroes are
a) 4, -5 b) 4, -3
c) -2, 5 d) 4, 3
44. The sum of zeroes of the polynomial $4x^2 - 9x + 2$ is [0.71]
a) $\frac{1}{4}$ b) $-\frac{9}{4}$
c) $\frac{2}{4}$ d) $\frac{9}{4}$
45. The reciprocal of the product of zeroes of the polynomial $x^2 - 9x + 20$ is [0.71]
a) 20 b) $\frac{1}{8}$
c) 5 d) $\frac{1}{20}$

Question No. 46 to 50 are based on the given text. Read the text carefully and answer the questions:

In an examination hall, students are seated at a distance of 2 due to the CORONA virus pandemic. Let three students sit at (7, 3) and (8, 5) respectively.



46. The distance between A and C is [1]
- i. $\sqrt{5}$ units
 - ii. $4\sqrt{5}$ units
 - iii. $3\sqrt{5}$ units
 - iv. none of the above
- a) Option (iv) b) Option (i)
c) Option (iii) d) Option (ii)
47. If an invigilator at point 7, lying on the straight line joining B and C such that it divides the distance between them in the ratio of 1 : 2. Then coordinates of I are [1]
- a) $(\frac{22}{3}, \frac{11}{3})$ b) (6, 1)
c) $(\frac{23}{3}, \frac{13}{3})$ d) (9, 1)
48. The mid-point of the line segment joining A and C is [1]
- a) $(\frac{11}{2}, 0)$ b) none of the above
c) (6, 1) d) (1, 6)
49. The ratio in which B divides the line segment joining A and C is [1]
- a) 3 : 1 b) none of these
c) 2 : 1 d) 1 : 2
50. The points A, B and C lie on [1]
- a) a straight line b) a scalene triangle
c) an equilateral triangle d) an isosceles triangle

Solution

Section A

1. (a) $\frac{17}{6}$

Explanation: $\frac{17}{6}$ has a non-terminal repeating decimal expansion.

$$\frac{17}{6} = 2.6333\dots$$

2. (d) 6

Explanation: For non-zero solution $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2} \Rightarrow \frac{3}{k} = \frac{5}{10} = \frac{0}{0}$

Taking, $\frac{3}{k} = \frac{5}{10} \Rightarrow k = \frac{3 \times 10}{5} = 6$

3. (d) 24 cm

Explanation: Since in isosceles triangle ABC, the altitudes bisect the opposite side. (The altitude and median coincide)

$$\therefore BD = \frac{BC}{2} = \frac{14}{2} = 7 \text{ cm Now, in } \triangle ABD,$$

$$AD = \sqrt{AB^2 - BD^2}$$

$$= \sqrt{(25)^2 - (7)^2}$$

$$= \sqrt{625 - 49}$$

$$= \sqrt{576} = 24 \text{ cm}$$

4. (a) $x = 3, y = 4$

Explanation: Divide throughout by xy and put $\frac{1}{x} = u$ and $\frac{1}{y} = v$ to get

$$4v + 6u = 3 \dots\dots(i)$$

$$\text{and } 8v + 9u = 5 \dots\dots(ii)$$

This gives $u = \frac{1}{3}$ and $v = \frac{1}{4}$. Hence, $x = 3$ and $y = 4$.

5. (d) $\frac{3}{160}$

Explanation: $\cos \theta = \frac{3}{5} = \frac{\text{Base}}{\text{Hypotenuse}}$

By Pythagoras Theorem, $(\text{Hypotenuse})^2 = (\text{Base})^2 + (\text{Alt.})^2$

$$\Rightarrow (5)^2 = (3)^2 + (\text{alt.})^2$$

$$\Rightarrow 25 = 9 + (\text{alt})^2 \Rightarrow (\text{alt})^2 = 25 - 9 = 16 = (4)^2$$

$$\text{Alt.} = 4$$

$$\text{Now, } \sin \theta = \frac{\text{Alt.}}{\text{Hypotenuse}} = \frac{4}{5}$$

$$\text{and } \tan \theta = \frac{\text{Alt.}}{\text{Base}} = \frac{4}{3}$$

$$\therefore \frac{\sin \theta \tan \theta - 1}{2 \tan^2 \theta} = \frac{\frac{4}{5} \times \frac{4}{3} - 1}{2 \times \left(\frac{4}{3}\right)^2} = \frac{\frac{16}{15} - 1}{2 \times \frac{16}{9}}$$

$$= \frac{\frac{1}{15}}{\frac{32}{9}} = \frac{1}{15} \times \frac{9}{32} = \frac{3}{160}$$

6. (a) 1

Explanation: We know that 6^n will end in 6

And 5^n will end in 5

Now, $6^n - 5^n$ always end with $6 - 5 = 1$

7. (a) both negative

Explanation: Given; $x^2 + 88x + 125 = 0$

$$D = (88)^2 - 4(1)(125)$$

$$D = 7244$$

Now,

$$x = \frac{-(88) \pm \sqrt{7244}}{2(1)}$$

$$\Rightarrow x = \frac{-88 \pm 2\sqrt{1811}}{2}$$

There roots are $x = -44 + \sqrt{1811}, -44 - \sqrt{1811}$

Which are both negative.

8. **(c)** 231 cm^2

Explanation: Area swept by minute hand in 60 minutes = πR^2

Area swept by it in 10 minutes

$$= \left(\frac{\pi R^2}{60} \times 10 \right) \text{ cm}^2 = \left(\frac{22}{7} \times 21 \times 21 \times \frac{1}{6} \right) \text{ cm}^2$$

$$= 231 \text{ cm}^2$$

9. **(a)** $5x^3 - 3x^2 - \sqrt{x} + 2$

Explanation: $5x^3 - 3x^2 - \sqrt{x} + 2$ is not a polynomial because each term of a polynomial should be a product of a constant and one or more variable raised to a positive, zero or integral power. Here \sqrt{x} does not satisfy the condition of being a polynomial.

10. **(d)** 25 : 49

Explanation: In $\triangle ABC$ and $\triangle DEF$,

$$\frac{AB}{DE} = \frac{BC}{EF} = \frac{AC}{DF} = \frac{5}{7}$$

Then, area ($\triangle ABC$) : area ($\triangle DEF$)

$$= \frac{AB^2}{DE^2} = \left(\frac{5}{7} \right)^2 = 25 : 49$$

11. **(c)** 0

Explanation: Elementary events are

(1, 1), (1, 2), (1, 3), (1, 4), (1, 5), (1, 6)

(2, 1), (2, 2), (2, 3), (2, 4), (2, 5), (2, 6)

(3, 1), (3, 2), (3, 3), (3, 4), (3, 5), (3, 6)

(4, 1), (4, 2), (4, 3), (4, 4), (4, 5), (4, 6)

(5, 1), (5, 2), (5, 3), (5, 4), (5, 5), (5, 6)

(6, 1), (6, 2), (6, 3), (6, 4), (6, 5), (6, 6)

\therefore Number of Total outcomes = 36

And Number of possible outcomes (sum of numbers appearing on die is 13) = 0

$$\therefore \text{Required Probability} = \frac{0}{36} = 0$$

12. **(a)** equal

Explanation: If we assume that a and b are equal and consider $a = b = k$

Then,

HCF (a, b) = k

LCM (a, b) = k

13. **(b)** 26

Explanation: Area of first circle of radius = $\frac{10}{2} = 5 \text{ cm}$

$$= \pi r^2 = \pi \times (5)^2 \text{ cm}^2 = 25\pi \text{ cm}^2$$

and area of second circle of radius = $\frac{24}{2} = 12 \text{ cm} = \pi(12)^2 \text{ cm}^2 = 144\pi \text{ cm}^2$

$$\therefore \text{Total area} = (25\pi + 144\pi) \text{ cm}^2 = 169\pi \text{ cm}^2$$

$$\therefore \text{Area of larger circle} = 169\pi \text{ cm}^2$$

$$\therefore \text{Radius} = \sqrt{\frac{\text{Area}}{\pi}} = \sqrt{\frac{169\pi}{\pi}} = \sqrt{169}$$


$$= 13 \text{ cm}$$

$$\therefore \text{Diameter} = 2 \times \text{radius} = 2 \times 13 = 26 \text{ cm}^2$$

14. **(d)** 2240 cm^2

Explanation: A chord AB makes an angle of 90° at the centre

Radius of the circle = 28 cm



Area of minor segment ACB

$$\begin{aligned}
 &= \pi r^2 \times \frac{\theta}{360^\circ} - \text{area of } \triangle AOB \\
 &= \pi r^2 \times \frac{90^\circ}{360^\circ} - \frac{1}{2} OA \times OB \\
 &= \frac{1}{4} \pi r^2 - \frac{1}{2} \times r^2 \\
 &= \frac{1}{4} \times \frac{22}{7} \times 28 \times 28 - \frac{1}{2} \times 28 \times 28 \\
 &= 616 - 392 \\
 &= 224 \text{ cm}^2
 \end{aligned}$$

\therefore Area of the major segment ADB

$$\begin{aligned}
 &= \text{Area of circle} - \text{area of minor segment} \\
 &= \pi r^2 - 224 = \frac{22}{7} \times 28 \times 28 - 224 \\
 &= 2464 - 224 \\
 &= 2240 \text{ sq. cm}
 \end{aligned}$$

15. (a) $AB^2 + AC^2 = BC^2$

Explanation: In triangle ADC, $AC^2 = AD^2 + CD^2$

In triangle ABD, $AB^2 = AD^2 + BD^2$

Adding both equations,

$$AC^2 + AB^2 = 2AD^2 + CD^2 + BD^2$$

$$\Rightarrow AC^2 + AB^2 = 2CD \cdot BD + CD^2 + BD^2 \text{ [Since } BD = CD = AD\text{]}$$

$$\Rightarrow AB^2 + AC^2 = (BD + CD)^2$$

$$\Rightarrow AB^2 + AC^2 = BC^2$$

16. (a) $\tan \theta$

Explanation: Here $\sqrt{(1 - \cos^2 \theta) \sec^2 \theta}$

$$= \sqrt{\sin^2 \theta \times \frac{1}{\cos^2 \theta}}$$

$$[\because 1 - \cos^2 \theta = \sin^2 \theta \text{ and } \sec^2 \theta = \frac{1}{\cos^2 \theta}]$$

$$= \sqrt{\frac{\sin^2 \theta}{\cos^2 \theta}}$$

$$= \sqrt{\tan^2 \theta}$$

$$= \tan \theta$$

17. (b) 6

Explanation: The given system of equations are

$$2x + 3y = 5$$

$$4x + ky = 10$$

For the equations to have infinite number of solutions, $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$

Here, we must have

$$\text{Therefore } \frac{2}{4} = \frac{3}{k} = \frac{5}{10}$$

$$\Rightarrow \frac{2}{4} = \frac{3}{k}$$

$$\Rightarrow 2k = 12$$

$$\Rightarrow k = \frac{12}{2}$$

$$\Rightarrow k = 6$$

18. (c) 0

Explanation: Elementary events are

(1, 1), (1, 2), (1, 3), (1, 4), (1, 5), (1, 6)

(2, 1), (2, 2), (2, 3), (2, 4), (2, 5), (2, 6)
 (3, 1), (3, 2), (3, 3), (3, 4), (3, 5), (3, 6)
 (4, 1), (4, 2), (4, 3), (4, 4), (4, 5), (4, 6)
 (5, 1), (5, 2), (5, 3), (5, 4), (5, 5), (5, 6)
 (6, 1), (6, 2), (6, 3), (6, 4), (6, 5), (6, 6)

∴ Number of Total outcomes = 36

And Number of possible outcomes (product of numbers appearing on die is 7) = 0

∴ Required Probability = $\frac{0}{36} = 0$

19. (d) composite number

Explanation: We have $7 \times 11 \times 13 + 13 = 13(77 + 1) = 13 \times 78$. Since the given number has 2 more factors other than 1 and itself, therefore it is a composite number.

20. (c) 228 cm^2

Explanation: Correct option: (b)

Side of a square = 20cm

Area of the square = $(20 \times 20) \text{ cm}^2 = 400 \text{ cm}^2$

Diagonal of square = $\sqrt{(20)^2 + (20)^2} = \sqrt{800} = 20\sqrt{2} \text{ cm}$

= Radius of the quadrant = $20\sqrt{2} \text{ cm}$

Area of a quadrant = $\frac{1}{4} \times 3.14 \times (20\sqrt{2})^2 = 628 \text{ cm}^2$

Thus, area of the shaded region = Area of a quadrant - Area of the square

= $(628 - 400) \text{ cm}^2$

= 228 cm^2

Section B


21. (b) 18 sq. units

Explanation: Here are the two solutions of each of the given equations. $x + 3y = 6$

x	0	3	6
y	2	1	0

$2x - 3y = 12$


x	0	3	6
y	-4	-2	0



∴ Area of triangle = $\frac{1}{2} \times \text{Base} \times \text{Height} = \frac{1}{2} \times 6 \times 6 = 18 \text{ sq. units}$

22. (a) 30°

Explanation:



$$\frac{AB}{AC} = \frac{BD}{DC}$$

$$\angle B = 70^\circ \quad \angle C = 50^\circ$$

But $\angle A + \angle B + \angle C = 180^\circ$ (Angles of a triangle)

$$\angle A = 180^\circ - (\angle B + \angle C)$$

$$= 180^\circ - (70^\circ + 50^\circ)$$

$$= 180^\circ - 120^\circ = 60^\circ$$

$$\frac{AB}{AC} = \frac{BD}{DC}$$


AD is the bisector of $\angle A$

$$\angle BAD = \frac{60}{2} = 30^\circ$$

23. (c) 4

Explanation:

Using the factor tree for prime factorisation, we have:



Therefore,

$$196 = 2 \times 2 \times 7 \times 7$$

$$196 = 2^2 \times 7^2$$

The exponents of 2 and 7 are 2 and 2 respectively.

Thus the sum of the exponents is 4.

24. (a) $\frac{m^2-1}{n^2-1}$

Explanation: Given: $\tan A = n \tan B$

$$\Rightarrow \frac{1}{\tan B} = \frac{n}{\tan A}$$

$$\Rightarrow \cot B = \frac{n}{\tan A}$$

And $\sin A = m \sin B$

$$\Rightarrow \frac{1}{\sin B} = \frac{m}{\sin A}$$

$$\Rightarrow \operatorname{cosec} B = \frac{n}{\sin A}$$

Now, $\operatorname{cosec}^2 B - \cot^2 B = 1$

$$\Rightarrow \frac{m^2}{\sin^2 A} - \frac{n^2}{\tan^2 A} = 1$$

$$\Rightarrow \frac{m^2}{\sin^2 A} - \frac{n^2 \cos^2 A}{\sin^2 A} = 1$$

$$\Rightarrow m^2 - n^2 \cos^2 A = \sin^2 A$$

$$\Rightarrow m^2 - n^2 \cos^2 A = 1 - \cos^2 A$$

$$\Rightarrow m^2 - 1 = (n^2 - 1) \cos^2 A$$

$$\Rightarrow \cos^2 A = \frac{m^2 - 1}{n^2 - 1}$$

25. (a) 6


Explanation: Given: $a_1 = 6, a_2 = 12, b_1 = 3, b_2 = c, c_1 = c - 3, c_2 = c$

Since the pair of given linear equations has infinitely many solutions,



$$\begin{aligned} \therefore \frac{a_1}{a_2} &= \frac{b_1}{b_2} = \frac{c_1}{c_2} \\ \Rightarrow \frac{6}{12} &= \frac{3}{c} = \frac{c-3}{c} \\ \text{Taking } \frac{6}{12} &= \frac{3}{c} \\ \Rightarrow c &= \frac{3 \times 12}{6} = 6 \end{aligned}$$

26. (b) 7 m



Let the ladder BC reaches the building at C.

Let the height of building where the ladder reaches be AC.

According to the question:

$$BC = 25 \text{ m}$$

$$AC = 24 \text{ m}$$

In right-angled triangle CAB, we apply Pythagoras theorem to find the value of AB.

$$BC^2 = AC^2 + AB^2$$


$$\Rightarrow AB^2 = BC^2 - AC^2 = 25^2 - 24^2$$

$$\Rightarrow AB^2 = 625 - 576 = 49$$

$$\Rightarrow AB = \sqrt{49} = 7 \text{ m}$$

27. (d) 8.8 cm.

Explanation: Let the two similar triangles be $\triangle ABC$ and $\triangle PQR$ such that $\text{ar}(\triangle ABC) = 121 \text{ cm}^2$ and $\text{ar}(\triangle PQR) = 64 \text{ cm}^2$. Let AM and PN be the respective medians of $\triangle ABC$ and $\triangle PQR$. Since the ratio of the areas of two similar triangles is equal to the ratio of the squares of their corresponding medians



$$\therefore \frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle PQR)} = \frac{AM^2}{PN^2} \Rightarrow \frac{121}{64} = \frac{(12.1)^2}{PN^2}$$

$$\Rightarrow PN^2 = \frac{146.41 \times 64}{121}$$

$$\Rightarrow PN = 8.8 \text{ cm}$$

28. (b) $-y_1 : y_2$

Explanation: Let a point A on x-axis divides the line segment joining the points $P(x_1, y_1)$ $Q(x_2, y_2)$ in the ratio $m_1 : m_2$ and

let co-ordinates of A be $(x, 0)$

$$\therefore 0 = \frac{m_1 y_2 + m_2 y_1}{m_1 + m_2} \Rightarrow 0 = m_1 y_2 + m_2 y_1$$

$$\Rightarrow m_1 y_2 = -m_2 y_1 \Rightarrow \frac{m_1}{m_2} = \frac{-y_1}{y_2}$$

\therefore Ratio is $-y_1 : y_2$

29. (a) $\frac{1}{2}$

Explanation: $\tan^2 45^\circ - \cos^2 30^\circ = x \sin 45^\circ \cos 45^\circ$

$$\Rightarrow (1)^2 - \left(\frac{\sqrt{3}}{2}\right)^2 = x \frac{1}{\sqrt{2}} \times \frac{1}{\sqrt{2}} \Rightarrow 1 - \frac{3}{4} = x \times \frac{1}{2}$$

$$\Rightarrow \frac{1}{4} = \frac{1}{2}x \Rightarrow x = \frac{1}{4} \times \frac{2}{1} = \frac{1}{2}$$

30. (b) 45 years

Explanation: Let my age and son's age be x and y years.

Given, $x = 3y$

$$x + 5 = \frac{5(y+5)}{2}$$

$$\Rightarrow 3y + 5 = \frac{5(y+5)}{2}$$

$$\Rightarrow 6y + 10 = 5y + 25$$

$$\Rightarrow y = 15$$

$$x = 3 \times 15 = 45$$

Hence, my age and son's age are 45 years and 15 years.

31. (b) irrational numbers

Explanation: All non-terminating and non-recurring decimal numbers are irrational numbers. A number is rational if and only if its decimal representation is repeating or terminating.


32. (b) other than 2 or 5 only

Explanation: A rational number can be expressed as a **non-terminating** repeating decimal if the denominator has the factors other than 2 or 5 only.

33. (b) $\frac{1-\cos\theta}{1+\cos\theta}$

Explanation: $(\operatorname{cosec}\theta - \cot\theta)^2 = \left(\frac{1}{\sin\theta} - \frac{\cos\theta}{\sin\theta}\right)^2 = \frac{(1-\cos\theta)^2}{\sin^2\theta} = \frac{(1-\cos\theta)^2}{(1-\cos^2\theta)} = \frac{(1-\cos\theta)}{(1+\cos\theta)}$

34. (b) $\frac{1}{4}\pi a^2$ sq units



Explanation:

According to the question,

Diameter of circle = side of a square

$$\Rightarrow d = a$$

$$\Rightarrow r = \frac{a}{2}$$

$$\text{Now, Area of the circle} = \pi r^2 = \pi \left(\frac{a}{2}\right)^2$$

$$\Rightarrow \text{Area of the circle} = \frac{1}{4}\pi a^2 \text{ sq. units}$$

35. (b) $\frac{3}{5}$

Explanation: Total number of balls in the bag = $4 + 6 = 10$.

Number of black balls = 6.

$$\therefore P(\text{getting a black ball}) = \frac{6}{10} = \frac{3}{5}$$

36. (d) 25 and 25

Explanation: Let number of Rs 1 coins = x

and number of Rs 2 coins = y

Now, by given conditions:

$$\text{Total number of coins} = x + y = 50 \dots(i)$$

Also, Amount of money with her = (Number of Rs 1 \times 1) + (Number of Rs 2 \times coin 2)

$$= x(1) + y(2) = 75$$

$$= x + 2y = 75 \dots(ii)$$

On subtracting Eq. (i) from Eq. (ii), we get

$$(x + 2y) - (x + y) = (75 - 50)$$

$$\text{So, } y = 25$$



Putting $y = 25$ we get $x = 25$.

Hence he has 25 one-rupee coins and 25 2-rupee coins.

37. (a) is non-terminating and non-recurring

Explanation: The value of $\pi = 3.141592653589\dots$

Therefore the value of π is not-repeating decimal, non-terminating and non-recurring numbers.

38. (b) 1

Explanation: Given: $\frac{1+\tan^2\theta}{\sec^2\theta}$

$$= \frac{\sec^2\theta}{\sec^2\theta} = 1$$

$$[\because \sec^2\theta = 1 + \tan^2\theta]$$

39. (b) $\frac{3}{10}$

Explanation: Total numbers are $\sum x_i = 10$

x	f
3	1
5	2
7	3
9	4

$$\text{Average} = \frac{3 \times 1 + 5 \times 2 + 7 \times 3 + 9 \times 4}{10}$$
$$= \frac{3 + 10 + 21 + 36}{10} = \frac{70}{10} = 7$$

$$\therefore m = 3$$

$$\therefore \text{Probability of average number} = \frac{3}{10}$$

40. (b) $\sqrt{2}$

Explanation: Distance between $(\cos\theta, \sin\theta)$ and $(\sin\theta, -\cos\theta)$

$$= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$
$$= \sqrt{(-\cos\theta - \sin\theta)^2 + (\sin\theta - \cos\theta)^2}$$
$$= \sqrt{1 + 1} = \sqrt{2} \{ \because \sin^2\theta + \cos^2\theta = 1 \}$$

Section C

41. (d) Quadratic polynomial

Explanation: Quadratic polynomial

42. (a) -3 and 3

Explanation: Since, the parabola intersects the x-axis at $x = -3$ and $x = 4$. So, zeroes of the polynomial are -3 and 4.

43. (b) 4, -3

Explanation: Let $f(x) = x^2 - x - 12$

$$= x^2 - 4x + 3x - 12 = (x + 3)(x - 4)$$

$$\text{Consider } f(x) = 0 \Rightarrow (x + 3)(x - 4) = 0 \Rightarrow x = 4, -3$$

44. (d) $\frac{9}{4}$

Explanation: Sum of zeroes = $-\frac{\text{Coefficient of } x}{\text{Coefficient of } x^2}$

$$= -\frac{(-9)}{4} = \frac{9}{4}$$

45. (d) $\frac{1}{20}$

Explanation: Product of zeroes = $\frac{20}{1} = 20$

$$\therefore \text{Reciprocal of product of zeroes} = \frac{1}{20}$$

46. (d) Option (ii)

Explanation: The distance between A and C




$$= \sqrt{(8-4)^2 + (5+3)^2} = \sqrt{4^2 + 8^2}$$

$$= \sqrt{16 + 64} = \sqrt{80} = 4\sqrt{5} \text{ units}$$

47. (a) $\left(\frac{22}{3}, \frac{11}{3}\right)$

Explanation: Let the coordinates of I be (x, y)



Then, by section formula,

$$x = \frac{1 \times 8 + 2 \times 7}{1+2} = \frac{8+14}{3} = \frac{22}{3}$$

$$\text{and } y = \frac{1 \times 5 + 2 \times 3}{1+2} = \frac{5+6}{3} = \frac{11}{3}$$

Thus, the coordinates of I is $\left(\frac{22}{3}, \frac{11}{3}\right)$

48. (c) (6, 1)

Explanation: The mid-point of A and C

$$= \left(\frac{8+4}{2}, \frac{5-3}{2}\right) = (6, 1)$$

49. (a) 3 : 1

Explanation: Let B divides the line segment joining A and C in the ratio k : 1. Then, the coordinates of B

will be $\left(\frac{8k+4}{k+1}, \frac{5k-3}{k+1}\right)$.

Thus, we have $\left(\frac{8k+4}{k+1}, \frac{5k-3}{k+1}\right) = (7, 3)$

$$\Rightarrow \frac{8k+4}{k+1} = 7 \text{ and } \frac{5k-3}{k+1} = 3$$

$$\text{Consider, } \frac{8k+4}{k+1} = 7 \Rightarrow 8k + 4 = 7k + 7 \Rightarrow k = 3$$

Hence, the required ratio is 3 : 1.

50. (a) a straight line

Explanation: \because B divides AC in the ratio 3 : 4.

\therefore A, B, C lie on a straight line.

