

Sample Question Paper - 34
Science (086)
Class- X, Session: 2021-22
TERM II

Time allowed : 2 hours

Maximum marks : 40

General Instructions :


- (i) All questions are compulsory.
- (ii) The question paper has three sections and 15 questions. All questions are compulsory.
- (iii) Section-A has 7 questions of 2 marks each; Section-B has 6 questions of 3 marks each; and Section-C has 2 case based questions of 4 marks each.
- (iv) Internal choices have been provided in some questions. A student has to attempt only one of the alternatives in such questions.

SECTION - A

- 1. (a) Name three alkali metals which form a Döbereiner's triad.
(b) Which among F, Cl, Br and I does not fit in the Döbereiner's triad?
- 2. State the reason why carbon can neither form C^{4+} cations nor C^{4-} anions, but forms covalent bonds.
- 3. Fusion of nucleus of sperm and nucleus of ovum marks the completion of fertilisation resulting in the formation of single-celled zygote. What happens when the egg is not fertilized in human females?
- 4. State two differences between budding in *Hydra* and budding in yeast.
- 5. If a trait A exists in 10% of a population of an asexually reproducing species and a trait B exists in 60% of the same population, which trait is likely to have arisen earlier?

OR

- (a) Name the information source for synthesizing protein in a cell and define it.
- (b) What is gene locus?
- 6. Three vertical conducting wires have the same amount of current flowing through them in the direction shown.




Given that the distance between the wires are the same, what is the direction of the resultant magnetic force acting on the middle wire?

OR

Four plotting compasses are placed near a bar magnet. You may ignore any effect of the Earth's magnetic field.

One compass appears approximately like this .



What is the possible position for the compass?

7. What will happen if all the deers are removed in the given food chain?
Plants → Deer → Tigers

OR


Producers always occupy the first trophic level in any food chain. Why?

SECTION - B


8. An element X (2, 8, 2) combines separately with NO_3^- , SO_4^{2-} and PO_4^{3-} ions. Write the formulae of the three compounds so formed. To which group of the periodic table does the element 'X' belong? Will it form covalent or ionic compound? Why?
9. (i) Define allotropes.
(ii) An allotrope of carbon has a molecular mass of 720. Discuss its structure.

OR

- (i) Two non-metals A and B combine with each other by sharing of electrons to form compound C.
(a) What is the nature of compound C?
(b) Will it dissolve in water or organic solvents?
(c) Will it be a good or bad conductor of electricity?
(d) Will it have high or low melting / boiling point?
(ii) Draw the Lewis dot structure of N_2 molecule.
10. List three distinguishing features between monohybrid and dihybrid cross.
11. Figure shows a battery of 12 V and internal resistance 0.6Ω connected to three resistors A, B and C. Find the current in each resistor.



12. (a) How does a solenoid behave like a bar magnet? Can you determine the north and south poles of a current carrying solenoid with the help of a bar magnet? Explain.
(b) A thick copper wire carries current due north placed parallel to and over a compass needle as shown in figure.




If the direction of current is changed from north to south, will the direction of compass needle changes?

OR

- (a) What happens when a current is passed through a solenoid?
- (b) What is the advantage of a solenoid over an ordinary coil?

13. Given pie-chart represents contribution of different gases to greenhouse effect.



- (a) Identify A, B, C and D in the given pie chart.
- (b) Explain briefly labelled part C.
- (c) Name the major sources of labelled part B.

SECTION - C

This section has 02 case-based questions (14 and 15). Each case is followed by 03 sub-questions (a, b and c). Parts a and b are compulsory. However, an internal choice has been provided in part c.

14. Gregor Johannes Mendel worked on garden pea plant (*Pisum sativum*) over a period of eight years and contributed in the field of inheritance and expression of such traits.
- (a) Why did Mendel chose garden pea for his experiments? Write two reasons.
 - (b) List two contrasting visible characters of garden pea that Mendel used for his experiment.
 - (c) Explain in brief how Mendel interpreted his results to show that the traits may be dominant or recessive.

OR

What will happen if both the characters present in F_1 generation pass together in F_2 generation?

15. The relationship between potential difference and current was first established by George Simon Ohm called Ohm's law. According to this law, the current through a metallic conductor is proportional to the potential difference applied between its ends, provided the temperature remain constant *i.e.* $I \propto V$ or $V = IR$; where R is constant for the conductor and it is called resistance of the conductor. Although Ohm's law has been found valid over a large class of materials, there do exist materials and devices used in electric circuits where the proportionality of V and I does not hold.
- (a) If both the potential difference and the resistance in a circuit are doubled, then what would be the effect on current?
 - (b) For a conductor obeying Ohm's law, draw the graph between V and I .
 - (c) When battery of 9 V is connected across a conductor and the current flows is 0.1 A, find the resistance. What would be the effect of increasing temperature on resistance?

OR

- (i) Out of 60 W and 40 W lamps, which one has higher electrical resistance when in use.
- (ii) What is the commercial unit of electric energy? Convert it into joules.

Solution

SCIENCE - 086

Class 10 - Science

1. (a) Li, Na and K form Dobereiner's triad as the atomic mass of sodium is roughly the average of the atomic masses of lithium and potassium.

(b) Cl, Br and I form Dobereiner's triad as the atomic mass of bromine is roughly the average of the atomic masses of chlorine and iodine. F does not fit in the Dobereiner's triad.

2. Ionic compounds are formed either by gaining or losing electrons from the outermost shells, but carbon which has four electrons in its outermost shell cannot form ionic bonds because

– If carbon forms ionic bonds by gaining four electrons to attain a noble gas configuration then it would be difficult for six protons in the nucleus to hold ten electrons.

– If carbon forms ionic bonds by loss of four electrons

then it would require a lot of energy to remove these electrons from outermost shell.

Due to these reasons, carbon forms covalent bonds by sharing the valence electrons.

Type of bonds formed in ionic compounds are called electrovalent bonds and the type of bonds formed in carbon compounds are called covalent bonds.

3. One egg is produced every month by one of the ovaries. The uterus prepares itself every month to receive and nurture the growing embryo. The lining thickens and is richly supplied with blood to nourish the growing embryo. If the egg is not fertilized, this lining is not needed any longer. So, the lining slowly breaks and comes out through the vagina as blood and mucus. This cycle takes place roughly every month and is known as menstruation which usually lasts for about two to eight days.

4.

	Budding in Hydra	Budding in Yeast
(i)	Buds produced are multicellular.	Buds produced are unicellular.
(ii)	Buds get detached from parent body soon.	Buds may remain attached to the parent body


5. In asexually reproducing organisms, there would be very minor differences between two generations due to small inaccuracies in DNA copying or due to some environmental factors or mutation. During reproduction, variations which occur in the first generation is inherited to the subsequent, thus is more frequent. Since, trait B occurs in more individuals, thus it is likely to have arisen earlier.

OR

(a) DNA in the nucleus of a cell is the information source for making proteins in the cell. DNA is the carrier of genetic information and it transmits the heredity from parents to next generation.

(b) Gene locus is the portion or region on chromosome representing a single gene. The alleles of a gene are present on the same gene locus on the homologous chromosomes.


6. By using right-hand rule on wire 1 and wire 2, it can be seen that the interaction between the two fields create opposite signs. This implies that there is an attractive force caused by the interaction of the fields.



The interaction between wire 2 and 3 is also similar as their current directions are the same as wire 1 and wire 2. This implies that wire 2 will be attracted to both wire 1 and wire 3. These two forces will have zero resultant force as the current flowing through three wires is of the same magnitude.

OR

Orientation of compass is as shown below.




7. In the given food chain, if the deers are removed, the population of plants will increase as deers are herbivorous and feed on plants whereas the population of tiger (that consume deer) will decrease, as food available for tiger would be less and it would lead to imbalance in the ecosystem.

OR

The first trophic level in a food chain is occupied by a producer, because these organisms can produce food with the help of sunlight and chlorophyll by a process called photosynthesis.

8. The element X (2, 8, 2) has two electrons in the valence shell, therefore, its valency is 2. The valencies of NO_3^- , SO_4^{2-} and PO_4^{3-} ions are 1, 2 and 3 respectively, therefore, the formulae of its compounds with NO_3^- , SO_4^{2-} and PO_4^{3-} ions are :




Element 'X' belongs to group 2 of the periodic table. Since element 'X' has two electrons in the valence shell, therefore, it can easily lose these two electrons to acquire the stable electronic configuration of the nearest noble gas (*i.e.* neon) to form X^{2+} ions. This ion then combines with NO_3^- , SO_4^{2-} and PO_4^{3-} ions to form ionic compounds.

9. (i) The different forms of an element having different physical properties and structures even though their chemical properties are the same are defined as allotropes.

(ii) Atomic mass of C = 12 and molecular mass of an allotrope of carbon = 720.


$$\therefore \text{No. of C atoms} = 720/12 = 60.$$

Thus, the allotrope of carbon is C_{60} , *i.e.*, Buckminster fullerene and its structure is similar to a soccer ball.



OR

- (i) (a) Covalent
- (b) Usually soluble in organic solvents
- (c) Bad conductor
- (d) Usually low melting / boiling point.



Lewis dot structure
of N_2 molecule

10. Differences between monohybrid and dihybrid cross are as follows:


	Monohybrid Cross	Dihybrid Cross
1.	It is study of the inheritance of a single pair of alleles.	It is the study of inheritance of two pair of alleles.
2.	It produces a phenotypic ratio of 3 : 1 in F_2 generation.	It produces a phenotypic ratio of 9 : 3 : 3 : 1 in F_2 generation.
3.	It produces genotypic ratio of 1 : 2 : 1 in F_2 generation.	It produces genotypic ratio of 1 : 2 : 1 : 2 : 4 : 2 : 1 : 2 : 1 in F_2 generation

11. The equivalent resistance of B and C is given by

$$\frac{1}{R} = \frac{1}{4 \Omega} + \frac{1}{6 \Omega} = \frac{6 \Omega + 4 \Omega}{(6 \Omega)(4 \Omega)} = \frac{10}{24 \Omega} = \frac{5}{12 \Omega}$$

or $R = \frac{12}{5} \Omega = 2.4 \Omega$

$$\text{Total resistance in circuit} = 1.0 \Omega + 2.4 \Omega + 0.6 \Omega = 4.0 \Omega$$



$$\text{Current in the circuit, } I = \frac{12 \text{ V}}{4 \Omega} = 3 \text{ A}$$

$$\text{The potential difference across both B and C} = 3.0 \text{ A} \times 2.4 \Omega = 7.2 \text{ V}$$

$$\text{Current through B, } I_1 = \frac{7.2 \text{ V}}{4.0 \Omega} = 1.8 \text{ A}$$

$$\text{Current through C, } I_2 = \frac{7.2 \text{ V}}{6 \Omega} = 1.2 \text{ A}$$

Thus, current through resistor A = 3.0 A

Current through resistor B = 1.8 A

Current through resistor C = 1.2 A

12. (a) When current flows through a solenoid, a magnetic field is set up around the solenoid which is similar to the magnetic field of a bar magnet.

The polarities of its ends depend upon the direction of current flowing through it. In order to determine the polarities of its ends, place it in a brass hook and suspend it with a long thread so that it moves freely. Bring north pole of a bar magnet near one of its ends. In case the solenoid moves towards the bar magnet, that end of the solenoid is a south pole and in case the solenoid moves away from the magnet, that end of the solenoid is its north pole. The polarity of the other end of the solenoid can similarly be determined.

(b) When the direction of current changes, the direction of magnetic field also changes.

OR

(a) When current is passed through a solenoid the magnetic field around is produced almost like a bar magnet.

(b) The advantage of solenoid over an ordinary coil is that inside a solenoid a uniform magnetic field is produced.

13. (a) In the given pie-chart, contribution of different gases to greenhouse effect is, A is CO₂ (60%), B is CH₄ (20%), C is CFCs (14%) and D is N₂O (6%).

(b) C is Chlorofluorocarbons, they (CFCs) are synthetic gaseous compounds of carbon and halogen which are odourless, non-toxic, noninflammable, chemically inert propellants used in aerosol cans and jet fuel, refrigerants in air conditioners and refrigerators.

(c) Flooded paddy fields, marshes and cattle are the major sources of CH₄.

14. (a) Mendel chose garden pea for his experiments because

(i) Pea plant is an annual plant with short life span.
 (ii) The flower structure of pea is such as to allow controlled breeding. Though plant is self pollinated, but it can be cross bred manually.

(b) Two contrasting characters used by Mendel for his experiment are as follows :

(i) Round/Wrinkled seeds
 (ii) Tall/Short plants


(c) When Mendel crossed two pea plants with a pair of contrasting characters, only one character appeared in all the members of F₁ progeny, the other was not expressed.

On selfing F₁, the hidden characters reappeared in just 25% of the offsprings and the other 75% shared the characters expressed in F₁.

Mendel concluded that the character which expresses itself in F₁ and in 75% of the individuals of F₂ is dominating while the other one is recessive.

OR

If both the characters (*i.e.*, dihybrid cross) present in F₁ generation pass together then they will not follow the law of segregation. If this happens in the traits of hybrid generation or F₁ generation (RrYy), then recessive trait will not be able to express its effect in the next generation and it will result in polyploidy.




15. (a) $V = IR$

So, $V' \rightarrow 2V, R' \rightarrow 2R$

$$I' = \frac{2V}{2R} = I$$

Current will remain same.

(b) Since, $V \propto I$. So, the graph is a straight line and passing through origin.



(c) Given : $V = 9\text{ V}, I = 0.1\text{ A}$

$$R = \frac{V}{I} = \frac{9}{0.1} = 90\ \Omega$$

On increasing temperature, resistance of conductor increases.

OR

(i) For the same applied voltage, $P \propto \frac{1}{R}$ or $R \propto \frac{1}{P}$

i.e., less the power of electrical device, higher is its electrical resistance. Therefore, a 40 W lamp has higher electrical resistance when in use.

(ii) Commercial unit of electrical energy : Kilowatt hour

$$1\text{ kW h} = 1000\text{ W h} = 1000 \frac{\text{J}}{\text{s}} \times 3600\text{ s} = 3600000\text{ J} = 3.6 \times 10^6\text{ J}$$