

## Inverse Trigonometric Functions

- If  $\sin y = x$ , then  $y = \sin^{-1}x$  (We read it as sine inverse  $x$ )

Here,  $\sin^{-1}x$  is an inverse trigonometric function. Similarly, the other inverse trigonometric functions are as follows:

- If  $\cos y = x$ , then  $y = \cos^{-1}x$
- If  $\tan y = x$ , then  $y = \tan^{-1}x$
- If  $\cot y = x$ , then  $y = \cot^{-1}x$
- If  $\sec y = x$ , then  $y = \sec^{-1}x$
- If  $\operatorname{cosec} y = x$ , then  $y = \operatorname{cosec}^{-1}x$
- The domains and ranges (principle value branches) of inverse trigonometric functions can be shown in a table as follows:

Function	Domain	Range (Principle value branches)
$y = \sin^{-1}x$	$[-1, 1]$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
$y = \cos^{-1}x$	$[-1, 1]$	$[0, \pi]$
$y = \tan^{-1}x$	$\mathbf{R}$	$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
$y = \cot^{-1}x$	$\mathbf{R}$	$(0, \pi)$
$y = \sec^{-1}x$	$\mathbf{R} - (-1, 1)$	$[0, \pi] - \left\{\frac{\pi}{2}\right\}$
$y = \operatorname{cosec}^{-1}x$	$\mathbf{R} - (-1, 1)$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$

- Note that  $y = \tan^{-1}x$  does not mean that  $y = (\tan x)^{-1}$ . This argument also holds true for the other inverse trigonometric functions.
- The principal value of an inverse trigonometric function can be defined as the value of inverse trigonometric functions, which lies in the range of principal branch.

**Example 1:** What is the principal value of  $\tan^{-1}(-\sqrt{3}) + \sin^{-1}(1)$ ?

**Solution:** Let  $\sin^{-1}(1) = z$

$$\Rightarrow \tan y = -\sqrt{3} = -\tan\left(\frac{\pi}{3}\right) = \tan\left(-\frac{\pi}{3}\right) \text{ and } \sin z = 1 = \sin \frac{\pi}{2}$$








We know that the ranges of principal value branch of  $\tan^{-1}$  and  $\sin^{-1}$  are  $(-\frac{\pi}{2}, \frac{\pi}{2})$  and  $[-\frac{\pi}{2}, \frac{\pi}{2}]$  respectively. Also,  $\tan(-\frac{\pi}{3}) = -\sqrt{3}$  and  $\sin(\frac{\pi}{2}) = 1$

Therefore, principal values of  $\tan^{-1}(-\sqrt{3}) = -\frac{\pi}{3}$  and  $\sin^{-1}(1) = \frac{\pi}{2}$

$$\therefore \tan^{-1}(-\sqrt{3}) + \sin^{-1}(1) = -\frac{\pi}{3} + \frac{\pi}{2} = \frac{\pi}{6}$$

- Graphs of the six inverse trigonometric functions can be drawn as follows:





- The relation  $\sin y = x \Rightarrow y = \sin^{-1} x$  gives  $\sin (\sin^{-1} x) = x$ , where  $x \in [-1, 1]$ ; and  $\sin^{-1}(\sin x) = x$ , where  $x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$

This property can be similarly stated for the other inverse trigonometric functions as follows:

- $\cos(\cos^{-1}x) = x, x \in [-1, 1]$  and  $\cos^{-1}(\cos x) = x, x \in [0, \pi]$
- $\tan(\tan^{-1}x) = x, x \in \mathbf{R}$  and  $\tan^{-1}(\tan x) = x, x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
- $\operatorname{cosec}(\operatorname{cosec}^{-1}x) = x, x \in \mathbf{R} - (-1, 1)$  and  $\operatorname{cosec}^{-1}(\operatorname{cosec} x) = x, x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$
- $\sec(\sec^{-1}x) = x, x \in \mathbf{R} - (-1, 1)$  and  $\sec^{-1}(\sec x) = x, x \in [0, \pi] - \left\{\frac{\pi}{2}\right\}$
- $\cot(\cot^{-1}x) = x, x \in \mathbf{R}$  and  $\cot^{-1}(\cot x) = x, x \in (0, \pi)$

- For suitable values of domains;

$$\sin^{-1}\left(\frac{1}{x}\right) = \operatorname{cosec}^{-1}x, x \in \mathbf{R} - (-1, 1)$$

$$\cos^{-1}\left(\frac{1}{x}\right) = \sec^{-1}x, x \in \mathbf{R} - (-1, 1)$$

$$\tan^{-1}\left(\frac{1}{x}\right) = \begin{cases} \cot^{-1}x, & x > 0 \\ \cot^{-1}x - \pi, & x < 0 \end{cases}$$

$$\operatorname{cosec}^{-1}\left(\frac{1}{x}\right) = \sin^{-1}x, x \in [-1, 1]$$

$$\sec^{-1}\left(\frac{1}{x}\right) = \cos^{-1}x, x \in [-1, 1]$$

$$\cot^{-1}\left(\frac{1}{x}\right) = \begin{cases} \tan^{-1}x, & x > 0 \\ \pi + \tan^{-1}x, & x < 0 \end{cases}$$

**Note:** While solving problems, we generally use the formulas  $\tan^{-1}\left(\frac{1}{x}\right) = \cot^{-1}x$  and  $\cot^{-1}\left(\frac{1}{x}\right) = \tan^{-1}x$  when the conditions for  $x$  (i.e.,  $x > 0$  or  $x < 0$ ) are not given

- For suitable values of domains;

- $\sin^{-1}(-x) = -\sin^{-1}x, x \in [-1, 1]$
- $\cos^{-1}(-x) = \pi - \cos^{-1}x, x \in [-1, 1]$
- $\tan^{-1}(-x) = -\tan^{-1}x, x \in \mathbf{R}$

- **$\operatorname{cosec}^{-1}(-x) = -\operatorname{cosec}^{-1}x, |x| \geq 1$**

- $\sec^{-1}(-x) = \pi - \sec^{-1}x, |x| \geq 1$
- $\cot^{-1}(-x) = \pi - \cot^{-1}x, x \in \mathbf{R}$

- For suitable values of domains;

$$\circ \sin^{-1}x + \cos^{-1}x = \frac{\pi}{2}, x \in [-1, 1]$$

$$\circ \tan^{-1}x + \cot^{-1}x = \frac{\pi}{2}, x \in \mathbf{R}$$

$$\circ \sec^{-1}x + \operatorname{cosec}^{-1}x = \frac{\pi}{2}, |x| \geq 1$$

- For suitable values of domains;

$$\tan^{-1}x + \tan^{-1}y = \begin{cases} \tan^{-1} \frac{x+y}{1-xy}, xy < 1 \\ \pi + \tan^{-1} \frac{x+y}{1-xy}, xy > 1 \end{cases}$$

$$\circ \tan^{-1}x + \tan^{-1}y = \tan^{-1} \frac{x-y}{1+xy}$$

**Note:** While solving problems, we generally use the

formula  $\tan^{-1}x + \tan^{-1}y = \tan^{-1} \frac{x+y}{1-xy}$  when the condition for  $xy$  is not given.

$$\bullet \text{ For } x \in [-1, 1], 2\tan^{-1}x = \sin^{-1} \frac{2x}{1+x^2}$$

$$\bullet \text{ For } x \in (-1, 1), 2\tan^{-1}x = \tan^{-1} \frac{2x}{1-x^2}$$

$$\bullet \text{ For } x \in [0, 1], 2\tan^{-1}x = \cos^{-1} \frac{1-x^2}{1+x^2}$$

**Example: 2** For  $x, y \in [-1, 1]$ , show that:  $\sin^{-1}x + \sin^{-1}y = \sin^{-1}(x\sqrt{1-y^2} + y\sqrt{1-x^2})$

**Solution:** We know that  $\sin^{-1}x$  and  $\sin^{-1}y$  can be defined only for  $x, y \in [-1, 1]$

Let  $\sin^{-1}x = a$  and  $\sin^{-1}y = b$

$\Rightarrow x = \sin a$  and  $y = \sin b$

Also,  $\cos a = \sqrt{1-x^2}$  and  $\cos b = \sqrt{1-y^2}$

We know that,  $\sin(a+b) = \sin a \cos b + \cos a \sin b$

$$\Rightarrow a+b = \sin^{-1} [x\sqrt{1-y^2} + y\sqrt{1-x^2}]$$

$$\Rightarrow \sin^{-1}x + \sin^{-1}y = \sin^{-1} [x\sqrt{1-y^2} + y\sqrt{1-x^2}]$$

**Example: 3** If  $\tan^{-1}(\frac{5}{6}) + \tan^{-1}(\frac{1}{11}) = x$ , then find  $\sec x$ .

**Solution:**

$$x = \tan^{-1}(\frac{5}{6}) + \tan^{-1}(\frac{1}{11}) = \tan^{-1} \left[ \frac{\frac{5}{6} + \frac{1}{11}}{1 - \frac{5}{6} \times \frac{1}{11}} \right]$$

We have

Using the identity  $\tan^{-1}x + \tan^{-1}y = \tan^{-1}\left(\frac{x+y}{1-xy}\right)$ , where  $x = \frac{5}{6}$  and  $y = \frac{1}{11}$

$$\therefore x = \tan^{-1}\left[\frac{\frac{55+6}{66}}{\frac{66-5}{66}}\right]$$

$$= \tan^{-1}1$$

$$= \frac{\pi}{4}$$

$$\sec x = \sec \frac{\pi}{4} = \sqrt{2}$$

**Example: 4**

Show that:  $3\tan^{-1}x = \tan^{-1}\left(\frac{3x-x^3}{1-3x^2}\right)$  where  $x \in \left(-\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}\right)$

**Solution:** We know that,

$$3\tan^{-1}x = \tan^{-1}x + 2\tan^{-1}x$$

$$= \tan^{-1}x + \tan^{-1}\frac{2x}{1-x^2}$$

$$= \tan^{-1}\left[\frac{x + \frac{2x}{1-x^2}}{1-x \times \frac{2x}{1-x^2}}\right]$$

$$= \tan^{-1}\left[\frac{\frac{3x-x^3}{1-x^2}}{\frac{1-3x^2}{1-x^2}}\right]$$

$$= \tan^{-1}\left(\frac{3x-x^3}{1-3x^2}\right)$$