


Introduction to Trigonometry

- **Trigonometric Ratio**



$$\sin \theta = \frac{\text{Opposite side}}{\text{Hypotenuse}} = \frac{AB}{AC}$$

$$\cos \theta = \frac{\text{Adjacent side}}{\text{Hypotenuse}} = \frac{BC}{AC}$$

$$\tan \theta = \frac{\text{Opposite side}}{\text{Adjacent side}} = \frac{AB}{BC}$$

$$\csc \theta = \frac{1}{\sin \theta} = \frac{\text{Hypotenuse}}{\text{Opposite side}} = \frac{AC}{AB}$$

$$\sec \theta = \frac{1}{\cos \theta} = \frac{\text{Hypotenuse}}{\text{Adjacent side}} = \frac{AC}{BC}$$

$$\cot \theta = \frac{1}{\tan \theta} = \frac{\text{Adjacent side}}{\text{Opposite side}} = \frac{BC}{AB}$$


$$\text{Also, } \tan \theta = \frac{\sin \theta}{\cos \theta}, \cot \theta = \frac{\cos \theta}{\sin \theta}$$

If one of the trigonometric ratios of an acute angle is known, then the remaining trigonometric ratios of the angle can be calculated.

Example:

If $\sin \theta = \frac{7}{25}$, then find the value of $\sec \theta(1 + \tan \theta)$.

Solution:



It is given that $\sin \theta = \frac{7}{25}$

$$\sin \theta = \frac{\text{AB}}{\text{AC}} = \frac{7}{25}$$

$\Rightarrow AB = 7x$ and $AC = 25x$, where x is some positive integer

By applying Pythagoras theorem in ΔABC , we get:

$$AB^2 + BC^2 = AC^2$$

$$\Rightarrow (7x)^2 + BC^2 = (25x)^2$$

$$\Rightarrow 49x^2 + BC^2 = 625x^2$$

$$\Rightarrow BC^2 = 625x^2 - 49x^2$$

$$\Rightarrow BC = \sqrt{576}x = 24x$$

$$\therefore \sec \theta = \frac{\text{Hypotenuse}}{\text{Adjacent side}} = \frac{25}{24}$$

$$\tan \theta = \frac{\text{Opposite side}}{\text{Adjacent side}} = \frac{7}{24}$$

$$\therefore \sec \theta (1 + \tan \theta) = \frac{25}{24} \left(1 + \frac{7}{24} \right) = \frac{25}{24} \times \frac{31}{24} = \frac{775}{576}$$

- Use trigonometric ratio in solving problem.

Example:

If $\tan \theta = \frac{3}{5}$, then find the value of $\frac{\sin \theta + \cos \theta}{\sin \theta - \cos \theta}$

Solution:

$$\frac{\sin \theta + \cos \theta}{\sin \theta - \cos \theta}$$

Take $\cos \theta$ common from numerator and denominator both

$$= \frac{\frac{\sin \theta}{\cos \theta} + 1}{\frac{\sin \theta}{\cos \theta} - 1}$$

$$= \frac{\tan \theta + 1}{\tan \theta - 1}$$

$$= \frac{\frac{3}{5} + 1}{\frac{3}{5} - 1}$$

$$= \frac{\frac{3+5}{5}}{\frac{3-5}{5}}$$

$$= \frac{8}{-2}$$

$$= -4$$


- Trigonometric Ratios of some specific angles

q	0	30°	45°	60°	90°
$\sin q$	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1
$\cos q$	1	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	0
$\tan q$	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	Not defined
$\operatorname{cosec} q$	Not defined	2	$\sqrt{2}$	$\frac{2}{\sqrt{3}}$	1
$\sec q$	1	$\frac{2}{\sqrt{3}}$	$\sqrt{2}$	2	Not defined
$\cot q$	Not defined	$\sqrt{3}$	1	$\frac{1}{\sqrt{3}}$	0

Example 1:

ΔABC is right-angled at B and $AB = 6$ m, $BC = \sqrt{12}$ m. Find the measure of $\angle A$ and $\angle C$.

Solution:



$$AB = 6 \text{ m}$$

$$BC = \sqrt{12} \text{ m} = 2\sqrt{3} \text{ m}$$

$$\tan C = \frac{\text{Opposite side}}{\text{Adjacent side}} = \frac{AB}{BC} = \frac{6}{2\sqrt{3}} = \sqrt{3}$$

$$\Rightarrow \tan C = \tan 60^\circ \quad \left[\because \tan 60^\circ = \sqrt{3} \right]$$

$$\Rightarrow \angle C = 60^\circ$$

$$\therefore \angle A = 180^\circ - (90 + 60) = 30^\circ$$

Example 2:

Evaluate the expression

$$4(\cos^3 60^\circ - \sin^3 30^\circ) + 3(\sin 30^\circ - \cos 60^\circ)$$

Solution:

$$4(\cos^3 60^\circ - \sin^3 30^\circ) + 3(\sin 30^\circ - \cos 60^\circ)$$

$$= 4 \left[\left(\frac{1}{2}\right)^3 - \left(\frac{1}{2}\right)^3 \right] + 3 \left(\frac{1}{2} - \frac{1}{2} \right)$$

$$= 4 \times 0 + 3 \times 0 = 0 + 0 = 0$$

- **Trigonometric Ratios of Complementary Angles**

$$\sin(90^\circ - \theta) = \cos \theta \quad \cos(90^\circ - \theta) = \sin \theta$$

$$\tan(90^\circ - \theta) = \cot \theta \quad \cot(90^\circ - \theta) = \tan \theta$$

$$\operatorname{cosec}(90^\circ - \theta) = \sec \theta \quad \sec(90^\circ - \theta) = \operatorname{cosec} \theta$$

Where θ is an acute angle.

Example 1: Evaluate the expression

$$\sin 28^\circ \sin 30^\circ \sin 54^\circ \sec 36^\circ \sec 62^\circ$$

Solution:

$$\begin{aligned}
& \sin 28^\circ \sin 30^\circ \sin 54^\circ \sec 36^\circ \sec 62^\circ \\
& = (\sin 28^\circ \sec 62^\circ)(\sin 54^\circ \sec 36^\circ) \sin 30^\circ \\
& = \{\sin 28^\circ \csc(90^\circ - 62^\circ)\} \{\sin 54^\circ \csc(90^\circ - 36^\circ)\} \sin 30^\circ \\
& = (\sin 28^\circ \csc 28^\circ)(\sin 54^\circ \csc 54^\circ) \sin 30^\circ \\
& = \left(\frac{1}{\sin 28^\circ} \right) \left(\frac{1}{\sin 54^\circ} \right) \times \frac{1}{2} \\
& = \frac{1}{2}
\end{aligned}$$

Example 2: Evaluate the expression

$$4\sqrt{3}(\sin 40^\circ \sec 30^\circ \sec 50^\circ) + \frac{\sin^2 34^\circ + \sin^2 56^\circ}{\sec^2 31^\circ - \cot^2 59^\circ}$$

Solution:

$$\begin{aligned}
& 4\sqrt{3}(\sin 40^\circ \sec 30^\circ \sec 50^\circ) + \frac{\sin^2 34^\circ + \sin^2 56^\circ}{\sec^2 31^\circ - \cot^2 59^\circ} \\
& = 4\sqrt{3} [\sec 30^\circ (\sin 40^\circ \sec 50^\circ)] + \frac{\sin^2 34^\circ + \sin^2(90^\circ - 56^\circ)}{\sec^2 31^\circ - \tan^2(90^\circ - 59^\circ)} \\
& \quad \left[\because \cos(90^\circ - \theta) = \sin \theta, \tan(90^\circ - \theta) = \cot \theta \right] \\
& = 4\sqrt{3} [\sec 30^\circ \sin 40^\circ \csc(90^\circ - 50^\circ)] + \frac{\sin 34^\circ + \cos^2 34^\circ}{\sec^2 31^\circ - \tan^2 31^\circ} \\
& = 4\sqrt{3} \left[\frac{2}{\sqrt{3}} \sin 40^\circ \csc 40^\circ \right] + \frac{1}{1} \\
& = 8 + 1 = 9
\end{aligned}$$

- **Trigonometric Identities**

1. $\cos^2 A + \sin^2 A = 1$
2. $1 + \tan^2 A = \sec^2 A$
3. $1 + \cot^2 A = \csc^2 A$

Example:

If $\cos \theta = \frac{5}{7}$, find the value of $\cot \theta + \csc \theta$

Solution:

We have, $\cos \theta = \frac{5}{7}$

Now, $\sin^2 \theta + \cos^2 \theta = 1$

$$\therefore \sin \theta = \sqrt{1 - \cos^2 \theta}$$

$$= \sqrt{1 - \left(\frac{5}{7}\right)^2}$$

$$= \sqrt{\frac{49-25}{49}} = \frac{2\sqrt{6}}{7}$$

$$\therefore \operatorname{cosec} \theta = \frac{7}{2\sqrt{6}}$$

$$\text{Also, } \cot \theta = \frac{\cos \theta}{\sin \theta}$$

$$= \frac{\frac{5}{7}}{\frac{2\sqrt{6}}{7}} = \frac{5}{2\sqrt{6}}$$

$$\therefore \cot \theta + \operatorname{cosec} \theta = \frac{5}{2\sqrt{6}} + \frac{7}{2\sqrt{6}}$$

$$= \frac{12}{2\sqrt{6}} = \frac{6}{\sqrt{6}} \times \frac{\sqrt{6}}{\sqrt{6}}$$

$$= \sqrt{6}$$

- **Use of trigonometric identities in proving relationships involving trigonometric ratio.**

Example: Prove the following identities

$$\tan^2 \theta + \cot^2 \theta + 2 = \sec^2 \theta \operatorname{cosec}^2 \theta$$

Solution:

We have

$$\begin{aligned}
 \text{LHS} &= \tan^2 \theta + \cot^2 \theta + 2 \\
 &= \tan^2 \theta + \cot^2 \theta + 2 \cdot \tan \theta \cdot \cot \theta \quad [\because \tan \theta \cdot \cot \theta = 1] \\
 &= \left(\tan \theta + \cot \theta \right)^2 \quad \left[\because a^2 + b^2 + 2ab = (a+b)^2 \right] \\
 &= \left(\frac{\sin \theta}{\cos \theta} + \frac{\cos \theta}{\sin \theta} \right)^2 \\
 &= \left(\frac{\sin^2 \theta + \cos^2 \theta}{\sin \theta \cdot \cos \theta} \right)^2 \\
 &= \left(\frac{1}{\sin \theta \cdot \cos \theta} \right)^2 \\
 &= \left(\sec \theta \cdot \csc \theta \right)^2 \\
 &= \sec^2 \theta \cdot \csc^2 \theta \\
 &= \text{RHS}
 \end{aligned}$$