## Linear equation in two variables:

An equation of the form, ax + by + c = 0, where a, b and c are constants, such that a and b are both not zero and x and y are variables is called a linear equation in two variables. For example, 2x + 3y + 10 = 0, 3x + 7y = 0

• Real life situations can be expressed mathematically as linear equations in two variables.

**Example:** The age of Ram is 3 more than twice the age of Mohan. Write a linear equation in two variables to represent this statement.

**Solution:** Let the age of Mohan be *x* years and the age of Ram be *y* years. Thus, the given condition can be expressed as y = 2x + 3

## Solution of a linear equation in two variables:

The values of the variables in a linear equation, which satisfy the equation are the solutions of that linear equation.

- A linear equation in two variables has infinitely many solutions.
- Solution of linear equation in two variables can be found by substitution method.

**Example:** Find two different solutions of the equation 4x + 5y = 20.

```
Solution: Given equation is 4x + 5y = 20.
If we take x = 0, we obtain:
4 \times 0 + 5y = 20
\Rightarrow 5y = 20
\Rightarrow y = 4
So, (0, 4) is a solution of the given equation.
```

If we take y = 0, we obtain:  $4x + 5 \times 0 = 20$   $\Rightarrow 4x = 20$   $\Rightarrow x = 5$ So, (5, 0) is a solution of the given equation.

- The geometrical representation of the linear equation, ax + by + c = 0, is a straight line.
- In order to represent a linear equation in two variables graphically, its two or three different points are calculated and then the corresponding points are plotted and joined on

Get More Learning Materials Here :





the coordinate plane.

**Example:** Represent x + 3y = 6 on a graph paper.

**Solution:** The given equation is x + 3y = 6

For x = 0,  $3y = 6 \Rightarrow y = \frac{6}{3} = 2$ For x = 3,  $3 + 3y = 6 \Rightarrow 3y = 3 \Rightarrow y = 1$ 

x	0	3
У	2	1

By plotting (0, 2) and (3, 1) on coordinate plane and then joining them, the given equation can be represented as:



- An equation of the form, *y* = *mx*, represents a line passing through the origin.
- Graphical solution of linear equation in two variables:

Every point on the graph of a linear equation in two variables is a solution of the linear equation and moreover, every solution of the linear equation is a point on the graph of the linear equation.

**Example:** A bag contains some Re 1 coins and some Rs 2 coins. The total worth of coins is Rs 45. Find the number of Re 1 coins, if there are 10 coins of Rs 2.

Solution: Let there be x coins of Re 1 and y coins of Rs 2.Thus, 1x + 2y = 45 $\Rightarrow x + 2y = 45$ This is the required linear equation of the given information. The three solutions of thisequation have been given in the tabular form as follows:x51525

Get More Learning Materials Here : 📕







From the above graph, it can be seen that the value of *x* corresponding to y = 10 is 25. Therefore, there are 25 coins of Re 1, if there are 10 coins of Rs 2.

- The graph of *x* = *a* is a straight line parallel to the *y*-axis, situated at a distance of *a* units from *y*-axis.
- The graph of *y* = *b* is a straight line parallel to the *x*-axis, situated at a distance of *b* units from *x*-axis.

**Example:** Represent the equation 2y + 5 = 0, on Cartesian plane.

Solution: 2y + 5 = 0  $\Rightarrow 2y = -5$  $\Rightarrow y = \frac{-5}{2} = -2.5$ , which is of the form y = b.

The graph of this equation can be drawn as follows:

Get More Learning Materials Here :







Get More Learning Materials Here : 📕



